

We are very grateful to the referees for their supportive comments on this work, and for their helpful suggestions for modifications and improvements. Responses to the comments are now provided (the original comments are shown in blue font), along with the revised manuscript (showing tracked changes).

A. Comments by Craig Taatjes

General comments:

This work assesses kinetics for the fast-moving field of carbonyl oxide chemistry. The evaluation is thorough and thoughtful and accurately captures and accounts for most of the questions and disagreements that arise from the available studies. The evaluation makes insightful recommendations of the key points for additional research (including branching fractions for some key reactions). The manuscript, although an experiment based evaluation, also points out the key role of advanced theoretical methods, which now can produce highly accurate kinetics values for experimentally inaccessible reactions. This is an excellent work that will be very useful to modelers and as a guide to experimental kineticists. There are just a few improvements that I would suggest, focused on the aspect I am most familiar with, reaction kinetics of carbonyl oxides.

We thank the referee for these positive comments on our evaluation; and for the helpful suggestions for improvements and clarifications, which are dealt with in the responses below.

Specific comments:

Comment A1: First, new investigations have continued since the evaluation that forms the basis of this paper, especially in the reaction kinetics of the stabilized carbonyl oxides. That is unavoidable. However, the authors may consider the newest investigations (PNAS 117 (18), 9733-9740 (2020)) of the kinetics of the C₄ carbonyl oxide methyl vinyl ketone oxide, because here theory and direct experiments suggest that the bimolecular reactions of the conjugated carbonyl oxides may differ from the reactions used as analogies in the evaluation. For example, direct kinetics measurements show that the reaction of methyl vinyl ketone oxide with SO₂ is a factor of three or so slower than the reactions of non-conjugated carbonyl oxides of similar size.

Response: We are grateful to the referee for highlighting the study of Caravan et al. (2020) on the kinetics of methyl vinyl ketone (MVK) oxide, which was published at around the time our paper was submitted. We had anticipated that such studies were likely imminent (page 11, line 4) and are pleased to see that important new work on structurally-complex sCIs is already emerging.

We have since expanded the evaluation to include the reactions of *syn*-MVK oxide (*E*-(CH=CH₂)(CH₃)COO) with SO₂ and HC(O)OH, data sheets for which (CGI_22 and CGI_30, respectively) are now available at <http://iupac.pole-ether.fr/>, and included in Supplement B of the revised manuscript. Our preferred values for these rate coefficients are thus based on the determinations of Caravan et al. (2020), and appear in Table 5 in the revised manuscript. The work is now also referred to in Sects. 6.1 and 6.2 of the revised manuscript, where the kinetics of structurally complex sCIs are discussed. Thus, in Sect. 6.1 (*Measurements of absolute rate coefficients for reactions of sCIs*), the following text appears (new or adjusted text in red font):

“As indicated in Sect. 5.2, methods for **the** production and spectroscopic characterization of **more complex isoprene-derived** species are emerging (e.g. Barber et al., 2018; Vansco et al., 2018; 2019), **and these have provided the basis for their direct kinetics study (Caravan et al., 2020).** However, direct experimental determinations of rate coefficients have not yet been reported for **larger** complex species (**> C₄**), **derived for example from monoterpenes and sesquiterpenes.**”

and in Sect. 6.2 (*Evaluation of rate coefficients for bimolecular reactions*):

“**With exception of the reactions of *E*-(CH=CH₂)(CH₃)COO with SO₂ and HC(O)OH (Caravan et al., 2020),** there are currently no direct kinetics determinations for reactions of the C₄ sCIs derived from isoprene.”

This has also led to a minor revision to the analysis presented in data sheet CGI_21 on isoprene-derived sCIs (included in Supplement B), which assessed the parameters assigned to the ensemble of unimolecular decomposition reactions and bimolecular reactions with SO₂, H₂O and (H₂O)₂, using the results of published chamber and slow-flow experiments (Sipilä et al., 2014; Newland et al., 2015; Nguyen et al., 2016).

Comment A2: Second, as the manuscript acknowledges, measuring thermal kinetics for unimolecular reactions of carbonyl oxides is often experimentally difficult because of the need to correct for competing bimolecular processes. In this case the theoretical characterizations, especially those for which microcanonical rate coefficients and the related tunneling parameters have been corroborated by direct energy-specific dynamics measurements, provide valuable information about the unimolecular processes (see Int. Rev. Phys. Chem. 39 (1), 1-33 (2020)). The evaluation already makes good use of

theory in interpreting the unimolecular kinetics, but the authors might consider emphasizing experimentally validated calculations (J. Chem. Phys. 146, 134307 (2017)) for acetone oxide ($k_{298} = 276 \text{ s}^{-1}$), where the tunneling-adjusted microcanonical rate coefficients match experiment over a wide energy range, and J. Chem. Phys. 145, 234308 (2016) that treats the unimolecular decay of Z-acetaldehyde oxide with similar tests of tunneling parameters against direct experiment ($k_{298} = 122 \text{ s}^{-1}$). These values are slightly lower than the recommendations. I note that the unimolecular decay kinetics for some methyl vinyl ketone oxide conformers (J. Am. Chem. Soc., 140 (34), 10866–10880 (2018)) have also been compared to energy-resolved dynamics measurements..

Response: We thank the referee for alerting us to the recent review of Stephenson and Lester (2020) on unimolecular decay dynamics of Criegee intermediates, which we now refer to in Sect. 7 (*Overall reactivity conclusions – comparison of experiment and theory*) in the revised manuscript.

We also agree that the studies of Fang et al. (2016; 2017) on Z-CH₃CHOO and (CH₃)₂COO, and Barber et al. (2018) on *syn*-MVK oxide (*E*-(CH=CH₂)(CH₃)COO), should have been cited in the manuscript in relation to the comparison of energy-resolved dynamics measurements with theoretical estimates. The following information therefore now also appears Sect. 7 (new text in red font):

“However, the body of experimental information now available for a series of sCIs allows the results of theoretical studies to be validated, and for the methods to be refined, optimised and extended. This has included comparison of experimental and theoretical unimolecular decay rates of a number of infra-red activated sCIs for a range of excitation energies (e.g. Fang et al., 2016; 2017; Barber et al., 2018).”

In addition to the main manuscript, it should be noted that the results of theoretical studies were already discussed (in the “*comments on preferred values*”) in all the unimolecular decay reaction data sheets in Supplement B, thereby providing an indication of where theoretical calculations have been experimentally validated. Reference to Barber et al. (2018), for example, thus also appears in data sheet CGI_21, relating to the unimolecular decay kinetics of *syn*-MVK oxide, *E*-(CH=CH₂)(CH₃)COO, where the following text appears:

“Barber et al. (2018) also report HO formation from the thermal decomposition of *E*-(CH=CH₂)(CH₃)COO, compatible with the expected pathway involving 1,4 H-atom transfer from the -CH₃ group to form a vinyl hydroperoxide intermediate, and at a rate that is consistent with theory.”

It should also be noted that the results of a large number of theoretical studies are taken into account in the comprehensive theoretical/SAR study of Vereecken et al. (2017), which is mainly used for the experiment/theory comparisons that we present in the manuscript.

Comment A3: Third, in the discussion of the atmospheric role of carbonyl oxides, are there other non-kinetics uncertainties (e.g., in the alkene source inventory) that should be mentioned? The development of sensitive measurement methods for carbonyl oxides (e.g., J. Am. Chem. Soc. 139 (38), 13387–13392 (2017)) that may eventually constrain the concentration of these intermediates in the field should possibly be mentioned as an important area for continued effort.

Response: The referee raises two valid points, both of which have been addressed in the revised manuscript. An additional section has now been included at the end of Supplement C to discuss uncertainties in the steady state calculations, including potentially missing sources and sinks. The following text has also been added to the introductory paragraph of Sect. 8 (*Impact of Criegee intermediates in atmospheric oxidation chemistry*):

“Although the calculations presented here aim to take account of the most important production and loss routes, the estimates are inevitably subject to potential omissions and uncertainties in the sources and sinks of the sCIs (as discussed further in Sect. C4), in addition to uncertainties associated with the kinetic parameters and sCI yields, as discussed in earlier sections and in the data sheets in Supplements A and B.”

Regarding measurement methods, the following text has now been included at the end of Sect. 8:

“In addition, the continued development of sensitive detection methods for sCIs (e.g. Berndt et al., 2017), that may eventually allow sCI concentrations to be measured in the field, would be valuable for evaluation and testing of the representation of sCI sources and sinks in atmospheric models.”

Comment A4: Finally – is there a reference to verify a unity quantum yield of O (1D) from UV excitation of carbonyl oxides (section 5)?

Response: We agree that further information is required here. We have therefore replaced the relevant sentence in Sect. 5 with the text shown below, specifically referring to information for CH₂OO:

“Photodissociation of CH₂OO from this excitation is reported to proceed with a quantum yield of unity (e.g. Ting et al., 2014), producing HCHO and either O(³P) or O(¹D). Production of O(¹D) has been reported to be dominant (e.g. Li et al., 2015), and exclusive in the long wavelength tail ($\lambda \geq 364 \text{ nm}$; Vansco et al., 2017).”

B. Comments by Referee 2

Opening comment:

The manuscript provides a comprehensive assessment of gas-phase Criegee intermediate chemistry and photochemistry. Kinetics measurements are summarized thoroughly for ozonolysis of a broad range of unsaturated VOCs, the reactions of stabilized Criegee intermediates with a selection of trace gases, and unimolecular decomposition reactions. Where appropriate, theoretical work is also referenced to support the assessment of experimental studies. Overall, this is an excellent and thorough summary of our current understanding of Criegee intermediate chemistry in the atmosphere.

I have only very minor suggestions for the authors.

Response: We thank the referee for these positive comments on our work, and for the suggestions for improvements, which are dealt with in the following responses.

Comment B1: First, the discovery of the UV spectrum of formaldehyde oxide that is attributed to Sheps [J. Phys. Chem. Lett. 4, 4201 (2013)] on page 10 should more properly be attributed to Beames et al. [J. Am. Chem. Soc. 134, 20045 (2012)].

Response: We thank the referee for pointing this out. Because the relevant text is discussing the application of UV absorption spectroscopy to the study of bimolecular reaction kinetics at ambient temperatures and pressures, we feel the study of the UV absorption spectrum (and kinetics) of Sheps (2013) at 295 K is a more appropriate reference at this point than the prior study of the UV photodissociation action spectrum of Beames et al. (2012) in a cooled supersonic expansion. However, the referee is quite correct that the “discovery” of the UV spectrum is more properly attributed to Beames et al. (2012). Noting that the sentence already specifies “absorption spectrum”, we have therefore replaced the word “discovery” with “characterization”.

As indicated below in our *additional changes* section, we have now also added reference to both Beames et al. (2012) and Sheps (2013) in an earlier section (*Sect. 5.2: UV spectra of stabilised Criegee intermediates*).

Comment B2: Second, while discussing on page 9 apparent discrepancies between measurements of the UV absorption spectrum the authors comment on the lack of detailed data on the temperature dependence of the cross sections. Foreman et al. [Phys. Chem. Chem. Phys. 17, 32539 (2015)] demonstrated that the spectra were independent of temperature over the range 276-357 K.

Response: The referee is correct that Foreman et al. (2015) report temperature-dependent measurements over the range 276-357 K in the long-wavelength tail for CH₂OO. However, the point we make remains valid, because this temperature range does not encompass the very low temperatures reached in molecular beams. We therefore now make this clear in the revised text at the appropriate point:

“... the lack of detailed data on the temperature dependence of the cross sections **over the required range** precludes firm conclusions to be drawn.”

As a result of the referee’s comment, we realised that the lack of the temperature dependence reported by Foreman et al. (2015) was not previously explicitly stated in the relevant data sheet, P33 (in Supplement B), and a statement to this effect has now therefore been included.

Comment B3: Third, the range of reactions of stabilized Criegee intermediates covered in the assessment is somewhat smaller than that compiled by Khan et al. [Environ. Sci.: Processes Impacts 20, 437 (2018)]. The authors may want to comment explicitly on why they have focused on the more limited set of reactions.

Response: The referee is correct that, although we clearly state (e.g. in Sect. 1 and Fig. 2) which sCIs are considered in the evaluation, the reaction coverage is less well explained and justified. In Sect. 1, the following text (slightly adjusted because of the recent data for *E*-(CH=CH₂)(CH₃)COO – see comment A1) provides a broad indication of the scope of the evaluation relating to the reactions of sCIs:

“The spectroscopy and kinetics recommendations for the sCI reactions are presented and discussed in Sects. 5 and 6. These include data for bimolecular and unimolecular reactions of selected sCIs of particular atmospheric relevance for which direct kinetic data have been reported, namely CH₂OO, *Z*- and *E*- CH₃CHOO, (CH₃)₂COO **and *E*-(CH=CH₂)(CH₃)COO**; and we also provide some discussion of the **complete set of C₄** intermediates formed from isoprene (see Fig. 2 for sCI structures)”

In Sect. 6.2 (*Evaluation of rate coefficients for bimolecular reactions*), we have now added the following explicit information on reaction coverage (new text in red font):

“Table 5 provides a summary of the preferred values of bimolecular reaction rate coefficients, with additional details given in the corresponding reaction data sheets in Supplement B. **As indicated in Sect. 1, the evaluation focuses on classes of reaction that are of particular significance in tropospheric chemistry. Where data are available, these include reactions with SO₂, NO₂, H₂O, (H₂O)₂, CH₃CHO, CH₃C(O)CH₃, CF₃C(O)CF₃, HC(O)OH, CH₃C(O)OH and**

CF₃C(O)OH for the set of sCIs, with some additional reactions also considered for CH₂OO. It is noted that data have been reported for some other classes of reaction (e.g. with alkenes and alcohols) that are uncompetitive under tropospheric conditions, and some relative rate data are also available for large sCIs not considered here, e.g. as summarised in the compilation reported by Khan et al. (2018).”

The referee is correct that there are some reaction classes that are included in the tabulations of Khan et al. (2018) which we do not consider, these being reactions that are uncompetitive under tropospheric conditions; and we are grateful for the opportunity of clarifying this in the revised manuscript.

It should be noted, however, that our evaluation also includes some additional reactions and information not provided by Khan et al. (2018). These include speciated recommendations for, and discussion of, the complete set of four isoprene-derived C₄ sCIs for unimolecular decomposition and bimolecular reaction with SO₂, H₂O and (H₂O) (i.e. 16 reactions), for which Khan et al. (2018) only tabulate a single bulk parameter (not significantly different from zero) for unimolecular decomposition of the C₄ set collectively (from Newland et al. (2015)). In addition, we evaluate the data and provide a recommended rate coefficient for each reaction considered. In contrast, Khan et al. (2018) tabulate most laboratory and some theoretical determinations of the rate coefficients to inform their review and discussion of the atmospheric chemistry of sCIs, but provide no recommendations or preferred values.

C. Additional changes

The chemistry of sCIs is a fast-moving field, and a number of new studies have been reported during the open discussion of the paper. Some of these have been raised by the referees, and have been dealt with as described above. In addition, we have responded to publication of other studies by implementing the following changes and additions:

(i) **Yields of stabilised Criegee intermediates:** Newland et al. (2020) recently reported a systematic study of total sCI yields, and speciated yields of stabilised CH₂OO and (CH₃)₂COO, from the reactions of O₃ with a series of alkenes. These have resulted in small changes to the recommended total sCI yields for propene and 2-methylpropene in Table 2, and a new entry for β-myrcene. This also alerted us to a prior study of Newland et al. (2018), that had been overlooked, resulting in further small changes to the recommended total sCI yields for β-pinene and limonene. The underlying information is also summarised and discussed in the relevant O₃ + alkene data sheets, which have been revised accordingly.

(ii) **Kinetics of CH₂OO decomposition and its reaction with HC(O)OH:** Peltola et al. (2020) have recently reported a kinetics study of CH₂OO. This was produced using a new source, namely the photolysis of CH₂IBr. As a result, the following change has been made to the text in Sect. 5.2 (*UV spectra of stabilised Criegee intermediates*):

“In experimental studies of UV-visible spectra, the series of C₁ – C₃ sCIs have **mainly** been formed by photolysis of the corresponding di-iodoalkane (via C-I bond fission), followed by the reaction of the iodoalkyl radical with O₂, e.g. in the case of CH₂OO (e.g. Beames et al., 2012; Sheps, 2013):



The formation of CH₂I (and subsequently CH₂OO) from the photolysis of CH₂IBr (via C-Br bond fission) has also recently been reported (Peltola et al., 2020), suggesting that bromo-iodoalkanes may also be used more widely as sCI precursors.”

The reported kinetic data for CH₂OO decomposition and its reaction with HC(O)OH have been included in data sheets CGI_12 and CGI_11. This has resulted in changes to the recommended rate coefficients, as shown in Table 5 of the revised manuscript. In the case of CH₂OO decomposition (CGI_12), the recommended rate coefficient has been significantly simplified to an upper limit value at 298 K and 1 bar only. This is also discussed in Sect 7 (*Overall reactivity conclusions – comparison of experiment and theory*), where the value is now compared with that calculated for the same conditions by Long et al. (2016). However, this revision does not change the main conclusion that CH₂OO decomposition is an unimportant loss process, compared with other routes, under tropospheric conditions

References

Barber, V. P., Pandit, S., Green, A. M., Trongsiwat, N., Walsh, P. J., Klippenstein, S. J. and Lester, M. I.: Four-carbon Criegee intermediate from isoprene ozonolysis: Methyl vinyl ketone oxide synthesis, infrared spectrum, and OH production, J. Am. Chem. Soc., 140, 10866-10880, <https://doi.org/10.1021/jacs.8b06010>, 2018.

Beames, J. M., Liu, F., Lu, L., and Lester, M. I.: Ultraviolet spectrum and photochemistry of the simplest Criegee intermediate CH₂OO J. Amer. Chem. Soc., 134, 20045-20048, <https://doi.org/10.1021/ja310603j>, 2012.

Berndt, T., Herrmann, H. and Kurtén, T.: Direct probing of Criegee intermediates from gas-phase ozonolysis Using chemical ionization mass spectrometry, *J. Am. Chem. Soc.*, 139, 13387–13392, 2017.

Caravan, R. L., Vansco, M. F., Au, K., Khan, M. A. H., Li, Y.-L. Winiberg, F. A. F., Zuraski, K., Lin, Y.-H., Chao, W., Trongsiwat, N., Walsh, P. J., Osborn, D. L., Percival, C. J., Lin, J.-J. M., Shallcross, D. E., Sheps, L., Klippenstein, S. J., Taatjes, C. A., and Lester, M. I.: *Proc. Nat. Acad. Sci.*, 117 (18), 9733-9740, doi: 10.1073/pnas.1916711117, 2020.

Fang, Y., Liu, F., Barber, V. P., Klippenstein, S. J., McCoy, A. B. and Lester, M. I.: Deep tunneling in the unimolecular decay of CH_3CHOO Criegee intermediates to OH radical products, *J. Chem. Phys.* 145, 234308, 2016.

Fang, Y., Liu, F., Barber, V. P., Klippenstein, S. J., McCoy, A. B. and Lester, M. I.: Tunneling effects in the unimolecular decay of $(\text{CH}_3)_2\text{COO}$ Criegee intermediates to OH radical products, *J. Chem. Phys.* 146, 134307, <https://doi.org/10.1063/1.4979297>, 2017.

Foreman, E. S., Kapnas, K. M., Jou, Y., Kalinowski, J., Feng, D., Gerber, R. B., and Murray, C.: High resolution absolute absorption cross sections of the $\text{B}^1\text{A}'\text{--X}^1\text{A}'$ transition of the CH_2OO biradical, *Phys. Chem. Chem. Phys.*, 17, 32539-32546, 2015.

Khan, M. A. H., Percival, C. J., Caravan, R. L., Taatjes, C. A. and Shallcross, D. E.: Criegee intermediates and their impacts on the troposphere, *Environ. Sci.: Processes Impacts*, 20, 437-453, 2018.

Li, H., Fang, Y., Beames, J. M. and Lester, M. I.: Velocity map imaging of O-atom products from UV photodissociation of the CH_2OO Criegee intermediate, *J. Chem. Phys.*, 142, 214312, <https://doi.org/10.1063/1.4921990>, 2015.

Long, B., Bao, J. L. and Truhlar, D.G.: Atmospheric chemistry of Criegee intermediates: Unimolecular reactions and reactions with water, *J. Am. Chem. Soc.* 138, 43, 14409-14422, <https://doi.org/10.1021/jacs.6b08655>, 2016.

Newland, M. J., Rickard, A. R., Vereecken, L., Muñoz, A., Ródenas, M., and Bloss, W. J.: Atmospheric isoprene ozonolysis: impacts of stabilised Criegee intermediate reactions with SO_2 , H_2O and dimethyl sulfide, *Atmos. Chem. Phys.*, 15, 9521-9536, <https://doi.org/10.5194/acp-15-9521-2015>, 2015b.

Newland, M. J., Rickard, A. R., Sherwen, T., Evans, M. J., Vereecken, L., Muñoz, A., Ródenas, M., and Bloss, W. J.: The atmospheric impacts of monoterpene ozonolysis on global stabilised Criegee intermediate budgets and SO_2 oxidation: experiment, theory and modelling, *Atmos. Chem. Phys.*, 18, 6095-6120, <https://doi.org/10.5194/acp-18-6095-2018>, 2018.

Newland, M. J., Nelson, B. S., Muñoz, A., Ródenas, M., Vera, T., Tárrega, J., and Rickard, A. R.: Trends in stabilisation of Criegee intermediates from alkene ozonolysis, *Phys. Chem. Chem. Phys.*, 22, 13698-13706, <https://doi.org/10.1039/D0CP00897D>, 2020.

Nguyen, T. B., Tyndall, G. S., Crounse, J. D., Teng, A. P., Bates, K. H., Schwantes, R. H., Coggon, M. M., Zhang, L., Feiner, P., Miller, D. O., Skog, K. M., Rivera-Rios, J. C., Dorris, M., Olson, K. F., Koss, A., Wild, R. J., Brown, S. S., Goldstein, A. H., de Gouw, J. A., Brune, W. H., Keutsch, F. N., Seinfeld, J. H. and Wennberg, P. O.: Atmospheric fates of Criegee intermediates in the ozonolysis of isoprene, *Phys. Chem. Chem. Phys.*, 18, 10241-10254, 2016.

Peltola, J., Seal, P., Inkilä, A., and Eskola, A.: Time-resolved, broadband UV-absorption spectrometry measurements of Criegee intermediate kinetics using a new photolytic precursor: unimolecular decomposition of CH_2OO and its reaction with formic acid, *Phys. Chem. Chem. Phys.*, 22, 11797-11808, <https://doi.org/10.1039/D0CP00302F>, 2020.

Sheps, L.: Absolute ultraviolet absorption spectrum of a Criegee intermediate CH_2OO , *J. Phys. Chem. Lett.*, 4, 4201-4205, 2013.

Sipilä, M., Jokinen, T., Berndt, T., Richters, S., Makkonen, R., Donahue, N. M., Mauldin III, R. L., Kurtén, T., Paasonen, P., Sarnela, N., Ehn, M., Junninen, H., Rissanen, M. P., Thornton, J., Stratmann, F., Herrmann, H., Worsnop, D. R., Kulmala, M., Kerminen, V.-M., and Petäjä, T.: Reactivity of stabilized Criegee intermediates (sCIs) from isoprene and monoterpene ozonolysis toward SO_2 and organic acids, *Atmos. Chem. Phys.*, 14, 12143-12153, <https://doi.org/10.5194/acp-14-12143-2014>, 2014.

Stephenson, T. A. and Lester, M. I.: Unimolecular decay dynamics of Criegee intermediates: Energy-resolved rates, thermal rates, and their atmospheric impact, *Int. Rev. Phys. Chem.*, 39 (1), 1-33, <https://doi.org/10.1080/0144235X.2020.1688530>, 2020.

Ting, W-L, Chen, Y-H., Chao, W., Smith, M.C., and Lin, J Jr -M.: The UV absorption spectrum of the simplest Criegee intermediate CH_2OO , *Phys. Chem. Chem. Phys.*, 16, 10438-10443, 2014.

Vansco, M. F., Li, H. and Lester, M. I.: Prompt release of O^1D products upon UV excitation of CH_2OO Criegee intermediates, *J. Chem. Phys.* 147, 013907, <https://doi.org/10.1063/1.4977987>, 2017.

Evaluated kinetic and photochemical data for atmospheric chemistry: Volume VII - Criegee intermediates

R. Anthony Cox¹, Markus Ammann², John N. Crowley³, Hartmut Herrmann⁴, Michael E. Jenkin⁵, V. Faye McNeill⁶, Abdelwahid Mellouki⁷, Jürgen Troe⁸, and Timothy J. Wallington⁹

¹ Centre for Atmospheric Science, Dept. of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EP, UK

² Laboratory of Radiochemistry and Environmental Chemistry, OFLB 103, Paul Scherrer Institut, 5232 Villigen, Switzerland

³ Max Planck Institute for Chemistry, Division of Atmospheric Chemistry, 55128 Mainz, Germany

⁴ Leibniz Institute for Tropospheric Research (TROPOS), Atmospheric Chemistry Dept. (ACD), 04318 Leipzig, Germany

⁵ Atmospheric Chemistry Services, Okehampton, Devon, EX20 4QB, UK

⁶ Department of Chemical Engineering, Columbia University, New York, NY 10027, USA

⁷ ICARE-CNRS, 1 C Av. de la Recherche Scientifique, 45071 Orléans CEDEX 2, France

⁸ Institute of Physical Chemistry, University of Göttingen, Tammannstr. 6, 37077 Göttingen, Germany

⁹ Ford Motor Company, Research and Advanced Engineering, Mail Drop RIC-2122, Dearborn, Michigan 48121-2053, USA

Correspondence to: M. E. Jenkin (atmos.chem@btinternet.com), J. N. Crowley (john.crowley@mpic.de) and R. A. Cox (rac26@cam.ac.uk)

Abstract. This article, the seventh in the series, presents kinetic and photochemical data sheets evaluated by the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation. It covers an extension of the gas phase and photochemical reactions related to Criegee intermediates previously published in ACP in 2006, and implemented on the IUPAC website up to 2020. The article consists of an introduction, description of laboratory measurements, discussion of rate coefficients for reactions of O₃ with alkenes producing Criegee intermediates, rate coefficients of unimolecular and bimolecular reactions and photochemical data for reactions of Criegee intermediates, and overview of the atmospheric chemistry of Criegee intermediates. Summary tables of the recommended kinetic and mechanistic parameters for the evaluated reactions are provided. Data sheets summarizing information upon which the recommendations are based are given in two files, provided as a Supplement to this article.

1 Introduction

Laboratory kinetic and mechanistic studies of the reactions of alkenes with ozone (O₃) have established that “Criegee intermediates” (CIs) produced from these reactions are potentially important oxidants in atmospheric chemistry (e.g. Calvert et al., 2000; Johnson and Marston, 2008; Taatjes et al., 2014). This followed the suggestion by Cox and Penkett (1971; 1972) that the rapid oxidation of SO₂ in the presence of reacting mixtures of O₃ and alkenes in air, was caused by production of a reactive intermediate, namely the peroxidic zwitterion, R₁R₂C=O⁺–O[–], proposed by Rudolf Criegee, based on studies of the liquid phase ozonolysis of alkenes (e.g. Criegee et al., 1954).

This has led to extensive study of the mechanisms of O_3 + alkene reactions, and of the chemistry of the CIs formed. It is well established that the reaction proceeds by initial cyclo-addition of O_3 across the $C=C$ double bond in an alkene to form an energy-rich primary ozonide (POZ), which rapidly decomposes to form either of two sets of CI and carbonyl compound, as shown in Fig. 1. The reaction is exothermic, leading to an excess of energy ($200\text{--}250\text{ kJ mol}^{-1}$) distributed between these reaction products. Some of this excess energy is deposited as internal energy in the nascent CIs which can promote unimolecular decomposition, or which can be lost by collisional energy transfer to other gas molecules, leading to formation of stabilised Criegee intermediates (sCIs), which can themselves react with other atmospheric trace species. The impact of the ozonolysis reaction on atmospheric oxidation chemistry is therefore influenced by the relative importance of prompt CI decomposition vs. formation of sCI. The recognition of the important distinction between the chemically activated CI, formed promptly in excited state, and the thermally equilibrated sCI has led to extensive experimental efforts to determine the yield of sCI (Y) formed from the ozonolysis of a variety of alkenes, as discussed further in Sect. 3. It is also well established that the decomposition of both CI and sCI leads to the formation of hydroxyl (HO) radicals and other radical products. Due to its important role in initiating the oxidation of organics (including alkenes), the formation of HO radicals has received particular attention, and this is discussed further in Sect. 4.

The mechanism in Fig. 1 shows that the CIs (and sCIs) formed from the ozonolysis of a simple alkene can each be formed as either of two stereo-isomers, with different orientations of the outer O atom relative to the substituent groups. This potentially has an important impact on the chemical pathways available, and their relative rates. The stereo-isomers have generally been distinguished using the terms *syn*- and *anti*-, to specify the orientation of the outer O atom relative to a particular substituent; although the use of the IUPAC *Z*- and *E*- nomenclature is becoming increasingly adopted (e.g. Vereecken et al., 2017). As shown in Fig. 1, the ozonolysis of a simple alkene containing four different substituents therefore produces four distinct CIs, with this number being systematically reduced in symmetrical alkenes (because the products of the two POZ fragmentation pathways are the same); or in alkenes possessing two identical substituents on the same carbon atom (because the stereo-isomerism in the CI is removed). However, the number of different CI isomers can also be increased if the alkene contains alkenyl substituents, as is the case for the CIs formed from the ozonolysis of dienes such as the C_4 species derived from isoprene, because of additional stereo-isomerism in the substituent group(s).

Based on current understanding of the mechanism of alkene ozonolysis (as illustrated in Fig. 1), the steady state concentration of a given stabilised Criegee intermediate, $[sCI_i]$, maintained by a balance between production and loss, is described by Eq. (1):

$$[sCI_i] = \frac{\sum_j (Y_{ij} \times k_{1j} \times [\text{alkene}_j] \times [O_3])}{(k_{di} + J_i + k_{2i}[H_2O] + k_{3i}[(H_2O)_2] + k_{4i}[SO_2] + k_{5i}[NO_2] + \dots + \dots)} \quad (1)$$

Here, k_{1j} is the rate coefficient for the reaction of O_3 with alkene_j , and Y_{ij} is the yield of sCI_i from that reaction. The numerator of Eq. (1) therefore quantifies the source term for formation of sCI_i from all relevant alkenes, and the denominator

quantifies the sum of the rates of the unimolecular and individual bimolecular loss processes for sCI_i, with the example contributors to the summation being based on the processes shown in Fig. 1. This illustrates that knowledge of the yields, rate coefficients and products for the component reactions is important for quantitative description of the chemical pathways controlling the atmospheric chemistry and impact of the given sCI_i. It also shows the importance of establishing how these parameters vary from one sCI to another, e.g. the structural-dependence of the rate coefficients for their unimolecular and bimolecular reactions.

Although rate constants have been determined accurately for a large number of O₃ + alkene reactions, using both direct and relative rate techniques, all kinetic data reported for sCI reactions prior to 2012 were based on indirect relative rate techniques. Many of these data were previously evaluated by the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (Atkinson et al., 2006). Since 2012, many new rate coefficients for sCI reactions have been reported using direct kinetic studies, providing a wealth of data for the elementary reaction kinetics and spectroscopy of sCI reactions. The current evaluation therefore addresses these reactions, extending substantially the scope of our former evaluation published in ACP in 2006. This includes a major extension of the scope of the evaluation to include rate coefficients of elementary reactions of selected sCIs, which have provided a better understanding of the atmospheric impact of sCI chemistry.

In this review we summarize the results of this evaluation activity, presenting in turn the recommended kinetic data for the key reactions in the above mechanism, using data for those species which are representative of the chemistry of the terrestrial atmosphere. The rate coefficients for O₃ + alkene initiation reactions are presented and discussed in Sect. 2, with reference to a series of detailed data sheets which are provided in Supplement A. Information on the sCI and HO radical yields from the ozone + alkene reactions is presented in Sects. 3 and 4, with additional discussion once again provided in the corresponding data sheets in Supplement A. The data sheets therefore each include summary information on the reported kinetics studies of the given reaction, and provide an overview of reported mechanistic information and product yields where available.

The spectroscopy and kinetics recommendations for the sCI reactions are presented and discussed in Sects. 5 and 6. These include data for bimolecular and unimolecular reactions of selected sCIs of particular atmospheric relevance for which direct kinetic data have been reported, namely CH₂OO, Z- and E- CH₃CHOO₂ ~~and~~ (CH₃)₂COO and E-(CH=CH₂)(CH₃)COO; and we also provide some discussion of the complete set of C₄ intermediates formed from isoprene (see Fig. 2 for sCI structures). These are predicted to be among the most important sCIs in tropospheric chemistry (Vereecken et al., 2017), and can also act as a systematic set of template species for representing the fates of some larger and more complex sCIs. Detailed data sheets for the sCI reactions are provided in Supplement B, providing supporting summary information and discussion. Finally, the recommended kinetics parameters are used to evaluate the relative importance of the different fates of the sCIs under representative atmospheric conditions in Sect. 7, and an overview of the impact of Criegee intermediates in atmospheric oxidation chemistry is given in Sect. 8.

2 Rate coefficients of O₃ + alkene reactions

The present evaluation considers the reactions of O₃ with 31 alkenes, including small (C₁ to C₄) alkenes, isoprene, monoterpenes and sesquiterpenes. This represents a substantial increase on those considered in our previous evaluation (Atkinson et al., 2006), which were limited to ethene, propene, isoprene and α -pinene. The reactions are listed in Table 1, along with the associated recommended rate coefficients. A detailed data sheet for each reaction is also provided in Supplement A. As discussed in detail previously (e.g. Calvert et al., 2000; 2015), the data indicate that the rate coefficients are highly sensitive to alkene structure, and depend on the degree of alkyl substitution of the unsaturated bond(s), on steric effects and on ring-strain effects in cyclic compounds. The lower tropospheric lifetimes of the alkenes, with respect to reaction with 20 ppb O₃, therefore cover several orders of magnitude, ranging from as short as 2-3 minutes for reactive species such as α -terpinene, β -caryophyllene and α -humulene, to about seven weeks or longer for camphene and longifolene. For the simple alkenes, the lifetimes range from about 30 minutes for the fully-substituted 2,3-dimethylbut-2-ene to about two weeks for ethene. With the exception of the least reactive compounds, removal by ozonolysis is expected to make a contribution for all the evaluated alkenes under lower tropospheric conditions, and is generally the dominant fate for those with rate coefficients in excess of about 10^{-15} cm³ molecule⁻¹ s⁻¹.

3 sCI yields from O₃ + alkene reactions

As described in Sect. 1 and Fig. 1, the chemically-activated CIs formed from the O₃ + alkene reactions may either decompose promptly, or lose energy by collisions with other molecules to form stabilised Criegee intermediates (sCIs). The sCIs have the potential to undergo reactions with other atmospheric trace gases leading to their oxidation and formation of characteristic products. Thus, it is important to quantify the yield (*Y*) of each sCI from each relevant precursor alkene if the impact of alkene ozonolysis and Criegee chemistry on oxidation processes is to be correctly represented in atmospheric mechanisms, or to allow the local steady-state concentration of the sCIs to be estimated by Eq. (1).

There have been extensive experimental efforts to determine *Y* for a variety of O₃ + alkene systems. The yields have generally been expressed as a fraction of the molar amount of O₃ reacted, and have been determined by reaction of the sCIs formed with an appropriate added scavenger reagent (e.g. SO₂, H₂O, HCHO, HCOOH, CF₃C(O)CF₃). The yield is determined either from quantitative analysis of a characteristic product of the sCI + scavenger reaction, or through measurement of the loss of the scavenger. The reported values of *Y* are therefore indirect measurements which have generally quantified the total yield of sCIs formed in a given alkene + O₃ system, with little or no information on the contributions of the component sCI species for asymmetric alkenes being reported. Table 2 gives a summary of the recommended values of *Y* for the evaluated O₃ + alkene reactions at 298 K and 1 bar, with additional details provided in the corresponding reaction data sheets in Supplement A. Values for selected other alkenes (*trans*-dec-5-ene and cyclohexene) are also given to help illustrate structural variations in *Y*, as discussed further below.

As indicated in Sect. 1, the stabilisation of the promptly formed chemically-activated CIs requires loss of internal energy through collisional energy transfer to other gas molecules. As a result, studies of a number of O_3 + alkene systems have shown that Y depends on pressure (e.g. Hatakayama et al., 1986; Drodz and Donahue, 2011; Hakala and Donahue, 2016; 2018; Campos-Pineda and Zhang, 2017), but with significant residual values at zero pressure. This indicates that a limiting
5 yield of sCIs (i.e. with internal energy below the threshold required for decomposition or isomerization) is typically formed directly from decomposition of the primary ozonide, POZ (although this is not illustrated in Fig. 1 for simplicity), with additional sCI formation resulting from collisional stabilisation of the chemically-activated CIs.

The value of Y is also expected to vary systematically with alkene structure, and the reported data display some logical structural trends that are consistent with theoretical treatments. These can be rationalized in terms of the nascent internal
10 energy in the Criegee intermediate, and how this is influenced by partitioning of the excess energy into either translational modes (i.e. recoil), internal energy of the carbonyl co-product, or into non-reactive vibrational or rotational modes within the Criegee intermediate (e.g. Choung et al., 2004; Drodz et al., 2011). The value of Y is therefore expected to increase with alkene size along a homologous series, as has been confirmed for a set of *trans*- symmetric alkenes between C_4 and C_{14} by Hakala and Donahue (2018). As a result, the values at 298 K and 1 bar for large acyclic alkenes (e.g. $Y = 1.0$ for *trans*-dec-5-
15 ene and *trans*-tetradec-7-ene) tend to be larger than those for the smaller acyclic alkenes, as listed in Table 2. Similarly, those for large cycloalkenes with endocyclic double bonds (e.g. $Y > 0.6$ for the C_{15} β -caryophyllene) tend to be larger than those for smaller species such as the C_{10} α -pinene ($Y = 0.18$) and the C_6 cyclohexene ($Y < 0.05$). It is also clear that the values for cycloalkenes with endocyclic double bonds are systematically lower than those for similarly sized acyclic alkenes (or cycloalkenes with exocyclic double bonds). This can be rationalised in terms of the excess energy being confined within a
20 single product possessing both Criegee and carbonyl functionalities for the cycloalkenes with endocyclic double bonds; whereas it can be dissipated into translational modes, and the internal energy of the carbonyl co-product, for acyclic alkenes (and for cycloalkenes with exocyclic double bonds).


It is also recognized that the indirect yield measurements are often subject to significant uncertainties, and variability between different measurement methods (e.g. see Hakala and Donahue, 2016), such that systematic trends can be masked.
25 Another important consideration is that exceptionally rapid unimolecular decomposition and isomerization reactions are predicted to be available for some sCIs, e.g. some of the C_4 species formed from O_3 + isoprene (Vereecken et al., 2017), as is discussed further in Sects. 4 and 7. In these cases it is possible that the bimolecular reactions with scavengers are unable to compete with the fast unimolecular processes, and consequently the reported yields for sCI formation may be underestimated.

30 4 HO radical yields from O_3 + alkene reactions

It has been established for several decades that the reactions of O_3 with alkenes lead to the formation of HO radicals (e.g. Finlayson et al., 1972; Donahue et al., 1998), and considerable attention has been given to quantifying HO radical yields for

many O₃ + alkene systems. In most of the reported studies, the yields have been determined indirectly by reaction of the HO formed with an appropriate added scavenger reagent (e.g. cyclohexane, 1,3,5-trimethylbenzene, butan-2-ol). The yield is then determined either from quantitative analysis of the yield of a characteristic product of the HO + scavenger reaction, or through measurement of the loss of the scavenger. However, direct detection methods (particularly LIF) have also been used to quantify HO yields in a number of studies (e.g. Donahue et al., 1998; Siese et al., 2000; Kroll et al., 2001a; 2001b; Malkin et al., 2010; Alam et al., 2013), providing unequivocal identification of HO as a product of O₃ + alkene reactions, and a means of confirming the validity and interpretation of the indirect methods (e.g. Malkin et al., 2010). Table 3 gives a summary of the recommended HO radical yields for the evaluated O₃ + alkene reactions at 298 K and 1 bar, with additional details provided in the corresponding reaction data sheets in Supplement A.

The formation of HO radicals can result from both prompt unimolecular decomposition of chemically-activated CIs, and from decomposition of thermally-equilibrated sCIs over longer timescales (e.g. as demonstrated in the time-resolved pressure dependence measurements of Kroll et al., 2001c). The most important mechanism forming HO is generally accepted to proceed by a 1,4 H shift isomerization to an excited vinyl hydroperoxide intermediate, which decomposes to form HO and a vinoxy or β-oxo alkyl radical; and this has been characterised for a variety of Criegee intermediates in theoretical studies (e.g. Vereecken et al., 2017). Using Z-CH₃CHOO as an example, the mechanism proceeds as shown below:



First proposed for (CH₃)₂COO by Niki et al. (1987), this mechanism is therefore potentially available for all di-substituted and Z- mono-substituted Criegee intermediates that possess a β-hydrogen atom, but is unavailable for CH₂OO and E- mono-substituted Criegee intermediates (e.g. E-CH₃CHOO), where the outer O atom of the CI moiety is not directed towards an organic group. In the cases where reaction (R6) is available, it is believed to be the dominant decomposition route for small (e.g. methyl- and ethyl-substituted) Criegee intermediates. The reported variation of HO yields with structure for simple small alkenes (e.g. as shown in Table 3) can therefore be broadly rationalized in terms of the combination of Criegee intermediates formed, and whether or not reaction (R6) is available; and this has provided the basis of simple structure-activity relationships (SARs) for HO yields from O₃ + alkene reactions (e.g. Rickard et al., 1999). Thus, those for fully methyl-substituted alkenes (e.g. 0.93 for 2,3-dimethyl-but-2-ene) tend to be systematically higher than those for partially methyl-substituted alkenes (e.g. 0.33 for *cis*-but-2-ene and 0.60 for *trans*-but-2-ene), which in turn are higher than that for the unsubstituted ethene (0.17).

Another key route involves initial rearrangement (1,3 ring-closure) to form a dioxirane intermediate (see Fig. 1). This provides an alternative unimolecular decomposition pathway for CH₂OO and E- mono-substituted Criegee intermediates

(e.g. *E*-CH₃CHOO), and is also calculated to be competitive for some *Z*- mono-substituted Criegee intermediates possessing oxygenated substituents (Vereecken et al., 2017). In the former case, it is likely to be significant only for the chemically-activated CI, [CH₂OO]*, because unimolecular loss of the thermally-equilibrated sCI, CH₂OO, is observed and calculated to be slow (see Sect. 7). The dioxirane intermediate isomerizes to form “hot” formic acid, [HC(O)OH]*, which can fragment

5 via a number of pathways:




10 Reaction (R7a) can therefore account for the small HO yield (0.17) resulting from the O₃ + ethene reaction (Table 3), and the set of product channels can also rationalize the observed formation of HO₂ (from the reactions of O₂ with H and HCO), CO and CO₂ (see data sheet OX_VOC5 in Supplement A). In the cases of *E*- mono-substituted Criegee intermediates (e.g. *E*-CH₃CHOO), isomerization via a dioxirane intermediate is again expected to be significant for chemically-activated CIs, and is also calculated to occur for thermally-equilibrated sCIs (e.g. Vereecken et al., 2017). For sCIs, however, it is in
15 competition with some particularly rapid bimolecular reactions under atmospheric conditions (see Sects. 7 and 8), and may therefore be of limited importance. In the case of the chemically activated [*E*-CH₃CHOO]*, the dioxirane intermediate isomerizes to form “hot” acetic acid, [CH₃C(O)OH]*, and the following pathways can rationalize the observed formation of HO₂, CO₂, CH₂=CO (ketene), CH₄ and CH₃OH from the reactions of O₃ with propene and *cis*- and *trans*-but-2-ene (see data sheets OX_VOC6 , OX_VOC17 and OX_VOC18 in Supplement A1).




In principal, formation of HO (and CH₃CO, or CH₃ and CO) may also occur, by a route analogous to reaction (R7a). The
25 possible contribution of the corresponding channel more generally for *E*- mono-substituted CIs is poorly characterised, although it is generally accepted to be only a minor source of HO compared with the 1,4 H shift isomerization route for the *Z*- conformers. The dioxirane route has also been reported to lead to the formation of stabilised acid, ester or lactone products, particularly in larger O₃ + alkene systems (e.g. Hakola et al., 1994; Griesbaum et al., 1998; Winterhalter et al., 2009; Nguyen et al., 2009a,b).

30 A number of other unimolecular reactions are available for Criegee intermediates possessing larger organic substituents, particularly those that are unsaturated (e.g. see Vereecken et al., 2017). For example, very rapid 1,5 ring-closure reactions are expected to dominate for *Z*-α,β-unsaturated sCIs, such as *Z*-(CH=CH₂)(CH₃)COO and *Z*-(C(CH₃)=CH₂)CHOO formed from O₃ + isoprene, e.g. for *Z*-(CH=CH₂)(CH₃)COO:



In the cases of carbonyl-substituted sCIs, such as those formed from cycloalkenes with endocyclic double bonds, the potential for (bicyclic) ring-closure to form intramolecular secondary ozonides is well established, e.g.:



15 These reactions have been characterised in a number of theoretical studies (e.g. Chuong et al., 2004; Nguyen et al., 2009b; Mackenzie-Rae et al., 2016; Vereecken et al., 2017; Long et al., 2019), with experimental evidence for their formation also reported (e.g. Winterhalter et al., 2009; ~~Vibenholt et al., 2009~~; Beck et al., 2011). In practice, however, these reactions are only observed (and calculated) to be significant for larger systems (e.g. sesquiterpene ozonolysis), where the Criegee intermediates are formed significantly stabilised, and the ring-closure reaction does not result in prohibitive ring-strain. Where these criteria are met, they are predicted to be rapid reactions that can compete with, or dominate over, other decomposition routes; and this is one factor contributing to the low HO yields reported for some sesquiterpenes (e.g. β -caryophyllene and α -humulene).

20

5 Structure and spectroscopy of Criegee intermediates

Assessment of the photolysis rates and product channels for sCIs requires quantitative data for the absorption cross sections and quantum yields for individual sCI species at atmospheric, actinic wavelengths, mainly in the UV and visible. The development of methods for creating specific sCIs in defined concentrations has allowed experimental investigation of their spectroscopy and structure, e.g. see the review by Osborn and Taatjes (2015). Moreover, advances in theoretical methods have also provided insight into the spectra and structure of sCIs, and quantum calculations have given further details of reaction mechanisms and product channels of sCI photolysis (e.g. Samanta et al., 2014).

25

The spectroscopic studies of sCIs have led to the recognition that they have a singlet ground electronic state whose dominant configuration is that of a zwitterion, and this is reflected in the large dipole moment of these species (Chhantyal-Pun et al., 2017a). The observed spectra of the $C_1 - C_3$ sCIs exhibit strong and broad absorptions centred in the near UV, with maximum cross sections of the order of $10^{-17} \text{ cm}^2 \text{ molecule}^{-1}$. These features, and their detailed rovibrational structures, are consistent with $B(1A') \leftarrow X(1A')$ transitions, i.e. intense $\pi^* \leftarrow \pi$ transitions analogous to the familiar UV spectrum of O_3 in

30

the Hartley/Huggins bands. Photodissociation of ~~sCIs-CH₂OO~~ from this excitation is reported to proceed with a quantum yield of unity (e.g. Ting et al., 2014), producing HCHO and either O(³P) or O(¹D). Production of O(¹D) has been reported to be dominant (e.g. Li et al., 2015), and exclusive in the long wavelength tail ($\lambda \geq 364$ nm; Vansco et al., 2017), yields O(¹D) and a carbonyl compound with a quantum yield of unity.

5 5.1 Conformers

As described in Sect. 1 and Fig. 1, Criegee intermediates with dissimilar substituents can exist as two possible conformers, denoted *E*- and *Z*-, which differ in the orientation of the outer O atom relative to the substituent groups. The simplest examples are *E*- and *Z*-CH₃CHOO (acetaldehyde oxide, see Fig. 2), for which conformer dependence has been demonstrated experimentally and theoretically in its spectra and its reaction rates and pathways. Calculations place *Z*-CH₃CHOO about 15
10 kJ mol⁻¹ lower in energy than *E*-CH₃CHOO (Kuwata et al., 2010), reflecting the zwitterionic character of the Criegee intermediate structure. Calculated energies of the *E*- and *Z*- conformers of CH₃CHOO are consistent with the spectral shift of $\lambda_{\text{max}}(\textit{Z-}) < \lambda_{\text{max}}(\textit{E-})$ shown in Fig. 3. This has enabled conformer-specific reactions to be investigated, using direct observation of the kinetics and products of the two conformers. The barrier to interconversion is substantial (~160 kJ mol⁻¹), and consequently *E*- and *Z*-CH₃CHOO act as distinct chemical species at atmospheric temperatures. The absence of rotation
15 is an important indicator of the zwitterionic character of the intermediate, as originally proposed by Criegee et al. (1954).

5.2 UV spectra of stabilised Criegee intermediates

In experimental studies of UV-visible spectra, the series of C₁ – C₃ sCIs have mainly been formed by photolysis of the corresponding di-iodoalkane (via C-I bond fission), followed by the reaction of the iodoalkyl radical with O₂, e.g. in the case of CH₂OO (e.g. Beames et al., 2012; Sheps, 2013):



The formation of CH₂I (and subsequently CH₂OO) from the photolysis of CH₂IBr (via C-Br bond fission) has also recently been reported (Peltola et al., 2020), suggesting that bromo-iodoalkanes may also be used more widely as sCI precursors.

Absolute cross sections at specific wavelengths have been derived by monitoring the laser UV-induced depletion of the sCIs, monitored, for example, by mass spectrometry, or by time resolved UV-absorption spectroscopy. The experimental data reveal some discrepancies regarding the shapes, structure, and intensities of the B(¹A') ← X(¹A') spectra determined using transient absorption spectroscopy, compared with laser-induced depletion techniques, determined under molecular beam conditions. This discrepancy has been attributed to the much lower temperatures reached in the molecular beams, compared to measurements at ambient temperature, but lack of detailed data on the temperature dependence of the cross sections over
25 the required range precludes firm conclusions to be drawn. The evaluation of the spectral data, and detailed discussion of the reported studies, is given for the C₁ – C₃ sCIs in the data sheets in Supplement Sect. B5. The recommendations for the

maximum absorption cross sections are given in Table 4, and the spectra are presented in Fig. 3. Using these data, representative lower tropospheric photolysis removal rates in the range $0.4\text{--}1.4\text{ s}^{-1}$ can be calculated for a solar zenith angle of 30° at the surface, based on the actinic flux estimates of Madronich, presented by Finlayson-Pitts and Pitts (2000). This indicates that loss by photolysis is likely only a minor or negligible loss process for sCIs in the lower atmosphere, compared with their collective removal by the thermal reactions discussed in the following section.

Novel methods for the production of the C_4 isoprene-derived sCIs have also been reported (Barber et al., 2018; Vansco et al., 2019), with *Z*- and *E*-($\text{CH}=\text{CH}_2$)(CH_3)COO formed from the photolysis of 1,3-di-iodobut-2-ene and *Z*- and *E*-($\text{C}(\text{CH}_3)=\text{CH}_2$)CHOO from the photolysis of 1,3-di-iodo-2-methylprop-1-ene, both in the presence of O_2 . This has allowed characterization of the UV-visible spectra of the unsaturated C_4 sCIs (Vansco et al., 2018; 2019), which are reported to be broader and shifted to longer wavelengths compared with those of the simple $C_1 - C_3$ sCIs as a result of the conjugation of the vinyl and carbonyl oxide groups (see detailed discussion in the data sheets in Supplement Sect. B5).

6 Rate coefficients for thermal reactions of sCIs

6.1 Measurements of absolute rate coefficients for reactions of sCIs

In recent years, numerous direct studies of the elementary reaction kinetics of sCIs have been reported. This has been made possible by two developments. First, the discovery of a novel fast photochemical source of sCIs, from the reactions of iodoalkyl radicals with O_2 , has allowed generation of specific sCIs, following the photolysis of the corresponding di-iodoalkane (and recently, bromo-iodoalkene); as shown above in Sect. 5 for the example of CH_2OO (reactions (R11) and (R12)). Second, direct time-resolved detection and measurements of sCI concentrations have been achieved using spectroscopic methods involving both tunable vacuum UV multiplexed photoionization mass spectrometry (MPIMS), and UV or IR absorption.

The first breakthrough in these developments for direct studies of sCI kinetics came from the work of Taatjes and co-workers, who used MPIMS to monitor the time-resolved decay of CH_2OO in the presence of bimolecular reaction partners such as SO_2 , NO and NO_2 (Welz et al., 2012; Taatjes et al., 2012). Subsequently it was shown that this technique for detection and production could be equally well applied to kinetics studies of the larger Criegee intermediates (e.g. Taatjes et al., 2013; Chhantyal-Pun et al., 2017b), so that structural effects on the basic oxidation rates and mechanisms could be explored directly.

The tunable light sources needed for MPIMS are not readily available for conventional laboratory rate constant measurements. The ~~discovery-characterization~~ of the strong absorption spectrum of CH_2OO in the mid UV (Sheps, 2013) offered a second, more flexible and sensitive (but less specific) detection method for following sCI kinetics, which has the advantage of the ability to monitor sCI kinetics at up to 1 bar pressure, appropriate for lower atmospheric conditions. The UV absorption method is also applicable to the larger Criegee intermediates, produced from the same source chemistry (e.g.

Sheps et al., 2014; Huang et al., 2015), and can provide kinetic/spectroscopic distinction of the *Z*- and *E*- conformers, where applicable.

The lower atmosphere contains many alkenes from both natural and man-made sources, which react with O₃ to form Criegee intermediates with a wide variety of structures. Earlier work on ozone-alkene reactions gave little clue on the structural dependence of sCI reactivity with trace gases such as SO₂, H₂O, NO₂ and organics, or of their unimolecular decomposition rates. Direct kinetic studies have provided new information on the reaction rate constants and mechanisms of C₁ – C₃ sCIs formed from ozonolysis of simple alkenes. As indicated in Sect. 5.2, methods for their production and spectroscopic characterization of more complex isoprene-derived species are emerging (e.g. Barber et al., 2018; Vansco et al., 2018; 2019), and these have providing-provided a-the basis for their direct kinetics study (Caravan et al., 2020). However-, although-direct experimental determinations of rate coefficients have not yet been reported for other, morelarger complex species (\geq C₄), derived for example from monoterpenes and sesquiterpenes.

6.2 Evaluation of rate coefficients for bimolecular reactions

As noted in Sect. 1, most of the information on the kinetics of sCI reactions up to 2006 was based on data obtained using indirect relative rate techniques. These were evaluated by the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation and published in ACP in 2006 (Atkinson et al., 2006). Since 2012, the direct kinetics studies described above have provided a wealth of new data on the elementary reaction kinetics and spectroscopy of sCIs. This has stimulated further competitive rate studies using static and slow-flow experiments in chambers to generate sCIs from O₃ + alkene reactions under atmospheric pressure and temperature conditions. In this evaluation, recommended rate coefficients are generally based on the results of direct kinetic studies of the sCIs, derived from di-iodoalkane precursors as described above (Sects. 5.2 and 6.1). However, the results of relative rate studies are also used to assess information on the kinetics, and to check for consistency of the kinetic data for C₁ – C₃ sCIs, when produced by alkene ozonolysis. Table 5 provides a summary of the preferred values of bimolecular reaction rate coefficients, with additional details given in the corresponding reaction data sheets in Supplement B. As indicated in Sect. 1, the evaluation focuses on classes of reaction that are of particular significance in tropospheric chemistry. Where data are available, these include reactions with SO₂, NO₂, H₂O, (H₂O)₂, CH₃CHO, CH₃C(O)CH₃, CF₃C(O)CF₃, HC(O)OH, CH₃C(O)OH and CF₃C(O)OH for the set of sCIs, with some additional reactions also considered for CH₂OO. It is noted that data have been reported for some other classes of reaction (e.g. with alkenes and alcohols) that are uncompetitive under tropospheric conditions, and some relative rate data are also available for large sCIs not considered here, e.g. as summarised in the compilation reported by Khan et al. (2018). With exception of the reactions of *E*-(CH=CH₂)(CH₃)COO with SO₂ and HC(O)OH (Caravan et al., 2020), tThere are currently no direct kinetics determinations for reactions of the C₄ sCIs derived from isoprene. In these cases, the recommendations are either inferred from those for the simpler C₂ and C₃ species and *E*-(CH=CH₂)(CH₃)COO, or adopted from reported theoretical studies (Vereecken et al., 2017; Chhantyal-Pun et al., 2017a). The performance of the ensemble of

bimolecular reactions with SO₂, H₂O and (H₂O)₂ (and unimolecular decomposition reactions) has been checked for consistency, using the results of published chamber and slow-flow experiments (Sipilä et al., 2014; Newland et al., 2015; Nguyen et al., 2016), as described in detail in data sheet CGI_21 (Supplement B). The results provide some confidence in these recommendations for use in practical applications, although the data also support some tolerance in the parameter values applied. Measurements of speciated sCI yields, and further direct kinetics studies of the rate coefficients and product channels for the reactions of the C₄ sCI isomers, are therefore required to allow these recommendations to be confirmed or refined.

6.3 Evaluation of rate coefficients for unimolecular decomposition

Table 6 shows a summary of preferred values of unimolecular decomposition rate coefficients, k_d , given in the corresponding data sheets in Supplement B. The evaluations are based on both consideration of direct time-resolved measurements, and those reported in relative rate experiments, where k_d is determined relative to the loss rate of sCI via a well-defined competing bimolecular reaction. In the former case, k_d is determined from observation of the decay kinetics of the Criegee intermediates themselves (or of a marker species) in the absence of a second reagent, or by extrapolation of the observed first order removal rate vs. reagent concentration plots to zero. Although unimolecular decomposition can make a major, or the dominant, contribution to such limiting first-order removal rates, other processes also need to be taken into account (e.g. wall loss, reaction with impurity gases or diffusive loss from the monitoring probe area), as is usually discussed in the individual studies. In some cases, therefore, direct kinetics studies can only provide upper limit estimates of unimolecular decomposition rates for sCIs, particularly when the decomposition rate is slow. Relative rate determinations can also be influenced by background loss processes for the sCI (e.g. reaction with impurity gases or products), and reported rate coefficient ratios need to be placed on an absolute basis using the rate coefficient for the competing, reference bimolecular reaction, which itself has an associated uncertainty. In the present evaluations, the reference (bimolecular) rate coefficients are all based on the preferred values given in Table 5.

As indicated above in Sect. 6.2, the recommendations for the unimolecular decomposition of C₄ sCIs derived from isoprene ozonolysis are adopted from the theoretical study of Vereecken et al. (2017). These recommendations have been assessed, along with the recommended bimolecular rate coefficients for reactions with SO₂, H₂O and (H₂O)₂, using the results of published chamber and slow-flow experiments (Sipilä et al., 2014; Newland et al., 2015; Nguyen et al., 2016), as described in detail in data sheet CGI_21 (Supplement B).

7 Overall reactivity conclusions – comparison of experiment and theory

In addition to the progressive increase in the availability of experimental data, there have been substantial advances in the theoretical treatment of structure and reaction kinetics of sCIs in the gas phase (e.g. Olzmann et al., 1997; Zhang et al., 2002;

Anglada et al., 2002; 2011; Ryzhkov and Ariya, 2004; Kuwata et al., 2010; Vereecken et al., 2012; 2017; Liu et al., 2014; Anglada and Solé, 2016; [Long et al., 2016](#); Vereecken and Nguyen, 2017; Chhantyal-Pun et al., 2017a; [Stephenson and Lester, 2020](#)). Theoretical studies provided particular guidance prior to the advances in sCI production and detection methods for direct kinetics measurements, as described above. However, the body of experimental information now available for a series of sCIs allows the results of theoretical studies to be validated, and for the methods to be refined, optimised and extended. [This has included comparison of experimental and theoretical unimolecular decay rates of a number of infra-red activated sCIs for a range of excitation energies \(e.g. Fang et al., 2016; 2017; Barber et al., 2018\).](#)

In this section, the recommended rate coefficients for unimolecular decomposition and bimolecular reactions of the C₁ – C₃ sCIs with H₂O and (H₂O)₂ at 298 K and atmospheric pressure are compared with those derived from theoretical calculations (see Table 7). [With the exception of the unimolecular decomposition of CH₂OO, the](#) calculated values are based on ~~those~~ [the high-pressure limiting values](#) reported in the comprehensive study of Vereecken et al. (2017), which presented theory-based structure-activity relationships (SARs) for 98 atmospherically relevant classes of sCI. Those for unimolecular decomposition reactions were derived from consideration of 14 reaction types (e.g. including the 1,4 H shift isomerization, 1,3 ring-closure and 1,5 ring-closure reactions discussed and illustrated in Sect. 4) for a benchmark series of sCIs containing key substituents, comprising a set of about 170 calculated rate coefficients. Those for the bimolecular reactions with H₂O and (H₂O)₂ were based on fitting theory-derived reactivity trends to a set of literature data, which included rate coefficients available at the time for the same set of C₁ – C₃ sCIs considered in the present evaluation. As a result, the absolute scaling of the theory-based SAR values cannot be considered to be entirely independent of our experimentally-based evaluations for the H₂O and (H₂O)₂ reactions, although comparison of the (considerable) reactivity variation across the series of sCIs is valid.

The comparisons shown in Table 7 demonstrate that the theory-based SAR rate coefficients reported by Vereecken et al. (2017) show a good level of consistency with our recommended rate coefficients for the reactions of the C₁ – C₃ sCIs. Both sets of parameters display a similar structural dependence across the series of sCIs. The rate coefficients for the bimolecular reactions with H₂O and (H₂O)₂ agree to within about a factor of about four (which is well within the combined uncertainties), where direct comparison is possible (i.e. for CH₂OO and *E*-CH₃CHOO); and where only an upper limit recommendation is possible in the present work (i.e. for *Z*-CH₃CHOO and (CH₃)₂COO), the SAR rate coefficient is fully compatible with that recommendation. This indicates a consistent structure-reactivity variation across the series, with systematically higher reactivities for the reactions of CH₂OO and *E*-CH₃CHOO with both H₂O and (H₂O)₂.

The unimolecular decomposition parameters recommended in the present study for *E*- and *Z*-CH₃CHOO and (CH₃)₂COO are in very good agreement with the theory-based SAR values; being consistent with dominant 1,3 ring-closure to form a dioxirane intermediate for *E*-CH₃CHOO, and dominant 1,4 H shift isomerization to form vinyl hydroperoxide intermediates for *Z*-CH₃CHOO and (CH₃)₂COO. In the case of CH₂OO, ~~however, the approximate upper limit rate coefficient at 298 K and atmospheric pressure recommended in the present work, based on extrapolation of higher temperature experimental data (see data sheet CGI_12), is two orders of magnitude lower than the theory-based SAR value~~ [consistent with the value calculated](#)

by Long et al. (2016), which is about a factor of four lower than the calculated high-pressure limiting value (see data sheet CGI_12), although both the recommended and calculated values are sufficiently slow for unimolecular decomposition to be uncompetitive under tropospheric conditions.

Also shown in Table 7 are representative lower tropospheric first-order loss rates (k_1) (at 298 K and atmospheric pressure) calculated for the same series of reactions using both sets of rate parameters; and for the bimolecular reactions with SO₂, based on the rate coefficients recommended here. The representative rates for the bimolecular reactions assume a mid-range relative humidity of 40 % (corresponding to [H₂O] $\approx 3 \times 10^{17}$ molecule cm⁻³ and [(H₂O)₂] $\approx 2 \times 10^{14}$ molecule cm⁻³), and an SO₂ concentration of 2.5×10^{10} molecule cm⁻³ (about 1 ppbv), broadly typical of urban background air. Although these concentrations are only representative, the associated values of k_1 nevertheless give a clear indication that unimolecular decomposition is the dominant loss route for *Z*-CH₃CHOO and (CH₃)₂COO, consistent of a widespread role for 1,4 H shift isomerization for *Z*- monoalkyl-substituted and dialkyl-substituted sCIs possessing a β -hydrogen atom. In contrast, bimolecular reactions with H₂O and/or (H₂O)₂ are the dominant loss routes for CH₂OO and *E*-CH₃CHOO, and likely most other *E*- monoalkyl-substituted sCIs. The values of k_1 also give an indication that loss due to reaction with SO₂ generally only makes a minor contribution to sCI removal, away from the immediate vicinity of SO₂ sources, such that their ambient concentration is mainly controlled by either decomposition or reaction with H₂O and/or (H₂O)₂.

The match between the experimentally-based recommendations presented here, and those derived from the theory-based SARs for this set of sCIs, gives some confidence that the SAR rate coefficients of Vereecken et al. (2017) provide a very reasonable basis for representing the structural dependence of the kinetic parameters for unimolecular decomposition and bimolecular reactions with H₂O and (H₂O)₂. In view of this, our corresponding recommendations for the more complex C₄ isoprene-derived species (*E*- and *Z*-(CH=CH₂)(CH₃)COO and *E*- and *Z*-(C(CH₃)=CH₂)CHOO) are currently adopted from Vereecken et al. (2017), as evaluated and discussed in detail in data sheet CGI_21 (Supplement B). The 298 K rate coefficients and representative first order loss rates (k_1) are also shown for these C₄ sCIs in Table 7. The values of k_1 clearly demonstrate that bimolecular reactions cannot compete with the very rapid unimolecular decomposition of *Z*-(CH=CH₂)(CH₃)COO and *Z*-(C(CH₃)=CH₂)CHOO (via 1,5 ring-closure) under atmospheric conditions (and indeed most reported experimental conditions); and that decomposition of *Z*-(CH=CH₂)(CH₃)COO (via 1,4 H shift isomerization) is also likely to be its major loss route, by virtue of its slow bimolecular reactions with H₂O and (H₂O)₂. In the case of *E*-(C(CH₃)=CH₂)CHOO, however, loss via unimolecular decomposition and bimolecular reactions with H₂O and (H₂O)₂ are predicted to occur at comparable rates, and experimental confirmation of the rate coefficients would be of particular value for this species.

8 Impact of Criegee intermediates in atmospheric oxidation chemistry

The kinetics and mechanistic information for sCI reactions recommended in the present evaluation provides the basis for representing the associated impact of alkene ozonolysis in atmospheric chemical mechanisms. The significance of sCIs as

atmospheric oxidants can be discussed in terms of Eq. (1), which defines the local balance between production and loss of sCIs, and hence their steady-state concentrations. As described in more detail in Supplement C, the parameters recommended in this evaluation (supplemented by data from other sources) have been used to calculate surface production rates, loss rates and steady-state concentrations of a series of sCIs for average ambient conditions representative of rural background, suburban background and urban kerbside (urban traffic) locations in the south-east UK in both winter and summer. The calculations make use of measured or inferred concentrations of a series of C₁ – C₆ alkenes, isoprene, α-pinene, limonene, O₃, NO₂, SO₂ and HC(O)OH; in conjunction with H₂O and (H₂O)₂ concentrations based on modelled temperature, and relative humidity data typical of the region. In this section, the key results are summarized and placed in context by comparison with reported results calculated for other locations, and in global modelling studies. Although the calculations presented here aim to take account of the most important production and loss routes, the estimates are inevitably subject to potential omissions and uncertainties in the sources and sinks of the sCIs (as discussed further in Sect. C4), in addition to uncertainties associated with the kinetic parameters and sCI yields, as discussed in earlier sections and in the data sheets in Supplements A and B.

8.1 sCI production rates

The formation of a series of 28 C₁ – C₁₀ sCIs from the speciation of 19 precursor alkenes was represented in the present calculations (see Supplement C for full details). The core set of C₁ – C₄ sCIs specifically considered in the present evaluation collectively makes an important contribution to the total production rate at each of the three locations; the respective winter and summer contributions lying in the ranges 88 – 91 % and ~~42–45~~ – ~~85~~86% (see Table 8). The speciation of this core set of sCIs is also presented in Fig. 6 for the example of the rural background location (Chilbolton Observatory). The formation of E- and Z-CH₃CHOO is most significant, because they are formed from propene and all the alk-2-enes in the applied speciation. They are also favoured because reaction with O₃ is a major (and sometimes the dominant) removal route for alk-2-enes (and other internal alkenes), because of their particularly rapid reactions with O₃ (see Table 1). CH₂OO also makes a notable contribution to the totals, because it is formed from all the alk-1-enes and isoprene. The higher alkyl-substituted sCIs make systematically lower contributions, primarily because their precursor alkenes have systematically lower abundances (see Table 8).

As also shown in Table 8, sCIs formed from biogenic hydrocarbons logically make an increased and important collective contribution under the summer conditions. This is particularly the case at the suburban background location (London Eltham), which has a mixture of trees and other vegetation in close proximity to the measurement site. The monoterpene (α-pinene and limonene) derived sCIs are calculated to be particularly significant, because ozonolysis generally makes a major contribution to the removal of endocyclic alkenes, by virtue of their particularly rapid reactions with O₃ (see Table 1). In contrast, the reaction of O₃ with isoprene is comparatively slow, its dominant removal reaction being with HO radicals. Therefore, the production rates of the isoprene-derived sCIs are approaching an order of magnitude lower than those of the

monoterpene-derived sCIs under the conditions represented here. However, isoprene-derived sCIs have been shown to make more important contributions globally, particularly in specific regions such as the Amazonian rain forest (e.g. Vereecken et al., 2017; Khan et al., 2018), as a result of the dominant contribution of isoprene to global biogenic VOC emissions.

8.2 sCI loss rates

5 Based on the information presented and discussed in Sects. 6 and 7, sCI removal by unimolecular decomposition and bimolecular reactions with H_2O , $(\text{H}_2\text{O})_2$, NO_2 , SO_2 and $\text{HC}(\text{O})\text{OH}$ was taken into account. Table 8 and Fig. 6 present information on the speciated and total first-order loss rates of the sCIs, and the contributions made by the series of removal reactions considered. Consistent with the analysis presented in Sect. 7, the results confirm that the major loss routes for most of the sCIs are either thermal decomposition, or reaction with $(\text{H}_2\text{O})_2$, supplemented by reaction with H_2O . As a result, these
10 reaction classes dominate total sCI removal under all conditions, with reaction with $(\text{H}_2\text{O})_2$ and H_2O being slightly more important in the winter, and thermal decomposition being slightly more important in the summer, based on the average of the three sites. As also indicated in Sect. 7, thermal decomposition tends to dominate the removal of *Z*- mono-substituted and di-substituted sCIs, with reaction with $(\text{H}_2\text{O})_2$ and H_2O dominating the removal of CH_2OO and *E*- mono-substituted sCIs.

The total first-order loss rates for the individual sCIs lie in approximate range 20 s^{-1} to $20,000 \text{ s}^{-1}$ for the full series of
15 considered conditions (see Supplement C). Those toward the low end of the range generally correspond to sCIs for which the dominant removal route is 1,4 H atom migration, occurring at only a modest rate (e.g. as in the cases of *Z*- CH_3CHOO and *E*- $(\text{CH}=\text{CH}_2)(\text{CH}_3)\text{COO}$), particularly under winter conditions; and for which the reactions with $(\text{H}_2\text{O})_2$ and H_2O are very slow. Those at the high end of the range generally correspond to *E*- monosubstituted sCIs (e.g. *E*- CH_3CHOO) for which the dominant removal reactions with $(\text{H}_2\text{O})_2$ and H_2O are very fast. The associated loss rates for selected sCIs in the former
20 category are sufficiently slow for removal by reaction with $\text{HC}(\text{O})\text{OH}$ and SO_2 (and to a lesser extent, reaction with NO_2) to make a notable contribution (e.g. *E*- $(\text{CH}=\text{CH}_2)(\text{CH}_3)\text{COO}$) in Fig. 6). In collective terms, however, these classes of reaction each makes only a small ($< 4 \%$) contribution to total sCI removal under the series of conditions considered here, as shown in Table 8.

8.3 Steady-state concentrations and speciation

25 The total first-order sCI loss rates indicated above correspond to individual sCI lifetimes lying in the range $50 \mu\text{s}$ to 50 ms , confirming that calculation of their concentrations using the steady-state approximation, described by Eq. (1), is valid. The resultant calculated steady-state concentrations of the core set of $\text{C}_1 - \text{C}_4$ sCIs for the rural background conditions are shown in Fig. 4, with the total sCI concentrations for the three scenarios given in Table 8. The totals calculated for the rural background conditions (389-386 molecule cm^{-3} and 384-375 molecule cm^{-3} for winter and summer, respectively) are broadly
30 consistent with the low annual average values simulated for the UK in the global modelling calculations of Vereecken et al.

(2017); and the concentrations calculated for the series of locations and conditions (up to about 1100 molecule cm⁻³) are comparable with those reported by Khan et al. (2018), based on similar UK calculations to those reported here.

The distributions of sCIs are generally dominated by a limited number of individual species, and show similarities to those reported elsewhere for locations with significant anthropogenic VOC emissions (e.g. Vereecken et al., 2017). Z-CH₃CHOO

5 is the most abundant sCI for all the considered scenarios, accounting for ~~76~~75-79 % of the totals for winter conditions, and ~~24~~25-77 % of the totals for summer conditions. This results from the combination of its high production rate (see Fig. 4) and its relatively slow removal rate (see Fig. 5), as discussed above. Its lowest contribution occurs for summer conditions at the suburban location, when the sCIs derived from α -pinene and limonene (and to a lesser extent, isoprene) are collectively dominant (67 %), as discussed in more detail in Supplement C. Given the relatively low biogenic hydrocarbon emission rates
10 in the UK, this result for a suburban location in London site gives a strong indication that biogenic hydrocarbon derived sCIs will dominate the global concentrations, as clearly demonstrated in the modelling studies presented by Vereecken et al. (2017) and Khan et al. (2018). Those studies reported important and widespread global contributions from sCIs derived from both isoprene and monoterpenes, although they report substantially different sCI concentrations. In the work of Vereecken et al. (2017), the rapid unimolecular decomposition rates calculated for many of the of sCIs (as also adopted in the present
15 work) strongly suppress the simulated concentrations compared with those reported by Khan et al. (2018). The resultant annual average sCI concentration at the surface maximizes at 7×10^3 molecule cm⁻³ over the Amazon basin, but is generally less than 2×10^3 molecule cm⁻³ over most of the globe (Vereecken et al., 2017). This further emphasizes the need for direct kinetics studies of a structurally diverse series of isoprene and terpene-derived sCIs, to help validate and refine the rate coefficients calculated in theoretical studies.

20 8.3 Oxidation of SO₂ and organic acids

The speciated sCI distributions have also been used to calculate the associated SO₂ oxidation rates. As shown in Table 8, the total oxidation rates are calculated to be between 0.006 % h⁻¹ and ~~0.031~~0.028 % h⁻¹. They broadly follow the simulated trend in total sCI concentrations, but also reflect that the rate coefficient values for individual species span almost an order of magnitude. These oxidation rates can be compared with a reference SO₂ oxidation rate of about 0.3 % h⁻¹ for reaction with
25 HO radicals at a concentration of 10⁶ molecule cm⁻³. This comparison is therefore consistent with the < 10 % annual average contribution to gas phase SO₂ oxidation for the UK, reported in the global modelling calculations of Vereecken et al. (2017) and Khan et al. (2018). The more widespread potential role of biogenic hydrocarbon derived sCIs in global SO₂ oxidation has also been considered in those modelling studies. Although the results possess some similarities, in terms of the relative regional variation, their role is much more limited in the Vereecken et al. (2017) calculations, because of the high calculated
30 decomposition rates applied to many of the sCIs, and the resultant suppression of sCI concentrations commented on above. Nevertheless, annual average contributions of up to about 70 % were still simulated for the terrestrial equatorial belt.

The largest bimolecular rate coefficients for sCI reactions that have been measured experimentally are those for reactions with organic acids such as HC(O)OH, CH₃C(O)OH and CF₃C(O)OH (see Table 5). HC(O)OH and CH₃C(O)OH are present in the troposphere in significant concentrations (e.g. Andreae et al., 1988; Millet et al., 2015; Bannan et al., 2017) due in part to their formation in the photochemical oxidation or ozonolysis of many VOCs, from both manmade and natural sources. As discussed above, and shown more widely in the calculation of Vereecken et al. (2017), the reactions with organic acids can make small but significant contributions to sCI removal in some regions. The total sCI concentrations calculated here suggest associated oxidation rates of (0.04 – 0.11) % h⁻¹ for HC(O)OH, and similar rates for CH₃C(O)OH. This can be compared with reference oxidation rates of about 0.16 % h⁻¹ and 0.25 % h⁻¹, for reaction of HC(O)OH and CH₃C(O)OH with HO radicals at a concentration of 10⁶ molecule cm⁻³. This indicates that reaction with sCIs makes an important contribution to the oxidation of these acids under the conditions considered here, with oxidation rates comparable to those via HO reaction calculated for equatorial regions in the global modelling study of Vereecken et al. (2017). It is noted that the reactions of larger sCIs and organic acids (e.g. derived from the ozonolysis of monoterpenes and sesquiterpenes) potentially forms highly oxidized, low-volatility products (hydroperoxyl-esters), which may play a role in secondary organic aerosol (SOA) formation (e.g. Tobias and Ziemann, 2001; Chhantyal-Pun et al., 2018).

As shown in Table 5, the reactions of sCIs with CF₃C(O)OH are particularly rapid. The total sCI concentrations calculated here suggest an associated oxidation rate of (0.1 – 0.3) % h⁻¹ for CF₃C(O)OH, compared with 0.25 % h⁻¹ for its reaction with HO radicals at a concentration of 10⁶ molecule cm⁻³. This demonstrates the potential importance of sCIs as gas phase oxidants for CF₃C(O)OH (and other perfluoro-carboxylic acids, C_nF_{2n+1}C(O)OH) over land masses. However, it is noted that the reaction of the resultant hydroperoxyl-fluoroester products with HO radicals probably reforms the perfluoro-carboxylic acids on a timescale of 1–2 days (Taatjes et al., 2019).

Current understanding of the atmospheric chemistry of sCIs therefore supports the original hypothesis of Cox and Penkett (1971; 1972) regarding their potential importance as atmospheric oxidants; identified from chamber measurements of SO₂ oxidation associated with alkene ozonolysis, and the observed effect of relative humidity on the oxidation rates. Considerable progress has since been made in the understanding the kinetics and mechanisms of alkene ozonolysis, resulting from both experimental and theoretical studies, with particular advances since the pioneering work of Taatjes and co-workers less than a decade ago (e.g. Welz et al., 2012; Taatjes et al., 2013) in the detection of sCIs and direct measurements of the kinetics of their reactions. However, significant uncertainties remain in some aspects of mechanistic understanding, including measurements of the yields of sCIs and their speciation in asymmetric alkene systems. The current evaluation has focused primarily on those sCIs for which direct kinetics measurements are available (i.e. CH₂OO, *E*-CH₃CHOO, *Z*-CH₃CHOO and (CH₃)₂COO), with some consideration also given to the C₄ intermediates formed from isoprene. Whilst these represent an important subset of atmospheric sCIs, it is recognized that an enormous variety of sCIs are generated, with particularly important global contributions from those generated from the ozonolysis of monoterpenes and sesquiterpenes. The concurrent progress in the theoretical treatment of the structure and reaction kinetics of sCIs in the gas phase has allowed the development of theory-based structure activity relationships (SARs) (Vereecken et al., 2017), which provide a basis for

representing the reactions of structurally complex sCIs in atmospheric mechanisms. As a result, there is a need for direct kinetics studies of a structurally diverse series of isoprene and terpene-derived sCIs, to help validate and refine the rate coefficients calculated in theoretical studies. In addition, the continued development of sensitive detection methods for sCIs (e.g. Berndt et al., 2017) that may eventually allow sCI concentrations to be measured in the field, would be valuable for evaluation and testing of the representation of sCI sources and sinks in atmospheric models.

Author contributions. All authors defined the scope of the work. RAC, MEJ and TJW developed and drafted the data sheets and manuscript. All authors reviewed, refined, and revised the manuscript and data sheets.

10 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements

The Chairman and members of the Task Group wish to express their appreciation to I.U.P.A.C. for the financial help that facilitated the preparation of this evaluation. We also acknowledge financial support from the following organisations: Department of Chemistry, University of Cambridge, the Swiss National Science Foundation, the Office Fédéral de
15 l'Education et de la Science, the Centre National de la Recherche Scientifique-Institut National des Sciences de l'Univers (CNRS-INSU), Orleans University, and Observatoire des Sciences de l'Univers en région Centre (OSUC). We thank Cathy Boone for developing and maintaining the website.

References

- Alam, M. S., Camredon, M., Rickard, A. R., Carr, T., Wyche, K. P., Hornsby, K. E., Monks, P. S. and Bloss, W. J.: Total
20 radical yields from tropospheric ethene ozonolysis, Phys. Chem. Chem. Phys., 13, 11002-11015, 2011.
- Alam, M. S., Rickard, A. R., Camredon, M., Wyche, K. P., Carr, T., Hornsby, K. E., Monks, P. S. and Bloss, W. J.: Radical product yields from the ozonolysis of short chain alkenes under atmospheric boundary layer conditions, J. Phys. Chem., 117, 12468-12483, 2013.
- Andreae, M. O., Talbot, R. W., Andreae, T. W. and Harriss, R. C.: Formic and acetic-acid over the central Amazon region,
25 Brazil .1. Dry season, J. Geophys. Res.-Atmospheres, 93, 1616-1624, 1988.
- Anglada, J. M. and Solé, A.: Impact of water dimer on the atmospheric reactivity of carbonyl oxides., Phys. Chem. Chem. Phys., 18, 17698-17712, 2016.
- Anglada, J. M., Aplincourt, P., Bofill, J. M. and Cremer, D.: Atmospheric formation of OH radicals and H₂O₂ from alkene ozonolysis under humid conditions, Chem. Phys. Chem., 3, 215-221, 2002.

- Anglada, J. M., González, J. and Torrent-Sucarrat, M.: Effects of the substituents on the reactivity of carbonyl oxides. A theoretical study on the reaction of substituted carbonyl oxides with water, *Phys. Chem. Chem. Phys.*, 13, 13034-13045, 2011.
- Aschmann, S. M., Arey, J. and Atkinson, R.: OH radical formation from the gas-phase reactions of O₃ with a series of terpenes, *Atmos. Environ.*, 36, 4347-4355, 2002.
- 5 Aschmann, S. M., Tuazon, E. C., Arey, J. and Atkinson, R.: Products of the gas-phase reaction of O₃ with cyclohexene, *J. Phys. Chem. A*, 107, 2247-2255, 2003.
- Aschmann, S. M., Arey, J., and Atkinson, R.: OH radical formation from the gas-phase reactions of O₃ with methacrolein and methyl vinyl ketone, *Atmos. Environ.*, 30, 2939-2943, 1996.
- Atkinson, R. and Aschmann, S. M.: OH radical production from the gas-phase reactions of O₃ with a series of alkenes under
10 atmospheric conditions, *Environ. Sci. Technol.*, 27, 1357-1363, 1993.
- Atkinson, R., Aschmann, S. M., Arey, J. and Shorees, B.: Formation of OH radicals in the gas phase reactions of O₃ with a series of terpenes, *J. Geophys. Res.*, 97, 6065-6073, 1992.
- Bannan, T. J., Booth, A. M., Le Breton, M., Bacak, A., Muller, J. B. A., Leather, K. E., Khan, M. A. H., Lee, J. D., Dunmore, R. E., Hopkins, J. R., Fleming, Z. L., Sheps, L., Taatjes, C. A., Shallcross, D. E. and Percival, C. J.: Seasonality of
15 formic acid (HCOOH) in London during the ClearfLo campaign, *J. Geophys. Res. Atmos.*, 122, 12488-12498, 2017.
- Barber, V. P., Pandit, S., Green, A. M., Trongsirawat, N., Walsh, P. J., Klippenstein, S. J. and Lester, M. I.: Four-carbon Criegee intermediate from isoprene ozonolysis: Methyl vinyl ketone oxide synthesis, infrared spectrum, and OH production, *J. Am. Chem. Soc.*, 140, 10866-10880, <https://doi.org/10.1021/jacs.8b06010>, 2018.
- Beames, J. M., Liu, F., Lu, L., and Lester, M. I.: Ultraviolet spectrum and photochemistry of the simplest Criegee
20 intermediate CH₂OO J. Amer. Chem. Soc., 134, 20045-20048, <https://doi.org/10.1021/ja310603j>, 2012.
- Beck, M., Winterhalter, R., Herrmann, F., and Moortgat, G. K.: The gas-phase ozonolysis of α-humulene, *Phys. Chem. Chem. Phys.*, 13, 10970-11001, 2011.
- Berndt, T., Böge, O and Stratmann, F: Gas-phase ozonolysis of alpha-pinene: gaseous products and particle formation, *Atmos. Environ.*, 37, 3933-3945, 2003.
- 25 Berndt, T., Jokinen, T., Sipilä, M., Mauldin, R. L., Herrmann, H., Stratmann, F., Junninen, H. and Kulmala, M.: H₂SO₄ formation from the gas-phase reaction of stabilized Criegee intermediates with SO₂: Influence of water vapour content and temperature, *Atmos. Environ.*, 89, 603-612, 2014.
- Berndt, T., Herrmann, H. and Kurtén, T.: Direct probing of Criegee intermediates from gas-phase ozonolysis using chemical ionization mass spectrometry, J. Am. Chem. Soc., 139, 13387-13392, 2017.

- Calvert, J. G., Mellouki, A., Orlando, J. J., Pilling, M. J., and Wallington, T. J.: The mechanisms of atmospheric oxidation of the alkenes, Oxford University Press, Oxford, ISBN 978-0-19513177-2, 2000.
- Calvert, J. G., Orlando, J. J., Stockwell, W. R., and Wallington, T. J.: The mechanisms of reactions influencing atmospheric ozone, Oxford University Press, Oxford, ISBN 978-0-19023302-0, 2015.
- Campos-Pineda, M. and Zhang, J.: Product yields of stabilized Criegee intermediates in the ozonolysis reactions of *cis*-2-butene, 2-methyl-2-butene, cyclopentene, and cyclohexene, *Sci. China Chem.*, 61, 850-856, <https://doi.org/10.1007/s11426-017-9229-0>, 2018.
- Campos-Pineda, M. and Zhang, J.: Low-pressure yields of stabilized Criegee intermediates CH_3CHOO and $(\text{CH}_3)_2\text{COO}$ in ozonolysis of *trans*-2-butene and 2,3-dimethyl-2-butene, *Chem. Phys. Lett.*, 683, 647-652, 2017.
- Caravan, R. L., Vansco, M. F., Au, K., Khan, M. A. H., Li, Y.-L., Winiberg, F. A. F., Zuraski, K., Lin, Y.-H., Chao, W., Trongsiriwat, N., Walsh, P. J., Osborn, D. L., Percival, C. J., Lin, J.-J. M., Shallcross, D. E., Sheps, L., Klippenstein, S. J., Taatjes, C. A., and Lester, M. I.: *Proc. Nat. Acad. Sci.*, 117 (18), 9733-9740, doi: 10.1073/pnas.1916711117, 2020.
- Chhantyal-Pun, R., McGillen, M. R., Beames, J. M., Khan, M. A. H., Percival, C. J., Shallcross, D. E., and Orr-Ewing, A. J.: Temperature-dependence of the rates of reaction of trifluoroacetic acid with Criegee intermediates *Angew. Chem. Int. Ed.*, 56, 9044-9047, 2017a.
- Chhantyal-Pun, R., Welz, O., Savee J. D., Eskola, A. J., Lee, E. P. F., Blacker, L., Hill, H. R., Ashcroft, M., Khan, M. A. H., Lloyd-Jones, G. C., Evans, L., Rotavera, B., Rotavera H., Osborn, D. L., Mok, D. K. W., Dyke, J. M., Shallcross, D. E., Percival, C. J., Orr-Ewing, A. J. and Taatjes, C. A.: Direct measurements of unimolecular and bimolecular reaction kinetics of the Criegee intermediate $(\text{CH}_3)_2\text{COO}$, *J. Phys. Chem. A*, 121, 4-15, <https://doi.org/10.1021/acs.jpca.6b07810>, 2017b.
- Chhantyal-Pun, R., Rotavera, B., McGillen, M. R., Khan, M. A. H., Eskola, A. J., Caravan, R. L., Blacker, L., Tew, D. P., Osborn, D. L., Percival, C. J., Shallcross, D. E. and Orr-Ewing A. J.: Criegee intermediate reactions with carboxylic acids: a potential source of secondary organic aerosol in the atmosphere, *ACS Earth Space Chem.*, 2, 833-842, 2018.
- Chew, A. A. and Atkinson, R.: OH radical formation yields from the gas-phase reactions of O_3 with alkenes and monoterpenes, *J. Geophys. Res.*, 101, 28649-28653, 1996.
- Chuong, B., Zhang, J. and Donahue, N. M.: Cycloalkene ozonolysis: collisionally mediated mechanistic branching, *J. Am. Chem. Soc.*, 126, 12363-12373, 2004.
- Cox, R. A. and Penkett, S. A.: Oxidation of atmospheric SO_2 by products of the ozone-olefin reaction, *Nature*, 230, 321-322, 1971.

- Cox, R. A. and Penkett, S. A.: Aerosol formation from sulphur dioxide in the presence of ozone and olefinic hydrocarbons, J. Chem. Soc., Faraday Trans. 1, 68, 1735-1753, 1972.
- Criegee, R., Blust, G. and Zinke, H.: Eine neuartige synthese von ozoniden, Chem. Ber., 87,766-768, 1954.
- Donahue, N. M., Kroll, J. H., Anderson, J. G. and Demerjian, K. L.: Direct observation of OH production from the ozonolysis of olefins, Geophys. Res. Lett., 25(1), 59-62, <https://doi.org/10.1029/97GL53560>, 1998.
- Drozd, G. T. and Donahue, N. M.: Pressure dependence of stabilized Criegee intermediate formation from a sequence of alkenes, J. Phys. Chem. A, 115, 4381-4387, 2011.
- Drozd, G. T., Kroll, J. and Donahue, N. M.: 2, 3-Dimethyl-2-butene (TME) ozonolysis: pressure dependence of stabilized Criegee intermediates and evidence of stabilized vinyl hydroperoxides, J. Phys. Chem. A, 115, 161-166, 2011.
- 10 [Fang, Y., Liu, F., Barber, V. P., Klippenstein, S. J., McCoy, A. B. and Lester, M. I.: Deep tunneling in the unimolecular decay of CH₃CHOO Criegee intermediates to OH radical products, J. Chem. Phys. 145, 234308, 2016.](#)
- [Fang, Y., Liu, F., Barber, V. P., Klippenstein, S. J., McCoy, A. B. and Lester, M. I.: Tunneling effects in the unimolecular decay of \(CH₃\)₂COO Criegee intermediates to OH radical products, J. Chem. Phys. 146, 134307, <https://doi.org/10.1063/1.4979297>, 2017.](#)
- 15 Fenske, J. D., Hasson, A. S., Paulson, S. E., Kuwata, K. T., Ho, A. and Houk, K. N.: The pressure dependence of the OH radical yield from ozone-alkene reactions, J. Phys. Chem. A, 104, 7821-7833, 2000.
- Finlayson, B. J., Pitts, J. N. and Akimoto, H.: Production of vibrationally excited OH in chemiluminescent ozone-olefin reactions, Chem. Phys. Lett., 12, 495-498, 1972.
- Finlayson-Pitts, B. J. and Pitts, J. N.: Chemistry of the upper and lower atmosphere: theory, experiments, and applications, Academic Press, doi:10.1016/B978-012257060-5/50010-1, 2000.
- 20 Forester, C. D. and Wells, J. R.: Hydroxyl radical yields from reactions of terpene mixtures with ozone, Indoor Air, 21, 400-409, 2011.
- Griesbaum, K., Miclaus, V., and Jung, I. C.: Isolation of ozonides from gas-phase ozonolyses of terpenes, Environ. Sci. Technol., 32, 647-649, 1998.
- 25 Hakala, J. P. and Donahue, N. M.: Pressure-dependent Criegee intermediate stabilization from alkene ozonolysis, J. Phys. Chem. A, 120, 2173-2178, 2016.
- Hakala, J. P. and Donahue, N. M.: Pressure stabilization of Criegee intermediates formed from symmetric trans-alkene ozonolysis, J. Phys. Chem. A, 122, 9426-9434, 2018.

- Hakola, H., Arey, J., Aschmann, S. M. and Atkinson, R.: Product formation from the gas-phase reactions of OH radicals and O₃ with a series of monoterpenes, *J. Atmos. Chem.*, 18, 75-102, 1994.
- Hasson, A. S., Orzechowska, G. and Paulson, S. E.: Production of stabilized Criegee intermediates and peroxides in the gas phase ozonolysis of alkenes 1. Ethene, *trans*-2-butene, and 2,3-dimethyl-2-butene, *J. Geophys. Res.-Atmos.*, 106, 34131-34142, 2001a.
- 5 Hasson, A. S., Ho, A. W., Kuwata, K. T. and Paulson, S. E.: Production of stabilized Criegee intermediates and peroxides in the gas phase ozonolysis of alkenes 2. Asymmetric and biogenic alkenes, *J. Geophys. Res.-Atmos.*, 106, 34143-34153, 2001b.
- Hasson, A. S., Chung, M. Y., Kuwata, K. T., Converse, A. D., Krohn, D. and Paulson, S. E.: Reaction of Criegee intermediates with water vapor - An additional source of OH radicals in alkene ozonolysis? *J. Phys. Chem. A*, 107, 6176-6182, 2003.
- 10 Hatakeyama, S. and Akimoto, H.: Reactions of Criegee intermediates in the gas-phase, *Res. Chem. Intermed.*, 20, 503-524, 1994.
- Hatakeyama, S., Kobayashi, H. and Akimoto, H.: Gas-phase oxidation of SO₂ in the ozone-olefin reactions, *J. Phys. Chem.*, 88, 4736-4739, 1984.
- Hatakeyama, S., Kobayashi, H., Lin, Z.-Y., Takagi, H. and Akimoto, H.: Mechanism for the reaction of CH₂OO with SO₂, *J. Phys. Chem.*, 90, 4131-4135, 1986.
- 15 Herrmann, F., Winterhalter, R., Moortgat, G. K. and Williams, J.: Hydroxyl radical (OH) yields from the ozonolysis of both double bonds for five monoterpenes, *Atmos. Environ.*, 44, 3458-3464, 2010.
- Horie, O. and Moortgat, G. K.: Decomposition pathways of the excited Criegee intermediates in the ozonolysis of simple alkenes, *Atmos. Environ.*, 25A, 1881-1896, 1991
- Horie, O., Schafer, C. and Moortgat, G. K.: High reactivity of hexafluoro acetone toward Criegee intermediates in the gas-phase ozonolysis of simple alkenes, *Int. J. Chem. Kinet.*, 31, 261-269, 1999.
- 20 Huang, H.-L., Chao, W. and Lin, J. J.-M.: Kinetics of a Criegee intermediate that would survive high humidity and may oxidize atmospheric SO₂, *Proc. Natl. Acad. Sci.*, 112(35), 10857-10862, doi: 10.1073/pnas.1513149112, 2015.
- Jenkin, M. E., Wyche, K. P., Evans, C. J., Carr, T., Monks, P. S., Alfarra, M. R., Barley, M. H., McFiggans, G. B., Young, J. C., and Rickard, A. R.: Development and chamber evaluation of the MCM v3.2 degradation scheme for β-caryophyllene, *Atmos. Chem. Phys.*, 12, 5275-5308, 2012.
- 25 Kan, C. S., Su, F., Calvert, J. G. and Shaw, J. H.: Mechanism of the ozone-ethene reaction in dilute N₂/O₂ mixtures near 1-atm Pressure, *J. Phys. Chem.*, 85, 2359-2363, 1981.
- Khan, M. A. H., Percival, C. J., Caravan, R. L., Taatjes, C. A. and Shallcross, D. E.: Criegee intermediates and their impacts on the troposphere, *Environ. Sci.: Processes Impacts*, 20, 437-453, 2018.

- Kroll, J. H., Hanisco, T. F., Donahue, N. M., Demerjian, K. L. and Anderson, J. G.: Accurate, direct measurements of OH yields from gas-phase ozone-alkene reactions using an in situ LIF Instrument, *Geophys. Res. Lett.*, 28(20), 3863-3866, <https://doi.org/10.1029/2001GL013406>, 2001a.
- Kroll, J. H., Clarke, J. S., Donahue, N. M., Anderson, J. G. and Demerjian, K. L.: Mechanism of HO_x formation in the gas-phase ozone-alkene reaction. 1. Direct, pressure-dependent measurements of prompt OH yields, *J. Phys. Chem. A*, 105, 1554-1560, 2001b.
- Kroll, J. H., Sahay, S. R., Anderson, J. G., Demerjian, K. L. and Donahue, N. M.: Mechanism of HO_x formation in the gas-phase ozone-alkene reaction. 2. Prompt versus thermal dissociation of carbonyl oxides to form OH, *J. Phys. Chem. A*, 105, 4446-4457, 2001c.
- 10 Kuwata, K. T., Hermes, M. R., Carlson, M. J. and Zogg, C. K.: Computational studies of the isomerization and hydration reactions of acetaldehyde oxide and methyl vinyl carbonyl oxide, *J. Phys. Chem. A*, 114, 9192-9204, 2010.
- Li, H., Fang, Y., Beames, J. M. and Lester, M. I.: Velocity map imaging of O-atom products from UV photodissociation of the CH₂OO Criegee intermediate, *J. Chem. Phys.*, 142, 214312, <https://doi.org/10.1063/1.4921990>, 2015.
- Long, B., Bao, J. L. and Truhlar, D.G.: Atmospheric chemistry of Criegee intermediates: Unimolecular reactions and reactions with water, *J. Am. Chem. Soc.* 138, 43, 14409-14422, <https://doi.org/10.1021/jacs.6b08655>, 2016.
- 15 Long, B., Bao, J. L. and Truhlar, D.G.: Rapid unimolecular reaction of stabilized Criegee intermediates and implications for atmospheric chemistry, *Nat. Commun.*, 10, 2003, <https://doi.org/10.1038/s41467-019-09948-7>, 2019.
- Mackenzie-Rae, F. A., Karton, A. and Saunders, S. M.: Computational investigation into the gas-phase ozonolysis of the conjugated monoterpene alpha-phellandrene, *Phys. Chem. Chem. Phys.*, 18, 27991-28002, 2016.
- 20 Malkin, T. L., Goddard, A., Heard, D. E. and Seakins, P. W.: Measurements of OH and HO₂ yields from the gas phase ozonolysis of isoprene, *Atmos. Chem. Phys.*, 10, 1441-1459, 2010.
- McGill, C. D., Rickard, A. R., Johnson, D. and Marston, G.: Product yields in the reactions of ozone with Z-but-2-ene, E-but-2-ene and 2-methylbut-2-ene, *Chemosphere*, 38, 1205-1212, 1999.
- Mihelcic, D., Heitlinger, M., Kley, D., Musgen P. and Volz-Thomas, A.: Formation of hydroxyl and hydroperoxy radicals in the gas-phase ozonolysis of ethene, *Chem. Phys. Lett.*, 301, 559-564, 1999.
- 25 Millet, D. B., Baasandorj, M., Farmer, D. K., Thornton, J. A., Baumann, K., Brophy, P., Chaliyakunnel, S., de Gouw, J. A., Graus, M., Hu, L., Koss, A., Lee, B. H., Lopez-Hilfiker, F. D., Neuman, J. A., Paulot, F., Pollack, I. B., Ryerson, T. B., Warneke, C., Williams, B. J. and Xu, J.: A large and ubiquitous source of atmospheric formic acid, *Atmos. Chem. Phys.*, 15, 6283-6304, 2015.

- Neeb, P., Horie, O. and Moortgat, G. K.: Gas-phase ozonolysis of ethene in the presence of hydroxylic compounds, *Int. J. Chem. Kinet.*, 28, 721-730, 1996.
- Neeb, P., Sauer, F., Horie, O. and Moortgat, G. K.: Formation of hydroxymethyl hydroperoxide and formic acid in alkene ozonolysis in the presence of water vapour, *Atmos. Environ.*, 31, 417-1423, 1997.
- 5 Neeb, P., Horie, O. and Moortgat, G. K.: The ethene-ozone reaction in the gas phase, *J. Phys. Chem. A*, 102, 6778-6785, 1998.
- Neeb, P. and Moortgat, G. K.: Formation of OH radicals in the gas-phase reaction of propene, isobutene, and isoprene with O₃: Yields and mechanistic implications *J. Phys. Chem. A*, 103, 9003-9012, 1999.
- Newland, M. J., Rickard, A. R., Alam, M. S., Vereecken, L., Muñoz, A., Ródenas, M. and Bloss, W. J.: Kinetics of stabilised Criegee intermediates derived from alkene ozonolysis: reactions with SO₂, H₂O and decomposition under boundary layer
- 10 conditions, *Phys. Chem. Chem. Phys.*, 17, 4076-4088, 2015a.
- Newland, M. J., Rickard, A. R., Vereecken, L., Muñoz, A., Ródenas, M., and Bloss, W. J.: Atmospheric isoprene ozonolysis: impacts of stabilised Criegee intermediate reactions with SO₂, H₂O and dimethyl sulfide, *Atmos. Chem. Phys.*, 15, 9521-9536, <https://doi.org/10.5194/acp-15-9521-2015>, 2015b.
- Newland, M. J., Rickard, A. R., Sherwen, T., Evans, M. J., Vereecken, L., Muñoz, A., Ródenas, M., and Bloss, W. J.: The
- 15 atmospheric impacts of monoterpene ozonolysis on global stabilised Criegee intermediate budgets and SO₂ oxidation: experiment, theory and modelling, *Atmos. Chem. Phys.*, 18, 6095-6120, <https://doi.org/10.5194/acp-18-6095-2018>, 2018.
- Newland, M. J., Nelson, B. S., Muñoz, A., Ródenas, M., Vera, T., Tárrega, J., and Rickard, A. R.: Trends in stabilisation of
- Criegee intermediates from alkene ozonolysis, *Phys. Chem. Chem. Phys.*, 22, 13698-13706, <https://doi.org/10.1039/D0CP00897D>, 2020.
- 20 Nguyen, T. L., Peeters, J. and Vereecken, L.: Theoretical study of the gas-phase ozonolysis of β-pinene (C₁₀H₁₆), *Phys. Chem. Chem. Phys.*, 11, 5643-5656, 2009a.
- Nguyen, T. L., Winterhalter, R., Moortgat, G., Kanawati, B., Peeters, J. and Vereecken, L.: The gas-phase ozonolysis of β-caryophyllene (C₁₅H₂₄). Part II: a theoretical study, *Phys. Chem. Chem. Phys.*, 11, 4173-4183, 2009b.
- Nguyen, T. B., Tyndall, G. S., Crounse, J. D., Teng, A. P., Bates, K. H., Schwantes, R. H., Coggon, M. M., Zhang, L., Feiner, P.,
- 25 Milller, D. O., Skog, K. M., Rivera-Rios, J. C., Dorris, M., Olson, K. F., Koss, A., Wild, R. J., Brown, S. S., Goldstein, A. H., de Gouw, J. A., Brune, W. H., Keutsch, F. N., Seinfeld, J. H. and Wennberg, P. O.: Atmospheric fates of Criegee intermediates in the ozonolysis of isoprene, *Phys. Chem. Chem. Phys.*, 18, 10241-10254, 2016.
- Niki, H., Maker, P. D., Savage, C. M. and Breitenbach, L. P.: Fourier Transform IR spectroscopic observation of propylene ozonide in the gas phase reaction of ozone-*cis*-2-butene-formaldehyde, *Chem. Phys. Lett.*, 46, 327-330, 1977.

- Niki, H., Maker, P. D., Savage, C. M., Breitenbach, L. P. and Hurley, M. D.: FTIR spectroscopic study of the mechanism for the gas-phase reaction between ozone and tetramethylethylene, *J. Phys. Chem.*, 91, 941-946, 1987.
- Olzmann, M., Kraka, E., Cremer, D., Gutbrod, R. and Andersson, S.: Energetics, kinetics, and product distributions of the reactions of ozone with ethene and 2,3-dimethyl-2-butene, *J. Phys. Chem. A*, 101, 9421-9429, 1997.
- 5 Orzechowska, G. and Paulson, S. E.: Production of OH radicals from the reactions of C₄-C₆ internal alkenes and styrenes with ozone in the gas phase, *Atmos. Environ.*, 36, 571-581, 2002.
- Osborn, D. L. and Taatjes, C. A.: The physical chemistry of Criegee intermediates in the gas phase, *Int. Rev. Phys. Chem.*, 34, 309-360, 2015.
- Paulson, S. E., Chung, M., Sen, A. D. and Orzechowska, G.: Measurement of OH radical formation from the reaction of ozone
10 with several biogenic alkenes, *J. Geophys. Res.-Atmos.*, 103, 25533-25539, 1998.
- Paulson, S. E., Fenske, J. D., Sen, A. D. and Callahan, T. W.: A novel small-ratio relative-rate technique for measuring OH formation yields from the reactions of O₃ with alkenes in the gas phase, and its application to the reactions of ethene and propene, *J. Phys. Chem. A*, 103, 2050-2059, 1999.
- 15 Peltola, J., Seal, P., Inkilä, A., and Eskola, A.: Time-resolved, broadband UV-absorption spectrometry measurements of Criegee intermediate kinetics using a new photolytic precursor: unimolecular decomposition of CH₂OO and its reaction with formic acid, *Phys. Chem. Chem. Phys.*, 22, 11797-11808, <https://doi.org/10.1039/D0CP00302F>, 2020.
- Presto, A. A. and Donahue, N. M.: Ozonolysis fragment quenching by nitrate formation: The pressure dependence of prompt OH radical formation, *J. Phys. Chem. A*, 108, 9096-9104, 2004.
- Qi, B., Yang, B., Wang, Z. Q., Yang, H. Y. and Liu, L.: Production of radicals in the ozonolysis of propene in air, *Sci. China, Ser. B: Chem.*, 52, 356-361, 2009.
20
- Ren, Y., Grosselin, B., Daële, V. and Mellouki, A.: Investigation of the reaction of ozone with isoprene, methacrolein and methyl vinyl ketone using the HELIOS chamber, *Faraday Discuss.*, 200, 289-311, 2017.
- Rickard, A. R., Johnson, D., McGill, C. D. and Marston, G.: OH yields in the gas-phase reactions of ozone with alkenes, *J. Phys. Chem. A*, 103, 7656-7664, 1999.
- 25 Ryzhkov, A. B. and Ariya, P. A.: A theoretical study of the reactions of parent and substituted Criegee intermediates with water and the water dimer, *Phys. Chem. Chem. Phys.*, 6, 5042-5050, 2004.
- Samanta, K., Beames, J. M., Lester M. I., and Subotnik, J. E.: Quantum dynamical investigation of the simplest Criegee intermediate CH₂OO and its O–O photodissociation channels, *J. Chem. Phys.* 141, 134303, <https://doi.org/10.1063/1.4894746>, 2014.

- Sheps, L.: Absolute ultraviolet absorption spectrum of a Criegee intermediate CH_2OO , *J. Phys. Chem. Lett.*, 4, 4201-4205, 2013.
- Sheps, L., Scully, A. M. and Au, K.: UV absorption probing of the conformer-dependent reactivity of a Criegee intermediate CH_3CHOO *Phys. Chem. Chem. Phys.*, 16, 26701-26706, 2014.
- 5 Shu, Y. and Atkinson, R.: Rate constants for the gas-phase reactions of O_3 with a series of terpenes and OH radical formation from the O_3 reactions with sesquiterpenes at 296 ± 2 K, *Int. J. Chem. Kinet.*, 26, 1193-1205, 1994.
- Siese, M., Becker, K. H., Brockmann, K. J., Geiger, H., Hofzumahaus, A., Holland, F., Mihelcic, D. and Wirtz, K.: Direct measurement of OH radicals from ozonolysis of selected alkenes: A EUPHORE simulation chamber study, *Environ. Sci. Technol.*, 35, 4660-4667, 2001.
- 10 Sipilä, M., Jokinen, T., Berndt, T., Richters, S., Makkonen, R., Donahue, N. M., Mauldin III, R. L., Kurtén, T., Paasonen, P., Sarnela, N., Ehn, M., Junninen, H., Rissanen, M. P., Thornton, J., Stratmann, F., Herrmann, H., Worsnop, D. R., Kulmala, M., Kerminen, V.-M., and Petäjä, T.: Reactivity of stabilized Criegee intermediates (sCIs) from isoprene and monoterpene ozonolysis toward SO_2 and organic acids, *Atmos. Chem. Phys.*, 14, 12143-12153, <https://doi.org/10.5194/acp-14-12143-2014>, 2014.
- [Stephenson, T. A. and Lester, M. I.: Unimolecular decay dynamics of Criegee intermediates: Energy-resolved rates, thermal rates, and their atmospheric impact, *Int. Rev. Phys. Chem.*, 39 \(1\), 1-33, <https://doi.org/10.1080/0144235X.2020.1688530>, 2020.](#)
- 15 Su, F., Calvert, J. G. and Shaw, J. H.: A FT-IR Spectroscopic study of the ozone-ethene reaction mechanism in O_2 -rich mixtures, *J. Phys. Chem.*, 84, 239-246, 1980.
- Taatjes, C. A., Welz, O., Eskola, A. J., Savee, J. D., Osborn, D. L., Lee, E. P. F., Dyke, J. M., Mok, D. W. K., Shallcross, D. E., and Percival, C. J.: Direct measurement of Criegee intermediate (CH_2OO) reactions with acetone, acetaldehyde, and
- 20 hexafluoroacetone, *Phys. Chem. Chem. Phys.*, 14, 10391-10400, 2012.
- Taatjes, C. A., Welz, C. A.; Eskola, A. J., Savee, J. D., Scheer, A. M., Shallcross, D. E., Rotavera, B., Lee, E. P. F., Dyke, J. M., Mok, D. K. W., Osborn, D. L., and Percival, C. J.: Direct measurements of conformer-dependent reactivity of the Criegee intermediate CH_3CHOO , *Science*, 340, 177-181, 2013.
- Taatjes, C.A., Khan, M. A. H., Eskola, A. J., Percival, C. J., Osborn, D. L., Wallington, T. J., and Shallcross, D. E.: Reaction with
- 25 Criegee intermediates: An important atmospheric fate of perfluorocarboxylic acids, *Environ. Sci. Technol.*, 53, 1245-1251, 2019.
- [Ting, W-L, Chen, Y-H., Chao, W., Smith, M.C., and Lin, J Jr -M.: The UV absorption spectrum of the simplest Criegee intermediate \$\text{CH}_2\text{OO}\$, *Phys. Chem. Chem. Phys.*, 16, 10438-10443, 2014.](#)
- Tobias, H. J. and Ziemann, P. J.: Kinetics of the gas-phase reactions of alcohols, aldehydes, carboxylic acids, and water with the C_{13} stabilized Criegee intermediate formed from ozonolysis of 1-tetradecene, *J. Phys. Chem. A*, 105 (25), 6129-6135, 2001.

Vansco, M. F., Li, H. and Lester, M. I.: Prompt release of O¹D products upon UV excitation of CH₂OO Criegee intermediates, J. Chem. Phys. 147, 013907, <https://doi.org/10.1063/1.4977987>, 2017.

Vansco, M. F., Marchetti, B. and Lester, M. I.: Electronic spectroscopy of methyl vinyl ketone oxide: A four-carbon unsaturated Criegee intermediate from isoprene ozonolysis, J. Chem. Phys., 149, 244309, <https://doi.org/10.1063/1.5064716>, 2018.

- 5 Vansco, M. F., Marchetti, B. Trongsiriat, N., Bhagde, T., Wang, G., Walsh, P. J., Klippenstein, S. J. and Lester, M. I.: Synthesis, electronic spectroscopy, and photochemistry of methacrolein oxide: A four-carbon unsaturated Criegee intermediate from isoprene ozonolysis, J. Am. Chem. Soc., 141 (38), 15058-15069, <https://doi.org/10.1021/jacs.9b05193>, 2019.

Vereecken, L., Harder, H. and Novelli, A.: The reaction of Criegee intermediates with NO, RO₂, and SO₂, and their fate in the atmosphere, Phys. Chem. Chem. Phys., 14, 14682-14695, <https://doi.org/10.1039/C2CP42300F>, 2012.

- 10 Vereecken, L., Novelli, A. and Taraborrelli, D.: Unimolecular decay strongly limits the atmospheric impact of Criegee intermediates, Phys. Chem. Chem. Phys., 19, 31599-31612, doi: 10.1039/c7cp05541b, 2017.

~~Vibenholt, A., Nørgaard, A. W., Clausen, P. A. and Wolkoff, P.: Formation and stability of secondary ozonides from monoterpenes studied by mass spectrometry, Chemosphere, 76, 572-577, 2009.~~

- 15 Welz, O., Savee, J. D., Osborn, D. L., Vasu, S. S., Percival, C. J., Shallcross, D. E. and Taatjes, C. A.: Direct kinetic measurements of Criegee intermediate (CH₂OO) formed by reaction of CH₂I with O₂, Science, 335, 204-207, 2012.

Winterhalter, R., Neeb, P., Grossmann, D., Kolloff, A., Horie, O., and Moortgat, G. K.: Products and mechanism of the gas phase-reaction of ozone with β-pinene, J. Atmos. Chem., 35, 165-197, 2000.

Winterhalter, R., Herrmann, F., Kanawati, B., Nguyen, T. L., Peeters, J., Vereecken, L., and Moortgat, G. K.: The gas-phase ozonolysis of β-caryophyllene (C₁₅H₂₄). Part I: an experimental study, Phys. Chem. Chem. Phys., 11, 4152-4172, 2009.

- 20 Yao, L., Ma, Y., Wang, L., Zheng, J., Khalizov, A., Chen, M., Zhou, Y., Qi, L., and Cui, F.: Role of stabilized Criegee intermediate in secondary organic aerosol formation from the ozonolysis of α-cedrene, Atmos. Environ., 94, 448-457, 2014.

Zhang, D., Lei, W. and Zhang, R.: Mechanism of OH formation from ozonolysis of isoprene: kinetics and product yields, Chem. Phys. Lett., 358, 171, 2002.







Figure 1: Mechanism of the reaction of ozone with alkenes, showing formation of Criegee intermediates (CIs). The general types of reaction available for stabilised Criegee intermediates (sCIs) are also illustrated for one example. The substituents R_1 to R_4 can be either H atoms or organic groups, although the illustrated sCI/CI vinyl hydroperoxide route is unavailable if $\text{R}_2 = \text{H}$. Note that in the case of endocyclic C=C bonds in cycloalkenes, the initially formed carbonyl and CI moieties are substituents of the same organic product.




CH₂OO




Z-CH₃CHOO




E-CH₃CHOO




(CH₃)₂COO




Z-(CH=CH₂)(CH₃)COO



E-(CH=CH₂)(CH₃)COO



Z-(C(CH₃)=CH₂)CHOO



E-(C(CH₃)=CH₂)CHOO

5

Figure 2: Structures of the stabilised Criegee intermediates considered in the present study, and the nomenclature assigned. In the cases of the di-substituted (isoprene-derived) C₄ intermediates, the *Z*- and *E*- notations specify the orientation of the first named substituent (which has the higher Cahn-Ingold-Prelog priority) relative to the CI moiety. The displayed rotamers of the C₄ intermediates are calculated to be in near equilibrium under atmospheric conditions (Vereecken et al., 2017), and are assumed to act as a single species in each case.

10




Figure 3: Recommended spectra for CH_2OO , $\text{Z-CH}_3\text{CHOO}$, $\text{E-CH}_3\text{CHOO}$ and $(\text{CH}_3)_2\text{COO}$.





Figure 4: Production rates (upper panel) and steady-state concentrations (lower panel) for the core set of C₁ – C₄ sCIs for representative rural background conditions in the south-east UK. Note that the information is presented on log scales, with cut-offs of 10 molecule cm⁻³ s⁻¹ and 1 molecule cm⁻³, respectively. The total sCI production rates and concentrations are given in Table 8. Results for an extended series of 28 sCIs, and for suburban and urban traffic conditions, are presented in Supplement C.



5 **Figure 5:** Contributions of the unimolecular and bimolecular removal routes for the core set of C₁ – C₄ sCIs under winter conditions (upper panel) and summer conditions (lower panel) for representative rural background conditions in the south-east UK. The data label shows the total removal rate (in s^{-1}) for the given sCI. Results for an extended series of sCIs, and for suburban and urban traffic conditions, are presented in Supplement C.

Table 1. Summary of recommended rate coefficients for reactions of O₃ with alkenes

Reaction ID	Alkene	k_{298} cm ³ molecule ⁻¹ s ⁻¹	$\Delta \log k_{298}$	$k(T)$ cm ³ molecule ⁻¹ s ⁻¹	<i>T</i> range	$\Delta(E/R)$ K
<i>Small alkene reactions – based on data sheets in Supplement A Sect. A1</i>						
Ox_VOC5	ethene	1.55×10^{-18}	± 0.08	$6.82 \times 10^{-15} \exp(-2500/T)$	180-360	± 100
Ox_VOC6	propene	1.05×10^{-17}	± 0.15	$5.77 \times 10^{-15} \exp(-1880/T)$	230-370	± 100
Ox_VOC16	but-1-ene	1.0×10^{-17}	± 0.08	$3.55 \times 10^{-15} \exp(-1750/T)$	220-370	± 200
Ox_VOC17	<i>cis</i> -but-2-ene	1.3×10^{-16}	± 0.05	$3.37 \times 10^{-15} \exp(-970/T)$	220-370	± 200
Ox_VOC18	<i>trans</i> -but-2-ene	2.0×10^{-16}	± 0.1	$7.0 \times 10^{-15} \exp(-1060/T)$	220-370	± 200
Ox_VOC15	2-methylpropene	1.15×10^{-17}	± 0.05	$2.92 \times 10^{-15} \exp(-1650/T)$	220-370	± 200
Ox_VOC41	2,3-dimethylbut-2-ene	1.1×10^{-15}	± 0.08	$3.0 \times 10^{-15} \exp(-300/T)$	220-370	± 200
Ox_VOC7	isoprene	1.28×10^{-17}	± 0.08	$1.05 \times 10^{-14} \exp(-2000/T)$	240-360	± 200
<i>Monoterpene reactions – based on data sheets in Supplement A Sect. A2</i>						
Ox_VOC8	α-pinene	9.6×10^{-17}	± 0.15	$8.22 \times 10^{-16} \exp(-640/T)$	240-370	± 300
Ox_VOC19	β-pinene	1.9×10^{-17}	± 0.25	$1.39 \times 10^{-15} \exp(-1280/T)$	290-370	± 300
Ox_VOC20	limonene	2.2×10^{-16}	± 0.1	$2.91 \times 10^{-15} \exp(-770/T)$	290-370	± 300
Ox_VOC21	camphene	5.0×10^{-19}	± 0.3	$9.0 \times 10^{-18} \exp(-860/T)$	285-315	± 500
Ox_VOC22	2-carene	2.4×10^{-16}	± 0.2			
Ox_VOC23	3-carene	4.9×10^{-17}	± 0.2			
Ox_VOC24	β-myrcene	4.7×10^{-16}	± 0.2	$2.69 \times 10^{-15} \exp(-520/T)$	290-320	± 300
Ox_VOC25	β-ocimene	5.1×10^{-16}	± 0.2	$4.15 \times 10^{-15} \exp(-625/T)$	290-320	± 300
Ox_VOC26	α-phellandrene	2.9×10^{-15}	± 0.2			
Ox_VOC27	β-phellandrene	5.2×10^{-17}	± 0.3			
Ox_VOC28	sabinene	8.3×10^{-17}	± 0.15			
Ox_VOC29	α-terpinene	1.9×10^{-14}	± 0.2			
Ox_VOC30	γ-terpinene	1.6×10^{-16}	± 0.3			
Ox_VOC31	terpinolene	1.6×10^{-15}	± 0.15			
<i>Sesquiterpene reactions – based on data sheets in Supplement A Sect. A3</i>						
Ox_VOC32	β-caryophyllene	1.2×10^{-14}	± 0.15			
Ox_VOC33	α-cedrene	no recommendation (see data sheet)				
Ox_VOC34	α-copaene	1.5×10^{-16}	± 0.3			
Ox_VOC35	α-farnesene	5.9×10^{-16}	± 0.3	$3.5 \times 10^{-12} \exp(-2590/T)$	290-320	± 500
Ox_VOC36	β-farnesene	5.6×10^{-16}	± 0.25	$1.5 \times 10^{-12} \exp(-2350/T)$	290-320	± 500
Ox_VOC37	α-humulene	1.2×10^{-14}	± 0.15			
Ox_VOC38	isolongifolene	1.0×10^{-17}	± 0.3			
Ox_VOC39	longifolene	$< 5 \times 10^{-19}$				
Ox_VOC40	valencene	no recommendation (see data sheet)				

Table 2. Summary of recommended total sCI yields (*Y*) from O₃ + alkene reactions at 298 K and 1 bar

Reaction ID ^a	alkene	<i>Y</i>	comments
<i>Small alkene reactions</i>			
Ox_VOC5	ethene	0.42 ± 0.10	(b)
Ox_VOC6	propene	0. 25 <u>30</u> ± 0.10	(c)
Ox_VOC16	but-1-ene	~ 0.27	(d)
Ox_VOC17	<i>cis</i> -but-2-ene	0.38 ± 0.10	(e)
Ox_VOC18	<i>trans</i> -but-2-ene	0.43 ± 0.10	(f)
Ox_VOC15	2-methylpropene	<u>0.21 ± 0.05</u> ~0.17	(g)
Ox_VOC41	2,3-dimethylbut-2-ene	0.38 ± 0.10	(h)
Ox_VOC7	isoprene	0.65 ± 0.10	(i)
<i>Monoterpene and sesquiterpene reactions</i>			
Ox_VOC8	α-pinene	0.18 ± 0.05	(j)
Ox_VOC19	β-pinene	0.5 <u>5</u> ± 0.1 <u>0</u>	(k)
Ox_VOC20	limonene	0. 32 <u>27</u> ± 0. 14 <u>10</u>	(l)
Ox_VOC21	camphene	~ 0.31	(d)
<u>Ox_VOC24</u>	<u>β-myrcene</u>	<u>0.46 ± 0.15</u>	<u>(m)</u>
Ox_VOC32	β-caryophyllene	> 0.6	(nn)
<i>Selected other reactions</i>			
-	cyclohexene	< 0.05	(oo)
-	<i>trans</i> -dec-5-ene	1.0	(pp)
-	<i>trans</i> -tetradec-7-ene	1.0	(qq)
Comments			
^a See corresponding data sheets in Supplement A for further information; ^b Based on Su et al. (1980), Kan et al. (1981), Hatekayama et al. (1984; 1986), Horie and Moortgat (1991), Neeb et al. (1996; 1998), Horie et al. (1999), Hasson et al. (2001a), Alam et al. (2011), Newland et al. (2015a; <u>2020</u>); ^c Based on Hatekayama et al. (1984), and Horie and Moortgat (1991) <u>and Newland et al. (2020)</u> ; ^d Based on Hasson et al. (2001b); ^e Based on Newland et al. (2015a); ^f Based on Berndt et al. (2014), Newland et al. (2015a; <u>2020</u>) and Hakala and Donahue (2018). Pressure dependence measurements suggest <i>Y</i> falls to ~25% <u>0.25</u> at 50 Torr (Hakala and Donahue, 2018); ^g Based on Hatekayama et al. (1986) <u>and Newland et al. (2020)</u> ; ^h Based on Berndt et al. (2014), Newland et al. (2015a) and Hakala and Donahue (2016). Pressure dependence measurements suggest <i>Y</i> falls to +20.12-45% <u>0.15</u> at zero pressure (e.g. Hakala and Donahue, 2016; Campos-Pineda and Zhang, 2017); ⁱ Based on Sipilä et al. (2014), Newland et al. (2015b) and Nguyen et al. (2016), as also discussed further in data sheet CGI_21 (Supplement B); ^j Based on Drozd and Donahue (2011), and Sipilä et al. (2014) <u>and Newland et al. (2018)</u> . Approximately linear pressure dependence observed by Drozd and Donahue (2011), with <i>Y</i> ≈ 0.05 at 110 Torr; ^k Based on Winterhalter et al. (2000) <u>and Newland et al. (2018)</u> with support from theoretical study of Nguyen et al. (2009a); ^l Based on Sipilä et al. (2014) <u>and Newland et al. (2018)</u> ; ^m <u>Based on Newland et al. (2020)</u> ; ⁿ Based on Winterhalter et al. (2009) with support from theoretical study of Nguyen et al. (2009b); ^o Based on Hatekayama et al. (1984), Drozd and Donahue (2011), who observed no stabilization at 550-640 Torr, and Campos-Pineda and Zhang (2018) who observed no stabilization at 10-20 Torr; ^p Based on Drozd and Donahue (2011). Full stabilisation observed at pressures above ~ 400 Torr, with <i>Y</i> falling at lower pressures to ~ 0.6 at 70 Torr; ^q Based on Hakala and Donahue (2018). Pressure dependence measurements suggest <i>Y</i> falls to ~ 0.35 at 50 Torr.			

Table 3. Summary of recommended HO yields for reactions of O₃ with alkenes at 298 K and 1 bar

Reaction ID ^a	alkene	HO yield	comments
<i>Small alkene reactions</i>			
Ox_VOC5	ethene	0.17 ± 0.05	(b)
Ox_VOC6	propene	0.36 ± 0.04	(c)
Ox_VOC16	but-1-ene	0.38 ± 0.18	(d)
Ox_VOC17	<i>cis</i> -but-2-ene	0.33 ± 0.07	(e)
Ox_VOC18	<i>trans</i> -but-2-ene	0.60 ± 0.06	(f)
Ox_VOC15	2-methylpropene	0.69 ± 0.15	(g)
Ox_VOC41	2,3-dimethylbut-2-ene	0.93 ± 0.14	(h)
Ox_VOC7	isoprene	0.26 ± 0.04	(i)
<i>Monoterpene reactions</i>			
Ox_VOC8	α-pinene	0.80 ± 0.10	(j)
Ox_VOC19	β-pinene	0.30 ± 0.06	(k)
Ox_VOC20	limonene	0.66 ± 0.04	(l)
Ox_VOC21	camphene	≤ 0.18	(m)
Ox_VOC22	2-carene	0.81 ± 0.11	(n)
Ox_VOC23	3-carene	0.86 ± 0.11	(n)
Ox_VOC24	β-myrcene	0.63 ± 0.09	(n)
Ox_VOC25	β-ocimene	0.55 ± 0.09	(n)
Ox_VOC26	α-phellandrene	0.29 ± 0.05	(o)
Ox_VOC27	β-phellandrene	0.14 ^{+0.07} _{-0.05}	(m)
Ox_VOC28	sabinene	0.33 ± 0.05	(n)
Ox_VOC29	α-terpinene	0.32 ± 0.06	(p)
Ox_VOC30	γ-terpinene	0.81 ± 0.11	(n)
Ox_VOC31	terpinolene	0.70 ± 0.08	(q)
<i>Sesquiterpene reactions</i>			
Ox_VOC32	β-caryophyllene	0.08 ± 0.03	(r)
Ox_VOC33	α-cedrene	0.65 ± 0.05	(s)
Ox_VOC34	α-copaene	0.35 ^{+0.18} _{-0.12}	(t)
Ox_VOC37	α-humulene	0.16 ± 0.06	(u)

Comments

^a See corresponding data sheets in Supplement A for further information; ^b Based on Atkinson et al. (1992), Paulson et al. (1999), Rickard et al. (1999), Mihelcic et al. (1999), Fenske et al. (2000) and Alam et al. (2011). Comparable pressure-independent yield (0.14) reported by Kroll et al. (2001a) over pressure range 13–80 mbar; ^c Based on Atkinson and Aschmann (1993), Neeb and Moortgat (1999), Paulson et al. (1999), Rickard et al. (1999), Aschmann et al. (2003), Qi et al. (2009) and Alam et al. (2013); ^d Based on Atkinson and Aschmann (1993), Paulson et al. (1999), Fenske et al. (2000) and Alam et al. (2013); ^e Based on Atkinson and Aschmann (1993), McGill et al. (1999), Orzechowska and Paulson (2002) and Alam et al. (2013); ^f Based on Atkinson and Aschmann (1993), McGill et al. (1999), Orzechowska and Paulson (2002), Hasson et al. (2003) and Alam et al. (2013); ^g Based on Atkinson and Aschmann (1993), Neeb and Moortgat (1999), Paulson et al. (1999), Rickard et al. (1999) and Alam et al. (2013); ^h Based on Chew and Atkinson (1996), Rickard et al. (1999), Fenske et al. (2000), Siese et al. (2001), Orzechowska and Paulson (2002), Aschmann et al. (2003) and Alam et al. (2013); ⁱ Based on Aschmann et al. (1996), Paulson et al. (1998), Neeb and Moortgat (1999), Malkin et al. (2010), Nguyen et al. (2016) and Ren et al. (2017); ^j Based on Atkinson et al. (1992), Chew and Atkinson (1996), Paulson et al. (1998), Rickard et al. (1999), Siese et al. (2001), Aschmann et al. (2002), Berndt et al. (2003), Presto and Donahue (2004) and Forester and Wells (2011); ^k Based on Atkinson et al. (1992) and Rickard et al. (1999); ^l Based on Aschmann et al. (2002), Herrmann et al. (2010) and Forester and Wells (2011); ^m Based on Atkinson et al. (1992); ⁿ Based on Aschmann et al. (2002); ^o Based on Herrmann et al. (2010); ^p Based on Aschmann et al. (2002) and Herrmann et al. (2010); ^q Based on Aschmann et al. (2002) and Herrmann et al. (2010); ^r Based on Shu and Atkinson (1994), Winterhalter et al. (2009) and Jenkin et al. (2012); ^s Based on Shu and Atkinson (1994) and Yao et al. (2014). Substantially lower yield, 0.090 ± 0.016, reported in the presence sCI scavengers, CH₃C(O)OH or SO₂, by Yao et al. (2014); ^t Based on Shu and Atkinson (1994); ^u Based on Shu and Atkinson (1994) and Beck et al. (2011).

Table 4. Summary of the recommended spectral data for C₁ - C₃ sCIs

Reaction ID ^a	reaction	σ_{max} cm ² molecule ⁻¹	$\Delta\sigma_{\text{max}}$ cm ² molecule ⁻¹	λ_{max} nm	λ range nm	ϕ^b
P33	CH ₂ OO + hν	1.23×10^{-17}	$\pm (0.18 \times 10^{-17})$	340	280-455	1.0
P34	Z-CH ₃ CHOO + hν	1.20×10^{-17}	$\pm (0.18 \times 10^{-17})$	323	300-430	1.0
	E-CH ₃ CHOO + hν	1.20×10^{-17}	$\pm (0.18 \times 10^{-17})$	360	300-430	1.0
P35	(CH ₃) ₂ COO + hν	1.75×10^{-17}	$\pm (0.53 \times 10^{-17})$	330	280-405	1.0
P36 ^c	CH ₃ CH ₂ CHOO + hν	no recommendation (see data sheet)		322	280-400	1.0

Comments

^a See corresponding data sheets in Supplement B Sect. B5 for further information.

^b ϕ is the photodissociation quantum yield.

^c Data sheet for CH₃CH₂CHOO included for completeness, although thermal reactions of this sCI are not included in the current evaluation.

Table 5. Summary of recommended rate coefficients for gas phase bimolecular reactions of sCIs

Reaction ID	Reaction	k_{298} $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	$\Delta \log k_{298}$	$k(T)$ $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	$T \text{ range}$	$\Delta(E/R)$ K
Reactions of CH_2OO – based on data sheets in Supplement B Sect. B1						
CGI_1	$\text{CH}_2\text{OO} + \text{SO}_2$	3.7×10^{-11}	± 0.05			
CGI_2	$\text{CH}_2\text{OO} + \text{NO}_2$	3×10^{-12}	± 0.5			
CGI_3	$\text{CH}_2\text{OO} + \text{NO}$	$< 6 \times 10^{-14}$				
CGI_4	$\text{CH}_2\text{OO} + \text{H}_2\text{O}$	2.8×10^{-16}	± 0.3			
	$\text{CH}_2\text{OO} + (\text{H}_2\text{O})_2$	6.4×10^{-12}	± 0.2	$7.35 \times 10^{-18} \exp(4076/T)$	280-325	± 500
CGI_5	$\text{CH}_2\text{OO} + \text{CH}_2\text{OO}$	7.4×10^{-11}	± 0.1			
CGI_6	$\text{CH}_2\text{OO} + \text{I}$	9.0×10^{-12}	± 0.3			
CGI_7	$\text{CH}_2\text{OO} + \text{CH}_3\text{C(O)H}$	$k_0 = 1.6 \times 10^{-29} [\text{M}]$ $k_\infty = 1.7 \times 10^{-12}$	± 0.2 ± 0.2			
CGI_8	$\text{CH}_2\text{OO} + \text{CH}_3\text{C(O)CH}_3$	3.4×10^{-13}	± 0.3			
CGI_9	$\text{CH}_2\text{OO} + \text{CF}_3\text{C(O)CF}_3$	3.2×10^{-11}	± 0.1			
CGI_11	$\text{CH}_2\text{OO} + \text{HC(O)OH}$	1.1×10^{-10}	± 0.1	<u>$1.52 \times 10^{-11} \exp(590/T)$</u>	<u>290-460</u>	<u>± 300</u>
CGI_10	$\text{CH}_2\text{OO} + \text{CH}_3\text{C(O)OH}$	1.3×10^{-10}	± 0.1			
CGI_23	$\text{CH}_2\text{OO} + \text{CF}_3\text{C(O)OH}$	3.3×10^{-10}	± 0.2	$3.8 \times 10^{-18} T^2 \exp(1620/T) + 2.5 \times 10^{-10}$	240-340	± 500
Reactions of Z- and E-CH_3CHOO – based on data sheets in Supplement B Sect. B2						
CGI_15	$\text{Z-CH}_3\text{CHOO} + \text{SO}_2$	2.6×10^{-11}	± 0.1			
	$\text{E-CH}_3\text{CHOO} + \text{SO}_2$	1.4×10^{-10}	± 0.3			
CGI_16	$\text{Z-CH}_3\text{CHOO} + \text{H}_2\text{O}$	$< 2 \times 10^{-16}$				
	$\text{E-CH}_3\text{CHOO} + \text{H}_2\text{O}$	1.3×10^{-14}	± 0.3			
	$\text{Z-CH}_3\text{CHOO} + (\text{H}_2\text{O})_2$	-				
	$\text{E-CH}_3\text{CHOO} + (\text{H}_2\text{O})_2$	4.4×10^{-11}	± 0.5			
CGI_17	$\text{Z-CH}_3\text{CHOO} + \text{NO}_2$	2.0×10^{-12}	± 0.15			
	$\text{E-CH}_3\text{CHOO} + \text{NO}_2$	2.0×10^{-12}	± 0.3			
CGI_26	$\text{Z-CH}_3\text{CHOO} + \text{HC(O)OH}$	2.5×10^{-10}	± 0.1			
	$\text{E-CH}_3\text{CHOO} + \text{HC(O)OH}$	5.0×10^{-10}	± 0.3			
CGI_27	$\text{Z-CH}_3\text{CHOO} + \text{CH}_3\text{C(O)OH}$	1.7×10^{-10}	± 0.15			
	$\text{E-CH}_3\text{CHOO} + \text{CH}_3\text{C(O)OH}$	2.5×10^{-10}	± 0.15			
Reactions of $(\text{CH}_3)_2\text{COO}$ – based on data sheets in Supplement B Sect. B3						
CGI_18	$(\text{CH}_3)_2\text{COO} + \text{SO}_2$	$k_\infty = 1.55 \times 10^{-10}$	± 0.15	$k_\infty = 4.23 \times 10^{-13} \exp(1760/T)$	280-305	± 500
CGI_19	$(\text{CH}_3)_2\text{COO} + \text{H}_2\text{O}$	$< 1.5 \times 10^{-16}$				
	$(\text{CH}_3)_2\text{COO} + (\text{H}_2\text{O})_2$	$< 1.3 \times 10^{-13}$				
CGI_20	$(\text{CH}_3)_2\text{COO} + \text{NO}_2$	2.1×10^{-12}	± 0.3			
CGI_28	$(\text{CH}_3)_2\text{COO} + \text{HC(O)OH}$	3.1×10^{-10}	± 0.1			
CGI_29	$(\text{CH}_3)_2\text{COO} + \text{CH}_3\text{C(O)OH}$	3.1×10^{-10}	± 0.1			
CGI_24	$(\text{CH}_3)_2\text{COO} + \text{CF}_3\text{C(O)OH}$	6.2×10^{-10}	± 0.2	$4.9 \times 10^{-18} T^2 \exp(1620/T) + 5.2 \times 10^{-10}$	240-340	± 500

Table 5 (continued). Summary of recommended rate coefficients for gas phase bimolecular reactions of sCIs

Reaction ID	Reaction	k_{298} cm ³ molecule ⁻¹ s ⁻¹	$\Delta \log k_{298}$	$k(T)$ cm ³ molecule ⁻¹ s ⁻¹	<i>T</i> range	$\Delta(E/R)$ K
<i>Reactions of C₄ intermediates from isoprene – based on data sheets in Supplement B Sect. B4</i>						
CGI_21 ^a	Z-(CH=CH ₂)(CH ₃)COO + SO ₂	1.554.2 × 10 ^{-10.11}	-	4.23 × 10⁻¹³ exp(1760/T)	-	-
	Z-(CH=CH ₂)(CH ₃)COO + H ₂ O	1.79 × 10 ⁻¹⁸	-	2.21 × 10 ⁻²¹ T ^{2.27} exp(-1858/T)	-	-
	Z-(CH=CH ₂)(CH ₃)COO + (H ₂ O) ₂	4.87 × 10 ⁻¹⁵	-	2.25 × 10 ⁻²¹ T ^{2.27} exp(493/T)	-	-
	<i>E</i> -(CH=CH ₂)(CH ₃)COO + SO ₂	see data sheet CGI_22				
	<i>E</i> -(CH=CH ₂)(CH ₃)COO + H ₂ O	7.89 × 10 ⁻²⁰	-	7.07 × 10 ⁻¹⁹ T ^{1.46} exp(-3132/T)	-	-
	<i>E</i> -(CH=CH ₂)(CH ₃)COO + (H ₂ O) ₂	3.06 × 10 ⁻¹⁶	-	7.63 × 10 ⁻¹⁹ T ^{1.45} exp(-675/T)	-	-
	Z-(C(CH ₃)=CH ₂)CHOO + SO ₂	2.6 × 10 ⁻¹¹	-		-	-
	Z-(C(CH ₃)=CH ₂)CHOO + H ₂ O	1.19 × 10 ⁻¹⁹	-	2.24 × 10 ⁻¹⁹ T ^{1.65} exp(-2989/T)	-	-
	Z-(C(CH ₃)=CH ₂)CHOO + (H ₂ O) ₂	4.39 × 10 ⁻¹⁶	-	2.42 × 10 ⁻¹⁹ T ^{1.64} exp(-548/T)	-	-
	<i>E</i> -(C(CH ₃)=CH ₂)CHOO + SO ₂	1.4 × 10 ⁻¹⁰	-		-	-
<u>CGI_22</u>	<u><i>E</i>-(CH=CH₂)(CH₃)COO + SO₂</u>	<u>4.2 × 10⁻¹¹</u>	<u>± 0.2</u>			
	<u><i>E</i>-(CH=CH₂)(CH₃)COO + HC(O)OH</u>	<u>3.1 × 10⁻¹⁰</u>	<u>± 0.3</u>			
<u>CGI_30</u>	<u><i>E</i>-(CH=CH₂)(CH₃)COO + HC(O)OH</u>	<u>3.1 × 10⁻¹⁰</u>	<u>± 0.3</u>			
	<u><i>E</i>-(CH=CH₂)(CH₃)COO + HC(O)OH</u>	<u>3.1 × 10⁻¹⁰</u>	<u>± 0.3</u>			
CGI_25	<i>E</i> -(CH=CH ₂)(CH ₃)COO + CF ₃ C(O)OH	7.3 × 10 ⁻¹⁰	± 0.3	4.9 × 10 ⁻¹⁸ T ² exp(1620/T) + 6.3 × 10 ⁻¹⁰	240 -340	± 500
	<i>E</i> -(C(CH ₃)=CH ₂)CHOO + CF ₃ C(O)OH	7.3 × 10 ⁻¹⁰	± 0.3	4.9 × 10 ⁻¹⁸ T ² exp(1620/T) + 6.3 × 10 ⁻¹⁰	240 -340	± 500

Comments: ^a Rate coefficients for SO₂ reactions with Z-(CH=CH₂)(CH₃)COO, Z-(C(CH₃)=CH₂)CHOO and *E*-(C(CH₃)=CH₂)CHOO are inferred from those for *E*-(CH=CH₂)(CH₃)COO, Z-CH₃CHOO and *E*-CH₃CHOO, respectively. Temperature-dependent rate coefficients for H₂O and (H₂O)₂ reactions are adopted from the theoretical/SAR methods reported by Vereecken et al. (2017), as presented in Supplement Tables 35 and 40 of that paper. Individual parameters are not currently assigned uncertainties, but performance of ensemble of reactions (also including sCI decomposition reactions) was tested against reported O₃ + isoprene product observations (see data sheet CGI_21, Supplement B).

Table 6. Summary of recommended rate coefficients for gas phase unimolecular reactions of sCIs

Reaction ID	Reaction	k_{298} s ⁻¹	$\Delta \log k_{298}$	$k(T)$ s ⁻¹	T range	$\Delta(E/R)$ K
<i>Based on data sheets in Supplement B Sects. B1-B4</i>						
CGI_12	CH ₂ OO + M	<u>≤ 0.2</u> (1 bar)	<u>± 1.0</u>	$k_0 = 3.2 \times 10^{-4} (T/298)^{-5.81} \exp(-12770/T) [M]$ $k_{\infty} = 1.4 \times 10^{13} (T/298)^{0.06} \exp(-10010/T)$ ($E_c = 0.447$)	450-650 450-650	± 500 ± 500
CGI_13	Z-CH ₃ CHOO + M E-CH ₃ CHOO + M	150 (1 bar) 60 (1 bar)	± 0.3 ± 0.5	$7.4 \times 10^6 \exp(-3220/T)$	275-320	± 700
CGI_14	(CH ₃) ₂ COO + M	400 (1 bar)	± 0.2	$7.2 \times 10^6 \exp(-2920/T)$	280-330	± 700
CGI_21 ^a	Z-(CH=CH ₂)(CH ₃)COO + M	13,600	-	$9.75 \times 10^8 T^{1.03} \exp(-5081/T)$	-	-
	E-(CH=CH ₂)(CH ₃)COO + M	51.3	-	$4.36 \times 10^{-67} T^{25.9} \exp(2737/T)$	-	-
	Z-(C(CH ₃)=CH ₂)CHOO + M	14,000	-	$2.58 \times 10^9 T^{0.87} \exp(-5090/T)$	-	-
	E-(C(CH ₃)=CH ₂)CHOO + M	30.2	-	$1.68 \times 10^{10} T^{1.02} \exp(-7732/T)$	-	-
Comments: ^a Temperature-dependent rate coefficients adopted from the theoretical/SAR methods reported by Vereecken et al. (2017), as presented in Supplement Table 31 of that paper (N.B. exponent of the pre-exponential factor changed from 9 to 8 in the case of Z-(CH=CH ₂)(CH ₃)COO, for consistency with 298 K rate coefficient reported by Vereecken et al. (2017)). Individual parameters are not currently assigned uncertainties, but performance of the ensemble of reactions (also including sCI reactions with SO ₂ , H ₂ O and (H ₂ O) ₂) was tested against reported O ₃ + isoprene product observations (see data sheet CGI_21, Supplement B).						

Table 7. Rate coefficients (k) and representative lower tropospheric first order loss rates (k_1) at 298 K for sCI bimolecular reactions with SO₂, H₂O and (H₂O)₂, and unimolecular decomposition. The present recommendations (IUPAC) are compared with calculated values using the theory-based SAR developed by Vereecken et al. (2017) (except where noted).

Reaction	k^a		$k_1 \text{ (s}^{-1}\text{)}^b$	
	IUPAC	SAR ^c	IUPAC	SAR ^c
Reactions of CH₂OO				
CH ₂ OO + SO ₂	3.7×10^{-11}	-	0.93	-
CH ₂ OO + H ₂ O	2.8×10^{-16}	8.63×10^{-16}	86	266
CH ₂ OO + (H ₂ O) ₂	6.4×10^{-12}	1.48×10^{-12}	1250	289
CH ₂ OO + M	<u>$< 2 \times 10^{-1} \text{ s}^{-1}$</u>	<u>$7.2 \times 10^{-2} \text{ s}^{-1}$</u> ^d	<u>< 0.2</u>	<u>0.072</u> ^d
Reactions of Z-CH₃CHOO				
Z-CH ₃ CHOO + SO ₂	2.6×10^{-11}	-	0.65	-
Z-CH ₃ CHOO + H ₂ O	$< 2 \times 10^{-16}$	6.84×10^{-19}	< 62	0.210
Z-CH ₃ CHOO + (H ₂ O) ₂	-	2.05×10^{-15}	-	0.401
Z-CH ₃ CHOO + M	$1.5 \times 10^2 \text{ s}^{-1}$	$1.37 \times 10^2 \text{ s}^{-1}$	150	137
Reactions of E-CH₃CHOO				
E-CH ₃ CHOO + SO ₂	1.4×10^{-10}	-	3.5	-
E-CH ₃ CHOO + H ₂ O	1.3×10^{-14}	2.33×10^{-14}	4000	7190
E-CH ₃ CHOO + (H ₂ O) ₂	4.4×10^{-11}	2.63×10^{-11}	8600	5150
E-CH ₃ CHOO + M	$6.0 \times 10^1 \text{ s}^{-1}$	$5.22 \times 10^1 \text{ s}^{-1}$	60	52.2
Reactions of (CH₃)₂COO				
(CH ₃) ₂ COO + SO ₂	1.55×10^{-10}	-	3.9	-
(CH ₃) ₂ COO + H ₂ O	$< 1.5 \times 10^{-16}$	7.40×10^{-18}	< 46	2.28
(CH ₃) ₂ COO + (H ₂ O) ₂	$< 1.3 \times 10^{-13}$	1.79×10^{-14}	< 25	3.51
(CH ₃) ₂ COO + M	$4.0 \times 10^2 \text{ s}^{-1}$	$5.01 \times 10^2 \text{ s}^{-1}$	400	501
Reactions of C₄ intermediates from isoprene				
Z-(CH=CH ₂)(CH ₃) <u>C</u> OO + SO ₂	<u>4.2×10^{-11}</u> ^{de}	-	<u>1.1</u>	-
Z-(CH=CH ₂)(CH ₃) <u>C</u> OO + H ₂ O	<u>fe</u>	1.79×10^{-18}	<u>ef</u>	0.551
Z-(CH=CH ₂)(CH ₃) <u>C</u> OO + (H ₂ O) ₂	<u>fe</u>	4.87×10^{-15}	<u>fe</u>	0.951
Z-(CH=CH ₂)(CH ₃) <u>C</u> OO + M	<u>fe</u>	$1.36 \times 10^4 \text{ s}^{-1}$	<u>fe</u>	13600
E-(CH=CH ₂)(CH ₃) <u>C</u> OO + SO ₂	<u>4.2×10^{-11}</u>	-	<u>1.1</u>	-
E-(CH=CH ₂)(CH ₃) <u>C</u> OO + H ₂ O	<u>fe</u>	7.89×10^{-20}	<u>fe</u>	0.0243
E-(CH=CH ₂)(CH ₃) <u>C</u> OO + (H ₂ O) ₂	<u>fe</u>	3.06×10^{-16}	<u>fe</u>	0.0599
E-(CH=CH ₂)(CH ₃) <u>C</u> OO + M	<u>fe</u>	$5.13 \times 10^1 \text{ s}^{-1}$	<u>fe</u>	51.3
Z-(C(CH ₃)=CH ₂)CHOO + SO ₂	2.6×10^{-11} ^{dg}	-	0.65	-
Z-(C(CH ₃)=CH ₂)CHOO + (H ₂ O) ₂	<u>fe</u>	1.19×10^{-19}	<u>fe</u>	0.0367
Z-(C(CH ₃)=CH ₂)CHOO + SO ₂	<u>fe</u>	4.39×10^{-16}	<u>fe</u>	0.0859
Z-(C(CH ₃)=CH ₂)CHOO + M	<u>fe</u>	$1.40 \times 10^4 \text{ s}^{-1}$	<u>fe</u>	14000
E-(C(CH ₃)=CH ₂)CHOO + SO ₂	1.4×10^{-10} ^{dg}	-	3.5	-
E-(C(CH ₃)=CH ₂)CHOO + (H ₂ O) ₂	<u>fe</u>	1.43×10^{-16}	<u>fe</u>	44.1
E-(C(CH ₃)=CH ₂)CHOO + SO ₂	<u>fe</u>	2.79×10^{-13}	<u>fe</u>	54.5
E-(C(CH ₃)=CH ₂)CHOO + M	<u>fe</u>	$3.02 \times 10^1 \text{ s}^{-1}$	<u>fe</u>	30.2
Comments: ^a Units of k are cm ³ molecule ⁻¹ s ⁻¹ , unless otherwise stated; ^b k_1 at 298 K and 1 bar determined for 40 % relative humidity ([H ₂ O] = 3.08×10^{17} molecule cm ⁻³ and [(H ₂ O) ₂] = 1.96×10^{14} molecule cm ⁻³), and for [SO ₂] = 2.5×10^{10} molecule cm ⁻³ (~1 ppbv); ^c <u>Except where noted, Based on high-pressure limiting T-dependence parameters given in Tables 31, 35 and 40 of Vereecken et al. (2017);</u> ^d <u>Based on the value at 1 bar reported by Long et al. (2016);</u> ^e <u>Value inferred from that for E-(CH=CH₂)(CH₃)COO;</u> ^f <u>IUPAC k value is adopted from Vereecken et al. (2017), and given entry is therefore identical to the SAR value shown;</u> ^g <u>Values for Z- and E-(C(CH₃)=CH₂)CHOO inferred from those for Z and E-CH₃CHOO, respectively.</u>				

Table 8. Summary of results for the representative ambient calculations (see Sect. 8 and Supplement C for further details).

	Rural background ^a		Suburban background ^b		Urban traffic ^c	
	winter	summer	winter	summer	winter	summer
sCI production rate (molecule cm ⁻³ s ⁻¹)	6.34 ×10 ⁴	8.70 ×10 ⁴	8.77 ×10 ⁴	2.42 ×10 ⁵	6.67 ×10 ⁴	2.17 ×10 ⁵
sCI concentration (molecule cm ⁻³)	386	375	536	1075	379	879
SO ₂ oxidation rate (% h ⁻¹)	0.006	0.007	0.008	0.028	0.006	0.009
HC(O)OH oxidation rate (% h ⁻¹)	0.04	0.04	0.05	0.11	0.03	0.08
<i>Contributions to total sCI loss</i>						
Unimolecular decomposition	45.7%	49.3%	46.7%	57.7%	43.8%	49.0%
Reaction with H ₂ O	9.3%	9.6%	8.9%	8.6%	8.4%	9.3%
Reaction with (H ₂ O) ₂	44.4%	38.2%	42.2%	29.1%	43.0%	37.0%
Reaction with SO ₂	0.3%	0.2%	1.3%	1.6%	1.2%	0.6%
Reaction with NO ₂	0.2%	0.1%	0.4%	0.1%	1.1%	0.6%
Reaction with HC(O)OH	0.2%	2.6%	0.5%	3.0%	2.4%	3.4%
<i>Contributions to sCI production rate</i>						
Core C ₁ -C ₄ set ^c	91.1%	75.9%	88.6%	44.7%	88.0%	85.6%
Others C ₃ -C ₅ (anthropogenic) ^d	8.9%	7.2%	11.4%	4.8%	12.0%	7.4%
Others C ₁₀ (biogenic) ^e	-	16.9%	-	50.5%	-	7.0%
<i>Contributions to sCI concentration</i>						
Core C ₁ -C ₄ set ^c	89.3%	68.2%	86.6%	35.4%	85.6%	83.4%
Others C ₃ -C ₅ (anthropogenic) ^d	10.7%	7.6%	13.4%	4.5%	14.4%	7.8%
Others C ₁₀ (biogenic) ^e	-	24.2%	-	60.1%	-	8.8%
<i>Contributions to SO₂ oxidation rate ^f</i>						
Core C ₁ -C ₄ set ^c	91.2%	38.5%	90.7%	13.6%	90.8%	63.6%
Others C ₃ -C ₅ (anthropogenic) ^d	8.8%	3.7%	9.3%	1.2%	9.2%	4.7%
Others C ₁₀ (biogenic) ^e	-	57.8%	-	85.2%	-	31.7%
<i>Contributions to HC(O)OH oxidation rate ^f</i>						
Core C ₁ -C ₄ set ^c	88.7%	65.1%	86.5%	32.7%	85.7%	82.0%
Others C ₃ -C ₅ (anthropogenic) ^d	11.3%	7.6%	13.5%	4.1%	14.3%	7.6%
Others C ₁₀ (biogenic) ^e	-	27.3%	-	63.2%	-	10.4%
Comments: ^a Based on data for the Chilbolton Observatory site (51.149617, -1.438228); ^b Based on data for the London Eltham site (51.452580, 0.070766); ^c Based on data for the London Marylebone Road site (51.522530, -0.154611); ^d The core C ₁ -C ₄ set comprises the sCIs shown in Fig. 2, for which evaluated rate parameters have been presented in Sect. 6; ^e Comprises <i>E</i> - and <i>Z</i> -C ₂ H ₅ CHOO, <i>E</i> - and <i>Z</i> - <i>n</i> -C ₃ H ₇ CHOO, <i>E</i> - and <i>Z</i> - <i>i</i> -C ₃ H ₇ CHOO, <i>E</i> - and <i>Z</i> - <i>n</i> -C ₄ H ₉ CHOO, <i>E</i> - and <i>Z</i> -(C ₂ H ₅)(CH ₃)COO, <i>E</i> - and <i>Z</i> -(CH=CH ₂)CHOO formed from the C ₄ -C ₆ alkenes considered (see Supplement C); ^f Comprises eight pinonaldehyde oxide and limononaldehyde oxide isomers formed from the monoterpenes α -pinene and limonene (see Supplement C); ^f Specifies the contribution to the total SO ₂ or HC(O)OH oxidation rate due to reaction with sCIs.						