Supplementary material for "From a polar to a marine environment: has the changing Arctic led to a shift in aerosol light scattering properties?" by Heslin-Rees et al. (2020)

Contents

5	1	Data availability	2
	2	Particle light scattering coefficients compared to ambient relative humidity	3
	3	Back trajectory calculations (method)	5
	4	The contribution of coarse mode particles to particle light scattering	6
	5	Results of trajectory analysis with a fixed sea ice	7
10	6	Annual air mass contribution	8
	7	The annual contribution of the scattering Ångström exponent with respect to the main air mass sectors	9
	8	Table of results	10

1 Data availability



Figure S1 shows the available data of nephelometer measurements used in this study.

Figure S1. Data coverage: Grey bars denotes the number of hourly data points provided by the nephelometer before the quality control procedure. The green bars denote the number of hourly data points used.

15 2 Particle light scattering coefficients compared to ambient relative humidity

Within this work, the effect of local cloud occurrence is explicitly addressed in the trend analysis. As a proxy for in-cloud data, the ambient relative humidity (RH) (see Fig. S2) is being used. Figure S2a shows that there is a slight influence from the ambient relative humidity on the RH at the sample outlet of the nephelometer. From Fig. S2b it is shown that increasing ambient RH results in increased σ_{sp} up to a point, after which the high levels of ambient RH associated with the presence of clouds begin

- to reduce the inlet collection efficiency. The point at which inlet efficiency is negatively impacted by the presence of clouds is approximately 95 %. The overall effect is deemed insignificant. Figure S2c shows that *alpha* declines with increasing ambient RH, suggesting that the aerosol particles take up water and thus increase in size despite the nephelometer measuring under dry conditions (i.e. sample outlet RH < 40%). After approximately 93% ambient RH, there is a slight upturn in α suggesting that the presence of clouds at the inlet acts to remove a proportion of larger particles. However, the median α for cloud-free conditions
- 25 (i.e. <95%) is larger suggesting in fact that the size distribution is shifted closer to the smaller particles; this observation is put down to the increasing uncertainties observed in the data.



Figure S2. Effect of ambient relative humidity (RH): (a) The RH at the sample outlet vs. the ambient RH. (b) The scattering coefficient (σ_{sp}) vs. the ambient (RH). (c) The Ångström exponent (α) vs. the ambient (RH). The data is placed in bins of size 200 according to ascending values of ambient RH. The medians for each consecutive bin are calculated for the variables. The dotted red line (at ambient RH=95%) displays the cut-off point, whereby in-cloud scavenging and inlet losses, influence σ_{sp} significantly. (RH threshold of 92% perhaps)

3 Back trajectory calculations (method)

Figure S3 presents a detailed description of how the residence times over the three different surface types (land, ice and open water) are calculated. Note that the proportion of time the back trajectories spend above the mixed layer is also presented, and furthermore does not count towards the surface residence times.



Figure S3. Example of the source attribution using back trajectory calculations: 168-h back trajectory arriving at the Zeppelin Observatory on 2008-10-05 at 13:00:00. Sections of the back trajectory are coloured as follows: above the mixed layer (purple), and within the mixed layer and above land (green), open water (blue), ice (white). The mean coordinates of the back trajectory, based only on the section that traverses through the ML, is denoted by the star. The red cross displays the location of the Zeppelin Observatory. The sea ice concentrations (SIC) at the time of arrival are displayed, based on monthly 1° x 1°-resolution satellite-derived Met Office (2006) data. The meridian 11°89'E and parallel of 78°91'N lines are presented for reference.

4 The contribution of coarse mode particles to particle light scattering

Coarse-mode particles contribute a significant proportion of the overall scattering observed at ZEP. To illustrate the importance of the coarse mode (here defined for particles above 1 μm), we have used particle size distribution measurements of fine and coarse mode from 2008 and Mie theory assuming dry spherical particles with a refractive index of m = 1.51 and a wavelength
of λ = 550 nm (details on size distribution measurements and Mie code are given in Zieger et al., 2010). Figure S4 shows the mean monthly average of the particle surface size distribution and the differential scattering coefficient (the integral will deliver σ_{sp}) for July to October. As can be seen, in 2008 the coarse mode contributed on average between 20 and 45 % (at λ = 550 nm) to the particle light scattering and as such is an important contributor to overall light scattering properties. In Zieger et al. (2010), the coarse mode was identified as sea salt using chemical and trajectory analysis.



Figure S4. Monthly averages of (a) the particle surface size distribution and (b) the differential scattering coefficient (at $\lambda = 550$ nm) calculated using the SMPS (scanning mobility particle sizer) and OPSS (optical particle size spectrometer) measurements from Zieger et al. (2010). The percent contribution to light scattering of the coarse mode (CM) is given in the legend of panel b.

40 5 Results of trajectory analysis with a fixed sea ice

45

Figure S5 shows the relative surface contribution of the air masses arriving at ZEP. To investigate the effect of retreating Arctic sea ice on the surface type contributions alone, the contributions are calculated by holding the mean SIC at the annual cycle for 1999, The solid line denotes the surface type contributions based on the monthly sea ice concentrations (SIC). The dashed line is the SIC fixed to the values for the year of 1999. It can be seen that the relative contribution from different surface types does not change significantly between the dashed and solid lines. This suggests that the retreat in sea ice is not the main contributor to the increase spend the back trajectories spend over the open water.



Figure S5. Relative surface contributions: the solid line represents the relative seasonal presence of each different surface type. The dashed line presents the same information but with sea ice concentrations set at 1999 values. Note the contributions do not include parts of the back trajectories above the mixed-layer (thus the sum is < 1).

6 Annual air mass contribution

The increased presence of south-west air masses in the summer is displayed in Fig. S6. The presence of SW air masses in the summer is consistent with previous studies (Freud et al., 2017; Dall'Osto et al., 2017). The dominant flow from the north during the winter and spring months is well documented by Stone et al. (2014).



Figure S6. Annual variation of the monthly regional air mass contributions determined from the trajectory analysis for the entire observation period (1999-2016).

7 The annual contribution of the scattering Ångström exponent with respect to the main air mass sectors

55

In Figure S7, the hourly median α values are presented for each region, and for every year of observation, as a set of normalised histograms. The shift in air mass contributions is displayed in the relative histogram areas (A). The same years that feature peaks in their open water contributions (i.e. 2006, 2011, and 2016 in Fig S5) display decreases in their respective α medians. α for the western air masses are distinctly lower, compared to the eastern originated air masses. The western air masses are also more bimodal, suggesting the presence of more coarse-mode particles. It needs to be noted that the number of hourly data points for each season is not the same; spring and winter comprise of much more of the data (Tables S1 and S4) and thus annual medians are weighted more in their favour.



Figure S7. Annual histograms for hourly scattering Ångström exponent (α) medians of the four source regions namely, North-east (NE), South-east (SE), North-west (NW) and South-west (SW). The histograms are normalised by the total of each respective year. The respective α medians for each region are given, along with the region's relative contribution (A (%)). The area left of the vertical dashed line, where ($\alpha < 1$), represents the approximate contribution from coarse-mode aerosol particles. Note the scale for the year 1999 is different.

8 Table of results

60 Tables S1, S2, S3 and S4 present the seasonal trends in the aerosol optical properties for both statistical tests (i.e. least mean square and seasonal Mann-Kendall test). The median for each season is also given, along with the number of hourly data points comprised of the points used in the trend analysis.

Table S1. Trends in scattering coefficient (σ_{sp}) for daily medians and seasonal medians: Theil-Sen Slope estimator (TS) is presented as the change in Mm⁻¹ per year and the relative change (% yr⁻¹). Least Mean Squares (LMS) is the percentage change based on a log transformation. For daily medians, the Man-Kendall test (MK test) is based on a "prewhitened" time series; all seasons denotes the result of the seasonal MK test. Decreasing (D) and increasing (I) statistically significant trends are signified.

	Period	Hourly points	Data points	$\sigma_{ m sp}$ (Mm ⁻¹)	$\frac{\mathbf{TS}}{(\mathbf{Mm}^{-1}\mathbf{yr}^{-1})}$	$\frac{\text{TS}}{(\% \text{yr}^{-1})}$	$\frac{LMS}{(\% yr^{-1})}$	МК	LMS
Daily	spring	14930	792	3.38	0.04	1.24	1.52	Ι	Ι
	summer	6875	470	1.11	0.03	2.66	1.81	Ι	Ι
	autumn	10137	630	1.49	0.07	4.44	3.50	Ι	Ι
	winter	13411	690	3.10	0.19	6.03	6.10	Ι	Ι
	total	45353	2582	2.30	0.06	2.43	2.90	Ι	Ι
Seasonal	spring	14930	14	3.40	0.01	0.22	-0.24	-	-
	summer	6875	15	1.09	0.02	2.06	0.76	-	-
	autumn	10137	12	1.64	0.09	5.24	3.74	Ι	Ι
	winter	13411	11	3.24	0.21	6.50	4.60	Ι	Ι
	total	45353	52	2.09	0.05	2.21	1.27	Ι	-

Table S2. Trends in backscattering coefficient (σ_{bsp}) for daily medians and seasonal medians: Theil-Sen Slope estimator (TS) is presented as the change in Mm⁻¹ per year and the relative change (% yr⁻¹). Least Mean Squares (LMS) is the percentage change based on a log transformation. For daily medians, the Man-Kendall test (MK test) is based on a "prewhitened" time series; all seasons denotes the result of the seasonal MK test. Decreasing (D) and increasing (I) statistically significant trends are signified.

	Period	Hourly points	Data points	$\sigma_{\mathrm{bsp}} \ (\mathrm{Mm}^{-1})$	$\frac{\text{TS}}{(\text{Mm}^{-1}\text{yr}^{-1})}$	$\frac{\text{TS}}{(\% \text{yr}^{-1})}$	$\frac{LMS}{(\% yr^{-1})}$	MK	LMS
Daily	spring	14704	791	0.44	0.01	1.32	0.34	Ι	-
	summer	6063	469	0.18	0.00	2.01	0.52	Ι	Ι
	autumn	9054	629	0.21	0.01	3.32	0.85	Ι	Ι
	winter	12781	691	0.39	0.02	5.27	1.86	Ι	Ι
	total	42602	2580	0.30	0.01	2.14	0.62	Ι	Ι
Seasonal	spring	14704	14	0.44	0.00	0.73	-0.01	-	-
	summer	6063	15	0.19	0.00	2.53	0.46	Ι	Ι
	autumn	9054	12	0.26	0.01	3.93	0.87	Ι	Ι
	winter	12781	11	0.42	0.02	5.78	1.44	Ι	-
	total	42602	52	0.30	0.01	2.30	0.43	Ι	-

Table S3. Trend in hemispheric backscattering fraction b (-) for daily medians and seasonal medians: Theil-Sen Slope estimator (TS) is presented as the change per year and the relative change (% yr⁻¹). Least Mean Squares (LMS) is the relative change based on a log transformation. For daily medians, the Man-Kendall test (MK test) is based on a "prewhitened" time series; all seasons denotes the result of the seasonal MK test. Decreasing (D) and increasing (I) statistically significant trends are signified.

	Period	Hourly points	Data points	b (-)	TS (yr ⁻¹)	$\frac{\text{TS}}{(\% \text{yr}^{-1})}$	$\frac{LMS}{(\% yr^{-1})}$	МК	LMS
Daily	spring	14490	791	0.1296	0.0000	0.0320	-0.0140	-	-
	summer	5654	467	0.1490	-0.0008	-0.5194	-0.0820	D	D
	autumn	8508	622	0.1323	-0.0012	-0.9020	-0.1540	D	D
	winter	12515	690	0.1206	-0.0009	-0.7734	-0.1470	D	D
	total	41167	2570	0.1301	-0.0004	-0.3249	-0.0710	D	D
Seasonal	spring	14490	14	0.1276	0.0009	0.7199	0.0610	-	-
	summer	5654	15	0.1464	-0.0005	-0.3575	-0.0620	-	-
	autumn	8508	12	0.1319	-0.0007	-0.5545	-0.1110	-	-
	winter	12515	11	0.1196	-0.0001	-0.0739	-0.1110	-	-
	total	41167	52	0.1318	-0.0002	-0.1674	-0.0400	-	-

Table S4. Trend in scattering Ångström exponent (-) for daily medians and seasonal medians: Theil-Sen Slope estimator (TS) is presented as the change per year and the relative change (% yr⁻¹). Least Mean Squares (LMS) is the percentage change based on a log transformation. For daily medians, the Man-Kendall test (MK test) is based on a "prewhitened" time series; all seasons denotes the result of the seasonal MK test. Decreasing (D) and increasing (I) statistically significant trends are signified.

	Period	Hourly points	Data points	α (-)	$\frac{\mathbf{TS}}{(\mathbf{yr}^{-1})}$	$\frac{\text{TS}}{(\% \text{yr}^{-1})}$	$\frac{LMS}{(\% yr^{-1})}$	MK	LMS
Daily	spring	14930	792	1.59	-0.06	-3.51	-4.89	D	D
	summer	6875	470	1.68	-0.07	-4.42	-4.91	D	D
	autumn	10137	630	1.17	-0.09	-7.29	-7.61	D	D
	winter	13411	690	1.30	-0.08	-5.84	-7.31	D	D
	total	45353	2582	1.44	-0.07	-4.92	-6.42	D	D
Seasonal	spring	14930	14	1.51	-0.05	-3.10	-1.58	D	D
	summer	6875	15	1.74	-0.07	-3.95	-2.56	D	D
	autumn	10137	12	1.20	-0.10	-8.68	-3.54	D	D
	winter	13411	11	1.22	-0.06	-4.86	-2.99	D	D
	total	45353	52	1.36	-0.07	-5.05	-2.65	D	D

References

- Dall'Osto, M., Beddows, D., Tunved, P., Krejci, R., Ström, J., Hansson, H.-C., Yoon, Y., Park, K.-T., Becagli, S., Udisti, R., et al.: Arctic sea
 ice melt leads to atmospheric new particle formation, Sci. Rep., 7, 3318, 2017.
 - Freud, E., Krejci, R., Tunved, P., Leaitch, R., Nguyen, Q. T., Massling, A., Skov, H., and Barrie, L.: Pan-Arctic aerosol number size distributions: seasonality and transport patterns, Atmos. Chem. Phys., 17, 8101–8128, 2017.
 - Met Office, H. C.: HadISST 1.1 Global sea-Ice coverage and SST (1870-Present), http://badc.nerc.ac.uk/view/badc.nerc.ac.uk, accessed: 2019-04-30, 2006.
- 70 Stone, R. S., Sharma, S., Herber, A., Eleftheriadis, K., and Nelson, D. W.: A characterization of Arctic aerosols on the basis of aerosol optical depth and black carbon measurements, Elementa., 2014.
 - Zieger, P., Fierz-Schmidhauser, R., Gysel, M., Ström, J., Henne, S., Yttri, K. E., Baltensperger, U., and Weingartner, E.: Effects of relative humidity on aerosol light scattering in the Arctic, Atmos. Chem. Phys., 10, 3875–3890, 2010.