

Supporting Information for

Tracing the evolution of morphology and mixing state of soot particles along with the movement of an Asian dust storm

Liang Xu¹, Satoshi Fukushima², Sophie Sobanska³, Kotaro Murata², Ayumi Naganuma², Lei Liu¹,
Yuanyuan Wang¹, Hongya Niu⁴, Zongbo Shi⁵, Tomoko Kojima⁶, Daizhou Zhang², Weijun Li^{1,*}

¹Department of Atmospheric Sciences, School of Earth Sciences, Zhejiang University, Hangzhou
310027, China

²Faculty of Environmental and Symbiotic Sciences, Prefectural University of Kumamoto,
Kumamoto 862-8502, Japan

³Institute of Molecular Sciences, UMR CNRS 5255, University of Bordeaux, 351 cours de la
libération, 33405 Talence, France

⁴Key Laboratory of Resource Exploration Research of Hebei Province, Hebei University of
Engineering, Handan 056038, Hebei, China

⁵School of Geography, Earth and Environmental Sciences, University of Birmingham, UK

⁶Department Earth and Environmental Science, Faculty of Advanced Science and Technology,
Kumamoto University, Kumamoto 860-8555, Japan

Contents of this file

Table S1, Figures S1 to S6

1. Information about samples and the Asian dust storm event

Table S1. Detailed information about sampling dates, times, meteorological conditions, and PM concentrations for samples.

	Sample ID	Date	Sampling Time (BJT, UTC+8)	Sampling duration Second	Temp °C	RH %	Pressure hPa	Wind Direction °	Wind speed m/s	PM ₁₀ / PM _{2.5} µg/m ³	Particle No.
Jinan (T1)	dust-1	3-18	10:27	60	-	-	-	179	3.9		
	dust-2	3-18	10:28	50	-	-	-	357	2.4	526/98	412
	dust-3	3-18	10:40	100	-	-	-	89	1.7		
Qingdao (T2)	dust-1	3-18	7:54	90	12.5	49.5	1013.3	278	0.7	490/140	486
	dust-2	3-18	20:00	120	8.4	79.3	1013.6	162	1.3	153/55	
Amakusa (T3)	dust-1	3-18	23:16	120	16.2	85.1	1013	121	1.1	63/52	887
	dust-2	3-19	6:36	60	13.7	92.2	1012.6	358	0.6	54/47	

The dust total concentration in Figure S1 clearly shows that at 08:00 on 2014/03/18 (BJT, UTC+8), the dust storm has already passed T1 and T2, heading eastwards (wind arrows) to reach T3. The mean sea level pressure and wind vectors in Figure S2 could also confirm the movement of this high pressure system (yellow) on 03/17, 03/18, and 03/19.

The air behind a cold front is colder and drier than the air in front. When the cold front passes through, the RH and temperature can drop, and the pressure can increase. Based on this and Figure S3-S5, we can define the time of the cold front passage. The cold front arrived T1 around 14:00 on 03/17, T2 at 17:00 on 03/17, and T3 at 17:00 on 03/18 (BJT, UTC+8). This is consistent with the arrival time derived from the PM concentration in Figure 1b.




Figure S1. Distributions of dust in East Asian region at 2014/03/18. Data is provided by the National Institute for Environment Studies and Kyushu University (<http://www-cfors.nies.go.jp/~cfors/>).




Figure S2. The mean sea level pressure (hPa) and wind vectors (knots) at 03/17, 03/18, and 03/19 at the three sampling sites. Data source: NOAA Air Resources Laboratory's (ARL) Real-time Environmental Applications and Display sYstem (READY) (Rolph et al., 2017).




Figure S3. Time series of surface weather conditions: relative humidity (RH), temperature (T), and pressure (P) at Jinan (T1). The start time of the cold front passage is marked by the solid line. Collection time of dust samples is marked by the dash lines. Data source: the National Oceanic and Atmospheric Administration (NOAA).




Figure S4. Time series of surface weather conditions: relative humidity (RH), temperature (T), and pressure (P) at Qingdao (T2). Data source: the National Oceanic and Atmospheric Administration (NOAA).




Figure S5. Time series of surface weather conditions: relative humidity (RH), temperature (T), and pressure (P) at Amakusa (T3).

2. The correlation between EVD and ECD obtained by AFM




Figure S6. The correlation of equivalent circle diameter (ECD) and equivalent volume diameter (EVD) at three sampling sites obtained by AFM.

$$A = \pi r^2 = \pi \times \left(\frac{d}{2}\right)^2 = \frac{\pi d^2}{4} \rightarrow d = \sqrt{\frac{4A}{\pi}} \quad (1)$$

$$V = \frac{4}{3} \pi r^3 = \frac{4}{3} \times \frac{\pi D^3}{8} = \frac{\pi D^3}{6} \rightarrow D = \sqrt[3]{\frac{6V}{\pi}} \quad (2)$$

where d is the equivalent circle diameter (ECD) and D the equivalent volume diameter (EVD) (Chi et al., 2015).

References

- Chi, J. W., Li, W. J., Zhang, D. Z., Zhang, J. C., Lin, Y. T., Shen, X. J., et al. (2015), Sea salt aerosols as a reactive surface for inorganic and organic acidic gases in the Arctic troposphere, *Atmospheric Chemistry and Physics*, 15(19), 11341-11353.
- Rolph, G., Stein, A., and Stunder, B. (2017), Real-time Environmental Applications and Display sYstem: READY, *Environmental Modelling & Software*, 95, 210-228.