Referee 1:

This paper presents the results of a systematic aerosol dispersion and air quality modeling exercise based on the 2018 Camp Fire in northern California. A careful analysis is given, using a combination of satellite and surface measurements over the duration of the fire. There is a great need to analyze and improve such models, to provide reliable air quality forecasting. This work is worthy of publication in ACP in my opinion. I hope the notes below offer some avenues for minor improvements.

We appreciate the reviewer's valuable comments and constructive suggestions. We have carefully revised the manuscript according to these comments. Point-by-point responses are provided below. The reviewer's comments are in black, our responses are in blue, and the quotes from our manuscript are in italics.

1. Lines 110 to 116. I'm not clear what values were selected for some of the parameters in Equation 1, and the degree to which uncertainty in these parameters affects the final model results. This is only partly addressed in Section 4.2; presenting what was learned about emissions process modeling in more detail might be helpful.

The physical meaning and values of the parameters in the wildfire emission have been further elaborated on Page 5:

"for a certain species η , α_{veg} is the carbon density (the mass of burnable above-ground biomass per unit area of vegetation), β_{veg} is the combustion factor, EF_{veg} is the emission factor by species and vegetation type, and a_{fire} is the burning area of each fire pixel. Vegetation type is generated from the MODIS data following IGBP land cover classification. Vegetation type-specific emission factors (EF_{veg}) and combustion factors (β_{veg}) are derived from Ward et al. (1992) and Andreae and Merlet (2001). Vegetation type-specific carbon density (α_{veg}) is based on Olson et al. (2000) and Houghton et al. (2001). "

2. Lines 122-123. Wildfires tend to have a very distinct diurnal cycle. Especially given the extensive modeling effort performed here, using a 13:30 local time sample as diurnally representative might not be the best assumption. There also might be MODIS FRP data at 10:30 am as well as nighttime sampling for this fire.

We agree with the reviewer that the diurnal cycle of wildfires can be important in simulating firerelated aerosol formation. However, the current fire emission module used in this study, the "PREP-CHEM-SRC-1.5", averages all fire detections from satellite within one day and does not provide diurnal information, even though the VIIRS satellite provides one daytime retrieval and one nighttime retrieval of active fire count. We have added a statement in the discussion on Page 13 "*Future studies are needed to further improve the present modeling framework to simulate wildfires. Some wildfires exhibit a distinct diurnal cycle, but the current fire preparation module does not utilize the time information of nighttime fire radiative power measurements by the polar-orbiting satellites*".

3. Lines 144-146. The Camp Fire reportedly also burned the town of Paradise, California between 8 and 10 November 2018. Does urban structure represent a land cover type that should be included in the simulation, as it can produce very different emissions from grassland or forest?

We have added a statement in the discussion on Page 13 "Also, the current land cover and vegetation type data are still relatively coarse in spatial resolution and classification accuracy, which cannot fully resolve a small town in a rural area. In fact, the Camp Fire reportedly burned the town of Paradise, California between 8 and 10 November 2018. The town of Paradise covered 11,614 acres which corresponds to about 7.6% of the total burned area. This discrepancy definitely contributes to the uncertainty in the fire emission preparation. Additional verification of input fire data sources, such

as FINN, and their implementation in the WRF-Chem plume rise model is needed for studies of the vertical structure".

4. Lines 274-278. Given the assumptions required to perform the TROPOMI ALH retrieval, it might be worth comparing the results with any height retrievals from MODIS/MAIAC (Lyapustin et al. 2019, doi:10.1109/LGRS.2019.2936332) or MISR or CALIPSO. The comparison might help quantify measurement uncertainty. Line 285. Is the TROPOMI ALH actually accurate to 100 m?

We have now clarified that the median error of TROPOMI over land is about 1.75 km (Nanda et al., 2020), so most model-obs. differences are within that retrieval bias. We do not find the plume height products about the Camp Fire from either MODIS/MAIAC or MISR in the public domain, so we leave this satellite intercomparison task for the future study. We have added a statement in the discussion on Page 13 "*The recent TROPOMI aerosol layer height product shows promise as an analytical tool, but requires further development of the method by which it can be directly compared to WRF-Chem. Given the assumptions required to perform the TROPOMI ALH retrieval, more research is needed to compare that product with any height retrievals from MODIS/MAIAC (Lyapustin et al. 2019), MISR (Kahn, 2020), and CALIPSO".*

5. Line 282. Figure 9 is first referenced after Figures 10, 11, and 12. Probably warrants renumbering.

We have re-ordered the figures as suggested.

6. Lines 337-339. There are notable uncertainties in the satellite estimates of smoke emissions. For example, the satellite results are not species-specific, relatively coarse pixel resolution contributes, etc. The factor of 5 adjustment from Archer-Nicholls et al. is not unusual, and is an indication of the underlying limitations.

We agree with the reviewer on the satellite product uncertainty and it has been thoroughly discussed in the manuscript.

7. Given the complexity of the problem, I understand why you perturb individual factors in this study. As you have built an advanced modeling capability to assess smoke dispersion for air quality applications, I'm wondering whether testing at least a few combinations of the main factors might yield some additional insights. There are likely some non-linear interactions among mechanisms, and if the goal is to improve air quality prediction, this might be important.

To address the reviewer's comment about the non-linearity of different factors in regulating the firerelated PM pollution, we have added a new experiment by jointly perturbing two chosen factors, i.e. emission flaming factor and aerosol radiative feedback. We compare the results from this joint perturbation experiment with those from each individual perturbation experiment and the linear sum of the two. A new Figure 14 is provided with a new paragraph of discussion.



Fig. 14. Comparison of the effects on PM2.5 simulations in the Bay Area from the individual factor perturbation experiments and the joint perturbation experiment.

We have added a new discussion on Page 12 "To test the linearity of different factors in regulating the fire-related PM pollution, we choose two factors, emission flaming factor and aerosol radiative feedback, and conduct a new experiment by jointly perturbing these two. We compare the results from this joint perturbing experiment with those from each individual perturbing experiment and the linear sum of the two in Figure 14. It shows that for the most times, the effect of joint perturbation is close to the sum of the two individual effects (the black line follows well with the black circles), indicating that the relatively good linearity and additivity holds between those two factors in a general sense. The exception occurs under the extreme conditions. During Nov. 14-18 when the plume was thick and PM2.5 concentration was highest in the Bay Area, the aerosol radiative feedback dominates, and the effect of joint perturbation is close to the aerosol radiative feedback.

We have also added a statement in the conclusion and discussion section on Page 13 "Given the complexity of the problem, here we mainly perturb individual factors in this study. Future studies can test different combinations of the main factors identified by the present study, which can yield additional insights about non-linear interactions among different processes related with fire emission and transport".

Referee 2:

A comprehensive study on surface-level air quality impacts of the 2018 Camp Fire is conducted using a combination of WRF-Chem numerical simulations, ground-based monitoring station observations of PM2.5, black carbon ,carbon monoxide and meteorology, and a suite of space-borne satellite measurements for three separate regions, including (i) close proximity to the fire, (ii) the Sacramento Metropolitan Area and (iii) the San Francisco Bay Area. Evaluation of model simulations against ground-based observations showed good agreement for surface-level wind fields, ambient temperatures, and temporal trends in downwind PM2.5 and black carbon concentrations. Comparison to satellite products demonstrated the ability of model simulations to replicate the general spatiotemporal structure and evolution of the wildfire plumes. Sensitivity analyses were performed to investigate the influence of key parameter perturbations on the accuracy of model predictions relative to the baseline control simulation, including (i) aerosol radiative feedback, (ii) boundary layer dynamics, (iii) plume rise and entrainment, (iv) fire inventory data, (v) emission rates and (vi) flaming versus smoldering partitioning. Results indicate greatest sensitivity to fire emission and boundary layer parameterizations. The main objective of these efforts is to assist improvement in air quality forecasting of wildfire events to ultimately protect human health and reduce economic impact.

Major Comments: The authors do a commendable job in the scale and scope of their simulations and analyses. There is little doubt these efforts will be of interest to the broader community and promote forward movement of this field. The reviewer recommends publication upon consideration of a few key points and minor revisions.

We appreciate the reviewer's valuable comments and constructive suggestions. We have carefully revised the manuscript according to these comments. Point-by-point responses are provided below. The reviewer's comments are in black and our responses are in blue.

1. Perhaps the most striking feature of these wildland-urban interface firestorms is the scale of destruction of the built environment, including Santa Rosa during the 2017 Napa/Sonoma wildfires, Redding during the 2018 Carr Fire, and Paradise during the 2018 Camp Fire studied here. Although wildfires have been studied for decades and there is vast literature characterizing biomass combustion emissions, there are large knowledge gaps in the composition and toxicity of these emissions when a nontrivial fraction of the burnt area includes built environment comprising a vast array of non-biomass related materials. There is clearly a paucity of the types of land cover and fire emissions data required to incorporate these considerations into model simulations, but the reviewer feels it is a key point of sufficient significance to merit inclusion in the manuscript, if only from a speculative perspective. This discussion could easily be incorporated into section 4.2 - Fire Emission Inventory - or as a standalone subsection. Is it possible to calculate what fraction of the burned area can be attributed to the built environment relative to the other landcover vegetation types for the days that Paradise burned? If so, then these data could be included in Figure 3. Presumably, a large fraction of the non-biomass related materials does not sustain flaming combustion but rather are subjected to high temperature pyrolysis analogous to smoldering, which impacts gas-particle partitioning, particle size and composition, and injection heights, and thus downwind simulated surface-level PM2.5 concentrations. A brief synopsis of these complexities and their impact on model performance would be beneficial.

As suggested by the reviewer, we now discuss the fact that there is a paucity of the types of land cover (especially residential area) and fire emissions data required to incorporate these considerations into model simulations in the beginning of Section 4.2, which serves as motivation to conduct emission perturbation experiments. We have estimated that the area of Paradise, covering 11,614 acres, corresponds to about 7.6% of the total burned area. We acknowledge that this contributes to the uncertainty in the fire emission preparation in the final discussion.

2. There is a surplus of figures in the manuscript (14 total), many of which are large multi-panel figures, and some effort should be made to condense these to a critical mass necessary for effective visual dissemination of results and conclusions. For example, Figures 1 and 2, although well crafted, are nonessential to reader comprehension and can easily be described in text. Similarly, Figures 4 and 5 do not elucidate additional clarification to what is already discussed in the manuscript and well summarized in Table 2. Furthermore, Figures 10-12 all support the same underlying fundamental conclusion: deviation of simulated wind fields from observation explains underprediction of downwind surface-level PM2.5 mass concentrations in the Bay Area for the period Nov. 14-16. Only one figure (10 or 11) is necessary to make the point.

As suggested, we have removed previous Figures 11-12 and retained Figure 10 only. Meanwhile, we prefer to keep Figs. 1 and 2. Fig. 1 shows the model domains that are important information for the reader. Fig. 2 educates the readers about how the plume rise model works, for which there is no previous literature/document available. Figs. 4 and 5 provide useful information about the prevailing winds and their temporal evolution during the Camp Fire, which are closely related with pollutant transport.

Minor Revisions:

1. Consider not abbreviating LSM (land surface model) in Table 1;

2. Figures 6, 7, 10, 11, and 14: change y-axis and color scales from ug/m² to ug/m³;

3. In Table 3, Bay Area normalized mean bias is missing percentage symbol (%);

4. Figures should be numbered in the order in which they are discussed within the text, but Figures 10-12 are discussed prior to Figure 9; please renumber figures.

We thank the reviewer for identifying these issues. All the above suggested revisions have been made.

Air Quality Impact of the Northern California Camp Fire of November 2018

Brigitte Rooney¹, Yuan Wang^{1,2*}, Jonathan H. Jiang², Bin Zhao³, Zhao-Cheng Zeng⁴, and John H. Seinfeld^{5*}

¹Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

5 ²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

³Pacific Northwest National Laboratory, Richland, WA, USA

⁴Joint Institute for Regional Earth System Science and Engineering, University of California, Los Angeles, CA, USA

⁵Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA

10

*Corresponding authors:

Yuan Wang: yuan.wang@caltech.edu

John Seinfeld: seinfeld@caltech.edu

15 Abstract

The Northern California Camp Fire that took place in November 2018 was one of the most damaging environmental events in California history. Here, we analyze ground-based station observations of airborne particulate matter that has a diameter < 2.5 micrometers (PM_{2.5}) across northern California and conduct numerical simulations of the Camp Fire using the Weather Research and Forecasting model online coupled with Chemistry (WRF-Chem).

- 20 Simulations are evaluated against ground-based observations of PM_{2.5}, black carbon, and meteorology, as well as satellite measurements, such as Tropospheric Monitoring Instrument (TROPOMI) aerosol layer height and aerosol index. The Camp Fire led to an increase in Bay Area PM_{2.5} to over 70 µg m⁻³ for nearly two weeks, with localized peaks exceeding 300 µg m⁻³. Using the Visible Infrared Imaging Radiometer Suite (VIIRS) high resolution fire detection products, the simulations reproduce the magnitude and evolution of surface PM_{2.5} concentrations, especially
- 25 downwind of the wildfire. The overall spatial patterns of simulated aerosol plumes and their heights are comparable with the latest satellite products from TROPOMI. WRF-Chem sensitivity simulations are carried out to analyze uncertainties that arise from fire emissions, meteorological conditions, feedback of aerosol radiative effects on meteorology, and various physical parameterizations, including the planetary boundary layer model and the plume rise model. Downwind PM_{2.5} concentrations are sensitive to both flaming and smoldering emissions over the fire, so
- 30 the uncertainty in the satellite derived fire emission products can directly affect the air pollution simulations downwind. Our analysis also shows the importance of land surface and boundary layer parameterization in the fire simulation, which can result in large variations in magnitude and trend of surface PM_{2.5}. Inclusion of aerosol radiative feedback moderately improves PM_{2.5} simulations, especially over the most polluted days. Results of this study can assist in the development of data assimilation systems as well as air quality forecasting of health exposures and economic impact studies.

1 Introduction

50

Wildfires have become increasingly prevalent in California. It has been reported that between 2007 and 2016, as many as 3672 fires occurred in California, consuming up to 434,667 acres (Pimlott et al., 2016). Increasingly, the population has expanded into high fire-risk areas and near wildland-urban interfaces (Brown et al., 2020). The intense
smoke consisting of airborne particulate matter of diameter < 2.5 micrometers (PM_{2.5}) associated with these fires leads to an increased risk of morbidity and mortality (Cascio, 2018). PM_{2.5} from wildfires consists of a spectrum of light scattering and absorptive particles largely comprising organic and black carbon. It is increasingly important to understand the cause and nature of wildfires as the number of extreme events and the length of the wildfire season continue to grow (Kahn, 2020; Shi et al., 2019). Fire-related studies have estimated exposures to PM_{2.5} based on ground-level monitoring-station measurements (Shi et al., 2019; Herron-Thorpe et al., 2014; Archer-Nicholls et al., 2015). Spatial coverage of such monitoring stations often tends to be scarce, especially in rural areas. Satellite remote sensing offers a powerful method to monitor air quality during fire events. One study used radiance measurements from the TROPOspheric Monitoring Instrument (TROPOMI) to derive atmospheric carbon monoxide and assess the resulting air quality burden in major cities due to emissions from the California wildfires from November 2018

Deleted:

(Schneising, et al., 2020). Ideally, analysis of fire events is based on a combination of satellite-based measurements

and ground-level observations to obtain spatial and temporal distributions of emissions. The Camp Fire of November 2018 was, to date, the deadliest and most destructive wildfire in California (Kahn, 2020; Brown et al., 2020). Originating along the Sierra Nevada mountain range, smoke from the fire spread across the Sacramento Valley to the San Francisco Bay Area. Peak levels of $PM_{2.5}$ in the San Francisco area exceeded 200 µg m⁻³ and remained above 50 µg m⁻³ for nearly two weeks.

55

Numerous studies have addressed wildfire events using a variety of model frameworks and data sources (Shi et al., 2019; Herron-Thorpe et al., 2014; Archer-Nicholls et al., 2015; Sessions et al., 2011). Shi et al. (2019) used the WRF-Chem model with Moderate Resolution Imaging Spectroradiometer (MODIS) and VIIRS fire data to study the

- 60 wildfire of December 2017 in Southern California. Herron-Thorpe et al. (2014) evaluated simulations of the wildfires in the Pacific Northwest of 2007 and 2008 using the Community Multi-scale Air Quality (CMAQ) model with fire emissions generated by the BlueSky framework and fire locations determined by the Satellite Mapping Automated Reanalysis Tool for Fire Incident Reconciliation (SMART-FIRE). That study suggested that underprediction of PM_{2.5} was the result of underestimated burned area as well as underpredicted secondary organic
- 65 aerosol (SOA) production and incomplete speciation of SOA precursors within the CMAQ model. Archer-Nicholls et al. (2015) simulated biomass burning aerosol during the 2012 dry season in Brazil using WRF-Chem and fire emissions prepared from MODIS. That study proposed that biases in the model were likely a result of uncertainty in the plume injection height and emissions inventory, as well as simulated aerosol sinks (e.g., wet deposition), and lack of inclusion of SOA production in the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC). Sessions
- 70 et al. (2011) investigated methods for injecting wildfire emissions using WRF-Chem. That study tested two fire data preprocessors: PREP-CHEM-SRC (included with WRF-Chem) and the Naval Research Laboratory's Fire Locating and Monitoring of Burning Emissions (FLAMBE), and three injection methods: the 1-D plume rise model within WRF-Chem, releasing emissions only within the planetary boundary layer, and releasing emissions between 3 and 5 km. That study compared results from simulating wildfires during the NASA Arctic Research of the Composition of
- 75 the Troposphere from Aircraft and Satellites (ARCTAS) field campaign in 2008 with satellite data. Sessions et al. (2011) found that differences in injection heights result in different transport pathways.

The present study is a comprehensive investigation of air quality impacts of the Camp Fire using a combined analysis of ground-based and space-borne observations and WRF-Chem simulations. Descriptions of the observation and model are presented in Section 2; model evaluation is presented in Section 3; results of analysis are given in Section 4, followed by discussion and conclusion in Section 5.

2 Model Description and Observational Data

The present study employs WRF-Chem (version 3.8.1) driven by the latest version of meteorological reanalysis data for initialization and boundary conditions. Fire emissions are determined by pairing active fire location data from VIIRS Satellite with the Brazilian Biomass Burning Emission Model (3BEM), which calculates species mass emissions from the burned biomass carbon density, combustion factors, emission factors, and the burning area. WRF-

85

80

Chem simulations are evaluated against EPA surface observations and TROPOMI satellite products.

2.1 WRF-Chem Configuration

90

The WRF-Chem simulation time period is 7 November 2018 (a day before the fire began) to 22 November 2018 (when the fire was 90% contained). We carried out simulations over two domains (Fig. 1): Domain 1 includes all of California at 8 km × 8 km horizontal resolution, while Domain 2 covers Northern California at 2 km × 2 km horizontal resolution. 49 vertical layers are used from the surface to 100 hPa with 50 m vertical resolution in the planetary boundary layer. The meteorological boundary and initial conditions for the outer domain are generated from the fifth generation of European Centre for Medium-range Weather Forecasts (ECMWF) Re-Analysis dataset (ERA5) at 30 km × 30 km resolution (Copernicus Climate Change Service, 2017). Chemical boundary and initial conditions for the outer domain are generated from the Model for Ozone and Related Chemical Tracers version 4 (MOZART-4) (University Corporation for Atmospheric Research, 2013).

95

We use physical options of the Noah Land-Surface Model (Tewari et al., 2004), the Mellor-Yamada-Janjic (MYJ) boundary layer scheme (Janjic, 1994), and the RRTM (longwave) and Dudhia (shortwave) radiative transfer schemes (Dudhia, 1989). Cumulus parameterization is not included. The second-generation Regional Acid Deposition
Model (RADM2) chemical mechanism coupled with the Modal Aerosol Dynamics model for Europe (MADE) and Secondary Organic Aerosol Model (SORGAM) (Zhao et al., 2011) are employed. Aerosol optical properties are calculated based on the volume approximation, for which the volume average of each aerosol species is used to calculate refractive indices (Jin, et al., 2015). Aerosol radiative feedbacks on meteorology and chemistry are included in the simulations.

105 We use the National Emission Inventory for anthropogenic emissions (US EPA, 2018). Biogenic emissions are calculated online using the Guenther scheme (Guenther et. al., 2006). Dust emissions are calculated online using the Goddard Chemical Aerosol Radiation Transport (GOCART) dust emission scheme with University of Cologne (UOC) modifications (Shao et al., 2011). Sea salt emissions are excluded. Technical details of wildfire emissions and the plume rise calculation are discussed in the next section.

110 2.2 Fire Emissions Inventory and Plume Rise Model

Wildfire emissions are generated using the PREP-CHEM-SRC v1.5 preprocessor (Freitas et al., 2011) employing the Brazilian Biomass Burning Emission Model (3BEM, Longo et al., 2010) with satellite data on detected fires. For each pixel with fire detected, the mass of emitted species is calculated by:

$$M^{[\eta]} = \alpha_{veg} \cdot \beta_{veg} \cdot EF_{veg}^{[\eta]} \cdot a_{fire}$$

for a certain species η, where α_{veg} is the carbon density (the mass of burnable above-ground biomass per unit area of vegetation), β_{veg} is the combustion factor, EF_{veg} is the emission factor by species and vegetation type, and a_{fire} is the burning area of each fire pixel. Vegetation type is generated from the MODIS data following IGBP land cover classification. Vegetation type-specific emission factors (EF_{veg}) and combustion factors (β_{veg}) are derived from Ward et al. (1992) and Andreae and Merlet (2001). Vegetation type-specific carbon density (α_{veg}) is based on Olson et al.
(2000) and Houghton et al. (2001). Active fire detection is retrieved from the VIIRS fire product with 375 m spatial

Formatted: Indent: First line: 0"

Deleted:

(1)

Moved down [1]: Active fire detection is retrieved from the VIIRS fire product with 375 m spatial resolution. Deleted: Carbon Moved (insertion) [1] 125 <u>resolution.</u> A limitation of the VIIRS fire count product is its relatively low temporal resolution. As a polar-orbiting satellite, VIIRS provides fire detection during the daytime only once (about 13:30 local time) at each location.

Deleted: In the absence of data to the contrary, we assume equal fire emission rates for each hour of the day.

The emission preprocessor generates a file formatted for WRF-Chem containing the smolderingphase surface emission fluxes of each species, the fire size for each vegetation type, and flaming factor. Flaming factor is the ratio of biomass consumed in the flaming phase to biomass consumed in the smoldering phase. The 17 IGBP land cover classes are aggregated into four main types: tropical forest, extratropical forest, savanna, and grassland. The size of the wildfire and phase of combustion play important roles in the structure of the plume and the vertical distribution of emissions. Wildfire combustion is generally considered to occur in two phases: smoldering and flaming. Emissions from the smoldering phase are allotted to the first layer of the computational grid, while those from the flaming phase are released at injection heights above the surface, as determined by the plume rise model

- 135 described below. Fire size determines the total surface heat flux, as well as the entrainment radius of the plume. Fire parameters are ascribed a daily temporal resolution and are distributed to the WRF-Chem domains. The fire parameters are then input to the plume rise model (Freitas et al., 2007, 2010). The plume rise model is a 1-dimensional model implemented in each WRF-Chem grid cell with an independent vertical grid resolution of 100 m. It calculates the maximum height to which a plume reaches and distributes emissions therein (Fig. 2). The plume top
- height, determined by the surface heat flux from the fire and the thermodynamic stability of the atmospheric environment, is defined as the height at which the in-plume parcel vertical velocity < 1 m s⁻¹. The plume rise model uses upper and lower bounds of heat fluxes determined by each land type to calculate the minimum and maximum plume top height. Flaming emissions are distributed equally to each vertical level within the injection layer with the following calculation: *Flaming Emission per Level = Smoldering Emission × Flaming Factor × DZ⁻¹*, where *DZ = Maximum Plume Top Height Minimum Plume Top Height Minimum Plume Top Height*. The model also accounts for entrainment, water balance,
- and internal gravity wave damping.

Figure 3 shows the fire size and particulate matter emissions produced from MODIS and VIIRS data. The Camp Fire burned primarily extratropical forest vegetation (which comprised 68% of the total burned area), followed by savanna (23% of total area). The flaming emission rate for species n from vegetation type v, is calculated by

(2)

150

130

At maximum, the carbon monoxide (CO) emission flux was 4.1×10^7 mol km⁻² hr⁻¹, and PM_{2.5} flux was 3.7×10^4 µg m⁻² s⁻¹. On average, 46% of the fuel burned is estimated to have been consumed during the flaming phase.

Flaming Phase Rate_{n,v} = $\sum_{Fire \ Cells} Area_v \cdot Smoldering Phase \ Flaming \ Factor_v$

155

Fire Inventory from NCAR (FINN) Version 1.5 (Wiedinmyer, 2011) is another fire emissions product that we will test in a sensitivity analysis. It is assembled for atmospheric chemistry models with a daily temporal resolution and a 1 km horizontal resolution. FINN is generated using satellite observations of active fires and land cover paired with emission factors and fuel loading estimates. The emissions are allocated to a diurnal cycle following WRAP (2005). FINN outputs the total wildfire emission flux, fire size, and land type fraction. As FINN does not include a smoldering-to-flaming phase ratio, the plume rise model calculates a ratio based on CO emissions.

2.3 Surface and Satellite Observations

The observational data include both ground-based measurements and satellite observations. Meteorological and surface concentration data were obtained from the NOAA's National Climatic Data Center (NCDC) and EPA Air Quality System (AQS), respectively. We focus on three areas: the region closest to the fire, the Sacramento Metro Area (population of 2.5 million), and the San Francisco Bay Area (population of 7 million). Hourly observations of wind speed at 10 m, wind direction at 10 m, temperature at 2 m, PM2.5, black carbon (BC), and CO are available for the sites shown in Fig. 1. We use level-2 products from the TROPOMI onboard the Copernicus Sentinel-5 Precursor satellite (S5P) to evaluate the spatial and vertical distribution of predictions. We compare TROPOMI aerosol layer height retrievals (3.5 km × 7 km) with the predicted WRF-Chem height of maximum PM2.5, and 170 ultraviolet aerosol index (UVAI, 3.5 km × 7 km) with the predicted WRF-Chem BC columns. The model results are sampled around 13:30 local time when S5P passes over California.

2.4 Control and Sensitivity Simulations

165

To investigate the effects of key model parameters on the ability to predict the atmospheric impact of the wildfire, we conduct a range of sensitivity simulations. As meteorology and atmospheric structure play important roles in plume 175 dynamics and the transport of particulate matter, we separately perturb the aerosol radiative feedback to meteorology, the planetary boundary layer parameterization, and the plume entrainment coefficient. To understand further the extent to which fire characteristics provided by satellite data can affect the simulations, we analyze the influence of fire data sources, the emission rate, and partitioning between smoldering phase and flaming phase emissions. A summary of these simulations is provided in Table 1.

- 180 Our evaluation focuses on the control simulation (S_CTRL). S_CTRL applies a factor of 3 to the smoldering emissions on 13 November and a factor of 2 to the smoldering emissions on 14-16 November due to the intermittent cloudy conditions over the northern California on those days. S CTRL uses the native flaming factor and fire size products, the default entrainment constant of 0.05, and the Mellor-Yamada-Janjic planetary boundary layer scheme. In the following scenarios, one parameter is individually perturbed from this configuration. S_EMRAW uses the 185 native emissions input with unaltered smoldering phase emissions, S NOAERO turns off the aerosol radiative feedback to meteorological fields, S_FCTX2 doubles the flame factor for the entire simulation period (thus increasing
- flaming phase emissions without changing the smoldering phase), S_ENTR reduces the entrainment coefficient within the plume rise model from 0.05 to 0.02, and S LSM employs an alternative land surface model and planetary boundary layer scheme. We perform another sensitivity simulation using FINN in place of VIIRS (S FINN).

190 **3** Evaluation of Fire Simulations

3.1 Meteorology

The three spatial areas of our interest differ significantly in topography and meteorology. Figure 4 shows the averaged wind observations and S CTRL predictions. S CTRL captures general wind patterns and achieves strong correlation with observed temperatures in each of the areas (Fig. 5). In the first few days of the Camp Fire, the foothills

- 195 and the Sacramento area experienced strong northerly winds, while the Bay Area experienced northeasterly winds, both predicted by the simulation. Other distinct features like those on 11 November near the fire and in the Bay Area are also reproduced by S_CTRL with some bias in timing. In the Bay Area, winds were typically southerly at speeds less than 2 m s⁻¹ and consistent through most of the simulation duration. In the relatively dry Sacramento Valley inland, winds were also predominantly southerly, but were calmer (< 1 m s⁻¹) and varied more than those on the coast.
- 200 After 11 November, the wind speeds were much slower. Coastal air regulates Bay Area temperatures, whereas the drier Sacramento area experiences a greater temperature range. S_CTRL also produced these relative characteristics, but, in general, generated faster winds and higher temperatures than those observed. A summary of model performance statistics is provided in Table 2. The complex terrain of the Bay Area and the Sierra Nevada Foothills near the fire location likely contribute to uncertainty in predicting meteorological parameters. Note that the 4-5 K mean biases in
- the regional surface temperature (T) are non-negligible. Fig. 5 shows that the largest biases mainly occur during the night when the hourly temperature reaches the minimum during the day (deviation from the 1:1 line), while the daytime temperature matches relatively well with the observations (close to the 1:1 line). However, it is important to realize that the nighttime temperature biases have little influence on the PM simulations we focus on. The temporal evolutions of observed PM near Sacramento and the Bay Area do not show a clear diurnal cycle (Fig. 6), nor do the modeled PM biases, as shown in the next section.

3.2 Surface-Level Particulate Matter

Figure 6 shows the predicted evolution of surface PM2.5 from AQS observations and S CTRL over the period of the wildfire. Within hours of the onset of the Camp Fire, observed PM2.5 concentrations in Sacramento and the San Francisco Bay Area (130 and 240 km downwind) increased from below the National Ambient Air Quality Standard 215 (NAAQS) 24-h average of 35 µg m⁻³ to 50 µg m⁻³. Both areas remained above the standard for more than a week, reaching values of three times the standard for multiple days. The region near the fire, Sacramento, and the San Francisco Bay Area were each out of attainment of the NAAQS 24-h average of PM2.5 for 11, 11, and 12 days, respectively, during 7-20 November, while S CTRL predicted 12, 11, and 11 days, respectively. Much of northern California did not return to attainment until 22 November when the wildfire reached 90% containment. Table 3 220 summarizes the ability of S_CTRL to reproduce observed values of surface PM2.5 in the three focus areas and at stations 27 and 28 in the Bay Area. The model prediction exhibits a mean bias of 64.8 µg m⁻³ in the region of the Camp Fire, -11.4 µg m⁻³ in Sacramento, and -16.8 µg m⁻³ in the Bay Area. Mean bias was smaller at some individual monitoring stations, such as Station 27 and 28 that has mean bias of -9.9 µg m⁻³ and -6.2 µg m⁻³, respectively. In the broader area near the fire, S_CTRL significantly overestimates surface PM2.5, reaching nearly 1 mg m-3 while observed 225 concentrations peaked closer to 300 ug m⁻³. However, S CTRL shows a similar temporal trend to that observed, capturing many peak times. The Sacramento area experienced maxima near 300 µg m3, while the Bay Area reached around 200 µg m⁻³. S CTRL shows good agreement of the magnitude and temporal evolution of surface PM2.5 in the Bay Area and Sacramento for most days, with the exception of 10 November and 14-16 November (to be discussed subsequently). Time series of observed and predicted surface CO and BC in the Bay Area are shown in Fig. 7. Again,

230 S_CTRL shows good agreement with the magnitude and trend of both species. While PM2.5 is largely underpredicted

Moved (insertion) [2]

in the period of 14-16 November, BC is over predicted by 5-10 µg m⁻³ at peaks. S CTRL also produces positive bias in surface CO over 16-18 November.

235

Error in surface PM2.5 can, in part, be attributed to error in the predicted wind fields. In the latter hours of 8 November near the Camp Fire, S_CTRL predicts southerly winds, while observations are steadily northerly, leading to some return of initially transported plume. Again, on 11 November, predicted winds show a dramatic reversal, and surface PM2.5 spikes. In Sacramento on 10 November, observed and predicted northerly winds at midday initially lead to increased PM2.5 concentrations, but winds swing southerly in the later hours. On 13 November, observed winds blow south and transport emissions to Sacramento, while S_CTRL predicts winds in the opposing direction, leading to an underprediction in PM2.5. However, error in predicted wind fields does not explain the substantial 240 underprediction of surface PM2.5 in the Bay Area over 14-16 November, as the station-averaged winds of the area do not show significant deviation from observations. We tested the Four-Dimensional Data Assimilation (FDDA) of large-scale horizontal wind from the ERA5, but it could not reduce the aforementioned biases in wind, possibly due to the fact that the observed wind patterns are driven by some mesoscale or even local-scale dynamics.

To study the structural evolution of the wildfire plume, we compare simulated total black carbon column with TROPOMI UVAI satellite retrievals (Fig. 8). TROPOMI UVAI is based on the difference between wavelength-245 dependent Rayleigh scattering observed in an atmosphere with aerosols and that of a modeled molecular atmosphere (Stein Zweers et al., 2018). This difference is measured in the UV spectral range where ozone absorption is small. A positive residual (red coloring) indicates the presence of UV-absorbing aerosols, like black carbon (BC), while a negative residual (blue coloring) indicates presence of non-absorbing aerosols. As WRF-Chem does not generate an 250 aerosol index parameter, we compare UVAI to total BC column, a significantly absorbing aerosol. Over the period of

- the simulation, broad characteristics and shape, as well as some more distinct features, of the Camp Fire plume are reproduced by S CTRL. Using similar input data sources and WRF-Chem configuration, but a simpler plume rise model, Shi et al. (2019) also capture the general shape of the plume, but underestimate aerosol magnitude. Discrepancies in S CTRL plume transport correlate to bias in surface PM2.5. On the first day of the fire, observations
- 255 show that strong winds in northern California drag the plume west, where steady coastal winds transported the plume south and inland again (Fig. 8). The dynamics creates a dense plume with two narrow stretches. S CTRL predictions of total BC column fail to capture the hook-shape present in the UVAI retrievals but reflect the two separate stretches of narrow plume. The simulation constrains one stretch to the valley, leading to overprediction of surface PM2.5 in Sacramento on 8 November (Fig. 6b). On 11 November, the simulation does not reproduce the second band of the
- 260 plume which wraps along the coast and towards San Francisco; rather, the plume remains more concentrated to the Sacramento Valley again. This leads to underprediction of surface PM2.5 in the Bay Area and overprediction in Sacramento (Fig. 6b and c). The narrow PM2.5 peaks of S CTRL on 14-16 November in Sacramento can likely be attributed to the more pronounced plume on 14 November and 16 November. A stark horizontal gradient of fire emissions could restrict accumulation of PM2.5 averaged over the Sacramento region.
- 265 To investigate the predicted decrease of surface PM2.5 in the Bay Area in the afternoon of 14 November, we individually analyze station 27 (Fig. 2). Figure 9 shows the vertical profile of S CTRL PM2.5 concentrations, the

Deleted: 10) and station 28 (Moved up [2]: Fig. Deleted: 11). Figures 10 and 11 show 270 observed and predicted surface PM_{2.5}, and the observed and predicted wind fields. Additionally, Fig. 10 shows the spatial distribution of PM2.5 and surface winds of observations (a) and predictions (b) at four times on 14 November. In the late morning at station 27, observed winds become northeasterly and PM2.5 spikes as more particle-laden air flows westward (Fig. 2). At the same time, S_CTRL winds also become northeasterly and PM2.5 increases accordingly. However, predicted winds reverse, and PM2.5 levels remain relatively low from midday 14 November to midday 15 275 November. This behavior emerges as part of a larger flow pattern in Fig. <u>10</u>. Throughout the morning of 14 November, the simulated wildfire plume approaches the Bay Area and is then driven back inland by a strong sea breeze in the afternoon, not present in the observational data. This behavior is demonstrated in the vertical profile of PM2.5 (Fig. 2a). A column of clean air flushing the Bay Area leads to a predicted bias of -50 µg m⁻³ on 15 November.

3.3 Aerosol Vertical Profile

280

The TROPOMI ALH retrieval represents vertically localized aerosol layers within the free troposphere in cloudfree conditions and is designed to capture aerosol layers produced by biomass burning aerosol (such as wildfires), volcanic ash, and desert dust (Apituley et al., 2019). ALH is retrieved based on the significant effect of aerosol vertical structure on the high spectral resolution observations in the O2-A band in the near-infrared (759 to 770 nm). The ALH algorithm includes a spectral fit estimation of reflectance across the O2 A band using the Optimal Estimation retrieval 285 method with primary fit parameters of aerosol layer mid pressure and aerosol optical thickness (de Graaf et al., 2019). The assumed aerosol profile is a single uniform scattering layer with a fixed pressure thickness, constant aerosol volume extinction coefficient, and constant aerosol single scatter albedo. The mid pressure of the layer, defined as the average of the top and bottom pressures, is converted to altitude with a temperature profile. This parameterization is best suited for aerosol profiles dominated by a sole elevated and optically thick aerosol layer, which is characteristic 290 of wildfire plumes.

We compare the satellite-derived aerosol layer height to WRF-Chem predictions of PM2.5 using two methods. We define the smoke aerosol layer with a PM2.5 threshold concentration of 3 µg m3. For the first method, the layer height is calculated as the average of heights at which PM2.5 is greater than the threshold. For the second method, these heights are weighted by BC mass. Figure 11 shows the satellite-derived layer height (a) and the S_CTRL model bias 295 of average heights (b) and mass weighted average heights (c). TROPOMI layer heights are generally 1 to 2 km and reach greater than 6 km in some instances. Using purely averaged heights, S CTRL typically overpredicts ALH by 100 to 400 m and remains within a smaller range than TROPOMI. S CTRL layer heights weighted by BC mass are lower, thus improving agreement with the satellite. Note that the reported retrieval bias in TROPOMI ALH is about 780 m for wildfire emission plumes and 1.75 km over land generally (Nanda et al., 2020), so the above model-satellite 300 differences in ALH are within the . Archer-Nicholls et al. (2015) and Sessions et al. (2011) also reported overpredicted aerosol layer heights using WRF-Chem when compared to airborne data and Multi-angle Imaging Spectro Radiometer MISR stereo heights, respectively. Using CMAQ, however, Herron-Thorpe et al. (2014) reported underpredicted heights when compared to Cloud-Aerosol Lidar with Orthogonal Polarization CALIOP products. Archer-Nicholls et al. (2015) found that error in plume injection height can contribute to error in surface PM, and that PM biases were dependent on vegetation type as carbon-density and heat release vary by vegetation. Location of the aerosol layer

305

Deleted: 12

Deleted: 10

Deleted: Station 28 exhibits similar behavior of an increase in PM25 with wind change, then a sharp drop as predicted winds deviate strongly northward. Deleted: 12 Deleted: also

Deleted: 10a and 11a

Deleted: 9

- 315 within the column likely also contributes to error in surface predictions of PM2.5 in this study, however, the current analysis is inconclusive. The assumption of a single, elevated aerosol layer used in the TROPOMI ALH derivation may not be characteristic of the vertical structure predicted by WRF-Chem. As seen in Fig. 9 and 10 and in the vertical profile near the wildfire, layers of aerosol are commonly present at the surface and exist as multiple nonlocalized layers. Sessions et al. (2011) also found that using the FLAMBE fire data preprocessor with emission injection heights 320 not constrained to the boundary layer resulted in better agreement with satellite products than PREP-CHEM-SRC.
- Consideration of the WRF vertical grid is also necessary when comparing surface level values. Further development of the analytic method used to evaluate WRF-Chem aerosol layer heights may provide insight into the behavior of the plume rise model and its vertical structure.

4 Sensitivity Simulation Analysis

325 We conduct sensitivity simulations to investigate the effects of various parameters on the ability of the WRF-Chem model to accurately predict downwind PM concentrations from wildfires. As meteorological conditions and related boundary structure play important roles in plume dynamics and the transport of PM, we separately test the aerosol feedback to meteorology and the land surface model. To understand the extent to which fire characteristics provided by satellite data can affect the simulation, we analyze the fire product sources (VIIRS versus FINN), the total 330 fire emissions, and the division between smoldering versus flaming phase emissions. To examine the influence of the plume rise model, we perturb a key parameter, the entrainment coefficient.

4.1 Aerosol Radiative Feedback to Meteorology

By absorbing and scattering solar radiation, aerosols can impact the radiative fluxes, cloud formation, and precipitation in the atmosphere (Wang et al., 2016; 2020), and, in turn, the meteorological conditions for aerosol formation, transport, and removal (Li et al., 2019). WRF-Chem has the option to couple aerosol-radiative direct effects 335 with meteorology simulation. S NOAERO uses the same input data and configuration as S CTRL, but disables the aerosol radiative feedback. Figure 12 shows the evolution of surface wind speed and temperature throughout the wildfire near the source (a), in Sacramento (b), and in the Bay Area (c). The aerosol radiative impact on simulated meteorology is more pronounced for surface temperature than wind. When aerosol radiative feedbacks are noticeable, 340 colder temperatures and calmer winds are found near the surface. Generally, feedbacks are more evident in the region closer to the fire sources with larger PM concentrations. Also, in the Bay Area, the largest changes in meteorology coincide with the largest differences in surface PM2.5 between the two scenarios (Fig. 2.3), which occurs when higher concentrations are predicted (10-11 November, 14-16 November). Consequently, the aerosol radiative feedback in WRF-Chem acts to stabilize the atmosphere, presumably due to the solar absorption by smoke aerosols and reduction 345 of radiation reaching the surface (Wang et al., 2013). When taking the entire time period into account, the overall smoke radiative effect on meteorology is relatively small in the downwind region, like the Bay Area, even when

4.2 Fire Emission Inventory

aerosol concentrations are high.

Deleted: 10 Deleted: 11

Deleted: 13

Deleted: 14

Deleted: aerosol

Currently fire emission inventories generally have large uncertainty. Although wildfires have been studied for 355 decades and there is vast literature characterizing biomass combustion emissions, there are large knowledge gaps in the composition of these emissions when a nontrivial fraction of the burnt area includes built environment comprising a vast array of non-biomass related materials. For the Camp Fire, there is a paucity of the types of burned land cover and fire emissions data required to incorporate these considerations into model simulations. WRF-Chem input fire files produced with VIIRS and PREP-CHEM-SRC include fire size, smoldering emission flux, and flaming factor. 360 Here, we test the sensitivity of predictions to different emission dataset (FINN (S FINN) versus VIIRS/MODIS), as well as emission injection parameters, such as the smoldering emission flux (S_EMRAW) and flaming factor (S_FCTX2). S_FINN produces very little aerosol, though it captures the timing of some peaks. The aerosol underestimation may be a result of bias in the emission inventory or an issue of its implementation in the plume rise model code, as FINN specifies total wildfire emissions rather than a smoldering and flaming distribution.

365 When VIIRS emission inventory is used, the total wildfire emission flux can be altered through two parameters: the smoldering emission flux at the surface and the flaming factor. Directly increasing the smoldering emission flux adds emissions to the surface layer and increases flaming phase emissions proportionally. Figure 13 shows the impact of doubling smoldering emissions on 13 November and tripling them during 14-16 November. These changes to the inventory more than double concentrations of surface PM2.5 in the area of the wildfire and increase concentrations in 370 the Bay Area by 20 to 60 µg m⁻³ during 14-16 November. Consequently, increasing input of total wildfire emissions

improves the agreement of predictions with observations in Sacramento and the Bay Area, suggesting that some uncertainty may stem from satellite fire products. This finding is supported by Archer-Nicholls et al. (2015), as they applied a factor of 5 to scale up the wildfire emissions in their simulations. By modifying the flaming factor, we perturb only the emissions injected aloft by the plume, as emissions higher in the atmosphere may allow for greater 375 transport downwind. By doubling the flaming factor over the full simulation duration, S FCTX2 recovers 10-35 µg m⁻³ in the Bay Area 14-16 November (Fig. <u>13c</u>), when S_CTRL substantially underpredicts PM_{2.5}.

4.3 Plume Rise Parameterization - Entrainment Coefficient

380

proportional to the plume radius (Freitas et al., 2010). Greater entrainment causes rapid cooling, such that near surface plume temperatures are only slightly warmer than the environment, lowering buoyancy and reducing the plume height. Larger wildfires generate less entrainment and reach higher injection heights. The parameterization also includes the effect of horizontal winds on entrainment. Strong wind shear can enhance entrainment and increase boundary layer mixing (Freitas et al., 2010). Archer-Nicholls et al. (2015) decreased the original entrainment coefficient (Freitas et al., 2007) from 0.1 to 0.05 to improve their simulations of a wildfire. As the Camp Fire developed rapidly and 385 intensely, we performed the sensitivity simulation S ENTR with a lower entrainment coefficient of 0.02 to allow for higher injection heights. However, entrainment perturbation resulted in less than 1% change in surface PM2.5 from S_CTRL. A possible reason is that the background winds were quite strong already, for which the entrainment coefficient played a limited role.

The plume rise model parameterizes entrainment as proportional to the plume vertical velocity and inversely

Deleted:

Deleted: 14c

Deleted: 14

We compare simulations using two different land surface models (LSM) which include the PBL schemes: the Noah LSM with Mellor-Yamada-Janjic (MYJ) PBL and the Pleim-Xiu LSM (referred to here as P-X) with the Asymmetric Convection Model 2 (ACM2) PBL (Janjic, 1994; Pleim and Xiu, 1995; Chen & Dudhia, 2001; Pleim, 2007). Land surface models simulate the heat and radiative fluxes between the ground and the atmosphere (Campbell et al., 2018). Noah LSM has four soil moisture and temperature layers, while the Pleim-Xiu LSM has two (Hu et al, 2014; Campbell et al., 2018). Both include a vegetation canopy model and vegetative evapotranspiration. The PBL scheme provides the boundary layer fluxes (heat, moisture, and momentum) and the vertical diffusion within the column. It uses boundary layer eddy fluxes to distribute surface fluxes and grows the PBL by entrainment. A key
feature of PBL schemes is the inclusion of local mixing (between adjacent layers) and/or nonlocal mixing (from the surface layer to higher layers). The MYJ scheme is a turbulent kinetic energy prediction, while the ACM2 scheme is a member of the diagnostic non-local class. MYJ solves for the total kinetic energy in each column from buoyancy and shear production, dissipation, and vertical mixing. ACM2 has two main components: a term for local transport by

small eddies and a term for nonlocal transport by large eddies. Coniglio et al. (2013) showed that the MYJ scheme
can undermix the PBL in locations upstream of convection in the presence of overly cool and moist conditions near the ground in the daytime, whereas ACM2 can result in an excessively deep PBL in evening. Pleim (AMS, 2007) also noted that ACM2 predicts the PBL profile of potential temperature and velocity with greater accuracy.

The use of P-X and ACM2 results in substantially different aerosol trends and plume evolution, the effects of which are largely location-dependent (Fig. <u>13</u>). Near the fire and in the Bay Area, S_LSM produces little similarity in surface PM_{2.5} magnitude and trend as compared to S_CTRL. S_LSM reduces PM_{2.5} concentrations by more than 50% in both areas for the majority of the simulation period. However, S_CTRL overpredicts PM_{2.5} near the wildfire, while S_LSM underpredicts but produces a more muted temporal pattern, similar to observations. In the Sacramento area, S_LSM generally predicts higher PM_{2.5} values with a distinct diurnal trend. Peaks are of similar magnitude to S_CTRL but displaced temporally. The topography of the Sacramento area is more uniform than the complex terrain of the Bay area as well as the foothills and canyons near the wildfire, likely contributing to the distinctions in the behavior of the two schemes. Moreover, the current sensitivity study stresses the importance of the parameterization of the land surface and the boundary layer. As shown here, the Noah LSM and MYJ scheme performs well for the broader region of northern California, whereas improvement near the wildfire itself may be attained with altered PBL parameterization.

420 4.4 Joint Perturbation

425

To test the linearity of different factors in regulating the fire-related PM pollution, we choose two factors, emission flaming factor and aerosol radiative feedback, and conduct a new experiment by jointly perturbing these two. We compare the results from this joint perturbing experiment with those from each individual perturbing experiment and the linear sum of the two in Figure 14. It shows that for the most times, the effect of joint perturbation is close to the sum of the two individual effects (the black line follows well with the black circles), indicating that the relatively

good linearity and additivity holds between those two factors in a general sense. The exception occurs under the extreme conditions. During Nov. 14-18 when the plume was thick and PM_{2.5} concentration was highest in the Bay

Deleted: 14

Deleted:

Deleted: 5

Area, the aerosol radiative feedback dominates, and the effect of joint perturbation is close to the aerosol radiative effect (the black line follows well with the blue dotted line).

5. Conclusions and Discussion

435

The record-breaking Camp Fire ravaged northern California for nearly two weeks. At a distance of 240 km downwind of the wildfire, Bay Area surface PM2.5 levels reached nearly 200 µg m-3 and remained over 70 µg m-3 over 7-22 November 2018. It is uncertain to what extent the current chemical transport models can reproduce the key features of this historical event. Here, we employ the WRF-Chem model to characterize the spatio-temporal PM concentrations across northern California and to investigate the sensitivity of predictions to key parameters of the model. The model utilizes satellite fire detection products with a resolution of 375 m and a biomass burning model to generate the fire emission inventory at near real time. We conduct model simulations at 2 km resolution. A wide range 440 of observational data is employed to evaluate the model performance, including ground-based observations of PM2.5, black carbon, and meteorology from EPA and NOAA stations, as well as satellite measurements, such as Tropospheric Monitoring Instrument (TROPOMI) aerosol layer height and aerosol index.

We focus on three geographic areas: the vicinity of the wildfire, Sacramento, and the San Francisco Bay Area. 445 The control experiment was able to simulate the general transport and extent of the plume as well as the magnitude and temporal evolution of surface PM2.5 in Sacramento and the Bay Area. Meanwhile, the control experiment substantially overpredicted surface PM2.5 near the fire, but captured the general evolution of the fire development. On the Pacific coast, the Bay Area was subject to significant sea breezes not observed during the time period of simulation. Owing to strong winds predicted from the ocean, a large negative bias existed in surface PM2.5. Increasing total wildfire 450 emissions (smoldering + flaming) and increasing flaming phase emissions alone each recovered some PM2.5 biases. Aerosol radiative feedback on meteorology acted to stabilize the atmosphere and slightly increased the PM2.5 concentration near the surface during most severe episodes. Hence, its inclusion modestly improves model performance. Our study shows that sources of downwind PM error stem primarily from the localized structure of the plume and uncertainty in fire emissions. Uncertainty of partitioning between smoldering and flaming phases may also

455 contribute to uncertainty in plume horizontal transport.

Future studies are needed to further improve the present modeling framework to simulate wildfires. Some wildfires exhibit a distinct diurnal cycle, but the current fire preparation module has not utilized the time information of the fire radiative power measurements by the polar-orbiting satellites. Also, the current land cover and vegetation type data are still relatively coarse in spatial resolution and classification accuracy, which cannot fully resolve a small 460 town in a rural area. In fact, the Camp Fire reportedly burned the town of Paradise, California between 8 and 10 November 2018. The town of Paradise covered 11,614 acres which corresponds to about 7.6% of the total burned area. This contributes to the uncertainty in the fire emission preparation. Additional verification of input fire data sources, such as FINN, and their implementation in the WRF-Chem plume rise model is needed for studies of the vertical structure. Deeper understanding of the role of plume dynamics and boundary layer parameterization on aerosol 465 concentrations downwind from wildfires will inform updates to forecast models like WRF-SFIRE-CHEM, which

Moved down [3]: The recent TROPOMI aerosol layer height product shows promise as an analytical tool, but requires further development of the method by which it can be directly compared to WRF-Chem.

Moved down [4]: Herron-Thorpe et al. (2014) noted that careful consideration must also be given to the vertical coordinates across models and satellite products, as discrepancies in reporting heights in reference to sea level, ground level, or the geoid can influence analyses.

475 couples WRF with a fire spread model and smoke dispersion simulation (Barbunzo 2019; Kochanski et al., 2013). Given the complexity of the problem, here we only perturb individual factors in this study. Future studies can test different combinations of the main factors identified by the present study, which can yield additional insights about non-linear interactions among different processes related with fire emission and transport.

The recent TROPOMI aerosol layer height product shows promise as an analytical tool, but requires further
 development of the method by which it can be directly compared to WRF-Chem. Given the assumptions required to perform the TROPOMI ALH retrieval, more research is needed to compare that product with any height retrievals from MODIS/MAIAC (Lyapustin et al. 2019), MISR, and CALIPSO. The intercomparison can help quantify measurement uncertainty. Herron-Thorpe et al. (2014) noted that careful consideration must also be given to the vertical coordinates across models and satellite products, as discrepancies in reporting heights in reference to sea level, ground level, or the geoid can influence analyses.

Code availability

WRF-Chem model code is available for download via the WRF website
 (<u>https://www2.mmm.ucar.edu/wrf/users/downloads.html</u>). The FINN utility is available for download via the NCAR Atmospheric Chemistry Observations & Modeling website (<u>http://bai.acom.ucar.edu/Data/fire</u>).

Data availability

US Environmental Protection Agency Air Quality System Data Mart (internet database) is available for download
 (https://www.epa.gov/airdata). NCDC data is available for download via the NCEI website (https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C00684/html#). TROPOMI data is available for download via the Copernicus Open Access Hub website (https://scihub.copernicus.eu/). ERA5 data is available for download via the Copernicus Climate Data Store website (https://cds.climate.copernicus.eu/). FINN emission data is available for download via the NCAR Atmospheric Chemistry Observations & Modeling website
 (http://bai.acom.ucar.edu/Data/fire).

Author contribution

Y.W., J.H.S., and J.H.J conceived and designed the research. Y.W., and B.R. performed the WRF-Chem simulations.
B.R., Y.W., and J.H.S. performed the data analyses and produced the figures. B. Z. provided technical support for fire
emission preparation. Z.C.Z. helped satellite data analyses. B.R., Y.W., and J.H.S wrote the paper. All authors contributed to the scientific discussions and preparation of the manuscript.

Competing Interests

The authors declare that they have no conflict of interest.

510

Moved (insertion) [3]

Moved (insertion) [4]

Acknowledgements

This study was supported by the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA. We thank Kristal R. Verhulst, Yi Yin, Don Longo, Gonzalo Ferrada, and Saulo Freitas for their support and discussion.

515

525

References

Andreae, M. O., and Merlet, P.: Emission of trace gases and aerosols from biomass burning. Global Biogeochemical Cycles, 15(4), 955–966, https://doi.org/10.1029/2000GB001382, 2001.

 Apituley, A., Pedergnana, M., Sneep, M., Pepijn Veefkind, J., Loyola, D., Landgraf, J., and Borsdorff, T.: Sentinel-5
 precursor/TROPOMI Level 2 Product User Manual Carbon Monoxide, SRON-S5P-LEV2-MA-002, http://www.tropomi.eu/sites/default/files/files/Sentinel-5P-Level-2-Product-User-Manual-Carbon-Monoxide_v1.00.02_20180613.pdf, 2018.

Apituley, A., Pedergnana, M., Sneep, M., Pepijn Veefkind, J., Loyola, D., Sanders, B., de Graaf, M.: Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Aerosol Layer Height, S5P-KNMI-L2-0022-MA, http://www.tropomi.eu/sites/default/files/fulls/55P-KNMI-L2-0022-MA-

Product_User_Manual_for_the_Sentinel_5_precursor_Aerosol_layer_height-1.3.2-20190926_signed.pdf, 2019.

Archer-Nicholls, S., Lowe, D., Darbyshire, E., Morgan, W.T., Bela, M.M., Pereira, G., Trembath, J., Kaiser, J.W., Longo, K.M., Freitas, S.R., Coe, H., and McFiggans, G.: Characterising Brazilian biomass burning emissiongs using WRF-Chem with MOSAIC sectional aerosol. Geosci. Model Dev., 8, 549-577, https://doi.org/10.5194/gmd-8-549-2015, 2015.

530 2015, 2

Barbuzano, J.: Wildfire smoke traps itself in valleys, Eos, 100, https://doi.org/10.1029/2019EO135955, 2019.

Brown, T., Leach, S., Wachter, B., Gardunio, B.: The Northern California 2018 Extreme Fire Season [in "Explaining Extremes of 2018 from a Climate Perspective"]. Bull. Amer. Meteor. Soc., 101 (1), S1–S4, https://doi.org/10.1175/BAMS-D-19-0275.1, 2020.

535 Campbell, P.C., Bash, J.O., Spero, T.L.: Updates to the Noah land surface model in WRF-CMAQ to improve simulated meteorology, air quality, and deposition. Journal of Advances in Modeling Earth Systems, 11(1), 231-256, https://doi.org/ 10.1029/2018MS001422, 2018.

Cascio, W. E.: Wildland fire smoke and human health, Science of the Total Environment, 624, 586-595, https://.doi.org/10.1016/j.scitotenv.2017.12.086, 2018.

540 Chen, F., and J. Dudhia, 2001: Coupling an advanced land surface-hydrology model with the Penn State-NCAR MM5 modeling system, part I, model implementation and sensitivity. Mon. Wea. Rev., 129. 586-585.

Copernicus Climate Change Service (C3S): ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store (CDS), https://cds.climate.copernicus.eu/cdsapp#!/home, last access: 22 May 2020, 2017.

545 Copernicus Sentinel data processed by ESA, Koninklijk Nederlands Meteorologisch Instituut (KNMI), Sentinel-5P TROPOMI Aerosol Index 1-Orbit L2 7km x 3.5km, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), https://doi.org/10.5270/S5P-0wafvaf. 2018.

550

560

570

Copernicus Sentinel data processed by ESA, Koninklijk Nederlands Meteorologisch Instituut (KNMI), Sentinel-5P TROPOMI Aerosol Layer Height 1-Orbit L2 7km x 3.5km, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), https://doi.org/10.5270/S5P-j7aj4gr, 2019.

Copernicus Sentinel data processed by ESA, Koninklijk Nederlands Meteorologisch Instituut (KNMI)/Netherlands Institute for Space Research (SRON), Sentinel-5P TROPOMI Carbon Monoxide CO Column 1-Orbit L2 7km x 7km, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), https://doi.org/10.5270/S5P-1hkp7rp, 2018.

555 De Graaf, M., de Haan, J.F., and Sanders, A.F.J.: TROPOMI ATBD of the Aerosol Layer Height, S5P-KNMI-L2-0006-RP, http://www.tropomi.eu/sites/default/files/files/publicSentinel-5P-TROPOMI-ATBD-Aerosol-Height.pdf, 2019.

Dudhia, J.: Numerical study of convection observed during the Winter Monsoon Experiment using a mesoscale twodimensional model, J. Atmos. Sci., 46, 3077–3107, https://doi.org/10.1175/1520-0469(1989)046<3077:NSOCOD>2.0.CO;2, 1989.

Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), Geosci. Model Dev., 3, 43-67, https://doi.org/10.5194/gmd-3-43-2010, 2010.

565 Freitas, S. R., Longo, K. M., Chatfield, R., Latham, D., Dias, M., Andreae, M. O., Prins, E., Santos, J. C., Gielow, R., and Carvalho, J. A.: Including the sub-grid scale plume rise of vegetation fires in low resolution atmospheric transport models, Atmos. Chem. Phys., 7(13), 3385–3398. https://doi.org/10.5194/acp-7-3385-2007, 2007.

Freitas, S. R., Longo, K. M., Trentmann, J., and Latham, D.: Technical note: Sensitivity of 1-D smoke plume rise models to the inclusion of environmental wind drag, Atmos. Chem. Phys., 10(2), 585–594. https://doi.org/10.5194/acp-10-585-2010, 2010.

Freitas, S. R., Longo, K. M., Alonso, M. F., Pirre, M., Marecal, V., Grell, G., Stockler, R., Mello, R. F., and Sánchez Gácita, M.: PREP-CHEM-SRC – 1.0: a preprocessor of trace gas and aerosol emission fields for regional and global atmospheric chemistry models, Geosci. Model Dev., 4, 419–433, https://doi.org/10.5194/gmd-4-419-2011, 2011.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., and Geron, C.: Estimates of global terrestrial isoprene
 emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmos. Chem. Phys., 6, 3181-3210, https://doi.org/10.5194/acp-6-3181-2006, 2006.

Herron-Thorpe, F. L., Mount, G. H., Emmons, L. K., Lamb, B. K., Jaffe, D. A., Wigder, N. L., Chung, S.H., Zhang, R., Woelfe, M.D., and Vaughan, J.K.: Air quality simulations of wildfires in the Pacific Northwest evaluated with surface and satellite observations during the summers of 2007 and 2008. Atmos. Chem. Phys., 14(22), 12,533–12,551. https://doi.org/10.5194/acp-14-12533-2014, 2014.

Houghton, R., Lawrence, K., Hackler, J., and Brown, S.: The spatial distribution of forest biomass in the Brazilian Amazon: A comparison of estimates, Global Change Biology, 7(7), 731–746. https://doi.org/10.1046/j.1365-2486.2001.00426.x, 2001.

580

Hu, Z., Zhong-Feng, X., Ning-Feng, Z., Ma, Z., Guo-Ping, L.: Evaluation of the WRF model with different land
 surface schemes: a drought event simulation in southwest China during 2009-10. Atmospheric and Oceanic Science
 Letters, 7(2) 168-173, doi: 10.3878/j.issn.1674-2834.13.007, 2014.

Janjic, Zavisa I.: The Step-Mountain Eta Coordinate Model: Further developments of the convection, viscous sublayer, and turbulence closure schemes, Mon. Wea. Rev., 122, 927–945, https://doi.org/10.1175/1520-0493(1994)122<0927:TSMECM>2.0.CO;2, 1994.

590 Janjic, Z. I.,: Nonsingular implementation of the Mellor-Yamada Level 2.5 Scheme in the NCEP Meso model. NCEP Office Note No. 437, 61 pp., 2002.

Jin, Q., Wei Z.-L., Yang B., Huang J.: Consistent response of Indian summer monsoon to Middle East dust in observations and simulations, Atmos. Chem. Phys., 15, 9897-9915 https://doi.org/10.5194/acp-15-9897-2015, 2015.

Lyapustin, A., Wang, Y., Korkin, S., Kahn, R., Winker, D., MAIAC Thermal Technique for Smoke Injection Height
 From MODIS, IEEE Geoscience and Remote Sensing Letters, 7(5) 730 - 734, 2020.

Li, Z., *et al.*, East Asian Study of Tropospheric Aerosols and Impact on Regional Cloud, Precipitation, and Climate (EAST-AIRCPC), J. Geophys. Res. 124 (23) 13026-13054, 2019.

Longo, K. M., Freitas, S. R., Andreae, M. O., Setzer, A., Prins, E., and Artaxo, P.: The Coupled Aerosol and Tracer Transport model to the Brazilian developments on the Regional Atmospheric Modeling System (CATT-BRAMS)—

600 Part 2: Model sensitivity to the biomass burning. Atmospheric Chemistry and Physics, 10(13), 5785–5795. https://doi.org/10.5194/acp-10-5785-2010, 2010.

Kahn, R. (2020), A global perspective on wildfires, Eos, 101, https://doi.org/10.1029/2020EO138260, 2020.

Kochanski, A., Beezley, D., Mandel, J., Clements, B.: Air pollution forecasting by coupled atmosphere-fire model WRF and SFIRE with WRF-Chem. ArXiv: 1304.7703 [physics.ao-ph], 2013.

605 Mlawer, E.J., Taubman, S.J., Brown, P.D., Iacono, M.J. and Clough, S.A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, J. Geophys. Res., 102, 16663–16682, https://doi.org/10.1029/97JD00237, 1997.

610

630

Nanda, S., de Graaf, M., Veefkind, J. P., Sneep, M., ter Linden, M., Sun, J., and Levelt, P. F.: A first comparison of <u>TROPOMI</u> aerosol layer height (ALH) to CALIOP data, Atmos. Meas. Tech., 13, 3043–3059, https://doi.org/10.5194/amt-13-3043-2020, 2020.

Olson, J., Watts, J., and Allison, L.: Major world ecosystem complexes ranked by carbon in live vegetation: A database, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA. 2000.

Pimlott, K., Laird, J., and Brown, E.G.: 2015 Wildfire Activity Statistics. California Department of Forestry and Fire Protection. Available at: https://www.fire.ca.gov/media/10061/2015_redbook_final.pdf.

615 Pleim, Jonathan E.: A Combined Local and Nonlocal Closure Model for the Atmospheric Boundary Layer. Part I: Model Description and Testing, J. Appl. Meteor. Climatol., 46, 1383–1395. doi:10.1175/JAM2539.1, 2007.

Pleim, J., A. and Xiu.: Development and testing of a surface flux and planetary boundary layer model for application in mesoscale models. Journal of Applied Meteorology, 34, 16-34, https://doi.org/10.1175/1520-0450-34.1.16, 1995.

 Sessions, W.R., Fuelberg, H.E., Kahn, R.A., and Winker, D.M.: An investigation of methods for injecting emissions
 from boreal wildfires using WRF-Chem during ARCTAS. Atmos. Chem. Phys., 11, 5719-5744, https://doi.org/ 10.5194/acp-11-5719-2011, 2011.

Shao, Y., Ishizuka, M., Mikami, M., and Leys, J.: Parameterization of size-resolved dust emission and validation with measurements, J. Geophys. Res. Atmos., 116, D08203, doi:10.1029/2010JD014527, 2011.

Shi, H., Jiang, Z., Zhao, B., Li, Z., Chen, Y., Gu, Y., Jiang, J.H., Lee, M., Liou, K.N., Neu, J., Payne, V., Su, H.,
Wang, Y., Marcin, W., and Worden, J.: Modeling study of the air quality impact of record-breaking Southern California wildfires in December 2017. Journal of Geophysical Research: Atmospheres, 124. https://doi.org/10.1029/2019JD030472, 2019.

Schneising, O., Buchwitz, M., Reuter, M., Bovensmann, H., and Burrows, J. P.: Severe Californian wildfires in November 2018 observed from space: the carbon monoxide perspective, Atmos. Chem. Phys., 20, 3317–3332, https://doi.org/10.5194/acp-20-3317-2020, 2020.

Strahler, A., Muchoney, D., Borak, J., Friedl, M., Gopal, S., Lambin, E., & Moody, A.: MODIS Land Cover Product Algorithm Theoretical Basis Document (ATBD). Version 5.0, available at https://modis.gsfc.nasa.gov/data/atbd/atbd_mod12.pdf, 1999.

Stein Zweers, D.: TROPOMI ATBD of the UV aerosol index, S5P-KNMI-L2-0008-RP, available at http://www.tropomi.eu/sites/default/files/files/S5P-KNMI-L2-0008-RP-TROPOMI_ATBD_UVAI-1.1.0-20180615_signed.pdf, 2018.

Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M.A., Mitchell, K., Ek, M., Gayno, G, Wegiel, J., and Cuenca, R.H.: Implementation and verification of the unified NOAH land surface model in the WRF model. 20th conference on weather analysis and forecasting/16th conference on numerical weather prediction, pp. 11–15. 2004.

640 University Corporation for Atmospheric Research: MOZART DOWNLOAD. https://www.acom.ucar.edu/wrfchem/mozart.shtml, 2018.

US Environmental Protection Agency: Air Quality System Data Mart [internet database] available via https://www.epa.gov/airdata, last access: 17 May 2020.

US Environmental Protection Agency: National Emissions Inventory (NEI). https://www.epa.gov/air-emissionsinventories/national-emissions-inventory-nei, 2018.

Wang, Y., Le, T., Chen, G., Yung, Y.L., Su, H., Seinfeld, J.H., Jiang, J.H.: Reduced European aerosol emissions suppress winter extremes over northern Eurasia, Nat. Climate Change, 10, 225–230, https://doi.org/10.1038/s41558-020-0693-4, 2020.

Wang, Y., Ma, P.-L., Jiang, J., Su, H., and Rasch, P.: Towards Reconciling the Influence of Atmospheric Aerosols
and Greenhouse Gases on Light Precipitation Changes in Eastern China, J. Geophys. Res. Atmos. 121(10), 5878–5887, https://doi.org/10.1002/2016JD024845, 2016.

Wang, Y., Khalizov. A., Levy, M., and Zhang, R.: New Directions: Light Absorbing Aerosols and Their Atmospheric Impacts, Atmos. Environ., 81, 713-715, https://doi.org/10.1016/j.atmosenv.2013.09.034, 2013.

Ward, D., Susott, R., Kauffman, J., Babbitt, R., Cummings, D., Dias, B., Holben, B.N., Kaufman, Y.J., Rasmussen,
R.A., and Setzer, A.W.: Smoke and fire characteristics for cerrado and deforestation burns in Brazil: BASE-B experiment. J. Geophys. Res., 97(D13), 14,601–14,619. https://doi.org/10.1029/92JD01218, 1992.

Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): A high resolution global model to estimate the emissions from open burning. Geoscientific Model Development, 4(3), 625–641. https://doi.org/10.5194/gmd-4-625-2011, 2011.

660 WRAP (Western Regional Air Partnership): 2002 Fire Emission Inventory for the WRAP Region- Phase II, Project No. 178-6, available at: http://www.wrapair.org/forums/fejf/tasks/FEJFtask7PhaseII.html, 2005.

Zhao, C., Liu, X., Leung, L., and Hagos, S.: Radiative impact of mineral dust on monsoon precipitation variability over West Africa. Atmos Chem. Phys., 11, 1879-1893, https://doi.org/10.5194/acp-11-1879-2011, 2011.

665

645

Deleted:

	~					
Table I	Summary	ot.	sensitivity	/ simii	lation	setun
I GOIC I	. Oummun	01	bononti vit,	onna.	iuuon	Secup

2	5	1								
Name	Fire Data	Smoldering Emissions	Flaming Factor	Entrainment Constant	Land Surface Model	Aerosol Radiative Feedback	 Delet	ed: LSM		
S_CTRL ¹	VIIRS	x3 Nov. 13, x2 Nov. 14-16	Native	0.05	Noah/MYJ	Yes				
S_EMRAW	VIIRS	Native	Native	0.05	Noah/MYJ	Yes				
S_NOAERO	VIIRS	x3 Nov. 13, x2 Nov. 14-16	Native	0.05	Noah/MYJ	No				
S_FCTX2	VIIRS	x3 Nov. 13, x2 Nov. 14-16	x2	0.05	Noah/MYJ	Yes				
S_ENTR	VIIRS	x3 Nov. 13, x2 Nov. 14-16	Native	0.02	Noah/MYJ	Yes				
S_LSM	VIIRS	x3 Nov. 13, x2 Nov. 14-16	Native	0.05	P-X/ACM2	Yes				
S_FINN	FINN	-	-	0.05	Noah/MYJ	Yes				
S_FCTX2andNOAERC	<u>VIIRS</u>	<u>x3 Nov. 13,</u> x2 Nov. 14-16	<u>x2</u>	<u>0.05</u>	<u>Noah/MYJ</u>	<u>No</u>				

 lScenario that agrees best with surface observations and is of primary focus in this study. Bold denotes parameter perturbed from the S_CTRL scenario.

V	Demonster	Near	S	Bay	Station	L	 Deleted: Station 28
Variable	Parameter	Source ¹	Sacramento.	Area ¹	27	•	Deleted Cells
Wind	Observation						Formatted Table
Speed ² (m s ⁻¹)	Mean	1.4 (0.2)	1.0 (0.2)	1.6 (0.7)	1.5		 Deleted: 0.7
	S_CTRL Mean	2.6 (0.3)	1.4 (0.4)	2.0 (0.7)	2.3	۲	 Deleted: 1.0
	Mean Bias	1.2	0.5	0.5	0.9	۲	 Deleted: 0.3
Wind Direction ³ (deg)	Observation Mean	360.0	338.2	73.9	68.9	۲	 Deleted: 148.8
	S_CTRL Mean	356.9	325.9	26.7	72.8	۲	 Deleted: 11.8
	Mean Bias	2.9	11.0	0.2	2.8	۲	 Deleted: 10.1
Temp (°C)	Observation Mean	8.2 (2.3)	10.1 (1.7)	10.8 (1.9)	9.9	۲	 Deleted: 8.1
	S_CTRL Mean	12.5 (3.6)	13.7 (1.4)	15.7 (1.2)	15.5	۲	 Deleted: 13.8
	Mean Bias	4.4	3.6	4.9	5.6	۲	 Deleted: 5.7

Table 2. Summary of meteorological model performance metrics for the simulation duration.

¹Area winds are averaged for 4 stations near source, 6 stations in Sacramento, and 12 stations in the Bay Area. Area temperatures are averaged for 10 stations near source, 7 in Sacramento, and 13 in the Bay Area. Standard deviation of station averages is noted in parenthesis.
²Mean wind speed is calculated as the average of the magnitude of the wind vector.
³Mean wind direction is calculated assuming a unity vector. 680

	Parameter	Near	Sacramento*	Bay Area*	Station 27	х	Deleted: Station 28
_		Source*		2			Deleted Cells
	Observation Mean	98.3 (39.7)	77.2 (24.9)	74.1 (5.4)	77.9	v	Formatted Table
	S CTRL Mean	163.1	65.8 (16.3)	57.2 (6.4)	68.1	۲	 Deleted: 69.8
		(108.5)		e , (e)			 Deleted: 63.6
	Mean Bias	64.8	-11.4	-16.8	-9.9	v	 Deleted: -6.2
	Normalized Mean Bias	76.5%	-17.4%	-23.1 <u>%</u>	-12.7%	τ	 Deleted: -8.9%

Table 3. Summary of model performance metrics for surface $PM_{2.5}(\mu g\ m^{-3})$ for the simulation duration.

695 *Area values are averaged for 5 stations near source, 7 stations in Sacramento, and 13 stations in the Bay Area. Standard deviation of station averages is noted in parenthesis.



Figure 1. Study domain (a) and observation station locations (b,c). Domain d01 covers the western US with a horizontal resolution of 8 km. Domain d02 is centered over northern California with a horizontal resolution of 2 km. AQS and NCDC observation sites are shown in panel b and panel c, where stations marked in green measure only PM_{2.5}, stations in blue measure wind and temperature, stations in orange measure both PM_{2.5} and meteorology, and stations in yellow measure temperature only. Additionally, BC and CO are measured at 8 and 12 sites in the Bay Area, respectively.



Figure 2. Plume rise model schematic. For each grid cell in which wildfire occurs, the plume rise model uses satellite fire products and the surrounding WRF-Chem environmental conditions to calculate two plume top heights by using the land-type dependent minimum and maximum wildfire heat fluxes. Smoldering phase emissions are allotted to the surface layer, while flaming phase emissions are distributed linearly aloft within the injection layers at a vertical resolution of 100 m.

r



Figure 3. Wildfire area by vegetation type in m² (a) and PM_{2.5} emission rate in µg s⁻¹ by combustion phase and species (b) input into WRF-Chem. The base inventory is produced from VIIRS and MODIS fire products using the PREP-CHEM-SRC processor and is employed by S_EMRAW. The control and remaining sensitivity simulations use an 720 inventory with triple emission flux of all species on 13 November and double during 13-16 November, shown here. About 59% of total PM2.5 emissions occur in the smoldering phase (darker colors in panel b). The total PM2.5 emitted is composed of 69.5% organic carbon and 4.5% black carbon. The Camp Fire burned primarily extratropical forest (purple) followed by savanna (yellow). Burning of extratropical forest generated the greatest fraction of emissions in the flaming phase at 44.2%, followed by savanna at 22.9% and tropical forest at 17.4%. Grassland emits only in the smoldering phase.



730

Figure 4. Comparison of AQS and NCDC wind observations (black) with S_CTRL predictions (red) averaged over the three areas of study: a) near the wildfire (N = 4), b) Sacramento (N = 6), and c) the Bay Area (N = 12). Arrows indicate the wind direction and their length represents wind speed. For reference, S_CTRL predicts maximum wind speeds of 8.7, 7.5, and 7.1 m s⁻¹ near the source, in Sacramento, and in the Bay Area, respectively. Paradise and the Sacramento areas experienced strong northerly winds during the first few days of the fire. S_CTRL generally predicted faster and more variable winds, but broader trends in Sacramento and the Bay Area were represented well.



Figure 5. Comparison of AQS and NCDC temperature observations versus S_CTRL predictions: a) near the wildfire
 (N = 10), b) Sacramento (N = 7), and c) the Bay Area (N = 13). The solid red lines show a linear regression fit, while the dotted black lines denote 1:1 simulations vs. observations. The simulations achieved a correlation coefficient R² of 0.61 near the fire, 0.72 in Sacramento, and 0.75 in the Bay Area.



Figure 6. Comparison of AQS surface PM_{2.5} observations (black) with S_CTRL predictions (red) averaged over the three areas of study: a) near the wildfire (N = 5), b) Sacramento (N = 7), and c) the Bay Area (N = 13). Shading indicates the standard deviation of the sampled stations. S_CTRL overpredicted PM_{2.5} in the region in the vicinity of the fire but performed well in the areas downwind.



Figure 7. Comparison of AQS surface black carbon (a, N = 5) and carbon monoxide (b, N = 12) observations (black)
 with S_CTRL predictions (red) at monitoring sites in the Bay Area. S_CTRL captures the temporal evolution of BC and CO and is close to observed values. BC peaks are often overpredicted. The greatest bias of BC and CO occurs during 16-18 November, likely due to the scale factor applied to emissions during 13-16 November.







Figure 9, Vertical profile of $PM_{2.5}$ (a), time series of surface $PM_{2.5}$ (b), winds (c; observations in black and predictions in red) at Station 27 in the Bay Area. The gray box highlights the timeframe of greatest model bias of surface $PM_{2.5}$. 755 Sharp increases in PM2.5 correlate with a switch to northeasterly winds that import fire emissions to the Bay Area. Large negative PM2.5 bias on 15 November occurs when S_CTRL deviates from observations and produces southerly winds which bring in clean air. This can be seen with the column of low level of PM on 15 November in (a).



31



790

Figure 10. Surface $PM_{2.5}$ and wind field on 14 November in the Bay Area of observations (a) and S_CTRL predictions (b). Note that the reference wind vector for S_CTRL is 2 m s⁻¹ while the reference is 1 m s⁻¹ for observations. While the plume encroaches on the Bay Area, a strong sea breeze develops midday, driving plumes back inland. This sea breeze is not present in observational data, leading to a large underprediction of surface $PM_{2.5}$



Deleted: Figure 11. Vertical profile of $PM_{2.5}$ (a), time series of surface $PM_{2.5}$ (b), winds (c; observations in black and predictions in red) at Station 28 in the Bay Area. This station experienced different wind and PM evolution compared to Station 27 in Figure 10. The gray shading highlights the timeframe of greatest model bias of surface $PM_{2.5}$ Sharp increases in $PM_{2.5}$ correlate with a switch to northerly winds that import emissions to the Bay Area. Large negative $PM_{2.5}$ bias on 15 November occurs when S_CTRL deviates from observations and produces stronger southerly winds. This can be seen with the column of reduced particulate matter on 15 November in (a).

Fage Dieak
[3]
Deleted:Page Break
Formatted: Default Paragraph Font, Font color: Text 1, Kern at 12 pt







835

840

Figure 12. Comparison of meteorology generated by S_CTRL (solid red) and S_NOAERO (in which aerosol effects do not feed back to the meteorology, dashed blue) over the three areas of study: a) near the wildfire, b) Sacramento, and c) the San Francisco Bay Area. Exclusion of the aerosol feedback has the greatest effect nearest the fire, where S_NOAERO increased wind and temperature by 9.8% and 9.7%, respectively, on average. The aerosol feedback mechanism has the least significance in the Bay Area, where S_NOAERO wind speed differs less than 2% and temperature differs 3.1% on average. The most pronounced changes occur during 14-16 November when S_CTRL significantly underpredicts surface PM2.5. In WRF-Chem, the feedback of aerosol-radiation interactions on meteorology act to stabilize the atmosphere, slow wind speeds, and increase PM concentrations.

Moved (insertion) [6]



Figure <u>13</u>. Time series of surface PM_{2.5} (μg m⁻³) predicted by the sensitivity simulations (Table 1) averaged for the three areas of study: a) near the wildfire (N = 5), b) Sacramento (N = 7), and c) the Bay Area (N = 13). S_ENTR is omitted from the figure as it resulted in less than 1% change from S_CTRL. In the Bay Area, S_FCTX2 generally predicted more surface PM_{2.5}, recovering 10-35 µg m⁻³ 14-16 November when S_CTRL significantly underpredicts PM_{2.5} compared to observations. S_EMRAW demonstrates the impact of increasing the emissions inventory for 13-16 November. In the Bay Area, using the unperturbed emissions inventory reduces PM_{2.5} by more than 30% over 14-16 November. The impact of the aerosol feedback mechanism on PM_{2.5} (S_NOAERO) is location dependent. Excluding the feedback to meteorology generally reduces PM_{2.5} near the wildfire and in the Bay Area, while increasing PM_{2.5} in Sacramento. Employing the ACM2 PBL scheme results in a vastly different temporal evolution with a distinct diurnal pattern (S_LSM). FINN input fire data produces very little PM_{2.5}.

Deleted: 14

35



Figure 14. Comparison of surface PM_{2.5} (µg m⁻³) predicted by the joint perturbation experiment S_FCTX2andNOAERO with individual perturbation experiments (Table 1). ΔS * is the difference between each perturbation experiment (S_FCTX2 in dotted blue, S_NOAERO in dotted yellow, S_FCTX2andNOAERO in solid black) and the control experiment (S_CTRL). Black circles plot the sum of the effects from S_FCTX2 and S_NOAERO. Generally, the impact of the joint perturbation is similar to the sum of the two individual effects.

Page 31: [1] Deleted	Wang, Yuan	9/19/20 2:57:00 PM
▼		
Page 31: [2] Deleted	Wang, Yuan	9/19/20 2:57:00 PM
▼		
Page 32: [3] Deleted	Wang, Yuan	9/19/20 2:57:00 PM