

**The Effect of Meteorological Conditions and Atmospheric Composition in the  
Occurrence and Development of New Particle Formation (NPF) Events in  
Europe**

**Dimitrios Bousiotis<sup>1</sup>, James Brean<sup>1</sup>, Francis Pope<sup>1</sup>, Manuel Dall'Osto<sup>2</sup>  
Xavier Querol<sup>3</sup>, Andres Alastuey<sup>3</sup>, Noemi Perez<sup>3</sup>, Tuukka Petäjä<sup>4</sup>  
Andreas Massling<sup>5</sup>, Jacob Klenø Nøjgaard<sup>5</sup>, Claus Nørdestrom<sup>5</sup>  
Giorgos Kouvarakis<sup>6</sup>, Stergios Vratolis<sup>7</sup>, Konstantinos Eleftheriadis<sup>7</sup>  
Jarkko V. Niemi<sup>8</sup>, Harri Portin<sup>8</sup>, Alfred Wiedensohler<sup>9</sup>, Kay Weinhold<sup>9</sup>, Maik Merkel<sup>9</sup>,  
Thomas Tuch<sup>9</sup> and Roy M. Harrison<sup>1a\*</sup>**

**<sup>1</sup>Division of Environmental Health and Risk Management  
School of Geography, Earth and Environmental Sciences  
University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom**

**<sup>2</sup>Institute of Marine Sciences, Passeig Marítim de la Barceloneta, 37-49 E-08003  
Barcelona, Spain**

**<sup>3</sup>Institute of Environmental Assessment and Water Research (IDAEA - CSIC)  
08034, Barcelona, Spain**

**<sup>4</sup>Institute for Atmospheric and Earth System Research (INAR) / Physics, Faculty of Science  
University of Helsinki, Finland**

**<sup>5</sup>Department for Environmental Science, Aarhus University, DK-400, Roskilde, Denmark**

**<sup>6</sup>Environmental Chemical Processes Laboratory (ECPL), Department of Chemistry, University of  
Crete, 70013, Heraklion, Greece**

**<sup>7</sup>Environmental Radioactivity Laboratory, Institute of Nuclear and Radiological Science &  
Technology, Energy & Safety, NCSR Demokritos, Athens, Greece**

**<sup>8</sup>Helsinki Region Environmental Services Authority (HSY),  
FI-00066 HSY, Helsinki, Finland**

**<sup>9</sup>Leibniz Institute for Tropospheric Research (TROPOS),  
Permoserstr. 15, 04318 Leipzig, Germany**

**<sup>a</sup>Also at: Department of Environmental Sciences / Center of Excellence in Environmental Studies,  
King Abdulaziz University, PO Box 80203, Jeddah, 21589, Saudi Arabia**

## 42 **ABSTRACT**

43 Although new particle formation (NPF) events have been studied extensively for some decades, the  
44 mechanisms that drive their occurrence and development are yet to be fully elucidated. Laboratory  
45 studies have done much to elucidate the molecular processes involved in nucleation, but this  
46 knowledge has yet to be conclusively linked to NPF events in the atmosphere. There is great  
47 difficulty in successful application of the results from laboratory studies to real atmospheric  
48 conditions, due to the diversity of atmospheric conditions and observations found, as NPF events  
49 occur almost everywhere in the world without always following a clearly defined trend of  
50 frequency, seasonality, atmospheric conditions or event development. The present study seeks  
51 common features in nucleation events by applying a binned linear regression over an extensive  
52 dataset from 16 sites of various types (combined dataset of 85 years from rural and urban  
53 backgrounds as well as roadside sites) in Europe. At most sites, a clear positive relation is found  
54 between the solar radiation intensity (up to  $R^2 = 0.98$ ), temperature (up to  $R^2 = 0.98$ ) and  
55 atmospheric pressure (up to  $R^2 = 0.97$ ) with the probability of NPF events, while relative humidity  
56 (RH) presents a negative relation (up to  $R^2 = 0.95$ ) with NPF event probability. Wind speed  
57 presents a less consistent relationship which appears to be heavily affected by local conditions.  
58 While some meteorological variables (such as the solar radiation intensity and RH) appear to have a  
59 crucial effect on the occurrence and characteristics of NPF events, especially at rural sites, it  
60 appears that their role becomes less marked when at higher average values.

61

62 The analysis of chemical composition data presents interesting results. Concentrations of almost all  
63 chemical compounds studied (apart from O<sub>3</sub>) and the Condensation Sink (CS) have a negative  
64 relationship with NPF event probability, though areas with higher average concentrations of SO<sub>2</sub>  
65 had higher NPF event probability. Particulate Organic Carbon (OC), Volatile Organic Compounds  
66 (VOCs) and particulate phase sulphate consistently had a positive relation with the growth rate of  
67 the newly formed particles. As with some meteorological variables, it appears that at increased  
68 concentrations of pollutants or the CS, their influence upon NPF probability is reduced.

69

## 70 1. INTRODUCTION

71 New Particle Formation (NPF) events are an important source of particles in the atmosphere  
72 (Merikanto et al., 2009; Spracklen et al., 2010), which are known to have adverse effects on human  
73 health (Schwartz et al., 1996; Politis et al., 2008; Kim, et al., 2015) as well as affecting the optical  
74 and physical properties of the atmosphere (Makkonen et al., 2012; Seinfeld and Pandis, 2012).  
75 While they occur almost everywhere in the world (Dall'Osto et al., 2018; Kulmala et al., 2017;  
76 O'Dowd et al., 2002; Wiedensohler et al., 2019; Chu et al., 2019; Kerminen et al., 2018), with some  
77 exceptions mentioned in the literature in forest (Lee et al., 2016; Pillai et al., 2013; Rizzo et al.,  
78 2010) or high-elevation sites (Bae et al., 2010; Hallar et al., 2016), great diversity is found in the  
79 atmospheric conditions within which they take place. Many studies have been done in a large  
80 number of different types of locations (urban, traffic, regional background) around the world and  
81 differences were found in both the seasonality and intensity of NPF events. To an extent this  
82 variability is due to the mix of conditions that are specific to each location, which blurs the general  
83 understanding of the conditions that are favourable for the occurrence of NPF events (Berland et al.,  
84 2017; Bousiotis et al., 2020). For example, solar radiation is considered as one of the most  
85 important factors in the occurrence of NPF events (Kulmala and Kerminen, 2008; Kürten et al.,  
86 2016; Pikridas et al., 2015; Salma et al., 2011), as it is needed for the photochemical reactions that  
87 lead to the formation of sulphuric acid (Petäjä et al., 2009; Cheung et al., 2013). Sulphuric acid is  
88 considered as the main component of the formation and growth of the initial clusters (Iida et al.,  
89 2008; Stolzenburg et al., 2020; Weber et al., 1995). Nevertheless, in many cases NPF events did not

90 occur in the seasons with the highest insolation (Park et al., 2015; Vratolis et al., 2019). Similarly,  
91 uncertainty exists over the effect of temperature (Yli-Juuti et al., 2020; Stolzenburg et al., 2018).  
92 Higher temperatures are considered favourable for the growth of the newly formed particles as  
93 increased concentrations of both Biogenic Volatile Organic Compounds (BVOCs) and  
94 Anthropogenic Volatile Organic Compounds (AVOCs) (Yamada, 2013; Paasonen et al., 2013) and  
95 their oxidation products (Ehn et al., 2014) are associated to the growth of the particles. The negative  
96 effect of increasing temperatures in increasing the energy barriers the clusters have to overcome to  
97 become stable and grow in size though should not be overlooked (Kürten et al., 2018; Zhang et al.,  
98 2012). This appears to be true in most cases, as higher growth rates are found in most cases in the  
99 local summer (Nieminen et al., 2018), although the actual importance of those VOCs in the  
100 occurrence of NPF events is still not fully elucidated, with oxidation mechanisms still under intense  
101 research (Tröstl et al., 2016; Wang et al., 2020). The effect of other meteorological variables is even  
102 more complex, with studies presenting mixed results on the effect of the wind speed and  
103 atmospheric pressure. Extreme values of those variables may be favourable for the occurrence of  
104 NPF events, as they are associated with increased mixing in the atmosphere, but at the same time  
105 suppress due to increased dilution of precursors (Brines et al., 2015; Rimnácová et al., 2011; Shen  
106 et al., 2018; Siakavaras et al., 2016), or favour them due to a reduced condensation sink (CS).

107

108 The effect of atmospheric composition on NPF events is also a puzzle of mixed results. While the  
109 negative effect of the increased CS on the occurrence of the events is widely accepted (Kalkavouras

110 et al., 2017 ; Kerminen et al., 2004; Wehner et al., 2007), cases are found when NPF events occur  
111 on days with higher CS compared to average conditions (Größ et al., 2018; Kulmala et al., 2005).  
112 Sulphur dioxide (SO<sub>2</sub>), which is one of the most important contributors to many NPF pathways, in  
113 most studies was found in lower concentrations on NPF event days compared to average conditions  
114 (Alam et al., 2003; Bousiotis et al., 2019), although there are studies that have reported the opposite  
115 (Woo et al., 2001; Charron et al., 2008). Additionally, in a combined study of NPF events in China,  
116 events were found to be more probable under sulphur-rich conditions rather than sulphur-poor  
117 (Jayaratne et al., 2017). Similar is the case with the BVOCs and AVOCs, which present great  
118 variability depending the area studied (Dai et al., 2017), and their contribution in the growth of the  
119 particles is not fully understood yet. Until recently, it was considered unlikely for NPF events, as  
120 they are considered in the present study (deriving from secondary formation not associated with  
121 traffic related processes such as dilution of the exhaust), to occur within the complex urban  
122 environment due to the increased presence of compounds, mainly associated with combustion  
123 processes, which would suppress the survival of the newly formed particles within this type of  
124 environment (Kulmala et al., 2017). Despite this, NPF events were found to occur within even the  
125 most polluted areas and sometimes with high formation and growth rates (Bousiotis et al., 2019;  
126 Yao et al., 2018).

127

128 It is evident that while a general knowledge of the role of the meteorological and atmospheric  
129 variables has been achieved, there is great uncertainty over the extent and variability of their effect

130 (and for some of them even their actual effect) in the mechanisms of NPF in real atmospheric  
131 conditions, especially in the more complex urban environment (Harrison, 2017). The present study,  
132 using an extensive dataset from 16 sites in six European countries, attempts to elucidate the effect  
133 of several meteorological and atmospheric variables not only in general, but also depending on the  
134 geographical region or type of environment. While studies with multiple sites have been reported in  
135 the past (Dall'Osto et al., 2018; Kulmala et al., 2005; Rivas et al., 2020), to the authors' knowledge  
136 this is the first study that focuses directly on the effect of these variables upon the probability of  
137 NPF events as well as the formation and growth rates of newly formed particles in real atmospheric  
138 conditions.

139

## 140 **2. DATA AND METHODS**

### 141 **2.1 Site Description and Data Availability**

142 The present study uses a total of more than 85 years of hourly data from 16 sites from six countries  
143 of Europe of various land usage and climates. It was considered very important that at least a rural  
144 and an urban site would be available from each country to study the differences between the  
145 different land usage on NPF events throughout Europe. The sites were chosen to cover the greatest  
146 possible extent of the European continent, with sites from both northern, central and southern  
147 Europe, as well as from western and eastern. The sites are located in the UK (London and Harwell),  
148 Denmark (Copenhagen greater area), Germany (Leipzig greater area), Finland (Helsinki and  
149 Hyytiälä), Spain (Barcelona and Montseny – a site in a mountainous area) and Greece (Athens and

150 Finokalia). Unfortunately, not all sites had available data for all the variables studied, which to an  
151 extent may bias some of the results. An extended analysis of the typical and NPF event conditions,  
152 seasonal variations and trends at these sites for the same period is found in other studies (Bousiotis  
153 et al., 2019; 2020). A list of the available data and a brief description for each site is found in Table  
154 1 (for the ease of reading the sites are named by the country of the site followed by the last two  
155 letters which refer to the type of site, being RU for rural/regional background, UB for urban  
156 background and RO for roadside site), while a map of the sites is found in Figure 1.

157

## 158 **2.2 Methods**

### 159 **2.2.1 NPF events selection**

160 NPF events were selected using the method proposed by Dal Maso et al (2005). An NPF event is  
161 identified by the appearance of a new mode or particles in the nucleation mode (smaller than 20 nm  
162 in diameter), which prevails for some hours and shows signs of growth. The events can then be  
163 classified into classes I and II according to the level of certainty, while class I events can be further  
164 classified to Ia and Ib. Events having both a clear formation of a new mode of particles in the  
165 smallest size bins available (thus excluding possible advected events) as well as a distinct and  
166 persistent growth of the new mode of particles for at least 3 hours were classified as Ia, while Ib  
167 consists of rather clear events that fail though by at least one of the criteria set. Additionally, for the  
168 roadside sites, a formation of particles in the nucleation mode accompanied by a significant increase  
169 of the concentrations of pollutants was not considered as an NPF event, as it may be associated with



170 mechanisms other than the secondary formation. In the present study, only the events of class Ia  
171 were considered with the additional criterion of at least 1 nm h<sup>-1</sup> growth for at least 3 hours.

172

### 173 **2.2.2 Calculation of condensation sink, growth rate, formation rate, and NPF event** 174 **probability**

175 The condensation sink (CS) is calculated according to the method proposed by Kulmala et al.,  
176 (2001) as:

177

$$178 \quad CS = 4\pi D_{vap} \sum \beta_M r N \quad (1)$$

179

180 where r and N is the radius and number concentration of the particles respectively and D<sub>vap</sub> is the  
181 diffusion coefficient calculated as (Poling et al., 2001):

182

$$183 \quad D_{vap} = 0.00143 \cdot T^{1.75} \frac{\sqrt{M_{air}^{-1} + M_{vap}^{-1}}}{P \left( D_{x,air}^{\frac{1}{3}} + D_{x,vap}^{\frac{1}{3}} \right)^2} \quad (2)$$

184

185 for T = 293 K and P = 1013.25 mbar. M and D<sub>x</sub> are the molar mass and diffusion volume for air and  
186 sulphuric acid. β<sub>M</sub> is the Fuchs correction factor calculated as (Fuchs and Sutugin, 1971):

187

$$\beta_M = \frac{1 + K_n}{1 + \left(\frac{4}{3a} + 0.377\right) K_n + \frac{4}{3a} K_n^2} \quad (3)$$

where  $K_n$  is the Knudsen number, calculated as  $K_n = 2\lambda_m/d_p$  where  $\lambda_m$  is the mean free path of the gas.

Growth rate (GR) is calculated as (Kulmala et al., 2012):

$$GR = \frac{D_{P_2} - D_{P_1}}{t_2 - t_1} \quad (4)$$

for the size range between the minimum available particle diameter up to 30 nm (50 nm for the UK sites due to the higher minimum particle size available). The time window used for the calculation of the growth rate was from the start of the event until a) growth stopped, b) GMD reached the upper limit set or c) the day ended.

The formation rate  $J$  was calculated using the method proposed by (Kulmala et al., 2012) as:

$$J_{dp} = \frac{dN_{dp}}{dt} + \text{Coag}S_{dp} \times N_{dp} + \frac{GR}{\Delta d_p} \times N_{dp} + S_{losses} \quad (5)$$

206 where  $\text{CoagS}_{dp}$  is the coagulation rate of particles of diameter  $d_p$ , calculated as (Kerminen et al.,  
 207 2001):

208

$$209 \quad \text{CoagS}_{dp} = \int K(d_p, d'_p) n(d'_p) dd'_p \cong \sum_{d'_p=d_p}^{d'_p=\max} K(d_p, d'_p) N_{dp} \quad (6)$$

210

211  $K(d_p, d'_p)$  is the coagulation coefficient of particles with diameters  $d_p$  and  $d'_p$ , while  $S_{\text{losses}}$  accounts  
 212 for additional loss terms (i.e. chamber wall losses), which are not applicable in the present study.  
 213 For the present study, the formation rate of particles of diameter of 10 nm was calculated for  
 214 uniformity (16 nm for the UK sites), though most sites had data for particle sizes below 10 nm.  
 215

216 The NPF probability, used instead of NPF frequency when modelled results are presented, was  
 217 calculated by the number of NPF event days divided by the number of days with available data in  
 218 the given group (temporal, variable range etc.). The results presented in this study were normalised  
 219 according to the data availability, as:

220

$$221 \quad NPF_{\text{probability}} = \frac{N_{\text{NPF event days for group of days } X}}{N_{\text{days with available data for group of days } X}}$$

222

223

### 224 2.2.3 Calculation of the gradient and intercept for the variables used

225 Due to the large datasets available and the great spread of the values, a direct comparison between a  
226 given variable and any of the characteristics associated with NPF events (NPF probability, growth  
227 rate and formation rate) always provided results with low statistical significance. As a result, an  
228 alternative method which can provide a reliable result without the dispersion of the large datasets  
229 was used in the present study, to investigate the relationships between the variables which are  
230 considered to be associated with the NPF events. For this, a timeframe which is more directly  
231 associated with the NPF events typically observed in the mid-latitudes was chosen. For NPF  
232 probability and GR the timeframe between 05:00 to 17:00 Local Time (LT) was chosen, which is  
233 considered the time when the vast majority of NPF events take place and further develop with the  
234 growth of the particles. For the formation rate a smaller timeframe was chosen, 09:00 to 15:00 LT  
235 which is  $\pm 3$  hours from the time of the maximum formation rate found for almost all sites (12:00  
236 LT). This was done to exclude as far as possible the effect of the morning rush at the roadside sites,  
237 as well as only to include the time window when the formation rate is mostly relevant to NPF  
238 events (negative values that are more probable outside this timeframe and are not associated with  
239 the formation of the particles would bias the results).

240

241 For the CS the timeframe 05:00 to 10:00 LT was chosen. This was done to avoid including the  
242 direct effect of the NPF events (the contribution of newly formed particles to CS), as well as to  
243 provide results for the conditions which either promote or suppress the characteristics studied,

244 which specifically for the CS are more important before the start of the events. The extreme values  
245 (very high or very low) which bias the results only carrying a very small piece (forming bins of very  
246 small size) of information were then removed, though 90% of the available data was used for all the  
247 variables. The data left was separated into smaller bins and a minimum of 10 bins was required for  
248 each variable (for example if the difference between the minimum and the maximum relative  
249 humidity (RH) is 70%, then 14 bins each with a range of 5% were formed). The variables of interest  
250 were then averaged for each bin and plotted, and a linear relation was considered for each one of  
251 them.

252

253 The gradient of these linear relations ( $a_N$ ,  $a_G$  and  $a_J$  for NPF probability, growth rate and formation  
254 rate  $J_{10}$  accordingly) found in this analysis should be used with great caution as apart from the  
255 atmospheric conditions (local and meteorological as well as atmospheric composition) it is also  
256 affected by the variable in question (e.g. a greater NPF probability will provide a greater gradient),  
257 resulting in giving the same trend for all the atmospheric variables tested; the sites with the higher  
258 values of these variables (NPF probability and formation rate) always had greater gradient values  
259 and vice versa. In order to remove the effect of the variable in question (NPF probability or  
260 formation rate – growth rate will provide an unreliable result as it is calculated in a different range  
261 for each site due to the lower available size of particles), the gradients were normalised by dividing  
262 them by their respective variable (e.g. divide the gradient of the NPF probability with the NPF  
263 frequency), providing with a new normalised slope ( $a_N^*$  for NPF probability or  $a_J^*$  for the

264 formation rate) that will have no significance other than its absolute value, which can be used for  
265 direct comparisons:

266 
$$a_N^* = \frac{a_N}{\text{NPF \%}}$$

267 Where  $a_N$  is the gradient of the relation between the given variable and NPF frequency (NPF %)  
268

269 
$$a_J^* = \frac{a_J}{J_{10}}$$

270 Where  $a_J$  is the gradient of the relation between the given variable and the formation rate of 10 nm  
271 particles  $J_{10}$  ( $J_{16}$  for the UK sites).

272

### 273 3. RESULTS

274 In this study NPF events are generally observed as particles grow from a smaller size (typically 3-  
275 16 nm depending on the size detection limit of instruments used) to 30 nm or larger. They therefore  
276 reflect the result both of nucleation, which creates new particles of 1-2 nm (not detected with the  
277 instruments used in this study), and growth to larger sizes. In analysing NPF events, we therefore  
278 consider three diagnostic features:

- 279 • the probability of events occurring (i.e. days with an event divided by total days with relevant  
280 data, depending on the variable and range studied),  
281 • the rate of particle formation at a given size ( $J_{10}$  in this case),

282 • the growth rate of particles from the lower measurement limit to 30 nm (or 50 nm for the UK  
283 sites).

284

285 From the analysis of the extended dataset a total of 1952 NPF events were extracted and studied.

286 The NPF frequency, growth and formation rate for each site is found in Table 2. The seasonal

287 variation of NPF events is found in Figure S14.

288

### 289 **3.1 Meteorological Conditions**

290 The gradients, coefficients of determination ( $R^2$ ) and the p-values (deriving from one-way ANOVA

291 test) from the analysis of the meteorological variables, as well as the average conditions of these

292 variables are found in Table 3. The results for each site and variable are found in figures S1 – S5.

293

#### 294 **3.1.1 Solar radiation intensity**

295 As mentioned earlier, solar radiation intensity is considered to be one of the most important

296 variables in NPF occurrence, as it contributes to the production of  $H_2SO_4$  which is a main

297 component of the initial clusters and participates in the early growth of the newly formed particles.

298 Hidy et al. (1994) reported up to six times higher  $SO_2$  oxidation rates into  $H_2SO_4$  in typical summer

299 conditions compared to winter. For almost all sites this relation is confirmed with very strong

300 correlations ( $R^2 > 0.75$ ) between the intensity of solar radiation and the probability for NPF events

301 to occur. The relationship between the solar radiation and NPF probability was positive at all sites

302 and only three sites (FINUB, SPARU and GREUB) presented weak correlations ( $R^2 < 0.40$ ).  
303 Weaker correlations were found for the southern European sites, which might be associated with the  
304 higher averages for solar radiation intensity, or the interference of other processes (such as  
305 coinciding with increased CS by recirculation of air masses (Carnerero et al., 2019)), possibly  
306 making it less of an important factor for these areas.  
307  
308 The relationship of solar radiation with the growth rate was weaker in all cases and did not present a  
309 clear trend. Only some rural background sites (GERRU, FINRU and GRERU) presented a strong  
310 correlation ( $R^2 > 0.50$ ). The relationship found in most cases was positive apart from two roadside  
311 sites (GERRO and UKRO) and two urban background sites (GREUB and UKUB), though due to  
312 the low  $R^2$  ( $< 0.10$ ) these results cannot be considered with confidence. It seems though that the  
313 solar radiation intensity is probably a more important factor at background sites rather than at  
314 roadside sites, where possibly local conditions (such as local emissions) are more important (Olin et  
315 al, 2020). Finally, the formation rate has a positive relationship with the solar radiation intensity,  
316 with relatively strong correlations in most areas ( $R^2 > 0.50$ ). The correlations were stronger at the  
317 rural background sites compared to the roadside sites, which further underlines the increased  
318 importance of this factor at this type of site. A negative relationship between the solar radiation  
319 intensity and the formation rate was found at the GRERU site but the  $R^2$  is very low ( $R^2 = 0.05$ ).  
320



321 Plotting the normalised gradients for NPF event probability  $a_N^*$  with the average solar radiation  
322 intensity at each site (Figure 2) a negative relationship is found ( $R^2 = 0.62$ ), with the southern areas  
323 (those with higher average solar intensity) having smaller  $a_N^*$  compared to those in higher latitudes  
324 (and thus with a lower average solar radiation). This may indicate that while solar radiation is a  
325 deciding factor in the occurrence of an NPF event, when in greater intensity its role becomes  
326 relatively less important, a finding that was also implied by Wonaschütz et al. (2015). Additionally,  
327 the  $a_I^*$  was found to be higher at all rural sites compared to their respective roadside sites (and  
328 urban background sites for all but the Greek and German ones), making it a more important factor  
329 at this type of site (Figure 3).

330

### 331 **3.1.2 Relative humidity**

332 Relative humidity is considered to have a negative effect on the occurrence of NPF events (Jeong et  
333 al., 2010; Hamed et al., 2011; Park et al., 2015; Dada et al., 2017; Li et al., 2019). While water in  
334 the atmosphere is one of the main compounds needed for the formation of the initial clusters either  
335 on the binary or ternary nucleation theory (Henschel et al., 2016; Korhonen et al., 1999; Mirabel  
336 and Katz, 1974), under atmospheric conditions it may also play a negative role suppressing the  
337 number concentrations of new particles by increasing aerosol surface area (Li et al. 2019).  
338 Consistent with this, a negative relationship of the RH with NPF probability was found for all the  
339 sites of this study with very high  $R^2$  for almost all of them ( $R^2 > 0.80$ ). This is not simple to  
340 interpret as solar radiation intensity, temperature, RH and CS are not independent variables, since

341 an increase in temperature of an air mass due to increased solar radiation will be associated with  
342 reduced RH, which in turn affects the CS. The sites in Greece presented lower  $R^2$  compared to the  
343 other sites while, GRERU was found to have the weakest correlation ( $R^2 = 0.22$ ). This may be due  
344 to the different seasonality of the events found for the Greek sites (being more balanced within a  
345 year), as there was increased frequency of NPF events for the seasons with higher RH compared to  
346 other sites, making it a less important factor for their occurrence. Growth rate on the other hand had  
347 a variable relationship, either positive or negative, with only a handful of background sites having  
348 strong correlations. The German background sites as well as FINRU, which were among the sites  
349 with the highest average RH (average RH for GERRU is 81.9%, GERUB is 78.7% and FINUB is  
350 80.1%) presented a negative relationship between the RH and growth rate. DENRU (average RH at  
351 75.7%) had a positive relationship, which might indicate that the relationship between these two  
352 variables may vary depending upon the RH range. Formation rate also appears to have a negative  
353 relationship with the RH, though this relationship was significant ( $R^2 > 0.40$ ) for only 6 sites, which  
354 once again in most cases are sites with higher RH average conditions. Along with the results of the  
355 growth rate this might indicate that the RH becomes a more important factor in the development of  
356 NPF events as its values increase.

357

358 The normalised gradients once again provide some additional information. Regarding the NPF  
359 probability, it is found that the  $a_N^*$  was more negative at rural sites compared to roadside sites. This  
360 indicates that the RH has a smaller effect at roadside sites, as other variables, such as the

atmospheric composition, are probably more important within the complex environment in this type of site. Additionally, the relationship between  $a_N^*$  and average RH at the sites had a negative relationship ( $R^2 = 0.46$ ), which further shows that the RH becomes a more important factor at higher values (Figure 4). Furthermore, at the sets of rural and roadside sites with  $R^2$  higher than 0.40 for the relation between RH and the formation rate (UK and German sites), it was found that the  $a_J^*$  was more negative at the rural sites which indicates that the RH is a more important factor at rural sites compared to their respective roadside sites.

### 3.1.3 Temperature

Temperature can have both a direct and indirect effect in the development of NPF events, as it is directly associated with the abundance of both biogenic and anthropogenic volatile carbon, which is an important group of compounds whose oxidation products can participate in nucleation itself (Lehtipalo et al., 2018; Rose et al., 2018), as well as in the growth of newly formed particles. It may also have a negative effect on the particle size distributions or number concentrations through other processes such as particle evaporation. Most of the sites of the present study presented a strong relationship of NPF probability with temperature, which in most cases was positive, though in many cases (such as the Danish, Finnish and Spanish sites – figures S2b, d and e) there seems to be a peak in the NPF probability at some temperature, after which a decline starts (though being at the higher end does not greatly affect the results). Sites with smaller  $R^2$  (weaker association with temperature), were mainly those that have a seasonal variation that favoured seasons other than summer. These

381 sites not only had weaker relationship of NPF probability with temperature, but in most cases had a  
382 negative relationship (background sites in Finland, Spain and Greece). The Finnish sites, having the  
383 lowest average temperatures and a sufficient amount of data below zero temperature, show at all  
384 three sites the possible presence of a peak in the NPF event probability for temperatures below zero  
385 (Figure S2d). This seems to be the cause of the weak relations found there and they seem to be  
386 associated with the formation rate  $J_{10}$ , which also seems to have an increasing trend below zero  
387 degrees (Figure S2p). This may depend on the nucleation mechanism occurring, as cluster  
388 evaporation rates of sulphuric acid clusters are sensitive to the ternary stabilising compound present  
389 (Olenius et. al., 2017), as well as the possible enhancement of growth mechanisms at lower  
390 temperatures (below 5°C) by other chemical compounds in the atmosphere (i.e. nitric acid and  
391 ammonia) as found by Wang et al., (2020). Laboratory experiments show that the characteristics of  
392 organic aerosol forming from alpha-pinene is governed by gas phase oxidation (e.g. Ye et al. 2019).  
393 In the real atmosphere, the higher temperature enhances the amount of biogenic vapours (e.g.  
394 Paasonen et al. 2013) and, although the oxidation can be more efficient at higher temperatures, the  
395 lower temperatures favour formation of more non-volatile compounds (Quéléver et al., 2019;  
396 Stolzenburg et al. 2018; Ye et al. 2019).

397

398 Growth rate had a more uniform trend, with almost all sites having a positive relationship with  
399 temperature (apart from GERRO, though with  $R^2 = 0.00$ ). This relationship was very strong for  
400 most sites ( $R^2 > 0.60$  for 10 sites), which is also confirming the summer peak found for the growth

401 rate at most of these sites in other studies (Bousiotis et al., 2020; 2019). A rather strong relationship  
402 ( $R^2 > 0.50$ ) with temperature was also found for the formation rate for most sites, and was positive  
403 for almost all sites (apart from FINRO with  $R^2 = 0.01$  and the Greek sites with  $R^2 < 0.47$ ). As with  
404 the NPF probability, in general the sites with a seasonal variation of events that favoured summer  
405 had the strongest relationship (high  $R^2$ ) of the temperature with formation rate, which might  
406 indicate that this variable, either through its direct or indirect effect is an important one for the  
407 seasonal variability of NPF events in a given area.

408  
409 The normalised gradients for this variable did not present a clear trend among the areas studied,  
410 other than presenting greater  $a_N^*$  for the sites with a summer peak in their NPF event seasonal  
411 variation. As with other meteorological variables, the importance of this variable became smaller  
412 with increased values in the average conditions for both the NPF probability (Figure 5) and  $J_{10}$ ,  
413 though these relationships were not significant (biased by the very low average temperatures and  
414 different behaviour of the variables at the Finnish sites, without which the relation becomes a lot  
415 clearer as indicated in Figure S13). The variation though within the sites of the same area (different  
416 sites in same country / region) appears to directly follow the variability of temperature, showing that  
417 the temperature directly affects the occurrence of NPF events when other meteorological factors  
418 remain constant, having a negative trend for all countries but Finland. The  $a_j^*$  though is found to be  
419 greater (positively or negatively) at the rural background sites than at the other two types of sites at

all areas studied, showing that it is a more important factor for the formation rate at this type of site compared to others (Figure 6).

#### **3.1.4 Wind speed**

Wind speed may have both a positive and a negative effect on the occurrence of NPF events. On one hand, it may promote NPF events by the increased mixing of the condensable compounds in the atmosphere as well as by reducing the CS. On the other hand, high wind speeds may suppress NPF events due to increased dilution. It should be considered that the variability found is also affected by the specific conditions found at each site. The wind speed measurements in many cases, especially in urban sites, can be biased by the local topography or specific conditions found at each site, thus representing the local conditions for this variable rather than the regional ones. Similarly, measurements of wind speed at well sited meteorological stations may be more representative of regional conditions, than of those affecting the sites of nucleation measurement. The sites in this study presented mixed results, both in the importance as well as the effect of the wind speed variability. Three different behaviours were found in the variation of NPF event probability and wind speed which appear to be associated with local conditions as they are almost uniformly found among the sites within close proximity. Some sites presented a steady increase of NPF event probability with wind speed (Danish sites, UKUB, FINRU, SPAUB and GRERU), while others were found to steadily decline with increasing wind speeds (German sites – it should be noted that the German sites are the only ones that are located at a great distance from the sea), while some

440 were found to reach a peak and then decline, which also leads to smaller  $R^2$  (UKRU, UKRO,  
441 SPARU and to a lesser extent GREUB – figures S4a, e and f). The reasons for these differences  
442 between the sites are very hard to distinguish as apart from the wind speed the origin and the  
443 characteristics of these air masses play a crucial role. Following this, it appears that NPF probability  
444 is very low or zero for wind speeds close to calm for the sites with an increasing trend (as well as  
445 those that have a peak and decline after), while the opposite is observed for the German sites where  
446 the maximum NPF probability is found for very low wind speeds (fig. S4c).

447  
448 Similarly, the effect of different wind speeds upon the growth rate also varied a lot, though it was  
449 found to be negative in all the cases where  $R^2$  was higher than 0.50 (UKUB, DENRU, DENRO,  
450 GERRU, GERUB and GREUB). Finally, the formation rate was found to have a significant  
451 correlation ( $R^2 > 0.40$ ) only at two sites (UKRO and DENRU), probably indicating that the  
452 variability of the wind speed either does not affect this variable or its effect is rather small.

453  
454 The normalised gradients did not have any notable relation to either the NPF probability or the  
455 formation rate further confirming that the effect of the different wind speeds is not due to its  
456 variability only, but it is also influenced by the characteristics of the incoming air masses as well as  
457 specific local conditions found at each site.

458

459

### 460 3.1.5 Pressure

461 In almost all the sites with available data (apart from the Spanish), the NPF probability presented a  
462 positive relationship with high significance at all types of sites. The greater significance found at the  
463 rural sites (apart from SPARU) indicates the increased importance of meteorological conditions in  
464 the occurrence of NPF events at this type of site. The growth rate also presented a similar picture,  
465 with positive relationships at all the background sites of this study except the ones in Greece ( $R^2 >$   
466  $0.71$ ) and FINUB (though with low  $R^2$  at  $0.02$ ). This is probably associated with the seasonal  
467 variation found in Greece where higher growth rates were found in summer, a period when  
468 increased wind speeds and lower atmospheric pressure was found due to the Etesians, a pressure  
469 system that develops in the region every summer (Kalkavouras et al., 2017). An interesting finding  
470 is the negative gradients found at all the roadside sites, though the significance of these results is  
471 relatively low ( $R^2 < 0.43$ ) and always lower compared to the rural sites. The effects of pressure  
472 above are not likely to be important. Once again however, this is not an independent variable and  
473 higher pressure in summer tends to be associated with higher insolation and temperatures and lower  
474 RH. Since most events occur in the warmer months of the year, this is probably the explanation for  
475 the apparent effects of pressure. The formation rate presented relationships of low significance ( $R^2$   
476  $< 0.47$ ) for the sites of this study. Due to this, pressure should not be an important factor for the  
477 formation rate at any type of site.

478



479 The normalised gradients did not present any clear trends, even for the NPF probability for which  
480 the results presented significant relations at almost all sites.

481

## 482 **3.2 Atmospheric Composition**

483 The gradients,  $R^2$  and p-values from the analysis of a number of air pollutants ( $\text{SO}_2$ ,  $\text{NO}_x$ ,  $\text{O}_3$ ,  
484 organic compounds, sulphate and ammonia) and the CS, as well as the average conditions of these  
485 variables are found in Table 4. The results for each site and variable are found in Figures S6 – S12.

486

### 487 **3.2.1 Sulphur dioxide ( $\text{SO}_2$ )**

488 Sulphur dioxide, as a precursor of  $\text{H}_2\text{SO}_4$ , is considered as one of the main components associated  
489 with the NPF process. According to nucleation theories and observations,  $\text{H}_2\text{SO}_4$  is the most  
490 important compound from which the initial clusters are formed, as well as one of the candidate  
491 compounds for the initial steps of particle growth (Kirkby et al., 2011; Nieminen et al., 2010; Sipila  
492 et al., 2010; Stolzenburg et al., 2020). As  $\text{H}_2\text{SO}_4$  in the atmosphere is produced from oxidation  
493 reactions of  $\text{SO}_2$  it would be expected that increased concentrations of the latter would be associated  
494 with increased values for all the variables associated with the NPF process. Contrary to this though,  
495 the relationship of  $\text{SO}_2$  concentrations with NPF probability was found to be negative at all the sites  
496 in this study with available data. This relationship was relatively strong ( $R^2 > 0.50$ ) in most areas  
497 with an increased significance at roadside sites compared to their respective rural sites. As this is a  
498 negative relationship, this may indicate that  $\text{SO}_2$  is in sufficient concentrations for  $\text{H}_2\text{SO}_4$  formation,

499 thus not suppressing the occurrence of NPF events, as well as showing that in increased  
500 concentrations, it is a more important factor (or surrogate for a factor) in preventing the occurrence  
501 of NPF events within the urban environment, as higher SO<sub>2</sub> is likely associated with increased co-  
502 emitted particle pollution and hence CS. The growth rate on the other hand, presented mixed results  
503 and the significance of the relationships is low in most cases, which makes these results unreliable.  
504 Finally, the relationship of SO<sub>2</sub> concentrations with the formation rate was found to be positive at  
505 all sites but SPARU and FINRU (which had the lowest concentrations across the sites with  
506 available data). The significance of this relationship was rather low ( $R^2 < 0.40$ ) for all but the  
507 roadside sites. This suggests that higher H<sub>2</sub>SO<sub>4</sub> concentrations favour greater formation rates (i.e.  
508 more particles can be formed), rather than necessarily promoting nucleation itself because of the  
509 competing effect of condensation onto the pre-existing particle population.

510  
511 The normalised gradients  $a_N^*$  were found to be more negative at the background sites compared to  
512 their respective roadside sites, as well as being less negative in the UK (where SO<sub>2</sub> is in greater  
513 abundance) compared to the other sites with relatively significant relationships. Plotting the average  
514 SO<sub>2</sub> concentrations with the normalised gradients  $a_N^*$  for the all sites (though not all had significant  
515 relations), a positive relationship with relatively high  $R^2$  (when the extreme values from  
516 Marylebone Road-UKRO are removed) is found which might indicate that while increased  
517 concentrations are a negative factor in NPF event occurrence at a given site, in general the sites with  
518 higher SO<sub>2</sub> concentrations on average present higher probability for NPF events (Figures 7a and

7b). This appears to be in agreement with Dall'Osto et al. (2018) who discussed the variable role of SO<sub>2</sub> depending on its concentrations. No significant relations were found for the values of  $a_j^*$  as in most cases these relationships were rather weak.

### 3.2.2 Nitrogen oxides or nitrogen dioxide (NO<sub>x</sub> or NO<sub>2</sub>)

NO<sub>x</sub> and NO<sub>2</sub> are directly associated with pollution, which can be a limiting factor for NPF events as it increases the CS and may suppress the events (An et al., 2015), though with the reduction of SO<sub>2</sub> concentrations achieved the last couple of decades, there is a possibility for oxidation products of NO<sub>x</sub> to become an important component for NPF (Wang et al., 2020). For almost all sites (apart from GRERU) with available data a negative relationship between the NPF probability and NO<sub>x</sub> concentrations (or NO<sub>2</sub> depending on the available data) was found. Similarly, for all the sites but SPARU and GRERU, the correlations were strong with  $R^2 > 0.43$ . The rural background sites had a weaker relationship between the two variables compared to the urban sites, which is probably associated with them having rather low concentrations and variability of NO<sub>x</sub> (or NO<sub>2</sub>), making the variations of this factor less important. Growth rate had weaker correlations with NO<sub>x</sub> and different trends between the sites, either being positive or negative. The variable effect of NO<sub>x</sub> on particle growth, shifting HOMs volatility, was previously discussed by Yan et al. (2020). While variability was found for the background sites, all roadside sites regardless of the strength of the relationship had a positive relation between NO<sub>x</sub> and the growth rate. This may indicate the different components associated with the growth process at each type of site which, as found in other studies,

539 can be related to compounds associated with combustion processes that take place within the urban  
540 environment (Guo et al., 2020; Wang et al., 2017a). The formation rate presents few cases of strong  
541 relationships, with variable trends (positive and negative). While much effort was made to isolate  
542 the effect of NPF events by taking a shorter time frame before the event, the effect of local pollution  
543 is still included, especially at the urban sites (which probably explains the positive effect found).

544

545 The normalised gradients do not provide a significant result for the relationship of this variable with  
546 either the probability of the events or the formation rate. The only noteworthy points are the more  
547 negative  $a_N^*$  at the rural background sites compared to the roadside sites in all the areas studied,  
548 which shows the increased importance of a clean environment for NPF events to occur in areas  
549 where condensable compounds are in lesser abundance, such as a rural environment. Additionally,  
550 the negative gradients found at all the roadside sites, which increases the confidence that the events  
551 extracted at the roadside sites are not pollution incidents but NPF events. However, it appears that  
552 traffic pollution favours higher particle growth rates, although the components responsible for this  
553 effect are unknown.

554

### 555 3.2.3 Ozone (O<sub>3</sub>)

556 Ozone is typically the result of atmospheric photochemistry and is itself a source of hydroxyl  
557 radical through photolysis, or ozonolysis of alkenes both during daytime and night-time (Fenske et  
558 al., 2000). It might therefore be expected to act as an indicator of photochemical activity which

559 promotes the oxidation of SO<sub>2</sub> and VOCs. Ozone concentrations may be directly related to the  
560 solar radiation intensity as well as the pollution levels in the area studied, and O<sub>3</sub> is considered as a  
561 positive factor in the occurrence of NPF events (Woo et al., 2001; Berndt et al., 2006). As with the  
562 solar radiation intensity, there is a strong relationship between O<sub>3</sub> concentration and the probability  
563 for NPF events. This positive relationship was found to be stronger for the sites in northern Europe  
564 ( $R^2 > 0.51$ ), while it was not significant ( $R^2 < 0.38$ ) for the sites in southern Europe (Spanish sites  
565 and GRERU), possibly indicating that O<sub>3</sub> is a less important factor at the southern sites. Specifically  
566 for the Spanish sites which have the highest average concentrations of O<sub>3</sub> with some extreme values  
567 (Querol et al., 2017), the relationship of O<sub>3</sub> concentrations with the NPF probability presents a  
568 unique trend (Figure S8d), having a clear peak then a steady decline at both sites (though at  
569 different O<sub>3</sub> concentrations), which is also responsible for the low correlations found (this trend  
570 seems to also occur at SPARU for the growth rate and to a lesser extent for the formation rate as  
571 well, though for different O<sub>3</sub> concentration ranges – figures S8i and n). The specific variability  
572 found at the Spanish sites was also studied by Carnerero et al., (2019). For sites with a marked  
573 seasonal variation in ozone, associations with NPF may be artefactual due to correlations with other  
574 variables such as temperature, RH and solar radiation intensity.

575  
576 Unlike the solar radiation intensity though, the growth rate presents a negative relationship at the  
577 sites where the relationship between these two variables was significant (UKRU, UKUB, DENUB  
578 and FINRU), which might either be an indication of a polluted background that may have a

negative effect in the growth of the newly formed particles (though the trends found for  $\text{NO}_x$  indicate differently) or specific chemical processes which cannot be identified due to the lack of detailed chemical composition data. A significant relationship between  $\text{O}_3$  and the formation rate was only found for two sites (UKRO and DENRO, though the trends become a lot clearer if some values are removed from the extreme lower or higher end). This way the relationships become strong, but positive, for some areas and negative for some others without any clear trend (type or location of the site,  $\text{O}_3$  concentrations etc.). No clear relationship between these two variables was found as the sites with strong relationship have both positive (DENRO) and negative (UKRO) relationships and as a result no confident conclusions can be drawn.

As the correlations found were strong the normalised gradients for NPF probability, when plotted against the average concentrations of  $\text{O}_3$ , present a negative correlation with relatively high  $R^2$  (0.64), indicating that the  $\text{O}_3$  is a more important factor in the occurrence of NPF events when in lower concentrations (Figure 8). Finally, though with a low level of confidence for the southern sites, the  $a_N^*$  were smaller at the southern sites compared to those in the north, up to one order of magnitude between FINRU (furthest north rural background) and GRERU (furthest south rural background).

### 599 3.2.4 Organic compounds

#### 600 3.2.4.1 Particulate organic carbon (OC)

601 Organic carbon (OC) compounds in the secondary aerosol typically enter the particles via  
602 condensational processes, with a role that becomes increasingly important as the size of the  
603 particles becomes larger (Nieminen et al., 2010; Zhang et al., 2012; Shrivastava et al., 2017).  
604 Particulate OC, the data for which is available in the present study, can be associated with pollution,  
605 especially in the urban environment. Only a few of the sites of the present study were found to have  
606 a relatively strong negative relationship ( $R^2 > 0.50$ ) of particulate OC with the NPF probability  
607 (UKUB, UKRO and DENRU). Regardless though of the strength of this relationship, all other sites  
608 (apart from FINRU) had a negative relationship between these two variables as well, consistent  
609 with increased concentrations of particulate OC being associated with increased pollution, which  
610 elevate the CS, suppressing the occurrence of NPF events. Growth rate on the other hand was found  
611 to have a positive relationship ( $R^2 > 0.40$ ) for most of the sites. This relationship appeared to be  
612 stronger (higher  $R^2$ ) at the roadside sites with available data compared to their respective rural  
613 background sites. The relationship between particulate OC and the growth rate was positive at all  
614 the sites with available data regardless of their significance showing that, despite its effect in the  
615 occurrence of NPF events, it is still a favourable variable for the growth of the particles. The  
616 formation rate was found to have a significant relationship with particulate OC concentrations at  
617 half of the sites with available data (UKUB, UKRO, DENRU, DENRO).

618

619 The normalised gradients for this variable did not present any noteworthy relations with either the  
620 type of site or the concentrations of OC at a given site.

621

#### 622 **3.2.4.2 Volatile organic compounds (VOCs)**

623 Many volatile organic compounds have been found to be associated with the NPF process. Benzene,  
624 toluene, ethylbenzene, m-p-xylene, o-xylene and trimethylbenzenes have been reported to be able to  
625 form Highly Oxygenated Organic Molecules (HOMs) in flow tubes (Wang et al., 2017a; Molteni et  
626 al., 2018), which may act as contributors to particle nucleation and/or growth. Xylenes, and to a  
627 lesser extent trimethylbenzenes, are the most efficient at forming HOMs. Benzene and toluene are  
628 less efficient and will form more volatile HOMs. These HOMs may all be too volatile to form new  
629 particles, though this is not yet confirmed. Chamber studies involving H<sub>2</sub>SO<sub>4</sub> and trimethylbenzene  
630 oxidation products were associated with high formation rates when measuring J<sub>1.5</sub> (Metzger et al.,  
631 2010). All these HOMs though will be sufficiently involatile to contribute to particle growth. Those  
632 with higher oxygen content or carbon number will be classed as LVOC and if they dimerise, they  
633 will form ELVOC (Bianchi et al., 2019). Monoterpenes can also form HOMs which drive both the  
634 formation (Ehn et al., 2014; Riccobono et al., 2014) and growth (Tröstl et al., 2016), while isoprene  
635 can act as a sink for hydroxyl radical (Kiendler-Scharr et al., 2009) and is not as effective in HOM  
636 and secondary organic aerosol formation compared to monoterpenes (McFiggans et al., 2019).

637



638 Volatile organic compound data were available for three of the sites of this study (Table S2). Two  
639 of the sites with VOC data were from the rural background and the roadside site in the UK. Most of  
640 the compounds are associated with combustion sources and were found to have a negative  
641 relationship with NPF event occurrence at both sites, with high  $R^2$  ( $R^2 > 0.50$ ) in most cases.  
642 Additionally, isoprene, which may have either biogenic or anthropogenic sources (Wagner and  
643 Kuttler, 2014) was also found to have a negative relationship with NPF event occurrence at  
644 Marylebone Road-UKRO, though with low  $R^2$  (0.07). This result is in line with the VOCs being  
645 strongly correlated with particulate OC (which presented a negative relationship with NPF event  
646 probability, as discussed in Section 3.2.4.1), as well as with the CS (which also presented a negative  
647 relationship with NPF event probability, as mentioned in Section 3.2.6), further associating these  
648 compounds with combustion emissions.

649

650 Growth rate was found to have a positive relationship with VOCs in almost all cases for both UK  
651 sites. Few exceptions were found (with only 1,3 butadiene having a relatively high  $R^2$ ) which  
652 presented a negative relationship with the growth rate in rural Harwell-UKRU. Finally, the  
653 formation rate presented a different behaviour between the two sites. At UKRU, the relationship  
654 was unclear in most cases, with a group of VOCs presenting a negative relationship with the  
655 formation rate (ethane, ethene, propane, 1,3 butadiene, toluene, ethylbenzene, o-xylene and 1,2,4  
656 trimethylbenzene – with  $R^2 > 0.40$ ), two VOCs presented a rather clear positive relationship with  
657 the formation rate (iso-pentane and 2-methylbenzene) and the rest of the VOCs had an unclear

658 relationship. At UKRO though, VOCs presented a positive relationship with the formation rate (for  
659 particles of diameter 16 nm). This is probably due to the fact that these VOCs are associated with  
660 pollution emissions (as mentioned earlier) and though a smaller time window was chosen to avoid  
661 including the effect of the morning rush hour traffic, this is very difficult in the traffic polluted  
662 environment of Marylebone Road.

663  
664 As Hyytiälä (FINRU) is a rural background site far from the direct effect of combustion emissions,  
665 different VOCs were measured, which mainly originate from biogenic sources rather than  
666 anthropogenic ones. The results were mixed and less clear compared to those from the UK sites  
667 (mainly due to the smaller dataset), and three groups were found depending on their relationship  
668 with NPF probability. The first group, including acetonitrile, acetic acid and methyl ethyl ketone  
669 (MEK) presented a slight positive relationship. The second group presented a negative relationship,  
670 with the VOCs in this group being monoterpenes, methacroleine, benzene, isoprene and toluene  
671 (only the last two have  $R^2 > 0.50$ ). Finally, the third group included VOCs that presented a peak and  
672 then a decline for higher concentrations including methanol, and acetone. Two groups of VOCs  
673 were found depending on their relationship with the growth rate. The ones with a positive  
674 relationship being methanol, acetonitrile, acetone, acetic acid, isoprene, methacroleine,  
675 monoterpenes and toluene, while acetaldehyde, MEK and benzene had a negative relationship, with  
676 relatively high  $R^2$  in most cases. Finally, the results with the formation rate were unclear with only a  
677 handful presenting weak ( $R^2 < 0.21$ ) positive (methanol, acetic acid and benzene) or negative

678 (MEK) relationships that do not appear to be significant. The normalised gradients cannot be used  
679 for VOCs as there are very few sites with available data.

680

### 681 **3.2.5 Sulphate ( $\text{SO}_4^{2-}$ )**

682 Sulphate ( $\text{SO}_4^{2-}$ ) is a major secondary constituent of aerosols. Secondary  $\text{SO}_4^{2-}$  aerosols largely arise  
683 from either gas phase reaction between  $\text{SO}_2$  and OH, or in the aqueous phase by the reaction of  $\text{SO}_2$   
684 and  $\text{O}_3$  or  $\text{H}_2\text{O}_2$ , or  $\text{NO}_2$  (Hidy et al., 1994). In environments where  $\text{SO}_4^{2-}$  chemistry is dominant  
685 (i.e. remote areas),  $\text{SO}_4^{2-}$  and ammonium (bi) sulphate ( $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{HSO}_4$ ) particles are a  
686 large relative contributor to aerosol mass, while this contribution is lower in environments where  
687 other emissions are also significant (i.e. urban areas where the secondary  $\text{NO}_3^-$  relative contribution  
688 is a lot higher). While not well established, a possible relationship of  $\text{SO}_4^{2-}$ -containing compounds  
689 and variables of NPF events was found in previous studies (Beddows et al., 2015; Minguillón et al.,  
690 2015; Wang et al., 2017b). In the present study, only a few sites had  $\text{SO}_4^{2-}$  data available, for  $\text{PM}_1$   
691 ( $\text{FINRU}$ ),  $\text{PM}_{2.5}$  (Danish sites) or  $\text{PM}_{10}$  (rest of the sites). While this data cannot be considered as  
692 directly associated with the ultrafine particles, for two sites with available AMS data for ultrafine  
693 particles, the direct comparison between  $\text{SO}_4^{2-}$  aerosol in PM and in the range of particles of about  
694 50 nm, very high correlations were found (results not included). For all the sites with available data  
695 the NPF probability presented a negative relationship. The significance of this relationship was  
696 found to be relatively high ( $R^2 > 0.50$ ) only for background sites (apart from GERRU, which has  
697 rather low concentrations and probably different mechanisms for the NPF events). Similarly, the

698 growth rate presented a significant relationship ( $R^2 > 0.40$ ) for the same background sites (apart  
699 from FINRU), though this relationship was found to be positive at all sites regardless of its  
700 significance. Finally, the formation rate did not present a clear trend as it was found to have both  
701 negative and positive relationships for different sites. This relationship was significant only for two  
702 rural sites (UKRU and DENRU) and as a result no conclusions can be reached.

703

704 The normalised gradients cannot be used for any analysis on sulphate as the measurements available  
705 are from different particle size ranges.

706

### 707 **3.2.6 Gaseous ammonia ( $\text{NH}_3$ )**

708 Ammonia ( $\text{NH}_3$ ) can be an important compound in the nucleation process according to the ternary  
709 theory (Kirkby et al., 2011; Napari et al., 2002). It was found that elevations in  $\text{NH}_3$  concentrations  
710 can lead to elevations to NPF rate (Lehtipalo et al., 2018) and it was also found to be an important  
711 factor for NPF event occurrence even when stronger bases are present in high concentrations  
712 (Glasoe et al., 2015). No significant variation was found though between event and non-event days  
713 in a previous study in Harwell - UKRU (Bousiotis et al., 2019). Data for gaseous ammonia was only  
714 available for UKRU and presented a positive relationship with NPF probability, until reaching a  
715 peak point. Further increase in  $\text{NH}_3$  concentrations presented a decline with NPF probability (Figure  
716 S11a), which might be due to its association with increased pollution levels. It presented a clear  
717 positive relationship with both the growth rate (though it also appears to decline at high

718 concentrations) and the formation rate, consistent with its well-established role in accelerating both  
719 of these processes (Kirkby et al. 2011; Stolzenburg et al., 2020).

720

### 721 **3.2.7 Condensation sink (CS)**

722 The CS is a measure of the rate at which molecules will condense onto pre-existing aerosols  
723 (Lehtinen et al., 2003). It is highly dependent on the number and size of the particles in the  
724 atmosphere and as a result it is expected to be affected by both the local emissions within the urban  
725 environment as well as the formation and growth of the particles due to NPF events. As a result, for  
726 the specific metric a time frame before the events are in full development was chosen (05:00 to  
727 10:00 LT) to avoid including the effect of the NPF events and provide a picture of the atmospheric  
728 conditions that preceded the NPF events. With this data, the NPF probability presented very strong  
729 relationships with the condensation sink. Two groups of sites were found though; those which had a  
730 positive relationship and those with a negative relationship. In the first group are the sites in  
731 Germany and Greece while all others had a negative relationship. This grouping follows the trend  
732 between the countries, the sites of which presented a greater or smaller CS on NPF event days  
733 (having positive or negative gradients respectively), though it is unknown what causes this  
734 behaviour (at the German sites and GREUB it may be associated with the very high formation rates  
735 on NPF event days). While the gradients from this analysis cannot be used for direct comparisons, a  
736 trend was found for which the gradients were more positive or negative at the rural sites compared

737 to their respective roadside sites, which might indicate the greater importance of the variability of  
738 the CS at the rural sites in the occurrence of NPF events.

739

740 The growth rate was positively correlated with the CS for most of the sites, with strong  
741 relationships ( $R^2 > 0.40$ ) for about half of them. As the CS is a metric of pre-existing particles, it is  
742 also associated with the level of pollution in a given area. The increased significance and gradient  
743 found at the rural sites probably indicates the importance of enhanced presence of condensable  
744 compounds in a cleaner environment, which in many cases are associated with the moderate  
745 presence of pollution. The formation rate was also found to have a positive relationship with the CS.  
746 This relationship was more significant at the roadside sites of this study, a result which to some  
747 extent is biased by the presence of increased traffic emissions found in the timeframe chosen. While  
748 to an extent, increased presence of condensable compounds can be favourable for greater formation  
749 rates, this result should be considered with great caution.

750

751 The normalised gradients  $a_N^*$  followed a similar trend as those found with the initial analysis. These  
752 gradients were found to be more positive or negative, depending on the trend of the given area, at  
753 the rural sites compared to their roadside sites. The urban background sites did not always have a  
754 uniform behaviour (though in UK, Denmark and Finland these were between the rural site and the  
755 roadside site), due to their more diverse character compared to the other two types of sites.

756

### 757 3.3 Association of the Effect of the Variables

758 The Pearson correlation coefficients for the variables studied on each site are found in Table S1.

759 The relatively strong relationship between the solar radiation intensity, temperature and O<sub>3</sub> found,  
760 as well as their anticorrelation with the RH may lead to the conclusion that not all these factors play  
761 a role in NPF events, but their visible effect is the result of their relationship with each other. There  
762 is a similar case with the association of the CS and NO<sub>x</sub> (or NO<sub>2</sub>), and OC, as well as SO<sub>2</sub>,  
763 especially at urban sites. However, the factors affect different outcomes differently, as for example  
764 the solar radiation intensity does not seem to be as important a factor for the growth rate as  
765 temperature, or O<sub>3</sub> does not seem to be strongly associated with either the formation or the growth  
766 rate. This is further established by the fact that some of these variables do not correlate well at the  
767 southern sites, but still appear to be associated with either the probability of NPF events or the  
768 growth or nucleation rate. The effects of all of these factors have been demonstrated in both  
769 laboratory and atmospheric studies in the past and were discussed earlier in this paper. By the  
770 analysis provided in the present study, the effect of each of these variables is further established,  
771 providing an association of each one of these variables with either the formation or the growth  
772 mechanism. However, RH does not seem to be a consistent factor in any mechanism, and it appears  
773 that its effect is dependent on location specific conditions, although it was the variable with the  
774 most consistent relation with NPF event probability at almost all sites.

775

776

### 777 3.4 Relationship to a previous multi-station European study

778 The findings of our study in respect of the background sites show many similarities with the  
779 conclusions drawn in the previous multi-station study in Europe by Dall'Osto et al. (2018) despite  
780 the two studies using several different sampling stations as well as some in common. Both studies  
781 point towards the influence of variables such as solar radiation intensity and CS upon the  
782 occurrence of NPF events. The previous study suggested that different compounds participate in the  
783 growth of the particles, depending on the area considered. Thus, for northern and southern sites the  
784 growth of the particles is suggested to be driven mainly by organic compounds, while for the sites  
785 in central Europe sulphate plays a more important role. These findings are confirmed by the present  
786 study, as the growth rate was found to correlate better with organic compounds for the rural sites in  
787 Finland and Greece, while  $\text{SO}_4^{2-}$  presented a stronger relationship with the growth rate for the  
788 Danish and German sites (the latter presented high gradient values but low  $R^2$  due to a decline at  
789 higher  $\text{SO}_4^{2-}$  concentrations – figure S10i, probably associated with NPF events being suppressed  
790 by increased pollution). The growth of the particles at the rural background site in the UK,  
791 characterised as “Overlap” in the previous study, was found to be strongly associated with both  
792 organic compounds and sulphate, consistent with it being in the central group.

793

794 The seasonality of NPF events at northern sites was hard to explain in the previous study, and the  
795 possible effect of low temperature was considered. In the present study, the Finnish background  
796 sites presented a double-peak relationship of NPF probability with temperature, with one of the



797 peaks being below zero degrees. This might point to the possibility of different compounds driving  
798 the events for different temperature ranges, as well as the increased nucleation rate of H<sub>2</sub>SO<sub>4</sub> at  
799 lower temperatures (Kirkby et al., 2011; Yan et al., 2018), which makes the occurrence of NPF  
800 events more probable at lower temperatures in a region with low SO<sub>2</sub> concentrations.

801

#### 802 **4. CONCLUSIONS**

803 The present study attempts to explain the effect of several meteorological and atmospheric variables  
804 on the occurrence and development of NPF events, by using a large-scale dataset. More than 85  
805 site-years of data from 16 sites from six countries in Europe were analysed for NPF events. A total  
806 of 1952 NPF events with consequent growth of the newly formed particles were extracted and with  
807 the use of binned linear regression, the relationship between three variables associated with NPF  
808 events (NPF event probability, formation and growth rate) with meteorological conditions and  
809 atmospheric composition was studied. Among the meteorological conditions, solar radiation  
810 intensity, temperature and atmospheric pressure presented a positive relationship with the  
811 occurrence of NPF events, either promoting the formation or growth rate. RH presented a negative  
812 relationship with NPF event probability which in most cases was associated with it being a limiting  
813 factor on particle formation at higher average values. Wind speed on the other hand presented  
814 variable results, appearing to depend on the location of the sites rather than their type. This shows  
815 that while wind speed can be a factor in NPF event occurrence, the origin of the incoming air  
816 masses also plays a very important role. In most cases, meteorological conditions, such as

817 temperature or RH appeared to be more important factors in NPF event occurrence at rural sites  
818 compared to urban sites, suggesting that NPF events are driven more by them at this type of site  
819 compared to urban environments and the more complex chemical interactions found there.  
820 Additionally, while some meteorological variables appeared to play a crucial role in the occurrence  
821 of NPF events, this role appears to become less important at higher values when a positive relation  
822 was found (or lower when a negative relation was found).  
823  
824 The results for the levels of atmospheric pollutants presented a more interesting picture as most of  
825 these, which appear to be either directly or indirectly associated with the NPF process were found to  
826 have negative relationships with NPF probability. This is probably due to the fact that increased  
827 concentrations of such compounds are associated with more polluted conditions, which are a  
828 limiting factor in the occurrence of NPF events, as was found with the negative relationship  
829 between the CS and NPF probability in most cases. Thus, SO<sub>2</sub>, NO<sub>x</sub> (or NO<sub>2</sub>), particulate OC and  
830 SO<sub>4</sub><sup>2-</sup> concentrations were negatively correlated with NPF probability in most cases. Average SO<sub>2</sub>  
831 concentrations appeared to correlate positively with the normalised NPF event probability gradients  
832 with a relatively significant correlation, indicating that while increasing concentrations have a  
833 negative impact in the occurrence of NPF events at a given site, in general sites with higher SO<sub>2</sub>  
834 concentrations have higher probability for NPF events. Conversely, these compounds in many cases  
835 had a positive relationship (not always though with high significance) with the other variables  
836 considered. Thus, particulate OC (and VOCs where data was available) and SO<sub>4</sub><sup>2-</sup> consistently had a

837 positive relationship with the growth rate, while SO<sub>2</sub> was positively associated with both the  
838 formation and growth rate in most cases. Finally, O<sub>3</sub> was positively correlated with NPF event  
839 probability at all sites in this study, though it presented variable results with the other two variables.  
840 As with some meteorological conditions it was found that at sites with increased concentrations of  
841 O<sub>3</sub>, its importance as a factor was decreased, which to some extent can be related with the high CS  
842 associated with peak summer O<sub>3</sub> days in southern Europe.

843

844 It should be noted that the variables considered are in many cases inter-related (e.g. temperature and  
845 RH) and this considerably complicates the interpretation in terms of causal factors. Large datasets  
846 are very useful in providing more uniform results by removing the possible bias of short period  
847 extremities, which may lead to wrong assumptions. This study, apart from providing insights into  
848 the effect of a number of variables on the occurrence and development of NPF events in  
849 atmospheric conditions across Europe, also shows the differences that climatic, land use and  
850 atmospheric composition variations cause to those effects. Such variations are probably the cause of  
851 the differences found among previous studies. Following from this, the importance of a high-  
852 resolution measurement network, both spatially and temporally is underlined, as it can help in  
853 elucidating the mechanisms of new particle formation in the real atmosphere.

854

855

856

857 **DATA ACCESSIBILITY**

858 Data supporting this publication are openly available from the UBIRA eData repository at  
859 <https://doi.org/>  
860

861 **AUTHOR CONTRIBUTIONS**

862 The study was conceived and planned by RMH who also contributed to the final manuscript, and  
863 DB who also carried out the analysis and prepared the first draft of the manuscript. AM, JKN, CN,  
864 JVN, HP, NP, AA, GK, SV and KE have provided with the data for the analysis. JB provided help  
865 with analysis of the data. FDP provided advice on the analysis. MDO, XQ and TP contributed to the  
866 final manuscript.

867  
868 **COMPETING INTERESTS**

869 The authors have no conflict of interests.

870  
871 **ACKNOWLEDGMENTS**

872 This work was supported by the National Centre for Atmospheric Science funded by the U.K.  
873 Natural Environment Research Council (R8/H12/83/011).

874

## 875 REFERENCES

- 876
- 877 Aalto, P., Hämeri, K., Becker, E. D. O., Weber, R., Salm, J., Mäkelä, J. M., Hoell, C., O'Dowd, C.  
 878 D., Karlsson, H., Hansson, H., Väkevä, M., Koponen, I. K., Buzorius, G. and Kulmala, M.: Physical  
 879 characterization of aerosol particles during nucleation events, *Tellus, Ser. B Chem. Phys. Meteorol.*,  
 880 53(4), 344–358, doi:10.3402/tellusb.v53i4.17127, 2001.
- 881
- 882 Alam, A., Shi, J. P. and Harrison, R. M.: Observations of new particle formation in urban air, *J.*  
 883 *Geophys. Res. Atmos.*, 108(D3), n/a-n/a, doi:10.1029/2001JD001417, 2003.
- 884
- 885 An, J., Wang, H., Shen, L., Zhu, B., Zou, J., Gao, J. and Kang, H.: Characteristics of new particle  
 886 formation events in Nanjing, China: Effect of water-soluble ions, *Atmos. Environ.*, 108, 32–40,  
 887 doi:10.1016/j.atmosenv.2015.01.038, 2015.
- 888
- 889 Bae, M.-S., Schwab, J. J., Hogrefe, O., Frank, B. P., Lala, G. G. and Demerjian, K. L.:  
 890 Characteristics of size distributions at urban and rural locations in New York, *Atmos. Chem. Phys.*  
 891 *Discuss.*, 10(1), 69–108, doi:10.5194/acpd-10-69-2010, 2010.
- 892
- 893 Beddows, D. C. S., Harrison, R. M., Green, D. C. and Fuller, G. W.: Receptor modelling of both  
 894 particle composition and size distribution from a background site in London, UK, *Atmos. Chem.*  
 895 *Phys.*, 15(17), 10107–10125, doi:10.5194/acp-15-10107-2015, 2015.
- 896
- 897 Berland, K., Rose, C., Pey, J., Culot, A., Freney, E., Kalivitis, N., Kouvarakis, G., Cerro, J. C., Mallet,  
 898 M., Sartelet, K., Beckmann, M., Bourriane, T., Roberts, G., Marchand, N., Mihalopoulos, N. and  
 899 Sellegri, K.: Spatial extent of new particle formation events over the Mediterranean Basin from  
 900 multiple ground-based and airborne measurements, *Atmos. Chem. Phys.*, 17(15), 9567–9583,  
 901 doi:10.5194/acp-17-9567-2017, 2017.
- 902
- 903 Berndt, T., Böge, O. and Stratmann, F.: Formation of atmospheric H<sub>2</sub>SO<sub>4</sub>H<sub>2</sub>O particles in the  
 904 absence of organics: A laboratory study, *Geophys. Res. Lett.*, 33(15), 2–6,  
 905 doi:10.1029/2006GL026660, 2006.
- 906
- 907 Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crounse, J. D.,  
 908 Wennberg, P. O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R.,  
 909 Thornton, J. A., Donahue, N., Kjaergaard, H. G. and Ehn, M.: Highly oxygenated organic molecules  
 910 (HOM) from gas-phase autoxidation involving peroxy radicals: A key contributor to atmospheric  
 911 aerosol, *Chem. Rev.*, 119, 3472–3509, doi:10.1021/acs.chemrev.8b00395, 2019.
- 912
- 913 Bigi, A. and Harrison, R. M.: Analysis of the air pollution climate at a central urban background site,  
 914 *Atmos. Environ.*, 44(16), 2004–2012, doi:10.1016/j.atmosenv.2010.02.028, 2010.

915 Birmili, W., Weinhold, K., Rasch, F., Sonntag, A., Sun, J., Merkel, M., Wiedensohler, A., Bastian,  
 916 S., Schladitz, A., Löschau, G., Cyrys, J., Pitz, M., Gu, J., Kusch, T., Flentje, H., Quass, U., Kaminski,  
 917 H., Kuhlbusch, T. A. J., Meinhardt, F., Schwerin, A., Bath, O., Ries, L., Wirtz, K. and Fiebig, M.:  
 918 Long-term observations of tropospheric particle number size distributions and equivalent black  
 919 carbon mass concentrations in the German Ultrafine Aerosol Network (GUAN), *Earth Syst. Sci. Data*,  
 920 8(2), 355–382, doi:10.5194/essd-8-355-2016, 2016.  
 921  
 922 Bousiotis, D., Pope, F. D., Beddows, D. C., Dall’Osto, M., Massling, A., Nøjgaard, J. K.,  
 923 Nørdrstrom, C., Niemi, J. V., Portin, H., Petäjä, T., Perez, N., Alastuey, A., Querol, X., Kouvarakis,  
 924 G., Vratolis, S., Eleftheriadis, K., Wiedensohler, A., Weinhold, K., Merkel, M., Tuch, T., and  
 925 Harrison, R. M.: An Analysis of New Particle Formation (NPF) at Thirteen European Sites, *Atmos.*  
 926 *Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-414>, in review, 2020.  
 927  
 928 Bousiotis, D., Osto, M., Beddows, D. C. S., Pope, F. D. and Harrison, R. M.: Analysis of new  
 929 particle formation (NPF) events at nearby rural, urban background and urban roadside sites, *Atmos.*  
 930 *Chem. Phys.*, 19, 5679–5694, 2019.  
 931  
 932 Brines, M., Dall’Osto, M., Beddows, D. C. S., Harrison, R. M., Gómez-Moreno, F., Núñez, L.,  
 933 Artíñano, B., Costabile, F., Gobbi, G. P., Salimi, F., Morawska, L., Sioutas, C. and Querol, X.: Traffic  
 934 and nucleation events as main sources of ultrafine particles in high-insolation developed world cities,  
 935 *Atmos. Chem. Phys.*, 15(10), 5929–5945, doi:10.5194/acp-15-5929-2015, 2015.  
 936  
 937 Carnerero, C., Pérez, N., Petäjä, T., Laurila, T. M., Ahonen, L. R., Kontkanen, J., Ahn, K. H.,  
 938 Alastuey, A. and Querol, X.: Relating high ozone, ultrafine particles, and new particle formation  
 939 episodes using cluster analysis, *Atmos. Environ.* X, 4(October), doi:10.1016/j.aeaoa.2019.100051,  
 940 2019.  
 941  
 942 Charron, A. and Harrison, R. M.: Primary particle formation from vehicle emissions during exhaust  
 943 dilution in the roadside atmosphere, *Atmos. Environ.*, 37(29), 4109–4119, doi:10.1016/S1352-  
 944 2310(03)00510-7, 2003.  
 945  
 946 Charron, A., Birmili, W. and Harrison, R. M.: Fingerprinting particle origins according to their size  
 947 distribution at a UK rural site, *J. Geophys. Res. Atmos.*, 113(7), 1–15, doi:10.1029/2007JD008562,  
 948 2008.  
 949  
 950 Charron, A., Degrendele, C., Laongsri, B. and Harrison, R. M.: Receptor modelling of secondary and  
 951 carbonaceous particulate matter at a southern UK site, *Atmos. Chem. Phys.*, 13(4), 1879–1894,  
 952 doi:10.5194/acp-13-1879-2013, 2013.  
 953  
 954

955 Cheung, H. C., Chou, C. C.-K., Huang, W.-R. and Tsai, C.-Y.: Characterization of ultrafine particle  
 956 number concentration and new particle formation in an urban environment of Taipei, Taiwan, *Atmos.*  
 957 *Chem. Phys.*, 13(17), 8935–8946, doi:10.5194/acp-13-8935-2013, 2013.  
 958  
 959 Chu, B., Kerminen, V., Bianchi, F., Yan, C., Petäjä, T. and Kulmala, M.: Atmospheric new particle  
 960 formation in China, *Atmos. Chem. Phys.*, 19, 115–138, doi:10.5194/acp-2018-612, 2019  
 961  
 962 Dada, L., Paasonen, P., Nieminen, T., Buenrostro Mazon, S., Kontkanen, J., Peräkylä, O.,  
 963 Lehtipalo, K., Hussein, T., Petäjä, T., Kerminen, V. M., Bäck, J. and Kulmala, M.: Long-term  
 964 analysis of clear-sky new particle formation events and nonevents in Hyytiälä, *Atmos. Chem. Phys.*,  
 965 17(10), 6227–6241, doi:10.5194/acp-17-6227-2017, 2017.  
 966  
 967 Dai, L., Wang, H., Zhou, L., An, J., Tang, L., Lu, C., Yan, W., Liu, R., Kong, S., Chen, M., Lee, S.  
 968 and Yu, H.: Regional and local new particle formation events observed in the Yangtze River Delta  
 969 region, China, *J. Geophys. Res.*, 122(4), 2389–2402, doi:10.1002/2016JD026030, 2017.  
 970  
 971 Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P. and Lehtinen, K. E.  
 972 J.: Formation and growth of fresh atmospheric aerosols: Eight years of aerosol size distribution data  
 973 from SMEAR II, Hyytiälä, Finland, *Boreal Environ. Res.*, 10(5), 323–336,  
 974 doi:10.1016/j.ijpharm.2012.03.044, 2005.  
 975  
 976 Dall’Osto, M., Beddows, D. C. S., Asmi, A., Poulain, L., Hao, L., Freney, E., Allan, J. D.,  
 977 Canagaratna, M., Crippa, M., Bianchi, F., De Leeuw, G., Eriksson, A., Swietlicki, E., Hansson, H.  
 978 C., Henzing, J. S., Granier, C., Zemankova, K., Laj, P., Onasch, T., Prevot, A., Putaud, J. P., Sellegri,  
 979 K., Vidal, M., Virtanen, A., Simo, R., Worsnop, D., O’Dowd, C., Kulmala, M. and Harrison, R. M.:  
 980 Novel insights on new particle formation derived from a pan-european observing system, *Sci. Rep.*,  
 981 8(1), 1–11, doi:10.1038/s41598-017-17343-9, 2018.  
 982  
 983 Dall’Osto, M., Querol, X., Alastuey, A., O’Dowd, C., Harrison, R. M., Wenger, J. and Gómez-  
 984 Moreno, F. J.: On the spatial distribution and evolution of ultrafine particles in Barcelona, *Atmos.*  
 985 *Chem. Phys.*, 13(2), 741–759, doi:10.5194/acp-13-741-2013, 2013.  
 986  
 987 Dall’Osto, M., Beddows, D. C. S., Pey, J., Rodriguez, S., Alastuey, A., M. Harrison, R. and Querol,  
 988 X.: Urban aerosol size distributions over the Mediterranean city of Barcelona, NE Spain, *Atmos.*  
 989 *Chem. Phys.*, 12(22), 10693–10707, doi:10.5194/acp-12-10693-2012, 2012.  
 990  
 991 Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F.,  
 992 Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I. H., Rissanen, M., Jokinen, T.,  
 993 Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B.,  
 994 Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A.,

995 Kerminen, V. M., Kulmala, M., Worsnop, D. R., Wildt, J. and Mentel, T. F.: A large source of low-  
 996 volatility secondary organic aerosol, *Nature*, 506(7489), 476–479, doi:10.1038/nature13032, 2014.  
 997  
 998 Fenske, J. D., Hasson, A.S., Paulson, S. E., Kuwata, K. T., Ho, A., Houk, K. N.: The Pressure  
 999 Dependence of the OH Radical Yield from Ozone Alkene Reactions *J Phys Chem A*, 104 7821, 2000  
 1000  
 1001 Fuchs, N. A. and Sutugin, A. G.: Highly dispersed aerosols, *Top. Curr. Aerosol Res.*, 1,  
 1002 doi:https://doi.org/10.1016/B978-0-08-016674-2.50006-6, 1971.  
 1003  
 1004 Glasoe, W. a, Volz, K., Panta, B., Freshour, N., Bachman, R., Hanson, D. R., Mcmurry, P. H. and  
 1005 Jen, C.: Sulfuric acid nucleation: An experimental study of the effect of seven bases, , 1933–1950,  
 1006 doi:10.1002/2014JD022730, 2015.  
 1007  
 1008 Größ, J., Hamed, A., Sonntag, A., Spindler, G., Manninen, H. E., Nieminen, T., Kulmala, M.,  
 1009 Hörrak, U., Plass-Dülmer, C., Wiedensohler, A., and Birmili, W.: Atmospheric new particle  
 1010 formation at the research station Melpitz, Germany: connection with gaseous precursors and  
 1011 meteorological parameters, *Atmos. Chem. Phys.*, 18, 1835–1861, https://doi.org/10.5194/acp-18-  
 1012 1835-2018, 2018.  
 1013  
 1014 Guo, S., Hu, M., Peng, J., Wu, Z., Zamora, M. L., Shang, D., Du, Z., Zheng, J., Fang, X., Tang, R.,  
 1015 Wu, Y., Zeng, L., Shuai, S., Zhang, W., Wang, Y., Ji, Y., Li, Y., Zhang, A. L., Wang, W., Zhang, F.,  
 1016 Zhao, J., Gong, X., Wang, C., Molina, M. J. and Zhang, R.: Remarkable nucleation and growth of  
 1017 ultrafine particles from vehicular exhaust, *Proc. Nat. Acad. Sci. U. S. A.*, 117(7), 3427–3432,  
 1018 doi:10.1073/pnas.1916366117, 2020.  
 1019  
 1020 Hallar, A. G., Petersen, R., McCubbin, I. B., Lowenthal, D., Lee, S., Andrews, E. and Yu, F.:  
 1021 Climatology of new particle formation and corresponding precursors at storm peak laboratory,  
 1022 *Aerosol Air Qual. Res.*, 16(3), 816–826, doi:10.4209/aaqr.2015.05.0341, 2016.  
 1023  
 1024 Hamed, A., Korhonen, H., Sihto, S. L., Joutsensaari, J., Jrvinen, H., Petäjä, T., Arnold, F.,  
 1025 Nieminen, T., Kulmala, M., Smith, J. N., Lehtinen, K. E. J. and Laaksonen, A.: The role of relative  
 1026 humidity in continental new particle formation, *J. Geophys. Res. Atmos.*, 116(3), 1–12,  
 1027 doi:10.1029/2010JD014186, 2011.  
 1028  
 1029 Harrison, R. M.: Urban atmospheric chemistry: A very special case for study, *npj Clim. Atmos. Sci.*,  
 1030 1(1), 5, doi:10.1038/s41612-017-0010-8, 2017.  
 1031  
 1032 Henschel, H., Kurtén, T., Vehkamäki, H.: Computational study on the effect of hydration on new  
 1033 particle formation in the sulfuric acid/ammonia and sulfuric acid/dimethylamine systems, *J. Phys.*  
 1034 *Chem. A* 2016, 120, 11, 1886–1896, 2016.



1035 Hidy, G. M.: Atmospheric sulfur and nitrogen oxides, Academic Press, ISBN: 9781483288666,  
 1036 1994  
 1037  
 1038 Hietikko, R., Kuuluvainen, H., Harrison, R. M., Portin, H., Timonen, H., Niemi, J. V and Rönkkö,  
 1039 T.: Diurnal variation of nanocluster aerosol concentrations and emission factors in a street canyon,  
 1040 Atmos. Environ., 189, 98–106, doi:10.1016/j.atmosenv.2018.06.031, 2018.  
 1041  
 1042 Iida, K., Stolzenburg, M. R., McMurry, P. H. and Smith, J. N.: Estimating nanoparticle growth rates  
 1043 from size-dependent charged fractions: Analysis of new particle formation events in Mexico City, J.  
 1044 Geophys. Res. Atmos., 113(5), 1–15, doi:10.1029/2007JD009260, 2008.  
 1045  
 1046 Järvi, L., Hannuniemi, H., Hussein, T., Junninen, H., Aalto, P., Hillamo, R., Mäkelä, T., Keronen, P.  
 1047 and Siivola, E.: The urban measurement station SMEAR III : Continuous monitoring of air pollution  
 1048 and surface – atmosphere interactions in Helsinki , Finland, 14(April), 86–109, 2009.  
 1049  
 1050 Jayaratne, R., Pushpawela, B., He, C., Li, H., Gao, J., Chai, F. and Morawska, L.: Observations of  
 1051 particles at their formation sizes in Beijing, China, Atmos. Chem. Phys., 17(14), 8825–8835,  
 1052 doi:10.5194/acp-17-8825-2017, 2017.  
 1053  
 1054 Jeong, C.-H. H., Evans, G. J., McGuire, M. L., Y.-W. Chang, R., Abbatt, J. P. D. D., Zeromskiene,  
 1055 K., Mozurkewich, M., Li, S.-M. M., Leaitch, W. R., Chang, R. Y.-W., Abbatt, J. P. D. D.,  
 1056 Zeromskiene, K., Mozurkewich, M., Li, S.-M. M. and Leaitch, W. R.: Particle formation and growth  
 1057 at five rural and urban sites, Atmos. Chem. Phys., 10(16), 7979–7995, doi:10.5194/acp-10-7979-  
 1058 2010, 2010.  
 1059  
 1060 Kalkavouras, P., Bossioli, E., Bezantakos, S., Bougiatioti, A., Kalivitis, N., Stavroulas, I.,  
 1061 Kouvarakis, G., Protonotariou, A. P., Dandou, A., Biskos, G., Mihalopoulos, N., Nenes, A. and  
 1062 Tombrou, M.: New particle formation in the southern Aegean Sea during the Etesians: Importance  
 1063 for CCN production and cloud droplet number, Atmos. Chem. Phys., 17(1), 175–192,  
 1064 doi:10.5194/acp-17-175-2017, 2017.  
 1065  
 1066 Kerminen, V., Lehtinen, K. E. J., Anttila, T., Kulmala, M., Lehtinen, K. E. J., Anttila, T. and Kulmala,  
 1067 M.: Dynamics of atmospheric nucleation mode particles : a timescale analysis, Tellus, 56B, 135–146,  
 1068 doi:10.3402/tellusb.v56i2.16411, 2004.  
 1069  
 1070 Kerminen, V. M., Pirjola, L. and Kulmala, M.: How significantly does coagulation scavenging limit  
 1071 atmospheric particle production?, J. Geophys. Res. Atmos., 106(D20), 24119–24125,  
 1072 doi:10.1029/2001JD000322, 2001.  
 1073  
 1074

1075 Kerminen, V. M., Kulmala, M., Worsnop, D. R., Wildt, J. and Mentel, T. F.: A large source of low-  
 1076 volatility secondary organic aerosol, *Nature*, 506(7489), 476–479, doi:10.1038/nature13032, 2014.  
 1077  
 1078 Ketzel, M., Wählin, P., Kristensson, A., Swietlicki, E., Berkowicz, R., Nielsen, O. J. and Palmgren,  
 1079 F.: Particle size distribution and particle mass measurements at urban, near-city and rural level in the  
 1080 Copenhagen area and Southern Sweden, *Atmos. Chem. Phys. Discuss.*, 3(6), 5513–5546,  
 1081 doi:10.5194/acpd-3-5513-2003, 2004.  
 1082  
 1083 Kiendler-Scharr, A., Wildt, J., Dal Maso, M., Hohaus, T., Kleist, E., Mentel, T. F., Tillmann, R.,  
 1084 Uerlings, R., Schurr, U. and Wahner, A.: New particle formation in forests inhibited by isoprene  
 1085 emissions, 461, 381–384, 2009.  
 1086  
 1087 Kim, K. H., Kabir, E. and Kabir, S.: A review on the human health impact of airborne particulate  
 1088 matter, *Environ. Int.*, 74, 136–143, doi:10.1016/j.envint.2014.10.005, 2015.  
 1089  
 1090 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes,  
 1091 L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas,  
 1092 G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A.,  
 1093 Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin,  
 1094 A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J.,  
 1095 Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld,  
 1096 J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A.,  
 1097 Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D.  
 1098 R., Baltensperger, U. and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in  
 1099 atmospheric aerosol nucleation, *Nature*, 476(7361), 429–435, doi:10.1038/nature10343, 2011.  
 1100  
 1101 Korhonen, P., Kulmala, M., Laaksonen, A., Viisanen, Y., McGraw, R. and Seinfeld, J. H.: Ternary  
 1102 nucleation of H<sub>2</sub>SO<sub>4</sub>, NH<sub>3</sub> and H<sub>2</sub>O in the atmosphere, *J. Geophys. Res.*, 104(D21), 26349–26353,  
 1103 1999.  
 1104  
 1105 Kulmala, M., Petäjä, T., Mönkkönen, P., Koponen, I. K., Dal Maso, M., Aalto, P. P., Lehtinen, K.  
 1106 E. J. and Kerminen, V.-M.: On the growth of nucleation mode particles: source rates of condensable  
 1107 vapor in polluted and clean environments, *Atmos. Chem. Phys. Discuss.*, 4(5), 6943–6966,  
 1108 doi:10.5194/acpd-4-6943-2004, 2005.  
 1109  
 1110 Kulmala, M. and Kerminen, V. M.: On the formation and growth of atmospheric nanoparticles,  
 1111 *Atmos. Res.*, 90(2–4), 132–150, doi:10.1016/j.atmosres.2008.01.005, 2008.  
 1112  
 1113 Kulmala, M., Kerminen, V.-M. M., Petäjä, T., Ding, A. J. and Wang, L.: Atmospheric gas-to-particle  
 1114 conversion: Why NPF events are observed in megacities?, *Faraday Discuss.*, 200, 271–288,

doi:10.1039/c6fd00257a, 2017.

Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A. and Kerminen, V. M.: Measurement of the nucleation of atmospheric aerosol particles, *Nat. Protoc.*, 7(9), 1651–1667, doi:10.1038/nprot.2012.091, 2012.

Kulmala, M., Petäjä, T., Mönkkönen, P., Koponen, I. K., Dal Maso, M., Aalto, P. P., Lehtinen, K. E. J. and Kerminen, V.-M.: On the growth of nucleation mode particles: source rates of condensable vapor in polluted and clean environments, *Atmos. Chem. Phys. Discuss.*, 4(5), 6943–6966, doi:10.5194/acpd-4-6943-2004, 2005.

Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen, P., Hämeri, K. and O’Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus, Ser. B Chem. Phys. Meteorol.*, 53(4), 479–490, doi:10.3402/tellusb.v53i4.16622, 2001.

Kürten, A., Li, C., Bianchi, F., Curtius, J., Dias, A., Donahue, N. M., Duplissy, J., Flagan, R. C., Hakala, J., Jokinen, T., Kirkby, J., Kulmala, M., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Onnela, A., Rissanen, M. P., Simon, M., Sipilä, M., Stozhkov, Y., Tröstl, J., Ye, P., and McMurry, P. H.: New particle formation in the sulfuric acid–dimethylamine–water system: reevaluation of CLOUD chamber measurements and comparison to an aerosol nucleation and growth model, *Atmos. Chem. Phys.*, 18, 845–863, <https://doi.org/10.5194/acp-18-845-2018>, 2018.

Kürten, A., Bergen, A., Heinritzi, M., Leiminger, M., Lorenz, V., Piel, F., Simon, M., Sitals, R., Wagner, A. C. and Curtius, J.: Observation of new particle formation and measurement of sulfuric acid, ammonia, amines and highly oxidized organic molecules at a rural site in central Germany, *Atmos. Chem. Phys.*, 16(19), 12793–12813, doi:10.5194/acp-16-12793-2016, 2016.

Lee, S.-H. H., Uin, J., Guenther, A. B., de Gouw, J. A., Yu, F., Nadykto, A. B., Herb, J., Ng, N. L., Koss, A., Brune, W. H., Baumann, K., Kanawade, V. P., Keutsch, F. N., Nenes, A., Olsen, K., Goldstein, A. and Ouyang, Q.: Isoprene suppression of new particle formation: Potential mechanisms and implications, *J. Geophys. Res. Atmos.*, 121(24), 14,621–14,635, doi:10.1002/2016JD024844, 2016.

Lehtinen, K. E. J., Korhonen, H., Dal Maso, M. and Kulmala, M.: On the concept of condensation sink diameter, *Boreal Environ. Res.*, 8(4), 405–411, 2003.

Lehtipalo, K., Yan, C., Dada, L., Bianchi, F., Xiao, M., Wagner, R., Stolzenburg, D., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bernhammer, A.,

1155 Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S. B., Chen, D., Chen, X., Dias, A., Dommen,  
 1156 J., Draper, D. C., Duplissy, J., Ehn, M., Finkenzeller, H., Fischer, L., Frege, C., Fuchs, C., Garmash,  
 1157 O., Gordon, H., Hakala, J., He, X., Heikkinen, L., Heinritzi, M., Helm, J. C., Hofbauer, V., Hoyle, C.  
 1158 R., Jokinen, T., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Piel, F., Sarnela, N., Schallhart,  
 1159 S., Schuchmann, S., Sengupta, K. and Simon, M.: Multicomponent new particle formation from  
 1160 sulfuric acid, ammonia, and biogenic vapors, (3), 1–10, 2018.  
 1161  
 1162 Li, X., Chee, S., Hao, J., Abbatt, J. P. D., Jiang, J. and Smith, J. N.: Relative humidity effect on the  
 1163 formation of highly oxidized molecules and new particles during monoterpene oxidation, *Atmos.*  
 1164 *Chem. Phys.*, 19(3), 1555–1570, doi:10.5194/acp-19-1555-2019, 2019.  
 1165  
 1166 Makkonen, R., Asmi, A., Kerminen, V. M., Boy, M., Arneth, A., Hari, P. and Kulmala, M.: Air  
 1167 pollution control and decreasing new particle formation lead to strong climate warming, *Atmos.*  
 1168 *Chem. Phys.*, 12(3), 1515–1524, doi:10.5194/acp-12-1515-2012, 2012.  
 1169  
 1170 McFiggans, G., Mentel, T. F., Wildt, J., Pullinen, I., Kang, S., Kleist, E., Schmitt, S., Springer, M.,  
 1171 Tillmann, R., Wu, C., Zhao, D., Hallquist, M., Faxon, C., Le Breton, M., Hallquist, Å. M., Simpson,  
 1172 D., Bergström, R., Jenkin, M. E., Ehn, M., Thornton, J. A., Alfarra, M. R., Bannan, T. J., Percival, C.  
 1173 J., Priestley, M., Topping, D. and Kiendler-Scharr, A.: Secondary organic aerosol reduced by mixture  
 1174 of atmospheric vapours, *Nature*, 565(7741), 587–593, doi:10.1038/s41586-018-0871-y, 2019.  
 1175 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J. and Carslaw, K. S.: Impact of  
 1176 nucleation on global CCN, *Atmos. Chem. Phys.*, 9(21), 8601–8616, doi:10.5194/acp-9-8601-2009,  
 1177 2009.  
 1178  
 1179 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J. and Carslaw, K. S.: Impact of  
 1180 nucleation on global CCN, *Atmos. Chem. Phys.*, 9(21), 8601–8616, doi:10.5194/acp-9-8601-2009,  
 1181 2009.  
 1182  
 1183 Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S. H., Weingartner, E., Riipinen,  
 1184 I., Kulmala, M., Spracklen, D. V., Carslaw, K. S. and Baltensperger, U.: Evidence for the role of  
 1185 organics in aerosol particle formation under atmospheric conditions, *Proc. Nat. Acad. Sci.*, 107(15),  
 1186 6646–6651, doi:10.1073/pnas.0911330107, 2010.  
 1187  
 1188 Minguillón, M. C., Brines, M., Pérez, N., Reche, C., Pandolfi, M., Fonseca, A. S., Amato, F.,  
 1189 Alastuey, A., Lyasota, A., Codina, B., Lee, H. K., Eun, H. R., Ahn, K. H. and Querol, X.: New particle  
 1190 formation at ground level and in the vertical column over the Barcelona area, *Atmos. Res.*, 164–165,  
 1191 118–130, doi:10.1016/j.atmosres.2015.05.003, 2015.  
 1192  
 1193 Mirabel, P. and Katz, J. L.: Binary homogeneous nucleation as a mechanism for the formation of  
 1194 aerosols, *J. Chem. Phys.*, 60(3), 1138–1144, doi:10.1063/1.1681124, 1974.

1195 Mølgaard, B., Birmili, W., Clifford, S., Massling, A., Eleftheriadis, K., Norman, M., Vratolis, S.,  
 1196 Wehner, B., Corander, J., Hämeri, K. and Hussein, T.: Evaluation of a statistical forecast model for  
 1197 size-fractionated urban particle number concentrations using data from five European cities, J.  
 1198 Aerosol Sci., 66, 96–110, doi:10.1016/j.jaerosci.2013.08.012, 2013.  
 1199  
 1200 Molteni, U., Bianchi, F., Klein, F., El Haddad, I., Frege, C., Rossi, M. J., Dommen, J. and  
 1201 Baltensperger, U.: Formation of highly oxygenated organic molecules from aromatic compounds,  
 1202 Atmos. Chem. Phys., 18(3), 1909–1921, doi:10.5194/acp-18-1909-2018, 2018.  
 1203  
 1204 Napari, I., Noppel, M., Vehkamäki, H. and Kulmala, M.: An improved model for ternary nucleation  
 1205 of sulfuric acid-ammonia-water, J. Chem. Phys., 116(10), 4221–4227, doi:10.1063/1.1450557, 2002.  
 1206  
 1207 Nieminen, T., Kerminen, V.-M., Petäjä, T., Aalto, P. P., Arshinov, M., Asmi, E., Baltensperger, U.,  
 1208 Beddows, D. C. S., Beukes, J. P., Collins, D., Ding, A., Harrison, R. M., Henzing, B., Hooda, R., Hu,  
 1209 M., Hörrak, U., Kivekäs, N., Komsaare, K., Krejci, R., Kristensson, A., Laakso, L., Laaksonen, A.,  
 1210 Leaitch, W. R., Lihavainen, H., Mihalopoulos, N., Németh, Z., Nie, W., O ’dowd, C., Salma, I.,  
 1211 Sellegri, K., Svenningsson, B., Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M.,  
 1212 Wiedensohler, A., Wu, Z., Virtanen, A., Kulmala, M., O ’dowd, C., Salma, I., Sellegri, K.,  
 1213 Svenningsson, B., Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler,  
 1214 A., Wu, Z., Virtanen, A., Kulmala, M., O ’dowd, C., Salma, I., Sellegri, K., Svenningsson, B.,  
 1215 Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler, A., Wu, Z., Virtanen,  
 1216 A. and Kulmala, M.: Global analysis of continental boundary layer new particle formation based on  
 1217 long-term measurements, Atmos. Chem. Phys. Discuss, 5194, 2018–304, doi:10.5194/acp-2018-304,  
 1218 2018.  
 1219  
 1220 Nieminen, T., Lehtinen, K. E. J. and Kulmala, M.: Sub-10 nm particle growth by vapor condensation-  
 1221 effects of vapor molecule size and particle thermal speed, Atmos. Chem. Phys., 10(20), 9773–9779,  
 1222 doi:10.5194/acp-10-9773-2010, 2010.  
 1223  
 1224 O’Dowd, C. D., Jimenez, J. L., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hämeri, K., Pirjola,  
 1225 L., Kulmala, M., Jennings, S. G. and Hoffmann, T.: Marine aerosol formation from biogenic iodine  
 1226 emissions, Lett. to Nat., 417(June), 1–5, doi:10.1038/nature00773.1.2.3.4.5.6.7.8.9.10., 2002.  
 1227  
 1228 Olenius, T., Halonen, R., Kurten, T., Henschel, H., Maatta, O. K., Ortega, I. K., Jen, C.,  
 1229 Vehkamäki, H. and Riipinen, I.: New particle formation from sulfuric acid amines: Comparison of  
 1230 monomethylamine, dimethylamine, and trimethylamine, J. Geophys. Res. Atmos., 7103–7118,  
 1231 doi:10.1002/2017JD026501, 2017.  
 1232  
 1233 Olin, M., Kuuluvainen, H., Aurela, M., Kalliokoski, J., Kuittinen, N., Isotalo, M., Timonen, H. J.,  
 1234 Niemi, J. V., Rönkkö, T., and Dal Maso, M.: Traffic-originated nanocluster emission exceeds

1235 H<sub>2</sub>SO<sub>4</sub>-driven photochemical new particle formation in an urban area, *Atmos. Chem. Phys.*, 20, 1–  
1236 13, <https://doi.org/10.5194/acp-20-1-2020>, 2020.

1237

1238 Paasonen, P., Asmi, A., Petäjä, T., Kajos, M. K., Äijälä, M., Junninen, H., Holst, T., Abbatt, J. P.  
1239 D., Arneth, A., Birmili, W., Van Der Gon, H. D., Hamed, A., Hoffer, A., Laakso, L., Laaksonen,  
1240 A., Richard Leaitch, W., Plass-Dülmer, C., Pryor, S. C., Räisänen, P., Swietlicki, E., Wiedensohler,  
1241 A., Worsnop, D. R., Kerminen, V. M. and Kulmala, M.: Warming-induced increase in aerosol  
1242 number concentration likely to moderate climate change, *Nat. Geosci.*, 6(6), 438–442,  
1243 doi:10.1038/ngeo1800, 2013.

1244

1245 Park, M., Yum, S. S. and Kim, J. H.: Characteristics of submicron aerosol number size distribution  
1246 and new particle formation events measured in Seoul, Korea, during 2004–2012, *Asia-Pacific J.*  
1247 *Atmos. Sci.*, 51(1), 1–10, doi:10.1007/s13143-014-0055-0, 2015.

1248

1249 Petäjä, T., Mauldin, R. L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M.,  
1250 Adamov, A., Kotiaho, T. and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest  
1251 site, *Atmos. Chem. Phys.*, 9(19), 7435–7448, doi:10.5194/acp-9-7435-2009, 2009.

1252

1253 Pikridas, M., Sciare, J., Freutel, F., Crumeyrolle, S., Von Der Weiden-Reinmüller, S. L., Borbon, A.,  
1254 Schwarzenboeck, A., Merkel, M., Crippa, M., Kostenidou, E., Psichoudaki, M., Hildebrandt, L.,  
1255 Engelhart, G. J., Petäjä, T., Prévôt, A. S. H., Drewnick, F., Baltensperger, U., Wiedensohler, A.,  
1256 Kulmala, M., Beekmann, M. and Pandis, S. N.: In situ formation and spatial variability of particle  
1257 number concentration in a European megacity, *Atmos. Chem. Phys.*, 15(17), 10219–10237,  
1258 doi:10.5194/acp-15-10219-2015, 2015.

1259

1260 Pillai, P., Khlystov, A., Walker, J. and Aneja, V.: Observation and analysis of particle nucleation at  
1261 a forest site in southeastern US, *Atmosphere (Basel)*, 4(2), 72–93, doi:10.3390/atmos4020072, 2013.

1262

1263 Poling, B. E., Prausnitz, J. M. and O’Connell, J. P.: The properties of gases and liquids, 5th ed.,  
1264 McGraw-Hill Education., 2001.

1265

1266 Politis, M., Pilinis, C. and Lekkas, T. D.: Ultrafine particles (UFP) and health effects. Dangerous.  
1267 Like no other PM? Review and analysis, *Glob. Nest J.*, 10(3), 439–452, 2008.

1268

1269 Quéléver, L. L. J., Kristensen, K., Normann Jensen, L., Rosati, B., Teiwes, R., Daellenbach, K. R.,  
1270 Peräkylä, O., Roldin, P., Bossi, R., Pedersen, H. B., Glasius, M., Bilde, M. and Ehn, M.: Effect of  
1271 temperature on the formation of highly oxygenated organic molecules (HOMs) from alpha-pinene  
1272 ozonolysis, *Atmos. Chem. Phys.*, 19(11), 7609–7625, doi:10.5194/acp-19-7609-2019, 2019.

1273

1274

1275 Querol, X., Gangoiti, G., Mantilla, E., Alastuey, A., Minguillón, M. C., Amato, F., Reche, C., Viana,  
 1276 M., Moreno, T., Karanasiou, A., Rivas, I., Pérez, N., Ripoll, A., Brines, M., Ealo, M., Pandolfi, M.,  
 1277 Lee, H. K., Eun, H. R., Park, Y. H., Escudero, M., Beddows, D., Harrison, R. M., Bertrand, A.,  
 1278 Marchand, N., Liasota, A., Codina, B., Olid, M., Udina, M., Jiménez-Esteve, B. B., Jiménez-Esteve,  
 1279 B. B., Alonso, L., Millán, M. and Ahn, K. H.: Phenomenology of high-ozone episodes in NE Spain,  
 1280 *Atmos. Chem. Phys.*, 17(4), 2817–2838, doi:10.5194/acp-17-2817-2017, 2017.  
 1281  
 1282 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J.,  
 1283 Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J.,  
 1284 Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A.,  
 1285 Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petäjä, T., Tsagkogeorgas, G., Vaattovaara,  
 1286 P., Viisanen, Y., Vrtala, A. and Wagner, P. E.: Oxidation Products of biogenic atmospheric particles,  
 1287 *Science*, 717, 717–722, doi:10.1126/science.1243527, 2014.  
 1288  
 1289 Rimnácová, D., Ždímal, V., Schwarz, J., Smolík, J. and Rimnác, M.: Atmospheric aerosols in suburb  
 1290 of Prague: The dynamics of particle size distributions, *Atmos. Res.*, 101(3), 539–552,  
 1291 doi:10.1016/j.atmosres.2010.10.024, 2011.  
 1292 Rose, C., Zha, Q., Dada, L., Yan, C., Lehtipalo, K., Junninen, H., Mazon, S. B., Jokinen, T., Sarnela,  
 1293 N., Sipilä, M., Petäjä, T., Kerminen, V. M., Bianchi, F. and Kulmala, M.: Observations of biogenic  
 1294 ion-induced cluster formation in the atmosphere, *Sci. Adv.*, 4(4), 1–10, doi:10.1126/sciadv.aar5218,  
 1295 2018.  
 1296  
 1297 Rivas, I., Beddows, D. C. S., Amato, F., Green, D. C., Järvi, L., Hueglin, C., Reche, C., Timonen, H.,  
 1298 Fuller, G. W., Niemi, J. V., Pérez, N., Aurela, M., Hopke, P. K., Alastuey, A., Kulmala, M., Harrison,  
 1299 R. M., Querol, X. and Kelly, F. J.: Source apportionment of particle number size distribution in urban  
 1300 background and traffic stations in four European cities, *Environ. Int.*, 135, 105345,  
 1301 doi:10.1016/j.envint.2019.105345, 2020.  
 1302  
 1303 Rizzo, L. V., Artaxo, P., Karl, T., Guenther, A. B. and Greenberg, J.: Aerosol properties, in-canopy  
 1304 gradients, turbulent fluxes and VOC concentrations at a pristine forest site in Amazonia, *Atmos.*  
 1305 *Environ.*, 44(4), 503–511, doi:10.1016/j.atmosenv.2009.11.002, 2010.  
 1306  
 1307 Salma, I., Borsòs, T., Weidinger, T., Aalto, P., Hussein, T., Dal Maso, M. and Kulmala, M.:  
 1308 Production, growth and properties of ultrafine atmospheric aerosol particles in an urban environment,  
 1309 *Atmos. Chem. Phys.*, 11(3), 1339–1353, doi:10.5194/acp-11-1339-2011, 2011.  
 1310  
 1311 Schwartz, J., Dockery, D. W. and Neas, L. M.: Is Daily Mortality Associated Specifically with Fine  
 1312 Particles?, *J. Air Waste Manag. Assoc.*, 46(10), 927–939, doi:10.1080/10473289.1996.10467528,  
 1313 1996.  
 1314

1315 Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics: From Air Pollution to Climate  
 1316 Change, 3rd Editio., John Wiley & Sons, Inc, New Jersey, Canada, 2012.  
 1317  
 1318 Shen, X., Sun, J., Kivekäs, N., Kristensson, A., Zhang, X., Zhang, Y., Zhang, L., Fan, R., Qi, X., Ma,  
 1319 Q. and Zhou, H.: Spatial distribution and occurrence probability of regional new particle formation  
 1320 events in eastern China, *Atmos. Chem. Phys.*, 18(2), 587–599, doi:10.5194/acp-18-587-2018, 2018.  
 1321  
 1322 Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C.,  
 1323 Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H.,  
 1324 Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A.,  
 1325 Zelenyuk, A. and Zhang, Q.: Recent advances in understanding secondary organic aerosol:  
 1326 Implications for global climate forcing, *Rev. Geophys.*, 55(2), 509–559,  
 1327 doi:10.1002/2016RG000540, 2017.  
 1328  
 1329 Siakavaras, D., Samara, C., Petrakakis, M. and Biskos, G.: Nucleation events at a coastal city during  
 1330 the warm period: Kerbside versus urban background measurements, *Atmos. Environ.*, 140, 60–68,  
 1331 doi:10.1016/j.atmosenv.2016.05.054, 2016.  
 1332  
 1333 Sipila, M., Berndt, T., Petaja, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin III,  
 1334 R. L., Hyvarinen, A. P., Lihavainen, H. and Kulmala, M.: The Role of Sulfuric Acid in  
 1335 Atmospheric Nucleation, *Science*, 327, 1243–1246, doi:10.1126/science.1180315, 2010.  
 1336  
 1337 Spracklen, D. V., Carslaw, K. S., Merikanto, J., Mann, G. W., Reddington, C. L., Pickering, S., Ogren,  
 1338 J. A., Andrews, E., Baltensperger, U., Weingartner, E., Boy, M., Kulmala, M., Laakso, L.,  
 1339 Lihavainen, H., Kivekäs, N., Komppula, M., Mihalopoulos, N., Kouvarakis, G., Jennings, S. G.,  
 1340 O'Dowd, C., Birmili, W., Wiedensohler, A., Weller, R., Gras, J., Laj, P., Sellegri, K., Bonn, B.,  
 1341 Krejci, R., Laaksonen, A., Hamed, A., Minikin, A., Harrison, R. M., Talbot, R. and Sun, J.: Explaining  
 1342 global surface aerosol number concentrations in terms of primary emissions and particle formation,  
 1343 *Atmos. Chem. Phys.*, 10(10), 4775–4793, doi:10.5194/acp-10-4775-2010, 2010.  
 1344  
 1345 Stolzenburg, D., Simon, M., Ranjithkumar, A., Kürten, A., Lehtipalo, K., Gordon, H., Ehrhart, S.,  
 1346 Finkenzeller, H., Pichelstorfer, L., Nieminen, T., He, X.-C., Brilke, S., Xiao, M., Amorim, A.,  
 1347 Baalbaki, R., Baccarini, A., Beck, L., Bräkling, S., Caudillo Murillo, L., Chen, D., Chu, B., Dada,  
 1348 L., Dias, A., Dommen, J., Duplissy, J., El Haddad, I., Fischer, L., Gonzalez Carracedo, L., Heinritzi,  
 1349 M., Kim, C., Koenig, T. K., Kong, W., Lamkaddam, H., Lee, C. P., Leiminger, M., Li, Z.,  
 1350 Makhmutov, V., Manninen, H. E., Marie, G., Marten, R., Müller, T., Nie, W., Partoll, E., Petäjä, T.,  
 1351 Pfeifer, J., Philippov, M., Rissanen, M. P., Rörup, B., Schobesberger, S., Schuchmann, S., Shen, J.,  
 1352 Sipilä, M., Steiner, G., Stozhkov, Y., Tauber, C., Tham, Y. J., Tomé, A., Vazquez-Pufleau, M.,  
 1353 Wagner, A. C., Wang, M., Wang, Y., Weber, S. K., Wimmer, D., Wlasits, P. J., Wu, Y., Ye, Q.,  
 1354 Zauner-Wieczorek, M., Baltensperger, U., Carslaw, K. S., Curtius, J., Donahue, N. M., Flagan, R.



1355 C., Hansel, A., Kulmala, M., Lelieveld, J., Volkamer, R., Kirkby, J., and Winkler, P. M.: Enhanced  
 1356 growth rate of atmospheric particles from sulfuric acid, *Atmos. Chem. Phys.*, 20, 7359–7372,  
 1357 <https://doi.org/10.5194/acp-20-7359-2020>, 2020.  
 1358

1359 Stolzenburg, D., Fischer, L., Vogel, A. L., Heinritzi, M., Schervish, M. and Simon, M., Wagner, A.  
 1360 C., Dada, L., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A.,  
 1361 Bianchi, F., Breitenlechner, M., Brilke, S., Buenorstro Mazon, S., Chen, D., Dias, A., Draper, D. C.,  
 1362 Duplissy, J., El Haddad, I., Finkenzeller, H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., He, X.,  
 1363 Helm., J., Hofbauer, V., Hoyle, C. R., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lampilahti, J.,  
 1364 Lawler, M., Lehtipalo, K., Leiminger, M., Mai, H., Mathot, S., Mentler, B., Molteni, U., Nie, W.,  
 1365 Nieminen, T., Nowak, J. B., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Quéléver, L. L. J.,  
 1366 Rissanen, M. P., Sarnela, N., Schallhart, S., Tauber, C., Tome, A., Wagner, R., Wang, M., Weitz,  
 1367 L., Wimmer, D., Xiao, M., Yan, C., Ye, P., Zha, Q., Baltensperger, U., Curtius, J., Dommen, J.,  
 1368 Flagan, R. C., Kulmala, M., Smith, J. N., Worsnop, D. R., Hansel, A., Donahue, N. M., Winkler, P.  
 1369 M.: Rapid growth of organic aerosol nanoparticles over a wide tropospheric temperature range,  
 1370 *PNAS* , 115(37), doi:10.1073/pnas.1807604115, 2018.  
 1371

1372 Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C.,  
 1373 Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J.,  
 1374 Adamov, A., Almeida, J., Bernhammer, A. K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S.,  
 1375 Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T.,  
 1376 Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lawler,  
 1377 M., Leiminger, M., Mathot, S., Möhler, O., Nieminen, T., Onnela, A., Petäjä, T., Piel, F. M.,  
 1378 Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipilä, M.,  
 1379 Smith, J. N., Steiner, G., Tomè, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D.,  
 1380 Winkler, P. M., Ye, P., Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I.,  
 1381 Worsnop, D. R., Donahue, N. M. and Baltensperger, U.: The role of low-volatility organic compounds  
 1382 in initial particle growth in the atmosphere, *Nature*, 533(7604), 527–531, doi:10.1038/nature18271,  
 1383 2016.  
 1384

1385 Vratolis, S., Gini, M. I., Bezantakos, S., Stavroulas, I., Kalivitis, N., Kostenidou, E., Louvaris, E.,  
 1386 Siakavaras, D., Biskos, G., Mihalopoulos, N., Pandis, S. N. N., Pilinis, C., Papayannis, A. and  
 1387 Eleftheriadis, K.: Particle number size distribution statistics at City-Centre Urban Background, urban  
 1388 background, and remote stations in Greece during summer, *Atmos. Environ.*, 213(May), 711–726,  
 1389 doi:10.1016/j.atmosenv.2019.05.064, 2019.  
 1390

1391 Wagner, P. and Kuttler, W.: Biogenic and anthropogenic isoprene in the near-surface urban  
 1392 atmosphere - A case study in Essen, Germany, *Sci. Total Environ.*, 475, 104–115,  
 1393 doi:10.1016/j.scitotenv.2013.12.026, 2014.  
 1394

1395 Wang, D., Fu, Q., Geng, F., Li, L., Wang, H., Qiao, L., Yang, X., Chen, J., Kerminen, V. M.,  
 1396 Petäjä, T., Worsnop, D. R., Kulmala, M. and Wang, L.: Atmospheric new particle formation from  
 1397 sulfuric acid and amines in a Chinese megacity, *Science*, 361(6399), 278–281,  
 1398 doi:10.1126/science.aao4839, 2018.  
 1399  
 1400 Wang, S., Wu, R., Berndt, T., Ehn, M. and Wang, L.: Formation of Highly Oxidized Radicals and  
 1401 Multifunctional Products from the Atmospheric Oxidation of Alkylbenzenes, ,  
 1402 doi:10.1021/acs.est.7b02374, 2017a.  
 1403  
 1404 Wang, Z., Wu, Z., Yue, D., Shang, D., Guo, S., Sun, J., Ding, A., Wang, L., Jiang, J., Guo, H., Gao,  
 1405 J., Cheung, H. C., Morawska, L., Keywood, M. and Hu, M.: New particle formation in China: Current  
 1406 knowledge and further directions, *Sci. Total Environ.*, 577, 258–266,  
 1407 doi:10.1016/j.scitotenv.2016.10.177, 2017b.  
 1408  
 1409 Wang, F., Ketzel, M., Ellermann, T., Wählin, P., Jensen, S. S., Fang, D. and Massling, A.: Particle  
 1410 number, particle mass and NO<sub>x</sub> emission factors at a highway and an urban street in Copenhagen,  
 1411 *Atmos. Chem. Phys.*, 10(6), 2745–2764, doi:10.5194/acp-10-2745-2010, 2010.  
 1412  
 1413 Wang, M., Kong, W., Marten, R., He, X. C., Chen, D., Pfeifer, J., Heitto, A., Kontkanen, J., Dada,  
 1414 L., Kürten, A., Yli-Juuti, T., Manninen, H. E., Amanatidis, S., Amorim, A., Baalbaki, R., Baccarini,  
 1415 A., Bell, D. M., Bertozzi, B., Bräkling, S., Brilke, S., Murillo, L. C., Chiu, R., Chu, B., De  
 1416 Menezes, L. P., Duplissy, J., Finkenzeller, H., Carracedo, L. G., Granzin, M., Guida, R., Hansel, A.,  
 1417 Hofbauer, V., Krechmer, J., Lehtipalo, K., Lamkaddam, H., Lampimäki, M., Lee, C. P.,  
 1418 Makhmutov, V., Marie, G., Mathot, S., Mauldin, R. L., Mentler, B., Müller, T., Onnela, A., Partoll,  
 1419 E., Petäjä, T., Philippov, M., Pospisilova, V., Ranjithkumar, A., Rissanen, M., Rörup, B., Scholz,  
 1420 W., Shen, J., Simon, M., Sipilä, M., Steiner, G., Stolzenburg, D., Tham, Y. J., Tomé, A., Wagner,  
 1421 A. C., Wang, D. S., Wang, Y., Weber, S. K., Winkler, P. M., Wlasits, P. J., Wu, Y., Xiao, M., Ye,  
 1422 Q., Zauner-Wieczorek, M., Zhou, X., Volkamer, R., Riipinen, I., Dommen, J., Curtius, J.,  
 1423 Baltensperger, U., Kulmala, M., Worsnop, D. R., Kirkby, J., Seinfeld, J. H., El-Haddad, I., Flagan,  
 1424 R. C. and Donahue, N. M.: Rapid growth of new atmospheric particles by nitric acid and ammonia  
 1425 condensation, *Nature*, 581(7807), 184–189, doi:10.1038/s41586-020-2270-4, 2020.  
 1426  
 1427 Weber, R. J., McMurry, P. H., Eisele, F. L. and Tanner, D. J.: Measurement of expected nucleation  
 1428 precursor species and 3-500-nm diameter particles at Mauna Loa Observatory, Hawaii, *J. Atmos.*  
 1429 *Sci.*, 52(12), 2242–2257, doi:10.1175/1520-0469(1995)052<2242:MOENPS>2.0.CO;2, 1995.  
 1430  
 1431 Wehner, B., Siebert, H., Stratmann, F., Tuch, T., Wiedensohler, A., Petäjä, T., Dal Maso, M. and  
 1432 Kulmala, M.: Horizontal homogeneity and vertical extent of new particle formation events, *Tellus*,  
 1433 *Ser. B Chem. Phys. Meteorol.*, 59(3), 362–371, doi:10.1111/j.1600-0889.2007.00260.x, 2007.  
 1434

1435 Wiedensohler, A., Ma, N., Birmili, W., Heintzenberg, J., Ditas, F., Andreae, M. O. and Panov, A.:  
 1436 Infrequent new particle formation over the remote boreal forest of Siberia, *Atmos. Environ.*, 200,  
 1437 167–169, doi:10.1016/j.atmosenv.2018.12.013, 2019.  
 1438  
 1439 Wonaschütz, A., Demattio, A., Wagner, R., Burkart, J., Zíková, N., Vodička, P., Ludwig, W., Steiner,  
 1440 G., Schwarz, J. and Hitzenberger, R.: Seasonality of new particle formation in Vienna, Austria -  
 1441 Influence of air mass origin and aerosol chemical composition, *Atmos. Environ.*, 118, 118–126,  
 1442 doi:10.1016/j.atmosenv.2015.07.035, 2015.  
 1443  
 1444 Woo, K. S., Chen, D. R., Pui, D. Y. H. H. and McMurry, P. H.: Measurement of Atlanta aerosol  
 1445 size distributions: Observations of lutrafine particle events, *Aerosol Sci. Technol.*, 34, 75–87,  
 1446 doi:10.1080/02786820120056, 2001.  
 1447  
 1448 Yamada, H.: Contribution of evaporative emissions from gasoline vehicles toward total VOC  
 1449 emissions in Japan, *Sci. Total Environ.*, 449, 143–149, doi:10.1016/j.scitotenv.2013.01.045, 2013.  
 1450  
 1451 Yan, C., Nie, W., Vogel, A. L., Dada, L., Lehtipalo, K., Stolzenburg, D. and Wagner, R.: Size-  
 1452 dependent influence of NO<sub>x</sub> on the growth rates of organic aerosol particles, , *Sci. Adv.*, 6, 1–10,  
 1453 2020.  
 1454  
 1455 Yan, C., Dada, L., Rose, C., Jokinen, T., Nie, W., Schobesberger, S., Junninen, H., Lehtipalo, K.,  
 1456 Sarnela, N., Makkonen, U., Garmash, O., Wang, Y., Zha, Q., Paasonen, P., Bianchi, F., Sipilä, M.,  
 1457 Ehn, M., Petäjä, T., Kerminen, V.-M., Worsnop, D. R. and Kulmala, M.: The role of H<sub>2</sub>SO<sub>4</sub>-  
 1458 NH<sub>3</sub> anion clusters in ion-induced aerosol nucleation mechanisms in the boreal forest, *Atmos.*  
 1459 *Chem. Phys.*, 18, 13231–13243, doi:10.5194/acp-18-13231-2018, 2018.  
 1460  
 1461 Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S. B.,  
 1462 Ehn, M., Paasonen, P., Sipilä, M., Wang, M., Wang, X., Xiao, S., Chen, H., Lu, Y., Zhang, B.,  
 1463 Wang, M., Chen, D., Xiao, M., Ye, Q., Stolzenburg, D., Hofbauer, V., Ye, P., Vogel, A. L.,  
 1464 Mauldin, R. L., Amorim, A., Baccarini, A., Baumgartner, B., Brilke, S., Dada, L., Dias, A.,  
 1465 Duplissy, J., Finkenzeller, H., Garmash, O., He, X. C., Hoyle, C. R., Kim, C., Kvashnin, A.,  
 1466 Lehtipalo, K., Fischer, L., Molteni, U., Petäjä, T., Pospisilova, V., Quéléver, L. L. J., Rissanen, M.,  
 1467 Simon, M., Tauber, C., Tomé, A., Wagner, A. C., Weitz, L., Volkamer, R., Winkler, P. M., Kirkby,  
 1468 J., Worsnop, D. R., Kulmala, M., Baltensperger, U., Dommen, J., El-Haddad, I. and Donahue, N.  
 1469 M.: Photo-oxidation of Aromatic Hydrocarbons Produces Low-Volatility Organic Compounds,  
 1470 *Environ. Sci. Technol.*, 54(13), 7911–7921, doi:10.1021/acs.est.0c02100, 2020.  
 1471  
 1472 Ye, J., Abbatt, J. P. D., Chan, A. W.H., Novel pathway of SO<sub>2</sub> oxidation in the atmosphere:  
 1473 reactions with monoterpene ozonolysis intermediates and secondary organic aerosol, *Atmos. Chem.*  
 1474 *Phys.*, 18, 5549–5565, 2018

1475 Yli-Juuti, T., Mohr, C. and Riipinen, I.: Open questions on atmospheric nanoparticle growth,  
1476 Commun. Chem., 3(1), 2–5, doi:10.1038/s42004-020-00339-4, 2020.  
1477  
1478 Zhang, R., Khalizov, A., Wang, L., Hu, M. and Xu, W.: Nucleation and growth of nanoparticles in  
1479 the atmosphere, Chem. Rev., 112(3), 1957–2011, doi:10.1021/cr2001756, 2012.  
1480  
1481

1482 **TABLE LEGENDS**

1483

1484 **Table 1:** Location and data availability of the sites.

1485

1486 **Table 2:** Frequency (and number of NPF events), growth and formation rate of NPF events.

1487

1488 **Table 3:** Normalised gradients (non-normalised for growth rate),  $R^2$  and p-values (- for values  
1489  $>0.05$ ) for the relationship between meteorological conditions and NPF event  
1490 variables.

1491

1492 **Table 4:** Normalised gradients (non-normalised for growth rate),  $R^2$  and p-values (- for values  
1493  $>0.05$ ) for the relationship between atmospheric composition variables and NPF  
1494 event variables.

1495

1496

1497 **FIGURE LEGENDS**

1498

1499 **Figure 1:** Map of the sites of the present study.

1500

1501 **Figure 2:** Relation of average downward incoming solar radiation ( $K_{\downarrow}$ ) and normalised  
1502 gradients  $a_N^*$ .

1503

1504 **Figure 3:** Normalised gradients  $a_J^*$  for  $K_{\downarrow}$  (\*UK sites are calculated with solar irradiance).

1505

1506 **Figure 4a:** Relationship of average relative humidity and normalised gradients  $a_N^*$ .

1507

1508 **Figure 4b:** Relationship of average relative humidity and normalised gradients  $a_N^*$  (SPAUB not  
1509 included).

1510

1511 **Figure 5:** Relationship of average temperature and normalised gradients  $a_N^*$ .

1512

1513 **Figure 6:** Normalised gradients  $a_J^*$  for temperature.

1514

1515 **Figure 7a:** Relationship of average  $SO_2$  concentrations and normalised gradients  $a_N^*$ .

1516

1517 **Figure 7b:** Relationship of average  $SO_2$  concentrations and normalised gradients  $a_N^*$  (UKRO  
1518 not included).

1519

1520 **Figure 8:** Relationship of average  $O_3$  concentrations and normalised gradients  $a_N^*$ .

**Table 1:** Location and data availability of the sites.

Site	Location	Available data	Meteorological data location	Data availability	Reference
<b>UKRU</b>	Harwell Science Centre, Oxford, 80 km W of London, UK (51° 34' 15" N; 1° 19' 31" W)	SMPS (16.6 - 604 nm, 76.5% availability), NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , OC, SO <sub>4</sub> <sup>2-</sup> , gaseous ammonia	On site	2009 - 2015	Charron et al., 2013
<b>UKUB</b>	North Kensington, 4 km W of London city centre, UK (51° 31' 15" N; 0° 12' 48" W)	SMPS (16.6 - 604 nm, 83.3% availability), NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , OC, SO <sub>4</sub> <sup>2-</sup>	Heathrow airport	2009 - 2015	Bigi and Harrison, 2010
<b>UKRO</b>	Marylebone Road, London, UK (51° 31' 21" N; 0° 9' 16" W)	SMPS (16.6 - 604 nm, 74.3% availability), NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , OC, SO <sub>4</sub> <sup>2-</sup>	Heathrow airport	2009 - 2015	Charron and Harrison, 2003
<b>DENRU</b>	Lille Valby, 25 km W of Copenhagen, (55° 41' 41" N; 12° 7' 7" E) (2008 – 6/2010) Risø, 7 km north of Lille Valby, (55° 38' 40" N; 12° 5' 19" E) (7/2010 – 2017)	DMPS and CPC (5.8 - 700 nm, 68.3% availability), NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , OC, SO <sub>4</sub> <sup>2-</sup>	H.C. Ørsted – Institute station	2008 – 2017	Ketzel et al., 2004
<b>DENUB</b>	H.C. Ørsted – Institute, 2 km NE of the city centre, Copenhagen, Denmark (55° 42' 1" N; 12° 33' 41" E)	DMPS and CPC (5.8 - 700 nm, 61.4% availability), NO <sub>x</sub> , O <sub>3</sub>	On site	2008 – 2017	Wang et al., 2010
<b>DENRO</b>	H.C. Andersens Boulevard, Copenhagen, Denmark (55° 40' 28" N; 12° 34' 16" E)	DMPS and CPC (5.8 - 700 nm, 65.7% availability), NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , OC, SO <sub>4</sub> <sup>2-</sup>	H.C. Ørsted – Institute station	2008 – 2017	Wang et al., 2010
<b>GERRU</b>	Melpitz, 40 km NE of Leipzig, Germany (51° 31' 31.85" N; 12° 26' 40.30" E)	TDMPS with CPC (4.8 - 800 nm, 87.2% availability), OC, SO <sub>4</sub> <sup>2-</sup>	On site	2008 – 2011	Birmili et al., 2016
<b>GERUB</b>	Tropos, 3 km NE from the city centre of Leipzig, Germany (51° 21' 9.1" N; 12° 26' 5.1" E)	TDMPS with CPC (3 - 800 nm, 90.4% availability)	On site	2008 – 2011	Birmili et al., 2016
<b>GERRO</b>	Eisenbahnstraße, Leipzig, Germany (51° 20' 43.80" N; 12° 24' 28.35" E)	TDMPS with CPC (4 - 800 nm, 68.3% availability)	Tropos station	2008 – 2011	Birmili et al., 2016
<b>FINRU</b>	Hyttiälä, 250 km N of Helsinki, Finland (61° 50' 50.70" N; 24° 17' 41.20" E)	TDMPS with CPC (3 – 1000 nm, 98.2% availability), NO <sub>x</sub> , SO <sub>2</sub> , O <sub>3</sub> , VOCs	On site	2008 – 2011 & 2015 – 2018	Aalto et al., 2001
<b>FINUB</b>	Kumpula Campus 4 km N of the city centre, Helsinki, Finland (60° 12' 10.52" N; 24° 57' 40.20" E)	TDMPS with CPC (3.4 - 1000 nm, 99.7% availability)	On site	2008 – 2011 & 2015 – 2018	Järvi et al., 2009
<b>FINRO</b>	Mäkelänkatu street, Helsinki, Finland (60° 11' 47.57" N; 24° 57' 6.01" E)	DMPS (6 - 800 nm, 90.0% availability), NO <sub>x</sub> , O <sub>3</sub>	Pasila station and on site	2015 – 2018	Hietikko et al., 2018
<b>SPARU</b>	Montseny, 50 km NNE from Barcelona, Spain (41° 46' 45" N; 2° 21' 29" E)	SMPS (9 – 856 nm, 53.7% availability), NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	On site	2012 - 2015	Dall'Osto et al., 2013
<b>SPAUB</b>	Palau Reial, Barcelona, Spain (41° 23' 14" N; 2° 6' 56" E)	SMPS (11 – 359 nm, 88.1% availability), NO <sub>2</sub> , SO <sub>2</sub> , O <sub>3</sub>	On site	2012 – 2015	Dall'Osto et al., 2012
<b>GRERU</b>	Finokalia, 70 km E of Heraklion, Greece (35° 20' 16.8" N; 25° 40' 8.4" E)	SMPS (8.77 - 849 nm, 85.0% availability), NO <sub>2</sub> , O <sub>3</sub> , OC	On site	2012 – 2018	Kalkavouras et al., 2017
<b>GREUB</b>	"Demokritos", 12 km NE from the city centre, Athens, Greece (37° 59' 41.96" N; 23° 48' 57.56" E)	SMPS (10 – 550 nm, 88.0% availability)	On site	2015 – 2018	Mølgaard et al., 2013

**Table 2:** Frequency (and number of NPF events), growth and formation rate of NPF events.

Site	Frequency of NPF events (%)	GR (nm h-1)	J <sub>10</sub> (N cm <sup>-3</sup> s <sup>-1</sup> )
UKRU	7.0 (160)	3.4*	8.69E-03**
UKUB	7.0 (156)	4.2*	1.42E-02**
UKRO	6.1 (120)	5.5*	3.75E-02**
DENRU	7.9 (176)	3.19	2.57E-02
DENUB	5.8 (116)	3.19	2.40E-02
DENRO	5.4 (117)	4.45	8.07E-02
GERRU	17.1 (164)	4.34	9.18E-02
GERUB	17.5 (169)	4.24	1.02E-01
GERRO	9.0 (62)	5.17	1.38E-01
FINRU	8.7 (190)	2.91	1.19E-02
FINUB	5.0 (110)	2.87	2.49E-02
FINRO	5.1 (49)	3.74	6.94E-02
SPARU	12 (68)	3.87	1.54E-02
SPAUB	13.1 (97)	3.71	2.12E-02
GRERU	6.5 (116)	3.68	4.90E-03
GREUB	8.5 (82)	3.4	4.41E-02

\* GR up to 50 nm calculated

\*\* J<sub>16</sub> calculated

1525

**Table 3:** Normalised gradients (non-normalised for growth rate),  $R^2$  and p-values (- for values  $>0.05$ ) for the relation between meteorological conditions and NPF event variables.

<b>Downward shortwave solar radiation <math>K_{\downarrow}</math> (<math>W\ m^{-2}</math>)</b>										
Site	$a_N^*$ ( $W^{-1}\ m^2$ )	$R^2$	p	$a_G$	$R^2$	p	$a_J^*$ ( $W^{-1}\ m^2$ )	$R^2$	p	Average
UKRU*	<b>1.21E-03</b>	0.94	<0.001	6.53E-05	0.11	-	<b>6.28E-04</b>	0.93	<0.001	443
UKUB*	<b>6.81E-04</b>	0.90	<0.001	-8.26E-05	0.10	-	1.49E-04	0.19	-	448
UKRO*	<b>8.69E-04</b>	0.98	<0.001	-7.75E-06	0.00	-	<b>2.66E-04</b>	0.64	<0.005	464
DENRU	<b>2.22E-03</b>	0.88	<0.001	4.24E-04	0.20	-	<b>1.38E-03</b>	0.64	<0.001	115
DENUB	<b>1.87E-03</b>	0.91	<0.001	1.47E-04	0.03	-	8.98E-04	0.48	<0.01	115
DENRO	<b>2.46E-03</b>	0.95	<0.001	1.27E-04	0.01	-	<b>6.77E-04</b>	0.50	<0.005	117
GERRU	<b>2.87E-03</b>	0.98	<0.001	<b>9.88E-04</b>	0.72	<0.01	<b>1.45E-03</b>	0.81	<0.001	130
GERUB	<b>3.18E-03</b>	0.97	<0.001	<b>7.28E-04</b>	0.51	<0.005	<b>1.53E-03</b>	0.69	<0.001	114
GERRO	<b>2.40E-03</b>	0.95	<0.001	-5.89E-04	0.09	-	<b>9.95E-04</b>	0.59	<0.005	114
FINRU	<b>2.63E-03</b>	0.76	<0.001	<b>1.01E-03</b>	0.57	<0.01	<b>2.04E-03</b>	0.82	<0.001	91.5
FINUB	1.38E-03	0.37	-	1.81E-04	0.08	-	8.99E-04	0.25	-	111
FINRO	<b>1.76E-03</b>	0.59	<0.005	9.15E-04	0.34	<0.005	4.45E-04	0.03	-	114
SPARU	3.46E-04	0.35	<0.05	5.68E-04	0.13	-	<b>1.97E-03</b>	0.74	<0.001	162
SPAUB	<b>5.92E-04</b>	0.58	<0.05	6.98E-04	0.23	-	<b>1.58E-03</b>	0.81	<0.001	180
GRERU	<b>4.10E-04</b>	0.52	<0.001	<b>7.14E-04</b>	0.55	<0.001	-6.30E-04	0.05	-	201
GREUB	3.49E-04	0.31	-	-1.10E-04	0.02	-	8.97E-04	0.34	<0.05	183

\* Global solar irradiation measurements in  $kJ\ m^{-2}$

<b>Relative Humidity (%)</b>										
Site	$a_N^*$ ( $\%^{-1}$ )	$R^2$	p	$a_G$	$R^2$	p	$a_J^*$ ( $\%^{-1}$ )	$R^2$	p	Average
UKRU	<b>-5.89E-02</b>	0.85	<0.001	1.69E-03	0.02	-	<b>-3.35E-02</b>	0.85	<0.001	79.7
UKUB	<b>-3.42E-02</b>	0.94	<0.001	8.23E-03	0.24	-	-5.66E-03	0.19	-	75.3
UKRO	<b>-5.09E-02</b>	0.85	<0.001	7.03E-03	0.25	-	-1.49E-02	0.46	<0.05	74.5
DENRU	<b>-3.90E-02</b>	0.95	<0.001	<b>9.42E-03</b>	0.74	<0.001	5.45E-04	0.00	-	75.7
DENUB	<b>-3.14E-02</b>	0.94	<0.001	3.64E-03	0.06	-	2.57E-03	0.00	-	75.7
DENRO	<b>-3.64E-02</b>	0.95	<0.001	-1.21E-02	0.22	-	-3.91E-03	0.10	-	75.7
GERRU	<b>-5.08E-02</b>	0.88	<0.001	<b>-1.30E-02</b>	0.72	<0.001	<b>-2.46E-02</b>	0.91	<0.001	81.9
GERUB	<b>-5.35E-02</b>	0.86	<0.001	<b>-6.34E-03</b>	0.67	<0.001	<b>-2.25E-02</b>	0.86	<0.001	78.7
GERRO	<b>-2.83E-02</b>	0.90	<0.001	3.98E-03	0.05	-	<b>-1.72E-02</b>	0.81	<0.001	78.7
FINRU	<b>-4.48E-02</b>	0.94	<0.001	<b>-7.07E-03</b>	0.65	<0.001	<b>-2.16E-02</b>	0.87	<0.001	80.1
FINUB	<b>-5.89E-02</b>	0.95	<0.001	1.04E-02	0.26	-	-6.52E-03	0.18	-	76.5
FINRO	<b>-3.34E-02</b>	0.92	<0.001	-1.47E-03	0.01	-	7.39E-03	0.10	-	71.1
SPARU	<b>-1.54E-02</b>	0.90	<0.001	-4.67E-03	0.08	-	-7.12E-03	0.14	-	66.4
SPAUB	<b>-4.84E-02</b>	0.93	<0.001	<b>2.43E+02</b>	0.50	<0.01	-9.83E-03	0.19	-	69.2
GRERU	-7.72E-03	0.22	-	1.06E-02	0.06	-	-1.83E-01	0.15	-	70.0
GREUB	<b>-1.42E-02</b>	0.62	<0.001	2.83E-03	0.06	-	4.85E-04	0.00	-	60.5



1530

Temperature (°C)										
Site	a <sub>N</sub> * (°C <sup>-1</sup> )	R <sup>2</sup>	P	a <sub>G</sub>	R <sup>2</sup>	P	a <sub>J</sub> * (°C <sup>-1</sup> )	R <sup>2</sup>	P	Average
UKRU	<b>1.10E-01</b>	0.93	<0.001	<b>7.85E-02</b>	0.94	<0.001	<b>8.72E-02</b>	0.84	<0.001	10.6
UKUB	<b>9.04E-02</b>	0.98	<0.001	<b>1.39E-01</b>	0.96	<0.001	<b>6.34E-02</b>	0.73	<0.005	11.8
UKRO	<b>8.22E-02</b>	0.98	<0.001	<b>3.51E-02</b>	0.52	<0.05	4.32E-02	0.44	<0.05	12.1
DENRU	<b>6.68E-02</b>	0.83	<0.001	1.54E-02	0.08	-	<b>6.68E-02</b>	0.92	<0.001	9.80
DENUB	<b>2.50E-02</b>	0.45	<0.05	2.40E-02	0.33	-	3.05E-02	0.45	<0.05	9.82
DENRO	<b>6.64E-02</b>	0.88	<0.001	3.51E-03	0.00	-	<b>2.96E-02</b>	0.58	<0.005	10.0
GERRU	<b>7.27E-02</b>	0.92	<0.001	<b>5.65E-02</b>	0.92	<0.001	<b>5.37E-02</b>	0.93	<0.001	10.3
GERUB	<b>8.20E-02</b>	0.93	<0.001	<b>3.38E-02</b>	0.62	<0.001	<b>4.28E-02</b>	0.54	<0.005	11.1
GERRO	<b>5.08E-02</b>	0.89	<0.001	-3.33E-03	0.00	-	1.61E-02	0.11	-	11.1
FINRU	-2.01E-02	0.17	-	<b>1.13E-01</b>	0.79	<0.001	<b>4.27E-02</b>	0.72	<0.001	4.79
FINUB	-4.21E-03	0.00	-	<b>7.42E-02</b>	0.83	<0.001	1.67E-02	0.28	-	6.52
FINRO	<b>6.24E-02</b>	0.65	<0.005	<b>9.28E-02</b>	0.87	<0.001	-1.09E-02	0.05	-	7.72
SPARU	-2.51E-02	0.41	<0.05	<b>1.23E-01</b>	0.92	<0.001	<b>9.11E-02</b>	0.71	<0.001	13.9
SPAUB	-3.43E-03	0.02	-	<b>6.67E-02</b>	0.66	<0.005	1.18E-02	0.08	-	18.2
GRERU	<b>-4.66E-02</b>	0.75	<0.001	<b>1.74E-01</b>	0.75	<0.001	-9.45E-02	0.47	<0.05	18.2
GREUB	-1.00E-02	0.25	-	<b>4.67E-02</b>	0.62	<0.005	-2.85E-02	0.20	-	17.6

Wind Speed (m s <sup>-1</sup> )										
Site	a <sub>N</sub> * (m <sup>-1</sup> s)	R <sup>2</sup>	P	a <sub>G</sub>	R <sup>2</sup>	P	a <sub>J</sub> * (m <sup>-1</sup> s)	R <sup>2</sup>	P	Average
UKRU	5.72E-02	0.20	-	-3.04E-02	0.07	-	6.87E-03	0.00	-	3.96
UKUB	<b>1.72E-01</b>	0.87	<0.001	<b>-1.91E-01</b>	0.71	<0.001	3.56E-03	0.00	-	4.16
UKRO	6.34E-02	0.19	-	3.21E-02	0.02	-	7.28E-02	0.45	<0.005	4.14
DENRU	<b>1.08E-01</b>	0.88	<0.001	<b>-2.33E-01</b>	0.74	<0.001	1.28E-01	0.44	<0.01	4.17
DENUB	<b>1.50E-01</b>	0.90	<0.001	-3.33E-02	0.10	-	8.31E-02	0.19	-	4.17
DENRO	<b>1.65E-01</b>	0.89	<0.001	-1.51E-01	0.49	<0.001	9.08E-03	0.00	-	4.16
GERRU	<b>-1.06E-01</b>	0.57	<0.005	<b>-2.26E-01</b>	0.83	<0.001	-5.32E-03	0.00	-	2.58
GERUB	<b>-1.27E-01</b>	0.52	<0.01	<b>-1.41E-01</b>	0.60	<0.005	-3.32E-02	0.04	-	2.33
GERRO	<b>-2.40E-01</b>	0.56	-	-2.54E-01	0.38	-	-1.30E-01	0.22	-	2.33
FINRU	<b>1.62E-01</b>	0.63	<0.005	-1.29E-01	0.16	<0.05	7.99E-02	0.07	-	1.31
FINUB	-3.17E-02	0.08	-	7.26E-02	0.20	<0.05	-9.74E-02	0.17	-	3.43
FINRO	<b>8.62E-02</b>	0.51	<0.05	-1.60E-01	0.32	<0.05	-1.86E-01	0.32	-	4.26
SPARU	-2.20E-02	0.02	-	3.80E-01	0.31	-	5.74E-02	0.02	-	0.94
SPAUB	<b>2.90E-01</b>	0.93	<0.001	7.71E-02	0.24	-	-5.90E-02	0.05	-	2.05
GRERU	<b>4.37E-02</b>	0.54	<0.001	1.01E-01	0.36	<0.005	1.73E-03	0.00	-	6.06
GREUB	-1.13E-01	0.47	<0.01	<b>-1.88E-01</b>	0.50	<0.005	-3.78E-02	0.01	-	1.87

Atmospheric Pressure (mbar)										
Site	$a_N^*$ (mbar <sup>-1</sup> )	R <sup>2</sup>	P	$a_G$	R <sup>2</sup>	P	$a_I^*$ (mbar <sup>-1</sup> )	R <sup>2</sup>	P	Average
UKRU	<b>4.26E-02</b>	0.83	<0.005	<b>3.93E-02</b>	0.58	<0.005	2.95E-02	0.47	<0.05	1007.7
UKUB	<b>1.90E-02</b>	0.50	-	1.17E-02	0.05	<0.05	4.16E-03	0.04	-	1011.7
UKRO	<b>6.33E-02</b>	0.95	<0.001	-1.21E-01	0.40	-	-2.98E-02	0.17	-	1012
GERRU	<b>5.10E-02</b>	0.97	-	<b>8.95E-02</b>	0.85	<0.001	2.16E-02	0.21	-	1007.0
GERUB	<b>6.27E-02</b>	0.97	-	<b>4.00E-02</b>	0.76	-	2.00E-02	0.37	<0.05	995.5
GERRO	<b>4.57E-02</b>	0.79	-	-9.61E-02	0.43	-	-2.80E-02	0.21	-	995.5
FINRU	<b>3.46E-02</b>	0.88	<0.001	<b>2.90E-02</b>	0.57	<0.001	1.05E-02	0.14	-	985.1
FINUB	<b>2.61E-02</b>	0.55	<0.005	-3.57E-03	0.02	-	4.38E-03	0.05	-	1004.4
FINRO	<b>4.91E-02</b>	0.70	-	-2.67E-02	0.17	-	1.43E-02	0.26	-	1008.8
SPARU	-2.02E-02	0.09	-	4.79E-02	0.14	-	2.89E-02	0.08	-	939.3
SPAUB	-2.83E-02	0.44	<0.05	1.86E-02	0.08	-	1.68E-02	0.21	-	1006.3
GRERU	6.00E-02	0.46	<0.001	-1.50E-01	0.73	-	8.14E-02	0.33	-	1014.5
GREUB	9.42E-03	0.10	<0.05	-1.00E-01	0.71	-	1.58E-02	0.04	-	1015.7

**Table 4:** Normalised gradients (non-normalised for growth rate), R<sup>2</sup> and p-values (- for values >0.05) for the relation between atmospheric composition variables and NPF event variables.

SO <sub>2</sub> (µg m <sup>-3</sup> )										
Site	a <sub>N</sub> * (µg <sup>-1</sup> m <sup>3</sup> )	R <sup>2</sup>	p	a <sub>G</sub>	R <sup>2</sup>	p	a <sub>J</sub> * (µg <sup>-1</sup> m <sup>3</sup> )	R <sup>2</sup>	p	Average
UKRU	-1.97E-01	0.38	<0.05	-6.17E-02	0.02	-	3.30E-01	0.06	-	1.64
UKUB	<b>-2.57E-01</b>	0.62	<0.001	1.93E-02	0.00	-	4.18E-01	0.40	-	2.04
UKRO	<b>-1.03E-01</b>	0.82	<0.001	6.90E-02	0.34	<0.01	<b>8.43E-02</b>	0.77	<0.001	7.46
DENRU	<b>-9.77E-01</b>	0.53	<0.05	2.84E+00	0.37	-	4.38E-01	0.09	-	0.52
DENRO	<b>-4.20E-01</b>	0.91	<0.001	<b>6.42E-01</b>	0.54	<0.005	<b>5.66E-01</b>	0.62	<0.001	0.97
FINRU	-5.66E-01	0.05	-	-1.42E+00	0.19	-	-6.30E-02	0.00	-	0.09
SPARU	<b>-3.62E-01</b>	0.74	<0.001	-1.33E-01	0.02	-	-3.55E-02	0.01	-	0.95
SPAUB	-2.93E-02	0.04	-	<b>4.12E-01</b>	0.59	-	1.07E-01	0.29	-	1.99

NO <sub>x</sub> or NO <sub>2</sub> (ppb)										
Site	a <sub>N</sub> * (ppb <sup>-1</sup> )	R <sup>2</sup>	p	a <sub>G</sub>	R <sup>2</sup>	p	a <sub>J</sub> * (ppb <sup>-1</sup> )	R <sup>2</sup>	p	Average
UKRU	<b>-4.99E-02</b>	0.67	<0.005	<b>4.52E-02</b>	0.58	<0.05	<b>-4.51E-02</b>	0.70	<0.005	11.7
UKUB	<b>-8.75E-03</b>	0.83	<0.001	-3.97E-04	0.00	-	-1.09E-02	0.43	<0.05	53.6
UKRO	<b>-3.22E-03</b>	0.72	<0.001	1.44E-03	0.39	<0.05	<b>2.19E-03</b>	0.66	<0.001	299
DENRU	-9.41E-02	0.43	<0.005	-4.89E-03	0.00	<0.001	<b>-6.47E-02</b>	0.55	<0.01	5.42
DENUB	<b>-4.99E-02</b>	0.68	<0.001	2.85E-02	0.26	-	8.55E-04	0.00	-	10.5
DENRO	<b>-5.10E-03</b>	0.75	<0.001	<b>1.10E-02</b>	0.69	<0.001	<b>8.33E-03</b>	0.88	<0.001	68.5
FINRU	<b>-7.27E-01</b>	0.54	<0.001	-2.74E-01	0.11	-	1.95E-01	0.05	-	0.72
FINRO	<b>-6.24E-03</b>	0.68	<0.001	1.70E-03	0.12	-	3.25E-03	0.03	-	88.1
SPARU*	-1.53E-02	0.05	-	2.54E-02	0.01	-	1.25E-01	0.21	-	3.26
SPAUB*	<b>-2.59E-02</b>	0.62	<0.005	<b>2.23E-02</b>	0.70	<0.001	2.57E-03	0.01	-	31.4
GRERU*	3.01E-01	0.19	-	<b>-1.40E+00</b>	0.75	<0.001	5.23E-01	0.13	-	0.52

\* NO<sub>2</sub> measurements

O <sub>3</sub> (ppb)										
Site	a <sub>N</sub> * (ppb <sup>-1</sup> )	R <sup>2</sup>	p	a <sub>G</sub>	R <sup>2</sup>	p	a <sub>J</sub> * (ppb <sup>-1</sup> )	R <sup>2</sup>	p	Average
UKRU	<b>2.27E-02</b>	0.88	<0.001	<b>-4.89E-02</b>	0.53	<0.005	-3.53E-03	0.01	-	54.4
UKUB	<b>1.37E-02</b>	0.87	<0.001	<b>-3.45E-02</b>	0.68	<0.001	-5.95E-03	0.05	-	39.3
UKRO	<b>7.46E-02</b>	0.95	<0.001	-1.06E-02	0.09	-	<b>-2.44E-02</b>	0.63	<0.005	16.2
DENRU	<b>4.97E-02</b>	0.92	<0.001	-1.32E-02	0.15	-	1.23E-02	0.08	-	30.1
DENUB	<b>5.85E-02</b>	0.84	<0.001	<b>-1.69E-02</b>	0.58	-	2.77E-02	0.32	<0.05	28.2
DENRO	<b>6.42E-02</b>	0.51	<0.05	1.39E-02	0.03	-	<b>3.24E-02</b>	0.91	<0.05	31.1
FINRU	<b>6.76E-02</b>	0.77	<0.05	<b>-4.23E-02</b>	0.60	-	3.92E-02	0.37	<0.05	27.4
FINRO	<b>2.38E-02</b>	0.91	<0.001	6.11E-03	0.24	-	-1.83E-02	0.29	-	37.1
SPARU	1.57E-02	0.02	-	4.34E-02	0.11	-	1.31E-02	0.31	-	75.9
SPAUB	7.99E-03	0.38	<0.05	-5.83E-03	0.30	-	-1.13E-03	0.01	-	54.9
GRERU	7.55E-03	0.04	-	3.68E-02	0.17	-	-3.01E-02	0.15	-	49.5

Particulate Organic Carbon (µg m <sup>-3</sup> )										
Site	a <sub>N</sub> * (µg <sup>-1</sup> m <sup>3</sup> )	R <sup>2</sup>	p	a <sub>G</sub>	R <sup>2</sup>	p	a <sub>J</sub> * (µg <sup>-1</sup> m <sup>3</sup> )	R <sup>2</sup>	p	Average
UKRU	-3.30E-02	0.00	-	1.13E+00	0.42	<0.005	2.13E-01	0.16	-	1.96
UKUB	<b>-2.76E-01</b>	0.59	<0.005	<b>6.63E-01</b>	0.58	<0.05	<b>2.19E-01</b>	0.55	<0.05	3.63
UKRO	<b>-3.78E-01</b>	0.89	<0.001	<b>8.12E-01</b>	0.57	<0.005	<b>4.60E-01</b>	0.75	<0.001	6.24
DENRU	<b>-4.44E-01</b>	0.75	<0.001	2.24E-01	0.11	-	<b>-3.17E-01</b>	0.68	<0.01	1.48
DENRO	-7.80E-02	0.11	-	<b>1.10E+00</b>	0.77	<0.005	<b>4.02E-01</b>	0.81	<0.005	2.59
GERRU	-1.26E-01	0.24	-	1.35E-01	0.09	-	3.14E-02	0.03	-	2.18
FINRU	2.27E-02	0.00	-	<b>3.39E-01</b>	0.60	<0.005	-3.46E-01	0.16	-	1.78
GRERU	-2.08E-01	0.11	-	7.87E-01	0.41	<0.05	8.94E-01	0.11	-	1.58

Sulphate (µg m <sup>-3</sup> )										
Site	a <sub>N</sub> * (µg <sup>-1</sup> m <sup>3</sup> )	R <sup>2</sup>	p	a <sub>G</sub>	R <sup>2</sup>	p	a <sub>J</sub> * (µg <sup>-1</sup> m <sup>3</sup> )	R <sup>2</sup>	p	Average
UKRU <sup>1</sup>	<b>-2.62E-01</b>	0.57	<0.001	<b>7.34E-01</b>	0.77	<0.001	7.99E-01	0.44	<0.05	1.97
UKUB <sup>1</sup>	<b>-3.57E-01</b>	0.89	<0.001	9.28E-01	0.44	<0.01	9.72E-01	0.16	-	1.58
UKRO <sup>1</sup>	-6.05E-02	0.24	-	3.04E-01	0.34	<0.05	-6.22E-02	0.04	-	1.98
DENRU <sup>2</sup>	-7.81E-01	0.34	<0.05	<b>1.02E+00</b>	0.60	<0.05	<b>-1.03E+00</b>	0.63	<0.01	0.52
DENRO <sup>2</sup>	-8.23E-01	0.28	-	1.99E+00	0.22	-	2.82E-01	0.12	-	0.55
GERRU <sup>1</sup>	-3.37E-02	0.00	-	5.89E-01	0.11	-	-4.89E-02	0.01	-	0.92
FINRU <sup>3</sup>	<b>-1.18E+00</b>	0.65	<0.001	2.35E-01	0.09	-	-2.53E-01	0.17	-	1.02


<sup>1</sup> Measurements in PM<sub>10</sub>

<sup>2</sup> Measurements in PM<sub>2.5</sub>


<sup>3</sup> Measurements in PM<sub>1</sub>

1550


Condensation Sink (s <sup>-1</sup> )										
Site	a <sub>N</sub> * (s)	R <sup>2</sup>	P	a <sub>G</sub>	R <sup>2</sup>	P	a <sub>J</sub> * (s)	R <sup>2</sup>	P	Average
UKRU	-2.28E+02	0.72	<0.001	2.64E+02	0.60	<0.001	7.58E+01	0.22	-	3.38E-03
UKUB	-1.66E+02	0.78	<0.001	2.49E+02	0.41	<0.05	1.73E+02	0.35	<0.05	7.41E-03
UKRO	-4.03E+01	0.75	<0.001	2.33E+01	0.18	-	8.94E+01	0.91	<0.001	2.12E-02
DENRU	-4.48E+01	0.91	<0.001	6.90E+01	0.49	<0.05	5.37E+01	0.24	-	9.46E-03
DENUB	-3.78E+01	0.75	<0.001	3.58E+01	0.25	-	1.55E+01	0.56	<0.005	1.42E-02
DENRO	-1.06E+01	0.73	<0.001	2.53E+01	0.56	<0.005	2.72E+01	0.79	<0.001	3.10E-02
GERRU	1.54E+02	0.86	<0.001	1.33E+02	0.56	<0.001	6.67E+01	0.63	<0.001	7.02E-03
GERUB	3.59E+01	0.56	<0.005	3.63E+01	0.17	-	4.74E+01	0.75	<0.001	9.11E-03
GERRO	3.89E+01	0.22	<0.05	-2.21E+01	0.03	<0.005	3.54E+01	0.45	<0.005	1.20E-02
FINRU	-1.80E+02	0.59	<0.005	4.01E+02	0.74	<0.001	4.98E+01	0.10	-	2.32E-03
FINUB	-1.51E+02	0.63	<0.005	8.14E+01	0.31	-	2.01E+02	0.41	<0.05	6.34E-03
FINRO	-6.99E+01	0.77	<0.001	-1.56E+01	0.05	-	2.42E+02	0.83	<0.001	8.96E-03
SPARU	-2.15E+02	0.65	<0.005	1.86E+01	0.00	-	8.60E+01	0.47	<0.05	5.49E-03
SPAUB	-1.18E+02	0.65	<0.005	3.74E+01	0.38	<0.05	9.51E+01	0.52	<0.01	1.00E-02
GRERU	4.33E+00	0.00	-	2.86E+02	0.70	<0.001	1.77E+02	0.56	<0.005	4.66E-03
GREUB	1.64E+02	0.65	<0.001	9.31E+01	0.28	<0.05	1.73E+02	0.83	<0.001	7.55E-03




**Figure 1:** Map of the sites of the present study.




**Figure 2:** Relationship of average downward incoming solar radiation ( $K\downarrow$ ) and normalised gradients  $a_N^*$ .



**Figure 3:** Normalised slopes  $a_j^*$  for  $K\downarrow$  (\*UK sites are calculated with solar irradiance).




1565 **Figure 4:** Relationship of average relative humidity and normalised gradients  $a_N^*$ .




**Figure 5:** Relationship of average temperature and normalised gradients  $a_N^*$ .




1570




**Figure 6:** Normalised gradients  $a_j^*$  for temperature.



**Figure 7a:** Relationship of average  $\text{SO}_2$  concentrations and normalised gradients  $a_N^*$ .



**Figure 7b:** Relationship of average  $\text{SO}_2$  concentrations and normalised gradients  $a_N^*$  (UKRO not included).



**Figure 8:** Relationship of average  $O_3$  concentrations and normalised gradients  $a_N^*$ .