

Changes in satellite retrievals of atmospheric composition over eastern China during the 2020 COVID-19 lockdowns

Robert D. Field^{1,2}, Jonathan E. Hickman¹, Igor V. Geogdzhayev^{1,2}, Kostas Tsigaridis^{1,3}, Susanne E. Bauer¹

¹NASA Goddard Institute for Space Studies, 2880 Broadway, New York, NY, USA, 10025

²Dept. of Applied Physics and Applied Mathematics, Columbia University 2880 Broadway, New York, NY, USA, 10025

³Center for Climate Systems Research, Columbia University, 2880 Broadway, New York, NY, USA, 10025

Correspondence to: Robert D. Field (robert.field@columbia.edu)

Abstract. We examined daily Level-3 satellite retrievals of AIRS CO, OMI SO₂ and NO₂, and MODIS AOD over eastern China to understand how COVID-19 lockdowns affected atmospheric composition. Changes in 2020 were strongly dependent on the choice of background period since 2005 and whether trends in atmospheric composition were accounted for. Over central east China during the January 23 - April 8 lockdown window, CO in 2020 was between 3% and 12% lower than average depending on the background period. 2020 CO was not consistently less than expected from trends beginning between 2005 and 2016 and ending in 2019 but was 3-4% lower than the background mean during the 2017-2019 period when CO changes had flattened. Similarly for AOD, 2020 was between 14% and 30% lower than averages beginning in 2005 and 14-17% lower compared to different background means beginning in 2016. NO₂ in 2020 was between 30% and 43% lower than the mean over different background periods and between 17% and 33% lower than what would be expected for trends beginning later than 2011. Relative to the 2016-2019 period when NO₂ had flattened, 2020 was 30-33% lower. Over southern China, 2020 NO₂ was between 23% and 27% lower than different background means beginning in 2013, the beginning of a period of persistently lower NO₂. CO over southern China was significantly higher in 2020 than what would be expected, which we suggest was partly because of an active fire season in neighbouring countries. Over central east and southern China, 2020 SO₂ was higher than expected, but this depended strongly on how daily regional values were calculated from individual retrievals and reflects background values approaching the retrieval detection limit. Future work over China, or other regions, needs to take into account the sensitivity of differences in 2020 to different background periods and trends in order to separate the effects of COVID-19 on air quality from previously occurring changes, or from variability in other sources.

1 Introduction

In an effort to control the spread of COVID-19, the Chinese government implemented a range of restrictions on movement. These led to reductions in industrial and other work related and personal activities starting January 23, 2020 in Wuhan, Hubei province, then extending to other cities and regions in the days that followed. On April 8, 2020, Wuhan was the last city to re-open after a complete lockdown that prevented most people from leaving their homes. These measures have been linked to changes in air quality. A network of surface monitoring stations in northern China observed 35% decreases in PM_{2.5} and 60% decreases in NO₂ concentrations during January 29 through February 29, as compared to the preceding three weeks; CO and SO₂ also declined (Shi and Brasseur, 2020). In and around Wuhan, decreases of NO₂ and PM_{2.5} were similar to regional changes, but there was a slight

39 increase in SO₂ concentrations (Shi and Brasseur, 2020). Observations by the Tropospheric Monitoring Instrument
40 (TROPOMI) showed large decreases in tropospheric NO₂ column densities over Chinese cities, on the order of
41 40% for February 11 to March 24 2020 compared to the same period in 2019, ranging from roughly 25% for cities
42 not affected by lockdown to 60% for Wuhan and Xi'an (Bauwens et al., 2020). Prospective simulations suggested
43 that meteorology may limit the effect of reduced emissions on PM_{2.5} concentrations, with Chinese cities
44 experiencing less than 20% reductions (Wang et al., 2020).

45

46 The goal of our study was to consider these changes against pollution trends in China using NASA Earth
47 Observing System data by combining several products to give a holistic view covering several emission sectors
48 that are responsible for the observed changes. Over the last 2 to 3 decades, air pollution in China appears to have
49 followed the pattern described by the Environmental Kuznets Curve (Selden and Song, 1994). This framework
50 describes a relationship in which economic growth is initially accompanied by an increase in air pollution, when
51 poverty remains widespread. But as growth continues, air pollution is expected to level off and decline as a
52 consequence of changes in social awareness of environmental degradation and the economic, political, and
53 technological capacity to limit it (Sarkodie and Strezov, 2019; Selden and Song, 1994).

54

55 Bottom-up and top-down assessments of air pollutant emissions and concentrations suggest that China has
56 followed this pattern during the era of satellite monitoring of atmospheric composition, with concentrations of
57 SO₂, NO₂, CO, and aerosol optical depth (AOD) mostly exhibiting marked and steady declines over the last
58 decade. In the case of NO₂, multi-instrument analyses, which extend the observational record beyond the lifetime
59 of a single instrument, depict a consistent regional picture of NO₂ trends in China since 1996 (Geddes et al.,
60 2016; Georgoulias et al., 2019; Wang and Wang, 2020; Xu et al., 2020). Column totals show an increasing trend
61 during the first part of the satellite record, but this trend is reversed sometime between 2010 and 2014 (Georgoulias
62 et al., 2019; Krotkov et al., 2016; Lin et al., 2019; Xu et al., 2020; Si et al., 2019; Shah et al., 2020). The trend reversal
63 has been attributed to a combination of emission control measures (Zheng et al., 2018a) and variations in economic
64 growth (Krotkov et al., 2016).

65

66 Bottom-up estimates suggest that SO₂ emissions peaked earlier, with declines starting around 2005, primarily as
67 a result of power and industrial pollution control measures as well as the elimination of small industrial boilers
68 (Sun et al., 2018; Zheng et al., 2018b). An earlier peak in SO₂ emissions is consistent with observations by multiple
69 satellite instruments, which revealed declines in SO₂ column densities since 2005 (Fioletov et al., 2016; Krotkov
70 et al., 2016; Wang and Wang, 2020; Zhang et al., 2017; Si et al., 2019).

71

72 AOD retrievals from the Along Track Scanning Radiometer instruments show a steady increase over southeastern
73 China from 1995 to 2005 (Sogacheva et al., 2020), and a decline since 2005 in the MODIS AOD (He et al., 2019).
74 The AOD peak has been argued to match either the ~2011 peak in NO₂ (Zheng et al., 2018b; Xie et al., 2019), the
75 ~2005 peak of SO₂, or to have occurred at some point in between (Ma et al., 2016), with more rapid decreases in
76 AOD after 2011 (Lin et al., 2018). The recent decrease in AOD is also seen in VIIRS retrievals (Sogacheva et al.,
77 2020). Most mitigation of direct PM_{2.5} emissions since 2010 was by industry, with residential emissions also
78 decreasing substantially (Zheng et al., 2018b). The decline in SO₂ emissions also exerted an important influence,

79 with the sulfate concentration of PM_{2.5} decreasing substantially between 2013 and 2017 (Shao et al., 2018),
80 reflecting the negative trend in SO₂ emissions.

81

82 The peak in concentrations of CO, which has an atmospheric lifetime ranging from weeks to months, is less easily
83 identified. Some studies suggest that trends have been negative potentially throughout the 21st century (Han et al.,
84 2018;Strode et al., 2016;Wang et al., 2018;Yumimoto et al., 2014;Zheng et al., 2018a), but others suggest that
85 emissions and/or column densities were increasing or flat during at least the first decade of the century (Sun et
86 al., 2018;Zhao et al., 2013;Zhao et al., 2012). The negative trend has been attributed largely to reductions in
87 emissions from industrial activity, as well as from residential and transportation sectors (Zheng et al.,
88 2018a;Zheng et al., 2018b).

89

90 In addition to these long-term trends, a number of air pollutants also exhibit strong seasonal variation in China.
91 Anthropogenic emissions of CO, SO₂, and PM_{2.5} are highest in winter, reflecting large variation in emissions from
92 the residential sector and, in the case of CO, increased emissions associated with cold-start processes in the
93 transportation sector (Li et al., 2017). Outflow of CO and AOD has a spring maximum, resulting from transport
94 of pollution, dust, and boreal biomass burning emissions (Han et al., 2018;Luan and Jaegle, 2013).

95

96 Changes in pollution over China have also come from short-term interventions. To improve air quality for the
97 2008 summer Olympics—a time when emissions in China were high and still increasing—the Chinese
98 government imposed a series of strict emissions control measures from July through September 21, 2008, which
99 were qualitatively similar to the emissions reductions expected to have accompanied the COVID-19 lockdown
100 (UNEP, 2009). As a result, NO₂ concentrations over Beijing were estimated to have declined by between 40%
101 and 60% based on satellite observations, with substantial but smaller reductions in surrounding cities often on the
102 order of 20% to 30% compared to previous years (Mijling et al., 2009;Witte et al., 2009). Regional reductions of
103 SO₂ and CO during the months of the games were estimated to be 13% and 19%, respectively (Witte et al., 2009).
104 These results are broadly consistent with on-road observations (Wang et al., 2009), but larger than some surface
105 observations comparing concentrations before and after the emission control measures were implemented (Wang
106 et al., 2010).

107

108 The COVID-related lockdowns provide a similar natural experiment to the 2008 Beijing Olympics but on the
109 other side of the Kuznets curve. The fact that the lockdowns occurred during years of decreasing air pollution
110 needs to be taken into account in attributing changes in atmospheric composition to COVID-19 lockdowns,
111 independent of the long-term trend. Following Chen et al.'s (2020) analysis of air quality improvements on
112 mortality which controlled for changes in air quality since 2016, in this study we determine whether changes in
113 2020 in satellite retrievals of CO, SO₂, NO₂ and AOD departed significantly from the expected declines associated
114 with the long-term decreases in concentrations resulting from pollution controls and technological change.

115 **2 Data and methods**

116 We used daily Level-3 (L3) retrievals from four different instruments on three different NASA Earth Observing
117 System satellites. The Atmospheric Infrared Sounder (AIRS) instrument aboard NASA's Aqua satellite is a 2300-

118 channel infrared grating spectrometer in a sun-synchronous orbit with northward equator crossing time of 1:30
119 PM. AIRS carbon monoxide (CO) profiles are retrieved with horizontal resolution of 45 km at nadir, in a swath
120 of width 30 fields-of-view or about 1600 km. The retrieval uses a cloud-clearing methodology providing CO with
121 sensitivity that peaks around 500 hPa, with ~0.8-1.2 degrees-of-freedom-of-signal for 50-70% of scenes. More
122 sampling and higher information content is obtained in clear scenes (Warner et al., 2013). We used the daily
123 version 6 (AIRS3STD.006) product.

124

125 The Ozone Monitoring Instrument (OMI) aboard NASA's Aura satellite was launched in July 2004, and has a
126 local equator-crossing time of roughly 13:45. OMI is a nadir-viewing spectrometer, which measures solar
127 backscatter in the UV-visible range (Krotkov, 2013). We used NASA's L3 tropospheric NO₂ column density
128 Standard Product v3 (OMNO2d_003), and the OMI Principal Components Analysis Planetary Boundary Layer
129 (PBL) SO₂ product (OMSO2e_003), which grid retrievals to 0.25° resolution (Krotkov et al., 2017; Li et al., 2013).
130 Both products are cloud-screened; only pixels that are at least 70% cloud-free are included in the NO₂ product,
131 and those that are at least 80% cloud-free are included in the SO₂ product. The NO₂ product relies on air mass
132 factors (AMFs) calculated with the assistance of an atmospheric chemical transport model and are sensitive to
133 model representations of emission, chemistry, and transport data. Instead of AMFs, the SO₂ product uses
134 spectrally-dependent SO₂ Jacobians, but can be interpreted as having a fixed AMF that is representative of
135 summertime conditions. We applied basic transient SO₂ plume filtering, excluding retrievals with SO₂ > 15 DU
136 (Wang and Wang, 2020).

137

138 Because our trend analysis uses a seasonal mean as the response variable, we assume that random errors cancel
139 out, leaving only systematic errors, which do not contribute to uncertainty in the trend analysis. Systematic errors
140 in the OMI NO₂ product have an uncertainty of 20% (McLinden et al., 2014) and are associated with AMFs and
141 tropospheric vertical column contents. The OMI NO₂ products use an implicit aerosol correction to account for
142 the optical effects of aerosols, but retrievals can be biased when aerosol loading is extreme (Castellanos et al.,
143 2015). Under these conditions, the OMI NO₂ retrieval is biased low by roughly 20 to 40% (Chimot et al., 2016).
144 Note that any aerosol-related error would have the potential effect of underestimating the magnitude of decreases
145 in NO₂ column densities when comparing 2020 to previous years. Additional bias in the NO₂ product may be
146 introduced due to the reliance on nearly cloud-free pixels, in which greater sunlight may induce higher
147 photochemical rates. For example, the current NO₂ product is biased roughly 30% low over the Canadian oil sands
148 (McLinden et al., 2014). The level-2 OMI- NO₂ product has been validated against in situ and surface-based
149 observations showing good agreement (Lamsal et al., 2014). The use of fixed Jacobians in the SO₂ product
150 introduces systematic errors of 50 to 100% for cloud-free observations (Krotkov et al., 2016).

151

152 Starting in 2007, the quality of level 1B radiance data for some OMI viewing directions has been affected, known
153 as the row anomaly. The L3 products used here exclude all pixels affected by the row anomaly from each
154 observation, but the locations of the row anomaly pixels were dynamic between 2007 and 2011, which could
155 affect any comparisons including those years. Since 2011, the pixels affected by the row anomaly problem are the
156 same, so comparisons for data only since 2011 are not affected by changes in the row anomaly.

157

158 Moderate Resolution Imaging Spectroradiometer (MODIS) sensors observe the Earth from polar orbit, from Terra
159 satellite since 2000 and from Aqua since mid 2002. In this study we use MODIS-derived AOD at 550nm obtained
160 by merging Dark Target and Deep Blue retrievals (Sayer et al., 2014). Specifically, we use the
161 Deep_Blue_Aerosol_Optical_Depth_550_Land_Mean field over land and the over ocean
162 AOD_550_Dark_Target_Deep_Blue_Combined_Mean the from Collection 6.1 L3 Gridded products MYD08 and
163 MOD08 (Hubanks et al., 2019), though very few retrievals over ocean are included in our analysis. L3 values are
164 computed on $1^\circ \times 1^\circ$ spatial grid from L2 AOD products with resolution of 10x10 km. Over land 66% of MODIS-
165 retrieved Dark-target AOD values were shown to be $\pm 0.05 \pm 0.15 * \text{AOD}$ AERONET-observed values, with high
166 correlation ($R = 0.9$) (Levy et al., 2010). Around 78% of the Deep Blue retrievals are within the expected error
167 range of $\pm 0.05 \pm 0.20 * \text{AOD}$ (Sayer et al., 2013). MODIS AOD data have been extensively used by the modeling
168 and remote sensing scientific communities and inter-compared with a wide range of satellite AOD products (see
169 Schutgens et al. (2020) and references therein).

170

171 We analyzed these retrievals over two large regions (Fig. 1). Central east China was comprised of Shaanxi, Hubei,
172 Anhui, Jiangsu, Shanxi, Henan, Hebei, Shandong, Beijing, and Tianjin provinces. Southern China was comprised
173 of Guizhou, Guangxi, Hunan, Jiangxi, Guangdong, Fujian and Zhejiang provinces. Daily mean quantities were
174 calculated across all valid retrievals falling within the provinces comprising the regions. For the OMI NO₂
175 columns, individual retrievals were weighted by the L3 ‘Weight’ field, which is proportional to the fraction of the
176 grid cell with higher-quality retrievals, identified as those have less than 30% cloud fraction and not affected by
177 the row anomaly problem. We also calculated the daily value from the median of all retrievals, to understand
178 whether individual high values (mainly SO₂) had any effect on the significance of trends or differences between
179 2020 and different background periods. Monthly averages were calculated from the daily regional averages, with
180 each day weighted in the monthly average by the number of valid retrievals so as to not overrepresent days with
181 little satellite coverage or significant cloud cover. The monthly data were used to visually identify COVID-19
182 related changes against background seasonality and trends since 2005.

183

184 We examined the difference in the distribution of daily data during the 2020 January 23 to April 8 lockdown
185 period to the same period during previous years since 2005. We compared 2020 to 2019, to different background
186 periods, and to the expected value for 2020 estimated from trends over different background periods. Given the
187 uneven nature of changes in atmospheric composition over different parts of China identified in previous studies,
188 background periods were defined for each possible starting year between 2005 and 2018 with each ending in
189 2019. Retrieved quantities in 2020 were compared to the background means over each period and to the value
190 expected for 2020 estimated from the linear trend over each period. We tested the significance of these differences
191 using bootstrap resampling (Efron and Gong, 1983) with a resampling size of 2000.

192

193 We also considered how the analysis depended on how the lockdown period was defined. Emissions and pollution
194 can decrease during the Chinese New Year holidays (Chen et al., 2020), which started as early as January 23 in
195 2012 and as late as February 19 in 2015, complicating COVID-19 related analyses of atmospheric composition
196 over China (Bauwens et al., 2020; Chen et al., 2020). The timing and extent of lockdowns also varied between
197 provinces and we assume that ‘slowdowns’ could have happened before or after stricter, official lockdowns. For

198 example, ground and air transportation remaining below lockdown levels nationally at least through April 14,
199 2020 (International Energy Agency, 2020). Excluding the holiday period from all years is a straightforward
200 approach to excluding any New Year holiday effects but will exclude simultaneous lockdown effects during the
201 initial, and presumably most strict, stages of the lockdown. Rather than specifying different combinations of New
202 Year holiday period and provincial-level lockdown timing, we used January 23-April 8 as our baseline period
203 (which will include all holiday periods since 2005), but examined the sensitivity of the statistics to the length of
204 the lockdown period, namely a longer lockdown period beginning one week earlier and one week later, and a
205 shorter lockdown period for February only. In interpreting the data, we put more confidence in 2020 differences
206 that were insensitive to these choices.

207 **3 Results**

208 **3.1 Regional patterns and seasonality**

209 Figure 2 shows the 2020 –2019 differences over China during the January 23-April 8 lockdown period for the
210 four satellite-retrieved quantities. There were decreases of 5-10 ppbv in AIRS CO over central east China (Fig.
211 2a) and increases of 20-25 ppbv over southern China in 2020 compared to 2019. The increase in southern China
212 is adjacent to a stronger positive CO anomaly over the upper Mekong regions of Myanmar, Thailand and Laos.
213 There were no coherent regional changes in OMI SO₂ (Fig. 2b), but rather smaller localized difference of either
214 sign. There were decreases in NO₂ (Fig. 2c) across central east China exceeding 8×10^{15} molec cm⁻² coincident
215 with the weaker decrease in CO. Over southern China, there were comparable differences over Guangdong
216 province, with smaller differences elsewhere. There was a decrease in MODIS AOD (Fig. 2d) in central-east
217 China coincident with the decreases in CO and NO₂, but smaller in magnitude. There was a region of higher AOD
218 in and northeast of the upper Mekong region coincident with the CO increase, both presumably because of biomass
219 burning.

220
221 To put the 2020/2019 difference maps in a longer-term and seasonal context, Figure 3 shows monthly averages
222 of the four retrieved quantities over central east China since 2005. There are seasonal CO peaks in March-April,
223 June and September, with the minima usually in November and December (Figure 3a). There has been a decrease
224 since 2005 in CO. The seasonal decrease from January to February in 2020 is similar to that which has occurred
225 occasionally before, but the CO during February and March 2020 was the lowest for that time of the year since
226 2005. By April, CO had returned to levels typical of 2015-2019. The main characteristics of the monthly SO₂ over
227 the region are that it has decreased since 2005 (Figure 3b), and that early 2020 SO₂ was within the range of recent
228 levels. There is a strong seasonal NO₂ cycle (Figure 3c), with a July-August minimum, and December-January
229 peak, which has been attributed to increased heating needs (Yu et al., 2017; Si et al., 2019) and longer chemical
230 lifetime owing to lower OH and RO₂ (Shah et al., 2020). NO₂ has also decreased since 2011, and during most
231 years, there is a departure from a smooth seasonal cycle in January and February associated with the Chinese New
232 Year holiday period. January and February 2020 NO₂ was considerably lower than previous years, increased
233 during March, and had recovered to typical, recent levels by April. AOD has consistent seasonal peaks in summer
234 which have been attributed to hygroscopic growth and agricultural residue burning (Filonchik et al., 2019), but
235 had less regular seasonality otherwise, and has decreased since 2011. AOD during February and particularly

236 March of 2020 were lower than recent years, but during which time there was considerable variability in the
237 monthly data.

238

239 Figure 4 shows the four retrieved quantities over southern China. There is a springtime maximum in CO (Fig. 4a),
240 a less regular maximum during September-January, and an annual minimum in July. The range of CO is similar
241 to central east China. CO over the last 5 years is lower than earlier in the record, and early 2020 CO was higher
242 than recent years. SO₂ (Fig. 4b) is lower than central east China and any seasonal cycle is also hard to identify.
243 The high June 2011 values are due to the Nabro eruption in Ethiopia (Fromm et al., 2014) which is still apparent
244 in the time series despite excluding individual SO₂ retrievals that are greater than 15 DU, and are due to a
245 combination of higher overall background values and individual retrievals with very high (> 10 DU) SO₂. NO₂
246 (Fig. 4c) is lower than over central east China, but both regions share a similar seasonality. NO₂ during January-
247 April 2020 was slightly lower than in 2019. AOD (Fig. 4d) has weak seasonal peaks in October, March and June,
248 has decreased since 2011, and 2020 fell within the range of 2015-2019.

249

250 **3.2 Central east China**

251 Figure 5 shows the CO, SO₂, NO₂ and AOD for January 23 – April 8 of each year over central east China as box
252 and whisker plots with the median, interquartile range and 2.5th and 97.5th percentiles over all daily mean data as
253 horizontal lines and the mean shown by the black dot. The associated statistics comparing 2020 and 2019 are
254 provided in Table 1, and comparing 2020 to longer background periods with and without trends accounted for as
255 supplementary Tables S1-S4. The AIRS CO is shown in Figure 5a. The variation during January 23 – April 8 of
256 each year is due to weather-related factors and observational error. The mean CO of 133.5 ppbv in 2020 was 3.2%
257 less than the 2019 mean of 137.9 ppbv, which was only marginally significant, having a 95% confidence interval
258 (-6.3% - 0.1%) close to spanning 0. During years prior, there were increases and decreases in CO from year to
259 year, but an overall decreasing trend since 2005. To quantify if the 2020 departure was significant against this
260 background, we compared the distribution of observed 2020 CO to the background average and to that which
261 might be expected given any trends over the background period. Because there was no obvious starting year for
262 the background period, we considered different periods starting in each year between 2005 and 2018 and ending
263 in 2019 (Fig. 6a, Table S1). The difference between 2020 and the background depended strongly on the starting
264 year of the background period, ranging from -11.5% lower than the 2005-2019 mean to -3.1% lower than over
265 2018-2019, but all were statistically significant. Significant trends over years beginning between 2005 and 2016
266 (shown in Fig. 6a by the red line and shading) ranged between -1.5 ppbv yr⁻¹ when starting in 2013 to -3.6 ppbv
267 yr⁻¹ if starting in 2016. The uncertainty in the trends increased for trends over shorter periods, and were,
268 unsurprisingly, insignificant by 2017, with the 95% confidence intervals of the trends spanning 0. The differences
269 between the observed 2020 mean and the value predicted from the trend (magenta line) varied inversely with the
270 trend and was always negative, but, except for 2009, had 95% confidence intervals (magenta shading) spanning
271 0, and therefore were not considered significant. For CO therefore, 2020 was significantly lower than the
272 background period mean but not consistently lower than predicted given the decreasing trend during the
273 background period, no matter how this period was defined. Results were similar for CO analysed closer to the
274 surface at 850 hPa (not shown), but where the retrieval has less sensitivity.

275

276 OMI SO₂ (Fig. 5b) fluctuated over 2005 to 2011 and declined steadily afterward during which variation also
277 declined, becoming narrower to a degree not seen in the CO. The 2020 mean of 0.057 was 95% higher than the
278 2019 mean of 0.031, but with a wide 95% confidence interval (15% - 250%). For different background periods
279 (Table S2), 2020 SO₂ ranged from 83% less than the 2005-2019 mean to 30% less than the 2016-2019 mean, with
280 insignificant differences compared to more recent periods. Trends varied significantly from to -0.03 yr⁻¹ over
281 2005-2019 to -0.06 DU yr⁻¹ over 2012-2019 (Fig. 6b), during which the trend could explain a maximum of 32%
282 of the variation in the data. For periods starting in 2007 and after, the observed 2020 mean was significantly higher
283 than predicted. Relative to the value predicted from the 2012-2019 trend of -0.06, the observed 2020 SO₂ was
284 200% higher; the large percent difference reflects a predicted value close to zero, and we note that the retrieved
285 SO₂ can be negative for individual values and averages (Li et al., 2013; Wang and Wang 2020). The observed
286 2020 SO₂ was much higher than expected from trends calculated over 2016-2019 when SO₂ was flat and with less
287 variability, but the low SO₂ approaching the detection limit over this period make these estimates not particularly
288 meaningful. Furthermore, the change in 2020 SO₂ was strongly dependent on whether daily values were calculated
289 from the mean or median of individual values over the region. For most background periods (Figure S1b), the
290 trends in the median values were still negative until 2015, but 2020 was only 8.4% higher than predicted from the
291 2012-2019 trend and not significantly different from expected for trends beginning later. This likely reflects the
292 greater influence of high individual retrieval values on the daily mean value compared to the median, even after
293 the basic filtering of transient SO₂ plumes.

294

295 OMI NO₂ (Fig. 5c) increased from 2005 to 2011 and decreased thereafter with an apparent flattening since 2016.
296 The 2020 mean NO₂ of 6.5×10^{15} molec cm⁻² was 32% less than the 2019 mean of 9.6×10^{15} molec cm⁻²; the
297 pronounced regional difference between 2020 and 2019 (Fig. 2c, Fig. 5c) in part reflects a 2019 uptick from 2018.
298 For different background periods (Table S3), 2020 NO₂ ranged from 43.3% less than the 2010-2019 mean to 30%
299 less than the 2018-2019 mean, with all differences significant. Trends were negative and significant for starting
300 years between 2007 and 2015 (Fig. 6c) with the strongest trend of -0.75×10^{15} molec cm⁻² yr⁻¹ for the period
301 beginning in 2011. 2020 NO₂ was significantly less than the predicted value for all background periods but varied
302 from 16.8% less than predicted from the 2011-2019 trend to 27.1% less than predicted from the 2015-2019 trend,
303 the last period when there was a significant, although weak, decrease.

304

305 MODIS AOD (Fig. 5d) was flat or slightly increasing from 2005 to 2011, decreasing thereafter and with a
306 flattening since 2016 similar to SO₂ and NO₂. The 2020 mean AOD of 0.41 was 14% less than the 2019 mean of
307 0.48, but this was not significant. For different background periods (Table S4), 2020 AOD ranged from 30.2%
308 less than the 2007-2019 mean to 14.2% less than the 2018-2019 mean, with confidence intervals for the
309 differences becoming closer to spanning 0 for more recent periods. Trends were negative and significant for
310 starting years between 2005 and 2014 (Fig. 6d), with the strongest decrease of 0.04 yr⁻¹ over the 2012-2019 period.
311 There was no significant difference between the observed and predicted 2020 mean for periods beginning in 2008
312 and later, when the trends were strongest, and which approached 0 after 2014.

313 3.3 Southern China

314 Figure 7 shows the distribution of daily CO, SO₂, NO₂ and AOD for January 23-April 8 of each year over southern
315 China. The associated statistics comparing 2020 and 2019 are provided in Table 2. AIRS CO (Fig. 7a) in 2020
316 was 144.7 ppbv, 13% higher than the 2019 mean of 128.5 ppbv which can be seen in an upward shift in the
317 distribution of the box plot. 2020 CO was between 4.4% and 8.8% greater than the background mean for periods
318 starting after 2014 (Table S5), but not significantly different otherwise. CO decreased significantly for periods
319 starting between 2005 and 2016 (Fig. 8a). When these trends are taken into account, 2020 CO was between 11.2%
320 and 18.7% greater than predicted, and in all cases were significant.

321
322 OMI SO₂ (Fig. 7b) fluctuated from 2005 until 2013 and flattened afterwards, driven by fewer high individual SO₂
323 values in later years, as in central east China. The 2020 mean of 0.003 DU was 116% higher than the 2019 mean
324 of -0.02 DU but also with a wide 95% confidence interval (24% - 223%). 2020 was less than the background
325 mean periods starting between 2005 and 2011 (Table S6), but not significantly different otherwise. SO₂ trends
326 were consistently negative for all periods (Fig. 8b), although not as strong as over central east China. Whether
327 2020 SO₂ was greater than predicted from trends depended more on the background period than over central east
328 China and were also not significantly different from predicted when daily values were calculated from the median
329 SO₂ of individual retrievals for any background period (Figure S2b).

330
331 OMI NO₂ (Fig. 7c) increased toward 2011 and 2012, declining after to 2005-2010 levels. The 2020 mean of
332 3.3×10^{15} molec cm⁻² was 22% less than the 2019 mean of 4.3×10^{15} molec cm⁻². For longer background periods,
333 2020 was between 22.9% and 30.6% less than the mean (Table S7), all of which were significant. NO₂ trends
334 were significantly negative when the start of the trend was calculated using years between 2007 and 2012, but not
335 otherwise (Fig. 8c). The 2020 NO₂ mean was significantly lower than predicted, except for when the trend was
336 estimated beginning in 2011 or 2018. A two-year trend cannot be interpreted meaningfully, especially without
337 considering meteorological differences. Visually, however, it is hard to tell if the 2020 NO₂ distribution represents
338 a COVID-related departure or a decrease comparable to changes during recent previous years, unlike over central
339 east China.

340
341 MODIS AOD (Fig. 7d) was comparable to NO₂ in its increase toward 2012, decrease thereafter and flattening
342 during more recent years. The 2020 mean AOD of 0.38 was 12% higher than the 2019 mean of 0.34, but with a
343 95% confidence interval (-7% - 34%) spanning 0. Similarly, 2020 was between 14% and 22% lower than during
344 background periods beginning from 2005 to 2012, but not for more recent periods (Table S8). The AOD trends
345 were significantly negative for all start years until 2015. The 2020 mean was between 32 and 47% higher than
346 predicted from trends for periods starting between 2010 and 2015, but was not different from predicted for trends
347 starting in other years.

348
349 For both regions and all quantities, the differences between observed and predicted values for 2020 were
350 insensitive to a longer lockdown period, or to whether the bootstrap resampling was weighted by the number of
351 valid retrievals each day. For a February-only lockdown period (Figures S3 and S4), the CO trends were more
352 significant when starting in later years, but the differences between the observed and expected values remained

353 insignificant over central east China. The SO₂ trends for different periods were similar. The 2020 SO₂ differences
354 from what would be expected approached 0 for later periods but were also not consistently different when the
355 median values of individual retrievals were used. Results for NO₂ were unaffected. The AOD 2020 difference
356 from what would be expected was stronger and technically significant, but still with a very wide confidence
357 interval and therefore difficult to interpret. We emphasize that while a February-only lockdown period is useful
358 for comparison, it is problematic in not including the New Year's holiday periods from all previous years.

359 **4 Discussion and conclusions**

360 The degree to which the COVID-19 lockdowns in China resulted in changes in atmospheric composition depended
361 strongly on the background period and whether existing trends were taken into account. For AIRS CO over central
362 east China, the 2020 mean was 3-12% lower compared to different background periods. Relative to mean CO
363 concentrations during periods beginning between 2005 and 2016, there were significant decreases in CO but CO
364 in 2020 was not consistently different from what would be expected from trends calculated over this period. These
365 longer-term declines in CO concentrations do appear to flatten out in recent years; assuming that the flat CO
366 during 2017-2019 would have persisted, we estimate a 3-4% reduction in CO in 2020 relative to that period. For
367 MODIS AOD, the 2020 mean was between 14% and 30% less than different background averages, but not
368 significantly different from what would be expected for trends beginning between 2008 and 2014. As with CO
369 trends, the negative AOD trends in the region also appear to flatten in recent years. Relative to the flat AOD over
370 2016-2019, 2020 AOD was 14-17% lower than the background mean; as with CO, this range would be the more
371 meaningful estimate of changes in 2020 if we assume that this flattening were to persist. 2020 SO₂ was
372 significantly lower than background averages calculated over most periods, ranging from 83% less than over
373 2005-2019 to 30% less than over 2016-2019. Compared to the 2012-2019 period when there were no significant
374 SO₂ increases, 2020 SO₂ was 200% greater than what would be expected based on a trend starting in 2021, but
375 only 8% greater when the median of daily retrievals was used, and not significantly different from expected
376 relative to the trends beginning later than 2012. SO₂ concentrations were relatively flat from 2016-2019; when
377 using 2016 as the first year of the trend, SO₂ was significantly higher than the expected value when calculated
378 from the mean of the daily SO₂ retrievals, but not significantly different when calculated from the median. We
379 note also that analyses of SO₂ and NO₂ that include years prior to 2012 may be affected by changes in observation
380 sample size due to changes in the OMI row anomaly.

381
382 OMI NO₂ in 2020 over central east China was consistently lower than the background average and expected value
383 from the trends. There was a 17% decrease in 2020 relative to the value expected from a trend calculated over
384 2011-2019, but a 30-33% decrease relative to the different background means since 2016 when the NO₂ was
385 relatively flat. Again assuming that this flattening were to persist, this latter range may be the more meaningful
386 baseline for the 2020 decrease. For reference, Bauwens et al. (2020) reported a ~40% drop in OMI NO₂ from
387 2019 to 2020 over cities affected by the lockdown using the QA4ECV retrieval (Boersma et al., 2018), and a
388 ~51% drop in NO₂ over the eight cities (Beijing, Jinan, Nanjing, Qingdao, Tianjin, Wuhan, Xi'an and Zhengzhou)
389 falling within our central east China region. Our analysis cannot be compared directly because we include non-
390 urban areas and define the lockdown period differently, but we do note that NO₂ during the same period in 2019

391 appeared to be anomalously high relative to the previous few years, which would make the decreases in 2020
392 appear more significant.

393

394 The modest decreases in CO and AOD over central east China were unexpected; given its high population density
395 and level of industrial activity, lockdowns may have been anticipated to lead to larger decreases. In the case of
396 MODIS AOD, these modest decreases were possibly due to contributions from other sources unaffected by
397 COVID-19 related lockdowns, limitations in the MODIS AOD retrieval under cloudy conditions, climatological
398 variability from other sources such as mineral dust, and meteorology favourable to secondary aerosol formation
399 which could have offset lower emissions (Wang et al., 2020). The 2020 increase in SO₂ is more difficult to
400 interpret because of the discrepancies between daily values calculated from the mean or median of individual
401 retrievals, but is broadly consistent with surface observations that find no significant change in in-situ surface SO₂
402 over Wuhan in the daily mean, and a slight increase in daytime SO₂ possibly associated with increased residential
403 heating and cooking (Shi and Brasseur, 2020).

404

405 Over southern China retrieved 2020 SO₂ was significantly lower than the background average only for periods
406 beginning between 2005 to 2011. Significant departures from expected trends were uneven when using the mean
407 value of daily retrievals and absent when using the median value. As with central East China, we conclude that
408 no significant changes could be robustly detected in 2020 SO₂. NO₂ in 2020 was between 23% and 32% less than
409 the background average for different periods. Here, the flattening in NO₂ beginning in 2013 is easier to identify
410 than over central east China because of the much higher NO₂ during the three years prior; 2020 NO₂ was 23-27%
411 less than different background means between 2013 and 2019. The more significant reductions in NO₂ in central
412 east China compared to the south is presumably due the former's greater population and industrialization, and
413 consequently higher pollution levels. This is consistent with Chen et al.'s (2020) detection of a larger 2020
414 decrease in surface NO₂ in Wuhan compared to Shanghai. Retrieved CO in 2020 was between 4 and 8% greater
415 than background averages beginning in 2014, but between 11% and 19% higher than what would be expected
416 given the decreasing trends over any period. AOD in 2020 was lower than background averages calculated starting
417 with years earlier than 2012, but higher or not significantly different from expected for trends calculated starting
418 in years after 2012.

419

420 The focus of this analysis is on whether satellite retrievals of atmospheric composition over 2020 departed
421 significantly from different background periods and expected values for 2020 when daily variability and trends
422 are accounted for, but it is useful at a preliminary stage to speculate as to how different emissions changes could
423 have contributed to 1) why NO₂ was robustly lower in 2020 over central east China compared to CO and AOD,
424 and 2) why CO and perhaps AOD were higher over southern China compared to what would be expected from
425 recent trends.

426

427 To understand why NO₂ differences over central east China were more significant than other quantities, Table 3
428 shows the emissions by sector for a representative set of constituents from the Community Emissions Data System
429 (CEDS) (Hoesly et al., 2018) over China for 2014, the most recent year available. Other bottom-up emissions
430 inventories will vary in absolute emissions amounts and their sector contributions, particularly for more recent

431 periods, but CEDS is the standard available emissions dataset available globally as a baseline for the next IPCC
432 assessment, in anticipation of assessing 2020 COVID-19 related changes to atmospheric composition in other
433 regions, and for modeling studies involving a transboundary transport component. Across all species, energy
434 production, industrial activity, transportation, residential/commercial/other (RCO), and waste disposal constitute
435 the bulk of the emissions. Based on activity data for the first quarter of 2020, energy demand across China declined
436 by 7% compared to 2019, and transportation sector activity declined by 50 to 75% in regions with lockdowns in
437 place (International Energy Agency, 2020). These sectors are direct or indirect sources of numerous pollutants,
438 including SO₂ (the precursor of sulfate aerosol), NO_x, CO, and primary anthropogenic aerosols classified broadly
439 as organic carbon (OC) and black carbon (BC). If we apply the 7% reduction in energy production and mid-point
440 62.5% reduction to transportation from the IEA, assume a 20% reduction in industrial emissions, 5% reduction in
441 waste emissions, no change in RCO (with commercial decreases offset by residential increases), this yields a 10%
442 reduction in BC, 5% reduction in OC, 14% reduction in SO₂, 14% reduction in CO and 21% reduction in NO₂.
443 The larger reduction in NO₂ relative to other emissions could partly explain why OMI NO₂ column density
444 changes over central east China were stronger than in the other retrievals.

445

446 Following Si et al.'s (2019) consideration of biomass burning as a pollution source in China alongside
447 anthropogenic sources, we considered transboundary smoke transport as a possible reason for the higher 2020 CO
448 over southern China, guided by higher CO over the Upper Mekong region in 2020 compared to 2019 (Fig. 2a)
449 and the predominant westerly flow during this time of year (Reid et al., 2013). Table 4 compares January 23-April
450 8 AIRS CO over southern China to CO emissions estimates from biomass burning from the Global Fire
451 Assimilation System (GFAS) (Kaiser et al., 2012) over the upper Mekong region (17° N to 25° N, 95° E to 105°
452 E) including parts of eastern Myanmar, northern Thailand, and northern Laos. From 2005 to 2020, variation in
453 GFAS CO over this region explained a moderate (32%) amount of variability in AIRS CO over southern China,
454 suggesting it as a non-negligible contributor to variation in CO concentration, and a contributor to higher CO
455 in 2020. This illustrates that, at a minimum, sources such as biomass burning smoke and dust that are less affected
456 by COVID-19 related measures will complicate attribution studies. To that end, modeling studies following Wang
457 et al. (2020) will be required to isolate emissions, meteorological and chemical drivers of changes in atmospheric
458 composition and their effects at a process level. With proper instrument-equivalent comparisons, modelling
459 studies will also help to identify the extent to which the lack of significant changes are due to retrieval limitations,
460 namely low sensitivity near the surface where differences would presumably be more pronounced, particularly
461 given remote emissions sources such as dust, biomass burning smoke and volcanic SO₂, which will arrive at higher
462 altitudes.

463

464 The key implication of our study is that interpreting differences in 2020 retrievals of atmospheric composition
465 depends strongly on how the background period is defined and whether trends over these periods are accounted
466 for. Not taking these into account could lead to misattribution of changes in air quality to COVID-19 lockdowns,
467 or, at a minimum, that whether differences in 2020 are significant depend on the choice of background period,
468 which is somewhat subjective. Leading up to 2020, there was an apparent flattening of decreasing trends beginning
469 earlier in the decade across the retrievals; the considerable variability in the data made identifying this flattening
470 easier in some cases than in others. We are more confident, for example, in our estimate of a 23-27% decrease in

471 2020 NO₂ over southern China relative to a flat background period than the 30-33% decrease over central East
472 China, where the recent variability was greater and the flattening less apparent. Revisiting this type of analysis in
473 the years when regional economies have fully recovered post COVID-19 will help to distinguish between further
474 decreases and flat trends and will lend themselves to using non-linear models in estimating the trends. We have
475 approached the issue by comparing data for 2020 to what would have been expected given recent trends and by
476 applying a single lockdown period to two large regions, with additional analyses to gauge the sensitivity of the
477 2020 differences to these choices. Other studies over China or elsewhere will inevitably use other approaches that
478 more explicitly account for seasonality, meteorology, and which relate changes in pollution over smaller areas
479 (e.g. single provinces or states) to region-specific lockdown measures and timing at a process level. Regardless
480 of the approach, however, it is important to consider recent trends and variability. In places where pollution has
481 decreased, not accounting for recent context could result in over-attribution of changes in pollution to COVID-
482 19. In places where pollution has increased, such as parts of South Asia, this could result in under-attribution.

483 **Code/data availability:** The MATLAB code used to process the satellite data is provided as Supplementary
484 Material. All source data are publicly available.

485 **Competing interests:** The authors have no competing interests.

486

487 **Author contribution:** All authors conceived of the study. RF, IG and KT conducted the data analysis. RF and
488 JH prepared the manuscript with contributions from all co-authors.

489

490 **Acknowledgements:** The authors thank two anonymous reviewers and the editor for their constructive feedback
491 which improved the interpretation of the data and conclusions drawn in the paper.

492

493

494 **References**

- 495 Bauwens, M., Compornolle, S., Stavrakou, T., Müller, J., van Gent, J., Eskes, H., Levelt, P. F., van der A., R.,
496 Veeffkind, J. P., Vlietinck, J., Yu, H., and Zehner, C.: Impact of coronavirus outbreak on NO₂ pollution assessed using
497 TROPOMI and OMI observations, *Geophysical Research Letters*, 2, 0-3, 10.1029/2020GL087978, 2020.
- 498 Boersma, K. F., Eskes, H. J., Richter, A., De Smedt, I., Lorente, A., Beirle, S., van Geffen, J., Zara, M., Peters, E.,
499 Van Roozendaal, M., Wagner, T., Maasakkers, J. D., van der A., R. J., Nightingale, J., De Rudder, A., Irie, H., Pinardi,
500 G., Lambert, J. C., and Compornolle, S. C.: Improving algorithms and uncertainty estimates for satellite NO₂
501 retrievals: results from the quality assurance for the essential climate variables (QA4ECV) project, *Atmospheric*
502 *Measurement Techniques*, 11, 6651-6678, 10.5194/amt-11-6651-2018, 2018.
- 503 Castellanos, P., Boersma, K. F., Torres, O., and de Haan, J. F.: OMI tropospheric NO₂ air mass factors over South
504 America: effects of biomass burning aerosols, *Atmospheric Measurement Techniques*, 8, 3831-3849, 10.5194/amt-8-
505 3831-2015, 2015.
- 506 Chen, K., Wang, M., Huang, C., Kinney, P. L., and Paul, A. T.: Air Pollution Reduction and Mortality Benefit during
507 the COVID-19 Outbreak in China, *Lancet Planetary Health*, 0-3, 10.1101/2020.03.23.20039842, 2020.
- 508 Chimot, J., Vlemmix, T., Veeffkind, J. P., de Haan, J. F., and Levelt, P. F.: Impact of aerosols on the OMI tropospheric
509 NO₂ retrievals over industrialized regions: how accurate is the aerosol correction of cloud-free scenes via a simple
510 cloud model?, *Atmospheric Measurement Techniques*, 9, 359-382, 10.5194/amt-9-359-2016, 2016.
- 511 Efron, B., and Gong, G.: A Leisurely Look at the Bootstrap, the Jackknife, and Cross-Validation, *American Statistician*,
512 37, 36-48, 10.2307/2685844, 1983.
- 513 Filonchyk, M., Yan, H. W., and Zhang, Z. R.: Analysis of spatial and temporal variability of aerosol optical depth
514 over China using MODIS combined Dark Target and Deep Blue product, *Theoretical and Applied Climatology*, 137,
515 2271-2288, 10.1007/s00704-018-2737-5, 2019.
- 516 Fioletov, V. E., McLinden, C. A., Krotkov, N., Li, C., Joiner, J., Theys, N., Carn, S., and Moran, M. D.: A global
517 catalogue of large SO₂ sources and emissions derived from the Ozone Monitoring Instrument, *Atmospheric Chemistry*
518 *and Physics*, 16, 11497-11519, 10.5194/acp-16-11497-2016, 2016.
- 519 Fromm, M., Kablick, G., Nedoluha, G., Carboni, E., Grainger, R., Campbell, J., and Lewis, J.: Correcting the record
520 of volcanic stratospheric aerosol impact: Nabro and Sarychev Peak, *Journal of Geophysical Research-Atmospheres*,
521 119, 10.1002/2014jd021507, 2014.
- 522 Geddes, J. A., Martin, R. V., Boys, B. L., and van Donkelaar, A.: Long-Term Trends Worldwide in Ambient NO₂
523 Concentrations Inferred from Satellite Observations, *Environmental Health Perspectives*, 124, 281-289,
524 10.1289/ehp.1409567, 2016.
- 525 Georgoulias, A. K., van der A., R. J., Stammes, P., Boersma, K. F., & Eskes, H. J.: Trends and trend reversal detection
526 in two decades of tropospheric NO₂ satellite observations, *Atmospheric Chemistry and Physics*, 19, 6269–6294,
527 <https://doi.org/10.5194/acp-2018-988>, 2019.
- 528 Han, H., Liu, J., Yuan, H. L., Jiang, F., Zhu, Y., Wu, Y., Wang, T. J., and Zhuang, B. L.: Impacts of Synoptic Weather
529 Patterns and their Persistency on Free Tropospheric Carbon Monoxide Concentrations and Outflow in Eastern China,
530 *Journal of Geophysical Research-Atmospheres*, 123, 7024-7046, 10.1029/2017jd028172, 2018.

- 531 He, Q. Q., Gu, Y. F., and Zhang, M.: Spatiotemporal patterns of aerosol optical depth throughout China from 2003 to
532 2016, *Science of the Total Environment*, 653, 23-35, 10.1016/j.scitotenv.2018.10.307, 2019.
- 533 Hoesly, R. M., Smith, S. J., Feng, L. Y., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L.,
534 Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z. F., Moura,
535 M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750-2014) anthropogenic emissions of reactive gases and
536 aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369-408,
537 10.5194/gmd-11-369-2018, 2018.
- 538 Hubanks, P., Platnick, S., King, M., and Ridgway, B.: MODIS Algorithm Theoretical Basis Document No. ATBD-
539 MOD-30 for Level-3 Global Gridded Atmosphere Products (08_D3, 08_E3, 08_M3) and Users Guide (Collection 6.0
540 & 6.1, Version 4.4, 20 Feb 2019), NASA Goddard Space Flight Center, Greenbelt, MD, 2019.
- 541 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz,
542 M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system
543 based on observed fire radiative power, *Biogeosciences*, 9, 527-554, 10.5194/bg-9-527-2012, 2012.
- 544 Krotkov, N. A.: OMI/Aura NO₂ Cloud-Screened Total and Tropospheric Column L3 Global Gridded 0.25 degree x
545 0.25 degree V3, NASA Goddard Space Flight Center, 2013.
- 546 Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela,
547 E. J., Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He,
548 H., Lu, Z., and Streets, D. G.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015,
549 *Atmospheric Chemistry and Physics*, 16, 4605-4629, 10.5194/acp-16-4605-2016, 2016.
- 550 Krotkov, N. A., Lamsal, L. N., Celarier, E. A., Swartz, W. H., Marchenko, S. V., Bucsela, E. J., Chan, K. L., Wenig,
551 M., and Zara, M.: The version 3 OMI NO₂ standard product, *Atmospheric Measurement Techniques*, 10, 3133-3149,
552 10.5194/amt-10-3133-2017, 2017.
- 553 Lamsal, L. N., Krotkov, N. A., Celarier, E. A., Swartz, W. H., Pickering, K. E., Bucsela, E. J., Gleason, J. F., Martin,
554 R. V., Philip, S., Irie, H., Cede, A., Herman, J., Weinheimer, A., Szykman, J. J., and Knepp, T. N.: Evaluation of OMI
555 operational standard NO₂ column retrievals using in situ and surface-based NO₂ observations, *Atmospheric
556 Chemistry and Physics*, 14, 11587-11609, 10.5194/acp-14-11587-2014, 2014.
- 557 Levy, R. C., Remer, L. A., Kleidman, R. G., Mattoo, S., Ichoku, C., Kahn, R., and Eck, T. F.: Global evaluation of
558 the Collection 5 MODIS dark-target aerosol products over land, *Atmospheric Chemistry and Physics*, 10, 10399-
559 10420, 10.5194/acp-10-10399-2010, 2010.
- 560 Li, C., Joiner, J., Krotkov, N. A., and Bhartia, P. K.: A fast and sensitive new satellite SO₂ retrieval algorithm based
561 on principal component analysis: Application to the ozone monitoring instrument, *Geophysical Research Letters*, 40,
562 6314-6318, 10.1002/2013gl058134, 2013.
- 563 Li, M., Zhang, Q., Kurokawa, J., Woo, J. H., He, K. B., Lu, Z. F., Ohara, T., Song, Y., Streets, D. G., Carmichael, G.
564 R., Cheng, Y. F., Hong, C. P., Huo, H., Jiang, X. J., Kang, S. C., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian
565 anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP,
566 *Atmospheric Chemistry and Physics*, 17, 935-963, 10.5194/acp-17-935-2017, 2017.
- 567 Lin, C. Q., Liu, G., Lau, A. K. H., Li, Y., Li, C. C., Fung, J. C. H., and Lao, X. Q.: High-resolution satellite remote
568 sensing of provincial PM 2.5 trends in China from 2001 to 2015. *Atmospheric Environment*, 180, 110-116,
569 <https://doi.org/10.1016/j.atmosenv.2018.02.045>, 2018.
- 570 Lin, N., Wang, Y. X., Zhang, Y., and Yang, K.: A large decline of tropospheric NO₂ in China observed from space
571 by SNPP OMPS, *Science of the Total Environment*, 675, 337-342, 10.1016/j.scitotenv.2019.04.090, 2019.

- 572 Luan, Y., and Jaegle, L.: Composite study of aerosol export events from East Asia and North America, *Atmospheric*
573 *Chemistry and Physics*, 13, 1221-1242, 10.5194/acp-13-1221-2013, 2013.
- 574 Ma, Z. W., Hu, X. F., Sayer, A. M., Levy, R., Zhang, Q., Xue, Y. G., Tong, S. L., Bi, J., Huang, L., and Liu, Y.:
575 *Satellite-Based Spatiotemporal Trends in PM_{2.5} Concentrations: China, 2004-2013*, *Environmental Health*
576 *Perspectives*, 124, 184-192, 10.1289/ehp.1409481, 2016.
- 577 McLinden, C. A., Fioletov, V., Boersma, K. F., Kharol, S. K., Krotkov, N., Lamsal, L., Makar, P. A., Martin, R. V.,
578 Veefkind, J. P., and Yang, K.: Improved satellite retrievals of NO₂ and SO₂ over the Canadian oil sands and
579 comparisons with surface measurements, *Atmospheric Chemistry and Physics*, 14, 3637-3656, 10.5194/acp-14-3637-
580 2014, 2014.
- 581 Mijling, B., van der A, R. J., Boersma, K. F., Van Roozendaal, M., De Smedt, I., and Kelder, H. M.: Reductions of
582 NO₂ detected from space during the 2008 Beijing Olympic Games, *Geophysical Research Letters*, 36,
583 10.1029/2009gl038943, 2009.
- 584 Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J. L., Campbell, J. R., Christopher, S. A., Di
585 Girolamo, L., Giglio, L., Holz, R. E., Kearney, C., Miettinen, J., Reid, E. A., Turk, F. J., Wang, J., Xian, P., Zhao, G. Y.,
586 Balasubramanian, R., Chew, B. N., Janjai, S., Lagrosas, N., Lestari, P., Lin, N. H., Mahmud, M., Nguyen, A. X., Norris, B.,
587 Oanh, N. T. K., Oo, M., Salinas, S. V., Welton, E. J., and Liew, S. C.: Observing and understanding the Southeast Asian
588 aerosol system by remote sensing: An initial review and analysis for the Seven Southeast Asian Studies (7SEAS) program,
589 *Atmospheric Research*, 122, 403-468, 10.1016/j.atmosres.2012.06.005, 2013.
- 590
591 Sarkodie, S. A., and Strezov, V.: A review on Environmental Kuznets Curve hypothesis using bibliometric and meta-
592 analysis, *Science of the Total Environment*, 649, 128-145, 10.1016/j.scitotenv.2018.08.276, 2019.
- 593 Sayer, A. M., Hsu, N. C., Bettenhausen, C., and Jeong, M. J.: Validation and uncertainty estimates for MODIS
594 Collection 6 "Deep Blue" aerosol data, *Journal of Geophysical Research-Atmospheres*, 118, 7864-7872,
595 10.1002/jgrd.50600, 2013.
- 596 Sayer, A. M., Munchak, L. A., Hsu, N. C., Levy, R. C., Bettenhausen, C., and Jeong, M. J.: MODIS Collection 6
597 aerosol products: Comparison between Aqua's e-Deep Blue, Dark Target, and "merged" data sets, and usage
598 recommendations, *Journal of Geophysical Research-Atmospheres*, 119, 13965-13989, 10.1002/2014jd022453, 2014.
- 599 Schutgens, N., Sayer, A. M., Heckel, A., Hsu, C., Jethva, H., de Leeuw, G., Leonard, P. J. T., Levy, R. C., Lipponen,
600 A., Lyapustin, A., North, P., Popp, T., Poulson, C., Sawyer, V., Sogacheva, L., Thomas, G., Torres, O., Wang, Y.,
601 Kinne, S., Schulz, M., and Stier, P.: An AeroCom/AeroSat study: Intercomparison of Satellite AOD Datasets for
602 Aerosol Model Evaluation, *Atmos. Chem. Phys. Discuss.*, 2020, 1-43, 10.5194/acp-2019-1193, 2020.
- 603 Selden, T. M., and Song, D. Q.: Environmental Quality and Development - is there a Kuznets Curve for Air-Pollution
604 Emissions?, *Journal of Environmental Economics and Management*, 27, 147-162, 10.1006/jeeem.1994.1031, 1994.
- 605 Shah, V., Jacob, D. J., Li, K., Silvern, R. F., Zhai, S. X., Liu, M. Y., Lin, J. T., and Zhang, Q.: Effect of changing NO_x
606 lifetime on the seasonality and long-term trends of satellite-observed tropospheric NO₂ columns over China,
607 *Atmospheric Chemistry and Physics*, 20, 1483-1495, 10.5194/acp-20-1483-2020, 2020.
- 608 Shao, P. Y., Tian, H. Z., Sun, Y. J., Liu, H. J., Wu, B. B., Liu, S. H., Liu, X. Y., Wu, Y. M., Liang, W. Z., Wang, Y.,
609 Gao, J. J., Xue, Y. F., Bai, X. X., Liu, W., Lin, S. M., and Hu, G. Z.: Characterizing remarkable changes of severe
610 haze events and chemical compositions in multi-size airborne particles (PM₁, PM_{2.5} and PM₁₀) from January 2013
611 to 2016-2017 winter in Beijing, China, *Atmospheric Environment*, 189, 133-144, 10.1016/j.atmosenv.2018.06.038,
612 2018.

- 613 Shi, X., and Brasseur, G. P.: The Response in Air Quality to the Reduction of Chinese Economic Activities during the
614 COVID-19 Outbreak, *Geophysical Research Letters*, 0-1, 10.1029/2020GL088070, 2020.
- 615 Si, Y. D., Wang, H. M., Cai, K., Chen, L. F., Zhou, Z. C., and Li, S. S.: Long-term (2006-2015) variations and relations
616 of multiple atmospheric pollutants based on multi-remote sensing data over the North China Plain, *Environmental*
617 *Pollution*, 255, 10.1016/j.envpol.2019.113323, 2019.
- 618 Sogacheva, L., Popp, T., Sayer, A. M., Dubovik, O., Garay, M. J., Heckel, A., Hsu, N. C., Jethva, H., Kahn, R. A.,
619 Kolmonen, P., Kosmale, M., de Leeuw, G., Levy, R. C., Litvinov, P., Lyapustin, A., North, P., Torres, O., and Arola,
620 A.: Merging regional and global aerosol optical depth records from major available satellite products, *Atmospheric*
621 *Chemistry and Physics*, 20, 2031-2056, 10.5194/acp-20-2031-2020, 2020.
- 622 Strode, S. A., Worden, H. M., Damon, M., Douglass, A. R., Duncan, B. N., Emmons, L. K., Lamarque, J.-F., Manyin,
623 M., Oman, L. D., Rodriguez, J. M., Strahan, S. E., and Tilmes, S.: Interpreting space-based trends in carbon monoxide
624 with multiple models, *Atmospheric Chemistry and Physics*, 16, 7285-7294, 10.5194/acp-16-7285-2016, 2016.
- 625 Sun, W., Shao, M., Granier, C., Liu, Y., Ye, C. S., and Zheng, J. Y.: Long-Term Trends of Anthropogenic SO₂, NO_x,
626 CO, and NMVOCs Emissions in China, *Earth's Future*, 6, 1112-1133, 10.1029/2018ef000822, 2018.
- 627 United Nations Environment Program (UNEP): Independent Environmental Assessment: Beijing 2008 Olympic
628 Games. Nairobi, Kenya, 2009.
- 629 Wang, M., Zhu, T., Zheng, J., Zhang, R. Y., Zhang, S. Q., Xie, X. X., Han, Y. Q., and Li, Y.: Use of a mobile laboratory
630 to evaluate changes in on-road air pollutants during the Beijing 2008 Summer Olympics, *Atmospheric Chemistry and*
631 *Physics*, 9, 8247-8263, 10.5194/acp-9-8247-2009, 2009.
- 632 Wang, P., Chen, K., Zhu, S., Wang, P., and Zhang, H.: Severe air pollution events not avoided by reduced
633 anthropogenic activities during COVID-19 outbreak, *Resources, Conservation and Recycling*, 158, 104814,
634 10.1016/j.resconrec.2020.104814, 2020.
- 635 Wang, P. C., Elansky, N. F., Timofeev, Y. M., Wang, G. C., Golitsyn, G. S., Makarova, M. V., Rakitin, V. S., Shtabkin,
636 Y., Skorokhod, A. I., Grechko, E. I., Fokeeva, E. V., Safronov, A. N., Ran, L., and Wang, T.: Long-Term Trends of
637 Carbon Monoxide Total Columnar Amount in Urban Areas and Background Regions: Ground- and Satellite-based
638 Spectroscopic Measurements, *Advances in Atmospheric Sciences*, 35, 785-795, 10.1007/s00376-017-6327-8, 2018.
- 639 Wang, T., Nie, W., Gao, J., Xue, L. K., Gao, X. M., Wang, X. F., Qiu, J., Poon, C. N., Meinardi, S., Blake, D., Wang,
640 S. L., Ding, A. J., Chai, F. H., Zhang, Q. Z., and Wang, W. X.: Air quality during the 2008 Beijing Olympics:
641 secondary pollutants and regional impact, *Atmospheric Chemistry and Physics*, 10, 7603-7615, 10.5194/acp-10-7603-
642 2010, 2010.
- 643 Wang, Y., and Wang, J.: Tropospheric SO₂ and NO₂ in 2012-2018: Contrasting views of two sensors (OMI and
644 OMPS) from space, *Atmospheric Environment*, 223, 10.1016/j.atmosenv.2019.117214, 2020.
- 645 Warner, J., Carminati, F., Wei, Z., Lahoz, W., and Attie, J. L.: Tropospheric carbon monoxide variability from AIRS
646 under clear and cloudy conditions, *Atmospheric Chemistry and Physics*, 13, 12469-12479, 10.5194/acp-13-12469-
647 2013, 2013.
- 648 Witte, J. C., Schoeberl, M. R., Douglass, A. R., Gleason, J. F., Krotkov, N. A., Gille, J. C., Pickering, K. E., and
649 Livesey, N.: Satellite observations of changes in air quality during the 2008 Beijing Olympics and Paralympics,
650 *Geophysical Research Letters*, 36, 10.1029/2009gl039236, 2009.

- 651 Xie, G. Q., Wang, M., Pan, J., and Zhu, Y.: Spatio-temporal variations and trends of MODIS C6.1 Dark Target and
652 Deep Blue merged aerosol optical depth over China during 2000-2017, *Atmospheric Environment*, 214,
653 10.1016/j.atmosenv.2019.116846, 2019.
- 654 Xu, J. H., Xie, H. M., Wang, K., Wang, J., and Xia, Z. S.: Analyzing the spatial and temporal variations in tropospheric
655 NO₂ column concentrations over China using multisource satellite remote sensing, *Journal of Applied Remote
656 Sensing*, 14, 10.1117/1.jrs.14.014519, 2020.
- 657 Yu, S. M., Yuan, J. G., and Liang, X. Y.: Trends and Spatiotemporal Patterns of Tropospheric NO₂ over China During
658 2005-2014, *Water Air and Soil Pollution*, 228, 10.1007/s11270-017-3641-9, 2017.
- 659 Yumimoto, K., Uno, I., and Itahashi, S.: Long-term inverse modeling of Chinese CO emission from satellite
660 observations, *Environmental Pollution*, 195, 308-318, 10.1016/j.envpol.2014.07.026, 2014.
- 661 Zhang, Y., Li, C., Krotkov, N. A., Joiner, J., Fioletov, V., and McLinden, C.: Continuation of long-term global SO₂
662 pollution monitoring from OMI to OMPS, *Atmospheric Measurement Techniques*, 10, 10.5194/amt-10-1495-2017,
663 2017.
- 664 Zhao, Y., Nielsen, C. P., McElroy, M. B., Zhang, L., and Zhang, J.: CO emissions in China: Uncertainties and
665 implications of improved energy efficiency and emission control, *Atmospheric Environment*, 49, 103-113,
666 10.1016/j.atmosenv.2011.12.015, 2012.
- 667 Zhao, Y., Zhang, J., and Nielsen, C. P.: The effects of recent control policies on trends in emissions of anthropogenic
668 atmospheric pollutants and CO₂ in China, *Atmospheric Chemistry and Physics*, 13, 487-508, 10.5194/acp-13-487-
669 2013, 2013.
- 670 Zheng, B., Chevallier, F., Ciais, P., Yin, Y., Deeter, M. N., Worden, H. M., Wang, Y. L., Zhang, Q., and He, K. B.:
671 Rapid decline in carbon monoxide emissions and export from East Asia between years 2005 and 2016, *Environmental
672 Research Letters*, 13, 10.1088/1748-9326/aab2b3, 2018a.
- 673 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C. P., Geng, G. N., Li, H. Y., Li, X., Peng, L. Q., Qi, J., Yan, L., Zhang,
674 Y. X., Zhao, H. Y., Zheng, Y. X., He, K. B., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as
675 the consequence of clean air actions, *Atmospheric Chemistry and Physics*, 18, 14095-14111, 10.5194/acp-18-14095-
676 2018, 2018b.
- 677

678 **Tables**

679

680 **Table 1. Summary statistics for central east China comparing 2020 and 2019 during January 23 – April 8.**

Variable	2020 mean	2019 mean	2020 % difference from 2019
CO (ppbv)	133.5 (130.3, 136.8)	137.9 (134.7, 141.3)	-3.2 (-6.3, 0.1)
SO₂ (DU)	0.057 (0.045, 0.070)	0.031 (0.018, 0.046)	95 (14.8, 249.6)
NO₂ (10 ¹⁵ molec cm ⁻²)	6.5 (5.8, 7.2)	9.6 (8.7, 10.5)	-32.1 (-42.1, -21.7)
AOD	0.41 (0.36, 0.46)	0.48 (0.41, 0.55)	-14.3 (-29.4, 3.1)

681

682

683 **Table 2. Same as Table 1, but for southern China.**

Variable	2020 mean	2019 mean	2020 % difference from 2019
CO (ppbv)	144.7 (139.6, 150.3)	128.5 (124.4, 132.8)	12.6 (7.2, 18.3)
SO₂ (DU)	0.003 (-0.01, 0.020)	-0.020 (-0.04, -0.001)	116 (24, 223)
NO₂ (10 ¹⁵ molec cm ⁻²)	3.3 (3.0, 3.7)	4.3 (3.9, 4.7)	-22.2 (-32.6, -10.4)
AOD	0.38 (0.34, 0.43)	0.34 (0.30, 0.39)	12 (-7, 34)

684

685 **Table 3. 2014 anthropogenic emissions estimates by sector (in %) over China, excluding biomass burning, from the**
 686 **Community Emissions Data System (CEDs) for a representative set of constituents: black carbon (BC), carbon monoxide**
 687 **(CO), ammonia (NH₃), nitrogen oxides (NO_x), organic carbon (OC) and sulfur dioxide (SO₂). Residential, commercial and**
 688 **other sectors are combined as RCO.**


	BC	CO	NH₃	NO_x	OC	SO₂
Agriculture	0	0	61.6	1.1	0	0
Energy	32.6	8	0.4	38.5	28.3	29.4
Industrial	12.7	41.8	6.5	33	5.1	57.3
Ground transportation	8.1	7.2	0.5	17.5	1.7	0.3
RCO	38.1	36.7	5.2	4.2	38.4	12.5
Solvents	0	0	0	0	0	0
Waste	8.5	6.3	25.8	5.2	26.5	0.4
Shipping	0	0	0	0.2	0	0.1
Aircraft	0	0	0	0.2	0	0

689

690 **Table 4. Bottom up biomass Global Fire Assimilation System (Kaiser et al., 2012) burning CO emissions estimates from the**
 691 **Upper Mekong region (17° N to 24° N, 95° E to 105° E) and AIRS CO over southern China from January 23 to April 8, for**
 692 **2005-2020.**


Year	GFAS CO	AIRS CO
	Upper Mekong (KT)	southern China 500 hPa (ppbv)
2005	7977	157
2006	8905	146
2007	15734	165
2008	4542	153
2009	9990	140
2010	14176	149
2011	3591	147
2012	11320	153
2013	8684	145
2014	8722	142
2015	8084	143
2016	9642	149
2017	3736	131
2018	3179	139
2019	6309	128
2020	7871	145

693



695

696 **Figure 1. Groupings of provinces for central east China and southern China.**



697
698
699
700

Figure 2. 2020-2019 differences during January 23 to April 8 over China in a) AIRS carbon monoxide (CO) at 500 hPa, b) OMI PBL sulfur dioxide (SO₂), c) OMI tropospheric nitrogen dioxide (NO₂) and d) Aqua MODIS aerosol optical depth (AOD).




Figure 3. Monthly mean a) AIRS CO, b) OMI PBL SO₂, c) OMI tropospheric NO₂ and d) MODIS AOD over central east China since 2005. As in Bauwens et al. (2020), each year starts in August to show any departure from the seasonal cycle during the January 23 to April 8 lockdown period, shown by the thin grey vertical lines.






Figure 4. Same as Figure 3, but for southern China.



710 **Figure 5. January 23-April 8 box plots over central East China for a) AIRS CO, b) OMI PBL SO₂, c) OMI tropospheric NO₂ and d) Aqua and Terra MODIS AOD from 2005 to 2020. The black box plots show the median, interquartile range and 2.5th and 97.5th percentiles over all daily data, with the mean shown by the black dot.**



715 **Figure 6. Dependence of trends (red) and difference between 2020 observations and predicted value (magenta) on detrending start year over central east China for a) AIRS CO, b) OMI PBL SO₂, c) OMI tropospheric NO₂ and d) MODIS AOD. The solid line shows the mean of the estimate for each year and the shading shows the 95% confidence interval.**




Figure 7. Same as Figure 5 but for southern China.




Figure 8. Same as Figure 6, but for southern China.