

Anonymous Referee #1

We thank the Reviewer for their positive and helpful suggestions. We respond to their comments below

- 5 -Figure 3: marking a horizontal line with the $p < 0.05$ significance would aid interpretation. Also in the paragraph that discusses these results, comment on the significance of the correlations.

Author Response

- 10 *Estimating a statistical significance for the pattern correlations in Figure 3 is very difficult, since it would require a calculation of spatial decorrelation length scales (zonal and meridional) in sea ice and SAT, in order to accurately estimate the degrees-of-freedom (I.e. number of independent spatial pints in the trend patterns); this is a much more complex task than estimating significance of temporal correlations (as we have done for Figure 5).*
- 15 *Whilst we agree that it would be a nice addition to have such a significance estimate, it would greatly increase the complexity of the Methods (as the Reviewer notes, the intuitive simplicity of the Method is a strength of the paper), and it is our opinion that this would not alter the conclusions.*

- Line 133. You conclude that JRA55 and 20CRv3 have the best representations of long-term change over the Southern Ocean. I would clarify the phrasing here that they have the best representation over the study period. As JRA55 goes back to 1958, and 20CRv3 back to 1836, this could send the message to an uncautious reader that this comment applies to earlier periods. Given the lower amounts of data going into both reanalyses in earlier period, we do not know whether this conclusion holds for earlier periods (and you obviously can't test it due to lack of earlier sea-ice data).
- 25

Author Response

- This is an important caveat, and we have amended the sentence to read as follows: 'Based on this analysis we would conclude that JRA55 and 20CRv3 have the best representations of change since the late 1970s over the polar Southern Ocean, under the assumption that SIC trends should be closely related to SAT trends. We note that this may not hold true for earlier periods which are unconstrained by satellite retrievals, and for which we do not have reliable sea ice observations.'*
- 30

- 35 -Line 134 and 158, write numbers < 10 in full 'Three' and 'Two'.
We have edited the manuscript accordingly

- Table 1: I think the '1980' in the reanalysis period box for MERRA 2 may be in bold

Author Response

- 40 *Bold years indicated the first and last year for which data is available from all the reanalyses (1980 constrained by MERRA2; 2010 constrained by ERA20C. However, on consideration we feel the bold type does not add to the information, and have removed the bold type, and amended the Table caption accordingly.*

- Figure 1 is labelled as Figure 2.

45 *Author Response*

Typo corrected

- In the actual Figure 2, panel e needs clearer description. Define what 'S-RIP' is, and also are the black dots/line observations?

50 *Author Response*

Figure 2e legend amended to include 'Observations' for black lines/markers. S-RIP abbreviation is now expanded to Figure 2 caption

Anonymous Referee #2

55

We thank the Reviewer for their positive and helpful suggestions. We respond to their comments below

60 1. In discussing Fig 1 in Section 1 (showing SAT trends in the reanalyses), I think a note should be made of the fact that ERA20C is reliant on surface observations alone, of which there are very few in the Southern Ocean, so it is perhaps not surprising that it appears as an outlier.

Author Response

We have added the following sentence to this section:

65 *'Much of this spread is due to differences in the forecast model and assimilation technique, but it should be noted that some of the products (ERA20C and 20CRv3) are not constrained by satellite data in order to give a consistent product over long historical periods; this is a major limitation in the remote Antarctic region.'*

70 2. It would be nice to see some measure of statistical significance in Fig 5 (for instance, as a gray shaded region). For 30 year time series I would guess that a lot of the correlations are not significant.

Author Response

75 *As noted in the Figure 5 caption, only SAT trends that are significant at the 90% confidence level are plotted, and only areas where both the SAT and SIC trends are statistically-significant at the 90% level (and the SAT and SIC trends are inconsistent) are hatched.*

80 3. It is remarked that JRA55 is perhaps the best in terms of consistency between SAT and SIC trends. However, it is notable that it is quite an outlier in terms of the seasonal cycle of ice growth, having stronger growth rates in the fall/early winter than other reanalyses of observations (Fig 2e). I think the paper would benefit from some discussion of this – i.e. is JRA55 more reliable in terms of SAT trends, but perhaps less so in its seasonal cycle?

Author Response

We have added the following sentences to our summary fo Figure 3:

85 *'We note that this may not hold true for earlier periods which are unconstrained by satellite retrievals, and for which we do not have reliable sea ice observations. An interesting point to note is that although JRA55 performs well with respect to this metric, it has the strongest sea ice bias in March and April. This raises the question of whether the mean state is a good indicator of performance in respect of variability or trends.'*

90 4. L62: particular -> particularly
Author Response
Typo corrected

95 5. Fig 2e is not labelled with an 'e'.
Author Response
Figure caption added to panel 2e

100 6. L151: 'Sat' -> 'SAT'
Author Response
Typo corrected

7. L181: This sentence doesn't make sense. I think 'have shown' needs to be removed
Author Response
105 *Correct, we have corrected this sentence which now reads:*
'Whilst it has been hypothesized from model simulations that due to complex ocean-sea ice feedbacks, a surface cooling may lead to a loss of sea ice (Zhang, 2007), we note that in this particular region the sea ice loss has been robustly attributed to increasing poleward airflow, which both dynamically constrains the ice extent and advects warm air to the region (Holland and Kwok, 2012; Hosking et al, 2013; Raphael et al, 2016), and would be expected to drive warmer SAT.'
110

Validation of reanalysis Southern Ocean atmosphere trends using sea ice data

115 William R. Hobbs^{1,2}, Andrew R. Klekociuk^{1,3}, Yuhang Pan⁴

¹Australian Antarctic Program Partnership, Institute of Marine and Antarctic Studies, Private Bag 129, University of Tasmania, Hobart, TAS 7001, Australia.

²ARC Centre of Excellence for Climate Extremes, Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, TAS 7001, Australia.

120 ³Antarctica and the Global System Program, Australian Antarctic Division, 203 Channel Highway, Kingston, TAS 7050, Australia.

⁴Independent Researcher

Correspondence to: William R. Hobbs (will.hobbs@utas.edu.au)

125

Abstract. Reanalysis products are an invaluable tool for representing variability and long-term trends in regions with limited in-situ data, and especially the Antarctic. A comparison of 8 different reanalysis products shows large differences sea level pressure and surface air temperature trends over the high latitude Southern Ocean, with implications for studies of the atmosphere's role in driving ocean-sea ice changes. In this study, we use the established close coupling between sea ice cover and surface temperature to evaluate these reanalysis trends using the independent, 30-year sea ice record from 1980-2010. We demonstrate that sea ice trends are a reliable validation tool for most months of the year, although the sea ice-surface temperature coupling is weakest in summer when the surface energy budget is dominated by atmosphere-to-ocean heat fluxes. Based on our analysis, we find that surface air temperature trends in JRA55 are most consistent with satellite-observed sea ice trends over the polar waters of the Southern Ocean.

130

135 1 Introduction

Atmospheric trends in the Southern high latitudes have global importance. Wind patterns are essential for driving the Southern Ocean overturning, which is responsible for most of the global ocean's uptake of anthropogenic heating, and approximately half its uptake of anthropogenic carbon (Frolicher *et al*, 2015). Local wind changes are a factor in the ocean melting of West Antarctic ice shelves (Lenaerts *et al*, 2017; Paolo *et al*, 2018; Dotto *et al*, 2019), with implications for global barostatic sea level rise (Dupont and Alley, 2005; Pritchard *et al*, 2012; Paolo *et al*, 2015), and polar winds are clearly related to observed Antarctic sea ice trends (Holland and Kwok, 2012). More immediately, variability in the Southern Annular Mode – the dominant mode of mid-high latitude Southern Hemisphere atmospheric variability – is thought to influence Australian rainfall (Meneghini *et al*, 2007), with implications for current and future droughts. Clearly, a reliable and accurate representation of high latitude Southern Hemisphere atmosphere trends is essential.

140

145 For this data-sparse region, atmospheric reanalysis products are the primary research tool for analyzing observed changes or as surface boundary conditions for ocean-sea ice models. However, there is a wide spread in surface atmosphere trends over the Southern Ocean amongst different reanalysis products, which introduces uncertainty when interpreting observed ocean and

150 sea ice trends (Marshall, 2003; Swart and Fyfe, 2012; Hobbs *et al.*, 2016). Reanalysis validation studies have attempted to address this uncertainty, but have largely been restricted to comparisons with long-term surface measurement sites, almost all of which are located near the Antarctic coast (e.g. Turner *et al.*, 2014); relatively few studies have been conducted for the sea-ice zone (Bracegirdle and Marshall, 2012; Jones *et al.*, 2016).

155 Figure 1 shows linear trends in 2m air temperature (SAT) and mean sea level pressure (MSLP) from eight commonly-used atmosphere reanalyses (summarized in Table 1), and clearly demonstrates this spread in trends. Much of this spread is due to differences in the forecast model and assimilation technique, but it should be noted that some of the products (ERA20C and 20CRv3) are not constrained by satellite data in order to give a consistent product over long historical periods; this is a major limitation in the remote Antarctic region. Some reanalyses show almost no warming at all in West Antarctica, whilst NCEP2 shows a warming over the entire sea ice zone. Station data show a distinct asymmetry in the long-term behavior of SAT between Antarctica's eastern and western hemispheres, with the statistical significance of trends depending on epoch, season and location. A general warming has occurred in recent decades in the Antarctic Peninsula and parts of West
160 Antarctica, with weaker mixed trends in East Antarctica (Marshall *et al.*, 2013; Nicolas and Bromwich, 2014; Turner *et al.*, 2014). Compared with station measurements, SAT trends in reanalyses show less consistency with spurious behavior in some regions, particularly in East Antarctica where surface stations are sparse (Bromwich *et al.*, 2013; Steig and Orsi, 2013; Wang *et al.*, 2016; Simmons *et al.*, 2017). However, there is generally better agreement between observations and reanalyses when interannual variability rather than trends are considered (e.g. Wang *et al.*, 2016).

165 There is a similar spread in MSLP trends; many of the reanalyses show the widely-reported deepening of the Amundsen Sea Low (Hosking *et al.*, 2013; Turner *et al.*, 2013; Raphael *et al.*, 2016) – although with some disagreement on magnitude and exact location – but by no means all of them. Additionally, there is a known spread amongst reanalyses in the magnitude of the Southern Annular Mode positive trend (Marshall, 2003; Swart and Fyfe, 2012). This raises the question of which representation is the most accurate, for interpreting recent historical changes in the atmosphere-ocean-cryosphere system of
170 the polar Southern Ocean.

There is a close link between sea ice cover and the atmosphere, both for interannual variability and at longer time scales (e.g. Comiso *et al.*, 2017). Atmospheric thermal advection modulates the rate of sea ice freeze/melt, and wind driven ice motion redistributes the existing sea ice. In the Southern Ocean the sea ice-atmosphere relationship tends to be stronger in the sea ice growth season and weaker in the melt season (Raphael and Hobbs, 2014; Schroeter *et al.*, 2017). This may be because
175 approximately half of the heat driving sea melt comes from the ocean (Gordon, 1981), diminishing the relative impact of the atmosphere. The atmosphere-sea ice relationship is particularly strong for surface air temperature (e.g. Comiso *et al.*, 2017), due to positive feedbacks: a colder air temperature leads to increased sea ice cover, which due to increased albedo and much reduced ocean-to-atmosphere heat flux can further reduce air temperature. In short, sea ice affects air temperature, and air temperature affects sea ice.

180 Previous studies have exploited this close relationship to study sea ice. Notably, King and Harangozo (1998) demonstrated a close link between Antarctic Peninsula station temperature and local sea ice changes, Massonnet *et al.* (2013) were able to

reproduce Antarctic sea ice variability in a model driven by SAT, and both Kusahara *et al* (2017) and Schroeter *et al* (2018) showed the important role that thermodynamic forcing has on Antarctic sea ice trends. This close coupling between SAT and sea ice concentration (SIC) indicates that the passive microwave sea ice record may be used as an independent validation of Reanalysis SAT trends, at least for the broad spatial patterns that are clearly different in Figure 1.

In this study, we perform just such an evaluation. We demonstrate that SIC and SAT variability are closely related for much of the year, except for the season of strongest sea ice melt. Based on that premise, we find that a number of reanalysis products have trends that are physically-consistent with independently-observed sea ice trends, with ERA5 showing a marginally-better agreement than other products. A smaller group of products are very obviously inconsistent with the sea ice trends, and should be avoided for studies of long-term change in the high-latitude Southern Hemisphere. We argue that the weak SAT-SIC relationship in summer is due to the direction of ocean-atmosphere heat flux in those months; since the net balance in the sea ice zone is *from* atmosphere *to* ocean, the surface energy budget is more a response to - rather than a driver of - the near-surface atmosphere.

195 **2 Data and Method**

We use monthly mean SIC from passive microwave satellite observations as the primary dataset for evaluating reanalysis SAT trends. Specifically, we use the Goddard-merged data from the NOAA/NSIDC climate data record for SIC, available on a 25 km x 25 km equal area grid (Meier *et al*, 2014).

We analyze monthly-mean SIC, SAT and MSLP from 8 publicly-available reanalysis products, which are summarized in Table 1. These products span a range of spatial resolutions, assimilation algorithms and analysis periods. For this study, we consider the period 1980-2010 inclusive, which is the longest period covered by all 8 reanalyses, constrained by MERRA2 (starting in January 1980) and ERA-20C (ending in December 2010), and matches the period of the SPARC Reanalysis Intercomparison Project (S-RIP: Fujiwara *et al*, 2017).

Although we consider the relationship between SAT and SIC at interannual timescales, our primary focus is on the 31-year trends of the analysis period, calculated by month using Ordinary Least Squares regression. To quantify the level of agreement between trend patterns for SIC and reanalysis SAT, we use an uncentered pattern correlation (i.e. without removing spatial means), applying a cosine-weighting to account for latitude-dependence of the grid area. To facilitate this, all variables were regridded onto a common 1° x 1° latitude/longitude grid using bilinear interpolation.

210 **3 Results**

3.1 Evaluation of SAT based on sea ice trends

Although sea ice trends are not themselves the focus of this work, except as an independent validation of reanalysis SAT, the observed SIC trends are shown in Figure 2 for illustrative purposes. (Note that while the trends are aggregated in Figure 2 into seasons defined by sea ice melt/growth, for the SAT validation we used monthly trends, shown in Supplemental Material).

215 The trend patterns are well-established and have been described in many previous studies (e.g. Parkinson and Cavalieri, 2012; Hobbs *et al.*, 2016; Comiso *et al.*, 2017), and can be broadly summarized as a decrease in the Amundsen and Bellingshausen Seas (60° – 120°W), with compensating increases in the western Ross Sea (150°E – 180°E), Weddell Sea and King Haakon VII Sea (50°W – 30°E). There is some seasonal dependence, and the Ross Sea is the only region that has statistically-significant trends in all seasons. From this SIC pattern, we would expect a warming SAT trend in the region of the Antarctic Peninsula, and a cooling elsewhere, a pattern that is expressed by some of the reanalyses in Figure 1, but by no means all.

To quantify the level of agreement between SIC and SAT trends, we calculated correlations for each season and reanalysis amongst observed SIC, reanalysis SIC, and reanalysis SAT trends (Figure 3). Most of the reanalyses use boundary sea ice conditions that match the passive microwave record reasonably-well, with trend pattern correlations consistently greater than 0.9 for many products, the best match being for ERA5 (Figure 3a). Both ERA-int and MERRA2 have a sea ice boundary condition that diverges somewhat from observations in late winter. The NCEP reanalyses (i.e. NCEP2 and CFSR), have a coupled, freely-evolving ocean-sea ice system, which explains the very low agreement with the satellite record compared to the products which use a prescribed sea ice condition.

Figure 3b shows the pattern correlation between reanalysis SAT trends and the trend of each reanalysis product's prescribed SIC trend. This serves as a test for the expectation that SAT and prescribed SIC trends should be internally-consistent, regardless of the actual SIC trend pattern. For most months and reanalyses this is indeed the case, with strong negative correlations for most of the year. NCEP2 is the exception and shows a positive correlation for much of the year, presumably because of its strong warming pattern in the sea ice zone (Figure 1) that is inconsistent with an increased sea ice cover in much of the Antarctic domain. The sea ice-SAT trend relationship is strongest in the main sea ice growth season (March-July), consistent with previous model and observational studies showing that the sea ice-atmosphere relationship is stronger during the growth season (Raphael and Hobbs, 2014; Schroeter *et al.*, 2017). The relationship is surprisingly weak in December and January, months combining high melt rates with relatively low sea ice cover. We explore this result in more detail in section 3.2, and note that for these months SIC trends may be a less reliable test for SAT trends.

Figure 3c shows the pattern correlations between observed SIC trends and reanalysis SAT, and therefore summarizes the evaluation of SAT in the reanalyses based on Antarctic sea ice trends. Other than for December-January when the SAT-SIC relationship is relatively weak, JRA55 and 20CRv3 have consistently the closest relationship between SAT and SIC trends, despite ERA5 having a closer correspondence between prescribed SIC and the satellite record (Figure 3a). ERA20C and ERA5 also have reasonable agreement for much of the year, but none of the ERA products have any SAT-SIC trend correlation in December-January. ERA-int also has a notably weak SAT-SIC relationship in August, which seems to be largely due to the disagreement between the observed SIC trend and that of ERA-int's sea ice (Figure 3a), since the correlation between ERA-int SIC and SAT is strongly negative in August (Figure 3b).

Based on this analysis we would conclude that JRA55 and 20CRv3 have the best representations of change since the late 1970s over the polar Southern Ocean, under the assumption that SIC trends should be closely related to SAT trends. We note that

Deleted: long-term

250 this may not hold true for earlier periods which are unconstrained by satellite retrievals, and for which we do not have reliable sea ice observations. An interesting point to note is that although JRA55 performs well with respect to this metric, it has the strongest sea ice bias in March and April. This raises the question of whether the mean state is a good indicator of performance in respect of variability or trends.

The summer SIC-SAT relationship is much weaker in summer for all the products, but in the ~~three~~ ECMWF products analyzed here (ERA5, ERA-int and ERA20C) this relationship completely disappears. We further tested the physical relationship between sea ice and air temperature by mapping the correlation coefficients between detrended, interannually-varying reanalysis SAT and SIC, for each calendar month (Supplemental Figures S25-S36). The results are similar to those for the trend pattern correlations, showing a strong negative correlation throughout the ice pack in most months, but which is weaker and more complex in summer. In mid-winter the correlations are concentrated at the sea ice edge, where sea ice variability is greatest. The reanalyses with very high SIC have limited correlations within the ice pack, since sea ice concentrations > 0.9 must have limited variance and therefore weak covariance with SAT. In the next section, we explore the weak SAT-SIC relationship in December-January in more detail.

Deleted: 3

3.2 Further exploration of summer SAT-sea ice relationship

265 The correlations show that generally the relationship between SIC and SAT 30-year trends is weaker in December and January (Figure 3b). It is worthwhile considering the physical reasons for this weak summer relationship between SAT and sea ice, which are strongly coupled for the rest of the year. We note that December and January are months with very strong sea ice melt (Figure 2e), due to a surface heat flux from atmosphere to ocean. By contrast, for most of the year (March-October), the heat flux is from ocean-to-atmosphere, driving the ocean cooling that allows sea ice to form, and for these months there is a positive relationship between the magnitude of cooling and the strength of the SAT-SIC correlation for the reanalyses (Figure 4), i.e. stronger cooling leads to a stronger negative SIC-SAT correlation. During these cold months, SAT is cooler where sea ice prevents a flux of heat from ocean-to-atmosphere and warmer over open water, which is the physical mechanism explain the negative correlation between SAT and SIC trends.

Deleted: Sat

During warmer months when the mean flux is from atmosphere-to-ocean, this relationship breaks down or even becomes negative (Figure 4), indicating that surface heat flux is no longer a connection between ice and SAT variability. At a first pass we might still expect a relationship between the surface atmosphere and sea ice, since this heat flux is important for melting the sea ice. However, we suggest that ~~two~~ factors combine to break the SAT-SIC coupling. The first is that, from the perspective of the atmosphere (and in particular, an uncoupled atmosphere-only model which is the basis for most of the reanalyses), when the flux is from atmosphere-to-ocean, the ocean is a passive sink of energy that is modulated by atmospheric processes, rather than an active driver of the surface atmosphere during periods of ocean-to-atmosphere energy transfer. We note that the CFSR – one of the only reanalyses to have a coupled rather than prescribed sea ice – has a stronger summer SAT-SIC correlation (Figure 3).

Deleted: 2

The second factor is the process of sea ice melt. Although the melt is largely driven by incoming solar radiation, there is in fact relatively little melt on top of the sea ice because of the high albedo of snow-covered sea ice (Gordon, 1981; Drinkwater and Xiang, 2000). Instead, areas of open water such as leads absorb solar radiation, warming the ocean mixed layer and melting the ice pack from beneath (Stammerjohn *et al.*, 2012). This means that impact of ice cover, which in summer mainly affects the surface air temperature through reflecting solar radiation, is spatially diffused by the ocean and so reduces the direct spatial relationship between solar radiation, atmosphere, and sea ice.

A third and final factor, that is not immediately evident, is the nature of the summer ice pack, which although small in area comprises a higher proportion of thick, wind-compacted sea ice at the coastline than other months, since this is the ice most likely to survive the spring melt. This thick, compacted sea ice is relatively insensitive to atmospheric warming (Enomoto and Ohmura, 1990; Massom *et al.*, 2008), and so the sea ice-atmosphere relationship is also weak. As a result of these factors, we argue that a weak summer relationship between SIC and SAT is expected from the physical conditions in high summer.

However, this does not explain why the ERA products have apparently no SIC-SAT trend correlation in summer. Further analysis of the spatial distribution of SIC and SAT trends in Figure 5 reveals a local inconsistency between summer sea ice trends and the ERA SAT trends; this inconsistency is shown by hatching, which shows where a sea ice reduction is accompanied by a local cooling, or *vice versa*. The ERA products all show, to a greater or lesser degree of statistical-significance, a cooling over the Amundsen-Bellinghshausen Seas (60°W-150°W), a region with an intense and well-observed loss of summer sea ice (e.g. Parkinson, 2019). None of the other products show a statistically-significant cooling in the same region. Whilst it has been hypothesized from model simulations that due to complex ocean-sea ice feedbacks, a surface cooling may lead to a loss of sea ice (Zhang, 2007), we note that in this particular region the sea ice loss has been robustly attributed to increasing poleward airflow, which both dynamically constrains the ice extent and advects warm air to the region (Holland and Kwok, 2012; Hosking *et al.*, 2013; Raphael *et al.*, 2016), and would be expected to drive warmer SAT. Furthermore, whilst there are no direct measurements of SAT in this region there are a number of station observation records on the west Antarctic Peninsula, and reconstructions of continental surface temperature; these all indicate a warming trend over the west Antarctic landmass (Steig *et al.*, 2009; Nicolas and Bromwich, 2014) in response to the same increase in warm northerly airflow that has reduced Amundsen-Bellinghshausen sea ice cover. This continental warming is most clearly evident in JRA55 (Figure 5), but clearly does not seem to be consistent with a surface cooling over the adjacent ocean. We therefore consider that the ERA products must be considered with some degree of caution, especially for studies of change in the West Antarctic region.

4. Conclusions

Using the known close relation between sea ice cover and surface air temperature in polar oceans, we use satellite-observed of Antarctic sea ice trends as an independent validation of reanalysis trends over the polar Southern Ocean. Based on this analysis, we find several reanalysis products that reproduce reasonable surface air temperature trends, with JRA55 showing the consistently highest agreement with observed sea ice throughout the year.

Deleted:

Deleted: have shown

Deleted: ,

Deleted: due to complex ocean-sea ice feedbacks

We find that the relationship between surface air temperature and sea ice concentration is strong for most months of the year except mid-summer (i.e. December and January). These are the only months of the year when the polar Southern Ocean is a sink rather than a source of the net surface heat flux, and although much of the heat is used to melt sea ice, the heat is distributed by ocean processes, and the direct spatial correlation between sea ice and air temperature is relatively weak.

Although all eight of the reanalysis products that we analyze here have a weaker air temperature -sea ice relationship in summer, the ECMWF reanalyses (ERA5, ERA-int and ERA20C) have no correlation in summer at all. These seems to be due to their representation of a surface cooling in the Amundsen and Bellingshausen Seas, which is not consistent with a robustly-observed local sea ice loss since the late 1970s, nor with independent reconstructions of land surface temperature, which show a warming on the west Antarctic region adjacent to the Amundsen-Bellingshausen Seas.

Acknowledgements

The authors thank the two Anonymous Reviewers for their helpful and constructive comments that have improved this manuscript. This work was supported in part by the Antarctic Climate and Ecosystems Cooperative Research Centre, and the Australian Research Council Antarctic Gateway Partnership. Data analysis and visualization was performed using the NCAR Command Language (<http://dx.doi.org/10.5065/D6WD3XH5>). The authors would also like to acknowledge the help and resources of the NCAR Research Data Archive (<https://rda.ucar.edu/>), and reanalysis.org in acquiring and processing the data.

Deleted: T

References

- Bracegirdle, T. J., and Marshall, G. J.: The Reliability of Antarctic Tropospheric Pressure and Temperature in the Latest Global Reanalyses, *J Climate*, 25, 7138-7146, 10.1175/JCLI-D-11-00685.1, 2012.
- Bromwich, D. H., Nicolas, J. P., Monaghan, A. J., Lazzara, M. A., Keller, L. M., Weidner, G. A., and Wilson, A. B.: Central West Antarctica among the most rapidly warming regions on Earth, *Nat Geosci*, 6, 139-145, 10.1038/Ngeo1671, 2013.
- Comiso, J. C., Gersten, R. A., Stock, L. V., Turner, J., Perez, G. J., and Cho, K.: Positive Trend in the Antarctic Sea Ice Cover and Associated Changes in Surface Temperature, *J Climate*, 30, 2251-2267, 10.1175/jcli-d-16-0408.1, 2017.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, L., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q J Roy Meteor Soc*, 137, 553-597, 10.1002/qj.828, 2011.

Dotto, T. S., Naveira Garabato, A. C., Bacon, S., Holland, P. R., Kimura, S., Firing, Y. L., Tsamados, M., Wählin, A. K., and
355 Jenkins, A.: Wind-Driven Processes Controlling Oceanic Heat Delivery to the Amundsen Sea, Antarctica, *J Phys Oceanogr*,
49, 2829-2849, 10.1175/JPO-D-19-0064.1, 2019.

Drinkwater, M. R., and Xiang, L.: Seasonal to interannual variability in Antarctic sea-ice surface melt, *Icee T Geosci Remote*,
38, 1827-1842, 10.1109/36.851767, 2000.

Dupont, T. K., and Alley, R. B.: Assessment of the importance of ice-shelf buttressing to ice-sheet flow, *Geophysical Research*
360 *Letters*, 32, L04503, 10.1029/2004gl022024, 2005.

Enomoto, H., and Ohmura, A.: The influences of atmospheric half-yearly cycle on the sea ice extent in the Antarctic, *Journal*
of *Geophysical Research: Oceans*, 95, 9497-9511, 10.1029/JC095iC06p09497, 1990.

Frolicher, T. L., Sarmiento, J. L., Paynter, D. J., Dunne, J. P., Krasting, J. P., and Winton, M.: Dominance of the Southern
Ocean in Anthropogenic Carbon and Heat Uptake in CMIP5 Models, *J Climate*, 28, 862-886, 10.1175/Jcli-D-14-00117.1,
365 2015.

Fujiwara, M., Wright, J. S., Manney, G. L., Gray, L. J., Anstey, J., Birner, T., Davis, S., Gerber, E. P., Harvey, V. L., Hegglin,
M. I., Homeyer, C. R., Knox, J. A., Krüger, K., Lambert, A., Long, C. S., Martineau, P., Molod, A., Monge-Sanz, B. M.,
Santee, M. L., Tegtmeier, S., Chabrilat, S., Tan, D. G. H., Jackson, D. R., Polavarapu, S., Compo, G. P., Dragani, R.,
Ebisuzaki, W., Harada, Y., Kobayashi, C., McCarty, W., Onogi, K., Pawson, S., Simmons, A., Wargan, K., Whitaker, J. S.,
370 and Zou, C. Z.: Introduction to the SPARC Reanalysis Intercomparison Project (S-RIP) and overview of the reanalysis
systems, *Atmos. Chem. Phys.*, 17, 1417-1452, 10.5194/acp-17-1417-2017, 2017.

Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G.,
Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim,
G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S.
375 D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2
(MERRA-2), *J Climate*, 30, 5419-5454, 10.1175/JCLI-D-16-0758.1, 2017.

Gordon, A. L.: Seasonality of Southern-Ocean Sea Ice, *J Geophys Res-Oceans*, 86, 4193-4197, DOI
10.1029/JC086iC05p04193, 1981.

Hersbach, H., de Rosnay, P., Bell, B., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Alonso-Balmaseda, M., Balsamo, G.,
380 Bechtold, P., Berrisford, P., Bidlot, J.-R., de Boissésou, E., Bonavita, M., Browne, P., Buizza, R., Dahlgren, P., Dee, D.,
Dragani, R., Diamantakis, M., Flemming, J., Forbes, R., Geer, A. J., Haiden, T., Hólm, E., Haimberger, L., Hogan, R., Horányi,
A., Janiskova, M., Laloyaux, P., Lopez, P., Muñoz-Sabater, J., Peubey, C., Radu, R., Richardson, D., Thépaut, J.-N., Vitart,
F., Yang, X., Zsótér, E., and Zuo, H.: Operational global reanalysis: progress, future directions and synergies with NWP, 2018.

Hobbs, W., Massom, R., Stammerjohn, S., Reid, P., Williams, G. D., and Meier, W. N.: A Review of Recent Changes in
385 Southern Ocean Sea Ice, their Drivers and Forcings, *Global and Planetary Change*, 143, 228-250,
10.1016/j.gloplacha.2016.06.008, 2016.

- Holland, P. R., and Kwok, R.: Wind-driven trends in Antarctic sea-ice drift, *Nat Geosci*, 5, 872-875, 10.1038/NGEO1627, 2012.
- 390 Hosking, J. S., Orr, A., Marshall, G. J., Turner, J., and Phillips, T.: The Influence of the Amundsen-Bellinghousen Seas Low on the Climate of West Antarctica and Its Representation in Coupled Climate Model Simulations, *J Climate*, 26, 6633-6648, 10.1175/JCLI-D-12-00813.1, 2013.
- Jones, J. M., Gille, S. T., Goosse, H., Abram, N. J., Canziani, P. O., Charman, D. J., Clem, K. R., Crosta, X., de Lavergne, C., Eisenman, I., England, M. H., Fogt, R. L., Frankcombe, L. M., Marshall, G. J., Masson-Delmotte, V., Morrison, A. K., Orsi, A. J., Raphael, M. N., Renwick, J. A., Schneider, D. P., Simpkins, G. R., Steig, E. J., Stenni, B., Swingedouw, D., and Vance, 395 T. R.: Assessing recent trends in high-latitude Southern Hemisphere surface climate, *Nature Climate Change*, 6, 917-926, 10.1038/nclimate3103, 2016.
- King, J. C., and Harangozo, S. A.: Climate change in the western Antarctic Peninsula since 1945: observations and possible causes, *Ann Glaciol*, 27, 571-575, 10.3189/1998AoG27-1-571-575, 1998.
- Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., Endo, H., 400 Miyaoka, K., and Takahashi, K.: The JRA-55 Reanalysis: General Specifications and Basic Characteristics, *J Meteorol Soc Jpn*, 93, 5-48, 10.2151/jmsj.2015-001, 2015.
- Kusahara, K., Williams, G. D., Massom, R., Reid, P., and Hasumi, H.: Roles of wind stress and thermodynamic forcing in recent trends in Antarctic sea ice and Southern Ocean SST: An ocean-sea ice model study, *Global and Planetary Change*, 158, 103-118, 10.1016/j.gloplacha.2017.09.012, 2017.
- 405 Lenaerts, J. T. M., Lhermitte, S., Drews, R., Ligtenberg, S. R. M., Berger, S., Helm, V., Smeets, C. J. P. P., van den Broeke, M. R., van de Berg, W. J., van Meijgaard, E., Eijkelboom, M., Eisen, O., and Pattyn, F.: Meltwater produced by wind-albedo interaction stored in an East Antarctic ice shelf, *Nature Climate Change*, 7, 58-+, 10.1038/Nclimate3180, 2017.
- Marshall, G. J.: Trends in the southern annular mode from observations and reanalyses, *J Climate*, 16, 4134-4143, 10.1175/1520-0442(2003)016, 2003.
- 410 Marshall, G. J., Orr, A., and Turner, J.: A Predominant Reversal in the Relationship between the SAM and East Antarctic Temperatures during the Twenty-First Century, *J Climate*, 26, 5196-5204, 10.1175/JCLI-D-12-00671.1, 2013.
- Massom, R. A., Stammerjohn, S. E., Lefebvre, W., Harangozo, S. A., Adams, N., Scambos, T. A., Pook, M. J., and Fowler, C.: West Antarctic Peninsula sea ice in 2005: Extreme ice compaction and ice edge retreat due to strong anomaly with respect to climate, *J Geophys Res-Oceans*, 113, 10.1029/2007jc004239, 2008.
- 415 Massonnet, F., Mathiot, P., Fichet, T., Goosse, H., Beatty, C. K., Vancoppenolle, M., and Lavergne, T.: A model reconstruction of the Antarctic sea ice thickness and volume changes over 1980-2008 using data assimilation, *Ocean Model*, 64, 67-75, 10.1016/j.ocemod.2013.01.003, 2013.
- Meier, W. N., Peng, G., Scott, D. J., and Savoie, M.: Verification of a new NOAA/NSIDC passive microwave sea-ice concentration climate record, *Polar Res*, 33, 21004, 10.3402/polar.v33.21004, 2014.

- 420 Meneghini, B., Simmonds, I., and Smith, I. N.: Association between Australian rainfall and the Southern Annular Mode, *Int J Climatol*, 27, 109-121, 10.1002/joc.1370, 2007.
- Nicolas, J. P., and Bromwich, D. H.: New Reconstruction of Antarctic Near-Surface Temperatures: Multidecadal Trends and Reliability of Global Reanalyses, *J Climate*, 27, 8070-8093, 10.1175/JCLI-D-13-00733.1, 2014.
- Paolo, F. S., Fricker, H. A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 327-331, 10.1126/science.aaa0940, 2015.
- 425 Paolo, F. S., Padman, L., Fricker, H. A., Adusumilli, S., Howard, S., and Siegfried, M. R.: Response of Pacific-sector Antarctic ice shelves to the El Niño/Southern Oscillation, *Nat Geosci*, 11, 121-126, 10.1038/s41561-017-0033-0, 2018.
- Parkinson, C. L., and Cavalieri, D. J.: Antarctic sea ice variability and trends, 1979-2010, *Cryosphere*, 6, 871-880, 10.5194/Tc-6-871-2012, 2012.
- 430 Parkinson, C. L.: A 40-y record reveals gradual Antarctic sea ice increases followed by decreases at rates far exceeding the rates seen in the Arctic, *Proceedings of the National Academy of Sciences*, 10.1073/pnas.1906556116, 2019.
- Poli, P., Hersbach, H., Dee, D. P., Berrisford, P., Simmons, A. J., Vitart, F., Laloyaux, P., Tan, D. G. H., Peubey, C., Thépaut, J.-N., Trémolet, Y., Hölm, E. V., Bonavita, M., Isaksen, I., and Fisher, M.: ERA-20C: An Atmospheric Reanalysis of the Twentieth Century, *J Climate*, 29, 4083-4097, 10.1175/JCLI-D-15-0556.1, 2016.
- 435 Pritchard, H. D., Ligtenberg, S. R., Fricker, H. A., Vaughan, D. G., van den Broeke, M. R., and Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484, 502-505, 10.1038/nature10968, 2012.
- Raphael, M. N., and Hobbs, W.: The influence of the large-scale atmospheric circulation on Antarctic sea ice during ice advance and retreat seasons, *Geophysical Research Letters*, 41, 5037-5045, 10.1002/2014gl060365, 2014.
- Raphael, M. N., Marshall, G. J., Turner, J., Fogt, R. L., Schneider, D., Dixon, D. A., Hosking, J. S., Jones, J. M., and Hobbs, W. R.: The Amundsen Sea Low: Variability, Change, and Impact on Antarctic Climate, *B Am Meteorol Soc*, 97, 111-121, 10.1175/bams-d-14-00018.1, 2016.
- Saha, S., Moorthi, S., Pan, H. L., Wu, X. R., Wang, J. D., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H. X., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y. T., Chuang, H. Y., Juang, H. M. H., Sela, J., Iredell, M., Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H. L., Yang, R. Q., Lord, S., Van den Dool, H., 440 Kumar, A., Wang, W. Q., Long, C., Chelliah, M., Xue, Y., Huang, B. Y., Schemm, J. K., Ebisuzaki, W., Lin, R., Xie, P. P., Chen, M. Y., Zhou, S. T., Higgins, W., Zou, C. Z., Liu, Q. H., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G., and Goldberg, M.: The Ncep Climate Forecast System Reanalysis, *B Am Meteorol Soc*, 91, 1015-1057, 10.1175/2010bams3001.1, 2010.
- Schroeter, S., Hobbs, W., and Bindoff, N. L.: Interactions between Antarctic sea ice and large-scale atmospheric modes in 450 CMIP5 models, *Cryosphere*, 11, 789-803, 10.5194/tc-11-789-2017, 2017.
- Schroeter, S., Hobbs, W., Bindoff, N. L., Massom, R., and Matear, R.: Drivers of Antarctic Sea Ice Volume Change in CMIP5 Models, *J Geophys Res-Oceans*, 123, 7914-7938, 10.1029/2018jc014177, 2018.

- Simmons, A. J., Berrisford, P., Dee, D. P., Hersbach, H., Hirahara, S., and Thépaut, J. N.: A reassessment of temperature variations and trends from global reanalyses and monthly surface climatological datasets, *Q J Roy Meteor Soc*, 143, 101-119, 455 10.1002/qj.2949, 2017.
- Slivinski, L. C., Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Giese, B. S., McColl, C., Allan, R., Yin, X., Vose, R., Titchner, H., Kennedy, J., Spencer, L. J., Ashcroft, L., Brönnimann, S., Brunet, M., Camuffo, D., Cornes, R., Cram, T. A., Crouthamel, R., Domínguez-Castro, F., Freeman, J. E., Gergis, J., Hawkins, E., Jones, P. D., Jourdain, S., Kaplan, A., Kubota, H., Blancq, F. L., Lee, T.-C., Lorrey, A., Luterbacher, J., Maugeri, M., Mock, C. J., Moore, G. W. K., Przybylak, R., 460 Pudmenzky, C., Reason, C., Slonosky, V. C., Smith, C. A., Tinz, B., Trewin, B., Valente, M. A., Wang, X. L., Wilkinson, C., Wood, K., and Wyszynski, P.: Towards a more reliable historical reanalysis: Improvements for version 3 of the Twentieth Century Reanalysis system, *Q J Roy Meteor Soc*, 145, 2876-2908, 10.1002/qj.3598, 2019.
- Stammerjohn, S., Massom, R., Rind, D., and Martinson, D.: Regions of rapid sea ice change: An inter-hemispheric seasonal comparison, *Geophysical Research Letters*, 39, 10.1029/2012gl050874, 2012.
- 465 Steig, E. J., Schneider, D. P., Rutherford, S. D., Mann, M. E., Comiso, J. C., and Shindell, D. T.: Warming of the Antarctic ice-sheet surface since the 1957 International Geophysical Year (vol 457, pg 459, 2009), *Nature*, 460, 766-766, Doi 10.1038/Nature08286, 2009.
- Steig, E. J., and Orsi, A. J.: The heat is on in Antarctica, *Nat Geosci*, 6, 87-88, 10.1038/ngeo1717, 2013.
- Swart, N. C., and Fyfe, J. C.: Observed and simulated changes in the Southern Hemisphere surface westerly wind-stress, 470 *Geophysical Research Letters*, 39, Art16711 10.1029/2012gl052810, 2012.
- Turner, J., Phillips, T., Hosking, J. S., Marshall, G. J., and Orr, A.: The Amundsen Sea low, *Int J Climatol*, 33, 1818-1829, 10.1002/joc.3558, 2013.
- Turner, J., Barrand, N. E., Bracegirdle, T. J., Convey, P., Hodgson, D. A., Jarvis, M., Jenkins, A., Marshall, G., Meredith, M. 475 P., Roscoe, H., Shanklin, J., French, J., Goosse, H., Guglielmin, M., Gutt, J., Jacobs, S., Kennicutt, M. C., Masson-Delmotte, V., Mayewski, P., Navarro, F., Robinson, S., Scambos, T., Sparrow, M., Summerhayes, C., Speer, K., and Klepikov, A.: Antarctic climate change and the environment: an update, *Polar Record*, 50, 237-259, 10.1017/S0032247413000296, 2014.
- Wang, Y., Zhou, D., Bunde, A., and Havlin, S.: Testing reanalysis data sets in Antarctica: Trends, persistence properties, and trend significance, *Journal of Geophysical Research: Atmospheres*, 121, 839-812,855, 10.1002/2016JD024864, 2016.
- 480 Zhang, J. L.: Increasing Antarctic sea ice under warming atmospheric and oceanic conditions, *J Climate*, 20, 2515-2529, 10.1175/Jcli4136.1, 2007

	Description	Citation	Reanalysis period	Spatial Resolution (lat x lon)	Algorithm
NCEP2	NCEP-DOE AMIP II Reanalysis	Kobayashi <i>et al</i> (2015)	1979-present	2.5° x 2.5°	3D-VAR
CFSR	NCEP Climate Forecast System Reanalysis	Saha <i>et al</i> (2010)	1979-present	0.5° x 0.5°	3D-VAR
MERRA2	Modern Era Retrospective Analysis for Research and Applications, version 2	Gelaro <i>et al</i> (2017)	1980 -present	0.5° x 0.625°	3D-VAR
20CRv3	National Oceanic and Atmospheric Administration - Cooperative Institute for Research in Environmental Sciences 20 th Century Reanalysis version 3	Slivinski <i>et al</i> (2019)	1836-2015	1° x 1°	Ensemble Kalman Filter
ERA5	European Centre for Medium Range Weather Forecasting (ECMWF) Reanalysis version 5	Hersbach <i>et al</i> (2018)	1979-present	0.25° x 0.25°	4D-VAR
ERA-20C	ECMWF 20 th Century Reanalysis	Poli <i>et al</i> (2016)	1900 -2010	0.25° x 0.25°	4D-VAR
ERA-int	ECMWF Interim Reanalysis	Dee <i>et al</i> (2011)	1979-2019	0.75° x 0.75°	4D-VAR
JRA55	Japanese 55-year Reanalysis	Kobayashi <i>et al</i> (2015)	1958-present	1.25° x 1.25°	4D-VAR


Formatted: Font: Not Bold

Formatted: Font: Bold

Deleted: Bold years in the reanalysis periods indicate the first and final year of the period for which output is available for all products.

485

Table 1: Summary of reanalysis products used in this study.



495 **Figure 1: 1980-2010 trends in annual-mean SAT (shading: °C/year) and MSLP (contour lines: positive trends in blue, negative trends in magenta, with contour spacing = 2.5 Pa/year), for eight individual reanalyses (refer to Table 1 for details). Trend patterns for each month are shown in the Supplemental Material as Figures S1-S12.**






Figure 2: Panels a-d show observed 1980-2010 Antarctic SIC trends by season (decade-1). Hatching indicates trends that are statistically-significant at the 0.05 significance level, and magenta lines show the climatological sea ice edge (defined by the 15% SIC isoline). Seasons are defined by total sea ice area (SIA) growth and melt ($dSIA/dt$), shown on panel e. Line colors in 2e follow the [SPARC Reanalysis Intercomparison Project \(S-RIP\)](#) standard.



505 **Figure 3: Uncentered pattern correlations by month between 1980-2010 SIC and SAT trends: a) correlations between the observed SIC trend and each reanalysis' SIC; b) correlations between each reanalysis SAT trend that reanalysis' SIC trend; c) correlation between reanalysis SAT trend and observed SIC trend. Differences between each reanalysis SIC and the Goddard merged SIC product for each month are shown in the Supplemental Material as Figures S13-S24.**



510

Figure 4: Net heat flux averaged over sea ice zone (x-axis: Wm⁻², where positive indicates net flux from atmosphere to ocean), against the pattern correlation between reanalysis SIC and Sat (i.e. values plotted in Figure 3b). Data points are colored by month with different markers for each reanalysis. Colored lines show the best-fit line for each month estimated by Ordinary Least Squares regression.

515




Figure 5: Panels a-h) 1980-2010 December-January mean reanalysis SAT trends over the climatological sea ice zone ($^{\circ}\text{C}/\text{year}$); only trends that are statistically-significant at the 90% level are shown. Hatched regions show where the sign of statistically-significant reanalysis SIC and SAT trends are the same (i.e. the unexpected result of both a warming (cooling) and an increase (decrease) in sea ice cover). The magenta line shows climatological sea ice edge.