

Large-scale synoptic drivers of co-occurring summertime ozone and PM_{2.5} pollution in eastern China

Lian Zong¹, Yuanjian Yang^{1,*}, Meng Gao², Hong Wang¹, Peng Wang³, Hongliang Zhang⁴, Linlin Wang⁵, Guicai Ning⁶, Chao Liu¹, Yubin Li¹, Zhiqiu Gao^{1,5}

5

1. School of Atmospheric Physics, Nanjing University of Information Science & Technology, Nanjing, China

2. Department of Geography, Hong Kong Baptist University, Hong Kong SAR, China

3. Policy Research Center for Environment and Economy, Ministry of Ecology and Environment of the People's Republic of China, Beijing, China

10

4. Department of environmental science and engineering, Fudan University, China

5. State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry (LAPC), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

6. Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong, China

15

* Correspondence to: Dr./Prof. Y. Yang (yyj1985@nuist.edu.cn)

Abstract

Surface ozone (O₃) pollution during summer (June–August) over eastern China has become more severe in recent years, resulting in a co-occurrence of surface O₃ and PM_{2.5} (particulate matter with aerodynamic diameter ≤ 2.5 μm in the air) pollution. However, the mechanisms regarding how the synoptic circulation pattern might influence this compound pollution remain unclear. In this study, we applied the T-mode principal component analysis (T-PCA) method to objectively classify the occurrence of four synoptic weather patterns (SWPs) over eastern China, based on the geopotential heights at 500 hPa during summer (2015–2018). These four SWPs over eastern China were closely related to the western Pacific subtropical high (WPSH), exhibiting significant intraseasonal and interannual variations. Based on ground-level air quality observations, remarkable spatial and temporal disparities of surface O₃ and PM_{2.5} pollution were also found under the four SWPs. In particular, there were two SWPs that were sensitive to compound pollution (Type 1 and Type 2). Type 1 was characterized by a stable WPSH ridge with its axis at about 22°N and the rain belt located in the south of the Yangtze River Delta (YRD); and Type 2 also exhibited WPSH dominance (ridge axis at $\sim 25^\circ\text{N}$), but with the rain belt (over the YRD) at a higher latitude compared to Type 1. In general, SWPs have played an important role as driving factors of surface O₃–PM_{2.5} compound pollution in a regional context. Our findings demonstrate the important role played by SWPs in driving regional surface O₃–PM_{2.5} compound pollution, in addition to the large quantities of emissions, and may also provide insights into the regional co-occurring high levels of both PM_{2.5} and O₃ via the effects of certain meteorological factors.

Keywords: synoptic weather pattern, ozone, PM_{2.5}, compound pollution, western Pacific subtropical high (WPSH)

1. Introduction

In recent years, China has been experiencing serious air pollution problems owing to its enormous emissions of polluting gases [e.g., sulfur dioxide, nitrogen dioxide (NO₂), etc.] and aerosol particulates [e.g., particulate matter with aerodynamic diameter ≤ 2.5 (10) μm in the air PM_{2.5} (PM₁₀), etc.] associated with its rapid economic development, industrialization and

urbanization, together with certain unfavorable meteorological conditions (Wang and Chen, 2016; Zhang et al., 2014; Zhang et al., 2016). In Particular, atmospheric compound pollution has become
50 serious (Li et al., 2019; Saikawa et al., 2017; Zhang et al., 2019a), especially for the economically developed and densely populated eastern urban agglomerations of China, such as the Beijing–Tianjin–Hebei (BTH), Yangtze River Delta (YRD) and Pearl River Delta (PRD) regions (Cai et al., 2017; Du et al., 2019; Ji et al., 2018; Li et al., 2020), exerting a severe threat in terms of public health, economy and society (Chen et al., 2019; Cohen et al., 2017; Day et al., 2017; Yim et al.,
55 2019).

In general, a significant diurnal variation of PM_{2.5} pollution has been observed, possibly due to obvious local emissions caused by industrial production and human activities related to daily living (Amil et al., 2016; Liu et al., 2019a). In particular, the pollution level tends to be higher during the morning and evening of a normal weekday, with a weakened effect found in the
60 afternoon, possibly caused by the co-effects of the boundary layer structure and anthropogenic emissions. There has also been a seasonal variation of PM_{2.5} pollution detected across China, indicating a higher level of pollution in winter than summer (Ye et al., 2018; Zhang and Cao, 2015). The PM_{2.5} level in China showed a steady increase from 2004 to 2007, and has since stabilized (Ma et al., 2016); however, there are still frequent PM_{2.5} pollution events in autumn and
65 winter (Song et al., 2017; Yang et al., 2018; Ye et al., 2018; Zhang et al., 2014). In the past few years, the PM_{2.5} concentration in China has decreased significantly as a result of measures introduced across the country that have reduced multi-pollutant emissions, adjusted the energy structure, and increased the supply of clean energy (Gui et al., 2019; Yang et al., 2020; Zhang et al., 2019b; Zhang et al., 2020). While PM_{2.5} is still one of the dominant air pollutants across China,
70 surface O₃ pollution in summer has also gradually risen to prominence. Several studies have even indicated that O₃ might replace PM_{2.5} as the primary air pollutant during summer (Li et al., 2019), which has caught the attention of researchers in recent years. For instance, Sun et al. (2016) showed that the observed summertime O₃ at Mt. Tai increased significantly by 1.7 ppbv yr⁻¹ for the month of June and by 2.1 ppbv yr⁻¹ for the months of July–August during the period 2003–
75 2015. Furthermore, an increase in the maximum daily 8-h average concentration of O₃ (MDA8 O₃) at an annual-average rate of 4.6%, was reported by Fan et al. (2020), albeit with a decrease in the

frequency of PM_{2.5} pollution.

The modulations of atmospheric circulation systems often lead to changes in meteorological elements, and thereby also affect the processes of pollutant formation, transmission and diffusion.

80 Furthermore, many studies have indicated that PM_{2.5} and O₃ pollution are strongly correlated with local meteorological factors such as temperature, relative humidity (RH) and wind speed (WS) (Huang et al., 2016; Miao et al., 2015; Shu et al., 2019; Tai et al., 2010). Miao et al. (2015) suggested that a low boundary layer height (BLH) and stable atmosphere would be an unfavorable condition for the dispersion of winter aerosol pollution over the BTH region. Zhang et al. (2017) 85 found that the majority of O₃ extremes occurred with daily maximum temperature (T_{max}) of between 300 K and 320 K, minimum a RH (RH_{min}) of less than 40%, and a minimum WS of less than 3 m s⁻¹, through an analysis of extreme O₃ and PM_{2.5} events from historical data (30 years for O₃ and 10 years for PM_{2.5}) in the United States. Furthermore, the number of annual extreme PM_{2.5} days was highly positively correlated with the extreme RH_{min}/T_{max} days, and the correlation 90 coefficient between PM_{2.5} and RH_{min} (T_{max}) was highest in urban and suburban (rural) regions. Shi et al. (2020) studied the sensitivity of O₃-8h (O₃ 8-hour moving average) and PM_{2.5} associated with meteorological parameters. Their study focused on the air pollution and meteorological conditions between January and July, 2013, with the results showing that temperature could have had the greatest impact on the daily maximum O₃-8h, while the PM_{2.5} sensitivities were negatively 95 (positively) correlated with temperature, WS, and BLH (absolute humidity) in most regions of China. Miao et al. (2015) showed that RH was high when aerosol pollution occurred in the BTH region. However, O₃ pollution in China is more frequent in summer, and the warm and humid flow brought by the East Asian summer monsoon (EASM) induces a hot and humid condition over the summer. Zhao et al. (2019) investigated the RH of O₃ pollution in Shijiazhuang between 15 June 100 and 14 July 2016, and found that the O₃ concentration was higher at moderate humidity (RH average during daytime from 10:00 to 17:00 LT was 40%–50%). Recently, Han et al. (2020) assessed the impacts of local and synoptic meteorological factors on the daily variability of surface O₃ over eastern China. Their study revealed that the meteorological factors could explain ~46% of the daily variations in summer surface O₃. In particular, synoptic factors contributed to 105 ~37% of the overall effects associated with the meteorological factors. Furthermore, six

predominant synoptic weather patterns (SWPs) were identified by the self-organizing map, and related results indicated a weak cyclonic system and southward prevailing wind induced the positive O₃ anomalies over eastern China. The abovementioned studies indicates that the variations of meteorological factors play a non-negligible role in air pollution. Therefore, classification of air pollution according to meteorological circulation has become particularly important, not least because of its worth when applied to air quality monitoring, forecasting and evaluation (Liu et al., 2019b; Ning et al., 2019; Yang et al., 2018; Zheng et al., 2015).

Since 1990s, it has become possible to objectively classify atmospheric circulation conditions using weather data such as geopotential height (GH), sea level pressure, WS and temperature, thus allowing the physical mechanism of extreme weather to be better understood and analyzed. Compared with subjective weather classification, the objective approach has been widely used in air pollution research (Miao et al., 2017, 2019; Ning et al., 2018). Miao et al. (2019), based on the daily 900 hPa GH fields during winter in Beijing, identified seven synoptic patterns using an objective approach, and found that the weak northwesterly prevailing winds and strong elevated thermal inversion layer, along with the local emissions of aerosols, play a decisive role in the formation of heavy pollution in Beijing. The authors also noted that southerly prevailing winds can transport pollutants emitted from southern cities to Beijing. Zheng et al. (2015) studied the relationship between regional pollution and the patterns of large-scale atmospheric circulation over eastern China in October from 2001 to 2010 and identified six pollution types and three clean types. Specifically, weather patterns such as a uniform surface pressure field in eastern China or a steady straight westerly in the middle troposphere, particularly when at the rear of an anticyclone at 850 hPa, were found to be typically responsible for heavy pollution events. Many studies have suggested a modulating effect of the East Asian summer monsoon (EASM) and western Pacific subtropical high (WPSH) on air quality over China (Li et al., 2018; Yin et al., 2019; Zhao et al., 2010). In particular, Li et al. (2018) applied RegCM4-CHEM to analyze the differences in ozone between three strong and weak monsoon years, and found that the concentrations of O₃ over the central and eastern part of China were higher in strong EASM years than in weak EASM years. The anomalous high-pressure system at 500 hPa, associated with downward dry, hot air and intense solar radiation can enhance the photochemical reactions to elevate the production of

tropospheric O₃ (Gong and Liao, 2019; Yin et al., 2019). Furthermore, Zhao & Wang (2017) and Yin et al. (2019) noted that the positive GH anomalies at high latitudes tended to significantly weaken the cold-air advection from the north resulting in locally high temperatures near the surface in northern China, while the WPSH could transport sufficient water vapor to the YRD region leading to a decrease in surface O₃. In addition, different subregions can exhibit various distributions of pollutants, even with identical emission scenarios (Li et al., 2019; Saikawa et al., 2017; Zhang et al., 2019a). Also, it is still unclear how the distribution of pollution responds locally to large-scale atmospheric circulation patterns. Due to the variability of local meteorological conditions under various synoptic weather types and the modulation of large-scale movement of the WPSH (Li et al., 2018; Wang et al., 2019; Yang et al., 2021; Zhao and Wang, 2017), the causes and consequences of meteorological factors for the formation of compound O₃–PM_{2.5} pollution could be complex. Overall, the mechanism by which the SWP modulates the characteristics of O₃–PM_{2.5} compound pollution has yet to be comprehensively described.

In this study, the SWPs corresponding to the co-occurrence of O₃ and PM_{2.5} pollution during summertime were analyzed, with a focus on eastern China (104°–135°E, 17°–53°N). Then, the synoptic causes of O₃–PM_{2.5} compound pollution, as well as O₃-only pollution, from the perspective of the objective classification of atmospheric circulation patterns, were revealed. The findings are expected to provide a scientific reference for the monitoring, forecasting and evaluation of summertime air pollution in eastern China.

2. Data and methods

The air quality data, including PM_{2.5}, NO₂, O₃, and O₃-8h, are from the national 24-h continuous air quality observation published by the China Environmental Monitoring Station (<http://www.cnemc.cn/>). Summer hourly data (2015–2018) for 1174 stations were retrieved from an observational network in eastern China (104°–135°E, 17°–53°N), which include the more prominent pollution areas in the eastern urban agglomeration, such as the BTH (113.5°–119.8°E, 36°–42.6°N), YRD (115.3°–122.6°E, 27.2°–34.5°N), PRD (112.5°–113.7°E, 21.3°–23.1°N), Guanzhong Plain [GZP (104.6°–112.2°E, 33.3°–36.8°N)], and Northeast Megalopolis [NEM (121.2°–131.0°E, 39.8°–47.3°N)] regions (the specific locations of stations and urban

agglomerations are shown in Fig. 1a). Surface meteorological data, such as Tmax, precipitation,
165 WS and RH from 611 meteorological observation stations, along with sounding data at 1400
Beijing time (BJT) from 64 stations and at 0800 BJT and 2000 BJT from 77 stations, in eastern
China, were obtained from the China National Meteorological Information Center of the China
Meteorological Administration (<http://data.cma.cn/site/index.html>). The BLH was calculated
according to the method given by Seidel et al. (2012) and Guo et al. (2016, 2019) (the detailed
170 method can be seen in the supplementary material), and the FLWD [frequency of light wind (< 2
 m s^{-1}) days, which can be defined as the ratio between the number of days with average daily WS
lower than 2 m s^{-1} and the total days of each pattern], precipitation frequency (PF, which can be
defined as the ratio of the number of rainy days to total days under each pattern), and MDA8 O_3
were also counted.

175 Additionally, for synoptic analysis of particulate matter and O_3 pollution in summer, we use
the GH field at 500 hPa, wind, and specific humidity field at 850 hPa from the NCEP/NCAR
(National Centers for Environmental Prediction/National Center for Atmospheric Research) daily
reanalysis dataset on a $2.5^\circ \times 2.5^\circ$ latitude/longitude grid during the study period. For further
analysis of the modulation of the co-occurrence of O_3 - $\text{PM}_{2.5}$ pollution by the boundary layer
180 structure in some local areas, we also used the BLH, uv-wind, vertical velocity, RH and
temperature fields of the fifth generation European Centre for Medium-Range Weather Forecasts
reanalysis (ERA5), which has a high spatiotemporal resolution ($0.25^\circ \times 0.25^\circ$, hourly;
<https://cds.climate.copernicus.eu/cdsapp#!/home>).

T-mode principal component analysis (T-PCA) is an objective mathematical computer-based
185 method that can be used to classify the synoptic circulation patterns of regional gridded data in the
troposphere at the lower level. Indeed, it is commonly regarded as the most promising weather
pattern classification method at present (Huth et al., 2008). Moreover, this approach has been
widely used in the studies of aerosols and O_3 pollution-related atmospheric circulation in China
(Miao et al., 2017, 2019; Ning et al., 2018, 2019). The T-PCA analysis module of the COST733
190 software (<http://cost733.met.no/>) developed by the European Scientific and Technical Research
Cooperation, was used to classify the synoptic circulation pattern based on the 500 hPa GH field.
The cost733class program is a FORTRAN software package consisting of several modules for

classification, evaluation and comparison of weather and circulation patterns. First, the weather data are spatially standardized and split into 10 subsets by T-PCA. Then the principal components (PCs) of weather information are estimated by applying singular value decomposition, and the PC score for each subset can be calculated after oblique rotation. Finally, the resultant subset with the highest sum will be selected by comparing 10 subsets according to contingency tables, and its types can be output as well (Miao et al., 2017; Philipp et al., 2014). To assess the performance of synoptic classification and determine the number of classes, the explained cluster variance (ECV) was selected in this study (Hoffmann & Schl Nzen, 2013; Ning et al., 2019; Philipp et al., 2014). Detailed information about the ECV method is provided in the supplementary material.

Based on the Ambient Air Quality Standards (GB3095-2012) issued by the Ministry of Ecology and Environment of the People's Republic of China, O₃ (PM_{2.5}) pollution occurs when the MDA8 O₃ (PM_{2.5} 24-h) concentration exceeds 160 (75) µg m⁻³. For a region, when haze occurs at more than 50% of the observed sites, the day can be defined as a haze day (Chen and Wang, 2015). In this study, we characterized regional pollution days as occurring when the average values of more than 50% of sites in this region exceeded the aforementioned thresholds. The specific standard limits of each pollution level are according to their concentration limits based on the Technical Regulation on Ambient Air Quality Index (on trial) (HJ633-2012) issued by the Ministry of Ecology and Environment of the People's Republic of China (Table S1).

Finally, in order to make it clear in the analysis of different weather types of O₃ and PM_{2.5} concentration change, we calculated the average distribution of O₃ and PM_{2.5}, as well as the meteorological conditions for each type, and the anomalous distribution of these variables (i.e., the average of O₃ and PM_{2.5} and the average of the meteorological conditions under the respective patterns minus the average during summertime of 2015–2018) were given as well. The statistical significance was tested with a 0.05 confidence level via analysis of variance, which enabled us to distinguish the significant differences of spatial distribution characteristics between O₃ and PM_{2.5} pollution under four SWPs.

3. Results

3.1 Spatial and temporal distribution of O₃ and PM_{2.5} during summer 2015–2018

Figure 1 shows the summer averaged MDA8 O₃ and PM_{2.5} concentrations at 1174 stations in the eastern region of China for the period of 2015–2018. Among these stations, the MDA8 O₃ concentration at most stations (795/1174) exceeded 100 µg m⁻³, of which 45 sites exceeded 160 µg m⁻³. The highest O₃ pollution was found in Zibo, Shandong, with a value of 181.5 µg m⁻³. The averaged PM_{2.5} at most sites (844/1174) was below 35 µg m⁻³, while it reached 62.6 µg m⁻³ in Handan, Hebei Province. On the whole, the MDA8 O₃ and PM_{2.5} in the BTH region and its surrounding areas was significantly higher than in other regions; and besides, the level of O₃ in some urban clusters, such as the PRD, YRD, GZP and NEM regions, was particularly higher than that of the surroundings, and thus, we focus on analyzing these key areas later in the paper.

Figures 2 and 3 respectively show the daily variations in pollution levels of O₃ and PM_{2.5}. In recent years, the reduced visibility of haze days has weakened the solar radiation reaching the ground and inhibited photochemical reactions from generating O₃ (Li et al., 2019; Zhang et al., 2015). As a result, the concentration of O₃ has continued to increase with the mitigation of PM_{2.5} pollution. During the study period, the number of days of O₃ pollution in the BTH, YRD, PRD, GZP, and NEM regions was 254, 133, 84, 165 and 96 respectively, while the number of days of PM_{2.5} pollution was only 93, 8, 0, 2 and 1, of which compound pollution occurred on 76, 7, 0, 2, and 0 days according to Chinese standards (the asterisks in Fig. 3 indicate the compound pollution events). China has implemented strict policies for emission control, and the effects of these policies have been remarkable. However, despite a decrease in PM_{2.5} in the last five years, there has also been an increase in O₃ pollution over China (Fan et al., 2020; Sun et al., 2016), “double-high” pollution reported on the weather scale has reduced. As the limit of PM_{2.5} concentration for pollution control is relatively loose in China, previous studies have tended to refer to the interim target 1 (IT-1) of the World Health Organization (WHO) as the standard threshold. Our study pushed forward to the next stage, i.e., we used the WHO’s IT-2 threshold (24-h average concentration of PM_{2.5} of 35 µg m⁻³) as our target limit to count the number of compound pollution days across each region. Based on this target, the number of pollution days for the five urban clusters was 194, 52, 16, 47, and 20, respectively (Fig. 3). These results indicate that, despite PM_{2.5} reductions, compound pollution events remain serious and deserve of public attention. Overall, the O₃ and PM_{2.5} concentrations in eastern China exhibit distinct intraseasonal

and interannual variations, indicating that, aside from the changes in emission sources (because it is considered that inter-seasonal and short-term changes in emission sources are not significant), they may also be regulated by meteorological conditions, which is further analyzed below.

3.2 Objective classification of large-scale synoptic circulation patterns in summer

To analyze the effect of meteorological conditions on the changes in O_3 and $PM_{2.5}$ concentrations, it is necessary to statistically analyze the large-scale weather circulation situation in summer. Existing studies have shown that the WPSH (500 hPa GH field with obvious anticyclonic characteristics, and downward flow around the center) in summer prominently regulates the weather and climate of East Asia (Lu, 2002), owing to its varying location, shape and intensity (Ding, 1994). Low-level southerly monsoonal flow forming at the periphery of an anomalously enhanced WPSH, along with the transportation of warm and humid air from the ocean to East Asia, might also be responsible for the asymmetric spatial distribution of ground-level O_3 [i.e., a decrease in southern China but an increase over northern China (Zhao & Wang, 2017)].

Therefore, we used the T-PCA method to objectively classify the weather circulation of the 500-hPa GH field in the summers of 2015–2018, from which we ultimately obtained four SWPs related to the movement and development of the WPSH (Wang et al., 2019; Yang et al., 2021). The western ridge point and northern boundary of the WPSH at 500 hPa in Type 1 were located at around $120^\circ E$ and $30^\circ N$, respectively (Fig. 4a and Table S2). The southwestern flow of this WPSH was able to transport water vapor to the YRD region, resulting in a southwestward prevailing wind across the YRD region and westward flow from the north of the WPSH forming a convergence area at 850 hPa. These conditions were also associated with high temperature and humidity during the summer in the Meiyu season, which is a climate phenomenon characterized by continuous cloudy and rainy days that generally occurs during June and July every year across the middle and lower reaches of the Yangtze river, Taiwan in China, central/southern Japan, and southern Korea. For Type 2, the westerly trough was able to deepen as the WPSH shifted northwards slightly from Type 1 or retreated southeast from Type 3 (Fig. 4b). The southwest wind from the South China Sea might have combined with the southerly wind in the eastern periphery of the WPSH. As a result,

southerly winds prevailed across southeastern China, while northern China was mainly controlled by the westerly trough. In comparison to Type 2, Type 3 was characterized by the boundary of the WPSH being at a higher latitude, with a westward extension (Fig. 4c), disintegrating a closed high-pressure monomer along the eastern coast of China, and the main body of the WPSH remained over the ocean (Figs. 4c and S4). This led to a condition that was completely controlled by the monomer of the WPSH over the YRD region, resulting in hot and dry weather at the end of the rainy season at the beginning of mid-summer. Figure. 4d indicates that the location of the WPSH monomer was more to the west and north compared with under the other SWPs, thus controlling northern China for a long time; the western ridge point was located at around 95°E and the northern boundary at around 40°N.

Figure 5 presents the daily and annual variations of the SWPs in the summers of 2015–2018. Usually, the advancement of the WPSH in eastern China occurs in June and July, while its gradual withdrawal occurs mainly in August; and in this respect, Type 1 and Type 2 represented normal WPSH characteristics during early and late summer. Type 3 and Type 4, however, reflected a split of the WPSH, which mainly occurred in late summer. Consequently, there were 167, 117, 52 and 32 days for Type 1, Type 2, Type 3 and Type 4 over the study period, respectively. Since the WPSH's movement is generally affected by the weather phenomena of its surrounding climatic systems (such as typhoons, the Tibetan high, etc.) (Ge et al., 2019; Liu and You, 2020; Shu et al., 2016; Wang et al., 2019), it may have resulted in a short-term southward retreat during the advancement of the WPSH (e.g., around 10 August 2018) and a short-term northward advancement during its process of retreat (e.g., 21 and 29 August 2016). For instance, tropical storm NEPARTAK generated at 0000 UTC (0800 BJT) 3 July 2016 over the western North Pacific and upgraded to a super typhoon at 1200 UTC (2000 BJT) 5 July 2016 (Fig. S5; see also Su et al., 2017). Due to the rapid movement of NEPARTAK to the northwest, the WPSH quickly decomposed a monomer and moved north. With the strengthening and landing of the typhoon, the monomer gradually collapsed. The SWP also underwent a transition from Type 2 to Type 4, and then to Type 1 (Figs. 4 and S5). In general, the WPSH is able to represent the evidence of intra-seasonal and interannual changes over China, which will inevitably modulate the weather, as well as climatic and environmental changes in eastern China.

3.3 O₃ and PM_{2.5} pollution characteristics under the four SWPs

3.3.1 Spatial characteristics

We calculated the averaged (Fig. S6) and anomalous (Fig. 6) spatial distributions of the MDA8 of O₃ and PM_{2.5} under the four SWPs. The O₃ concentration was relatively high in the area north of the Yangtze River under Type 1, and the high values of MDA8 O₃ were mainly concentrated in the North China Plain (NCP) region, with a total of 100 stations surpassing 160 µg m⁻³. Type 2 O₃ pollution was slightly weaker than that for Type 1, and the MDA8 O₃ at the 72 sites exceeded 160 µg m⁻³. The O₃ high-value areas lay mainly in the NCP, GZP and YRD regions under Type 4, and there were 37 stations with concentrations larger than 160 µg m⁻³. Of the four SWPs, the lowest overall MDA8 O₃ occurred under Type 3, with only one site exceeding 160 µg m⁻³ (Figs. S6a-d). It was also found that the regions that experienced significant positive deviations of MDA8 O₃ from the summer mean were as follows: the BTH, YRD and NEM regions under Type 1; the BTH and GZP regions under Type 2; the central of the YRD and PRD regions under Type 3, and the YRD, GZP and PRD regions under Type 4 (Figs. 6a-d).

Analogously, Figs. 6e-h shows the anomaly and significance of difference in PM_{2.5} under the four weather types, presented as positive anomalies in the south of the BTH and YRD regions under Type 1, in the BTH, GZP and PRD regions under Type 2, and in the GZP and PRD regions under Type 4. Due to the obvious seasonal variations of PM_{2.5} concentration (higher in winter and lower in summer) (Liu et al., 2019a; Miao et al., 2015), no site exceeded 75 µg m⁻³ for the averaged PM_{2.5} concentration. Even so, the level of PM_{2.5} in the BTH region was still significantly higher under the four types than that for other urban agglomerations (Figs. S6e-h).

3.3.2 Pollution pattern differences in key areas

Air pollution is principally found in dense urban areas such as the BTH and YRD regions (Gui et al., 2019; Han et al., 2019), so we took the BTH, PRD, YRD, GZP and NEM regions in the eastern region as key areas, counted the daily anomalies and average variation of O₃ and PM_{2.5} in each key region under the different weather patterns (Figs. 7 and S7), and calculated the over-limit ratio in those key regions via the ‘stations × days’ statistics (see Table 1). The diurnal variation of

O₃ was more obvious, peaking at about 1500 BJT, while contrasting diurnal variations of PM_{2.5} were found for different regions. According to Fig. 7 and Table 1, the following characteristics could be identified for different urban clusters: (1) in the BTH region, the O₃ concentrations of Type 1 and Type 2 were relatively high, with their over-limit rates reaching 47.1% and 54.2%, and the PM_{2.5} pollution rates reaching 18.8% and 16.3%, respectively; (2) in the PRD region, the over-limit rates and concentrations of O₃ and PM_{2.5} were similar under the four SWPs; (3) in the YRD region, the O₃ pollution over-limit rate presents as Type 1 > Type 4 > Type 2 > Type 3, PM_{2.5} pollution largely appeared under Type 1, and both O₃ and PM_{2.5} under Type 1 were higher than those for the other types; (4) in the GZP region, the O₃ pollution frequency was higher under Type 2 and Type 4, and PM_{2.5} pollution occurred more frequently under Type 2; and (5) in the NEM region, O₃ pollution was always found under Type 1, Type 2 and Type 4, but the over-limit rate was no more than 15% and PM_{2.5} pollution under Type 1 was more than under Type 2.

In summary, Type 1 was prone to the formation of O₃–PM_{2.5} compound pollution (that is, when the ground MDA8 O₃ concentration exceeded 160 µg m⁻³, the PM_{2.5} concentration also exceeded 75 µg m⁻³) in the area from the BTH to northern YRD regions (Fig. S6), which can be denoted as “BTH–NYRD O₃–PM_{2.5} compound pollution”. In detail, Figure S8 shows the number and probability of occurrence of compound pollution days at each site in summer during 2015–2018, indicating that a high occurrence probability (maximum values approaching 46.7%) of compound pollution appeared over the NCP (to the north of 32°N), and that approximately 55.6% of compound pollution occurrence days at all sites occurred under Type 1. Similarly, Type 2 can also be denoted as “BTH O₃–PM_{2.5} compound pollution” with compound pollution occurrence days accounting for 33.8%, Type 3 as “BTH–YRD–PRD O₃-only pollution”, and Type 4 as “BTH–GZP–YRD–PRD O₃-only pollution” (Fig.12).

3.4 Analysis of local meteorological factors

To explore the meteorological causes of O₃ and PM_{2.5} pollution, we analyzed the distribution of the average and anomalies for Tmax, RH, PF, BLH and FLWD under the four SWPs (Figs. S9, S10, 8 and 9). Under the influence of the EASM, over 80% of the stations experience high temperatures (Tmax > 27°C) under each SWP, although the anomaly of Tmax under Type 1 (early

summer) presented as negative (Fig. 8a). Type 1 was characterized by humid condition in the southern area and dry conditions in the northern region owing to an extensive southwestern flow of the WPSH, resulting in a rain belt found in southeastern coastal areas such as the PRD and YRD regions. Type 2 was associated with meridional flow and dry and wet anomalies in northern China, resulting in a rain band located over the central areas between the BTH and YRD regions owing to the northern advancement of the WPSH compared with Type 1. Furthermore, there was higher RH for most of the study sites under Type 3 and Type 4, possibly because of the shifted rain belt in the BTH and NEM regions under Type 3 once the northern boundary of the WPSH reached 37.5°N, and an occurrence of heavy precipitation across the western PRD region as well as central areas between the BTH and YRD regions under Type 4 (Fig. S9).

In terms of their anomalous spatial distributions, the positive anomalies of Tmax were located in the southern region under Type 3 and most of the eastern region under Type 4; and since Type 1 always appeared in early summer, most areas were negative (Figs. 8a–d). For RH, Types 2, 3 and 4 were negative for the south and positive in the north, while the opposite was true under Type 1 (Figs. 8e–h). PF was characterized by positive anomalies in the area south of the Yangtze River under Type 1, in the YRD region under Type 2, in the BTH and NEM regions under Type 3, and in the area between the BTH and YRD regions under Type 4 (Figs. 8i–l). As can be seen from Fig. 9, when the BLH at 1400 BJT had a positive anomaly, on the contrary FLWD had a negative anomaly (e.g., BTH in Type 1), which indicates that the higher BLH and the lower FLWD, the more conducive it was to the diffusion of pollutants; and conversely, a lower BLH and higher FLWD (such as BTH under Type 2) did not support the diffusion of pollutants. After further inspection of Fig. S10, we found that the YRD region under Type 1, the YRD under Type 2, the BTH and PRD regions under Type 3 and 4 had shallow BLHs and high FLWDs, which was detrimental to the transportation of pollution in these areas, thus corresponding to high levels of pollution under these weather patterns. However, there was also more serious pollution in some higher BLH areas, such as in the BTH region under Type 1, which we discuss next.

3.5 Potential implications of NO₂

The photochemical production of O₃ mainly involve emissions of volatile organic compounds

and NO_x from anthropogenic, biogenic and biomass burning sources (Deng et al., 2019; Gvozdić et al., 2011; Sillman, 2002). The photochemical reaction of NO , NO_2 , and O_3 in the troposphere forms a closed system (Yu et al., 2020), and this photochemical cycle of NO_x and O_3 is the basis of photochemical processes in the troposphere. Oxidant (Ox , $\text{Ox} = \text{O}_3 + \text{NO}_2$), a conservative
400 quantity over short time scales, is defined as a parameter to evaluate the photochemical processes and, due to the unstable nature of NO , it can quickly react with the equivalent amount of O_3 to generate NO_2 (Kley et al., 1994). In order to compare the photochemical reaction efficiency of the five urban clusters under the different SWPs, Fig. 10 presents the daily variations of NO_2 and Ox . As we can see, the daily variations of NO_2 showed two peaks during a day, including a first peak
405 in the morning and a second peak associated with traffic emissions in the evening (Xie et al., 2016; Yu et al., 2020). As we found the lowest point of NO_2 at 1500 BJT, and NO_2 can be photolyzed to produce O_3 during the day, we assumed that this particular time was the peak time for O_3 formation across the study areas. As NO_2 is consumed through a photochemical reaction with the involvement of other precursors to produce a large amount of O_3 , Ox can form a peak in the
410 afternoon. In particular, abundant sunlight in summer is beneficial to the photochemical reaction process, but since most parts of eastern China are under a subtropical climate with the same period of rain and heat, the existence of the rainy season will inevitably inhibit the summertime photochemical process. Under the different SWPs, the photochemical reaction over each area bore an obvious relationship with the rain belt. For example, the rainy season in the BTH and NEM
415 areas mainly occurred under Type 3, and the Ox of Type 3 in this area was significantly lower than under the other SWPs.

4. Discussion

In the last section, we discussed how the SWPs and local meteorological factors modified the
420 summertime O_3 and $\text{PM}_{2.5}$ pollution. However, how did the boundary layer structure interact with the co-occurrence of O_3 and $\text{PM}_{2.5}$ pollution? To address this question, we conducted some further analysis as follows. As mentioned, the co-occurrence of O_3 and $\text{PM}_{2.5}$ pollution mainly took place in the BTH–NYRD areas under Type 1 and in the BTH area under Type 2. Lower WS and its negative anomalies at a lower boundary layer over the BTH–NYRD under Type 1 and over the

425 BTH area under Type 2, may not have enhanced the diffusion of air pollutants (Fig. S11). In contrast, the moderate RH and its negative anomalies might have favored the formation of compound pollution. Downward vertical motion and negative anomalies might also have stabilized the atmospheric characteristic of the boundary layer (Fig. S12). Furthermore, we summarized the boundary layer structure, precipitation, and ground-level wind flow across the BTH region. Based on the characteristics, we separately defined Type 1 and Type 2 into clean (the concentrations of both O_3 and $PM_{2.5}$ were less than level of pollution) and compound pollution periods (Figs. 11 and S13-S14). In particular, Type 1 had significantly warmer temperatures over the boundary layer during the compound pollution periods of the BTH region, as compared with the clean periods. The daytime BLH under the compound pollution condition was also higher than that under the clean condition. In addition, there were different directions of prevailing winds during the two periods. The prevailing southerly winds during the compound pollution period may have driven the transportation of air pollutants from the southern plains, resulting in more serious pollution (Fig. 11; see also. Miao et al. (2020) also proposed another mechanism—that is, the synoptic southerly warm advections at the top of PBL, can strengthen the elevated thermal inversion layer and suppress the development of the PBL, causing worse pollution. Co-influenced by the topographical effect of the northern mountainous areas and the boundary layer structure, air pollutants could be trapped in the BTH region. In comparison, although there was a southerly prevailing wind in the BTH region (Figs. 11 and S14), the rain belt also being located in the southern area of the BTH might have led to the potential removal of $PM_{2.5}$ (Fig. 9j). Therefore, compound pollution across the BTH region might mainly have been due to local emissions of air pollutants.

In summary, the different SWPs modulated the regional variability of summertime O_3 and $PM_{2.5}$ via changes to the local meteorological conditions as follows:

(1) Type 1: Under the conditions of high temperatures ($T_{max} > 27^\circ C$), moderate humidity (RH ~60%), and low PF, photochemical reactions were greatly promoted to cause severe O_3 pollution. Meanwhile, the BTH–NYRD areas were located in front of the westerly trough, under the influence of the warm and humid air of the WPSH, and so the hygroscopic growth of fine particulates potentially caused a certain amount of $PM_{2.5}$ pollution (Li et al., 2017; Zhang et al.,

2016), becoming O_3 – $PM_{2.5}$ compound pollution (Fig. 12). In addition, the prevailing southerly
455 winds in the boundary layer were able to transport the pollutants emitted from southern cities to
the BTH, atmospheric stratification was stable when the air mass was sinking (Miao et al., 2019b;
Figs. 11 and S12), and compound pollution may have been especially severe. Although a relatively
higher BLH occurred in the BTH region, the prevailing southerly winds in the boundary layer
served to further increase the pollution.

460 (2) Type 2: O_3 pollution was severe under the meteorological conditions of high
temperatures, moderate humidity, and weak precipitations. The $PM_{2.5}$ in the BTH region, which
was located in front of the westerly trough, was high since the shallow boundary layer and low
wind frequency were unfavorable for the diffusion of pollutants. Therefore, O_3 – $PM_{2.5}$ compound
pollution was also rather frequent (Fig. 12).

465 (3) Type 3: High temperatures, low humidity, and weak precipitations over the YRD region
tended to generate a large amount of O_3 , while the positive BLH and negative FLWD anomalies
were unfavorable to O_3 accumulation. On the other hand, summer typhoon activities might have
weakened the WPSH intensity over the YRD region, leading to the eastward retreat and northward
shift of the WPSH. As a result, the high WS across coastal areas was able to ease the ground-level
470 O_3 pollution (Shu et al., 2016). For the BTH and PRD regions, the high PF tended to suppress the
production of O_3 .

(4) Type 4: High temperatures, medium-high humidity and weak precipitations in the GZP
and PRD regions were able to cause O_3 – $PM_{2.5}$ compound pollution, but the $PM_{2.5}$ pollution in both
regions was not heavy, possibly in relation to local lower emissions of pollutants. Under the
475 control of the WPSH, there were strong photochemical reactions at high temperatures and little
rainfall in some eastern regions (such as the northern BTH, YRD), which was also conducive to
 O_3 generation (Fig. 12). Meanwhile, relative to Type 1, O_3 pollution was lighter in the BTH, due to
the differences of RH, BLH and FLWD.

It is important to note that our work contains a few limitations and uncertainties. Although
480 T-PCA, an objective classification method, was chosen in this study, there were still some
subjective decisions made, e.g., the number of SWPs (Huth et al., 2008). In the present work, we
selected four SWPs based on both the larger ECV and greater ΔECV to furthest reduce the

subjective impact. Nevertheless, at a large scale, the present four SWPs were closely associated with intraseasonal movements of the WPSH, because the WPSH is one of the most important components of the present large-scale SWPs in summertime (Zhao and Wang, 2017). In addition, note that short-term disturbances induced by typhoons with specific pattern were not excluded. The quick passage of a typhoon in summer could lead to various atmospheric processes (e.g., precipitation, large-scale subsidence) and pollution levels (Deng et al., 2019), which should be explored in future work. Besides, although this study emphasizes the important impacts of large-scale synoptic drivers of co-occurring summertime O_3 and $PM_{2.5}$ pollution in eastern China, the presences of $PM_{2.5}$ may play a role in radiation forcing to reduce O_3 . Indeed, the interaction between O_3 and $PM_{2.5}$ deserving of further exploration in future work to better comprehend the mechanism of O_3 – $PM_{2.5}$ compound pollution.

5 Conclusions

In this study, T-PCA, an objective classification method, was applied to classify the 500-hPa weather circulation pattern into four SWPs in the summers of 2015–2018. It was found that these four SWPs were closely related to the development of the WPSH. The spatial and temporal distribution characteristics of O_3 and $PM_{2.5}$ pollution in eastern China under the four SWPs were analyzed to regulate and differentiate O_3 and $PM_{2.5}$ pollution in key areas. We found two synoptic patterns were prone to leading to the co-occurrence of O_3 and $PM_{2.5}$ pollution: in BTH–NYRD areas under Type 1, and the BTH area under Type 2 were associated with the double high levels of O_3 and $PM_{2.5}$. The probabilities of compound pollution at all sites under Type 1, 2, 3, and 4 were 54.3%, 33.8%, 6.8%, and 5.1% respectively.

The Type 1 weather pattern appeared frequently in early summer, with a stable WPSH ridge axis at about 22°N, and the warm and humid air brought by the WPSH reached the area south of the Yangtze River, where a hot and humid Meiyu season was formed, with the high humidity suppressing the photochemical reaction of O_3 generation. Meanwhile, the north of China was controlled by a low-pressure trough at 500 hPa with high temperatures and little rain. The hygroscopic growth of $PM_{2.5}$ occurred in the corresponding area in front of the trough, with a small amount of water vapor transported by the WPSH, causing compound pollution of O_3 and

PM_{2.5} in the BTH–NYRD regions. In addition, the prevailing southerly winds in the boundary layer were able to transport the pollutants emitted from southern cities to the BTH region, and the atmospheric stratification was stable when the air mass was sinking. Thus, the compound pollution was potentially severe. In general, the synoptic circulation in the boundary layer might be responsible for the concentration of pollutants under this SWP.

Under Type 2, the WPSH shifted northwards from Type 1 or retreated southwards from Type 3 or Type 4, to 32.5°N, with the meridional deepening of the East Asian major trough at 500 hPa, and thus warm and humid airstreams were brought to the northern China (e.g., the BTH region), gradually elevating temperatures and humidity. Although the positive RH anomaly promoted the hygroscopic growth of PM_{2.5}, water vapor absorbed solar radiation leading by contrast to reduced O₃ formation. As a result, the probability of double high levels of O₃ and PM_{2.5} under Type 2 was less than under Type 1, and the extent of compound pollution under Type 2 was also narrowed, which was mainly located in the BTH area. On the other hand, weak precipitation, shallow boundary layer and low WS in the BTH area tended to create favorable conditions for the maintenance of pollution. In spite of the southerly winds over the BTH area, the precipitation in southern cities reduced pollutant concentrations and horizontal transportation. Meteorological factors might have been responsible for the accumulation of compound pollution.

In general, the location of the WPSH was found to be tightly associated with O₃ pollution in eastern China, and the changes in meteorological conditions in different regions affected by the WPSH induced significant regional differences in O₃ and PM_{2.5} pollution. On the one hand, appropriate warm and moist flow brought by the WPSH promoted hygroscopic growth of the fine particulate matter in some local areas (i.e., the BTH–NYRD areas under Type 1 and the BTH area under Type 2), resulting in increased of PM_{2.5} concentrations; whilst on the other hand, transboundary O₃ and PM_{2.5} were simultaneously transported to these local areas, which may have contributed to the formation of co-occurring surface O₃ and PM_{2.5} pollution. More importantly, the effects of various large-scale weather circulation patterns on O₃–PM_{2.5} compound pollution and their corresponding physical and chemical processes, have been clarified, which has important scientific reference value in terms of summer air-quality forecasts, as well as assessment and policy-making services.

Data availability

Hourly PM_{2.5}, NO₂, O₃, and O₃-8h data are published by the China Environmental Monitoring Station (<http://www.cnemc.cn/>). Surface meteorological data, such as Tmax, precipitation, WS and RH, and radiosonde data can be obtained from the China National Meteorological Information Center of the China Meteorological Administration (<http://data.cma.cn/site/index.html>). The NCEP/NCAR daily reanalysis dataset can be download from <https://psl.noaa.gov/data/gridded/data.ncep.reanalysis.html>. The ERA5 hourly reanalysis dataset can be derived from <https://cds.climate.copernicus.eu/cdsapp#!/home>.

Author contributions

L. Zong: Methodology, Data Curation, Formal Analysis, Writing—Original draft preparation, Results Discussion, Writing—Reviewing and Editing; Y. Yang: Conceptualization, Methodology, Formal Analysis, Results Discussion, Writing—Reviewing and Editing; M. Gao, H. Wang, P. Wang, L. Wang, H. Zhang, G. Ning, C. Liu, Y. Li, Z. Gao: Results Discussion, Comments, Writing—Reviewing and Editing.

Competing interests

The authors declare that they have no conflict of interests.

Acknowledgments

This study was jointly funded by the National Key Research and Development Program of China (2018YFC1506502) and the National Natural Science Foundation of China (41871029).

References

- Amil, N., Latif, M. T., Khan, M. F. and Mohamad, M.: Seasonal variability of PM_{2.5} composition and sources in the Klang Valley urban-industrial environment, *Atmos. Chem. Phys.*, 16(8), 5357–5381, doi:10.5194/acp-16-5357-2016, 2016.
- Cai, W., Li, K., Liao, H., Wang, H. and Wu, L.: Weather conditions conducive to Beijing severe

- 570 haze more frequent under climate change, *Nat. Clim. Chang.*, 7(4), 257–262, doi:10.1038/nclimate3249, 2017.
- Chen, C., Saikawa, E., Comer, B., Mao, X. and Rutherford, D.: Ship Emission Impacts on Air Quality and Human Health in the Pearl River Delta (PRD) Region, China, in 2015, With Projections to 2030, *GeoHealth*, 3(9), 284–306, doi:10.1029/2019GH000183, 2019.
- 575 Chen, H. and Wang, H.: Haze days in North China and the associated atmospheric circulations based on daily visibility data from 1960 to 2012, *J. Geophys. Res.*, 120(12), 5895–5909, doi:10.1002/2015JD023225, 2015.
- Cohen, A. J., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., 580 Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope, C. A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C. J. L. and Forouzanfar, M. H.: Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015, *Lancet*, 389(10082), 1907–1918, doi:10.1016/S0140-6736(17)30505-6, 2017.
- 585 Day, D. B., Xiang, J., Mo, J., Li, F., Chung, M., Gong, J., Weschler, C. J., Ohman-Strickland, P. A., Sundell, J., Weng, W., Zhang, Y. and Zhang, J. J.: Association of ozone exposure with cardiorespiratory pathophysiologic mechanisms in healthy adults, *JAMA Intern. Med.*, 177(9), 1344–1353, doi:10.1001/jamainternmed.2017.2842, 2017.
- Deng, Y., Li, J., Li, Y., Wu, R. and Xie, S.: Characteristics of volatile organic compounds, NO₂, 590 and effects on ozone formation at a site with high ozone level in Chengdu, *J. Environ. Sci. (China)*, 75(2), 334–345, doi:10.1016/j.jes.2018.05.004, 2019.
- Ding, Y.: The Summer Monsoon in East Asia, in *Monsoons over China*, pp. 1–90., 1994.
- Du, Y., Wan, Q., Liu, H., Liu, H., Kapsar, K. and Peng, J.: How does urbanization influence PM 2.5 concentrations? Perspective of spillover effect of multi-dimensional urbanization impact, 595 *J. Clean. Prod.*, 220, 974–983, doi:10.1016/j.jclepro.2019.02.222, 2019.
- Fan, H., Zhao, C. and Yang, Y.: A comprehensive analysis of the spatio-temporal variation of urban air pollution in China during 2014–2018, *Atmos. Environ.*, 220(November), 117066, doi:10.1016/j.atmosenv.2019.117066, 2020.

Ge, J., You, Q. and Zhang, Y.: Effect of Tibetan Plateau heating on summer extreme precipitation
600 in eastern China, *Atmos. Res.*, 218, 364–371, doi:10.1016/j.atmosres.2018.12.018, 2019.

Gong, C. and Liao, H.: A typical weather pattern for ozone pollution events in North China, *Atmos.*
Chem. Phys., 19(22), 13725–13740, doi:10.5194/acp-19-13725-2019, 2019.

Gui, K., Che, H., Wang, Y., Wang, H., Zhang, L., Zhao, H., Zheng, Y., Sun, T. and Zhang, X.:
Satellite-derived PM_{2.5} concentration trends over Eastern China from 1998 to 2016:
605 Relationships to emissions and meteorological parameters, *Environ. Pollut.*, 247, 1125–1133,
doi:10.1016/j.envpol.2019.01.056, 2019.

Guo, J., Miao, Y., Zhang, Y., Liu, H., Li, Z., Zhang, W., He, J., Lou, M., Yan, Y., Bian, L. and Zhai,
P.: The climatology of planetary boundary layer height in China derived from radiosonde and
reanalysis data, *Atmos. Chem. Phys.*, 16(20), 13309–13319, doi:10.5194/acp-16-13309-2016,
610 2016.

Guo, J., Li, Y., Cohen, J. B., Li, J., Chen, D., Xu, H., Liu, L., Yin, J., Hu, K. and Zhai, P.: Shift in
the Temporal Trend of Boundary Layer Height in China Using Long-Term (1979–2016)
Radiosonde Data, *Geophys. Res. Lett.*, 46(11), 6080–6089, doi:10.1029/2019GL082666,
2019.

615 Gvozdić, V., Kovač-Andrić, E. and Brana, J.: Influence of Meteorological Factors NO₂, SO₂, CO
and PM₁₀ on the Concentration of O₃ in the Urban Atmosphere of Eastern Croatia, *Environ.*
Model. Assess., 16(5), 491–501, doi:10.1007/s10666-011-9256-4, 2011.

Han, H., Liu, J., Shu, L., Wang, T. and Yuan, H.: Local and synoptic meteorological influences on
daily variability in summertime surface ozone in eastern China, *Atmos. Chem. Phys.*, 20(1),
620 203–222, doi:10.5194/acp-20-203-2020, 2020.

Hoffmann, P. and Heinke SchlüNzen, K.: Weather pattern classification to represent the urban heat
island in present and future climate, *J. Appl. Meteorol. Climatol.*, 52(12), 2699–2714,
doi:10.1175/JAMC-D-12-065.1, 2013.

Huang, X., Ding, A., Liu, L., Liu, Q., Ding, K., Nie, W., Xu, Z., Chi, X., Wang, M., Sun, J., Guo,
625 W. and Fu, C.: Effects of aerosol-radiation interaction on precipitation during
biomass-burning season in East China, *Atmos. Chem. Phys. Discuss.*, (April), 1–37,
doi:10.5194/acp-2016-272, 2016.

- Huth, R., Beck, C., Philipp, A., Demuzere, M., Ustrnul, Z., Cahynová, M., Kyselý, J. and Tveito, O. E.: Classifications of atmospheric circulation patterns: Recent advances and applications, *Ann. N. Y. Acad. Sci.*, 1146, 105–152, doi:10.1196/annals.1446.019, 2008.
- 630 Ji, X., Yao, Y. and Long, X.: What causes PM_{2.5} pollution? Cross-economy empirical analysis from socioeconomic perspective, *Energy Policy*, 119(April), 458–472, doi:10.1016/j.enpol.2018.04.040, 2018.
- Kley, D., Geiss, H. and Mohnen, V. A.: Tropospheric ozone at elevated sites and precursor emissions in the United States and Europe, *Atmos. Environ.*, 28(1), 149–158, doi:10.1016/1352-2310(94)90030-2, 1994.
- 635 Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K. H., Zhang, Q. and Zhai, S.: A two-pollutant strategy for improving ozone and particulate air quality in China, *Nat. Geosci.*, 12(11), 906–910, doi:10.1038/s41561-019-0464-x, 2019.
- 640 Li, M., Wang, L., Liu, J., Gao, W., Song, T., Sun, Y., Li, L., Li, X., Wang, Y., Liu, L., Daellenbach, K. R., Paasonen, P. J., Kerminen, V. M., Kulmala, M. and Wang, Y.: Exploring the regional pollution characteristics and meteorological formation mechanism of PM_{2.5} in North China during 2013–2017, *Environ. Int.*, 134(November 2019), 105283, doi:10.1016/j.envint.2019.105283, 2020.
- 645 Li, S., Wang, T., Huang, X., Pu, X., Li, M., Chen, P., Yang, X. Q. and Wang, M.: Impact of East Asian Summer Monsoon on Surface Ozone Pattern in China, *J. Geophys. Res. Atmos.*, 123(2), 1401–1411, doi:10.1002/2017JD027190, 2018.
- Li, Z., Guo, J., Ding, A., Liao, H., Liu, J., Wang, T., Xue, H., Zhang, H. and Zhu, B.: Aerosol and boundary-layer interactions and impact on air quality, *Natl. Sci. Rev.*, 4(6), 810–833, doi:10.1093/nsr/nwx117/4191281, 2017.
- 650 Liu, H., Tian, H., Zhang, K., Liu, S., Cheng, K., Yin, S., Liu, Y., Liu, X., Wu, Y., Liu, W., Bai, X., Wang, Y., Shao, P., Luo, L., Lin, S., Chen, J. and Liu, X.: Seasonal variation, formation mechanisms and potential sources of PM_{2.5} in two typical cities in the Central Plains Urban Agglomeration, China, *Sci. Total Environ.*, 657, 657–670, doi:10.1016/j.scitotenv.2018.12.068, 2019a.
- 655 Liu, J. and You, Q.: A diagnosis of the interannual variation of the summer hydrometeor based on

ERA-interim over Eastern China, *Atmos. Res.*, 231(October 2018), 104654, doi:10.1016/j.atmosres.2019.104654, 2020.

Liu, N., Zhou, S., Liu, C. and Guo, J.: Synoptic circulation pattern and boundary layer structure associated with PM_{2.5} during wintertime haze pollution episodes in Shanghai, *Atmos. Res.*, 228(46), 186–195, doi:10.1016/j.atmosres.2019.06.001, 2019b.

Lu, R.: Indices of the Summertime Western North Pacific Subtropical High, *Adv. Atmos. Sci.*, 19(6), 1004–1028, doi:10.1007/s00376-002-0061-5, 2002.

Ma, Z., Hu, X., Sayer, A. M., Levy, R., Zhang, Q., Xue, Y., Tong, S., Bi, J., Huang, L. and Liu, Y.: Satellite-based spatiotemporal trends in PM_{2.5} concentrations: China, 2004–2013, *Environ. Health Perspect.*, 124(2), 184–192, doi:10.1289/ehp.1409481, 2016.

Miao, Y., Hu, X.-M., Liu, S., Qian, T., Xue, M., Zheng, Y. and Wang, S.: Seasonal variation of local atmospheric circulations and boundary layer structure in the Beijing-Tianjin-Hebei region and implications for air quality., *J. Adv. Model. Earth Syst.*, 7, 1602–1626, doi:10.1002/2015ms000522, 2015.

Miao, Y., Guo, J., Liu, S., Liu, H., Li, Z., Zhang, W. and Zhai, P.: Classification of summertime synoptic patterns in Beijing and their associations with boundary layer structure affecting aerosol pollution, *Atmos. Chem. Phys.*, 17(4), 3097–3110, doi:10.5194/acp-17-3097-2017, 2017.

Miao, Y., Liu, S. and Huang, S.: Synoptic pattern and planetary boundary layer structure associated with aerosol pollution during winter in Beijing, China, *Sci. Total Environ.*, 682, 464–474, doi:10.1016/j.scitotenv.2019.05.199, 2019.

Miao, Y., Che, H., Zhang, X. and Liu, S.: Relationship between summertime concurring PM_{2.5} and O₃ pollution and boundary layer height differs between Beijing and Shanghai, China, *Environ. Pollut.*, doi:10.1016/j.envpol.2020.115775, 2020.

Ning, G., Wang, S., Yim, S. H. L., Li, J., Hu, Y., Shang, Z., Wang, J. and Wang, J.: Impact of low-pressure systems on winter heavy air pollution in the northwest Sichuan Basin, China, *Atmos. Chem. Phys.*, 18(18), 13601–13615, doi:10.5194/acp-18-13601-2018, 2018.

Ning, G., Yim, S. H. L., Wang, S., Duan, B., Nie, C., Yang, X., Wang, J. and Shang, K.: Synergistic effects of synoptic weather patterns and topography on air quality: a case of the

Sichuan Basin of China, *Clim. Dyn.*, 53(11), 6729–6744, doi:10.1007/s00382-019-04954-3, 2019.

Philipp, A., Beck, C., Esteban, P., Krennert, T., Lochbihler, K., Spyros, P., Pianko-kluczynska, K., Post, P., Alvarez, R., Spekat, A. and Streicher, F.: Cost733 user guide., 2014.

690 Saikawa, E., Kim, H., Zhong, M., Avramov, A., Zhao, Y., Janssens-Maenhout, G., Kurokawa, J. I., Klimont, Z., Wagner, F., Naik, V., Horowitz, L. W. and Zhang, Q.: Comparison of emissions inventories of anthropogenic air pollutants and greenhouse gases in China, *Atmos. Chem. Phys.*, 17(10), 6393–6421, doi:10.5194/acp-17-6393-2017, 2017.

Seidel, D. J., Zhang, Y., Beljaars, A., Golaz, J. C., Jacobson, A. R. and Medeiros, B.: Climatology
695 of the planetary boundary layer over the continental United States and Europe, *J. Geophys. Res. Atmos.*, 117(17), 1–15, doi:10.1029/2012JD018143, 2012.

Shi, Z., Huang, L., Li, J., Ying, Q., Zhang, H. and Hu, J.: Sensitivity Analysis of the Surface Ozone and Fine Particulate Matter to Meteorological Parameters in China, *Atmos. Chem. Phys. Discuss.*, 2020, 1–29, doi:10.5194/acp-2020-173, 2020.

700 Shu, L., Xie, M., Wang, T., Gao, D., Chen, P., Han, Y., Li, S., Zhuang, B. and Li, M.: Integrated studies of a regional ozone pollution synthetically affected by subtropical high and typhoon system in the Yangtze River Delta region, China, *Atmos. Chem. Phys.*, 16(24), 15801–15819, doi:10.5194/acp-16-15801-2016, 2016.

Shu, L., Wang, T., Xie, M., Li, M., Zhao, M., Zhang, M. and Zhao, X.: Episode study of fine
705 particle and ozone during the CAPUM-YRD over Yangtze River Delta of China: Characteristics and source attribution, *Atmos. Environ.*, 203(July 2018), 87–101, doi:10.1016/j.atmosenv.2019.01.044, 2019.

Sillman, S.: Chapter 12 The relation between ozone, NO_x and hydrocarbons in urban and polluted
rural environments, *Dev. Environ. Sci.*, 1(C), 339–385, doi:10.1016/S1474-8177(02)80015-8,
710 2002.

Song, C., Wu, L., Xie, Y., He, J., Chen, X., Wang, T., Lin, Y., Jin, T., Wang, A., Liu, Y., Dai, Q., Liu, B., Wang, Y. nan and Mao, H.: Air pollution in China: Status and spatiotemporal variations, *Environ. Pollut.*, 227, 334–347, doi:10.1016/j.envpol.2017.04.075, 2017.

Su, H., Qian, C., Gu, H. and Wang, Q.: The Impact of Tropical Cyclones on China in 2016, *Trop.*

- 715 Cyclone Res. Rev., 5(1–2), 1–11, doi:10.6057/2017TCRRh1.01, 2017
- Sun, L., Xue, L., Wang, T., Gao, J., Ding, A., Cooper, O. R., Lin, M., Xu, P., Wang, Z., Wang, X.,
Wen, L., Zhu, Y., Chen, T., Yang, L., Wang, Y., Chen, J. and Wang, W.: Significant increase of
summertime ozone at Mount Tai in Central Eastern China, *Atmos. Chem. Phys.*, 16(16),
10637–10650, doi:10.5194/acp-16-10637-2016, 2016.
- 720 Tai, A. P. K., Mickley, L. J. and Jacob, D. J.: Correlations between fine particulate matter (PM_{2.5})
and meteorological variables in the United States: Implications for the sensitivity of PM_{2.5} to
climate change, *Atmos. Environ.*, 44(32), 3976–3984, doi:10.1016/j.atmosenv.2010.06.060,
2010.
- Wang, H. J. and Chen, H. P.: Understanding the Recent Trend of Haze Pollution in Eastern China:
725 Roles of Climate Change, *Atmos. Chem. Phys. Discuss.*, 2016(January), 1–18,
doi:10.5194/acp-2015-1009, 2016.
- Wang, T., Zhong, Z., Sun, Y. and Wang, J.: Impacts of tropical cyclones on the meridional
movement of the western Pacific subtropical high, *Atmos. Sci. Lett.*, 20(5), 1–8,
doi:10.1002/asl.893, 2019.
- 730 Xie, M., Zhu, K., Wang, T., Chen, P., Han, Y., Li, S., Zhuang, B. and Shu, L.: Temporal
characterization and regional contribution to O₃ and NO_x at an urban and a suburban site in
Nanjing, China, *Sci. Total Environ.*, 551–552(x), 533–545,
doi:10.1016/j.scitotenv.2016.02.047, 2016.
- Yang, Y., Wang, R., Chen, F., Liu, C., Bi, X. and Huang, M.: Synoptic weather patterns modulate
735 the frequency, type and vertical structure of summer precipitation over Eastern China: A
perspective from GPM observations, *Atmos. Res.*, 249, doi:10.1016/j.atmosres.2020.105342,
2021.
- Yang, Y., Zheng, X., Gao, Z., Wang, H., Wang, T., Li, Y., Lau, G. N. C. and Yim, S. H. L.:
Long-Term Trends of Persistent Synoptic Circulation Events in Planetary Boundary Layer
740 and Their Relationships With Haze Pollution in Winter Half Year Over Eastern China, *J.*
Geophys. Res. Atmos., 123(19), 10,991–11,007, doi:10.1029/2018JD028982, 2018.
- Yang, Y., Zheng, Z., Yim, S. Y. L., Roth, M., Ren, G., Gao, Z., Wang, T., Li, Q., Shi, C., Ning, G.
and Li, Y.: PM_{2.5} Pollution Modulates Wintertime Urban Heat Island Intensity in the

- Beijing-Tianjin-Hebei Megalopolis, China, *Geophys. Res. Lett.*, 47(1), 0–3,
745 doi:10.1029/2019GL084288, 2020.
- Ye, W. F., Ma, Z. Y. and Ha, X. Z.: Spatial-temporal patterns of PM_{2.5} concentrations for 338
Chinese cities, *Sci. Total Environ.*, 631–632, 524–533, doi:10.1016/j.scitotenv.2018.03.057,
2018.
- Yim, S. H. L., Wang, M., Gu, Y., Yang, Y., Dong, G. and Li, Q.: Effect of Urbanization on Ozone
750 and Resultant Health Effects in the Pearl River Delta Region of China, *J. Geophys. Res. Atmos.*, 124(21), 11568–11579, doi:10.1029/2019JD030562, 2019.
- Yin, Z., Cao, B. and Wang, H.: Dominant patterns of summer ozone pollution in eastern China and
associated atmospheric circulations, *Atmos. Chem. Phys.*, 19(22), 13933–13943,
doi:10.5194/acp-19-13933-2019, 2019.
- 755 Yu, S., Yin, S., Zhang, R., Wang, L., Su, F., Zhang, Y. and Yang, J.: Spatiotemporal
characterization and regional contributions of O₃ and NO₂: An investigation of two years of
monitoring data in Henan, China, *J. Environ. Sci. (China)*, 90(November), 29–40,
doi:10.1016/j.jes.2019.10.012, 2020.
- Zhang, C., Liu, C., Hu, Q., Cai, Z., Su, W., Xia, C., Zhu, Y., Wang, S. and Liu, J.: Satellite UV-Vis
760 spectroscopy: implications for air quality trends and their driving forces in China during
2005–2017, *Light Sci. Appl.*, 8(1), doi:10.1038/s41377-019-0210-6, 2019a.
- Zhang, H., Wang, Y., Park, T. W. and Deng, Y.: Quantifying the relationship between extreme air
pollution events and extreme weather events, *Atmos. Res.*, 188, 64–79,
doi:10.1016/j.atmosres.2016.11.010, 2017.
- 765 Zhang, Q., Zheng, Y., Tong, D., Shao, M., Wang, S., Zhang, Y., Xu, X., Wang, J., He, H., Liu, W.,
Ding, Y., Lei, Y., Li, J., Wang, Z., Zhang, X., Wang, Y., Cheng, J., Liu, Y., Shi, Q., Yan, L.,
Geng, G., Hong, C., Li, M., Liu, F., Zheng, B., Cao, J., Ding, A., Gao, J., Fu, Q., Huo, J., Liu,
B., Liu, Z., Yang, F., He, K. and Hao, J.: Drivers of improved PM_{2.5} air quality in China
from 2013 to 2017, *Proc. Natl. Acad. Sci. U. S. A.*, 116(49), 24463–24469,
770 doi:10.1073/pnas.1907956116, 2019b.
- Zhang, R. H., Li, Q. and Zhang, R. N.: Meteorological conditions for the persistent severe fog and
haze event over eastern China in January 2013, *Sci. China Earth Sci.*, 57(1), 26–35,

doi:10.1007/s11430-013-4774-3, 2014.

- 775 Zhang, W., Wang, H., Zhang, X., Peng, Y., Zhong, J., Wang, Y. and Zhao, Y.: Evaluating the contributions of changed meteorological conditions and emission to substantial reductions of PM_{2.5} concentration from winter 2016 to 2017 in Central and Eastern China, *Sci. Total Environ.*, 716, 136892, doi:10.1016/j.scitotenv.2020.136892, 2020.
- Zhang, Y. L. and Cao, F.: Fine particulate matter (PM 2.5) in China at a city level, *Sci. Rep.*, 5(2014), 1–12, doi:10.1038/srep14884, 2015.
- 780 Zhang, Z., Zhang, X., Gong, D., Quan, W., Zhao, X., Ma, Z. and Kim, S. J.: Evolution of surface O₃ and PM_{2.5} concentrations and their relationships with meteorological conditions over the last decade in Beijing, *Atmos. Environ.*, 108, 67–75, doi:10.1016/j.atmosenv.2015.02.071, 2015.
- Zhang, Z., Zhang, X., Gong, D., Kim, S. J., Mao, R. and Zhao, X.: Possible influence of atmospheric circulations on winter haze pollution in the Beijing-Tianjin-Hebei region, northern China, *Atmos. Chem. Phys.*, 16(2), 561–571, doi:10.5194/acp-16-561-2016, 2016.
- 785 Zhao, C., Wang, Y., Yang, Q., Fu, R., Cunnold, D. and Choi, Y.: Impact of East Asian summer monsoon on the air quality over China: View from space, *J. Geophys. Res. Atmos.*, 115(9), 1–12, doi:10.1029/2009JD012745, 2010.
- 790 Zhao, W., Tang, G., Yu, H., Yang, Y., Wang, Y., Wang, L., An, J., Gao, W., Hu, B., Cheng, M., An, X., Li, X. and Wang, Y.: Evolution of boundary layer ozone in Shijiazhuang, a suburban site on the North China Plain, *J. Environ. Sci. (China)*, 83, 152–160, doi:10.1016/j.jes.2019.02.016, 2019.
- Zhao, Z. and Wang, Y.: Influence of the West Pacific subtropical high on surface ozone daily variability in summertime over eastern China, *Atmos. Environ.*, 170, 197–204, doi:10.1016/j.atmosenv.2017.09.024, 2017.
- 795 Zheng, X. Y., Fu, Y. F., Yang, Y. J. and Liu, G. S.: Impact of atmospheric circulations on aerosol distributions in autumn over eastern China: Observational evidence, *Atmos. Chem. Phys.*, 15(21), 12115–12138, doi:10.5194/acp-15-12115-2015, 2015.

Table 1. Over-limit ratio and concentration of MDA8 O₃ and PM_{2.5} calculated via ‘stations × days’ statistics in key urban clusters under four SWPs

Urban cluster	Month	Type1						Type2						Type3						Type4					
		Stas × days		MDA O ₃		PM _{2.5}		Stas × days		MDA O ₃		PM _{2.5}		Stas × days		MDA O ₃		PM _{2.5}		Stas × days		MDA O ₃		PM _{2.5}	
				OLR	Con	OLR	Con			OLR	Con	OLR	Con			OLR	Con	OLR	Con			OLR	Con	OLR	Con
		OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con	OLR	Con
BTH	6	6416	57.3%	172.5	16.7%	49.2	122	91.8%	209.6	3.3%	46.0	59	62.7%	176.0	32.2%	67.1	0	0	0	0	0	0	0	0	
	7	1716	25.6%	134.2	21.2%	51.8	3681	54.2%	165.6	22.5%	56.9	1356	33.8%	141.8	12.9%	46.7	577	43.5%	153.7	8.8%	49.5	49.5	49.5		
	8	1188	18.9%	118.5	6.9%	33.1	2805	35.3%	143.1	14.7%	44.0	1671	18.9%	121.2	10.1%	42.5	1268	31.9%	144.3	6.0%	39.7	39.7	39.7		
	6-8	9320	46.6%	158.6	16.3%	47.6	365	54.2%	169.8	18.8%	51.2	3086	26.3%	131.3	11.7%	44.8	1845	35.5%	147.3	6.9%	42.8	42.8	42.8		
YRD	6	19329	26.2%	127.2	4.4%	36.8	11098	14.3%	107.0	2.2%	44.5	181	12.7%	102.4	0.6%	32.0	0	0	0	0	0	0	0		
	7	5207	24.5%	119.8	5.3%	38.6	8459	22.0%	120.1	0.6%	28.0	4135	14.5%	112.6	0.1%	24.0	1743	30.9%	139.5	0.3%	30.0	30.0	30.0		
	8	3593	25.5%	127.6	0.3%	28.6	19922	18.3%	113.8	3.0%	33.3	4993	16.9%	116.7	0.0%	23.7	3817	14.7%	114.8	0.1%	25.1	25.1	25.1		
	6-8	28129	25.8%	125.9	4.0%	36.1	13754	17.4%	112.5	1.7%	30.6	9309	15.8%	114.6	0.1%	24.0	5560	19.7%	122.5	0.1%	26.7	26.7	26.7		
PRD	6	5327	5.2%	80.1	0	16.8	101	31.7%	146.9	0	24.9	48	0	77.6	0	13.2	0	0	0	0	0	0	0		
	7	1431	20.3%	107.2	0.6%	25.3	3076	7.6%	81.8	0.1%	19.1	1079	13.3%	92.4	0	19.3	461	8.0%	82.1	0	17.9	17.9	17.9		
	8	977	25.4%	122.5	1.1%	27.9	2316	20.6%	108.5	0.6%	28.0	1376	15.5%	103.4	0.3%	24.0	993	17.9%	111.4	0.1%	25.1	25.1	25.1		
	6-8	7735	10.5%	90.5	0.2%	19.8	5493	13.5%	94.3	0.3%	23.0	2503	14.3%	98.2	0.2%	21.8	1454	14.8%	102.1	0.1%	22.8	22.8	22.8		
GZP	6	1549	42.9%	147.6	0.4%	33.2	38	52.6%	165.3	0	30.4	19	26.3%	150.8	0	34.7	0	0	0	0	0	0	0		
	7	1879	40.7%	145.0	1.9%	33.7	1168	47.4%	160.3	0	30.2	432	24.3%	133.3	1.2%	31.7	173	26.6%	144.1	0	26.4	26.4	26.4		
	8	537	28.5%	134.0	1.6%	31.6	850	36.8%	146.3	1.6%	34.2	530	28.9%	143.1	1.9%	37.2	394	38.6%	154.9	1.3%	39.9	39.9	39.9		
	6-8	368	15.2%	112.9	0.9%	32.4	2056	43.1%	154.6	2.1%	35.9	981	26.8%	138.9	1.5%	34.7	567	34.9%	151.6	0.9%	35.8	35.8	35.8		
NEM	6	13126	17.9%	121.0	1.7%	26.4	243	64.2%	175.3	0.4%	31.1	120	13.3%	123.3	0.8%	37.6	0	0	0	0	0	0	0		
	7	3422	8.9%	106.8	2.3%	25.2	7338	17.0%	120.7	1.3%	28.1	2722	5.7%	93.0	0.7%	20.8	1141	15.1%	122.4	0.2%	24.3	24.3	24.3		
	8	2341	4.4%	92.6	1.0%	20.1	5520	6.3%	91.0	1.3%	21.0	3282	5.6%	98.3	0.9%	23.2	2507	7.8%	97.8	0.2%	19.5	19.5	19.5		
	6-8	18889	14.6%	114.9	1.7%	25.4	13101	13.4%	109.2	1.3%	25.2	6124	5.8%	96.4	0.8%	22.4	3648	10.1%	105.5	0.2%	21.0	21.0	21.0		

Notes: stas × days, stations × days; OLR, Over-limit ratio; Con, Concentration (μg m⁻³).




Fig. 1. Average concentration of MDA8 O₃ (a) and PM_{2.5} (b) in eastern China during the summers of 2015–2018. Stations and key urban clusters (black boxes) are shown in (a).




Fig. 2. Time series of MDA8 O₃ pollution levels in key urban clusters.




Fig.3. Time series of PM_{2.5} pollution levels in key urban clusters. Black dots indicate the compound events. Asterisks indicate the compound events under the Chinese standard (WHO interim target 1, IT-1), and the circles indicate the compound events under WHO IT-2.

810




Fig. 4. 850 hPa water vapor flux ($WVF = V * q/g$, where q is specific humidity, g is gravitational acceleration, V is horizontal wind; vectors; see scale arrow in the bottom right in units of $5 \text{ g cm}^{-1} \text{ hPa}^{-1} \text{ s}^{-1}$) and 500-hPa GH (contours; see scale bar at bottom in units of gpm) patterns based on objective classification (see text for details). The white framed area is the area of eastern China, the number in the upper-right corner of each panel indicates the frequency of occurrence of each pattern type, and the black line in each panel presents the ridge axis of the WPSH.




Fig. 5. Time series of synoptic circulation patterns.

825




Fig. 6. The MDA8 O₃ (a-d) and PM_{2.5} (e-h) anomaly under the four SWPs, where the sites marked with a '+' indicates the Analysis of Variance passes the significance level of 0.05.




Fig. 7. Daily variations of O_3 and $PM_{2.5}$ anomalies under the four SWPs in key urban clusters.




Fig. 8. As in Fig. 6 but for Tmax (a–d), RH (e–h), and PF (i–l). The black solid line presents the rain belt of each SWP.




Fig. 9. As in Fig. 6 but for the BLH at 1400 BJT (a–d) and FLWD (e–h).




Fig. 10. Daily variations of NO₂ (a–e) and Ox (f–j) under four SWPs in key urban clusters.




Fig. 11. Daily variations of horizontal wind, potential temperature and BLH in the BTH area during clean and compound pollution periods under Type 1 and Type 2 (a, b, e, f). The vertical cross-section of u-wind, w-wind and potential temperature for the same situation in the BTH region (c, d, g, h). The w-wind is multiplied by 100 when used. The data are from ERA5 reanalysis.




Fig. 12. Schematic diagrams describing the relationships between the WPSH, four SWPs and summertime O_3 and $PM_{2.5}$ pollution in various regions.