



Large contribution of organics to condensational growth and formation of cloud condensation nuclei (CCN) in remote marine boundary layer

5 Guangjie Zheng^{1,2}, Chongai Kuang², Janek Uin², Thomas Watson², and Jian Wang^{1,2*}

¹ Center for Aerosol Science and Engineering, Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, Missouri, USA

² Environmental and Climate Science Department, Brookhaven National Laboratory, Upton, New York, USA

10 *Correspondence to:* J.W. (jian@wustl.edu)

Abstract.

Marine low clouds strongly influence global climate, and their radiative effects are particularly susceptible to the concentration of cloud condensation nuclei (CCN). One major source of CCN is condensational growth of pre-CCN particles, and sulfate has long been considered the major condensing species in remote marine boundary layer. While some studies suggested that secondary organic species can contribute to the particle growth, its importance remains unclear. Here we present the first long-term observational evidence that organics play an important role in particle growth over remote oceans. To the contrary of traditional thinking, sulfate dominated condensational growth for only a small (~18%) fraction of the 62 observed growth events, even fewer than the organic-dominated events (24%). During most (58%) growth events, the major condensing species included both organics and sulfate. Potential precursors of the secondary organics are volatile organic compounds from ocean biological activities and those produced by the air-sea interfacial oxidation. Our results indicate that the condensation of secondary organics contributes strongly to the growth of pre-CCN particles, and thereby the CCN population over remote oceans.

15
20



1. Introduction

25 Marine low clouds play an important role in global climate system (Wood, 2012), and their properties and radiative effects are very sensitive to the concentration of cloud condensation nuclei (CCN) (Carslaw et al., 2013; Rosenfeld et al., 2019). Condensational growth of pre-CCN particles (i.e., particles that are too small to form cloud droplets) (Hoppel et al., 1990; Pierce and Adams, 2006) is one major source of CCN in remote marine boundary layer (MBL) (Pierce and Adams, 2006; Yu and Luo, 2009; Sanchez et al., 2018), and is likely the dominant one in late-spring to fall (Zheng et al., 2018). Over open ocean, 30 dimethyl sulfide (DMS) is the dominant biogenic volatile organic compound (VOC). The major oxidation products of DMS are sulfur dioxide (SO₂) and methanesulfonic acid (MSA) (Andreae et al., 1985). Further oxidation of SO₂ produces sulfuric acid (H₂SO₄), which readily condenses onto existing particles and participates in the formation of new particles (Kulmala et al., 2000). It has long been recognized that sulfate produced from DMS oxidation is a major species for particle condensational growth in the remote marine environment (Sanchez et al., 2018). Earlier studies (Willis et al., 2016; Kerminen and Wexler, 35 1997; Karl et al., 2011) suggest that MSA may also contribute to the growth of pre-CCN particles and thus the formation of CCN. However, the effect of MSA condensation on marine CCN concentration remains unclear. Model simulated effects range from negligible (e.g., a few percent) to significant (~20%) depending on the assumption of MSA volatility and the geographic location (Hodshire et al., 2019).

40 It has been suggested that in the remote MBL, secondary organics produced from two types of non-DMS VOCs can contribute substantially to particle condensational growth. The first type of VOCs, including isoprene, monoterpenes, and aliphatic amines (Facchini et al., 2008; Dall'Osto et al., 2012; Willis et al., 2017), is related to ocean biological activities, and SOA produced from these VOCs are positively correlated with MSA (Dall'Osto et al., 2012; Willis et al., 2016; Kim et al., 2017; Willis et al., 2017). While the mixing ratios of isoprene and monoterpenes are typically quite low over open oceans (Hu et al., 45 2013) due to their weak emissions, on rare occasions, elevated monoterpene mixing ratios up to ~100 ppt were observed (Kim et al., 2017), possibly due to enhanced microorganism growth as a result of nutrient replenishment (Kim et al., 2017). The second type of VOCs are produced by the oxidation reactions at the air-sea interface, especially when the sea surface microlayer is enriched in organic surfactants (Mungall et al., 2017; Brüggemann et al., 2018). These water-soluble organics can come from phytoplankton, but can also be from other sources, including other autotrophs and atmospheric depositions 50 (Wurl et al., 2011). Therefore, this type of oceanic VOCs and thus SOA formed may not correlate with MSA (Wurl et al., 2011; Mungall et al., 2017; Brüggemann et al., 2018).

At present, the contribution of secondary organics to the growth of pre-CCN particles in the MBL and the seasonal variation of this contribution remain unclear, largely due to the scarcity of the pre-CCN particle composition measurements. Existing 55 studies of pre-CCN growth in the MBL were typically within relatively short time periods (i.e., about 1-month) (Dall'Osto et al., 2012; Willis et al., 2016; Kim et al., 2017; Mungall et al., 2017; Willis et al., 2017; Vaattovaara et al., 2006; Modini et al.,



2009; Bzdek et al., 2014; Lawler et al., 2014; Swan et al., 2016), and were often conducted in coastal regions (Vaattovaara et al., 2006; Modini et al., 2009; Dall'Osto et al., 2012; Bzdek et al., 2014; Lawler et al., 2014; Swan et al., 2016) with substantial influences from continental emissions. Here we present the first long-term observational constraint on the importance of secondary organics to the growth of pre-CCN particles in remote MBL. Hygroscopicity of size-classified particles was characterized over a period of 14 months in the Eastern North Atlantic. By taking advantage of the contrasting hygroscopicity values of sulfate, MSA, and other secondary organic species, we constrain and identify the major species that are responsible for the growth of pre-CCN particles. Our results show that the organics represent an important or even the dominant condensing species during ~80% of growth events.

65 2. Measurements and datasets

The Eastern North Atlantic (ENA) atmospheric observatory was established by the Atmospheric Radiation Measurement (ARM) Climate Research Facility (<https://www.arm.gov/capabilities/observatories/ena>) in October 2013. This remote oceanic site, located on Graciosa Island, Azores, Portugal (39° 5' 30" N, 28° 1' 32" W, 30.48 m above mean sea level) (Mather and Voyles, 2013) straddles the boundary between the subtropics and mid-latitudes in the eastern North Atlantic. The ENA is a region of persistent but diverse marine low clouds, the albedo and precipitation of which are highly susceptible to perturbations of aerosol properties (Wood, 2012; Carslaw et al., 2013). Air masses arriving at this site can originate from North America, Northern Europe, the Arctic, and the Atlantic (Wood et al., 2015; Wang et al., 2016; Zheng et al., 2018). The routine measurements at the ENA site include meteorological parameters, trace gases mixing ratios, and aerosol and cloud properties (Zheng et al., 2018). The relevant routine measurements used in this study are summarized in section 2.3.

75 From June 2017 to Aug. 2018, the Aerosol and Cloud Experiments in the Eastern North Atlantic (ACE-ENA) campaign (Wang et al., 2016) was conducted in the Azores to investigate the aerosol-cloud interactions in the remote marine boundary layer (MBL). As a key part of this campaign, additional aerosol measurements were carried out at the ENA site, including aerosol size distribution and size-resolved CCN activated fractions (Mei et al., 2013c; Thalman et al., 2017). The instruments and calibration procedures are detailed elsewhere (Zheng et al., 2020), and are briefly described below. The data from these measurements are available at <https://www.arm.gov/research/campaigns/aaf2017ace-ena>.

2.1 Size distribution measurements and mode fittings

Aerosol size distribution was measured by a scanning mobility particle analyzer (SMPS, Model 3938, TSI Incorporated, Shoreview, MN, USA). Dry (RH < 25%) aerosol number size distribution ranging from 10 to 470 nm in particle diameter was measured every 8 minutes. In addition, a condensation particle counter (CPC, Model 3772, TSI Incorporated, Shoreview, MN, USA) was operated side-by-side to measure the total aerosol number concentrations (CN) concurrently. The measured aerosol number size distributions are fitted as a sum of up to three lognormal modes. Based on the fitted mode geometric mean



90 diameters ($D_{p,n}$), the fitted modes are classified as the nucleation mode ($D_{p,n} < 20$ nm), the Aitken mode ($20 < D_{p,n} < \sim 80$ nm), the accumulation mode ($\sim 80 < D_{p,n} < \sim 300$ nm), and the sea spray aerosol mode ($D_{p,n} > \sim 300$ nm) (Quinn et al., 2017; Zheng et al., 2018).

2.2 Size-resolved CCN activated fraction measurements

The size-resolved CCN measurement system (SCCN) consists of a Differential Mobility Analyzer (DMA, TSI Inc., Model 3081) coupled to a CPC (TSI Inc., Model 3010) and a cloud condensation nuclei counter (CCNC, Droplet Measurement Technologies, Boulder, CO) (Frank et al., 2006; Moore et al., 2010; Petters et al., 2007; Mei et al., 2013b). This system
95 measures the activated fraction (i.e., the fraction of particles that activate and form cloud droplets) of size-classified particles as a function of super-saturation (Thalman et al., 2017). During the ACE-ENA campaign, the DMA stepped through 6 dry particle diameters ($D_{p,SCCN}$) of 40, 50, 75, 100, 125, and 150 nm. At each $D_{p,SCCN}$, the super-saturation level inside the CCNC was varied by changing the flow rate and/or temperature gradient ΔT . The corresponding supersaturation levels, ranging from 0.07% to 1.34% at 298 K, were calibrated using ammonium sulfate particles following established procedures (Lance et al.,
100 2013; Mei et al., 2013a; Thalman et al., 2017). An entire measurement cycle through the 6 particle diameters took between 1~2 h, depending on particle number concentration. Temperature dependence of CCNC supersaturation (Rose et al., 2008; Thalman et al., 2017) and the effect of multi-charged particles (Thalman et al., 2017) are taken into consideration. The particle hygroscopicity parameter under supersaturated conditions, κ_{CCN} (Petters and Kreidenweis, 2007), is derived from the activated fraction spectrum and the corresponding particle diameter (Lance et al., 2013; Mei et al., 2013a; Thalman et al., 2017).

105 2.3 Other relevant datasets used in this study

Routine measurements at the ENA site used in this study include the non-refractory submicron aerosol (NR-PM₁) composition (organics, sulfate, nitrate, ammonium, and chloride) characterized by an Aerosol Chemical Speciation Monitor (ACSM; Aerodyne Research, Inc., Billerica, MA, USA) (Watson, 2017) and particle hygroscopic growth measured by a Humidified Tandem Differential Mobility Analyzer (HTDMA, Brechtel Manufacturing Inc., CA, USA) (Uin, 2016). The HTDMA
110 measures aerosol hygroscopic growth factor under $\sim 80\%$ RH at 5 particle diameters (50, 100, 150, 200 and 250 nm), from which the aerosol hygroscopicity under sub-saturated conditions (κ_{GF}) is derived (Petters and Kreidenweis, 2007).

Gas-phase SO₂ and MSA concentrations are from the Modern-Era Retrospective Analysis for Research and Applications, version 2 (MERRA-2) reanalysis data (Gelaro et al., 2017), at the grid corresponding to the ENA site.

115 3. Derivation of the hygroscopicity parameter of condensing species

Continuous growth of Aitken mode particles is identified from the aerosol size distribution time series. As a result of the condensational growth, aerosol chemical compositions and thus the hygroscopicity of Aitken mode particles are expected to



120 evolve with time during the growth events. As potential condensing species (Table S1) have contrasting hygroscopicity parameters, the variation of hygroscopicity parameter during the growth events can therefore be used to infer the major condensing species.

3.1 Matching aerosol size modes with the hygroscopicity measurements

Here we detail the procedure to correlate the aerosol size distribution with the SCCN measurements. The same procedure is also applied to correlate the aerosol size distribution with the HTDMA measurements. The CCN activated fraction spectrum was measured at 6 fixed sizes ($D_{p, SCCN}$). As the size of Aitken mode particles evolves continuously during the growth events, we first determine if the hygroscopicity of the growing Aitken mode can be captured by the measurement at one of the six $D_{p, SCCN}$ using the following two criteria. The first criterion is that $D_{p, SCCN}$ (e.g., 40 nm, 50nm, or 75 nm) is within one geometric standard deviation (σ) of the Aitken mode diameter, i.e., $D_{p, n} \sigma^{-1} < D_{p, SCCN} < D_{p, n} \sigma$ (Fig. 1a). For example, at time t_0 , $D_{p, SCCN}$ (40 nm) is within one σ range of the Aitken mode diameter (i.e., dark blue shaded area in Fig. 1a), and the κ_{CCN} value measured at 40 nm is considered representative of the Aitken mode (solid blue curve in Fig. 1a). In contrast, at a later time t_1 , the Aitken mode grew to larger sizes (dash blue curve in Fig. 1a), and 40 nm became smaller than $D_{p, n} \sigma^{-1}$ (light blue shaded area in Fig. 1a). Therefore, κ_{CCN} measured at 40 nm no longer represents the hygroscopicity of the Aitken mode at t_1 . The second criterion is that particle concentration at $D_{p, SCCN}$ is dominated by the Aitken mode only (Fig. 1b), i.e., over 95% of the particles at the measured $D_{p, SCCN}$ is contributed by the Aitken mode. As an example, both the Aitken mode (blue curve) and the accumulation mode (red curve) contribute to the number size distribution at $D_{p, SCCN}$ (black dash line, Fig. 1b). Although $D_{p, SCCN}$ is within one σ of the Aitken mode diameter, the contribution of Aitken mode is less than 95% at this size (orange curve in Fig. 1b). Therefore, measurement at $D_{p, SCCN}$ is not deemed as representative of the Aitken mode due to the substantial contribution from accumulation mode particles. Only data points that meet both criteria are selected, as illustrated in Fig. 1c. Figure 2a gives an example of the time series of Aitken mode diameter and paired κ_{CCN} value during a growth event.

3.2 Derivation of the hygroscopicity of condensed species (κ_c) during growth events

140 The derivation is applied to condensational growth events when there are sufficient number (> 6 points) of κ measurements that satisfy both criteria described in M2.1. For each condensational growth event selected (e.g., Fig. 2a), the average hygroscopicity parameter of condensing species, κ_c , is derived based on the following three assumptions. Here, κ represents either the hygroscopicity derived from SCCN (i.e., κ_{CCN}) or HTDMA (i.e., κ_{GF}) data.

145 The first assumption is that the change in particle volume (diameter) is due to the condensational growth only, namely:

$$V_c = \Delta V = V_1 - V_0 = (\pi/6) D_{p1}^3 - (\pi/6) D_{p0}^3 \quad (1)$$

where V is the particle volume and D_p is the particle diameter. Hereinafter we use X_1 and X_0 to denote the corresponding particle property X after and before the condensational growth, respectively, and X_c refers to the property X of the condensed species. The second assumption is that the aerosol κ follows the volume-weighted mixing law (Petters and Kreidenweis, 2007):



150
$$\kappa_1 = \kappa_0 (V_0 / V_1) + \kappa_c (V_c / V_1) \quad (2)$$

The third assumption is that the growth rate is identical for particles of the same size, and thus the relative position of any given particle in the accumulative size distribution is maintained throughout the growth. Let CDF_0 and CDF_1 denote the particle cumulative size distributions before and after the particle growth, and D_{p0} and D_{p1} represent particle diameters before and after particle growth, respectively. The number of particles smaller than D_{p1} following particle growth should be the same as the
155 number of particles smaller than D_{p0} prior to the growth event (Fig. 2b):

$$CDF_1(D_{p1}) = CDF_0(D_{p0}) \quad (3)$$

For each particle size (i.e., D_{p1}) measured during the growth events, the original particle size (i.e., D_{p0}) is derived from Eq. (3).

The volume fraction of condensed species, $f_{V, \text{cond}}$, is given by:

$$f_{V, \text{cond}} = V_c / V_1 = 1 - V_0 / V_1 = 1 - (D_{p0} / D_{p1})^3 \quad (4)$$

160 By combining Eq. 1-4, we have:

$$\kappa_1 = (\kappa_c - \kappa_0) f_{V, \text{cond}} + \kappa_0 \quad (5)$$

Both κ_1 and $f_{V, \text{cond}}$ are from the measurements as described above. Therefore, κ_c and κ_0 can be derived from the linear fitting of κ_1 vs. $f_{V, \text{cond}}$ for each growth event (e.g., Fig. 2c), where κ_0 is the intercept, and κ_c is the sum of slope and intercept. The method described here was applied to both SCCN and HTDMA measurements, and κ_c derived are referred to as $\kappa_{c, \text{CCN}}$ and $\kappa_{c, \text{GF}}$
165 hereinafter, respectively.

4. Constraining the major condensing species in remote MBL

Figure 3 shows two examples of the identified growth events, with the dominant condensing species being sulfate and organics, respectively. While the measured κ_{CCN} of the Aitken mode particles (i.e., pre-CCN particles that are below ~ 80 nm) are similar (~0.45) at the start of both events, the variations of κ_{CCN} with growing particle size show opposite trends. For the July case
170 (Fig. 3a,b), κ_{CCN} increased with the volume fraction of condensed species ($f_{V, \text{cond}}$, Fig. 3b), indicating that the hygroscopicity of the condensed species, $\kappa_{c, \text{CCN}}$, exceeds that of the original particles. The derived $\kappa_{c, \text{CCN}}$ value is 0.7, which is typical of sulfates (Table S1). In contrast, during the September growth event (Fig. 3c,d), κ_{CCN} decreased as the particles grew. The derived $\kappa_{c, \text{CCN}}$ value is ~0.3, indicating organics as the dominant condensing species. We note that $\kappa_{c, \text{CCN}}$ is derived from the volume-weighted mixing law (Petters and Kreidenweis, 2007) (i.e., ideal Zdanovskii, Stokes, and Robinson (ZSR) mixing).
175 Organic surfactants may facilitate CCN activation by lowering surface tension of growing droplets (Ovadnevaite et al., 2017). In scenarios when particles contain organic surfactants, particle hygroscopicity κ_{CCN} may be greater than the simple volume average of participating species. As a result, the derived $\kappa_{c, \text{CCN}}$ value based on the volume-weighted mixing law may be overestimated, therefore leading to an underestimation of the contribution of organics to the particle condensational growth.

180 A total of 62 growth events are identified during the 14-month campaign (Fig. 4). These events are classified into 3 categories according to the derived $\kappa_{c, \text{CCN}}$ value (Table S1): (1) low hygroscopicity (i.e., $\kappa_{c, \text{CCN}} < 0.45$) indicating organics as the dominant



condensing species, (2) high hygroscopicity (i.e., $\kappa_{c,CCN} > 0.65$) with acidic sulfate (i.e., H_2SO_4 or NH_4HSO_4) dominating the particle condensational growth, and (3) intermediate hygroscopicity value (i.e., $0.45 < \kappa_{c,CCN} < 0.65$), when $(NH_4)_2SO_4$ and/or mixtures of organics and acidic sulfate contribute to the particle growth.

185 5. Monthly distributions of the dominant condensing species

The monthly distribution of the identified growth events and the dominant condensing species are shown in Fig. 4. Relatively more events were observed during the summer seasons due to favorable synoptic conditions. In summer, there is a stronger influence by the Azores High while the influence from mid-latitude cyclones and the corresponding wet scavenging are much weaker (Zheng et al., 2018). The distribution of the event categories shows that, contrary to the conventional thinking, 190 NH_4HSO_4/H_2SO_4 dominated the condensational growth during only 18% of the growth events. This is less than the events dominated by organics at 24%. The majority (58%) of the growth events exhibit intermediate $\kappa_{c,CCN}$ values, suggesting that $(NH_4)_2SO_4$ or a mixture of organics and sulfate are responsible for the particle condensational growth.

To further constrain the condensing species for the intermediate $\kappa_{c,CCN}$ category, we compare the $\kappa_{c,CCN}$ value with the 195 hygroscopicity under sub-saturated conditions ($\kappa_{c,GF}$), which is derived from measured particle hygroscopic growth (section 3). For $(NH_4)_2SO_4$, the difference between $\kappa_{c,CCN}$ and $\kappa_{c,GF}$ is relatively small (within 20%) (Petters and Kreidenweis, 2007), while the difference is usually substantially larger (Wex et al., 2009; Rastak et al., 2017; Petters et al., 2009; Pajunoja et al., 2015; Ovadnevaite et al., 2011; Massoli et al., 2010) for organic species. The large difference has been attributed to the solution non-ideality (Petters et al., 2009), the formation of hydrogels (Ovadnevaite et al., 2011), and the solubility and phase states 200 (Pajunoja et al., 2015; Rastak et al., 2017). One example of the intermediate $\kappa_{c,CCN}$ category is shown in Fig. S1. For this case, the derived $\kappa_{c,CCN}$ and $\kappa_{c,GF}$ values are 0.59 and 0.45, respectively (Fig. S1). The difference is close to the measurement uncertainty (i.e., 20%), and therefore the major condensing species for this example is classified as $(NH_4)_2SO_4$.

Figure 5 compares the values of $\kappa_{c,CCN}$ and $\kappa_{c,GF}$ for all available events in the intermediate $\kappa_{c,CCN}$ category. For most of these 205 events, $\kappa_{c,GF}$ is at least 20% lower than $\kappa_{c,CCN}$, indicating organics likely played an important role in particle condensational growth. In addition, chemical composition of sub-micron non-refractory aerosol (NR-PM₁; aerodynamic diameters below 1 μm) indicates an ammonium-poor condition over the ENA (color bar in Fig. 5), typical of remote marine environment (Adams et al., 1999). Therefore, sulfate is not fully neutralized as $(NH_4)_2SO_4$. These evidences suggest that during most of the intermediate- $\kappa_{c,CCN}$ events, the condensed species are a mixture of sulfates and organics instead of dominated by $(NH_4)_2SO_4$. 210 Based on a κ_{CCN} value of 0.9 for acidic sulfates (H_2SO_4 and/or NH_4HSO_4 , Table S1), the average contribution of organics during the intermediate- $\kappa_{c,CCN}$ events ranges from 42% and 63%, depending on the κ_{CCN} values of organics assumed (0.1~0.36; Table S1). Therefore, organics played an important role during the intermediate- $\kappa_{c,CCN}$ events and dominated the particle



condensational growth for the low- $\kappa_{c,CCN}$ category. Together, these two categories represent a total of ~80% of the growth events and occurred throughout the year.

215 6. Sources of the condensing organics

Given the importance of secondary organics to particle condensational growth, the potential sources of the condensing organics are investigated by examining the air mass origins (SI S1). Here we classify the origin of air mass during the growth events into four types: (1) continental air masses from North America or Europe, (2) the Arctic, (3) the subtropical, and (4) the mid-latitude Atlantic. Note that an air mass is denoted as continental if it passed over the North America or Europe, so the non-
220 continental types represent the air masses that had stayed over oceans or clean continental areas (i.e., Arctic region) for at least 10 days (SI S1).

Growth events of mid-latitude Atlantic or Arctic type were observed exclusively from May to September, a period that coincides with the phytoplankton blooms in mid-latitude Atlantic or Arctic, but not the subtropics (Sapiano et al., 2012). For
225 these events, $\kappa_{c,CCN}$ is anti-correlated with MSA/SO₂ ratio (Fig. 6a), which is from MERRA-2 reanalysis data (section 2.3). As fixed yields of SO₂ and MSA from DMS oxidation are assumed in MERRA-2 data (Chin et al., 2000; Randles et al., 2017), a lower MSA/SO₂ ratio suggests other SO₂ sources in addition to DMS oxidation contribute to these events. These other sources could include volcanic emissions and combustion products from international shipping (Randles et al., 2017). As MSA is a
230 the condensed organics are likely SOA produced from VOCs emitted from ocean biological activities (e.g., phytoplankton blooms). The value of $\kappa_{c,CCN}$ is not correlated with the NR-PM₁ organic/sulfate ratio (Fig. 6b), suggesting different sources of the condensed species in pre-CCN and the accumulation mode particle composition.

Among the remaining growth events, only four of them are subtropical cases, which occurred outside the bloom periods.
235 During the other events, air masses were potentially influenced by continental emissions (Fig. S2). For these events, $\kappa_{c,CCN}$ is instead positively correlated with MSA/SO₂ ratio (Fig. 6c), indicating that secondary organics formed from phytoplankton-emitted VOCs likely played a minor role in the observed particle condensational growth. The $\kappa_{c,CCN}$ value generally decreases with increasing NR-PM₁ organic / sulfate ratios (Fig. 6d), suggesting that the formation of SOA led to increased organic fraction for both pre-CCN and accumulation mode particles. Possible sources of the condensed organics include SOA
240 generated from long-range transported continental VOCs and VOCs released by the sea-surface microlayer oxidation that are not directly related to phytoplankton emissions.

As continentally emitted VOCs are removed by oxidation during long-range transport, it is expected that in-situ SOA production from these VOCs is low and plays a minor role in particle condensational growth over the remote oceans (Kelly et



245 al., 2019; D'Andrea et al., 2013). On the other hand, aromatic compounds were detected in pre-CCN particles in clean air
masses at a coastal site (Lawler et al., 2014), indicating potential contribution of SOA from anthropogenic VOCs with long
lifetime. Oxidation reactions at the air-sea interface can produce VOCs, which lead to subsequent SOA formation (Mungall et
al., 2017; Brüggemann et al., 2018). This VOC source is present all-year round, even during winter when there is little
biological activity in the ocean (Brüggemann et al., 2018). Therefore, the secondary organics produced via this pathway can
250 contribute to the growth of pre-CCN particles outside the biologically active seasons of the ocean.

7. Conclusions

In summary, we show that during all seasons, secondary organics play an important role in the condensational growth of pre-
CCN particles, and by extension, the formation of CCN in the remote marine boundary layer. The secondary organic species
likely derive from a variety of precursors, including VOCs produced from marine biogenic activity, continentally emitted
255 VOCs with long lifetime that survive the long-range transport, and VOCs formed by oxidation at the air-sea interface. Current
global models typically assume that sulfates dominate the particle growth over remote oceans, and therefore may substantially
underestimate the formation of CCN by condensational growth in remote marine boundary layer.

260

Data availability. All data used in this study are available at <https://www.arm.gov/research/campaigns/aaf2017ace-ena> and
<https://www-air.larc.nasa.gov/missions/naames/index.html>.

Author contributions. J.W. and G.Z. designed the study. J.W. G.Z., C.K., J.U., and T.W carried out the measurements, G.Z.
265 and J.W. conducted the analysis and wrote the manuscript with contributions from all authors.

Competing interests. The authors declare no competing interests.

Acknowledgments. The research was conducted with funding from the Atmospheric System Research (ASR) program (Award
270 No. DE-SC0020259), Office of Biological and Environmental Research (OBER) of the United States Department of Energy.
We acknowledge additional support by the Atmospheric Radiation Measurement (ARM) Climate Research Facility, a user
facility of the United States Department of Energy, Office of Science, sponsored by the Office of Biological and Environmental
Research.



275 References

- Adams, P. J., Seinfeld, J. H., and Koch, D. M.: Global concentrations of tropospheric sulfate, nitrate, and ammonium aerosol simulated in a general circulation model, *Journal of Geophysical Research: Atmospheres*, 104, 13791-13823, 10.1029/1999jd900083, 1999.
- Andreae, M. O., Ferek, R. J., Bermond, F., Byrd, K. P., Engstrom, R. T., Hardin, S., Houmère, P. D., LeMarrec, F., Raemdonck, H., and Chatfield, R. B.: Dimethyl sulfide in the marine atmosphere, *Journal of Geophysical Research: Atmospheres*, 90, 12891-12900, 10.1029/JD090iD07p12891, 1985.
- Brüggemann, M., Hayeck, N., and George, C.: Interfacial photochemistry at the ocean surface is a global source of organic vapors and aerosols, *Nature Communications*, 9, 2101, 10.1038/s41467-018-04528-7, 2018.
- Bzdek, B. R., Lawler, M. J., Horan, A. J., Pennington, M. R., DePalma, J. W., Zhao, J., Smith, J. N., and Johnston, M. V.: Molecular constraints on particle growth during new particle formation, *Geophysical Research Letters*, 41, 6045-6054, 10.1002/2014gl060160, 2014.
- Carslaw, K., Lee, L., Reddington, C., Pringle, K., Rap, A., Forster, P., Mann, G., Spracklen, D., Woodhouse, M., and Regayre, L.: Large contribution of natural aerosols to uncertainty in indirect forcing, *Nature*, 503, 67, 2013.
- Chang, R. Y. W., Slowik, J. G., Shantz, N. C., Vlasenko, A., Liggio, J., Sjostedt, S. J., Leaitch, W. R., and Abbatt, J. P. D.: The hygroscopicity parameter (κ) of ambient organic aerosol at a field site subject to biogenic and anthropogenic influences: relationship to degree of aerosol oxidation, *Atmos. Chem. Phys.*, 10, 5047-5064, 10.5194/acp-10-5047-2010, 2010.
- Chin, M., Rood, R. B., Lin, S.-J., Müller, J.-F., and Thompson, A. M.: Atmospheric sulfur cycle simulated in the global model GOCART: Model description and global properties, *Journal of Geophysical Research: Atmospheres*, 105, 24671-24687, 10.1029/2000jd900384, 2000.
- D'Andrea, S. D., Häkkinen, S. A. K., Westervelt, D. M., Kuang, C., Levin, E. J. T., Kanawade, V. P., Leaitch, W. R., Spracklen, D. V., Riipinen, I., and Pierce, J. R.: Understanding global secondary organic aerosol amount and size-resolved condensational behavior, *Atmos. Chem. Phys.*, 13, 11519-11534, 10.5194/acp-13-11519-2013, 2013.
- Dall'Osto, M., Ceburnis, D., Monahan, C., Worsnop, D. R., Bialek, J., Kulmala, M., Kurtén, T., Ehn, M., Wenger, J., Sodeau, J., Healy, R., and O'Dowd, C.: Nitrogenated and aliphatic organic vapors as possible drivers for marine secondary organic aerosol growth, *Journal of Geophysical Research: Atmospheres*, 117, 10.1029/2012jd017522, 2012.
- Facchini, M. C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Ceburnis, D., Flanagan, R., Nilsson, E. D., de Leeuw, G., Martino, M., Woeltjen, J., and O'Dowd, C. D.: Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates, *Geophysical Research Letters*, 35, n/a-n/a, 10.1029/2008GL034210, 2008.
- Frank, G. P., Dusek, U., and Andreae, M. O.: Technical note: A method for measuring size-resolved CCN in the atmosphere, *Atmos. Chem. Phys. Discuss.*, 6, 4879-4895, 10.5194/acpd-6-4879-2006, 2006.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Silva, A. M. d., Gu, W., Kim,



- G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419-5454, 10.1175/jcli-d-16-0758.1, 2017.
- Hodshire, A. L., Campuzano-Jost, P., Kodros, J. K., Croft, B., Nault, B. A., Schroder, J. C., Jimenez, J. L., and Pierce, J. R.: The potential role of methanesulfonic acid (MSA) in aerosol formation and growth and the associated radiative forcings, *Atmos. Chem. Phys.*, 19, 3137-3160, 10.5194/acp-19-3137-2019, 2019.
- Hoppel, W. A., Fitzgerald, J. W., Frick, G. M., Larson, R. E., and Mack, E. J.: Aerosol size distributions and optical properties found in the marine boundary layer over the Atlantic Ocean, *Journal of Geophysical Research: Atmospheres*, 95, 3659-3686, 10.1029/JD095iD04p03659, 1990.
- Hu, Q.-H., Xie, Z.-Q., Wang, X.-M., Kang, H., He, Q.-F., and Zhang, P.: Secondary organic aerosols over oceans via oxidation of isoprene and monoterpenes from Arctic to Antarctic, *Sci Rep-Uk*, 3, 2280, 10.1038/srep02280, 2013.
- Johnson, G. R., Ristovski, Z., and Morawska, L.: Method for measuring the hygroscopic behaviour of lower volatility fractions in an internally mixed aerosol, *Journal of Aerosol Science*, 35, 443-455, <https://doi.org/10.1016/j.jaerosci.2003.10.008>, 2004.
- Karl, M., Gross, A., Pirjola, L., and Leck, C.: A new flexible multicomponent model for the study of aerosol dynamics in the marine boundary layer, *Tellus B: Chemical and Physical Meteorology*, 63, 1001-1025, 10.1111/j.1600-0889.2011.00562.x, 2011.
- Kelly, J. M., Doherty, R. M., O'Connor, F. M., Mann, G. W., Coe, H., and Liu, D.: The roles of volatile organic compound deposition and oxidation mechanisms in determining secondary organic aerosol production: a global perspective using the UKCA chemistry-climate model (vn8.4), *Geosci. Model Dev.*, 12, 2539-2569, 10.5194/gmd-12-2539-2019, 2019.
- Kerminen, V.-M., and Wexler, A. S.: Growth behavior of the marine submicron boundary layer aerosol, *Journal of Geophysical Research: Atmospheres*, 102, 18813-18825, 10.1029/97jd01260, 1997.
- Kim, M. J., Novak, G. A., Zoerb, M. C., Yang, M., Blomquist, B. W., Huebert, B. J., Cappa, C. D., and Bertram, T. H.: Air-Sea exchange of biogenic volatile organic compounds and the impact on aerosol particle size distributions, *Geophysical Research Letters*, 44, 3887-3896, 10.1002/2017gl072975, 2017.
- Kulmala, M., Pirjola, L., and Mäkelä, J. M.: Stable sulphate clusters as a source of new atmospheric particles, *Nature*, 404, 66-69, 10.1038/35003550, 2000.
- Lance, S., Raatikainen, T., Onasch, T. B., Worsnop, D. R., Yu, X. Y., Alexander, M. L., Stolzenburg, M. R., McMurry, P. H., Smith, J. N., and Nenes, A.: Aerosol mixing state, hygroscopic growth and cloud activation efficiency during MIRAGE 2006, *Atmos. Chem. Phys.*, 13, 5049-5062, 10.5194/acp-13-5049-2013, 2013.
- Lawler, M. J., Whitehead, J., O'Dowd, C., Monahan, C., McFiggans, G., and Smith, J. N.: Composition of 15–85 nm particles in marine air, *Atmos. Chem. Phys.*, 14, 11557-11569, 10.5194/acp-14-11557-2014, 2014.
- Massoli, P., Lambe, A. T., Ahern, A. T., Williams, L. R., Ehn, M., Mikkilä, J., Canagaratna, M. R., Brune, W. H., Onasch, T. B., Jayne, J. T., Petäjä, T., Kulmala, M., Laaksonen, A., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Relationship between



- aerosol oxidation level and hygroscopic properties of laboratory generated secondary organic aerosol (SOA) particles, *Geophysical Research Letters*, 37, 10.1029/2010gl045258, 2010.
- Mather, J. H., and Voyles, J. W.: The Arm Climate Research Facility: A Review of Structure and Capabilities, *Bulletin of the American Meteorological Society*, 94, 377-392, 10.1175/bams-d-11-00218.1, 2013.
- 345 Mei, F., Hayes, P. L., Ortega, A., Taylor, J. W., Allan, J. D., Gilman, J., Kuster, W., de Gouw, J., Jimenez, J. L., and Wang, J.: Droplet activation properties of organic aerosols observed at an urban site during CalNex-LA, *Journal of Geophysical Research: Atmospheres*, 118, 2903-2917, 10.1002/jgrd.50285, 2013a.
- Mei, F., Hayes, P. L., Ortega, A. M., Taylor, J. W., Allan, J. D., Gilman, J. B., Kuster, W. C., de Gouw, J. A., Jimenez, J. L., and Wang, J.: Droplet activation properties of organic aerosols observed at an urban site during CalNex-LA, *Journal of*
- 350 *Geophysical Research*, 118, 2903-2917 10.1002/jgrd.50285, 2013b.
- Mei, F., Setyan, A., Zhang, Q., and Wang, J.: CCN activity of organic aerosols observed downwind of urban emissions during CARES, *Atmos. Chem. Phys.*, 13, 12155-12169, 10.5194/acp-13-12155-2013, 2013c.
- Modini, R. L., Ristovski, Z. D., Johnson, G. R., He, C., Surawski, N., Morawska, L., Suni, T., and Kulmala, M.: New particle formation and growth at a remote, sub-tropical coastal location, *Atmos. Chem. Phys.*, 9, 7607-7621, 10.5194/acp-9-7607-
- 355 2009, 2009.
- Moore, R. H., Nenes, A., and Medina, J.: Scanning Mobility CCN Analysis—A Method for Fast Measurements of Size-Resolved CCN Distributions and Activation Kinetics, *Aerosol Science and Technology*, 44, 861-871, 10.1080/02786826.2010.498715, 2010.
- Mungall, E. L., Abbatt, J. P. D., Wentzell, J. J. B., Lee, A. K. Y., Thomas, J. L., Blais, M., Gosselin, M., Miller, L. A.,
- 360 Papakyriakou, T., Willis, M. D., and Liggiio, J.: Microlayer source of oxygenated volatile organic compounds in the summertime marine Arctic boundary layer, *Proceedings of the National Academy of Sciences*, 114, 6203-6208, 10.1073/pnas.1620571114, 2017.
- Ovadnevaite, J., Ceburnis, D., Martucci, G., Bialek, J., Monahan, C., Rinaldi, M., Facchini, M. C., Berresheim, H., Worsnop, D. R., and O'Dowd, C.: Primary marine organic aerosol: A dichotomy of low hygroscopicity and high CCN activity,
- 365 *Geophysical Research Letters*, 38, n/a-n/a, 10.1029/2011GL048869, 2011.
- Ovadnevaite, J., Zuend, A., Laaksonen, A., Sanchez, K. J., Roberts, G., Ceburnis, D., Decesari, S., Rinaldi, M., Hodas, N., Facchini, M. C., Seinfeld, J. H., and O' Dowd, C.: Surface tension prevails over solute effect in organic-influenced cloud droplet activation, *Nature*, 546, 637-641, 10.1038/nature22806, 2017.
- Pajunoja, A., Lambe, A. T., Hakala, J., Rastak, N., Cummings, M. J., Brogan, J. F., Hao, L., Paramonov, M., Hong, J., Prisle,
- 370 N. L., Malila, J., Romakkaniemi, S., Lehtinen, K. E. J., Laaksonen, A., Kulmala, M., Massoli, P., Onasch, T. B., Donahue, N. M., Riipinen, I., Davidovits, P., Worsnop, D. R., Petäjä, T., and Virtanen, A.: Adsorptive uptake of water by semisolid secondary organic aerosols, *Geophysical Research Letters*, 42, 3063-3068, 10.1002/2015gl063142, 2015.
- Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, 10.5194/acp-7-1961-2007, 2007.



- 375 Petters, M. D., Prenni, A. J., Kreidenweis, S. M., and DeMott, P. J.: On Measuring the Critical Diameter of Cloud Condensation Nuclei Using Mobility Selected Aerosol, *Aerosol Science and Technology*, 41, 907-913, 10.1080/02786820701557214, 2007.
- Petters, M. D., Wex, H., Carrico, C. M., Hallbauer, E., Massling, A., McMeeking, G. R., Poulain, L., Wu, Z., Kreidenweis, S. M., and Stratmann, F.: Towards closing the gap between hygroscopic growth and activation for secondary organic aerosol – Part 2: Theoretical approaches, *Atmos. Chem. Phys.*, 9, 3999-4009, 10.5194/acp-9-3999-2009, 2009.
- 380 Pierce, J. R., and Adams, P. J.: Global evaluation of CCN formation by direct emission of sea salt and growth of ultrafine sea salt, *Journal of Geophysical Research: Atmospheres*, 111, 10.1029/2005jd006186, 2006.
- Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction of marine cloud condensation nuclei made up of sea spray aerosol, *Nature Geosci*, advance online publication, 10.1038/ngeo3003, 2017.
- Randles, C., Da Silva, A., Buchard, V., Colarco, P., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., and
385 Hair, J.: The MERRA-2 aerosol reanalysis, 1980 onward. Part I: System description and data assimilation evaluation, *Journal of Climate*, 30, 6823-6850, 2017.
- Rastak, N., Pajunoja, A., Acosta Navarro, J. C., Ma, J., Song, M., Partridge, D. G., Kirkevåg, A., Leong, Y., Hu, W. W., Taylor, N. F., Lambe, A., Cerully, K., Bougiatioti, A., Liu, P., Krejci, R., Petäjä, T., Percival, C., Davidovits, P., Worsnop, D. R., Ekman, A. M. L., Nenes, A., Martin, S., Jimenez, J. L., Collins, D. R., Topping, D. O., Bertram, A. K., Zuend, A., Virtanen,
390 A., and Riipinen, I.: Microphysical explanation of the RH-dependent water affinity of biogenic organic aerosol and its importance for climate, *Geophysical research letters*, 44, 5167-5177, 10.1002/2017GL073056, 2017.
- Rose, D., Gunthe, S. S., Mikhailov, E., Frank, G. P., Dusek, U., Andreae, M. O., and Pöschl, U.: Calibration and measurement uncertainties of a continuous-flow cloud condensation nuclei counter (DMT-CCNC): CCN activation of ammonium sulfate and sodium chloride aerosol particles in theory and experiment, *Atmos. Chem. Phys.*, 8, 1153-1179, 10.5194/acp-8-1153-
395 2008, 2008.
- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 363, eaav0566, 10.1126/science.aav0566, 2019.
- Sanchez, K. J., Chen, C.-L., Russell, L. M., Betha, R., Liu, J., Price, D. J., Massoli, P., Ziemba, L. D., Crosbie, E. C., Moore, R. H., Müller, M., Schiller, S. A., Wisthaler, A., Lee, A. K. Y., Quinn, P. K., Bates, T. S., Porter, J., Bell, T. G., Saltzman, E.,
400 S., Vaillancourt, R. D., and Behrenfeld, M. J.: Substantial Seasonal Contribution of Observed Biogenic Sulfate Particles to Cloud Condensation Nuclei, *Sci Rep-Uk*, 8, 3235, 10.1038/s41598-018-21590-9, 2018.
- Sapiano, M. R. P., Brown, C. W., Schollaert Uz, S., and Vargas, M.: Establishing a global climatology of marine phytoplankton phenological characteristics, *Journal of Geophysical Research: Oceans*, 117, 10.1029/2012jc007958, 2012.
- Schmale, J., Henning, S., Decesari, S., Henzing, B., Keskinen, H., Sellegri, K., Ovadnevaite, J., Pöhlker, M. L., Brito, J.,
405 Bougiatioti, A., Kristensson, A., Kalivitis, N., Stavroulas, I., Carbone, S., Jefferson, A., Park, M., Schlag, P., Iwamoto, Y., Aalto, P., Äijälä, M., Bukowiecki, N., Ehn, M., Frank, G., Fröhlich, R., Frumau, A., Herrmann, E., Herrmann, H., Holzinger, R., Kos, G., Kulmala, M., Mihalopoulos, N., Nenes, A., O'Dowd, C., Petäjä, T., Picard, D., Pöhlker, C., Pöschl, U., Poulain, L., Prévôt, A. S. H., Swietlicki, E., Andreae, M. O., Artaxo, P., Wiedensohler, A., Ogren, J., Matsuki, A., Yum, S. S.,



- Stratmann, F., Baltensperger, U., and Gysel, M.: Long-term cloud condensation nuclei number concentration, particle number
410 size distribution and chemical composition measurements at regionally representative observatories, *Atmos. Chem. Phys.*, 18,
2853-2881, 10.5194/acp-18-2853-2018, 2018.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley &
Sons, 2016.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric
415 Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 10.1175/bams-
d-14-00110.1, 2015.
- Swan, H. B., Crough, R. W., Vaattovaara, P., Jones, G. B., Deschaseaux, E. S. M., Eyre, B. D., Miljevic, B., and Ristovski, Z.
D.: Dimethyl sulfide and other biogenic volatile organic compound emissions from branching coral and reef seawater: potential
sources of secondary aerosol over the Great Barrier Reef, *Journal of Atmospheric Chemistry*, 73, 303-328, 10.1007/s10874-
420 016-9327-7, 2016.
- Tang, M., Guo, L., Bai, Y., Huang, R.-J., Wu, Z., Wang, Z., Zhang, G., Ding, X., Hu, M., and Wang, X.: Impacts of
methanesulfonate on the cloud condensation nucleation activity of sea salt aerosol, *Atmospheric Environment*, 201, 13-17,
<https://doi.org/10.1016/j.atmosenv.2018.12.034>, 2019.
- Thalman, R., de Sá, S. S., Palm, B. B., Barbosa, H. M. J., Pöhlker, M. L., Alexander, M. L., Brito, J., Carbone, S., Castillo, P.,
425 Day, D. A., Kuang, C., Manzi, A., Ng, N. L., Sedlacek Iii, A. J., Souza, R., Springston, S., Watson, T., Pöhlker, C., Pöschl,
U., Andreae, M. O., Artaxo, P., Jimenez, J. L., Martin, S. T., and Wang, J.: CCN activity and organic hygroscopicity of aerosols
downwind of an urban region in central Amazonia: seasonal and diel variations and impact of anthropogenic emissions, *Atmos.*
Chem. Phys., 17, 11779-11801, 10.5194/acp-17-11779-2017, 2017.
- Uin, J.: 3002 Humidified Tandem Differential Mobility Analyzer Instrument Handbook, DOE Office of Science Atmospheric
430 Radiation Measurement (ARM) Program ..., 2016.
- Vaattovaara, P., Huttunen, P. E., Yoon, Y. J., Joutsensaari, J., Lehtinen, K. E. J., O'Dowd, C. D., and Laaksonen, A.: The
composition of nucleation and Aitken modes particles during coastal nucleation events: evidence for marine secondary organic
contribution, *Atmos. Chem. Phys.*, 6, 4601-4616, 10.5194/acp-6-4601-2006, 2006.
- Wang, J., Dong, X., and Wood, R.: Aerosol and Cloud Experiments in Eastern North Atlantic (ACE-ENA) Science Plan, DOE
435 Office of Science Atmospheric Radiation Measurement (ARM) Program ..., 2016.
- Watson, T. B.: Aerosol chemical speciation monitor (ACSM) instrument handbook, DOE Office of Science Atmospheric
Radiation Measurement (ARM) Program ..., 2017.
- Wex, H., Petters, M. D., Carrico, C. M., Hallbauer, E., Massling, A., McMeeking, G. R., Poulain, L., Wu, Z., Kreidenweis, S.
M., and Stratmann, F.: Towards closing the gap between hygroscopic growth and activation for secondary organic aerosol:
440 Part 1 – Evidence from measurements, *Atmos. Chem. Phys.*, 9, 3987-3997, 10.5194/acp-9-3987-2009, 2009.



- Willis, M. D., Burkart, J., Thomas, J. L., Köllner, F., Schneider, J., Bozem, H., Hoor, P. M., Aliabadi, A. A., Schulz, H., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Growth of nucleation mode particles in the summertime Arctic: a case study, *Atmos. Chem. Phys.*, 16, 7663-7679, 10.5194/acp-16-7663-2016, 2016.
- Willis, M. D., Köllner, F., Burkart, J., Bozem, H., Thomas, J. L., Schneider, J., Aliabadi, A. A., Hoor, P. M., Schulz, H.,
445 Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Evidence for marine biogenic influence on summertime Arctic aerosol, *Geophysical Research Letters*, 44, 6460-6470, 10.1002/2017gl073359, 2017.
- Wood, R.: Stratocumulus Clouds, *Monthly Weather Review*, 140, 2373-2423, 10.1175/mwr-d-11-00121.1, 2012.
- Wood, R., Wyant, M., Bretherton, C. S., Rémillard, J., Kollias, P., Fletcher, J., Stemmler, J., Szoeki, S. d., Yuter, S., Miller, M., Mechem, D., Tselioudis, G., Chiu, J. C., Mann, J. A. L., O'Connor, E. J., Hogan, R. J., Dong, X., Miller, M., Ghate, V.,
450 Jefferson, A., Min, Q., Minnis, P., Palikonda, R., Albrecht, B., Luke, E., Hannay, C., and Lin, Y.: Clouds, Aerosols, and Precipitation in the Marine Boundary Layer: An Arm Mobile Facility Deployment, *Bulletin of the American Meteorological Society*, 96, 419-440, 10.1175/bams-d-13-00180.1, 2015.
- Wurl, O., Wurl, E., Miller, L., Johnson, K., and Vagle, S.: Formation and global distribution of sea-surface microlayers, *Biogeosciences*, 8, 121-135, 10.5194/bg-8-121-2011, 2011.
- 455 Yu, F., and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691-7710, 10.5194/acp-9-7691-2009, 2009.
- Zheng, G., Wang, Y., Aiken, A. C., Gallo, F., Jensen, M. P., Kollias, P., Kuang, C., Luke, E., Springston, S., Uin, J., Wood, R., and Wang, J.: Marine boundary layer aerosol in the eastern North Atlantic: seasonal variations and key controlling processes, *Atmos. Chem. Phys.*, 18, 17615-17635, 10.5194/acp-18-17615-2018, 2018.
- 460 Zheng, G., Sedlacek, A. J., Aiken, A. C., Feng, Y., Watson, T. B., Raveh-Rubin, S., Uin, J., Lewis, E. R., and Wang, J.: Long-range transported North American wildfire aerosols observed in marine boundary layer of eastern North Atlantic, *Environment International*, 139, 105680, <https://doi.org/10.1016/j.envint.2020.105680>, 2020.



465 Figures






Figure 1. Matching aerosol Aitken mode with hygroscopicity measurements at fixed particle diameters. The black dash lines indicate the selected particle size (i.e., $D_{p,SCCN}$) at which the hygroscopicity parameter κ is derived. The shaded areas indicate one σ range from the fitted lognormal mode diameter, $D_{p,n}$.

470



475 **Figure 2. Derivation of $\kappa_{e,CCN}$ from size-resolved CCN measurement during an example condensational growth event.** (a) Mode diameter and hygroscopicity κ_{CCN} of the growing Aitken mode. The black circles are fitted mode diameter, $D_{p,n}$, and the shaded area indicate the one σ range of the fitted mode. Black line shows the increasing trend of $D_{p,n}$, which is used to identify growth events. (b) Derivation of the original particle diameter (D_{p0}) at the beginning of the growth event from particle diameter after growth (D_{p1}) using cumulative particle size distributions. (c) Derivation of $\kappa_{e,CCN}$ through linear fitting of κ_{CCN} versus $f_{V,cond}$.



480 **Figure 3. Examples of pre-CCN particle growth dominated by (a,b) acidic sulfates and (c,d) organics, respectively, as observed in 2017.** (a)(c) Examples of growth events identified from the time series of measured aerosol size distribution. The black circles indicate lognormal-fitted Aitken mode diameter, and the black lines indicate the growth of the mode diameter (see section 3). (b)(d) Particle hygroscopicity κ_{CCN} as a function of the volume fraction of condensed species in the growing particles ($f_{V,cond}$). $f_{V,cond}$ increases as particles grow by condensation. The value of $\kappa_{c,CCN}$ is derived from the variation of κ_{CCN} with $f_{V,cond}$ (section 3).

485






Figure 4. Monthly distribution of observed condensational growth events and the category of dominant condensing species during the ACE-ENA campaign.



490

Figure 5. Comparison of $\kappa_{c,CCN}$ and $\kappa_{c,GF}$ values for the intermediate $\kappa_{c,CCN}$ category, colored by the measured molar ratios of $\text{NH}_4^+ / \text{SO}_4^{2-}$.



495

Figure 6. Potential sources of the condensing organics. Correlations of the derived $\kappa_{c,CCN}$ with (a,c) MSA/SO₂ ratio and (b,d) non-refractory PM₁ organics/SO₄²⁻ ratio, for (a,b) the clean air masses from mid-latitude Atlantic or Arctic, and (c,d) the continental or subtropical air masses. Numbers shown in (a, b) indicate the month in which the growth events occurred.