

1
2
3
4 **Atmospheric gas-phase composition over the Indian Ocean**
5
6

7 Susann Tegtmeier¹, Christa Marandino², Yue Jia¹, Birgit Quack², Anoop S. Mahajan³
8
9

10 ¹Institute of Space and Atmospheric Studies, University of Saskatchewan, Saskatoon, Canada
11

12 ²GEOMAR Helmholtz Centre for Ocean Research Kiel, 24105 Kiel, Germany
13

14 ³Center for Climate Change Research, Indian Institute of Tropical Meteorology, Pune,
15 411016, India
16

17 Corresponding author: Susann Tegtmeier (susann.tegtmeier@usask.ca)
18

Abstract

The Indian Ocean is coupled to atmospheric dynamics, and chemical composition via several unique mechanisms, such as the seasonally varying monsoon circulation. During the winter monsoon season, high pollution levels are regularly observed over the entire northern Indian Ocean, while during the summer monsoon, clean air dominates the atmospheric composition, leading to distinct chemical regimes. The changing atmospheric composition over the Indian Ocean can interact with oceanic biogeochemical cycles and impact marine ecosystems, resulting in potential climate feedbacks.

Here, we review current progress in detecting and understanding atmospheric gas-phase composition over the Indian Ocean and its local and global impacts. The review considers results from recent Indian Ocean ship campaigns, satellite measurements, station data and information on continental and oceanic trace gas emissions. The distribution of all major pollutants and greenhouse gases shows pronounced differences between the landmass source regions and the Indian Ocean with strong gradients over the coastal areas. Surface pollution and ozone are highest during the winter monsoon over the Bay of Bengal and the Arabian Sea coastal waters due to air mass advection from the Indo-Gangetic Plain and continental outflow from Southeast Asia. We observe, however, that unusual types of wind patterns can lead to pronounced deviations of the typical trace gas distributions. For example, the ozone distribution maxima shift to different regions under wind scenarios that differ from the typical seasonal transport patterns. The distribution of greenhouse gases over the Indian Ocean shows many similarities when compared to the pollution fields, but also some differences of the latitudinal and seasonal variations resulting from their long lifetimes and biogenic sources. Mixing ratios of greenhouse gases such as methane show positive trends over the Indian Ocean, but long-term changes of pollution and ozone, and in particular how they are driven by changing emissions and transport patterns, require further investigation in the future. Although we know that changing atmospheric composition and perturbations within the Indian Ocean affect each other, the impacts of atmospheric pollution on oceanic biogeochemistry and trace gas cycling is severely understudied. We highlight potential mechanisms, future research topics and observational requirements that need to be explored in order to fully understand such interactions and feedbacks in the Indian Ocean region.

1. Introduction

Over the Indian Ocean, intense anthropogenic pollution from Southeast Asia mixes with pristine oceanic air. The interplay of the polluted continental and the clean oceanic air masses, and the resulting atmospheric composition are determined by distinct seasonal circulation patterns. The large-scale monsoon circulations in combination with anthropogenic emissions from southern Asia lead to seasonally contrasting chemical regimes over the Indian Ocean. As the anthropogenic emissions include relatively large contributions from biofuel/biomass combustion and incomplete industrial burning, the atmospheric composition during polluted periods shows unique characteristics when compared to other regimes. The complex mixture of chemical constituents and large-scale transport patterns can have a profound influence on oceanic processes, stratospheric composition, and neighbouring regions such as the Mediterranean and Africa. This article provides a review of the recent progress in our understanding of the atmospheric gas-phase composition over the Indian Ocean and how it impacts the ocean and upper atmosphere. This article is part of the special issue ‘Understanding the Indian Ocean system: past, present and future’.

1.1 Region

The Indian Ocean is the world's third largest ocean covering 19.8% of the water on the Earth's surface. In contrast to the Pacific and Atlantic Oceans, it does not stretch from pole to pole, but is enclosed on three sides by major landmasses and an archipelago. The Indian Ocean is centred on the Indian Peninsula, which also forms the northern border together with Iran, Pakistan, and Bangladesh. In the west, the Indian Ocean is bounded by East Africa and the Arabian Peninsula, while the eastern and southern boundaries are set by Southeast Asia, Australia, and the Southern Ocean.

The countries bordering the Indian Ocean are home to one-third of the world's population accounting for approximately 2.5 billion people (Roser et al., 2013). The economies of many Indian Ocean countries are expanding rapidly, with India being the fastest growing major economy in the world. Similarly, many Indian Ocean countries show a rapid population growth, which is expected to further increase in the future. Given the quickly growing populations and industries, the Indian Ocean is becoming a pivotal zone of strategic political competition. At the same time, the Indian Ocean hosts a large variety of marine ecosystems including coral reefs, seagrass beds, and mangrove forests. Anthropogenic activities along the coastlines and climate change threaten biodiversity in the Indian Ocean, which contains 25% of the Earth's biodiversity hotspots (Mittermeier et al., 2011).

Growing populations also lead to rapidly increasing anthropogenic emissions. Burning conditions are often poorly controlled, as for instance during biofuel burning in cook-stoves, and fossil fuel burning in vehicles (Li et al., 2017). Together with burning of coal and other fossil fuels for energy production, this leads to large emissions of man-made trace species including greenhouse gases and ozone precursors (e.g., Lawrence, 2004). In addition, primary aerosols, such as soot and dust, and precursors of secondary aerosols are released in relatively large amounts. As a result, air pollution is a serious health issue in many Indian Ocean countries, leading to increases in respiratory and cardiovascular problems (Rajak and Chattopadhyay, 2019). The intense pollution has also been linked to regional weather impacts, such as changes of rainfall patterns and decreasing crop harvests (e.g., Bollasina et al., 2011, Li et al., 2016).

1.2 Seasons

Seasonal changes of atmospheric transport patterns are the main driver of Indian Ocean chemical regimes and lead to periods of intense anthropogenic pollution alternating with periods of clean oceanic air. The South Asian monsoon circulation, the strongest monsoon system in the world, dominates the regional meteorology of the Indian subcontinent. The seasonal reversal of the winds is coupled to a strong annual cycle of precipitation with very wet summer and dry winter conditions (Chang, 1967). Being the dominant driver of the annual cycle of rainfall, the South Asian monsoon controls the water and food security of the region and, thus, the well-being and prosperity of large populations.

The monsoon system also has a strong impact on the atmospheric composition over the Indian Ocean. During the winter monsoon from December to February, continental aerosols as well as man-made trace species and their reaction products dominate the chemical regime (Lelieveld et al., 2001). A layer of air pollution is visible on satellite pictures as a haze of brown colour hanging over much of South Asia and the Indian Ocean. This so-called Indian Ocean brown cloud has been suggested to impact regional climate by masking greenhouse gas induced surface warming (Ramanathan et al., 2005) and to affect monsoon rainfall.

In contrast, clean air dominates the atmospheric composition over the Indian Ocean during the summer monsoon from June to September, leading to a completely different chemical regime. Atmospheric pollutant levels over the Indian Ocean are low and typical open ocean background

conditions can be observed (Lawrence and Lelieveld, 2010). While boreal summer conditions prevent the anthropogenic pollution from spreading across the Indian Ocean, an anticyclonic circulation, centred at 200 to 100 hPa, offers an efficient pathway for continental pollution into the global upper troposphere and lower stratosphere (e.g., Randel et al., 2010; Lelieveld et al., 2018).

Finally, during the monsoon transition periods from April to May and September to October, the offshore pollution is less strong, due to weaker air mass transport from Southeast Asia and Africa over the Indian Ocean (Sahu et al., 2006).

1.3 Early work

The largest international scientific study exploring the impact of South Asian emissions on the composition of the atmosphere over the Indian Ocean, the Indian Ocean Experiment (INDOEX), took place during the winter monsoon 1999, with some pilot campaigns conducted during 1996–1997. During the multiplatform field campaign, surprisingly high pollution was detected over the entire northern Indian Ocean all the way to the Intertropical Convergence Zone (ITCZ). Scientific studies revealed that the nature of the pollution was considerably different from that in Europe or North America, with strongly enhanced carbon monoxide concentrations related to widespread biofuel use and agricultural burning (e.g., Lelieveld et al., 2001). Other large pollution sources based on fossil fuel combustion and biomass burning were linked to high loads of sunlight-absorbing aerosols with potential consequences for the regional atmospheric energy balance (Ramanathan et al., 2002). A few years before INDOEX, the Joint Global Ocean Flux Study (JGOFS) India investigated the factors controlling carbon fluxes in the Arabian Sea and led to estimates of CO₂ emissions to the atmosphere for this region (Sarma et al., 2003).

The INDOEX findings, presented in many scientific publications, have drawn attention to this region and several projects and campaigns followed over the next decade. The Bay of Bengal Experiment (BOBEX) research cruise during February to March 2001 detected high ozone and pollution levels over the Bay of Bengal and linked them to transport from the continent (Naja et al., 2004; Lal et al., 2006). The southern Indian Ocean was explored during the Pilot Expedition to the Southern Ocean (PESO) research cruise from January to April 2004, which revealed much cleaner air masses with smaller aerosol loadings in the region south of the ITCZ (Pant et al., 2009). Other research cruises such as the Bay of Bengal Processes Studies (BOBPS) during September to October 2002, investigated oceanic productivity and nutrients in relation to air-sea exchange of climate active gases (Sardessai et al., 2007). A detailed overview of research cruises, island measurements and aircraft campaigns investigating the atmosphere over the Indian Ocean is given in Lawrence and Lelieveld (2010). The authors provide a comprehensive review of the state of the art at this time by bringing together observational and modelling studies.

1.4 Scope and organization of this study

Here we will focus on recent progress in the field by giving an overview of results ~~obtained~~ published after 2010. We will synthesize the current understanding of Indian Ocean gas phase atmospheric composition and explore how it is driven by emission sources, transport, and chemistry. Our region of interest encloses the Indian Ocean from 30°S to its northern continental borders and across its whole longitudinal range. It also includes neighbouring land mass such as parts of East Africa, the Arabian Peninsula, South Asia, Southeast Asia, and Australia as depicted in Figure 1. This review focuses on three groups of atmospheric gases 1) ozone and pollutants - carbon monoxide (CO), nitrogen oxides (NO_x), sulphur dioxide (SO₂), ammonia (NH₃) and mercury; 2) greenhouse gases - methane (CH₄), nitrous oxide (N₂O),

carbon dioxide (CO₂) and carbonyl sulphide (COS) and 3) volatile organic compounds (VOCs) and short-lived biogenic gases - dimethylsulphide (DMS), isoprene and halogen compounds.

Section 2 provides an overview of the physical processes in the atmosphere and ocean that are relevant for the atmospheric composition. Section 3 will introduce all campaigns and measurements that are the basis for scientific studies published after 2010 and are discussed here. Regional sources and sinks of greenhouse gases, pollution and biogenic trace gases will be given in Section 4. Short introductions to all gases listed above, including their role in the atmosphere, can also be found in section 4. The focus of section 5 is on our current knowledge of the atmospheric composition over the Indian Ocean and how it is driven by physical processes and by regional sources. We will present a synthesis of the scientific progress made after 2010 in Section 6, where we will also discuss the global and local impacts of the Indian Ocean atmospheric composition. An outlook and a summary of current knowledge gaps are given in Section 7. A key for all abbreviations used in this paper is provided in Appendix A.

2 Physical processes

2.1 Atmospheric processes

The South Asian monsoon circulation dominates the transport patterns and regional meteorology over the Indian Ocean. Strong seasonal circulation changes give rise to three main meteorological regimes: the summer monsoon from June to September, the winter monsoon from November to March and the transition periods from April to May and from the end of September to October. Surface flow patterns as well as seasonal transport regimes are described in the following subsections, while a discussion of intraseasonal and interannual variability can be found in the supplementary materials section A.

Near-surface flow patterns

A detailed picture of the near-surface flow patterns is provided in Fig. 1 in the form of seasonal mean surface wind fields and sea level pressure derived from 2018/2019 ERA-Interim reanalysis data. Seasonal mean plots here and in the rest of the paper are shown for core monsoon and transition periods, i.e., June to August (JJA) for the summer monsoon, December to February (DJF) for the winter monsoon, April to May (Apr-May) for the boreal spring transition period and October (Oct) for the boreal autumn transition period. The equatorial and northern Indian Ocean (north of 10°S) are dominated by seasonally reversing monsoon winds (Schott and McCreary, 2001; Schott et al., 2009). Southeast winds occur during the summer monsoon with the low-pressure system of the ITCZ shifted north of 15°N (Fig. 1a), while northwest winds occur during the winter monsoon with the low-pressure belt situated south of the equator (Fig. 1b). Over the southern Indian Ocean (south of 10°S), steady southeast trades prevail during all seasons, but reach further northward during the northern summer.

The seasonally reversing monsoon winds and inter-hemispheric pressure gradients over the equatorial Indian Ocean are a striking feature different from the other tropical oceans, where sustained easterly winds are found along the equator. In contrast, equatorial winds over the Indian Ocean are westerlies during the monsoon transition periods (Fig. 1c and 1d) and show a weak westerly annual mean component (Lamb and Hastenrath, 1979). These equatorial westerlies are driven by an interplay of an eastward pressure gradient along the equator, the latitudinal position of the flow recurvature and the strength of the trade winds (Hastenrath and Polzin, 2004).

MSLP and Surface Wind




Figure 1. Mean surface level pressure (MSLP) and surface wind for (a) summer monsoon 2018 (June – August), (b) winter monsoon 2018/2019 (December – February), (c) spring transition 2018 (April – May) and (d) autumn transition 2018 (October) periods from ERA-Interim.

Summer and winter monsoon

During the summer monsoon, steady onshore winds transport air from the ocean over to the continent, where it results in deep convection and the well-known Indian summer monsoon rains. Air masses experiencing fast upward transport in convective updraughts converge in the upper troposphere forming a high-pressure system. The associated anticyclone circulation is tied to the outflow of the deep convection and is situated directly over the highly polluted southern Asia. As a result, distinct tracer anomalies have been observed in the anticyclone indicating strong upward transport of pollution from the surface (e.g., Randel et al., 2010). Given the dynamical confinement of tropospheric tracers and aerosols in the anticyclone, the Asian monsoon system provides a potentially efficient pathway from the surface to the tropical upper troposphere and lower stratosphere. However, a significant fraction of the pollution can be removed from the air mass before entering the stratosphere due to lightning driven OH reactions in monsoonal convection (Lelieveld et al., 2018).

During the winter monsoon, the prevailing north-easterly winds reverse the meteorological situation. There is little rain over southern Asia marking this as the ‘dry season’ and the missing convection chemically disconnects the surface layer from the upper troposphere (Kunhikrishnan et al., 2004). Instead, pollution outflow occurs in the marine boundary layer (MBL) via offshore winds towards the northern Indian Ocean down to the equator. Primary MBL flow channels have been identified in the western Arabian Sea, the eastern Arabian Sea

just off the Indian west coast, the western Bay of Bengal and Southeast Asia (Krishnamurti 1997a; 1997b; Verver et al., 2001).

Winter monsoon flow patterns are further complicated by effects of the land-sea breeze, which lofts coastal air masses above the MBL (Simpson and Raman, 2004). The associated offshore flow above the MBL transports air masses over the coastal oceans where they constitute the so-called 'elevated layer'. Due to the relatively rapid outflow, the elevated layer provides an additional effective mechanism for pollution transport from the continents towards the Indian Ocean (Lawrence and Lelieveld, 2010). As a result, outflow during the winter monsoon occurs in two distinct layers, namely the pollutant plume within the MBL (up to 800–1000m) and the elevated layer (1–3km). Once over the northern Indian ocean, the north-easterly trade winds transport the air masses towards the ITCZ, typical within 7–10 days (Ethe' et al., 2002). Similarly, over the southern Indian Ocean, south-easterly winds transport pristine boundary layer air masses northwards. At the ITCZ, these trade wind flows converge, and associated convection transports the air upwards into the upper troposphere (Iyengar et al., 1999).

Over the western part of the tropical Indian Ocean, the ITCZ has been observed to occur simultaneously in two bands on either side of the equator forming the so-called double ITCZ (Meenu et al., 2007) throughout the year. Based on cloud characteristics and outgoing longwave radiation, the most preferred latitudes for the northern and southern bands of the ITZC were found to be around 5°N and 7.5°S to 10°S.

2.2 Oceanic processes transport

The physical oceanography of the northern Indian Ocean reflects the seasonal changes of the monsoon cycle, including impacts on currents, thermohaline circulation, sea surface temperature (SST), salinity, and upwelling events, while the equatorial and southern Indian Ocean do not experience this influence (~~Fig-2~~). For more details about general Indian Ocean physical processes, please see Schott et al. (2009) and references therein, as well as Phillips et al. (2021) from this issue and supplementary material B. In the following sections, we concentrate on how physical oceanography affects salinity, SST, and biological productivity, which in turn play a major role in controlling air-sea exchange and atmospheric composition, in particular for biogenic trace gases such as bromoform, DMS, and isoprene.





Figure 2. Schematic representation of identified oceanic currents during the summer monsoon (a) and winter monsoon (b). Current branches indicated are the South Equatorial Current (SEC), South Equatorial Countercurrent (SECC), Northeast and Southeast Madagascar Current (NEMC and SEMC), East African Coastal Current (EACC), Somali Current (SC), Southern Gyre (SG) and Great Whirl (GW) and associated upwelling wedges (green shades), Southwest and Northeast Monsoon Currents (SMC and NMC), South Java Current (SJC), East Gyral Current (EGC), and Leeuwin Current (LC). The subsurface return flow of the supergyre is shown in magenta. Depth contours shown are for 1000 m and 3000 m (grey). Updated representations are from SMC01; red vectors (Me) show directions of meridional Ekman transports. ITF indicates Indonesian Throughflow (from Schott et al., 2009, copyright 2009 by the American Geophysical Union, reproduced with permission).

Salinity, SSTs and productivity

On seasonal time scales, freshwater input due to rainfall and river discharge is important for the salinity balance in the Bay of Bengal and horizontal advection related to the monsoon plays a dominant role in the north Indian Ocean (Rao and Sivakumar 2003; Da-Allada et al., 2015). In the southwestern tropical Indian Ocean, the freshwater flux due to precipitation is a major control on the salinity (Da-Allada et al., 2015). The rainfall over the Indian Ocean shows a general migration to the summer hemisphere following sunlight and warm SSTs, highlighting their strong coupling. Conversely, latent heat loss caused by cool, dry air from the Asian continent leads to strong wintertime cooling in the northern Arabian Sea. The strong summertime cooling in parts of the Arabian Sea instead is a combined result of latent heat loss caused by the strong south-westerly winds and upwelling/offshore advection from the Somali and Omani coasts. During boreal summer, upwelling induced cooling off Somalia prevents atmospheric convection from the western Arabian Sea. From the eastern Arabian Sea to the South China Sea, north of the Equator, high SSTs promote atmospheric deep convection (Schott et al., 2009).

The oceanic upwelling caused by strong monsoonal winds supplies nutrients to the surface layer, where they support elevated rates of primary productivity. This occurs mainly in the Arabian Sea, the Somali Basin, along the Indian coast and the northern Bay of Bengal, especially during summer months. The seasonal reversals in the boundary currents of the northern Indian Ocean, including the seasonal switching from upwelling to downwelling circulations, have important biogeochemical and ecological impacts that modify primary productivity, nutrient stoichiometry, oxygen concentrations and phytoplankton species composition (Hood et al., 2017). In addition, transient upwelling due to seasonal variations of currents and mesoscale variability can give rise to episodically high levels of primary production throughout the Indian Ocean coastal waters.

2.3 Long-term changes

Indian Ocean warming

The Indian Ocean has warmed steadily over the past century, with an SST increase of 1°C during 1951–2015, markedly higher than the global average SST warming of 0.7°C, over the same period (Du and Xie, 2008; Han et al., 2014; Krishnan et al., 2020). Overall, this Indian Ocean-averaged warming rate is broadly consistent across observational products (Dong et al. 2014; Yao et al., 2016) and historical simulations from the Coupled Model Intercomparison Project - Phase 5 (CMIP5). It can be largely attributed to anthropogenic forcing rather than natural external forcing, such as volcanic and solar variations (e.g., Dong et al., 2014). It has been shown that the basin wide warming due to increasing greenhouse gases is slowed down by the indirect effects of anthropogenic aerosol (Dong and Zhou, 2014). In addition to anthropogenic forcing, the sustained warming over the Indian Ocean warm pool region is caused by local ocean–atmosphere coupled mechanisms, with their relative roles being debated (e.g., Dong et al., 2014; Du and Xie, 2008). The importance of the Indian Ocean in the global ocean heat budget was not recognised until the hiatus period at the beginning of the 21st century, during which the abrupt increase of the upper Indian Ocean heat content served as a major sink of the excessive heat entering the Earth system (e.g., Nieves et al., 2015).

The Indian Ocean warming is not spatially homogeneous in both models and observations. The western tropical Indian Ocean has been warming for more than a century, at a rate faster than any other region of the tropical oceans and is the largest contributor to the overall trend in the global mean SST (Roxy et al., 2014). Positive ~~sea surface temperature~~SST anomalies in the western part of the Indian ocean, have increased markedly since 1950, while negative events have reduced (Cai et al., 2009). The warming of the generally cool western Indian Ocean against the rest of the tropical warm pool region (Roxy et al., 2014, 2015) and corresponding changes

of the zonal SST gradient (Saha et al., 2014) have been both proposed as plausible explanations for the observed decrease of Indian monsoon rainfall over the last three decades. In addition, they have the potential to alter the marine food webs in this biologically productive region.

The Indian Ocean warming is projected to further increase over the course of the 21st century in response to unabated greenhouse gas emissions. By the end of the 21st century, strongest warming in the Arabian Sea and western equatorial Indian Ocean is consistently projected in CMIP models, which could yield more Arabian Sea cyclones and further decrease monsoonal rains (Gopika et al., 2020).

Summer monsoon and precipitation

There are large uncertainties related to variability in the South Asian summer monsoon in a changing climate. Several studies debate whether the monsoon is weakening or strengthening, as well as the mechanisms driving the changes (Roxy et al., 2015). According to a review by Singh et al. (2019a), both observational and modelling studies have determined that the potentially weakened monsoon is due to a combination of forcings, such as land use and irrigation changes, increased greenhouse gas as well as anthropogenic aerosols. Roxy et al. (2015) provide compelling evidence that Indian Ocean warming potentially weakens the land-sea thermal contrast and dampens the summer monsoon Hadley circulation, leading to reduced rainfall over parts of South Asia.

The Indian Ocean is one of the greatest moisture sources accounting for nearly one-third of the total net transport of water toward the continents (Bengtsson, 2010). Both remote and local SST anomalies can induce hydrological cycle changes over the Indian ocean affecting local and remote precipitation. ~~An increase in evaporation is due to the robust warming of SSTs during recent decades, via changes of the near surface specific humidity gradient, near surface wind and the turbulent exchange coefficient (Yu, 2007; Richter and Xie, 2008).~~ Han et al. (2019) showed that the moisture sources (evaporation minus precipitation) in the tropical central-eastern and south-western Indian Ocean experienced a significant increase during boreal summer between 1979 to 2016. In addition, there has been a significant reduction in the annual frequency of tropical cyclones and their associated rainfall over the northern Indian Ocean since the middle of the twentieth century (Krishnan et al., 2020). In contrast, the frequency of very severe cyclonic storms during the autumn transition season has increased significantly during the last two decades. At the same time, an enhanced rainfall contribution has occurred due to a higher precipitation efficiency (Singh et al., 2016; 2019b) possibly leading to a dry atmosphere. Further changes of the Indian summer monsoon rainfall are expected for the future, however current model projections give contradicting results (e.g., Roxy et al., 2015; Zou and Zhou, 2013).

Changes in surface wind are expected to be moderate for the first half of the 21st century, with a noticeable decline of wind speed over the tropical Indian Ocean due to reduced thermal land-sea contrasts. The southern extratropical region and Southern Ocean, on the other hand, show a significant strengthening of the wind fields by the end of the twenty-first century (Mohan and Bhaskaran, 2019). Not well represented ocean-atmosphere feedback and coarse model resolutions, however, are known to lead to large uncertainties in model estimates of wind speed changes (Annamalai et al. 2017; Mohan and Bhaskaran, 2019).

Salinity and productivity

Du et al. (2015) noted freshening in the south-eastern tropical Indian Ocean starting in the mid-1990s. Idealised model experiments suggest that multidecadal changes of subsurface ocean salinity during 1950–2000 were due to isopycnal migration related to ocean surface warming (Lago et al. 2016). However, the enhanced precipitation in the Maritime Continent and the

strengthening of the Indonesian Throughflow are thought to be the likely causes of the freshening trend in the south-east Indian Ocean since early 2000s (Llovel and Lee 2015; Hu and Sprintall, 2017).

While Behrenfeld et al. (2006) indicate a reduction in net primary productivity over most of the tropics as a result of surface thermal stratification, they have suggested an increase in primary productivity for the western Indian Ocean from 1998 to 2004. Recent biogeochemical simulations of the Arabian Sea ecosystem also predict that a projected intensification of monsoon winds strongly increases the ecosystem productivity, thereby amplifying the oxygen biological consumption and intensifying the oxygen minimum zone (OMZ) at depth (Lachkar et al., 2018). At the same time, the near-surface will experience increased ventilation due to the predicted stronger winds. On the contrary, a review in this issue summarizes evidence indicating a significant, but small, reduction in primary production in the northern Indian Ocean (Löscher et al., 2021). ~~Included, is a study using chlorophyll data and Earth system model simulations over a larger region of the western Indian Ocean, which points out a~~ An alarming decrease of up to 20% in marine phytoplankton during the past six decades has been identified in the western Indian Ocean (Roxy et al., 2016). ~~The authors suggest that this decrease is driven by enhanced ocean stratification due to the rapid warming in the Indian Ocean, which and~~ suppressed ~~ds~~ nutrient mixing from subsurface layers. Gregg and Rousseaux (2019) also conclude from the assimilation of ocean colour satellite data (1998–2015) into an ocean biogeochemical model, that the decline in global ocean primary productivity of 2.1% per decade is mainly driven by the northern and equatorial Indian Ocean. Any changes to biological processes have the potential to alter trace gas cycling in the surface ocean.

3. Campaigns, station data and satellite measurements

A few Indian Ocean coastal or island stations have been operated as part of long-term scientific measurement programmes or operational air quality networks providing limited area observations. In addition, intensive ship and aircraft campaigns have been conducted for detailed investigations of atmospheric processes during short episodes. These data can be complemented by satellite observations of the tropospheric composition, which provide the large-scale picture for a number of substances, albeit often with limited vertical resolution and reduced accuracy for individual measurements. In this section, we will give an overview of campaigns, station data and satellite measurements that have been applied to study the atmospheric composition over the Indian Ocean over the last decade.

3.1 Campaigns and station data

Over the last decades, chemical, physical, and biogeochemical processes occurring in and above the Indian Ocean have been explored during various field campaigns. Here, all campaigns that have contributed to the recent progress in the field and led to publications after 2010 are summarized in Table 1. It should be noted that the ICARB multi-platform field experiment consisted of three phases, with the first phase exploring post winter monsoon composition in 2006 (ICARB), a second phase taking place during the winter monsoon 2008/2009 (referred to as W_ICARB) supplemented by aircraft measurements and a third phase during the winter monsoon of 2018 (referred to as ICARB-218).

Table 1. Summary of campaigns in the Indian Ocean for the 21st century.

Campaign	Time	Region	Objective	Reference
ARMEX (Arabian Sea Monsoon Experiment)	2002 Jun – Aug	Arabian Sea	Study of reactive halides released from sea salt aerosols and their role in ozone chemistry	Ali et al., 2009
PESO (Pilot Expedition to the Southern Ocean)	2004 Jan – Apr	Southern Indian Ocean	Multi-disciplinary expeditions to understand the forcing mechanisms behind widely geographically separated climate change.	Pandey et al. (2006)
ISOE 1-11 (1st-11th Indian Southern Ocean Expedition)	Since 2004	Southern Indian Ocean	Identify role and response of Southern Ocean to the regional and global climate variability (emissions of trace gases since ISOE 8)	ISOE Reports Mahajan et al. (2019a, b) Inamdar et al. (2020)
ICARB (Integrated Campaign for Aerosols, gases and Radiation Budget)	2006 Mar – May; 2008/09 Dec – Jan; 2018 Jan – Feb	Indian mainland, northern Indian Ocean, Bay of Bengal	Characterize the physico-chemical properties and radiative effects of atmospheric aerosols and trace gases over the Indian landmass and the adjoining oceanic regions.	Moorthy et al. (2008) David et al., 2011
Sagar Kanya cruise during the CTCZ (Continental Tropical Convergence Zone) experiment	2009 Jul – Aug	Bay of Bengal	Understanding the gaseous atmospheric boundary layer composition over the Bay of Bengal during the summer monsoon season	Girach et al. (2017)
Campaign aboard the Ocean Research Vessel Sagar Kanya (SK-277)	2010 Oct – Nov	Bay of Bengal	Analyze atmospheric composition over the Bay of Bengal and how it is driven by air mass origin from Indian Ocean, Southeast Asia, and the Indian subcontinent	Mallik et al. (2013)
OASIS (Organic VSLs and their air sea exchange from the Indian Ocean to the Stratosphere)	2014 Jul – Aug	West Indian Ocean	Investigate oceanic emissions of very short-lived substances and their transport and chemistry from the tropical Indian Ocean to the atmosphere, in particular to the stratosphere	Fiehn et al. (2017) Zavarsky et al. (2018b)
OMO (Oxidation Mechanism Observations aircraft campaign)	2015 Jul – Aug	Indian Ocean and the Mediterranean	Identify atmospheric impacts of associated air pollution emissions at regional and global scales during the South Asian summer monsoon	Lelieveld et al. (2018)
IIOE-2 (Second International Indian Ocean Expedition)	2015 – 2020	Indian Ocean	Advance Indian Ocean initiatives and projects addressing emerging scientific issues of the Indian Ocean in the 21st century	Hood et al., 2016

AQABA (Air Quality and climate change in the Arabian BASin)	2017 Jul – Aug	Mediterranean and Arabian Peninsula	Study air quality and climate change in the Arabian Basin	Bourtsoukidis et al., 2019
--	-------------------	--	--	-------------------------------

In addition to dedicated campaigns, some island and coastal stations have conducted long-term measurements that provide valuable information about the atmospheric composition over the Indian Ocean.

CO₂ surface flask measurements from the Cape Rama site on the Indian coastline have been used to analyse the distribution and variability of CO₂ over this region for 2009–2012 (Nalini et al., 2018). Measurements of CH₄, another important greenhouse gas, and the pollutant CO are available from ground-based in situ cavity ring-down spectroscopy analysers and Fourier transform infrared spectrometers at two sites on Reunion Island in the southern Indian Ocean (Zhou et al., 2018). These multi-annual time series (2011–2017) allowed to investigate the impact of emissions from biomass burning in Africa and South America on atmospheric pollutant levels over the Indian Ocean. CO surface data are also available from the NOAA/ESRL Global Monitoring Division network station in Mahé (Wai et al., 2014).

In situ tropospheric ozone measurements have been collected from 2003 to 2007 from balloon-borne electrochemical concentration cell sensors launched above Ahmedabad in western India (Lal et al., 2014). The continuous dataset enabled studies of the impact of transport processes on the seasonal cycle and on the vertical distribution of ozone. The observation site in Trivandrum situated on the southwest coast of India collected measurements of nitrogen oxides with a chemiluminescence NO_x analyser from 2007 to 2009 (David and Nair, 2011).

3.2 Satellite measurements

Satellite measurements of atmospheric composition over the Indian Ocean have provided valuable information over the last decades that allowed for studies of the overall distribution and long-term changes of key substances. Most instruments used today apply passive remote sensing with observations being mainly done in nadir geometry. Here we will give a short overview of satellite instruments that provide measurements used in scientific studies of the Indian Ocean atmosphere. In addition, we have compiled plots of the seasonal CO, NO₂ and CH₄ surface distribution for this review article and will describe the respective satellite measurements more in detail.

Pollutants (NO₂) from OMI and TROPOMI

The Ozone Monitoring Instrument (OMI) is a key instrument onboard NASA's Aura satellite. OMI is a nadir-viewing, wide-field-imaging spectrometer that measures backscattered radiances at a spectral resolution of 0.42–0.63 nm (Levelt et al., 2006). Its wide field-of-view of 114° with a swath width of 2600 km yields daily global coverage with a spatial resolution of 13 km×24 km (Liu et al., 2010). OMI measures ozone profiles as well as other key air quality components such as NO₂, SO₂, and aerosol characteristics. In this article, we use the OMI tropospheric NO₂ column product from 2003 to 2020 to analyse long-term changes over different coastal and open ocean regions of the Indian Ocean (Section 6).

The TROPospheric Monitoring Instrument (TROPOMI) is a nadir-viewing imaging spectrometer on board the Copernicus Sentinel-5 Precursor satellite, which was launched in October 2017 for a mission of seven years. The satellite has a sun-synchronous orbit achieving near full-surface coverage on a daily basis. The TROPOMI instrument contains four spectrometers with three covering the ultraviolet-near infrared and one for the shortwave

infrared range. Key atmospheric species observed by TROPOMI include ozone, NO₂, SO₂, CO and aerosol properties. The TROPOMI tropospheric NO₂ column product (Boersma et al., 2018) shows improved spatial resolution ~~compared to previous version~~^{over previous}. The NO₂ retrieval algorithm is based on the NO₂ DOMINO retrieval previously used for OMI spectra with improvements made for all retrieval steps. In this article, we use the TROPOMI Level 2 NO₂ tropospheric column data product to show its distribution and seasonal variations (Section 5.1).

Pollutants (CO) from MOPITT

The Measurements of Pollution in the Troposphere (MOPITT) instrument is onboard NASA's Earth Observing System Terra spacecraft, measuring tropospheric CO since March 2000. The satellite is in a sun-synchronous polar orbit of 705 km allowing the instrument to make measurements in a 612 km cross-track scan with a footprint of 22 km × 22 km providing global coverage every 3 days. The MOPITT measurements provide vertical profiles and total columns of CO, which are useful to analyse the distribution, transport, sources and sinks on a global scale. CO retrieval products are generated with an iterative optimal-estimation-based retrieval algorithm based on the MOPITT calibrated radiances and a priori knowledge of CO variability. The recently released version 8 (V8) products benefit from updated spectroscopic information used in the radiative transfer model and improved methods for radiance bias corrections (Deeter et al., 2019). In this article, we use MOPITT V8 Level 3 monthly data (Near and Thermal Infrared Radiances, JIR) with day and night retrievals averaged to analyse the seasonal variation of surface CO distribution (Section 5.1).

Greenhouse gases (CH₄ and CO₂) from GOSAT

The Greenhouse Gases Observing Satellite (GOSAT/Ibuki) is a sun-synchronous polar orbit satellite that measures CO₂ and CH₄ from the stratosphere to the Earth's surface. The retrieval precision for CO₂ is smaller than 3.5 ppm (Yoshida et al., 2011) utilising the Thermal and Near Infrared Sensor for Carbon Observation – Fourier Transform Spectrometer, which operates in the shortwave and thermal emission bands. The GOSAT Level 3 product at a horizontal resolution of 2.5° × 2.5° has data gaps over the globe including a major portion of the Indian region during the monsoon season due to its limitation in retrieving CO₂ in the presence of clouds. This is rectified in the level 4 product that uses the Atmospheric Tracer Transport Model to incorporate ground-based observations and achieves a better distribution of CO₂ over the Indian Ocean (Nalini et al., 2018).

Observations from the Thermal and Near-infrared Sensor for carbon Observation-Fourier Transform Spectrometer (TANSO-FTS) onboard GOSAT in the thermal infrared (TIR) provide CH₄ profile information. While the sensitivity of TIR CH₄ observations is relatively low near the surface, the GOSAT/TANSO-FTS TIR instrument has been shown to have sufficient sensitivity to provide CH₄ information at the top of the boundary layer for the Indian subcontinent and Indian ocean region (Belikov et al., 2021). In this article we use GOSAT/TANSO-FTS CH₄ Version 1 (Level 2 V1) CH₄ data at 800 hPa averaged over 2009–2014 to analyse the seasonal variation of surface CH₄ distribution (Section 5.2).

Pollutants (NO₂) from GOME and SCIAMACHY

The Global Ozone Monitoring Experiment (GOME) is a UV/Visible spectrometer on the European polar sun-synchronous orbiting satellite ERS-2, launched in April 1995. It measures in 230–800 nm wavelength range, with a spectral resolution of 0.2–0.4 nm, and obtains global coverage at the equator after 3 days (Burrows et al. 1999). Problems with tape storage on ERS-2 led to the replacement of GOME by the Scanning Imaging Absorption Spectrometer for Atmospheric Chartography (SCIAMACHY), which was launched in 2002 on the European ENVISAT platform. It measures in the spectral range 240–2380 nm (Bovensmann et al., 1999).

Both instruments provide measurements of the mean columnar amount of tropospheric NO₂ ~~and allowed to facilitating~~ studies of its variations and long-term changes over the Indian subcontinent (Ghude et al., 2013; Mahajan et al., 2015a).

4. Regional sources and sinks

Atmospheric composition over the Indian Ocean is known to be impacted by the trace gas outflow from the surrounding continental land masses, long range transport and regional oceanic air-sea fluxes (Lawrence and Lelieveld, 2010). Here, we describe the distribution, seasonality and trends of continental and oceanic trace gas emissions important for the atmospheric composition over the Indian Ocean. Our study region includes East Africa, the Middle East, South Asia, East Asia, and Southeast Asia and is depicted in Fig. 23.

We use the latest versions of the Emissions Database for Global Atmospheric Research (EDGAR) to present continental pollution and greenhouse gas emissions over the last two decades. For air pollutants, EDGAR v5.0_AP for the period 1970–2015 is available (Crippa et al., 2020) and for greenhouse gases EDGAR v5.0_GHG for the period 1970–2015 (Crippa et al., 2019) can be used. The EDGAR datasets include continental emissions from the energy sector (i.e., power industry), industrial processes (i.e., manufacturing, industrial combustion), the transport sector (i.e., road transport, aviation), residential sources (small-scale combustion and waste treatment), and agriculture. Exhausts from ship engines as one of the major sources of air pollution over the open ocean are also included in the EDGAR emissions. The datasets are given at a high spatial resolution of 0.1° x 0.1°. The results shown in this section focus on the main pollutants CO, NO_x, and SO₂, and the greenhouse gases CH₄, N₂O, and CO₂. We also briefly discuss mercury emissions. The most recent year for which data is given (year 2015 for both air pollutants and greenhouse gases) is used here to present emissions strength and patterns representative for the last decade. Emission changes are calculated for the time period 2000–2015 and are shown in relative terms compared to the emissions in 2000. Emissions are averaged over East Africa, the Middle East, South Asia, East Asia, and Southeast Asia for a direct comparison of the regional contributions and the text and tables.

The ocean is an important source and sink to/from the atmosphere for many of the same gases mentioned above, as well as other climate- and chemically active compounds, such as DMS, isoprene, and halogen species. Below we will describe the net ocean fluxes of CO, CH₄, CO₂, N₂O, VOCs, DMS, isoprene, and bromoform (CHBr₃) as obtained from recent publications, placing special attention on monsoon related variability.

4.1 Pollutants

Among atmospheric pollutants, CO is considered to be one of the most important gases as it is highly toxic at elevated concentrations. Due to its intermediate lifetime of a few months (Seinfeld and Pandis, 2006), CO is much more variable in the troposphere than other atmospheric constituents with longer lifetimes and often used as a transport tracer. CO has an indirect radiative effect, since it scavenges the hydroxyl radical (OH), the cleaning agent of the atmosphere that otherwise would destroy the greenhouse gases CH₄ and O₃ (Daniel and Solomon, 1998). Another important pollutant is the family of the nitrogen oxides (NO_x) consisting of nitrogen dioxide (NO₂) and nitrogen oxide (NO). Tropospheric NO_x acts as a precursor for a number of harmful secondary air pollutants such as ozone and particulate matter and plays a role in the formation of acid rain. Breathing in raised levels of NO₂ can cause respiratory problems independently of negative health effects of other secondary pollutants. Once transported into the stratosphere, reactive odd-nitrogen species destroy ozone, in particular in the middle stratosphere near the ozone maximum (Portmann et al., 2012). SO₂ is another key component of gaseous air pollution. As for NO₂, exposure to SO₂ can harm the

human respiratory system. In addition, SO_2 can react with other compounds in the atmosphere to form small particles that contribute to particulate matter pollution. If oxidised within airborne water droplets, SO_2 produces sulphuric acid, which can be transported by wind over many hundreds of kilometres and deposited as acid rain. Atmospheric NH_3 is a pollutant which plays an important role in the formation of particulate matter, as well as in acidification and eutrophication of ecosystems (Lelieveld et al., 2015; Bauer et al., 2016).

The distributions of CO , NO_x , and SO_2 emissions are shown in Figures 23, 34 and 45, respectively. One of the common features of the spatial distribution of these emissions is that they generally coincide with the population distribution, such that high emissions appear in the densely populated areas. In East Asia, high-emission areas include northern China, the Yangtze River delta, Sichuan Basin, Korea, and Japan (not shown in the Figure). In South Asia, high emissions are distributed throughout northern India, Nepal, the southern point of India, and Bangladesh. In Southeast Asia, high emissions appear around some major cities including Bangkok, Hanoi, and Ho Chi Minh City, as well as Java. Similar to Southeast Asia, high-emission regions in East Africa are also around major cities like Kampala, Nairobi, and Addis Ababa. In the Middle East, high-emission regions are distributed around the Persian Gulf. Among the different source regions, East Asia and South Asia are main emitters. In 2015, the two regions accounted for 41% (East Asia) and 27% (South Asia) of the total CO emissions discussed here. East Asia is also a large emitter of NO_x (54%) and SO_2 (57%).

It is well known that pollution sources from Asia are characterized by inefficient combustion processes during biofuel and fossil fuel burning. For instance, the burning of biofuels such as wood, dung, and agricultural waste accounts for 18% of all CO emissions in Southeast Asia. Globally it only accounted for ~9% of all CO emissions in 2015, highlighting the role of biofuel burning in regions around the Indian Ocean. Inefficient combustion processes also occur during fossil fuel burning at lower temperatures and result in relatively low NO_x emissions and higher CO/CO_2 ratios, when compared to other industrialized areas around the globe. The incomplete fossil fuel combustion from the residential sector and road transportation are the two main sources contributing to the CO production, accounting for 29.5% and 29.0% of all CO emissions in our study region in 2015.




Figure 23. Annual mean CO emissions for 2015 (a) and relative change with respect to 2000 (b) from EDGAR V5.0_AP.




Figure 34. Same as Figure 23, but for NO_x.




Figure 45. Same as Figure 23, but for SO₂.

NO_x emissions mainly stem from high temperature combustion. Energy production, manufacturing industries and road transportation caused 30.7%, 25.8%, and 25.6% of all NO_x emissions in our study region in 2015, respectively. Manufacturing industries and energy (electricity and heat production) are also the two main contributors to the SO₂ emissions, accounting for 41% and 40%, respectively, of all SO₂ emissions in our study region in 2015. As per the 2000 Asian emission inventory, India has the second highest SO₂ emission (14%) after China (65%) with coal-burning power plants contributing to around half (47%) of the emissions in India (Kurokawa et al., 2013). About 40% of the thermal plants in India are located over the Indo-Gangetic Plains causing relatively high SO₂ emissions from this region (Fig. 45; Aswini et al., 2020). In addition, ship traffic leads to anthropogenic NO_x and SO₂ emissions directly over the open ocean with emissions concentrated along the major shipping lanes (e.g., Franke et al., 2009). In general, NO_x ship emissions can lead to substantial ozone enhancements and in turn to higher OH concentrations (Endresen et al., 2003).

Over the period 2000-2015, the emissions of all pollutants increased in almost all regions around the Indian Ocean, with CO emissions changing from 275.9 Tg yr⁻¹ to 350.3 Tg yr⁻¹, NO_x from 35.3 Tg yr⁻¹ to 58.3 Tg yr⁻¹, and SO₂ from 41.8 Tg yr⁻¹ to 61.1 Tg yr⁻¹. Between 2000 and 2015, CO emissions increased particularly along the Mekong River, north of the Persian Gulf, in Afghanistan and East Africa, while CO emissions in most regions of East Asia decreased despite a comparably low overall increase (15%, Table 2, Fig. 23b). NO_x emission increases

show a different pattern and are relatively high in most regions around the Indian Ocean, with peaks in East Asia, South Asia and East Africa (Fig. 34b). SO₂ emission changes show a similar distribution as the NO_x changes, with peaks along the Mekong River and in East Africa (Fig. 45b). Ship traffic in the Indian ocean has seen the largest increase worldwide between 1992 and 2012, especially on well-defined shipping lanes, such as the Red Sea-Arabian Gulf-Asia route or the Asia-Cape Town route (Tournadre, 2014). The overall increase of the pollutant emissions shows pronounced variations from region to region (Table 2) with the highest rate of all three pollutants emission increases found in South Asia.

In particular, for the time period after 2012, satellite measurements have shown pronounced regional SO₂ and NO₂ pollution changes. A decrease of SO₂ pollution from the North China Plain has been noted since 2011 as a result of government efforts, while SO₂ and NO₂ emissions from India have continued to grow at a fast rate (Krotkov et al., 2016). Recent emission estimates suggest that during 2013–2017, anthropogenic emissions from China decreased by 23 % for CO, 21 % for NO_x, and 59 % for SO₂ over this period as a consequence of the implementation of active clean air policies (Zheng et al., 2018).

Table 2. Emissions of CO, NO_x, and SO₂ from different regions in 2015 and their increase with respect to 2000.

	Emission in 2015 (Increase with respect to 2000)				
	East Africa	Middle East	South Asia	East Asia	Southeast Asia
CO	26.48 Tg yr ⁻¹ (40%)	16.76 Tg yr ⁻¹ (18%)	93.97 Tg yr ⁻¹ (50%)	144.21 Tg yr ⁻¹ (15%)	68.91 Tg yr ⁻¹ (25%)
NO _x	0.89 Tg yr ⁻¹ (76%)	7.27 Tg yr ⁻¹ (65%)	12.47 Tg yr ⁻¹ (101%)	31.39 Tg yr ⁻¹ (56%)	6.28 Tg yr ⁻¹ (54%)
SO ₂	0.43 Tg yr ⁻¹ (67%)	7.55 Tg yr ⁻¹ (54%)	13.09 Tg yr ⁻¹ (120%)	34.63 Tg yr ⁻¹ (29%)	5.42 Tg yr ⁻¹ (40%)

Unfortunately, measurements of oceanic CO emissions from the Indian Ocean are sparse. We only know of unpublished data sets (D. Arevalo-Martinez, personal communication) from one GEOMAR campaign (OASIS) and a series of NASA-SAGA cruises (www.saga.pmel.noaa.gov, eastern open Indian Ocean, summer 1987). Net fluxes covering the northern to southern extent of the Indian Ocean range from ~0.1 to ~1.4 Mg km⁻² yr⁻¹, as CO is always supersaturated in the surface ocean (Conte et al., 2019 and references therein), and values are similar to ship emissions but considerably smaller than continental emissions (Fig. 23). CO is produced in the surface ocean from organic material photochemistry and biological processes (Conte et al., 2019). Available data from the western Indian Ocean suggests that the most significant meridional gradients occur due to open ocean upwelling at 5°S–10°S. CO emissions are high from 5°–15°S, but to the north and south of this region, emissions decrease to zero with seasonal variations occurring due to upwelling changes. In the eastern Indian Ocean, seasonal variability is expected in association with surface productivity changes in the Seychelles–Chagos Thermocline Ridge. However, no seasonal cycle can be detected in available measurements from this region, and it is not clear if this is a real feature or caused by

the lack of data. Additional variability is expected in coastal regions, since large amounts of seasonally discharged runoff supply terrestrial organic material that serves as a precursor to CO₂ marine photoproduction.

The atmospheric pollutant mercury is transported around the globe as gaseous elemental mercury, eventually oxidizing to divalent mercury. The latter is known to deposit to the surface from where it can be taken up into food webs and be transformed to highly toxic species endangering humans and ecosystems (Selin et al., 2007). Atmospheric mercury is released from anthropogenic activities, such as coal-fired power plants, metal smelting, and waste incineration (Pacyna et al., 2005; Streets et al., 2005). Emissions associated with artisanal and small-scale gold mining account for almost 38% of the global total emission (UN-Environment, 2019). Mercury is also emitted from the oceans, soils, terrestrial vegetation, and biomass burning. These ‘natural’ emissions include some anthropogenic fraction related to the recycling of previously deposited mercury (Mason and Sheu, 2002). Based on 2015 inventories, Asia is responsible for a large part of the emissions (49%), which primarily stem from East and Southeast Asia. While emissions in North America and the European Union have shown moderate decreases, increased economic activity, notably in Asia, and the use and disposal of mercury-added products have led to a global increase of approximately 20% between 2010 and 2015 (UN-Environment, 2019).

For NH₃, East Asia and South Asia are the two main contributors, which account for 38.9% and 32.3% of the total emissions, respectively (not shown here). From 2000 to 2015, emissions of NH₃ in the regions around the Indian Ocean documented by EDGAR increased by 22.5%. Agricultural activities dominate the ammonia emissions, with about 56.7%, 18.4% of the emissions from the sectors of direct soil emission and manure management. Besides, long-term satellite measurements (van Damme et al., 2018) show other hotspots of ammonia emission not well represented in EDGAR inventory, most of which are associated with either high-density animal farming or industrial fertilizer production.

4.2 Greenhouse gases

CO₂ concentrations have been increasing steadily over the last decades reaching a new annual global mean record high in 2019 of 409.8 ± 0.1 ppm (Blunden and Arndt, 2020). Due to its high atmospheric abundance and long atmospheric lifetime, CO₂ is the most important of Earth’s long-lived greenhouse gases. In addition to its impact on climate, CO₂ is responsible for ocean acidification as it produces carbonic acid when it dissolves in the ocean. CH₄ is also a very effective greenhouse gas and the second-largest contributor to anthropogenic radiative forcing since preindustrial times after CO₂. In the troposphere, CH₄ acts to reduce the atmosphere’s oxidizing capacity. It has a relatively short atmospheric lifetime of about 12.4 years (Myhre et al., 2013) and exhibits a strong seasonal cycle as well as a distinct gradient across the equator. Despite its relatively low atmospheric concentrations, N₂O is the third anthropogenic greenhouse gas after CO₂ and CH₄ in terms of radiative forcing (Ciais et al., 2014). Due to its long atmospheric lifetime of about 116 years (Prather et al., 2015) and large infrared absorption capacity per molecule, N₂O is a much more efficient greenhouse gas than CO₂ with a global warming potential of 265 over a 100-year time span. In the stratosphere, reaction with O(¹D) leads to the production of NO (Seinfeld and Pandis, 2006), which is involved in chemical ozone depletion. As a consequence, N₂O has been estimated to be the main emitted ozone-depleting substance of the 21st century (Ravishankara et al., 2009; Butler et al., 2016).

Anthropogenic greenhouse gas emissions in the regions surrounding the Indian Ocean generally correspond to economic activities. As the largest emerging economies, East Asia and South Asia are the main emitters of CH₄ (Fig. 56) and N₂O (Fig. 67) with emission centres in the Indo-Gangetic Plain, northern China and Java. In 2015, East Asia and South Asia accounted for 37%

and 26% of the total CH₄ emission, as well as 43% and 26% of the total N₂O emission discussed here. Among the regions surrounding the Indian Ocean, East Asia is also the largest CO₂ emitter causing 68% of the total CO₂ emissions in our study region in 2015 (Fig. 78).

Atmospheric CH₄ has anthropogenic and natural sources, with the latter including natural wetlands, livestock, termites, hydrates and forest fires. Anthropogenic sources account for the majority of all emissions and can be split into biogenic and non-biogenic sectors. Almost a quarter (23%) of the CH₄ emissions in our study region stems from enteric fermentation (livestock farming), which acts as the primary source in South Asia and East Africa. Rice cultivation in Asia is responsible for 19% of CH₄ emissions in our study region causing a systematic seasonal pattern with peak emissions during the fully-grown stage in September and October (Pathak et al., 2005). Other main sources of CH₄ are solid fuels (17%), mainly from East Asia and Southeast Asia, and oil and gas production (14%), mainly from the Middle East.

N₂O emissions are linked to the biogeochemical cycle of nitrogen and are thus impacted by anthropogenic use of fertilizer and industrial activities that lead to the atmospheric deposition of reactive nitrogen (e.g., Davidson, 2009). More than half of the N₂O emissions in our study region (56%) are directly from managed soils and can be quite heterogeneous with spatial patterns revealing hot spots in agricultural areas in China and the Indo-Gangetic Plains (Ito et al., 2018; Fig. 67). Furthermore, the N₂O emissions from managed soils are characterized by a pronounced seasonal cycle and interannual variability, primarily in response to meteorological conditions and nitrogen inputs. In particular, N₂O emissions are correlated with soil moisture (Raut et al., 2015), leading to strongly enhanced emissions in South Asia during summertime when high precipitation events occur.




Figure 56. Annual mean CH₄ emissions for 2015 (a) and relative change with respect to 2000 (b) from EDGAR V5.0_GHG.

Similar to the air pollutants discussed above, the overall CH₄ and N₂O emissions increased significantly over the period 2000–2015 from 135.7 Tg yr⁻¹ to 182.4 Tg yr⁻¹ and from 2.80 Tg yr⁻¹ to 3.51 Tg yr⁻¹, respectively. Increasing CH₄ emission in South Asia (Fig. 56b) have been linked to increased rice cultivation area, natural wetlands, and warmer climate (Tian et al., 2015). Increasing N₂O emissions (Fig. 67b) are believed to stem from intensified crop production and nitrogen fertilizer use as well as higher air temperatures (Raut et al., 2015). While not being the main emitter, East Africa is the region with the fastest increase of CH₄ and N₂O emissions among the regions discussed here (Fig. 56b and 67b, Table 3). A recent study suggested that east African wetlands could account for up to a third of the spike in global CH₄ emissions between 2010 and 2016, with most of this coming from the South Sudanese wetland, one of the largest freshwater ecosystems in the world (Lunt et al., 2019).

Table 3. Emissions of CH₄, N₂O, and CO₂ from different regions in 2015 and their increase in respect to 2000.

Emission in 2015 (Increase respect to 2000)					
	East Africa	Middle East	South Asia	East Asia	Southeast Asia
CH ₄	12.33 Tg yr ⁻¹ (50%)	21.47 Tg yr ⁻¹ (50%)	47.65 Tg yr ⁻¹ (24%)	68.08 Tg yr ⁻¹ (34%)	32.83 Tg yr ⁻¹ (38%)
N ₂ O	0.36 Tg yr ⁻¹ (51%)	0.20 Tg yr ⁻¹ (21%)	0.91 Tg yr ⁻¹ (36%)	1.52 Tg yr ⁻¹ (15%)	0.52 Tg yr ⁻¹ (26%)
CO ₂	69.3 Tg yr ⁻¹ (137%)	2062.30 Tg yr ⁻¹ (93%)	2565.10 Tg yr ⁻¹ (125%)	13041.02 Tg yr ⁻¹ (127%)	1430.79 Tg yr ⁻¹ (79%)




Figure 67. Same as Figure 56 but for N₂O

For CO₂, the majority of the emissions in our study region stem from East Asia related to two main sectors: electricity and heat production (41%) and manufacturing industries (23%). Over the period 2000–2015, the CO₂ emissions in our study region more than doubled from 8790 Tg yr⁻¹ to 19168 Tg yr⁻¹. Especially in East Asia and South Asia, they grew at very fast rates with increases of 127% and 125%, respectively. Despite the apparent policy breakthrough leading to the Paris Agreement in 2015, CO₂ emissions from fossil fuel and industry have continued to increase over the recent years. According to the latest estimates from the Global Carbon Project, the expected growth of global emissions in 2019 will be almost entirely due to China and India with expected annual growth rates of 2.6% and 1.8%, respectively (Peters et al., 2020).

The ocean is also a source and sink of greenhouse gases. Compared to the terrestrial sources, the ocean is just a minor contributor to the atmospheric CH₄, accounting for 1–13% of the global atmospheric CH₄ budget (Saunois et al., 2016). The concentration of CH₄ in the Indian Ocean is characterized by a sharp decrease offshore (Naqvi et al., 2010a). Due to the large geographical changes in surface saturation and wind speed, the sea-to-air flux of CH₄ varies strongly in the northern Indian Ocean. Highest emissions were observed during the southwest monsoon in the Arabian Sea (~64 μmol m⁻² d⁻¹), and the estimated overall CH₄ emission from Arabian Sea amounts to 0.1–0.2 Tg yr⁻¹ (Naqvi et al., 2005), which is much smaller than the total terrestrial emissions mentioned above (~182 Tg yr⁻¹ in 2015).

Unlike CH₄, the ocean is a major source of N₂O, accounting for at least one third of global N₂O emissions (Bange, 2006). Intense N₂O emissions are usually found in upwelling regions with OMZs (Codispoti, 2010), such as in the Arabian Sea and Bay of Bengal (not shown). The South Asian monsoon drives intense seasonal changes of upwelling in both OMZs, thus affecting the regional N₂O productivity and emissions. The upwelling in the Arabian Sea is most intense during the South Asian summer monsoon, leading to high oceanic N₂O production (Naqvi et al., 2010a) and emissions of 0.34–0.99 Tg N₂O yr⁻¹ (Naqvi et al., 2010b), representing 2–31% global oceanic N₂O emissions (Suntharalingam et al., 2019). However, the estimate by Sudheesh et al. (2016) in the south-eastern Arabian Sea, based on the measurements from Mangalore and Kochi, is almost four times lower than previous estimates. Raes et al. (2016) proposed that the south-eastern Indian Ocean could be both a source and sink of N₂O, suggesting great uncertainty of the oceanic emission of N₂O in the Indian Ocean. The upwelling driven by the summer monsoon also occurs in the Bay of Bengal, however, it is attenuated by the intense precipitation and pronounced freshwater discharge from the Ganges, yielding lower N₂O productivity (Singh and Ramesh, 2015) and much smaller emission (~0.03–0.11 Tg N₂O yr⁻¹, Naqvi et al., 1994, 2010b). There is some indication that increased nitrogen deposition, due to anthropogenic perturbation, already influences the air-sea flux of N₂O in the northern Indian Ocean and will continue to do so into the future (Suntharalingam et al., 2019). Due to sparse measurements, it is impossible to compare the oceanic N₂O emission of the Indian Ocean with the terrestrial emissions directly. However, the emissions from the Arabian Sea alone are about 9.6–28% of the terrestrial emissions of the study region (~3.5 Tg yr⁻¹ in 2015), suggesting the importance of the oceanic source in the Arabian Sea region. Enhanced N₂O sea-to-air fluxes were also found in a zonal band between 5°S and 10°S as a result of wind-driven upwelling during the OASIS research cruise in 2014 (Ma et al., 2018).

While the global oceans act as a net sink of CO₂, absorbing about 25% of the annual anthropogenic CO₂ emission (Le Quéré et al., 2018), the air-sea exchange of CO₂ varies at different spatial and temporal scales. The northern Indian Ocean is a net source of CO₂ to the atmosphere, while the southern Indian Ocean is a net sink (e.g., Sarma et al., 2013). Based on the inorganic carbon data collected in the Indian Ocean in 1995, for example, Bates et al. (2006) suggested that the BoB is a net oceanic source of atmospheric CO₂. Takahashi et al. (2009) constructed the climatological mean distribution for the sea surface water pCO₂ over the global oceans based on the observations from 1970 and 2007, and they suggested that the BoB is near equilibrium for CO₂. By examining numerical results produced by various ocean biogeochemical models and different atmospheric inversions, Sarma et al. (2013) argued that the BoB is a small annual net sink region for the atmospheric CO₂. Other studies report that on an annual scale, specific regions in the Bay of Bengal emit between ~1.61 and 2.45 ± 0.49 Mg CO₂ yr⁻¹ km⁻² (Dixit et al., 2019; Ye et al., 2019). Significant impacts of tropical cyclones (Ye et al., 2019), biological productivity (Chakraborty et al., 2018) and freshwater discharge (Sarma et al., 2011) on the CO₂ air-sea exchange have been suggested for the Bay of Bengal. If compared to anthropogenic emissions of more than 500 Mg CO₂ yr⁻¹ km⁻² over large areas (Fig. 78a), the contribution of the northern Indian Ocean to the atmospheric CO₂ is relatively low. Analysing the Surface Ocean CO₂ Atlas (SOCATv2019; Bakker et al., 2016) demonstrates that in the Indian Ocean only a few CO₂ measurements are available for the last decade, especially when compared with the 1990s (Fig. 78c and d), making it impossible to assess the long-term changes of CO₂ air-sea exchange in this region. Northern Indian Ocean CO₂ flux variability may be influenced by ENSO events (Valsala and Maksyutov, 2013) and hydrographic features (i.e., eddies, Valsala and Murtugudde, 2015).




Figure 78. Annual mean CO₂ emissions for 2015 (a) and relative change with respect to 2000 (b) from EDGAR V5.0_GHG. Surface water fCO₂ observations (color shading, unit: μatm) over the Indian Ocean for 1990–2000 (c) and 2010–2019 (d) from SOCATv2019 (Bakker et al., 2016).

During the OASIS campaign in the western Indian Ocean (Zavarsky et al., 2018a), both positive and negative CO₂ fluxes were observed based on the direct eddy covariance flux technique. South of the equator, average values were $0.2 \text{ Mg day}^{-1} \text{ km}^{-2}$ and $-0.28 \text{ Mg day}^{-1} \text{ km}^{-2}$, respectively, making this region a net sink of CO₂. These results are consistent with Chen et al. (2011), who found significant spatial and temporal variability in the southern Indian Ocean carbon sink. However, by comparing campaigns that occurred from 1999–2000 to those from 2004–2005, Chen et al. (2011) imply that the sink of the southern Indian Ocean is weakening. A decadal variability analysis from 1991–2007 of dissolved CO₂ in surface seawater in the southern Indian Ocean (20° – 55°S) suggests that it increased at a faster rate than atmospheric CO₂ (Metzl, 2009), indicating that the ocean carbon sink weakened. The authors suggested that the reduction was related to variability in the Southern Annular Mode. The weakening of Indian Ocean carbon sink has also been found in a recent modelling study (DeVries et al. 2019). The southern Indian Ocean CO₂ fluxes are driven by both the solubility and biological pumps, with indications that their variability is driving by ENSO forcing (Valsala et al., 2013).

Carbonyl sulphide (COS) is another important long-lived trace gas that acts as a greenhouse gas in the troposphere and as the main precursor of aerosols in the stratosphere (Brühl et al. 2012; Kremser et al. 2016). The ocean is the main source of COS to the atmosphere, previously estimated at $441\text{--}542 \text{ Gg COS yr}^{-1}$, globally, but a revision of the vegetation sinks has led to the hypothesis that the ocean source might be stronger than previously calculated. The missing ocean source is hypothesised to be in the tropics (e.g., Suntharalingam et al., 2008; Glatthor et al., 2015). Launois et al. (2015) modelled oceanic concentrations and emissions for this region that are approximately an order of magnitude higher, but the values do not agree with the albeit sparse COS measurements that exist in the Indian Ocean. A recent measurement and modelling

study on the OASIS campaign has shown that, in fact, the ocean source of COS is not higher than previously determined (Lennartz et al., 2017). Daily integrated air-sea fluxes computed for the southern Indian Ocean ranged between -0.045 and -0.000375 g COS km⁻², indicating that the Indian Ocean may be a net sink for COS. In addition, COS is produced in the atmosphere from DMS and carbon disulphide (CS₂) oxidation, both of which are emitted from the ocean (Chin and Davis, 1993; Watts, 2000). These pathways increase the ocean source of COS indirectly, but do not account for the full missing ocean source (Lennartz et al., 2017). Campbell et al. (2015) and Lennartz et al. (2017) point to anthropogenic emissions of COS from Asia to close the gap and, indeed, Lee and Brimblecombe (2016) find twice as much COS in the atmosphere from anthropogenic emissions than previously thought. They report that anthropogenic COS emissions account for approximately one third of global emissions and originate from the paper industry and biofuel and coal combustion. Another study suggests that COS emission from domestic use coal combustion only in China would be at least 57.2 ± 10.5 Gg COS yr⁻¹, an order of magnitude greater than recent estimates of COS emissions from the total coal combustion in China (Du et al., 2016).

4.3 VOCs and short-lived gases DMS, isoprene and bromoform

Volatile organic compounds (VOCs)

Volatile organic compounds (VOCs), such as alkanes, alkenes, and alkynes, are highly reactive throughout the troposphere, influencing oxidants (OH, O₃, NO₃) (Williams et al. 2010), peroxyacetyl nitrate (PAN), a reservoir for reactive nitrogen compounds, and forming secondary organic aerosols (SOA) (Singh et al., 1995; Blando and Turpin 2000). Oxygenated VOCs play a special role in upper tropospheric OH production (e.g., Singh et al., 1995).

The Middle East is a hotspot of VOC emissions, such as ethane and propane, from oil and gas production. Interestingly, a recent study found that in addition deep water masses can be a large source of VOCs that is comparable with total anthropogenic emissions from individual Middle East countries (Bourtsoukidis et al., 2020). Other biogenic sources of VOCs include surrounding forested areas (Duflot et al., 2019). Formaldehyde (HCHO) is emitted via biomass burning and fuel combustion. In addition, it has a secondary source via reactions between OH and CH₄. Ship fuel combustion has recently been targeted a large source (Gopikrishnan et al. 2021). This was first postulated by Marbach et al. (2009) using satellite measurements of HCHO and shown again by De Smedt et al. (2021).

In addition to the Bourtsoukidis et al. (2020) study cited above, the Indian Ocean has been shown to be a main source of VOCs at other specific locations, such as at the Maïdo Observatory (Duflot et al., 2019). Evidence of alkene emissions from the Indian Ocean and specifically the Bay of Bengal was presented by Sahu et al. (2010; 2011) by demonstrating that the ratios of ethene/propene are in line with fresh oceanic emissions (2.3 ppt/ppt). Alkene emissions seem to be controlled by dissolved organic carbon, wind speed, and sunlight. The eastern Indian Ocean may be a source of biogenic VOCs, leading to the production of acids (diacids, oxocarboxylic acids, and α -dicarbonyls) in the atmosphere (Yang et al., 2020). The highly productive Arabian Sea OMZ area has been associated with enhanced production of alkanes and alkenes related to *Trichodesmium* and *Thalassiosira* phytoplankton species (Tripathi et al., 2020).

1 Dimethylsulphide (DMS)

2 Marine DMS is the largest source of biogenic sulphur to the atmosphere. DMS is produced
3 from the algal derived precursor, dimethylsulphoniopropionate (DMSP), which is cleaved by
4 marine microbes to form DMS. Only a small fraction of this DMS is released to the atmosphere.
5 The seminal CLAW hypothesis proposed a feedback loop between marine biogenic DMS
6 production, emissions, and climate, via aerosol and cloud formation (Charlson et al., 1987),
7 triggering decades of research on DMS cycling in the ocean and emission to the atmosphere.
8 Lana et al. (2011) is the most comprehensive and up-to-date monthly DMS concentration and
9 flux climatology resulting from this large body of research. Unfortunately, measurements in the
10 Indian Ocean are sparse and most values in the climatology are interpolated, with only 6271
11 non-uniformly spaced data points over 40 years available in the Indian Ocean.

12 DMS emissions exhibit clear seasonality, with the highest fluxes, basin wide, evident during
13 the summer monsoon period. According to Lana et al. (2011), the summertime values in the
14 Indian Ocean are a global hotspot for DMS emissions. The largest values are found in the
15 Arabian Sea (Fig. 89). High biological productivity associated with the upwelling areas off
16 north-east Africa and the Arabian Sea are strongly correlated with the monsoon cycle (Yoder
17 et al., 1993). Combined with the strong, steady winds during the summer monsoon, fluxes of
18 DMS reach their peak. Lowest fluxes are computed during the spring transition period, likely
19 associated with low productivity and low wind speeds. Year round, there is a relatively large
20 flux area around 15°S, which migrates north and south according to the summer hemisphere
21 and is related to biogenic processes in the upwelling. Winds are always relatively high in this
22 region throughout the year, which also enhances the flux (Fig. 1). Maximum emissions of over
23 650 kg DMS k⁻² yr⁻¹ for the Arabian Sea (Lana et al., 2011) translate to slightly larger S flux to
24 the atmosphere from DMS than that from SO₂ ship emissions (approximately 335 kg S k⁻² yr⁻¹
25 from DMS and 250 kg S k⁻² yr⁻¹ from SO₂).

26 Gali et al. (2018) used satellite-based proxies to estimate the DMS concentration climatology
27 and reported that the Lana et al. (2011) climatology overestimates the DMS in the Indian Ocean
28 region by 25–50% in all the seasons. DMS direct flux measurements using the eddy covariance
29 technique and ocean concentration measurements were performed during the OASIS campaign
30 in order to compare directly with the Lana climatology in the western tropical Indian Ocean
31 (Zavarsky et al., 2018a). The oceanic DMS concentrations were found to be lower than those
32 in the climatology, but the difference was more pronounced south of 16°S where measured
33 values were a third of those in the climatology. North of 16°S, the measured ocean
34 concentrations were in better agreement with those in the climatology until the vicinity of the
35 Maldives, where they again were lower by a factor of three. The measured fluxes, subsequently,
36 were lower than the climatology for the region by approximately 60% on average. This was
37 attributed to lower measured oceanic concentrations, as well as lower measured wind speeds
38 than used in the climatology and a different gas transfer parameterisation. The directly derived
39 gas transfer parameterisation was linearly dependent on wind speed, while the climatology uses
40 a quadratic wind speed dependence. Nonetheless, the Indian Ocean appears to be a hotspot for
41 DMS emissions during the summer monsoon, but with more likely sulphur loading to the
42 atmosphere on the order of half of that from SO₂ ship emissions.




Figure 89. DMS emission for (a) summer monsoon (June – August), (b) winter monsoon (December – February), (c) spring transition (April – May) and (d) autumn transition (October) periods from Lana et al. (2011).

Isoprene

Isoprene (2-methyl-1,3-butadiene) is a biogenic VOC and accounts for half of the total global biogenic VOCs in the atmosphere (Guenther et al., 2012). Most is emitted from terrestrial vegetation (400–600 TgC yr⁻¹, Guenther et al., 2006; Arneth et al., 2008). The ocean source strength is much lower, and the magnitude is debated (Carlton et al., 2009), with most estimates lower than 1 TgC yr⁻¹ (Palmer and Shaw, 2005; Arnold et al., 2009; Gantt et al., 2009; Booge et al., 2016). It is known from laboratory studies that phytoplankton produce isoprene (Exton et al., 2013 and references therein), but only a few studies have performed direct measurements of marine isoprene concentrations worldwide.

Emitted isoprene affects the oxidative capacity of the atmosphere through ozone and OH interactions and is a source for SOAs (Carlton et al., 2009). Due to the short atmospheric lifetime of minutes to a few hours, terrestrial isoprene does not reach the atmosphere over much of the ocean and, thus, marine emissions of isoprene could play an important role in SOA formation on regional and seasonal scales, especially in association with increased production during phytoplankton blooms (Hu et al., 2013). Furthermore, isoprene SOA yields increase under acid-catalysed particle phase reaction in low-NO_x conditions, which dominate over open oceans regions (Surratt et al., 2010), and which are significantly higher than that during neutral aerosol experiments (Henze and Seinfeld, 2006). Here we provide data from a published modelling study from the OASIS campaign with input variables from 2014 to assess seasonal isoprene fluxes to the atmosphere from the Indian Ocean (Booge et al., 2016; 2018).




Figure 940. Same as Figure 89 but for isoprene from Booge et al. (2016).

Isoprene fluxes to the atmosphere change seasonally, with the highest values computed during the summer monsoon over the entire Indian Ocean extent (Fig. 940). The summer values are the second highest values, globally, during that season, following the Southern Ocean. Lowest isoprene emissions in the Indian Ocean are found in the spring transition period. This seasonal pattern is similar to the DMS emissions pattern. Computed fluxes during the winter monsoon are high in the northern region of the Indian Ocean, especially in the Arabian Sea. A belt of relatively high isoprene fluxes can be seen in the autumn transition period around 15°S, but the values are lower than the highs seen during the summer (basin wide) and winter (Arabian Sea) monsoon season. Unlike DMS, this is only visible in the summer and autumn seasons. Isoprene production rates are phytoplankton functional type dependent and are driven further by light, SST, salinity, and nutrients (Booge et al., 2018). High light and high SST favour higher production, while high salinity and high nutrients lead to lower production. The combination of the direct influence of wind speeds on fluxes and the interaction of the environmental factors and isoprene production leads to the seasonal patterns.

Halogens

Halogenated very short-lived substances (VSLs, with lifetimes shorter than 6 months) from the ocean, such as bromoform (CHBr_3), dibromomethane (CH_2Br_2) and methyl iodide (CH_3I), contribute to atmospheric halogen loading and ozone depletion (Engel and Rigby et al., 2019). The oceanic CHBr_3 surface concentrations are spatially and temporally highly variable. Natural production of CHBr_3 involves marine organisms such as macroalgae and phytoplankton (Gschwend et al., 1985), while CH_2Br_2 is formed in parallel and correlates with CHBr_3 in water and air (Tokarczyk and Moore, 1994). A recent study suggests that heterotrophic processes in

the ocean can increase the flux of CH_2Br_2 from the sea to the atmosphere (Mehlmann et al., 2020). Enhanced emissions of brominated VSLs coincide with biologically active equatorial and coastal upwelling regions (Quack et al., 2007) and the distribution of macroalgae and anthropogenic sources along the coasts (Carpenter and Liss, 2000; Maas et al., 2020). Iodinated VSLs such as CH_3I , on the other hand, show elevated oceanic abundances in the subtropical gyre regions in agreement with identified production by photochemical reactions (Richter and Wallace, 2004).

Various CHBr_3 emission inventories have been derived from the extrapolation of measurement-based data (Ziska et al., 2013; Fiehn et al., 2017), oceanic modelling (Stemmler et al., 2015), top-down atmospheric modelling approaches (Liang et al., 2010) and a data-oriented machine-learning algorithm (Wang et al., 2019). Overall, large differences between CHBr_3 emission inventories exist with the observation-based, bottom-up emissions (Ziska et al., 2013) being most consistent with atmospheric measurements in the tropics (Hossaini et al., 2013). All inventories agree on the tropical Indian Ocean being a productive source region of CHBr_3 .




Figure 104. Bromoform (CHBr_3) emission for (a) summer monsoon (June – August), (b) winter monsoon (December – February), (c) spring transition (April – May) and (d) autumn transition (October) periods based on static surface concentrations and ERA-Interim meteorology for 2014 (Fiehn et al., 2018).

Here we show the most recent bottom-up CHBr_3 emission inventory (Fiehn et al., 2018) based on two campaigns in the marginal seas (Yamamoto et al., 2001; Roy et al., 2011), one campaign in the open Indian Ocean (Fiehn et al., 2017) and extrapolations of measurements from other oceans (Ziska et al., 2013) in Fig. 104. The emission inventory is based on static surface

concentration maps generated from atmospheric and oceanic surface ship-borne in-situ measurements collected within the HalOcAt (Halocarbons in the ocean and atmosphere) database project (<https://halocat.geomar.de>, last access July 2020). While the concentration maps do not provide any temporal variability, the emission parameterisation is based on monthly mean meteorological ERA-Interim data allowing for relative emission peaks related to maxima in the horizontal wind ~~and SST fields and sea surface temperature~~.

The CHBr_3 emissions peak along the Northern Hemisphere (NH) coastlines due to macroalgae production and anthropogenic sources ($3.3\text{--}6.6\text{ kg yr}^{-1}\text{ km}^{-2}$), the central Bay of Bengal (up to $11.1\text{ kg yr}^{-1}\text{ km}^{-2}$) and the southern tropical Indian Ocean ($2.2\text{ kg yr}^{-1}\text{ km}^{-2}$). The coastal emissions in the Indian Ocean of the bottom-up inventory presented here agree well with other emission estimates (Fiehn et al., 2018). High emissions along the coasts of Somalia and Oman due to coastal upwelling detected in biogeochemical modelling studies (Stemmler et al., 2015) are not captured here due to missing CHBr_3 measurements in this biogeochemical regime. The emissions show a pronounced seasonal cycle with a peak during the summer monsoon period (Fig. 104) due to higher wind speeds over the whole Indian Ocean during this time of year (Fig. 1).

Once destroyed in the atmosphere, brominated VSLs contribute to the family of inorganic bromine (Br_y) consisting of bromine radicals such as bromine monoxide (BrO) and non-radical reservoir species such as HBr and HOBr . Inorganic bromine has also other anthropogenic and natural sources including methyl bromide, which is a product of biomass burning, leaded fuel combustion, plant and marsh emissions, as well as soil fumigation (Mano and Andreae, 1994). Inorganic bromine can also be released when sea salt is exposed to the atmosphere from young sea-ice surfaces, frost flowers, snowpack, seawater and marine aerosols (Hay et al., 2007).

Similar to bromine, inorganic iodine such as iodine oxide (IO) is produced through the degradation of its organic precursor CH_3I and other short-lived iodinated VSLs. The primary source of iodine to the marine boundary layer, however, are believed to be inorganic iodine emissions at the ocean surface from reactions of dissolved iodide with deposited gas-phase ozone (Carpenter et al., 2013). Once in the atmosphere, inorganic iodine plays an important role for the boundary layer chemistry by influencing the oxidising capacity through catalytically removing O_3 and altering the HO_x and NO_x balance.

5. Atmospheric composition

During the South Asian summer monsoon, clean air dominates the atmospheric composition over the Indian Ocean, leading to a completely different chemical regime than observed during wintertime. As the ITCZ moves over the Indian subcontinent (Waliser and Gautier, 1993), air mass transport via steady winds is directed from the ocean towards the land and anthropogenic pollutants are confined to the continents. During this period, the intense summer monsoon rainfalls also effectively remove many soluble species from the continental boundary layer. During the winter monsoon and transitional months, the wind pattern is reversed with continental air masses being transported towards and over the open ocean environment. This leads to an increase in the anthropogenic origin trace gases and aerosol loading over the Indian Ocean. Based on studies and campaigns that took place before 2010, Lawrence and Lelieveld (2010) provided a comprehensive review of atmospheric composition over the Indian Ocean. Here we summarize their most important findings and report on new datasets and results that have emerged since then.

5.1 Pollution and O₃

The surface distributions of anthropogenic pollutants and ozone levels are often highly correlated as the majority of tropospheric ozone formation occurs when CO, NO_x, and other ozone precursor gases react in the atmosphere in the presence of sunlight. Tropospheric ozone is a greenhouse gas with a relatively short lifetime on the order of several weeks and is therefore considered a near-term climate forcer. The direct radiative forcing of ozone is estimated to be $0.40 \pm 0.20 \text{ W m}^{-2}$ (IPCC, 2013) and additional indirect radiative forcings can result from its impact on vegetation, carbon uptake and methane lifetime (e.g., Lombardozzi et al., 2015; Fiore et al., 2008). Ozone also acts as an environmental pollutant detrimental to human health, and crop and ecosystem productivity (e.g., Monks et al., 2015). Changes of ozone precursor emissions, low-frequency climate variability and long-term anthropogenic climate change all impact the quantity and distribution of tropospheric ozone (e.g., Barnes et al., 2016).

The distribution and variations of pollutants and ozone over the Indian Ocean have been investigated during individual campaigns (e.g., ICARB) and can be derived from continuous in-situ (e.g., ozone profiling at Indian coastal stations) and satellite measurements (e.g., OMI and MOPITT). The various data sets offer different advantages, allowing for a wide range of applications. While the in-situ measurements are characterised by higher resolution and lower uncertainties, they are usually limited in space (stations) or time (campaigns). The satellite measurements on the other hand, offer comprehensive spatial coverage and extend over longer time periods, but suffer from limited vertical resolution and higher uncertainties. Most available Indian Ocean campaigns and studies investigate O₃, CO and NO_x. As a result, the following sections focus on the distribution and variability of these three gases, while SO₂ and mercury are only discussed briefly.

Carbon monoxide (CO) and nitrogen oxides (NO_x)

The distribution of the major pollutant CO over the Indian Ocean is well known from MOPITT satellite measurements (e.g., Ghude et al., 2011; Srivastava et al., 2012a). Seasonal mean surface values for 2014/15–2018/19 show a clear gradient between CO over the Indian Ocean and over the landmass source regions (Fig. 112). Highest CO surface values over the Indian Ocean occur during the winter monsoon with multiannual mean mixing ratios of around 150 ppb over the open ocean and 350–400 ppb over Bay of Bengal coastal waters (Fig. 112b). CO mixing ratios over marine regions of more than 150 ppb are considered as polluted continental air (Nowak et al., 2004), and during the winter monsoon most parts of the NH Indian ocean fall into this category. South of 5°S, mixing ratios drop below 100 ppb during winter and can be considered as part of the pristine oceanic regime.




Figure 112. Surface carbon monoxide (CO) mixing ratios from MOPITT (coloured shading) and surface wind from ERA-Interim (black arrows) for (a) summer monsoon 2014 – 2018 (June – August), (b) winter monsoon 2014/2015 – 2018/2019 (December – February), (c) spring transition 2014 – 2018 (April – May) and (d) autumn transition 2014 – 2018 (October) periods.

The tropospheric distribution of NO_x is relatively similar to that of CO, however with a larger variability on small scales and steeper gradients between the continental source regions and the Indian Ocean. Figure 123 shows the tropospheric NO_2 column from TROPOMI, which, due to the fast photochemical cycling between NO and NO_2 , can be taken as a robust measure for concentrations of nitrogen oxides. Seasonal mean values for 2018/19 show a clear gradient of NO_2 , with maxima over the landmass source regions and minimum values over the equatorial western Indian Ocean (Fig. 123). Coastal gradients are particularly pronounced around the Indian coastline during the autumn transition period and winter monsoon. One clear difference in comparison to the CO distribution is the appearance of enhanced NO_2 columns along the shipping lanes, in particular along the major route at around 6°N over the Bay of Bengal from the southern tip of India to Malaysia.

The INDOEX campaign showed that the winter monsoon season is characterized by strongly enhanced CO abundances over the northern Indian Ocean, with values comparable to air downwind of Europe and North America (Lawrence and Lelieveld, 2010). Based on simultaneous measurements of methyl cyanide (CH_3CN) and model simulations, biomass burning in South and Southeast Asia and subsequent transport of the polluted air towards the Indian Ocean was identified as a major source of CO (Lelieveld et al., 2001). Over the open Indian Ocean, southward transport along with chemical processing, dilution, and surface deposition causes a strong north-south pollution gradient. In the boundary layer, the contrast

between polluted NH and pristine Southern Hemisphere (SH) air results in a sharp gradient across the ITCZ (Lawrence et al., 2003), while the upper troposphere was identified as a region for interhemispheric exchange of these air masses (Williams et al., 2002).




Figure 123. Tropospheric nitrogen dioxide (NO₂) column from TROPOMI for (a) summer monsoon 2019 (June – August), (b) winter monsoon 2018/2019 (December – February), (c) spring transition 2019 (April – May) and (d) autumn transition 2019 (October) periods.

Recent studies of atmospheric composition during the winter monsoon period have focused on the Bay of Bengal where the ship-based measurements were conducted during the second part of ICARB for December 2008–January 2009. Mixing ratios of NO₂ and CO show a distinct spatial pattern with elevated values in the head (northern part) and the south-east Bay of Bengal due to transport from adjoining landmasses (Asatar and Nair, 2010; David et al., 2011). The in-situ measurements reveal highest CO surface mixing ratios of 379 ± 58 ppb over the south-east Bay of Bengal, which exceed the MOPITT multiannual mean values (Fig. 112b). Similar to CO, the NO₂ surface distribution exhibits prominent highs over the head and the south-east Bay of Bengal, with lower values in the south (David et al., 2011). Analysis of back trajectories and airflow patterns revealed that the high pollution over the head of the Bay of Bengal (15°N–21°N) was advected from the Indo-Gangetic Plain, which is one of the major CO emission centres (Fig. 23). The high pollution over the south-east Bay of Bengal, on the other hand, was attributed to the effects of continental outflow from Southeast Asia, a transport pattern discernible from the wind patterns in Fig. 112. Airborne measurements conducted during the same campaign confirm the higher pollution over the northern and south-eastern part of the Bay of Bengal (Srivastava et al., 2012b). Analyses of the acetylene/CO ratio suggested that air mass

1 samples taken over the northern Bay of Bengal were less chemically aged compared to samples
2 taken during the other flights.

3 Enhanced wintertime CO values of up to 365 ppbv were also found in the Arabian Sea near the
4 southwest coast of India, while lower levels of around 50 ppb prevailed over the equatorial
5 Indian Ocean during ICARB-2018 (Girach et al., 2020a). Consistent with previous campaign
6 results, the authors linked the enhanced surface values to the impact of South Asian outflow.
7 The study also highlighted the impact of convective outflow from Southeast Asia in the upper
8 troposphere on the pollution levels. Enhanced CO in the upper troposphere (300–200 hPa) over
9 the equatorial IO was suggested to result from anomalous westerlies associated with a disturbed
10 Walker cell and the influence of African forest fires.

11 Seasonal variations of NO_x at a surface station in Trivandrum, situated on the south-west coast
12 of India, also show highest values during the winter monsoon season due to land-based
13 emissions and transport patterns (David and Nair, 2011). While the strongest NO_x emissions
14 are located in East Asia, enhanced emissions also exist in southern India (Fig. 34) contributing
15 to the wintertime NO_x maxima over the Indian coastline (see also Fig. 123). Ship based
16 observations of NO₂ show a strong decreasing gradient in concentrations from the sub-continent
17 towards the open ocean. Observations made on the IIOE-2 expedition observed peak values of
18 854 ± 223 ppt close to Goa, but the values were below the detection limit of 120 ppt away from
19 the coast. A similar gradient is evident from the satellite retrieved tropospheric NO₂ with the
20 emissions of NO_x over the sub-continent leading to higher values close to the coast (Fig. 123;
21 Mahajan et al., 2019a).

22 The impact of anthropogenic emissions during the winter monsoon is evident over large parts
23 of the Indian ocean, as demonstrated by surface measurements from the island of Mahé in the
24 western Indian Ocean just south of the equator. Seasonal variations of CO values peak in
25 January–February as a result of the long-range transport of anthropogenic emissions from India
26 during this time of year (Wai et al., 2014). Reunion Island around 19°S, on the other hand, is
27 much less impacted by the winter monsoon due to the ITCZ functioning as a transport barrier.
28 Seasonal variations of CO measurements at two stations on Reunion Island peak in September–
29 November, primarily driven by the emissions from biomass burning in Africa and South
30 America (Zhou et al., 2018). The seasonal variations of the CO distribution given by the
31 MOPITT satellite observations (Fig. 112) are consistent with the findings of the in-situ
32 measurement studies discussed above.

33 During the summer monsoon, CO mixing ratios over the Arabian Sea, Bay of Bengal and the
34 Indian Ocean drop mostly below 80 ppb, except for over coastal waters in the Bay of Bengal
35 with values of up to 150 ppb (Fig 12). As the ITCZ moves north over the Indian subcontinent,
36 the large-scale air flow is directed from the Indian Ocean towards the land, preventing the
37 spread of pollution in the maritime environment. In addition, convective uplifting and mixing
38 also contribute to lower CO abundances over the Bay of Bengal (Girach and Nair, 2014a). The
39 coastal regions of Bangladesh and eastern India are exceptions, as they are impacted by strong
40 westerlies transporting polluted air from southern India and Sri Lanka resulting in elevated CO
41 mixing ratios larger than 150 ppb (Fig. 112a). In-situ measurements in 2018 conducted as part
42 of IIOE-2 confirmed elevated CO levels up to 164 ppb downwind of Chennai, a coastal
43 megacity in East India (Girach et al., 2020b). In addition, the impact of shipping emissions was
44 detected near 6°N with higher values of CO (up to 120 ppb) in the otherwise pristine airmasses
45 of the southern Bay of Bengal. The elevated pollution along the eastern India coastline is in
46 stark contrast to the western India coastline, with very low CO mixing ratios of well below 100
47 ppb. The overall prevalence of clean oceanic background air during the summer monsoon
48 season was confirmed by the low CO values detected over the equatorial Indian Ocean during
49 the OMO campaign in July/August 2015 (Tomsche et al., 2019). Vertical profiles of CO were

1 derived during flights over Gan (Maldives), which is situated close to the equator and, thus,
2 influenced by air from the southern Indian Ocean. The CO profiles over Gan with values of
3 ~60 ppb were found to be lower than the NH background and the Asian monsoon CO profiles
4 for all altitudes below 8 km (Fig. 2 in Tomsche et al., 2019). During the AQABA ship campaign
5 in summer 2017, low pollution levels over the Arabian Sea were confirmed by measurements
6 of relatively low NO_x mixing ratios (0.19 ppb, Tadic et al., 2019).

7 The transition periods during boreal spring and autumn are characterized by a large spatial and
8 temporal variability of the pollution distribution, which shifts between the clean summer regime
9 and the polluted winter regime. Mean CO mixing ratios in the Bay of Bengal and Arabian Sea
10 range between 100 and 200 ppb, while the equatorial and southern Indian Ocean are dominated
11 by clean oceanic air with CO values below 100 ppb (Fig 12c and d). ICARB measurements
12 during March–April 2006 show higher CO values in the Bay of Bengal compared to the Arabian
13 Sea and higher CO values towards the coast, as expected with continental pollution being the
14 main source of the observed values (Aneesh et al., 2008). Transport regimes start to shift in
15 March when the outflow from India into the Arabian Sea and from Southeast Asia into the Bay
16 of Bengal begins to weaken. This shift leads to slowly decreasing pollution, as observed during
17 ICARB where CO values over the northern Bay of Bengal were found to be around 234 ppb
18 during March (Srivastava et al., 2012a), thus being lower than the above discussed winter
19 monsoon values. In April, offshore flow from the Indian subcontinent or Southeast Asia has
20 weakened further leading to very little pollution transport towards the ocean. The southern Bay
21 of Bengal is dominated by south-westerly winds (see also Fig 12c) carrying cleaner marine air,
22 as evident from CO mixing ratios of 88 ppb obtained during ICARB April measurements
23 (Srivastava et al., 2012a).

24 The boreal autumn transition period, also referred to as post-summer monsoon, marks the onset
25 of the polluted winter regime after the withdrawal of the monsoon winds. During this period, a
26 sudden increase of pollutant levels can be expected, in particular in coastal regions and over the
27 Bay of Bengal. During the SK-277 ship campaign, a large spatial heterogeneity of pollution
28 was observed over the Bay of Bengal (Mallik et al., 2013) reflecting the direct impact of air
29 masses being advected from different source regions in South Asia. Highest CO levels were
30 found in air masses originating in Southeast Asia with signatures of biomass and biofuel
31 burning. Continental pollution sources for NO_x were further enhanced by regional sources
32 possibly from ship emissions over the Bay of Bengal, which contains the international shipping
33 corridor connecting the southern tip of India and the Strait of Malacca.

34 **Sulfur dioxide (SO₂), mercury and ammonia**

35 In comparison to other pollutants, the SO₂ characterisation above the Indian Ocean is sparse.
36 Data available from the recent ICARB campaign in 2018, nonetheless, can be used to assess
37 the influence of anthropogenic SO₂ in the marine atmosphere over the Indian Ocean by using
38 non-sea salt sulfate aerosol (SO₄²⁺) as a proxy for SO₂. Aswini et al. (2020) show the presence
39 of SO₄²⁺ in an intense pollution plume over the Arabian Sea and the Indian Ocean during the
40 winter monsoon. Meteorological conditions during this season are favourable for SO₂ to SO₄²⁺
41 conversion. This is proposed to take place through photochemical oxidation of SO₂ by the OH
42 radical in the gas phase and through oxidation of SO₂ by H₂O₂ and O₃ in the aqueous phase
43 (Seinfeld and Pandis, 2006). About two-thirds of the total tropospheric SO₄²⁺ formation is
44 thought to occur through aqueous phase reactions (e.g., Warneck, 1999).

45 SO₂ has both anthropogenic and natural sources (i.e. oxidation of DMS), but comparison with
46 previous research cruise studies conducted nearly two decades ago shows a more than two-fold
47 increase in the concentration of nss-sulphate aerosols over the continental outflow region in the
48 Arabian Sea that appears to be due to anthropogenic SO₂ (Aswini et al., 2020). Despite
49 decreasing SO₂ emissions in East Asia since 2010, emissions are rapidly increasing in South

Asia, by about 10% per year with the Indo-Gangetic Plains being a major source region (Lelieveld et al., 2019, Sec. 4). In order to further explore the contribution of anthropogenic SO₂ to nss-sulphate aerosol formation, the ratio of methanesulphonic acid (MSA), due solely to DMS oxidation, to SO₄²⁺ was computed. The average MSA/nss-SO₄²⁺ ratio during the ICARB campaign was 3.1 × 10³. Over the more pristine sections of the cruise, the average ratio was 4.7 × 10³. In remote marine regions, ratios of 0.065 have been found (Savoie and Prospero, 1989). The lower MSA/nss-SO₄²⁺ ratio implies that most of the nss-SO₄²⁺ is from anthropogenic sources (Aswini et al., 2020).

Under the framework of the Global Mercury Observation System (GMOS) project, a mercury monitoring station was set up on Amsterdam Island, a remote and small island located in the southern Indian Ocean (Sprovieri et al., 2016). Observations of gaseous elemental mercury (GEM), reactive gaseous mercury (RGM) and particle-bound mercury (PBM) from this station over several years were reported by Angot et al. (2014). GEM concentrations were found to be remarkably steady, with an average hourly mean concentration of 1.03 ± 0.08 ng m⁻³, and show a small seasonal cycle. In comparison, the high altitude GMOS site in Kodaikanal located in southern India shows significantly higher GEM concentrations of 1.54 ± 0.20 ng m⁻³ in 2013, possibly related to different long-range transport patterns and closer proximity of anthropogenic sources (Sprovieri et al., 2016).

RGM and PBM concentrations at Amsterdam Island were also very low (0.34 pgm⁻³ and 0.67 pgm⁻³, respectively), but displayed a strong seasonal variability (ranging between the detection limit and 4.07 pgm⁻³ and 12.67 pgm⁻³, respectively). Analysis showed that, despite the remoteness of the island, long-range transport from southern Africa contributed significantly to the GEM and PBM budgets from July to September when biomass burning peaks in Africa (Angot et al., 2014). During these periods, the higher GEM concentrations observed at Amsterdam Island were comparable to those recorded at other tropical stations distributed around the globe. During periods of lower GEM concentrations, on the other hand, values of less than 1 ng m⁻³ were found to be characteristic for air masses from the southern Indian Ocean and Antarctic continent (Sprovieri et al., 2016).

Observations of ambient ammonia (NH₃) are rare in the Indian Ocean region. Recent measurements of NH₄ in aerosols over the Indian Ocean have been related to gaseous NH₃ emissions from land (either anthropogenic or crustal sources, Aryasree et al., 2015; Aswini et al., 2020). Earlier direct observations in 1980 by Ayers and Gras (1980) in the Southern Indian Ocean showed a range between 2.2 and 4.4 nmol m⁻³. Later observations in the same region of the ocean showed a lower range of 0.3 to 2.1 nmol m⁻³ (mean of 1.1 nmol m⁻³) (Norman and Leck, 2005). In the northwest Arabian Sea, observations of NH₃ were first carried out in 1999 reporting higher concentrations in the coastal environment 2.5 to 5.6 nmol m⁻³ (mean 3.8 nmol m⁻³) as compared to the remote open ocean environment 0.4 to 1.8 nmol m⁻³ (mean 1 nmol m⁻³), showing the importance of continental fluxes to the ambient marine NH₃ loading (Gibb and Mantouura, 1999a; 1999b). Similar to the Southern Indian Ocean, later observations by Norman and Leck (2005) again reported a lower range between 0.05 and 0.2 nmol m⁻³ (0.1 nmol m⁻³). The reason for this discrepancy between the two studies is not clear and was not discussed in the study by Norman and Leck (2005).

Closer to the Indian coast, observations of NH₃ have been made in the Bay of Bengal over five studies and show much higher concentrations as compared to the central or southern Indian Ocean. Khemani et al. (1987) reported high concentrations of NH₃ in the coastal region in the range between 117.6 and 211.8 nmol m⁻³ (mean 158.8 nmol m⁻³). Later, Carmichael et al. (2003) also reported high NH₃ concentrations at two coastal sites (Bhubneswar: mean 288.2 nmol m⁻³ and Berhampur: mean 329.4 nmol m⁻³). These observations show a west-east positive gradient

close to the western coast of the Bay of Bengal, which is most likely driven by the local transport of NH_3 . Further to the northwest, Biswas et al. (2005) made observations of ambient NH_3 close to the Sundarban mangrove forest which is one of the largest river deltas in the world. They saw highly elevated levels, ranging from 105.2 to 675.0 nmol m^{-3} (mean 265.2 nmol m^{-3}). The most recent reports in the literature were measurements done during the winter phase of ICARB in the Bay of Bengal, which reported an average concentration of ranging between 11.7 and 441.2 nmol m^{-3} (mean 281.2 nmol m^{-3}), with higher concentrations observed closer to the coast and the lower concentrations observed in the open ocean environment (Sharma et al., 2012). Unfortunately, no seasonal information is available due to the lack of continuous observations through the entire year in the marine environment.

Ozone

Ozone distribution over the Indian Ocean is largely determined by the abundance of precursor gases, transport patterns and chemical processing. During the winter monsoon, the southern Asian outflow brings ozone precursors such as CO , NO_x and VOCs from their source regions in South and East Asia (Section 4) into the marine environment. Within the outflow, substantial photochemical production of ozone occurs due to high pollution levels and strong tropical solar radiation, which is particularly effective under cloud-free conditions. As a result, relatively high ozone mixing ratios of 60–70 nmol/mol have been observed off the coast of India and over the Bay of Bengal up to a few hundred kilometres downwind (Lawrence and Lelieveld, 2010 and references therein). Similarly, high concentrations of up to 80 nmol/mol have also been reported off the western Indian coast in the Arabian Sea (Lal and Lawrence, 2001). Once produced, ozone can be transported to the remote marine environment, where local dominant sources of ozone precursor gases are absent. As a result, the marine boundary layer is considered as an ideal place to study the processes that drive ozone photochemistry (e.g., Monks et al., 1998). Tropospheric ozone is also influenced by downward transport from the stratosphere (e.g., Ganguly and Tzanis, 2011) and deposition to surfaces (Graedel and Crutzen, 1992). Photodissociation of ozone leads to increasing OH levels particularly within the highly humid marine boundary layer affecting the chemistry of the tropical marine environment and highlighting the importance of continental outflow over cleaner oceanic regions.

A continuous dataset of O_3 vertical profiles at Ahmedabad, a city in western India, shows a clear annual cycle dominated by the wind patterns (Lal et al., 2014). The lowest ozone (~20 ppb) was observed near the surface during September, which is at the end of the summer monsoon. Clean air from the ocean and removal of precursors due to monsoon rains drive this reduction. Model simulations showed that the lower tropospheric (<3 km) O_3 during the summer monsoon was transported from the Indian Ocean via the east coast of Africa and the Arabian Sea. Observations of mid-tropospheric ozone are highest (70–75 ppb) during April–June and lowest (40–50 ppb) during winter due to the impact of long-range transport from North Africa, North America, and the stratosphere. Balloon-borne O_3 measurements during 2011–2014 at the southwest coast of India confirmed the highly dynamic nature of tropospheric O_3 profiles (Ajayakumar et al., 2019). The variations in their vertical structure and short-term changes can be attributed to photochemistry as well as meteorological conditions in form of synoptic scale systems, long range transport and stratospheric intrusions. Unfortunately, such continuous measurements are not available at any site off the Indian subcontinent, but model simulations suggest that a similar annual cycle can also be expected in the northern Indian Ocean environment (Lal et al., 2014).

During the winter monsoon, measurements of ozone in the Bay of Bengal have been carried out as a part of the ICARB campaign (David et al., 2011). The marine boundary layer showed large variations in the mixing ratios of ozone and its precursor gases with similar spatial patterns (Fig. 134a) pointing to the same source regions, but different relative contributions. In the head

1 and south-east of the Bay of Bengal, mixing ratios of ozone (61 ± 7 ppb and 53 ± 6 ppb) and
2 precursors (discussed earlier) were found to be very high. Air mass back trajectories originated
3 from the Indo-Gangetic Plain and Southeast Asian countries, respectively, both characterized
4 by high tropospheric ozone and NO₂ values. In the south/south-western part of the Bay of
5 Bengal, the ozone mixing ratios were low, and the back trajectories originated from coastal
6 regions. Here, the longer transit times over the marine environment could have resulted in the
7 OH-driven destruction of ozone and precursors as well as in changes of surface level mixing
8 ratios due to updrafts and downdrafts (David et al., 2011). Over this oceanic region,
9 photochemical production involving NO₂ did not play a major role for the ozone production,
10 while the high water vapour acted as a sink.

11 Observations during the spring transition season of near-surface ozone in the Bay of Bengal
12 were carried out as part of the first ICARB phase during March and April 2006 (Fig. 134b).
13 Over the northern Bay of Bengal, higher ozone mixing ratios (~30 ppb, Nair et al., 2011) were
14 found to be well correlated with precursor gases indicating their co-located sources (Srivastava
15 et al., 2012a). Similar to the winter monsoon conditions, north-westerly winds transported large
16 amounts of anthropogenic pollutants from the Indo-Gangetic Plain to the northern part of the
17 Bay of Bengal causing the elevated pollution levels. The middle and southern Bay of Bengal
18 were more influenced by the open ocean environment and showed lower ozone mixing ratios
19 (~10–15 ppb, Nair et al., 2011). For both regions, the surface ozone mixing ratios were
20 anticorrelated with the boundary layer height. Ozone sonde measurements taken during the
21 same campaign revealed an elevated plume between 1 and 3 km over the northern Bay of
22 Bengal with very high ozone mixing ratios of 60–90 ppb (Srivastava et al., 2011; Lal et al.,
23 2013). The plume was also characterised by higher temperature and lower specific humidity
24 compared to the background marine air over other regions. This elevated layer sandwiched
25 between the marine boundary layer and the free troposphere can be attributed to the advection
26 of land air mass as a consequence of land-sea breeze circulation near the coast or long-range
27 transport. Ozone profiles in the first 3 km over the Arabian Sea are around 10 ppb lower than
28 over the Bay of Bengal (Lal et al., 2013). Above this height, ozone levels are about 20 ppb
29 higher over the Arabian Sea possibly due to intrusions of ozone-rich air from upper levels.

30 Observations over the Arabian sea during the spring transition period revealed a completely
31 different picture. Ozone and pollutants showed significantly lower mixing ratios when
32 compared to the northern Bay of Bengal, suggesting the larger impact of cleaner marine air
33 (Srivastava et al., 2012a). This argument is further supported by the poor inter-correlations of
34 ozone and the various precursors. Based on simulations with the global chemical transport
35 model MOZART, the authors hypothesize that loss of ozone due to halogen chemistry might
36 be an additional reason for the relatively low ozone abundances over the Arabian Sea.
37 Latitudinal gradients of the gases were slightly negative most likely due to some transport of
38 polluted air from southern India over the southern Arabian Sea and continuous dilution of
39 pollution with time over this region. Such negative pollution gradients over the Arabian sea
40 during the transition periods were confirmed by the CO distribution obtained from MOPITT
41 (Srivastava et al., 2012a; Fig. 112c and d).




Figure 134. Surface ozone mixing ratios in the Bay of Bengal observed during a) boreal spring (March – April 2006, ICARB) and b) winter monsoon (December – January 2009, ICARB). From David et al., 2011, copyright 2011, reproduced with permission.

During the summer monsoon period, unusually low levels of ozone (9 ppb) were reported for the Arabian Sea during the ARMEX campaign in 2002 (Ali et al., 2009). Based on MOZART model simulations, the authors attribute the low ozone abundances to chemical loss driven by reactive halogens from sea salt, which can account for up to 30% of the total ozone loss ($2.15 \text{ ppb day}^{-1}$). In addition, increased marine surface reactivity facilitated enhanced dry deposition of ozone and thus also contributed to the observed low ozone levels ($\sim 0.12 \text{ ppb day}^{-1}$). Summer monsoon ozone measurements close to the Arabian Peninsula are available from the AQABA ship campaign in 2017. Consistent with the low pollution levels detected in this region, ozone mixing ratios were relatively low (22.5 ppb) representing remote MBL conditions (Tadic et al., 2019). Net ozone production rates did not significantly deviate from zero and indicate weak net ozone production of 5 ppb day^{-1} . In contrast, measurements during the rest of the campaign in the Red Sea, Oman Gulf and Arabian Gulf revealed strongly enhanced tropospheric ozone with larger ozone production rates of up to 28 ppb day^{-1} . The sensitivity of ozone production to the prevailing conditions was assessed by defining ozone production regimes in terms of the OH reactivities of VOCs and NO_x . Based on this analysis, the relatively clean air of the Arabian Sea was found to be due to partially NO_x -limited ozone production in this region (Pfannerstill et al., 2019).

Summer monsoon measurements in the Bay of Bengal report enhanced ozone and pollution levels over the coastal regions, in particular, downwind of the tropical megacity Chennai. The Copernicus Atmosphere Monitoring Service (CAMS) model simulations suggest that in the Chennai plume, VOC-limited ozone production dominates in the morning and gradual shifts to a NO_x -limited regime in the afternoon (Girach et al., 2020b).

During the autumn transition season, simultaneous measurements of O_3 and precursors were made over the Bay of Bengal in October – November 2010. These measurements revealed large variability in O_3 (11–60 ppb) with maximum values found in the northern Bay of Bengal. Back trajectory analysis led to similar conclusion as the above mentioned ICARB study, revealing the influence of pollution plumes from the India-Bangladesh region and Southeast Asia in addition to long-range transport of pristine marine air from the Indian Ocean (Mallik et al.,

2013). Diurnal variations found in surface ozone were also attributed to the interplay of these two processes.

In addition to the studies focusing on the Bay of Bengal and the Arabian Sea, observations of O₃ were made along two ship tracks from the east and west of the Indian subcontinent and heading towards the open oceans during December-March as a part of the ISOE 8 and IIOE-2 in 2014 and 2015. The ISOE 8 campaign started from Chennai on the east coast and headed towards the Southern Ocean, while IIOE-2 started in Goa and went across the Arabian sea and the north equatorial Indian Ocean towards Mauritius. Both campaigns show a strong reduction in the ozone concentrations from ~50 ppb to ~5–10 ppb within a few degrees off the Indian coast (Mahajan et al. 2019a; 2019b), demonstrating that continental emissions are the main drivers behind the high ozone concentrations close to the Indian subcontinent.

Diurnal variations of ozone in the marine boundary layer are driven by effects of large-scale horizontal advection from source regions, entrainment of ozone rich air from higher altitudes, photochemical production involving precursor gases, and loss mechanism involving OH radicals. Over the last decade, significant progress has been made in understanding how the different processes determine diurnal ozone variations based on in-situ observations corroborated by chemistry simulations from box models, regional forecast models and chemical reanalysis.

In polluted air masses, daytime photochemical production of ozone from anthropogenic precursors such as NO_x has been observed (Nair et al., 2011; Mallik et al., 2013; Girach et al., 2020b). Such photochemical production only occurs if a sufficient amount of precursor gases is present and leads to peak ozone values around noon. Ozone daytime buildup has been identified in the downwind of the shipping lane over the Bay of Bengal (Girach et al., 2020b) and close to coastal cities (Nair et al., 2011).

In pristine air masses over the Indian Ocean with low pollution levels and high water vapour, photochemical destruction of surface ozone occurs during daytime (Nair et al., 2011; Mallik et al., 2013; Girach et al., 2020b). Photolysis of ozone produces O(1D), which reacts with H₂O and produces OH radicals with the latter causing the destruction of ozone (Pitts and Pitts, 2000). Increase in the absolute water vapour content was found to coincide with the decrease in ozone and vice versa (Nair et al., 2011). During the night-time, ozone rich air can be entrained from free troposphere into the marine boundary layer allowing ozone recovery and leading to next day daytime destruction. Box model simulations have highlighted the importance of such entrainment processes given their high sensitivity to various background levels of ozone (Mallik et al., 2013).

In the remote marine regions, ozone is known to show ‘virtually no diurnal variation’ (Liu et al., 1983) due to the scarcity of precursor gases and the lack of exchange between the ozone-rich free troposphere and the ozone-poor boundary layer. The absence of in situ photochemical production of ozone has been observed during various Indian Ocean campaigns (e.g., Sahu et al., 2006; Lal et al., 2007; Girach et al., 2017; Girach et al., 2018). Such absence of daytime ozone build-up suggests that the observed spatio-temporal variations in surface ozone can be attributed to in situ photochemical build-up in air masses moving from coastal polluted regions towards the Indian Ocean (Ojha et al., 2012; Girach et al., 2018). Ozone simulations with the Weather Research and Forecasting model with Chemistry (WRF-Chem) along air mass trajectories suggest mean en-route ozone production in the outflow towards the Bay of Bengal (Girach et al., 2017).

Table 4. O₃ mean values [ppb] and latitudinal gradients reported for various campaigns in the Bay of Bengal, Arabian Sea and Indian Ocean. Ozone maximum values are given in the marine boundary layer (MBL) and in the elevated layer (EL), as available.

Season	Year and campaign	Region	O ₃ [ppb]	O ₃ maxima [ppb]	Latitudinal O ₃ gradient [ppb/°]	Reference
Winter monsoon	1999 INDOEX	Indian Ocean Arabian Sea 15°S–20°N	21.5 ± 3.5 43.9 ± 7.9	80 – 100 (EL)	1.5 – 2	Lelieveld et al. (2001) Lal et al. (2006)
	2001 BOBEX I	Bay of Bengal 13°N–20°N	42 ± 12	64 (MBL)	1.5	Lal et al. (2006)
	2008/2009 ICARB	Bay of Bengal 8°N–21°N	48 ± 8	65 (MBL)	2.1	David et al. (2011)
	2003 BOBEX II	Bay of Bengal 4°N–19°N	34 ± 6	50 (MBL)	1.4	Lal et al. (2007)
	2014 ISOE-8	Bay of Bengal Indian Ocean 11.5°N–55°S	range: 5 – 53	53 (MBL)		Mahajan et al. (2019a)
	2015 IIOE-2	Arabian Sea Indian Ocean 11.5°N–20°S	range: 10 – 52	52 (MBL)		Mahajan et al. (2019b)
Spring transition	2006 ICARB	Bay of Bengal 6°N–21°N	18	30 (MBL)	1.3 ± 0.1	Nair et al. (2011)
		Bay of Bengal 6°N–21°N	28.3 ± 14.4	55 (MBL) 80 (EL)	5.4 ± 0.9 (>12°N)	Srivastava et al. (2012a)
		Arabian Sea 9°N–22°N	19.8 ± 4.1	26 (MBL) 60 (EL)	-0.4 ± 0.2	Srivastava et al. (2012a)
Summer monsoon	2002 ARMEX	Arabian Sea 10°N–17°N	9 ± 5	15 (MBL)		Ali et al. (2009)
	2017 AQABA	Arabian Sea 12°N–23°N	22.5	35 (MBL)		Tadic et al. (2019)

	2018 HIOE-2	Bay of Bengal 7°N–17°N	27.5 ± 6.3	45 (MBL)		Girach et al. (2020b)
Summer monsoon and autumn transition	2002 BOBPS	Bay of Bengal 7°N–20°N	27 ± 6	43 (MBL)	1.3	Sahu et al. (2006)
Autumn transition	SK-277 2010	Bay of Bengal 8°N–18°N	41 ± 9	60	3.95	Mallik et al. (2013)

Available ozone measurements from campaigns conducted in the Bay of Bengal, Arabian Sea, and Indian Ocean over the last decades have been summarized in Table 4. The measurements reveal some clear spatial and seasonal patterns with highest ozone mixing ratios during the winter monsoon, in particular in the Bay of Bengal and Arabian Sea. On average, ozone abundances decrease during the spring transition period and increase again after the summer monsoon. Nearly all campaigns detected highest ozone values close to the northern continental landmasses, reflecting direct impact of air masses originating from different pollution source regions in South Asia and the Indo-Gangetic Plain. However, the situation can also be reversed, with higher levels of ozone observed over the central Bay of Bengal, not near the coastal regions (Lal et al., 2006), due to different types of wind patterns and possible titration by fresh emissions of NO. Sharp latitudinal gradients ranging from 1.3 to 5.4 ppb O₃/° with increasing ozone towards northern landmasses were identified during all campaigns except for ICARB measurements in the Arabian Sea during the spring transition period. The strength of the gradients depends on the season, with sharper gradients during the winter and sometimes transition periods, and on the latitudinal extent, with sharper gradients closer to the coastlines. Air-sea breeze triggered transport of polluted air masses in the elevated layer can lead to substantially higher ozone values above the MBL, as demonstrated during the campaigns where ozonesonde measurements were available. Direct comparisons of ozone values in the Bay of Bengal and the Arabian Sea are in most cases not possible, except for the ICARB spring campaign during which ozone peaked in the Bay of Bengal.

5.2 Greenhouse gases

Greenhouse gases are largely emitted from anthropogenic activities over the continents, as are most of the atmospheric pollutants discussed above (Section 4). Therefore, greenhouse gases and pollution show a similar spatial distribution with higher values towards the coast and lower values over the open ocean. However, greenhouse gases have in general longer lifetimes resulting in overall smaller spatial gradients. In addition, they often have large sources from the terrestrial biosphere of the NH midlatitudes, which can impact their seasonal cycle over coastal and open ocean regions. In the following section, we discuss the distribution and variability of the greenhouse gases CH₄, N₂O, CO₂ and COS.

Methane (CH₄)

In-situ observations collected over the last two decades have significantly contributed to our current understanding of the large-scale CH₄ distribution. During the summer monsoon season, near-surface CH₄ mixing ratios of 1830 ± 140 ppb were detected over the northern Bay of Bengal during the 2009 CTCZ campaign (Girach et al., 2017). Back trajectory simulations linked the enhanced values over the Bay of Bengal to emissions from central and northern India.

1 Ship data collected during the winter monsoon shows decreasing CH₄ mixing ratios from the
2 northern (1910 ppb) to southern Bay of Bengal (1750 ppb) and slightly smaller abundances
3 over the Arabian Sea (1700 to 1800 ppb) (Srivastava et al., 2012a). Station data for 1993–2002
4 from the west coast of India revealed large seasonal variations with maximum values of up to
5 1980 ppb at the beginning of the winter monsoon season (Bhattacharya et al., 2009) related to
6 the interplay of seasonal variations in emissions and transport. CH₄ production from Asian rice
7 cultivation maximises during the autumn transition season when the plants are fully grown
8 (Section 4.2). At the same time, the onset of the winter regime starts to allow for air mass
9 transport from continental regions with significant anthropogenic influence. During the winter
10 monsoon, these transport patterns intensify bringing air masses with elevated CH₄ over the
11 Indian Ocean.

12 The three-dimensional distribution of CH₄ above the boundary layer over the Arabian Sea, Bay
13 of Bengal and Indian Ocean can be investigated based on GOSAT satellite measurements (Fig.
14 145; CH₄ at 800 hPa). CH₄ is higher towards the coast and lower over the open ocean due to
15 the proximity of the source rich land regions. In particular, the fossil fuel mining areas in the
16 Arab region are believed to contribute to enhanced CH₄ over the northern Arabian Sea (Kavitha
17 and Nair, 2019). Overall, latitudinal CH₄ variations between 20°N and 30°S of up to 6% are
18 much smaller than latitudinal variations of CO over the same region of up to 150%. Consistent
19 with in-situ surface measurements, CH₄ at the top of the boundary layer of the eastern Arabian
20 Sea and Bay of Bengal maximises during the autumn transition and winter monsoon seasons.
21 Peak values of over 2000 ppb appear over the northern Bay of Bengal and along the Indian west
22 coast. The overall spatial distribution does not change much with height and at all altitudes
23 between 850 – 200 hPa highest CH₄ values are found over the northern Arabian Sea and
24 northern Bay of Bengal (Kavitha and Nair, 2019). Similarly, the monthly variations of CH₄
25 mixing ratios over all oceanic regions remain more or less similar at all altitudes with a peak
26 during the autumn transition season.

27




Figure 145. Methane (CH₄) mixing ratios at 800 hPa from GOSAT for (a) summer monsoon 2009 – 2014 (June – August), (b) winter monsoon 2009/2010 – 2014/2015 (December – February), (c) spring transition 2009 – 2014 (April – May) and (d) autumn transition 2009 – 2004 (October) periods.

Over the equatorial Indian Ocean, CH₄ values at the top of boundary layer are relatively low during all seasons with values mostly below 1800 ppb, except for the winter season during which values can reach 1870 ppb (Fig. 145b). The OMO aircraft campaign in July/August 2015 confirmed low CH₄ values detected by the satellite for the summer monsoon. Observations over Gan (Maldives) show average CH₄ mixing ratios of 1778.3 ± 19.5 ppb during the summer period being significantly lower than in the NH background or the Asian monsoon profiles (Tomsche et al., 2019). CH₄ mixing ratios increase with height due to inter-hemispheric transport mixing NH air masses into the SH at higher altitudes. As the NH air masses have experienced convective uplift from the boundary layer, they advect higher CH₄ mixing ratios into the SH (Bergamaschi et al., 2013).

The SH Indian Ocean has been found to be much less impacted by the winter monsoon compared to the NH Indian Ocean due to the ITCZ functioning as a transport barrier. In situ measurements from Reunion Island around 19°S, suggest the opposite seasonal cycle with lowest CH₄ values in December–February and highest CH₄ values in August–September (Zhou et al., 2018). Here, the CH₄ variability is not dominated by nearby land sources, but rather seasonal variations of OH radicals drive the seasonal cycle of CH₄.

Nitrous oxide (N₂O)

Measurements of N₂O over the Indian Ocean are sparse. The distribution of N₂O derived during a ship campaign in the Arabian Sea, equatorial Indian Ocean, and southwest part of the Bay of

Bengal during the autumn transition period (October–November 2004) reveals clear latitudinal gradients (Mandal et al., 2006). Near the coast, N₂O values of 312 ppb were found, and they increased towards the open ocean, with a maximum of 432 ppb at the equator. In the equatorial SH, N₂O decreased sharply to 312 ppb and did not change substantially during the rest of the cruise along the equator and in the southwest part of the Bay of Bengal. Given the sparse data it is difficult to determine if the peak values around the equator are caused by local oceanic emissions or are related to long-range transport from other sources.

Carbon dioxide (CO₂)

CO₂ satellite-based observations over the Indian Ocean reveal strong seasonal and latitudinal gradients (Nalini et al., 2018; Nayak et al., 2011). During most seasons, larger CO₂ values over the south and south-east Asian land masses are accompanied by pronounced latitudinal gradients over the coastal regions and lower CO₂ values over the ocean. This distribution has been confirmed by ship-based observations during the summer monsoon (Kumar et al., 2014) and autumn transition season (Mandal et al., 2006). The measurements showed elevated CO₂ with large variability near the coastal region and relatively low CO₂ with correspondingly lower variability over the open ocean.

CO₂ over the Bay of Bengal and Arabian Sea is characterized by a pronounced seasonal cycle, with highest values in boreal spring and lowest values in autumn (Nalini et al., 2018). Satellite observations show that during winter, a band of relatively low CO₂ mixing ratios (376–378 ppm) extends over the northern Indian Ocean between 0°–25°N in the west-east direction, where the smallest values exist over the western Arabian Sea (Nayak et al., 2011). During boreal spring, the CO₂ in this region increases, in particular over the Bay of Bengal. Seasonality over the southern Indian Ocean shows a different pattern. Similarly, to the northern part, a belt of low CO₂ is found during boreal winter between 10°S and 25°S. During boreal spring, however, this band of relatively low CO₂ values remains prominent (Nayak et al., 2011), highlighting the potential role of this region as an oceanic CO₂ sink (Section 4.2). In addition, Valsala et al. (2020) used 60 years of reanalysis data to identify that atmospheric mixing ratios over the southeastern tropical region are driven by the Indian Ocean Dipole.

Lagrangian transport simulations, carried out to analyse the seasonal cycle at two Indian coastal stations, suggest that the seasonality of the CO₂ distribution over the Indian Ocean is driven by atmospheric circulation changes combined with the seasonal cycle of CO₂ over land regions (Nalini et al., 2018). While dominant marine contributions during the summer monsoon result in slowly decaying values, the winter and spring monsoon season is characterized by sources from the eastern continental land masses, leading to maximum values over the coastal stations during spring. The equatorial Indian Ocean, on the other hand, shows only a weak seasonal cycle when compared to Indian coastal locations, the Bay of Bengal, and the Arabian Sea. Here, contributions from oceanic regions dominate during all seasons with smaller continental contributions during the winter period. Overall, the anthropogenic emissions over the continental land masses appear not to be a large source for CO₂ in the open Indian Ocean environment.

Carbonyl sulfide (COS)

There are only three published datasets of ship or land-based observations of COS mixing ratios over the Indian Ocean. The measurements were made over the course of a year at Amsterdam Island (Mihalopoulos et al., 1991), on a research cruise in the eastern and southern Indian Ocean in November (Inomata et al., 2006), and on the OASIS research cruise in the western tropical Indian Ocean in July (Lennartz et al., 2017, see section 3). All measured COS values are either at or below the global mean COS atmospheric mixing ratio of ~549 ppt and exhibit no clear seasonal or latitudinal pattern (Lennartz et al., 2020). Satellite measurements and inverse

modelling studies show elevated values of the atmospheric COS mixing ratio over the Indian Ocean, but the values are not as high as those over the Pacific Ocean and Maritime Continent at similar latitudes (Glatthor et al., 2015; Kuai et al., 2015). The elevated values have led to the hypothesis that the northern Indian Ocean is one of the missing source regions (Kuai et al., 2015). Given the lack of compelling evidence from surface ocean measurements, however, it does not appear that the Indian Ocean is the location of the missing source. Instead, anthropogenic sources around Southeast Asia could be responsible, as discussed in section 4.

5.3 VOCs and short-lived gases DMS, isoprene and halogens

VOCs

The VOC mixing ratios over the Bay of Bengal are higher during the winter monsoon than during the summer monsoon for most VOCs, but with clear north-south gradients all year round (Sahu et al., 2011). Alkenes show an opposite seasonality to the rest of the major VOCs, with higher mixing ratios during the summer monsoon and no clear latitudinal gradients. For the other major VOCs, latitudinal gradients during the summer monsoon are caused by the transport of pollutants from peninsular India to the northern Bay of Bengal and clean marine air over southern Bay of Bengal. Latitudinal gradients during the winter monsoon are ascribed to dilution and increased photochemical loss towards the southern Bay of Bengal. However, measurements along the coasts during both seasons showed no clear latitudinal trend and were influenced by photochemistry and stagnant air flow. Seasonal changes in the distribution of VOCs in the surface layer over the Bay of Bengal resemble changes in wind speed (Sahu et al., 2010). Diurnal variability of alkenes is evident in summer, but not winter, and there is evidence that cyclones and convective activities in the summer season impact their mixing ratios. High mixing ratios are also found over the Arabian Gulf (Bourtsoukidis et al., 2020). VOC measurements at the Maïdo observatory in the southern Indian Ocean over two years reveal VOCs in the surface layer from anthropogenic sources (38%), biomass burning (33%), and both primary (15%) and secondary (14%) biogenic sources (Verreyken et al., 2021).

Regarding HCHO, Gopikrishnan et al. (2021) used satellite measurements for the period 2005–2014 and found large amounts along the ship routes in the Indian Ocean. The values were double the natural background without the presence of ships. The trend in HCHO concentrations over the Indian Ocean shipping lanes is about 0.008×10^{15} molec./cm²/year, where the HCHO values and trends from the northern Indian Ocean routes are comparable to those in the Panama Canal, Mediterranean Sea and Strait of Malacca.

DMS and isoprene

DMS in the Arabian Sea is projected to act as a relatively large oceanic source of sulfur to the atmosphere during the Asian summer monsoon season (Lana et al., 2011). These high emissions have been confirmed by recent ship-based measurements during the AQABA campaign, identifying elevated DMS values in the atmosphere ranging from 100 to 500 ppt (Edtbauer et al., 2020). Given the low atmospheric background pollution and the elevated DMS mixing ratios, the latter was found among the ten most important OH sinks with a reactivity of 0.02 s⁻¹ (Pfannerstill et al., 2019). During the OASIS campaign, similar atmospheric mixing ratios of approximately 20–300 ppt were measured over the western tropical Indian Ocean. The atmospheric distribution patterns in time and space during OASIS generally matched measured fluxes (Zavarsky et al., 2018b). During this study, the directly measured DMS fluxes, as well as calculated isoprene fluxes and sea spray fluxes, were correlated with satellite-derived aerosol products over the region. Many of the correlations, which took into account the influence of regional transport as computed FLEXPART trajectories, were statistically significant. The aerosol product distribution more closely resembled the trace gas fluxes than the sea spray flux

distributions. Thus, sea spray appears to be more of a minor precursor than the trace gases. This is supported by Quinn et al. (2017) who found a 30% contribution of sea spray to cloud condensation nuclei in the tropics. These results illustrate the importance of the regional coupling between marine-derived gaseous precursors and aerosol products in the remote MBL, which can give rise to local feedback processes (Zavarsky et al., 2018b).

The longest continuous observations of atmospheric DMS in the Indian Ocean were obtained at the Amsterdam Islands (37.8°S, 77.5°E) in the southern Indian Ocean (Scaire et al., 2000). Measurements from 1990 to 1999 revealed atmospheric DMS ranging from 5 to 1900 ppt with a clear seasonal cycle. Maximum DMS values in January were on average 20 times higher than minimum values in July-August. These strong seasonal variations are not caused by atmospheric transport patterns but are linked to a similar cycle in DMS concentration in seawater induced by enhanced phytoplanktonic activity during the boreal summer. Model runs using the Lana et al. (2011) climatology show a good match with these observations confirming the dominant impact of oceanic processes on atmospheric DMS concentrations over the open Indian Ocean (Mahajan et al., 2015b).

Atmospheric mixing ratios of isoprene were also measured during the OASIS campaign. The mean measured mixing ratio was 2.5 ± 1.5 ppt (Booge et al., 2016), which is in agreement with atmospheric measurements in other remote open ocean regions (Shaw et al., 2003). Booge et al. (2016) used a top-down approach to calculate isoprene emissions in order to compare with the bottom-up flux estimates using a box model, in which the only source of isoprene for the air was assumed to be the sea-to-air flux (emission), the atmospheric lifetime was assumed to be determined by reaction with OH (chemical loss, 1 h) and assuming air values to be zero at the start. Computed atmospheric mixing ratios were 45 times lower than measured. In order to calculate values consistent with measured mixing ratios, isoprene emissions must be more than one order of magnitude greater than those computed using the bottom-up estimate based on measured oceanic isoprene levels (section 4). This result agrees with isoprene emissions computed with a numerical model by Luo and Yu (2010). One possible explanation could be that production in the surface microlayer (SML) is not taken into account with the bottom-up approach. Ciuraru et al. (2015) showed that isoprene can be produced photochemically by surfactants in an organic monolayer directly at the air-sea interface. SML surfactant enrichment has been observed (Wurl et al., 2011), which could result in about two orders of magnitude larger isoprene fluxes than the highest fluxes calculated during the OASIS campaign. Further field measurements targeting isoprene production in the SML could be a step forward in reconciling the ocean source of isoprene to the atmosphere.

Halogens

The Indian Ocean is an important source region for halogenated VSLs such as CHBr_3 , CH_2Br_2 and CH_3I . During the OASIS ship campaign in summer 2014 in the west Indian Ocean, high VSL emissions with pronounced hotspots were detected (Fiehn et al., 2017). The prevailing atmospheric mixing ratios, however, were generally low. The atmospheric mixing ratios of CHBr_3 showed an overall mean of 1.20 ± 0.35 ppt. Elevated mixing ratios larger than 1.5 ppt were found in the equatorial region coinciding with lower wind speeds and maximum values larger than 2 ppt were detected close to islands, where coastal sources appear to influence the atmosphere. Atmospheric mixing ratios of CH_2Br_2 varied little around the mean value of 0.9 ppt and showed a similar pattern to the CHBr_3 mixing ratios. The CH_3I mixing ratios showed relatively large variations with a mean of 0.8 ppt (Fiehn et al., 2017).

Reactive halogen species, such as iodine oxide (IO) and bromine oxide (BrO), that result from VSLs and other sources (Section 4.3), were found to be below the detection limit or very low over the Indian Ocean. In December 2000, measurements in the southern Indian Ocean were carried out as preparative study for an intensive field campaign within the ELCID4 project with

ship track between Reunion Islands, Amsterdam Islands, Corzet Islands and Kerguelen Islands. IO and BrO were both below the detection limit of the instrument, and hence upper limits of 4 ppt were reported for both species in the Indian Ocean (Hönninger, 2002). A field campaign in the Maldives also reported upper limits for the BrO vertical columns (3×10^{14} molecules cm^{-2}) (Ladstätter-Weissenmayer et al., 2007). Observations on the ISOE and IIOE-2 cruises reconfirmed that BrO was below the detection limit, with a lower upper limit of 2 ppt on cruises starting from both the east and west of the Indian subcontinent (unpublished data). However, ship-based observations have confirmed the presence of iodine oxide in the marine atmosphere, although at low levels (<1 ppt). At these levels, the effect of halogen chemistry on ozone destruction in the Indian Ocean MBL is significantly smaller than the Atlantic MBL, where reactive halogen species can result in as much as 45% of the total ozone loss (Read et al., 2008). Additionally, the outflow of NO_x from the Indian sub-continent can titrate iodine chemistry through the formation of IONO_2 , which further reduces the impact of iodine on atmospheric chemistry (Mahajan et al., 2019a, 2019b). Another land-based campaign observed between 2.4–3.1 ppt of IO at the Maldives (Oetjen, 2009). These results suggest that, although the reactive species are low, they could contribute to local oxidative chemistry, considering that they are higher than levels observed in the Atlantic MBL where they contribute towards significant surface ozone loss (Mahajan et al., 2010).

6 Long-term changes

Long-term changes of atmospheric dynamics, oceanic processes and anthropogenic emissions impact the atmospheric composition over the Indian Ocean with the growth of traffic and industries being expected to lead to increasing air pollution. Available trend studies of pollution and greenhouse gases often make ~~often~~ use of long-term satellite measurements.

Despite the increasing emissions, CO abundances have been found to decrease in a number of studies. Srivastava et al. (2012a), using monthly mean CO at 900 hPa obtained from MOPITT, have observed lower offshore pollution during the transition periods than expected based on modelling studies with The Model for OZone And Related Tracers (MOZART) model. They suggest that temporal dilution of pollutants is the main reason for this mismatch. A study focusing on regions further south in the Indian Ocean indicates that CO was decreasing (1.8% per year over Madagascar and 1.7% per year over Reunion Island) over the years 2005 to 2009. The main drivers behind this decrease were identified as the La Niña between 2006 and 2008 and the reduction in biomass burning emissions in southern Africa (Tohir et al., 2015). Decreasing CO abundance in the lower troposphere was also found by Girach and Nair (2014b) using MOPITT CO data for 2000–2014. The authors attributed the negative trend of around 2% per year partly due to an increase in lower tropospheric water vapour and ozone and partly to a strengthening of convective activity, uplifting CO to higher altitudes. As a result of the latter, upper tropospheric CO was found to increase with a long-term trend of up to 3.2% per year. A recent study conducted as part of ICARB observed lower CO levels (200 ± 43 ppbv) than those during the INDOEX campaign (229 ± 40 ppbv) confirming decreasing levels of CO over the IO (Girach et al, 2020). As demonstrated in section 4, CO emissions in most regions surrounding the Indian Ocean have increased, which highlights the importance of the interplay of emission, dynamics, and chemistry for trace gas variability. In particular, CO emissions along the Mekong River, north of the Persian Gulf, in Afghanistan and East Africa have shown pronounced growth rates. Future studies of the CO trends over the Indian Ocean over longer time periods are needed in order to understand how these changing emissions, together with changing atmospheric dynamics and chemistry, impact the CO concentrations in the marine atmosphere.

The variation of the mean total columnar amount of tropospheric NO_2 has been studied in detail over the Indian subcontinent. Mahajan et al. (2015a) have reported an average increase of 2.20

1 $\pm 0.73 \text{ \% yr}^{-1}$ using four different satellites across India. For OMI, this rate of increase from
2 2004–2013 was $2.79 \pm 0.23 \text{ \% yr}^{-1}$. This compares well to the rate of $2.9 \pm 1.9 \text{ \% yr}^{-1}$ reported
3 by Ghude et al. (2013), even though their study focused on urban locations. Studies focusing
4 on the atmosphere over the Indian Ocean are relatively scarce, but Tournadre (2014) reported
5 a 50% increase along the Sri Lanka-Sumatra-China shipping lane. The OMI tropospheric
6 column data from 2003–2020 shows an increasing trend in the Arabian sea, close to the Indian
7 coast ($0.83 \pm 0.24 \text{ \% yr}^{-1}$; $t_B=3.41$), but along the eastern coast in the Bay of Bengal the trend
8 is not significant ($0.44 \pm 0.29 \text{ \% yr}^{-1}$; $t_B=1.53$). In the remote Indian ocean, OMI observations
9 show an increasing trend ranging between $0.76\text{--}1.87 \text{ \% yr}^{-1}$, although the open ocean trends
10 might not be accurate considering the low columnar densities close to or below the detection
11 limit of the satellite instruments (unpublished data).

12 Long-term changes of CH_4 have been estimated from AIRS satellite measurements over the
13 2003–2015 time period (Kavitha and Nair, 2019). Over the Arabian Sea and Bay of Bengal, a
14 consistent positive trend ranging from 2 ppb year^{-1} to 6 ppb year^{-1} was found at all pressure
15 levels comparable to trends over Indian land regions. Interestingly, the trend was larger at
16 higher altitudes ($<500 \text{ hPa}$) and maximized at the 300 hPa – 150 hPa level. The authors
17 hypothesized that the changes of growth rate with altitude are related to increased convective
18 activity uplifting CH_4 , leading to a smaller growth rate at the lower levels and higher growth at
19 the upper levels.

20 Based on OMI satellite data, a positive long-term trend in annual mean ozone over the Arabian
21 Sea and northern Bay of Bengal was detected for the ten-year period from 2000 to 2009 (David
22 and Nair, 2013). The southern Bay of Bengal is an exception to this and shows a negative long-
23 term change for annual mean ozone and the annual minimum with the latter representing
24 background conditions. Overall, quantifying long-term changes of ozone is an ongoing
25 challenge given its high variability and the scarcity of long-term measurement stations, so that
26 currently no updated, reliable trend estimates are available.

27 Observations of reactive halogen species over the Indian Ocean do not display any significant
28 long-term trends, with BrO values below the stated upper limits of 2 ppt across observations
29 made over two decades. In the case of IO, annual observations since 2014 during the ISOE ship
30 campaigns have also not identified a significant difference, although the time period is less than
31 one decade and hence changes would not be expected beyond instrumental accuracy and natural
32 variation. Laboratory studies indicate a strong link between ozone concentrations in the MBL
33 and emissions of iodine species from the ocean surface (Carpenter et al., 2013, section 6.3).
34 Modelling studies suggest that ocean emissions of iodine species have increased on a global
35 scale over the last few decades driven by an increase in ozone concentrations due to
36 anthropogenic emissions (Saiz-Lopez et al., 2014; 2015). This increase has been observed
37 indirectly through an increase in the concentrations of iodine species in ice cores in the Alps
38 and in the Arctic (Cuevas et al., 2018; Legrand et al., 2018). However, in the absence of long-
39 term ozone observations or iodine fluxes in the Indian Ocean region, it is difficult to quantify
40 the change in iodine emissions.

41 The examples listed above make clear that not only the increasing anthropogenic emissions,
42 but also atmospheric dynamics, chemistry, and ocean-atmosphere interactions, are important
43 for the long-term changes of the atmospheric composition over the Indian Ocean. In particular,
44 changes of transport patterns and convective uplifting can amplify or dampen the emission-
45 driven changes of greenhouse gases and pollution. Furthermore, long-term changes of the
46 frequency of modes of interannual variability can mask long-term changes of atmospheric
47 composition if the analysed period is relatively short (e.g., Tohir et al., 2015). In addition,
48 physical, chemical, and biological processes in the Indian Ocean, as well as dynamically driven

changes of the air-sea gas exchange, can be expected to impact the long-term trends of atmospheric composition.

7 Impact on the upper atmosphere and ocean

The changing atmospheric composition over the Indian Ocean can influence remote regions like the stratosphere via convective transport pathways (Section 7.1). It can also interact with the ocean by impacting biogeochemical cycles and marine ecosystems (Section 7.2), which can, in turn, feedback on the overlying atmosphere. Here, we present new insights into how atmospheric pollution over the Indian Ocean can impact other Earth system components.

7.1 Impact on the stratosphere

Marine trace gases emitted from the Indian Ocean can be transported into the stratosphere via convection directly above the Indian Ocean or via the summer monsoon convection and anticyclone (e.g., Fiehn et al., 2017; Hanumanthu et al., 2020). Recent studies have highlighted the role of the Indian Ocean as a source region for stratospheric VSLS entrainment and similar mechanisms could also apply to other marine traces gases.

~~Regional and global modelling studies have shown that stratospheric entrainment of oceanic VSLSs from the Indian Ocean occurs mainly via localized convection (Tegtmeier et al., 2020; Fiehn et al., 2018). Modelled CHBr₃ entrainment displays a global maximum over the Bay of Bengal, the Arabian Sea, and the southern tip of India (Tegtmeier et al., 2020). The main source regions for this global maximum include the Indian Ocean during boreal summer and the tropical west Pacific Ocean during boreal winter. Based on the OASIS campaign data in the west Indian Ocean and atmospheric modelling, Fiehn et al. (2017) have shown confirmed that, from the west Indian Ocean boundary layer, VSLSs can be transported from the boundary layer into the tropical tropopause layer and eventually into the stratosphere. All three studies highlight that the importance of the Indian Ocean sources for the stratospheric halogen budget depends on the regional and seasonal strength of emissions and the transit time in preferred transport regimes. On very short timescales, While the main entrainment occurs via convection above the Indian Ocean lifting oceanic trace gases towards the tropopause, Fiehn et al. (2017) suggest that, On longer timescales, the Asian summer monsoon may also play a role. The monsoon circulation can transports the oceanic VSLSs towards India and the Bay of Bengal. From where they VSLSs are lifted with the monsoon convection and reach stratospheric levels in the south-eastern part of the Asian monsoon anticyclone, which is generally the preferred transport regime during the boreal summer. StratoClim aircraft measurements of CH₂Br₂ in the UTLS above the Asian monsoon confirm these findings (Adcock et al., 2020). They show CH₂Br₂ abundances very similar to tropical background values, indicating that while some VSLS enter the stratosphere via the monsoon circulation, it is not a hotspot for entrainment of oceanic short-lived species. The stratospheric bromine injection from the tropical Indian Ocean and west Pacific depends critically on the seasonality and spatial distribution of the VSLS emissions (Fiehn et al., 2018). The main oceanic source regions for the stratosphere include the Arabian Sea and Bay of Bengal in boreal summer and the tropical west Pacific Ocean in boreal winter. For the OASIS case study in the western tropical Indian Ocean, the projected CH₂Br₂ entrainment was very high due to high surface emissions, while the entrainment of CHBr₃ and CH₃I was found to be relatively small compared to case studies in the tropical west Pacific Ocean (Fiehn et al., 2017; Tegtmeier et al., 2012). Overall, the Indian Ocean is a strong source of VSLSs for the stratosphere with the main entrainment occurring via localized convection, and modelled CHBr₃ shows a global maximum over India, the Bay of Bengal, and the Arabian Sea (Tegtmeier et al., 2020)~~

Besides contributing to stratospheric halocarbons, during the summer monsoon, the Indian Ocean and surrounding areas are potentially important source regions for stratospheric entrainment of other naturally produced gases such as N₂O and COS (Ma et al., 2018; Lennartz et al., 2017). The open Indian Ocean is in general a rather weak, but likely perennial and far-reaching (in terms of atmospheric transport) source of several trace gases to the atmosphere. However, it is unclear to what extent the atmospheric mixing ratios at a given location in the north Indian Ocean might correspond to air masses transported from the southwest Indian Ocean, where they have been enriched in oceanic trace gases (Ma et al., 2018). Such knowledge is necessary for quantifying the total contribution of the Indian Ocean emissions to the stratosphere and should be addressed in future studies. Potentially important feedback may occur if, for example, more N₂O is emitted to the atmosphere as a result of over fertilization (see section 6.3) and subsequently makes its way to the stratosphere. As N₂O contributes to stratospheric ozone depletion, resulting radiation changes at the ocean surface could feed back on biological and chemical processes.

7.2 Impact on the ocean

Changes of the atmospheric composition over the Indian Ocean, together with changing transport patterns and river inputs, affect oceanic production and biogeochemical trace gas cycling, which in turn feedback on the overlying atmosphere (e.g., Rixen et al, 2020 this issue). Satellite-derived time-series measurements indicate that the atmospheric pollution of some gases and annual aerosol load, including ship emissions, are increasing over the northern Indian Ocean and especially over the Bay of Bengal (Hsu et al., 2012; Tournadre, 2014). River inputs lead to low salinities in the surface mixed layer, resulting in strong stratification that dampens coastal upwelling and traps nutrients in the subsurface (Prasanna Kumar et al., 2002, Rixen et al., 2006). Other concerns include how potentially changing monsoon intensity will affect air mass trajectories (Goes et al., 2005), which influence deposition to the sea surface over the Indian Ocean and that Indian Ocean warming exacerbates ocean acidification (e.g., in the Arabaian Sea, Sreeush et al., 2019).

It has been shown that atmospheric deposition can have both favourable (nutrifying, e.g., Rahav et al., 2018) and unfavourable effects (toxic, e.g., Paytan et al., 2009, Jordi et al., 2012, Wang et al., 2015) on biological processes. Further, it has been shown that mixing natural and anthropogenic sources of gases and aerosols may alter the properties and subsequent functionality of the deposited materials (increased bioavailability, e.g., Herut et al., 2016, Krom et al., 2016, Jickells et al., 2017, Mahowald et al., 2018). Biogeochemical changes due to atmospheric deposition in the SML can be important for trace gas cycling and air-sea exchange, for example through reactions with deposited ozone (Zhou et al. 2014, Chiu et al. 2017, Mungall et al. 2017, see below) or changes to the heterotrophic community (Astrahan et al. 2016). Some modelling studies show that atmospheric deposition will lead to less CO₂ uptake and less ocean acidification (GESAMP 2012), while iron deposition doubles primary production (Guieu et al., 2019). The bioaccumulation of persistent organic pollutants and mercury in fish are of prevailing concern for fish stocks and human health (Brooks et al., 2019). In turn, marine-derived volatile forms of iodine, sulfur, and selenium are essential to recycle the elements onto the land and can be important for human health (Rayman, 2000; Fuge and Johnson, 2015).

To date, the anthropogenically-derived nitrogen input to the northern Indian Ocean, supplied by both atmospheric deposition and riverine fluxes (Jickells et al., 2017), has significantly increased in recent decades. This external nutrient source has the potential to affect the vulnerable biogeochemistry, by increasing overfertilization of the surface waters. Overfertilization can increase primary productivity, remineralization in the water column,

enhanced ocean deoxygenation and thus N_2O production via denitrification and nitrification cycles. As a consequence, the air-sea flux of N_2O increases (Suntharalingam et al., 2019), which can be expected to continue in the future.

Another example for the complex interaction between natural and anthropogenic processes in marine surface water and the atmosphere is the deposition of tropospheric ozone, and its reaction with marine iodide, which is linked to biological productivity and increases the input of inorganic iodine to the atmosphere (Carpenter et al., 2013). This major source of reactive iodine species in the marine atmosphere, in turn, forms a significant sink for atmospheric ozone through catalytic destruction. In polluted regions, however, the anthropogenic NO_x inactivates the reactive species to reservoir species (Mahajan, 2019b), thus volatile organoiodine compounds (Martino et al., 2009) or organobromine compounds (Kornmüller, 2007) are formed by ozone deposition. Especially in the polluted coastal regions of the northern landmasses, their formation adds to the release of organohalogens formed during industrial disinfection processes (Maas et al., 2020; 2021). Unexpected formation of ozone from reactive bromine and iodine species can occur in the marine boundary layer under, near coastlines and ship plumes, under VOC-limited conditions associated with high nitrogen oxide concentrations (Shechner and Tas, 2017).

8 Summary and current knowledge gaps

The atmospheric composition over the Indian Ocean is determined by a complex array of atmospheric transport patterns, chemistry, anthropogenic emissions, and atmosphere-ocean interactions. Emissions of pollutants and greenhouse gases in the regions surrounding the Indian Ocean generally correspond to population densities and economic activities with emission centres in the Indo-Gangetic Plain, northern China, and Java. The distribution of all major pollutants and greenhouse gases shows pronounced differences between the landmass source regions and the Indian Ocean, with strong gradients over the coastal areas. During the winter monsoon, the north-south pollution gradients continue over the open Indian Ocean driven by southward transport along with chemical processing, dilution, and surface deposition. In the boundary layer, the contrast between polluted NH and pristine SH air results in a sharp gradient across the ITCZ, where interhemispheric exchange of these air masses occurs in the upper troposphere. During the summer monsoon, clean air dominates the atmospheric composition over the northern Indian Ocean, leading to a different chemical regime with low atmospheric pollutant levels. Over the last decade, new knowledge of the atmospheric composition and related processes over the Indian Ocean has been derived from intensive ship and aircraft campaigns, coastal station measurements and satellite data. Here, we provide a synthesis of the scientific progress made after 2010 taking into account all observational data, updated emission inventories and modelling studies presented in Sections 3 to 7.

8.1 Pollution and O_3

Over the last decade, campaign and station measurements have revealed new findings of the distribution of pollution and ozone with some clear spatial and seasonal patterns. Over the Bay of Bengal strongly elevated pollution and ozone were detected in the head and the south-eastern part during winter and attributed to advection from the Indo-Gangetic Plain and continental outflow from Southeast Asia, respectively. Post-winter observations show similar maxima in the head, but significantly lower values in middle and southern Bay of Bengal due to open ocean influence. A completely different picture was found over the Arabian sea, where during spring transition period clean marine air results in significantly lower pollution when compared to the northern Bay of Bengal. Recent studies have also revealed that the impact of

anthropogenic pollution can extend far over the Indian Ocean. Remote locations such as the island of Mahé just south of the equator show CO maxima during winter that result from the long-range transport of anthropogenic emissions from India. Pollution levels further south such as over Reunion Island are driven by the emissions from biomass burning in Africa and South America. In addition to continental pollution source, ship emissions are known to impact the Indian Ocean air masses leading to enhanced NO_x abundances. Black carbon aerosols also show enhancements over Bay of Bengal shipping lanes, and thus similar signatures for CO could be expected. However, no studies have so far reported enhanced CO abundances in close proximity to Indian Ocean shipping lanes prompting the need for further investigations.

Given the strongly increasing pollutant emissions from the surrounding continents, similar trends might be expected in pollutants over the Indian Ocean. Comparisons of current observations and previous data obtained two decades earlier have demonstrated the growing influence of anthropogenic SO₂ on the marine atmosphere over the Indian Ocean. This pronounced increase has occurred despite the decreasing SO₂ emissions in East Asia and is most likely related to rapidly increasing emissions in South Asia. For tropospheric NO₂ column data an increasing trend was detected to be significant in the Arabian sea, close to the Indian coast. Lower tropospheric CO, on the other hand, was found to decrease, due to increasing water vapour and ozone as well as a strengthening of convective activity, uplifting CO to higher altitudes.

Scientific studies conducted over the last 10 years have added many details to our picture of the spatial distribution in ozone. Latitudinal gradients with increasing ozone towards northern landmasses were found to depend on the season, with sharper gradients during the winter and transition periods, and on the latitudinal extent, with sharper gradients closer to the coastlines. It has been noted that local wind patterns can lead to changes of the ozone seasonal cycle and pronounced deviations of the ozone distribution causing complex temporal and spatial variations. Diurnal variations of boundary layer ozone were found to follow different regimes with daytime photochemical production from anthropogenic precursors occurring only in polluted air masses. In pristine air masses, the absence of in situ photochemical production leads to either no diurnal ozone variation or photochemical destruction during daytime and entrainment of ozone rich from the free troposphere during nighttime. Despite the notable progress, a complete picture of the seasonal, latitudinal, and vertical ozone variations as well as reliable and updated trend estimates are ~~currently momentarily~~ not available.

8.2 Greenhouse gases

Emissions of CH₄, N₂O, and CO₂ in the regions surrounding the Indian Ocean have increased steadily over the last decade with East Asia often being the largest emitter. Important recent findings in this area have revealed high, and increasing, CH₄ emissions from east African wetlands. The ocean also acts as an important source/sink, especially for N₂O (source), CO₂ (sink), and OCS (source), and its role in regulating atmospheric budgets must be understood. Campaign data collected in 1999–2000 and 2004–2005 suggests that the CO₂ sink in the southern Indian Ocean is weakening, a conclusion further supported by a recent modelling study. Despite such alarming findings, we report here a drastic decrease in oceanic CO₂ observations over the last decades that must be rectified.

The distribution of greenhouse gases over the Indian Ocean shows many similarities when compared to the pollution fields, presenting higher values towards the coast due to the proximity of the source rich land regions. However, longer atmospheric lifetimes and active air-sea exchange also lead to some clear differences, such as the latitudinal variations of greenhouse gases being much smaller than the latitudinal variations of the shorter-lived pollutants. In

1 addition, greenhouse gas emissions are characterized by pronounced seasonal variations
2 because of their biogenic and agricultural sources. Over the northern Indian Ocean, CH₄
3 maximises during the autumn transition and winter monsoon following the seasonal cycle of
4 rice cultivation, while CO₂ is characterized by highest values in boreal spring due to biogenic
5 sources from the eastern continental land masses. Seasonality over the southern Indian Ocean
6 shows different patterns with seasonal variations of OH radicals leading to CH₄ maxima during
7 summer and air-sea exchange leading to relatively constant CO₂ levels. Long-term changes of
8 CH₄ show positive trends that maximized in the upper troposphere potentially driven by
9 increased convective activity.

10 Over the last decade, important progress has been made in understanding the interplay of
11 greenhouse gases and aerosols in driving physical trends in the region. It is clear now that the
12 Indian Ocean continues to warm due to greenhouse warming partially dampened by
13 anthropogenic aerosols. This warming has been shown to have serious consequences for the
14 monsoon and extreme weather in this region as well as for the global heat budget, as highlighted
15 by the recent hiatus period at the beginning of the 21st century. The warming is expected to
16 increase throughout the 21st century in response to continuing greenhouse gas emissions, with
17 the strongest warming in the Arabian Sea and western equatorial Indian Ocean consistently
18 projected in CMIP models.

20 **8.3 Short-lived gases DMS, isoprene and halogens**

21 For marine short-lived gases such as DMS, isoprene and halogenated VSLs, the Indian Ocean
22 is a hotspot emission region. Before 2010, short lived gases were some of the least sampled
23 compounds in the Indian Ocean. This review shows that over the last 10 years significant
24 improvements have been made in addressing the lack of data through multiple campaigns by
25 groups from all over the world.

26 For DMS, the number of datapoints available in the Indian Ocean has increased by 30–40%
27 since 2010. These new measurements can provide critical input for improving DMS emission
28 estimates, which are a major uncertainty in the currently available emission climatology. While
29 the total data are still too sparse to identify a long-term trend in DMS, proxy-based estimations
30 suggest an increase in global oceanic DMS concentrations, including the Indian Ocean. New
31 ship campaign data showed that oceanic sulfur fluxes in the form of DMS emissions are the
32 major precursor of aerosols in the summer monsoon remote MBL, with the possibility to
33 significantly contribute to cloud condensation nuclei in the tropical Indian Ocean.

34 In the case of isoprene, new campaigns conducted over the last 10 years have increased our
35 understanding of isoprene emissions, highlighting aspects such as the Arabian Sea isoprene
36 emissions peak in the oxygen minimum zone. An important new finding revealed that fluxes of
37 DMS and isoprene show a significant positive correlation with aerosol, suggesting a regional
38 influence of marine emissions.

39 New observations since 2014 have confirmed the ubiquitous presence of reactive iodine and
40 bromine species in the Indian Ocean, which interact with air pollutants from the Indian
41 subcontinent and need to be considered for regional air quality modelling. While the levels of
42 reactive halogen species were overall low, iodine chemistry was found to be responsible for up
43 to 25% ozone destruction in the northern Indian Ocean marine boundary layer. The improved
44 availability of reactive halogen dataset in the Indian Ocean demonstrates how important regular
45 in-situ observations are for our understanding of the regional atmospheric chemistry and
46 composition.

8.4 Knowledge gaps

Long-term changes ~~in the~~ atmospheric composition over the Indian Ocean are driven by increasing anthropogenic emissions, but also by atmospheric dynamics such as changes of transport patterns and convection. Furthermore, physical, chemical, and biological processes in the Indian Ocean can be expected to play a role for the long-term trends of atmospheric composition. Possible changes in stratification, mixing, microbial speciation, and primary productivity will influence trace gases in the atmosphere via modified oceanic emissions. Similarly, ocean acidification can impact biogenic trace gas production and exchange. In addition to changes of oceanic trace gas production, most of the observed and predicted atmospheric long-term changes can influence air-sea gas exchange. For example, more rainfall and intensified cyclone activities could lead to more turbulence at the sea surface and higher fluxes of gases. In most cases, the direction and magnitude of such changes are currently unclear and future studies are required to link oceanic changes to observed changes of atmospheric composition.

Our current understanding of the Indian Ocean is mostly based on sporadic ship campaigns and remote sensing data. Details of the large-scale features such as the seasonal cycles over the individual ocean basins, variations of the latitudinal gradients, the vertical distributions and long-term changes for several trace gases are not well known, and nor are all the processes well understood. Large uncertainties remain in the current emission inventories that are used in models and data for validation of model simulations is sparse. Dedicated long-term measurement stations, such as they exist in the Atlantic (e.g., Cape Verde Atmospheric Observatory), Pacific (e.g., Mauna Loa Observatory) or the Southern Ocean (e.g., Pointe Bénédicte station), are required in order to link the distribution of trace gases over the Indian Ocean to transport patterns and chemical regimes and to investigate feedback and forcing mechanisms. In addition to long term stations, it is imperative to maintain a repository of the past and future observations made in the Indian Ocean. Identifying long term trends, which are crucial to understanding the processes and impacts of climate change, are hindered by the lack of a consolidated database.

Interactions between the Indian Ocean and the atmosphere are bidirectional, and it has become clear that the changing atmospheric composition can impact many oceanic processes, leading to feedback mechanisms. Severe atmospheric pollution, known as the ‘South Asian Brown Cloud’ transports high levels of gaseous pollutants and aerosols (including dust) containing various nutrients and toxic substances to the Indian Ocean. This review only covers gas-phase composition over the Indian Ocean, however, the complex interactions between anthropogenic and marine trace gases and aerosols are also highly important for atmosphere-ocean feedback mechanisms and climate impacts. Interactions between aerosols and trace gases can form new particles, thus contributing to cloud formation, and wet deposition in the ~~clear-clean~~ marine atmosphere, as well as influence storm patterns and tropical cyclones. Such dynamical effects together with the additional supply of pollution and aerosols will impact biogeochemical and biological processes in the Indian Ocean. The connection between ocean biogeochemistry and atmospheric deposition involves several issues of importance for society, including global and regional pollution, the health of the ocean, fisheries, ocean fertilization, and carbon sequestration, all related to the UN Sustainable Development Goals. However, the impacts of atmospheric pollution and dust on the Indian Ocean’s biogeochemistry, trace gas cycling, and potential climate feedbacks, are severely understudied. Further research is needed to understand how sources, transport, reactivity, and the atmosphere-ocean feedback mechanisms interact. Such understanding is required in order to predict future changes, to assess if such changes can have harmful effects on the environment and to find pathways which strengthen atmosphere, ocean and community resilience in the Indian ocean region and globally.

Appendix A: Major abbreviations and terms

AQABA	Air Quality and climate change in the Arabian Basin
AIRS	Atmospheric Infrared Sounder
ARMEX	Arabian Sea Monsoon Experiment
BOB	Bay of Bengal
BOBEX	Bay of Bengal Experiment
BOBPS	Bay of Bengal Processes Studies
CMIP	Coupled Model Intercomparison Project
CTCZ	Continental Tropical Convergence Zone experiment
EACC	East African Coastal Current
EDGAR	Emissions Database for Global Atmospheric Research
EMC	East Madagasear Current
ENSO	El Niño–Southern Oscillation
GMOS	Global Mercury Observation System
GOME	Global Ozone Monitoring Experiment
GOSAT	Greenhouse Gases Observing Satellite
ICARB	Integrated Campaign for Aerosols, gases and Radiation Budget
IGP	Indo-Gangetic Plains
IIOE-2	2nd International Indian Ocean Expedition program
INDOEX	Indian Ocean Experiment
IO	Indian Ocean
ISOE	Indian Southern Ocean Expedition
ITF	Indonesian Throughflow
ITZC	Intertropical Convergence Zone
MBL	Marine Boundary Layer
MJO	Madden-Julian Oscillation
MSLP	Mean surface level pressure
MOPITT	Measurements of Pollution in the Troposphere
NH	Northern Hemisphere
NMHC	Non-Methane Hydrocarbon
NMVOC	Non-Methane Volatile Organic Compound
OASIS	Organic VSLs and their air sea exchange from the Indian Ocean to the Stratosphere
OMI	Ozone Monitoring Instrument
OMO	Oxidation Mechanism Observations
OMZ	Oxygen Minimum Zone
OVOC	Oxidized Volatile Organic Compound
PESO	Pilot Expedition to the Southern Ocean
SCIAMACHY	Scanning Imaging Absorption Spectrometer for Atmospheric Chartography
SEC	South Equatorial Current
SECC	South Equatorial Countercurrent
SH	Southern Hemisphere
SICC	South Indian Countercurrent
SML	Surface Microlayer
SOA	Secondary Organic Aerosols
SST	Sea Surface Temperature
TROPOMI	TROPOspheric Monitoring Instrument
VOC	Volatile Organic Compound
VSLs	Very Short-Lived Substances

Data availability.

EDGAR pollutant and greenhouse gas emission data are available from <https://edgar.jrc.ec.europa.eu>, SOCAT CO₂ data are available from <https://www.socat.info/index.php/data-access/>, MOPITT V8 Level 3 CO data are available from <https://www2.acom.ucar.edu/mopitt>, GOSAT CH₄ profiles are available at <http://www.gosat.nies.go.jp/en/>, and TROPOMI Level 2 NO₂ data are available from <https://scihub.copernicus.eu/>.

Author contributions.

ST coordinated the writing processes. The paper was written jointly by all co-authors with ST and CM designing the review and contributing to all sections. YJ contributed to sections 3 and 5; BQ contributed to sections 2, 7 and 8; AM contributed to sections 5 and 6.

Competing interests.

The authors declare that they have no conflict of interest.

Special issue statement.

This article is part of the special issue “Understanding the Indian Ocean system: past, present and future (BG/ACP/OS/SE inter-journal SI)”. It is not associated with a conference.

Acknowledgments

We thank Jonathan Williams, Liji M. David, Imran A. Girach, Vinu Valsala, Hermann Bange, Sinikka Lennartz, and Damian Arevalo for helpful discussions of the manuscript and Debora Griffin for assisting with the TROPOMI data. The Surface Ocean CO₂ Atlas (SOCAT) is an international effort, endorsed by the International Ocean Carbon Coordination Project (IOCCP), the Surface Ocean Lower Atmosphere Study (SOLAS) and the Integrated Marine Biosphere Research (IMBeR) program, to deliver a uniformly quality-controlled surface ocean CO₂ database. The many researchers and funding agencies responsible for the collection of data and quality control are thanked for their contributions to SOCAT. IITM is funded by the Ministry of Earth Sciences, Government of India.

References

[Adcock, K. E., Fraser, P. J., Hall, B. D., Langenfelds, R. L., Lee, G., Montzka, S. A., Oram, D. E., Röckmann, T., Stroh, F., Sturges, W. T., Vogel, B., and Laube, J. C.: Aircraft-Based Observations of Ozone-Depleting Substances in the Upper Troposphere and Lower Stratosphere in and Above the Asian Summer Monsoon, J. Geophys. Res.-Atmos., 126, e2020JD033137, <https://doi.org/10.1029/2020JD033137>, 2021.](#)

Ajayakumar, R. S., Nair, P. R., Girach, I. A., Sunilkumar, S. V., Muhsin, M., and Chandran, P. R. S.: Dynamical nature of tropospheric ozone over a tropical location in Peninsular India: Role of transport and water vapor. *Atmospheric Environment*, 218, 117018, 2019.

Ali, K., Beig, G., Chate, D. M., Momin, G. A., Sahu, S. K., and Safai, P. D.: Sink mechanism for significantly low level of ozone over the Arabian Sea during monsoon, *J. Geophys. Res.*, 114, D17306, doi:10.1029/2008JD011256, 2009.

Alory, G., and Meyers, G.: Warming of the upper equatorial Indian Ocean and changes in the heat budget (1960–99). *J. Climate*, 22 (1), 93–113. doi:10.1175/2008jcli2330.1, 2009.

- 1
2 Aneesh, V.R., Mohankumar, G. and Sampath, S.: Spatial distribution of atmospheric carbon
3 monoxide over Bay of Bengal and Arabian Sea: Measurements during pre-monsoon period of
4 2006. *J Earth Syst Sci* 117, 449–455, <https://doi.org/10.1007/s12040-008-0044-8>, 2008.
5
- 6 Angot, H., Barret, M., Magand, O., Ramonet, M., and Dommergue, A.: A 2-year record of
7 atmospheric mercury species at a background Southern Hemisphere station on Amsterdam
8 Island. *Atmos. Chem. Phys.*, 14(20), 11461–11473. [https://doi.org/10.5194/acp-14-11461-](https://doi.org/10.5194/acp-14-11461-2014)
9 2014, 2014.
- 10
11 Annamalai, H., Taguchi, B., McCreary J.P., Nagura, M., and Miyama T.: Systematic errors in
12 south Asian monsoon simulation: importance of Equatorial Indian Ocean processes. *J. Climate*,
13 30, 8159–817, 2017.
- 14
15 Arneth, A., Monson, R. K., Schurgers, G., Niinemets, Ü., and Palmer, P. I.: Why are estimates
16 of global terrestrial isoprene emissions so similar (and why is this not so for monoterpenes)?,
17 *Atmos. Chem. Phys.*, 8, 4605–4620, 10.5194/acp-8-4605-2008, 2008.
- 18
19 Arnold, S. R., Spracklen, D. V., Williams, J., Yassaa, N., Sciare, J., Bonsang, B., Gros, V.,
20 Peeken, I., Lewis, A. C., Alvain, S., and Moulin, C.: Evaluation of the global oceanic isoprene
21 source and its impacts on marine organic carbon aerosol, *Atmos. Chem. Phys.*, 9, 1253–1262,
22 10.5194/acp-9-1253-2009, 2009.
- 23
24 Aryasree, S., Nair, P.R., Girach, I.A. et al. Winter time chemical characteristics of aerosols over
25 the Bay of Bengal: continental influence. *Environ Sci Pollut Res* **22**, 14901–14918.
26 <https://doi.org/10.1007/s11356-015-4700-7>, 2015.
- 27
28 Asatar, G. I. and Nair., P. R.: Spatial distribution of near-surface CO over Bay of Bengal during
29 winter: role of transport, *J. Atmos. Sol.-Terr. Phy.*, 72, 1241–1250,
30 doi:10.1016/j.jastp.2010.07.025, 2010.
- 31
32 Asharaf, S., and Ahrens, B.: Indian summer monsoon rainfall processes in climate change
33 scenarios. *J. Climate*, 28(13), 5414–5429, 2015.
- 34
35 Astrahan P., Herut B., Paytan A., and Rahav E.: The Impact of Dry Atmospheric Deposition
36 on the Sea-Surface Microlayer in the SE Mediterranean Sea: An Experimental Approach,
37 *Frontiers in Marine Science*, 3, 222, doi=10.3389/fmars.2016.00222, 2016.
- 38
39 Aswini, A. R., Hegde, P., Aryasree, S., Girach, I. A., and Nair, P. R.: Continental outflow of
40 anthropogenic aerosols over Arabian Sea and Indian Ocean during wintertime: ICARB-2018
41 campaign, *Sci. Total Environ.*, 712, 135214, <https://doi.org/10.1016/j.scitotenv.2019.135214>,
42 2020.
- 43
44 Ayers, G. P. and Gras, J. L.: Ammonia gas concentration over the Southern Ocean, *Nature*, 284,
45 539–540, 1980.
- 46
47 Bakker, D. C. et al.: A multi-decade record of high quality fCO₂ data in version 3 of the Surface
48 Ocean CO₂ Atlas (SOCAT). *Earth Syst. Sci. Data*, 8: 383–413. doi:10.5194/essd8-383-2016,
49 2016.
- 50

- 1 Bange, H. W., Ramesh, R., Rapsomanikis, S., and Andreae, M.: Methane in surface waters of
2 the Arabian Sea, *Geophys. Res. Lett.*, 25, 3547–3550, 1998.
- 3
- 4 Bange, H. W.: New Directions: The importance of the oceanic nitrous oxide emissions, *Atmos.*
5 *Environ.*, 40, 198–199, 2006.
- 6
- 7 Barnes, E. A., Fiore, A. M. and Horowitz, L. W.: Detection of trends in surface ozone in the
8 presence of climate variability. *J. Geophys. Res.*, 121: 6112–6129. DOI:
9 10.1002/2015JD024397, 2016.
- 10
- 11 Bauer, S. E., Tsigaridis, K., and Miller, R.: Significant atmospheric aerosol pollution caused by
12 world food cultivation, *Geophys. Res. Lett.*, 43, 5394–5400,
13 <https://doi.org/10.1002/2016GL068354>, 2016.
- 14
- 15 Behrenfeld, M. J., O'Malley, R. T., Siegel, D. A., McClain, C. R., Sarmiento, J. L., Feldman,
16 G. C., Milligan, A. J., Falkowski, P. G., Letelier, R. M., and Boss, E. S.: Climate-driven trends
17 in contemporary ocean productivity. *Nature*, 444(7120):752–755, 2006.
- 18
- 19 Belikov, D.A., Saitoh, N., Patra, P.K., Chandra, N.: GOSAT CH₄ Vertical Profiles over the
20 Indian Subcontinent: Effect of a Priori and Averaging Kernels for Climate Applications.
21 *Remote Sens.* 13, 1677. <https://doi.org/10.3390/rs13091677>, 2021.
- 22
- 23 Bengtsson, L.: The global atmospheric water cycle, *Environ. Res. Lett.*, 5, 025002.
24 <http://dx.doi.org/10.1088/1748-9326/5/2/025002>, 2010.
- 25
- 26 Bergamaschi, P., S. Houweling, A. Segers, M. Krol, C. Frankenberg, R. A. Scheepmaker, E.
27 Dlugokencky, S. C. Wofsy, E. A. Kort, C. Sweeney, T. Schuck, C. Brenninkmeijer, H. Chen,
28 V. Beck, C. Gerbig: Atmospheric CH₄ in the first decade of the 21st century: Inverse modeling
29 analysis using SCIAMACHY satellite retrievals and NOAA surface measurements, *J. Geophys.*
30 *Res.*, 118, 7350– 7369, doi:10.1002/jgrd.50480, 2013.
- 31
- 32 Bhattacharya, S. K., Borole, D. V., Francy, R. J., Allison, C. E., Steele, L. P., Krummel, P.,
33 Langenfelds, R., Masarie, K. A., Tiwari, Y. K., and Patra, P. K.: Trace gases and CO₂ isotope
34 records from Cabo de Rama, India. *Current Science*, 97(9), 1336–1344,
35 <http://www.jstor.org/stable/24109728>, 2009.
- 36
- 37 Biswas, H., Chatterjee, A., Mukhopadhyaya, S. K., De, T. K., Sen, S., and Jana, T. K.: Estimation
38 of ammonia exchange at the land-ocean boundary condition of Sundarban mangrove northeast
39 coast of Bay of Bengal, India, *Atmos. Environ.*, 39, 4489– 4499, 2005.
- 40
- 41 Blando, J.D., Turpin, B.J.: Secondary organic aerosol formation in cloud and fog droplets: a
42 literature evaluation of plausibility, *Atmos. Environ.*, 34, 1623–1632, 2000.
- 43
- 44 Blunden, J. and D. S. Arndt, Eds.: State of the Climate in 2019. *Bull. Amer. Meteor. Soc.*,
45 101(8), S1–S429, doi: 10.1175/2020BAMSStateoftheClimate.1, 2020.
- 46
- 47 Boersma, K. F., Eskes, H. J., Richter, A., De Smedt, I., Lorente, A., Beirle, S., van Geffen, J.
48 H. G. M., Zara, M., Peters, E., Van Roozendaal, M., Wagner, T., Maasakkers, J. D., van der A,
49 R. J., Nightingale, J., De Rudder, A., Irie, H., Pinardi, G., Lambert, J.-C., and Compernolle, S.
50 C.: Improving algorithms and uncertainty estimates for satellite NO₂ retrievals: results from the

quality assurance for the essential climate variables (QA4ECV) project, *Atmos. Meas. Tech.*, 11, 6651–6678, <https://doi.org/10.5194/amt-11-6651-2018>, 2018.

Bollasina, M. A., Ming, Y. and Ramaswamy, V., Anthropogenic Aerosols and the Weakening of the South Asian Summer Monsoon, *Science*, 334, 6055, 502-505, 2011.

Booge, D., Marandino, C. A., Schlundt, C., Palmer, P. I., Schlundt, M., Atlas, E. L., Bracher, A., Saltzman, E. S., and Wallace, D. W. R.: Can simple models predict large-scale surface ocean isoprene concentrations?, *Atmos. Chem. Phys.*, 16, 11807-11821, 10.5194/acp-16-11807-2016, 2016.

Booge, D., Schlundt, C., Bracher, A., Endres, S., Zäncker, B., and Marandino, C. A.: Marine isoprene production and consumption in the mixed layer of the surface ocean – a field study over two oceanic regions, *Biogeosciences*, 15, 649-667, 10.5194/bg-15-649-2018, 2018.

Bourtsoukidis, E., Ernle, L., Crowley, J. N., Lelieveld, J., Paris, J. D., Pozzer, A., Walter, D., and Williams, J.: Non-methane hydrocarbon (C2–C8) sources and sinks around the Arabian Peninsula, *Atmos. Chem. Phys.*, 19, 7209-7232, 10.5194/acp-19-7209-2019, 2019.

Bourtsoukidis, E., Pozzer, A., Sattler, T. Matthaïos, V.N., Ernle, L. Edtbauer, A., Fischer, H., Könemann, T., Osipov, S., Paris, J.-D., Pfannerstill, E. Y., Stönnner, C., Tadic, I., Walter, D., Wang, N., Lelieveld, J., Williams, J.: The Red Sea Deep Water is a potent source of atmospheric ethane and propane. *Nat. Commun.*, -, 447, <https://doi.org/10.1038/s41467-020-14375-0>, 2020.

Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission Objectives and Measurement Modes, *J. Atmos. Sci.*, 56, 127-150, 10.1175/1520-0469(1999)056<0127:Smoamm>2.0.Co;2, 1999.

Brooks, S. D., T. D. Jickells, P. S. Liss, D. C. O. Thornton, and R. Zhang: Biogeochemical Coupling between Ocean and Atmosphere—A Tribute to the Lifetime Contribution of Robert A. Duce. *J. Atmos. Sci.*, 76, 3289–3298, <https://doi.org/10.1175/JAS-D-18-0305.1>, 2019.

Brühl, C., Lelieveld, J., Crutzen, P. J., and Tost, H.: The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate, *Atmos. Chem. Phys.*, 12, 1239-1253, 10.5194/acp-12-1239-2012, 2012.

Butler, A.H., J.S. Daniel, R.W. Portmann, A.R. Ravishankara, P.J. Young, D.W. Fahey, and K.H. Rosenlof: Diverse policy implications for future ozone and surface UV in a changing climate. *Environ. Res. Lett.*, 11: 064017, 2016.

Burrows, J. P., Weber, M., Buchwitz, M., Rozanov, V., Ladstätter-Weissenmayer, A., Richter, A., DeBeek, R., Hoogen, R., Bramstedt, K., Eichmann, K.-U., Eisinger, M., and Perner, D.: The Global Ozone Monitoring Experiment (GOME): Mission Concept and First Scientific Results, *J. Atmos. Sci.*, 56, 151-175, 10.1175/1520-0469(1999)056<0151:Tgomeg>2.0.Co;2, 1999.

Cai, W. et al.: Increased frequency of extreme Indian Ocean Dipole events due to greenhouse warming. *Nature* 510, 254–258, 2014.

- 1 Cai, W., A. Sullivan, and T. Cowan: Climate change contributes to more frequent consecutive
2 positive Indian Ocean Dipole events. *Geophys. Res. Lett.*, 36, L19783,
3 doi:10.1029/2009GL040163, 2009.
- 4
- 5 Campbell, J. E., Whelan, M. E., Seibt, U., Smith, S. J., Berry, J. A., and Hilton, T. W.:
6 Atmospheric carbonyl sulfide sources from anthropogenic activity: Implications for carbon
7 cycle constraints, *Geophys. Res. Lett.*, 42, 3004-3010, 10.1002/2015gl063445, 2015.
- 8
- 9 Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol
10 (SOA) formation from isoprene, *Atmos. Chem. Phys.*, 9, 4987-5005, 10.5194/acp-9-4987-
11 2009, 2009.
- 12
- 13 Carmichael, G. R., Ferm, M., Thongboonchoo, N., Woo, J. H., Chan, L. Y., Murano, K., Viet,
14 P. H., Mossberg, C., Bala, R., Boonjawat, J., Upatum, P., Mohan, M., Adhikary, S. P., Shrestha,
15 A. B., Pinaar, J. J., Brunke, E. B., Chen, T., Jie, T., Guoan, D., Peng, L. C., Dhiharto, S.,
16 Harjanto, H., Jose, A. M., Kimani, W., Kirouane, A., Lacaus, J.-P., Richard, S., Barturen, O.,
17 Cerda, J. C., Athayde, A., Tavares, T., Cotrina, J. S., and Bilici, E.: Measurements of sulfur
18 dioxide, ozone and ammonia concentration in Asia, Africa and South America using passive
19 samplers, *Atmos. Environ.*, 37, 1293–1308, 2003.
- 20
- 21 Carpenter, L. J., and Liss, P. S.: On temperate sources of bromoform and other reactive organic
22 bromine gases, *J. Geophys. Res.*, 105(D16), 20539– 20547, doi:10.1029/2000JD900242, 2000.
- 23
- 24 Carpenter, L. J., MacDonald, S. M., Shaw, M. D., Kumar, R., Saunders, R. W., Parthipan, R.,
25 Wilson, J. and Plane, J. M. C.: Atmospheric iodine levels influenced by sea surface emissions
26 of inorganic iodine, *Nat. Geosci.*, 6(2), 108–111, doi:10.1038/ngeo1687, 2013.
- 27
- 28 Chakraborty, K., V. Valsala, G. V. M. Gupta and V. V. S. S. Sarma, (2018): Dominant
29 biological control over upwelling on pCO₂ in sea east of Sri Lanka, *J. Geophysical Res.*,
30 doi.org/10.1029/2018JG004446.
- 31
- 32 Chang, J.-H.: The Indian Summer Monsoon, *Geogr. Rev.*, vol. 57, no. 3, 373–396, 1967.
- 33
- 34 Charlson, R., Lovelock, J., Andreae, M., and Warren, S. G.: Oceanic phytoplankton,
35 atmospheric sulphur, cloud albedo and climate. *Nature* 326, 655–661.
36 <https://doi.org/10.1038/326655a0>, 1987.
- 37
- 38 Chen, L., Xu, S., Gao, Z., Chen, H., Zhang, Y., Zhan, J., and Li, W.: Estimation of monthly air-
39 sea CO₂ flux in the southern Atlantic and Indian Ocean using in-situ and remotely sensed data,
40 *Remote Sens. Environ.*, 115, 1935-1941, <https://doi.org/10.1016/j.rse.2011.03.016>, 2011.
- 41
- 42 Chen, J., and Bordoni, S.: Early summer response of the East Asian summer monsoon to
43 atmospheric CO₂ forcing and subsequent sea surface warming, *J. Climate*. 295431–46, 2016.
- 44
- 45 Chin, M. and Davis, D. D.: Global sources and sinks of OCS and CS₂ and their distributions,
46 *Global Biogeochem. Cy.*, 7, 321–337, 1993.
- 47
- 48 Chiu, R., Tinel, L., Gonzalez, L., Ciuraru, R., Bernard, F., George, C., and Volkamer, R.: UV
49 photochemistry of carboxylic acids at the air-sea boundary: A relevant source of glyoxal and
50 other oxygenated VOC in the marine atmosphere, *Geophys. Res. Lett.*, 44, 1079– 1087,
51 doi:10.1002/2016GL071240, 2017.

Ciais, P., Dolman, A. J., Bombelli, A., Duren, R., Peregon, A., Rayner, P. J., Miller, C., Gobron, N., Kinderman, G., Marland, G., Gruber, N., Chevallier, F., Andres, R. J., Balsamo, G., Bopp, L., Bréon, F. M., Broquet, G., Dargaville, R., Battin, T. J., Borges, A., Bovensmann, H., Buchwitz, M., Butler, J., Canadell, J. G., Cook, R. B., DeFries, R., Engelen, R., Gurney, K. R., Heinze, C., Heimann, M., Held, A., Henry, M., Law, B., Luyssaert, S., Miller, J., Moriyama, T., Moulin, C., Myneni, R. B., Nussli, C., Obersteiner, M., Ojima, D., Pan, Y., Paris, J. D., Piao, S. L., Poulter, B., Plummer, S., Quegan, S., Raymond, P., Reichstein, M., Rivier, L., Sabine, C., Schimel, D., Tarasova, O., Valentini, R., Wang, R., van der Werf, G., Wickland, D., Williams, M., and Zehner, C.: Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system, *Biogeosciences*, 11, 3547-3602, 10.5194/bg-11-3547-2014, 2014.

Ciuraru, R., Fine, L., Pinxteren, M. V., D'Anna, B., Herrmann, H., and George, C.: Unravelling New Processes at Interfaces: Photochemical Isoprene Production at the Sea Surface, *Environ. Sci. Technol.*, 49, 13199–13205, doi:10.1021/acs.est.5b02388, 2015.

Codispoti, L. A.: Interesting times for marine N₂O. *Science*, 327(5971), 1339–1340. <https://doi.org/10.1126/science.1184945>, 2010.

Conte, L., Szopa, S., Séférian, R., and Bopp, L.: The oceanic cycle of carbon monoxide and its emissions to the atmosphere, *Biogeosciences*, 16, 881–902, <https://doi.org/10.5194/bg-16-881-2019>, 2019.

Crippa, M., Oreggioni, G., Guizzardi, D., Muntean, M., Schaaf, E., Lo Vullo, E., Solazzo, E., Monforti-Ferrario, F., Olivier, J.G.J., Vignati, E.: Fossil CO₂ and GHG emissions of all world countries - 2019 Report, EUR 29849 EN, Publications Office of the European Union, Luxembourg, ISBN 978-92-76-11100-9, doi:10.2760/687800, JRC117610, 2019.

Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global Atmospheric Research, *Sci. Data*, 7, 121, 10.1038/s41597-020-0462-2, 2020.

Cuevas, C. A., Maffezzoli, N., Corella, J. P., Spolaor, A., Vallelonga, P., Kjær, H. A., Simonsen, M., Winstrup, M., Vinther, B., Horvat, C., Fernandez, R. P., Kinnison, D., Lamarque, J.-F., Barbante, C. and Saiz-Lopez, A.: Rapid increase in atmospheric iodine levels in the North Atlantic since the mid-20th century, *Nat. Commun.*, 9(1), 1452, doi:10.1038/s41467-018-03756-1, 2018.

Da-Allada, C.Y., Gaillard, F. and Kolodziejczyk, N.: Mixed-layer salinity budget in the tropical Indian Ocean: seasonal cycle based only on observations. *Ocean Dynam.*, 65, 845–857, <https://doi.org/10.1007/s10236-015-0837-7>, 2015.

Daniel, J. S., and Solomon, S.: On the climate forcing of carbon monoxide, *J. Geophys. Res.*, 103(D11), 13249–13260, doi:10.1029/98JD00822, 1998.

David, L. M., Girach, I. A., and Nair, P. R.: Distribution of ozone and its precursors over Bay of Bengal during winter 2009: role of meteorology, *Ann. Geophys.*, 29, 1613–1627, <https://doi.org/10.5194/angeo-29-1613-2011>, 2011.

- David, L. M., and Nair, P. R.: Diurnal and seasonal variability of surface ozone and NO_x at a tropical coastal site: Association with mesoscale and synoptic meteorological conditions, *J. Geophys. Res.*, 116, D10303, doi:10.1029/2010JD015076, 2011.
- David, L. M. and Nair, P. R.: Tropospheric column O₃ and NO₂ over the Indian region observed by Ozone Monitoring Instrument (OMI): Seasonal changes and long-term trends, *Atmos. Environ.*, 65, 25–39, <https://doi.org/10.1016/j.atmosenv.2012.09.033>, 2013.
- Davidson, E. A.: The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nat. Geosci.*, 2:659–662. <https://doi.org/10.1038/NGEO608>, 2009.
- Deeter, M. N., Edwards, D. P., Francis, G. L., Gille, J. C., Mao, D., Martínez-Alonso, S., Worden, H. M., Ziskin, D., and Andreae, M. O.: Radiance-based retrieval bias mitigation for the MOPITT instrument: the version 8 product, *Atmos. Meas. Tech.*, 12, 4561–4580, [10.5194/amt-12-4561-2019](https://doi.org/10.5194/amt-12-4561-2019), 2019.
- De Smedt, I., Pinardi, G., Vigouroux, C., Compernelle, S., Bais, A., Benavent, N., Boersma, F., Chan, K.-L., Donner, S., Eichmann, K.-U., Hedelt, P., Hendrick, F., Irie, H., Kumar, V., Lambert, J.-C., Langerock, B., Lerot, C., Liu, C., Loyola, D., PETERS, A., Richter, A., Rivera Cárdenas, C., Romahn, F., Ryan, R. G., Sinha, V., Theys, N., Vlietinck, J., Wagner, T., Wang, T., Yu, H., and Van Roozendaal, M.: Comparative assessment of TROPOMI and OMI formaldehyde observations and validation against MAX-DOAS network column measurements, *Atmos. Chem. Phys.*, 21, 12561–12593, <https://doi.org/10.5194/acp-21-12561-2021>, 2021.
- DeVries, T., Le Quéré, C., Andrews, O., Berthet, S., Hauck, J., Ilyina, T., Landschützer, P., Lenton, A., Lima, I. D., Nowicki, M., Schwinger, J., and Séférian, R.: Decadal trends in the ocean carbon sink, *Proc. Natl. Acad. Sci. USA*, 116, 11646–11651, [10.1073/pnas.1900371116](https://doi.org/10.1073/pnas.1900371116), 2019.
- Dixit, A., L, K., Bharti, R., and Mahanta, C.: Net Sea–Air CO₂ Fluxes and Modeled Partial Pressure of CO₂ in Open Ocean of Bay of Bengal, *IEEE J. Sel. Top. Appl.*, 12, 2462–2469, [10.1109/JSTARS.2019.2902253](https://doi.org/10.1109/JSTARS.2019.2902253), 2019.
- Dong L, Zhou T and B. Wu: Indian Ocean warming during 1958–2004 simulated by a climate system model and its mechanism. *Clim. Dynam.*, 42, 203–217, doi.org/10.1007/s00382-013-1722-z (2014), 2014.
- Dong, L. and T. Zhou: The Indian Ocean sea surface temperature warming simulated by CMIP5 models during the twentieth century: Competing forcing roles of GHGs and anthropogenic aerosols. *J. Clim.* 27, 3348–3362, 2014.
- Du, Y. and S. P. Xie: Role of atmospheric adjustments in the tropical Indian Ocean warming during the 20th century in climate models. *Geophys. Res. Lett.*, 35, L08712, 2008.
- Du, Y., Zhang, Y., Feng, M., Wang, T., Zhang, N., and S.E. Wijffels: Decadal trends of the upper ocean salinity in the tropical Indo–Pacific since mid-1990s. *Sci. Rep.*, 5:16050. doi.org/10.1038/srep16050, 2015.

- 1 Du, Q., Zhang, C., Mu, Y., Cheng, Y., Zhang, Y., Liu, C., Song, M., Tian, D., Liu, P., Liu, J.,
2 Xue, C., and Ye, C.: An important missing source of atmospheric carbonyl sulfide: Domestic
3 coal combustion, *Geophys. Res. Lett.*, 43, 8720-8727, 10.1002/2016gl070075, 2016.
- 4
- 5 Dufлот, V., Tulet, P., Flores, O., Barthe, C., Colomb, A., Deguillaume, L., Väitilingom, M.,
6 Perring, A., Huffman, A., Hernandez, M. T., Sellegri, K., Robinson, E., O'Connor, D. J.,
7 Gomez, O. M., Burnet, F., Bourrianne, T., Strasberg, D., Rocco, M., Bertram, A. K., Chazette,
8 P., Totems, J., Fournel, J., Stamenoff, P., Metzger, J.-M., Chabasset, M., Rousseau, C.,
9 Bourrianne, E., Sancelme, M., Delort, A.-M., Wegener, R. E., Chou, C., and Elizondo, P.:
10 Preliminary results from the FARCE 2015 campaign: multidisciplinary study of the forest–gas–
11 aerosol–cloud system on the tropical island of La Réunion, *Atmos. Chem. Phys.*, 19, 10591–
12 10618, <https://doi.org/10.5194/acp-19-10591-2019>, 2019.
- 13
- 14 Edtbauer, A., Stönnner, C., Pfannerstill, E. Y., Berasategui, M., Walter, D., Crowley, J. N.,
15 Lelieveld, J., and Williams, J.: A new marine biogenic emission: methane sulfonamide
16 (MSAM), DMS and DMSO₂ measured in air over the Arabian Sea, *Atmos. Chem. Phys.*
17 *Discuss.*, <https://doi.org/10.5194/acp-2019-1021>, in review, 2020.
- 18
- 19 Endresen, Ø., Sørsgård, E., Sundet, J.K., Dalsøren, S.B., Isaksen, I.S.A., Berglen, T.F., and
20 Gravir, G.: Emission from international sea transportation and environmental impact, *Journal*
21 *of Geophysical Research*, 108, 4560. doi:10.1029/2002JD002898, 2003.
- 22
- 23 Engel, A., Rigby, M., Burkholder, J. B., Fernandez, R. P. Froidevaux, L., Hall, B. D., Hossaini,
24 R., Saito, T., Vollmer, M. K., Yao, B.: Update on ozone-depleting substances (ODSs) and other
25 gases of interest to the Montreal Protocol. In *Scientific assessment of ozone depletion: 2018*,
26 Doherty, S. J., Means, T., Stewart, B. C., McCarrick, A., Dailey-Fisher, D., Reiser, A. M., Eds.;
27 Global ozone research and monitoring project, Vol. 58; WMO (World Meteorological
28 Organization): Geneva; pp 1.1-1.87, 2019.
- 29
- 30 Ethé, C., Basdevant, C., Sadourny, R., Appu, K. S., Harenduprakash, L., Sarode, P. R., and
31 Viswanathan, G.: Air mass motion, temperature, and humidity over the Arabian Sea and
32 western Indian Ocean during the INDOEX intensive phase, as obtained from a set of
33 superpressure drifting balloons, *J. Geophys. Res.*, 107, 8023, doi:10.1029/2001JD001120,
34 2002.
- 35
- 36 Exton, D. A., Suggett, D. J., McGenity, T. J., and Steinke, M.: Chlorophyll-normalized isoprene
37 production in laboratory cultures of marine microalgae and implications for global models,
38 *Limnol. Oceanogr.*, 58, 1301-1311, 10.4319/lo.2013.58.4.1301, 2013.
- 39
- 40 Fadnavis, S., Semeniuk, K., Schultz, M. G., Mahajan, A. S., Pozzoli, L., Sonbawane, S., and
41 Kiefer, M.: Transport pathways of peroxyacetyl nitrate in the upper troposphere and lower
42 stratosphere from different monsoon systems during the summer monsoon season. *Atmos.*
43 *Chem. Phys.*, 15(14), 11477–11499. <https://doi.org/10.5194/acp-15-11477-2015>, 2015.
- 44
- 45 Fiehn, A., Quack, B., Hepach, H., Fuhlbrügge, S., Tegtmeier, S., Toohey, M., et al.: Delivery
46 of halogenated very short-lived substances from the west Indian Ocean to the stratosphere
47 during the Asian summer monsoon. *Atmos. Chem. Phys.*, 17(11), 6723–6741.
48 <https://doi.org/10.5194/acp-17-6723-2017>, 2017.
- 49
- 50 Fiehn, A., Quack, B., Stemmler, I., Ziska, F., and Krüger, K.: Importance of seasonally resolved
51 oceanic emissions for bromoform delivery from the tropical Indian Ocean and west Pacific to

- the stratosphere. *Atmos. Chem. Phys.*, 18(16), 11,973–11,990. <https://doi.org/10.5194/acp-18-11973-2018>, 2018.
- Fiore, A.M., West, J.J., Horowitz, L.W., Naik, V. and Schwarzkopf, M.D.: Characterizing the tropospheric ozone response to methane emission controls and the benefits to climate and air quality. *J. Geophys. Res.*, 113(D08): 307. DOI: 10.1029/2007JD009162, 2008.
- Franke, K., Richter, A., Bovensmann, H., Eyring, V., Jöckel, P., Hoor, P., and Burrows, J. P.: Ship emitted NO₂ in the Indian Ocean: comparison of model results with satellite data, *Atmos. Chem. Phys.*, 9, 7289–7301, <https://doi.org/10.5194/acp-9-7289-2009>, 2009.
- Fuge, R., and C. C. Johnson: Iodine and human health, the role of environmental geochemistry and diet, a review. *Appl. Geochem.*, 63, 282–302, 2015.
- Galí, M., Levasseur, M., Devred, E., Simó, R., and Babin, M.: Sea-surface dimethylsulfide (DMS) concentration from satellite data at global and regional scales. *Biogeosciences*, 15(11), 3497–3519. <https://doi.org/10.5194/bg-15-3497-2018>, 2018.
- Ganguly, N.D. and Tzanis, C.: Study of Stratosphere-troposphere exchange events of ozone in India and Greece using ozonesonde ascents. *Met. Apps*, 18: 467-474. doi:10.1002/met.241, 2011.
- Gantt, B., Meskhidze, N., and Kamykowski, D.: A new physically-based quantification of marine isoprene and primary organic aerosol emissions, *Atmos. Chem. Phys.*, 9, 4915-4927, 10.5194/acp-9-4915-2009, 2009.
- GESAMP: Rep. Stud. GESAMP 84: 69, 2012.
- Ghude, S. D., Beig, G., Kulkarni, P. S., Kanawade, V. P., Fadnavis, S., Remedios, J. J., and Kulkarni, S. H.: Regional CO pollution over the Indian-subcontinent and various transport pathways as observed by MOPITT, *Int. J. Remote Sens.*, 32, 6133–6148, 2011.
- Ghude, S.D., Kulkarni, S.H., Jena, C., Pfister, G.G., Beig, G., Fadnavis, S., van der A, R.J.: Application of satellite observations for identifying regions of dominant sources of nitrogen oxides over the Indian Subcontinent. *J. Geophys. Res.*, 118, 1075e1089. <http://dx.doi.org/10.1029/2012JD017811>, 2013.
- Gibb, S. W. and Mantoura, R. F. C.: Ocean-atmosphere exchange and atmospheric speciation of ammonia and methylamines in the region of the NW Arabian Sea, *Global Biogeochem. Cycles*, 13, 161–178, 1999a.
- Gibb, S. W., Mantouura, R. F. C., and Liss, P. S.: Ocean-atmosphere exchange and speciation of ammonia and methylamines in the region of the NW Arabian Sea, *Global Biogeochem. Cycles*, 13, 161–178, 1999b.
- Girach, I. A. and Nair, P. R.: On the vertical distribution of carbon monoxide over Bay of Bengal during winter: role of water vapour and vertical updrafts, *J. Atmos. Sol. Terr. Phys.*, 117, 31–47, doi:10.1016/j.jastp.2014.05.003, 2014a.
- Girach, I. A. and Nair, P. R.: Carbon monoxide over Indian region as observed by MOPITT, *Atmos. Environ.*, 99, 599–609, <https://doi.org/10.1016/j.atmosenv.2014.10.019>, 2014b.

- 1
- 2 Girach, I. A., Ojha, N., Nair, P. R., Pozzer, A., Tiwari, Y. K., Kumar, K. R., and Lelieveld, J.:
3 Variations in O₃, CO, and CH₄ over the Bay of Bengal during the summer monsoon season:
4 shipborne measurements and model simulations, *Atmos. Chem. Phys.*, 17, 257–275,
5 <https://doi.org/10.5194/acp-17-257-2017>, 2017.
- 6
- 7 Girach, I. A., Ojha, N., Nair, P. R., Tiwari, Y. K., and Kumar, K. R., Variations of trace gases
8 over the Bay of Bengal during the summer monsoon. *Journal of Earth System Science*, doi:
9 10.1007/s12040-017-0915-y, 2018.
- 10
- 11 Girach, I. A., Nair, P. R., Ojha, N., and Sahu, S. K.: Tropospheric carbon monoxide over the
12 northern Indian Ocean during winter: Influence of inter-continental transport, *Climate*
13 *Dynamics*, 54, 5049–5064. <https://doi.org/10.1007/s00382-020-05269-4>, 2020a.
- 14
- 15 Girach, L. A., N. Tripathi, P.R. Nair, L.K. Sahu, N. Ojha: O₃ and CO in the South Asian outflow
16 over the Bay of Bengal: impact of monsoonal dynamics and chemistry, *Atm. Environ*, 233, p.
17 117610, 10.1016/j.atmosenv.2020.117610, 2020b.
- 18
- 19 Glatthor, N., Höpfner, M., Baker, I. T., Berry, J., Campbell, J. E., Kawa, S. R., Krysztofiak, G.,
20 Leyser, A., Sinnhuber, B.-M., Stiller, G. P., Stinecipher, J., von Clarmann, T.: Tropical sources
21 and sinks of carbonyl sulfide observed from space, *Geophys. Res. Lett.*, 42, 10,082–10,090,
22 doi:10.1002/2015GL066293, 2015.
- 23
- 24 Goes, J.I., P.G. Thoppil, H. Gomes, and J.T. Fasullo: Warming of the Eurasian landmass is
25 making the Arabian Sea more productive, *Science*, 308 (5721), 545-547,
26 10.1126/science.1106610, 2005.
- 27
- 28 Gopika, S., Izumo, T., Vialard, J., Lengaigne, M., Suresh, I. and M. R. R. Kumar: Aliasing of
29 the Indian Ocean externally-forced warming spatial pattern by internal climate variability,
30 *Clim. Dynam.*, 54, 1-2, 1093-1111, 2020.
- 31
- 32 Gopikrishnan, G. S., and Kuttippurath, J.: A decade of satellite observations reveal significant
33 increase in atmospheric formaldehyde from shipping in Indian Ocean, *Atmos. Environ.*, 246,
34 118095, <https://doi.org/10.1016/j.atmosenv.2020.118095>, 2021
- 35
- 36 Graedel, T. E. and Crutzen, P. J.: *Atmospheric Change: an Earth System Perspective*, W. H.
37 Freeman, New York, 1992.
- 38
- 39 Gregg, W. W., and Rousseaux, C. S.: Global ocean primary production trends in the modern
40 ocean color satellite record (1998–2015). *Environ. Res. Lett.* 2019;14:1–9, 2019.
- 41
- 42 Gschwend, P. M., MacFarlane, J. K. and Newman, K. A.: Volatile halogenated organic
43 compounds released to seawater from temperate marine macroalgae, *Science* 227, 1033–1035,
44 1985.
- 45
- 46 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of
47 global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and
48 Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.
- 49
- 50 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and
51 Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1

- (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, 10.5194/gmd-5-1471-2012, 2012.
- Guerreiro, C. V., Baumann, K.-H., Brummer, G.-J. A., Fischer, G., Korte, L. F., Merkel, U., Sá, C., de Stigter, H., and Stuut, J.-B. W.: Coccolithophore fluxes in the open tropical North Atlantic: influence of thermocline depth, Amazon water, and Saharan dust, *Biogeosciences*, 14, 4577–4599, <https://doi.org/10.5194/bg-14-4577-2017>, 2017.
- Guieu, C., O. Aumont, A. Paytan, L. Bopp, C. S. Law, N. Mahowald, E. P. Achterberg, E. Maraño, B. Salihoglu, A. Crise, T. Wagener, B. Herut, K. Desboeufs, M. Kanakidou, N. Olgun, F. Peters, E. Pulido-Villena, A. Tovar-Sanchez and C. Völker: The significance of the episodic nature of atmospheric deposition to Low Nutrient Low Chlorophyll regions, *Global Biogeochem. Cy.*, 28, 1179–1198, doi:10.1002/2014GB004852, 2014.
- Guieu, C., A., Azhar, M., Aumont, O., Mahowald, N., Levy, M., Ethé, C., and Lachkar, Z.: Major impact of dust deposition on the productivity of the Arabian Sea. *Geophys. Res. Lett.*, 46, 6736–6744. <https://doi.org/10.1029/2019GL082770>, 2019.
- Han, W., and J. P. McCreary: Modelling salinity distributions in the Indian Ocean, *J. Geophys. Res.*, 106, 859 – 877, 2001.
- Han, W. et al.: Indian Ocean decadal variability: a review. *Bull. Am. Meteorol. Soc.* 95, 1679–1703, 2014.
- Han, Z., Su, T., Zhang, Q., Wen. Q., and G. Feng: Thermodynamic and dynamic effects of increased moisture sources over the Tropical Indian Ocean in recent decades, *Clim. Dynam.*, 53:7081–7096, 2019.
- Hanumanthu, S., Vogel, B., Müller, R., Brunamonti, S., Fadnavis, S., Li, D., Ölsner, P., Naja, M., Singh, B. B., Kumar, K. R., Sonbawne, S., Jauhiainen, H., Vömel, H., Luo, B., Jorge, T., Wienhold, F. G., Dirksen, R., and Peter, T.: Strong day-to-day variability of the Asian Tropopause Aerosol Layer (ATAL) in August 2016 at the Himalayan foothills, *Atmos. Chem. Phys.*, 20, 14273–14302, <https://doi.org/10.5194/acp-20-14273-2020>, 2020.
- Hastenrath, S. and Polzin, D., Dynamics of the surface wind field over the equatorial Indian Ocean. *Q.J.R. Meteorol. Soc.*, 130: 503–517. doi:10.1256/qj.03.79, 2004.
- Hastenrath, S., Zonal Circulations over the Equatorial Indian Ocean. *J. Climate*, 13, 2746–2756, [https://doi.org/10.1175/1520-0442\(2000\)013<2746:ZCOTEI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<2746:ZCOTEI>2.0.CO;2), 2000.
- Hay, T., Kreher, K. and Riedel, K.: Bromine explosions and Antarctic ozone. *Water and Atmosphere*, 15, pp. 12–13, 2007.
- Henze, D. K., and Seinfeld, J. H.: Global secondary organic aerosol from isoprene oxidation, *Geophys. Res. Lett.*, 33, 10.1029/2006gl025976, 2006.
- Herut B., Rahav E., Tsagaraki T. M., Giannakourou A., Tsiola A., Psarra S., Lagaria A., Papageorgiou N., Mihalopoulos N., Theodosi C. N., Violaki K., Stathopoulou E., Scoullou M., Krom M. D., Stockdale A., Shi Z., Berman-Frank I., Meador T. B., Tanaka T., and Paraskevi P.: The Potential Impact of Saharan Dust and Polluted Aerosols on Microbial Populations in

the East Mediterranean Sea, an Overview of a Mesocosm Experimental Approach, *Frontiers in Marine Science*, 3, 226, doi=10.3389/fmars.2016.00226, 2016.

Hönninger, G.: Halogen Oxide Studies in the Boundary Layer by Multi Axis Differential Optical Absorption Spectroscopy and Active Longpath-DOAS. University of Heidelberg, 2002.

Hood, R. R., Urban, E. R., McPhaden, M. J., Su, D., and Raes, E.: The 2nd International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Basin, *Limnol. Oceanogr. Bulletin*, 25, 117-124, 10.1002/lob.10149, 2016.

Hood R. R., Beckley, L. E. and Wiggert, J. D.: Biogeochemical and ecological impacts of boundary currents in the Indian Ocean. *Prog. Oceanogr.*, 156: 290–325, 2017.

Hopkins, F. E. and Archer, S. D.: Consistent increase in dimethyl sulfide (DMS) in response to high CO₂ in five shipboard bio assays from contrasting NW European waters, *Biogeosciences*, 11, 4925–4940, <https://doi.org/10.5194/bg-11-4925-2014>, 2014.

Hopkins, F. E., Turner, S. M., Nightingale, P. D., Steinke, M., Bakker, D., and Liss, P. S.: Ocean acidification and marine trace gas emissions, *P. Natl. Acad. Sci. USA*, 107, 760–765, 2010.

Hossaini, R., Mantle, H., Chipperfield, M. P., Montzka, S. A., Hamer, P., Ziska, F., Quack, B., Krüger, K., Tegtmeier, S., Atlas, E., Sala, S., Engel, A., Bönisch, H., Keber, T., Oram, D., Mills, G., Ordóñez, C., Saiz-Lopez, A., Warwick, N., Liang, Q., Feng, W., Moore, F., Miller, B. R., Maréchal, V., Richards, N. A. D., Dorf, M., and Pfeilsticker, K.: Evaluating global emission inventories of biogenic bromocarbons, *Atmos. Chem. Phys.*, 13, 11819-11838, 10.5194/acp-13-11819-2013, 2013.

Hsu, N. C., Gautam, R., Sayer, A. M., Bettenhausen, C., Li, C., Jeong, M. J., Tsay, S.-C., and Holben, B. N.: Global and regional trends of aerosol optical depth over land and ocean using SeaWiFS measurements from 1997 to 2010, *Atmos. Chem. Phys.*, 12, 8037–8053, <https://doi.org/10.5194/acp-12-8037-2012>, 2012.

Hu, Q. H., Xie, Z. Q., Wang, X. M., Kang, H., He, Q. F., and Zhang, P.: Secondary organic aerosols over oceans via oxidation of isoprene and monoterpenes from Arctic to Antarctic, *Sci Rep*, 3, 2280, 10.1038/srep02280, 2013.

Hu, S., and Sprintall, J.: Observed strengthening of interbasin exchange via the Indonesian seas due to rainfall intensification. *Geophys. Res. Lett.* 44, 1448–1456. doi: 10.1002/2016GL072494, 2017.

Inamdar, S., Tinel, L., Chance, R., Carpenter, L. J., Sabu, P., Chacko, R., Tripathy, S. C., Kerkar, A. U., Sinha, A. K., Bhaskar, P. V., Sarkar, A., Roy, R., Sherwen, T., Cuevas, C., Saiz-Lopez, A., Ram, K., and Mahajan, A. S.: Estimation of Reactive Inorganic Iodine Fluxes in the Indian and Southern Ocean Marine Boundary Layer, *Atmos. Chem. Phys. Discuss.*, 2020, 1-57, 10.5194/acp-2019-1052, 2020.

Inomata, Y., Hayashi, M., Osada, K., and Iwasaka, Y.: Spatial distributions of volatile sulfur compounds in surface seawater and overlying atmosphere in the northwestern Pacific Ocean, Eastern Indian Ocean, and Southern Ocean, *Global Biogeochem. Cy.*, 20, GB2022, <https://doi.org/10.1029/2005gb002518>, 2006.

- 1
- 2 IPCC. Climate Change 2013: The Physical Science Basis. In: Stocker, TF, Qin, D, Plattner, G-
- 3 K, Tignor, M, Allen, SK, Boschung, J, Nauels, A, Xia, Y, Bex, V and Midgley, PM (eds.),
- 4 Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental
- 5 Panel on Climate Change, 1535. Cambridge University Press, Cambridge, United Kingdom and
- 6 New York, NY, USA, 2013.
- 7
- 8 Ito, A., Nishina, K., Ishijima, K. et al. Emissions of nitrous oxide (N₂O) from soil surfaces and
- 9 their historical changes in East Asia: a model-based assessment. *Prog Earth Planet Sci* 5, 55,
- 10 <https://doi.org/10.1186/s40645-018-0215-4>, 2018.
- 11
- 12 Iyengar, G. R., V. S. Prasad, and K. J. Ramesh, Circulation characteristic associated with Inter
- 13 Tropical Convergence Zone during northern winter, *Curr. Sci.*, 76, 903–906, 1999.
- 14
- 15 Jensen, T. G.: Cross-equatorial pathways of salt and tracers from the northern Indian Ocean:
- 16 Modelling results, *Deep Sea Res., Part II*, 50, 2111 – 2128, 2003.
- 17
- 18 Jickells, T. D., E. Buitenhuis, K. Altieri, A. R. Baker, D. Capone, R. A. Duce, F. Dentener, K.
- 19 Fennel, M. Kanakidou, J. LaRoche, K. Lee, P. Liss, J. J. Middelburg, J. K. Moore, G. Okin, A.
- 20 Oschlies, M. Sarin, S. Seitzinger, J. Sharples, A. Singh, P. Suntharalingam, M. Uematsu, L. M.
- 21 Zamora: A reevaluation of the magnitude and impacts of anthropogenic atmospheric nitrogen
- 22 inputs on the ocean. *Global Biogeochem. Cy.*, 31, 289–305, [https://](https://doi.org/10.1002/2016GB005586)
- 23 doi.org/10.1002/2016GB005586, 2017.
- 24
- 25 Jordi, A., G. Basterretxea, A. Tovar-Sánchez, A. Alastuey, and X. Querol: Copper aerosols
- 26 inhibit phytoplankton growth, *Proc. Natl. Acad. Sci. USA*, 109 (52) 21246-21249; DOI:
- 27 [10.1073/pnas.1207567110](https://doi.org/10.1073/pnas.1207567110), 2012.
- 28
- 29 Kavitha, M. and P. R. Nair: Satellite-retrieved vertical profiles of methane over the Indian
- 30 region: impact of synoptic-scale meteorology, *Int. J. Remote Sens.*, 40:14, 5585-5616, DOI:
- 31 [10.1080/01431161.2019.1580791](https://doi.org/10.1080/01431161.2019.1580791), 2019.
- 32
- 33 Khemani, L. T., Momin, G. A., and Singh, G.: Variation in trace gases concentrations in
- 34 different environments in India, *PA- GEOPH.*, 125, 151–158, 1987.
- 35
- 36 Kitoh, A., Endo, H., Krishna Kumar, K., Cavalcanti, IFA, Goswami, P. and Zhou, T.: Monsoons
- 37 in a changing world: A regional perspective in a global context. *J. Geophys. Res.*, 118(8):3053–
- 38 3065, 2013.
- 39
- 40 Kim, P. S., Jacob, D. J., Liu, X., Warner, J. X., Yang, K., Chance, K., Thouret, V., and Nedelec,
- 41 P.: Global ozone–CO correlations from OMI and AIRS: constraints on tropospheric ozone
- 42 sources, *Atmos. Chem. Phys.*, 13, 9321–9335, <https://doi.org/10.5194/acp-13-9321-2013>,
- 43 2013.
- 44
- 45 Kornmüller, A.: Review of fundamentals and specific aspects of oxidation technologies in
- 46 marine waters. *Water Sci Technol*, 55 (12): 1–6. doi: <https://doi.org/10.2166/wst.2007.379>,
- 47 2007.
- 48
- 49 Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C.,
- 50 Toohey, M., Stenke, A., Schwarz, J. P., Weigel, R., Fueglistaler, S., Prata, F. J., Vernier, J.-P.,
- 51 Schlager, H., Barnes, J. E., Antuña-Marrero, J.-C., Fairlie, D., Palm, M., Mahieu, E., Notholt,

- 1 J., Rex, M., Bingen, C., Vanhellemont, F., Bourassa, A., Plane, J. M. C., Klocke, D., Carn, S.
- 2 A., Clarisse, L., Trickl, T., Neely, R., James, A. D., Rieger, L., Wilson, J. C., and Meland, B.:
- 3 Stratospheric aerosol—Observations, processes, and impact on climate, *Rev. Geophys.*, 54,
- 4 278–335, 10.1002/2015rg000511, 2016.
- 5
- 6 Krishnamurti, T. N., Jha, B., Rasch, P. J., and Ramanathan, V.: A high resolution global
- 7 reanalysis highlighting the winter monsoon, Part I, Reanalysis fields, *Met. Atmos. Phys.*, 64,
- 8 123–150, 1997a.
- 9
- 10 Krishnamurti, T. N., Jha, B., Rasch, P. J., and Ramanathan, V.: A high resolution global
- 11 reanalysis highlighting the winter monsoon, Part II, transients and passive tracer transports,
- 12 *Met. Atmos. Phys.*, 64, 151–171, 1997b.
- 13
- 14 Krishnan, R. et al.: Assessment of Climate Change over the Indian Region, Springer Singapore,
- 15 DOI 10.1007/978-981-15-4327-2, 2020.
- 16
- 17 Krom M. D., Shi Z., Stockdale A., Berman-Frank I., Giannakourou A., Herut B., Lagaria A.,
- 18 Papageorgiou N., Pitta P., Psarra S., Rahav E., Scoullos M., Stathopoulou E., Tsiola A., and
- 19 Tsagaraki T. M.: Response of the Eastern Mediterranean Microbial Ecosystem to Dust and Dust
- 20 Affected by Acid Processing in the Atmosphere, *Frontiers in Marine Science*, 3, 133
- 21 doi=10.3389/fmars.2016.00133, 2016.
- 22
- 23 Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V.,
- 24 Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt,
- 25 P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z., and Streets, D. G.: Aura OMI
- 26 observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmos. Chem.*
- 27 *Phys.*, 16, 4605–4629, 10.5194/acp-16-4605-2016, 2016.
- 28
- 29 Kuai, L., Worden, J. R., Campbell, J. E., Kulawik, S. S., Li, K.-F., Lee, M., Weidner, R. J.,
- 30 Montzka, S. A., Moore, F. L., Berry, J. A., Baker, I., Denning, A. S., Bian, H., Bowman, K.
- 31 W., Liu, J., and Yung, Y. L.: Estimate of carbonyl sulfide tropical oceanic surface fluxes using
- 32 Aura Tropospheric Emission Spectrometer observations, *J. Geophys. Res.*, 120, 11012–11023,
- 33 <https://doi.org/10.1002/2015JD023493>, 2015.
- 34
- 35 Kumar, K.R., Tiwari, Y.K., Valsala, V., and R. Murtugudde: On understanding the land–ocean
- 36 CO₂ contrast over the Bay of Bengal: A case study during 2009 summer monsoon. *Environ Sci*
- 37 *Pollut Res* 21, 5066–5075, <https://doi.org/10.1007/s11356-013-2386-2>, 2014.
- 38
- 39 Kunhikrishnan, T., Lawrence, M. G., von Kuhlmann, R., Richter, A., Ladstaetter, A., and
- 40 Burrows, J. P.: Analysis of tropospheric NO_x over Asia using the Model of Atmospheric
- 41 Transport and Chemistry (MATCH-MPIC) and GOME-satellite observations, *Atmos.*
- 42 *Environ.*, 38, 581–596, 2004.
- 43
- 44 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T.,
- 45 Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian
- 46 regions during 2000–2008: Regional Emission inventory in ASia(REAS) version 2, *Atmos.*
- 47 *Chem. Phys.*, 13, 11019–11058, doi:10.5194/acp-13-11019-2013, 2013.
- 48
- 49 Lachkar, Z., Levy, M. and Smith, S.: Intensification and deepening of the Arabian Sea oxygen
- 50 minimum zone in response to increase in Indian monsoon wind intensity. *Biogeosciences* 15,
- 51 159–186, 2018.

- Ladstätter-Weissenmayer, A., Altmeyer, H., Bruns, M., Richter, A., Rozanov, A., Rozanov, V., Wittrock, F., and Burrows, J. P.: Measurements of O₃, NO₂ and BrO during the INDOEX campaign using ground based DOAS and GOME satellite data, *Atmos. Chem. Phys.*, 7, 283–291, 2007.
- Lago, V., Wjffels, S.E., Durack, P.J., Church, J.A., Bindoff, N.L., and Marsland, S.J.: Simulating the role of surface forcing on observed multidecadal upper-ocean salinity changes. *J. Clim.*, 29:5575–5588. <https://doi.org/10.1175/JCLI-D-15-0519.1>, 2016.
- Lal, S., Chand, D., Sahu, L. K., Venkataramani, S., Brasseur, G., and Schultz, M. G.: High levels of ozone and related gases over the Bay of Bengal during winter and early spring of 2001, *Atmos. Environ.*, 40, 1633–1644, 2006.
- Lal, S. and Lawrence, M. G.: Elevated mixing ratios of surface ozone over the Arabian Sea, *Geophys. Res. Lett.*, 28, 1487–1490, 2001.
- Lal, S., Sahu, L. K., and Venkataramani, S.: Impact of transport from the surrounding continental regions on the distributions of ozone and related trace gases over the Bay of Bengal during February 2003, *J. Geophys. Res.*, 112, D14302, doi:10.1029/2006JD008023, 2007.
- Lal, S., Venkataramani, S., Srivastava, S., Gupta, S., Mallik, C., Naja, M., Sarangi, T., Acharya, Y. B., and Liu, X.: Transport effects on the vertical distribution of tropospheric ozone over the tropical marine regions surrounding India, *J. Geophys. Res.*, 118, 1513–1524, <https://doi.org/10.1002/jgrd.50180>, 2013.
- Lal, S., Venkataramani, S., Chandra, N., Cooper, O. R., Brioude, J., and Naja, M.: Transport effects on the vertical distribution of tropospheric ozone over western India, *J. Geophys. Res.*, 119, 10012–10026, 10.1002/2014jd021854, 2014.
- Lamb, P. J., and Hastenrath, S., *Climatic Atlas of the Indian Ocean: Surface Climate and Atmospheric Circulation*. Madison: The University of Wisconsin Press, 1979.
- Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cy.*, 25, 10.1029/2010gb003850, 2011.
- Latif, M. and T.P. Barnett: Interactions of the Tropical Oceans. *J. Climate*, 8, 952–964, [https://doi.org/10.1175/1520-0442\(1995\)008<0952:IOTTO>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<0952:IOTTO>2.0.CO;2), 1995.
- Launois, T., Belviso, S., Bopp, L., Fichot, C. G., and Peylin, P.: A new model for the global biogeochemical cycle of carbonyl sulfide; Part 1: Assessment of direct marine emissions with an oceanic general circulation and biogeochemistry model, *Atmos. Chem. Phys.*, 15, 2295–2312, 10.5194/acp-15-2295-2015, 2015.
- Lawrence, M. G., Rasch, P. J., von Kuhlmann, R., Williams, J., Fischer, H., de Reus, M., Lelieveld, J., Crutzen, P. J., Schultz, M., Stier, P., Huntrieser, H., Heland, J., Stohl, A., Forster, C., Elbern, H., Jakobs, H., and Dickerson, R. R.: Global chemical weather forecasts for field campaign planning: predictions and observations of large-scale features during MINOS,

CONTRACE, and INDOEX, *Atmos. Chem. Phys.*, 3, 267–289, doi:10.5194/acp-3- 267-2003, 2003.

Lawrence, M.G., Export of Air Pollution from Southern Asia and its Large-Scale Effects. In: Stohl A. (eds) *Air Pollution. The Handbook of Environmental Chemistry*, vol 4G. Springer, Berlin, Heidelberg, 2004.

Lawrence and Lelieveld, Atmospheric pollutants outflow from southern Asia: a review, *Atmos. Chem. Phys.*, 10, pp. 11017-11096, 2010.

Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneeth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozzi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, *Earth Syst. Sci. Data*, 10, 2141-2194, 10.5194/essd-10-2141-2018, 2018.

Lee, C.-L., and Brimblecombe, P.: Anthropogenic contributions to global carbonyl sulfide, carbon disulfide and organosulfides fluxes, *Earth-Sci. Rev.*, 160, 1-18, <https://doi.org/10.1016/j.earscirev.2016.06.005>, 2016.

Legrand, M., McConnell, J. R., Preunkert, S., Arienzo, M., Chellman, N., Gleason, K., Sherwen, T., Evans, M. J. and Carpenter, L. J.: Alpine ice evidence of a three-fold increase in atmospheric iodine deposition since 1950 in Europe due to increasing oceanic emissions, *Proc. Natl. Acad. Sci.*, 115(48), 12136–12141, doi:10.1073/pnas.1809867115, 2018.

Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367-371, 10.1038/nature15371, 2015.

Lelieveld, J., Crutzen, P. J., Ramanathan, V., Andreae, M. O., Brenninkmeijer, C. A. M., Campos, T., Cass, G. R., Dickerson, R. R., Fischer, H., de Gouw, J. A., Hansel, A., Jefferson, A., Kley, D., de Laat, A. T. J., Lal, S., Lawrence, M. G., Lobert, J. M., Mayol-Bracero, O. L., Mitra, A. P., Novakov, T., Oltmans, S. J., Prather, K. A., Reiner, T., Rodhe, H., Scheeren, H. A., Sikka, D., and Williams, J.: The Indian Ocean experiment: Widespread air pollution from South and Southeast Asia, *Science*, 291, 1031–1036, 2001.

Lelieveld, J., Bourtsoukidis, E., Brühl, C., Fischer, H., Fuchs, H., Harder, H., Hofzumahaus, A., Holland, F., Marno, D., Neumaier, M., Pozzer, A., Schlager, H., Williams, J., Zahn, A., and Ziereis, H.: The South Asian monsoon – Pollution pump and purifier, *Science*, 361, 270–273, <https://doi.org/10.1126/science.aar2501>, 2018.

Lelieveld, J., Klingmüller, A., Pozzer, R.T., Burnett, A., Haines and V. Ramanathan: Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *Proc. Natl.*

Acad. Sci. USA, 116 (15), 7192-7197, doi:www.pnas.org/cgi/doi/10.1073/pnas.1819989116, 2019.

Lennartz, S. T., Marandino, C. A., von Hobe, M., Cortes, P., Quack, B., Simo, R., Booge, D., Pozzer, A., Steinhoff, T., Arevalo-Martinez, D. L., Kloss, C., Bracher, A., Röttgers, R., Atlas, E., and Krüger, K.: Direct oceanic emissions unlikely to account for the missing source of atmospheric carbonyl sulfide, *Atmos. Chem. Phys.*, 17, 385–402, <https://doi.org/10.5194/acp-17-385-2017>, 2017.

Lennartz, S. T., Marandino, C. A., von Hobe, M., Andreae, M. O., Aranami, K., Atlas, E., Berkelhammer, M., Bingemer, H., Booge, D., Cutter, G., Cortes, P., Kremser, S., Law, C. S., Marriner, A., Simó, R., Quack, B., Uher, G., Xie, H., and Xu, X.: Marine carbonyl sulfide (OCS) and carbon disulfide (CS₂): a compilation of measurements in seawater and the marine boundary layer, *Earth Syst. Sci. Data*, 12, 591–609, <https://doi.org/10.5194/essd-12-591-2020>, 2020.

Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Malkki, A., Huib, V., Johan de, V., Stammes, P., Lundell, J. O. V., and Saari, H.: The ozone monitoring instrument, *IEEE T. Geosci. Remote Sens.*, 44, 1093–1101, <https://doi.org/10.1109/tgrs.2006.872333>, 2006.

Li, Z., Lau, W.K., Ramanathan, V., Wu, G., Ding, Y., Manoj, M.G., Liu, J., Qian, Y., Li, J., Zhou, T., Fan, J., Rosenfeld, D., Ming, Y., Wang, Y., Huang, J., Wang, B., Xu, X., Lee, S.-S., Cribb, M., Zhang, F., Yang, X., Zhao, C., Takemura, T., Wang, K., Xia, X., Yin, Y., Zhang, H., Guo, J., Zhao, P., Sugimoto, N., Babu, S. S., and Brasseur, G. P.: Aerosol and monsoon climate interactions over Asia, *Rev. Geophys.*, 54, 866–929, 2016.

Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935–963, <https://doi.org/10.5194/acp-17-935-2017>, 2017.

Li, T., Wang, B., Chang, C.P., and Y. Zhang: A theory for the Indian Ocean dipole–zonal mode. *J. Atmos. Sci.*, 60(17):2119–2135, 2003.

Li, Y., Han W., A. Hu, G. A. Meehl, and F. Wang: Multidecadal changes of the upper Indian Ocean heat content during 1965–2016. *J. Climate*, 31, 7863–7884, <https://doi.org/10.1175/JCLI-D-18-0116.1>, 2018.

Liang, Q., Stolarski, R. S., Kawa, S. R., Nielsen, J. E., Douglass, A.R., Rodriguez, J. M., Blake, D. R., Atlas, E. L., and Ott, L. E.: Finding the missing stratospheric Br_y: a global modeling study of CHBr₃ and CH₂Br₂, *Atmos. Chem. Phys.*, 10, 2269–2286, <https://doi.org/10.5194/acp-10-2269-2010>, 2010.

Liu, S.C., McFarland, M., Kley, D., Zafiriou, O., Huebert, B.: Tropospheric NO_x and O₃ budgets in the equatorial Pacific. *Journal of Geophysical Research* 88, 1360e1368, 1983.

Liu, X., Bhartia, P. K., Chance, K., Spurr, R. J. D., and Kurosu, T. P.: Ozone profile retrievals from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 10, 2521–2537, <https://doi.org/10.5194/acp-10-2521-2010>, 2010.

- 1 Llovel, W. and Lee, T.: Importance and origin of halosteric contribution to sea level change in
2 the southeast Indian Ocean during 2005–2013. *Geophys. Res. Lett.*, 42, 1148–1157, 2015.
- 3
- 4 Löscher, C. R.: Reviews and syntheses: Trends in primary production in the Bay of Bengal – is
5 it at a tipping point?, *Biogeosciences*, 18, 4953–4963, [https://doi.org/10.5194/bg-18-4953-](https://doi.org/10.5194/bg-18-4953-2021)
6 2021, 2021.
- 7
- 8 Lombardozi, D, Levis, S, Bonan, G, Hess, P. G. and Sparks, J. P.: The influence of chronic
9 ozone exposure on global carbon and water cycles. *J. Climate*, 28: 292–305. DOI:
10 10.1175/JCLI-D-14-00223.1, 2015.
- 11
- 12 Lunt, M. F., Palmer, P. I., Feng, L., Taylor, C. M., Boesch, H., and Parker, R. J.: An increase
13 in methane emissions from tropical Africa between 2010 and 2016 inferred from satellite data,
14 *Atmos. Chem. Phys.*, 19, 14721–14740, <https://doi.org/10.5194/acp-19-14721-2019>, 2019.
- 15
- 16 Luo, G. and Yu, F.: A numerical evaluation of global oceanic emissions of α -pinene and
17 isoprene, *Atmos. Chem. Phys.*, 10, 2007–2015, doi:10.5194/acp-10-2007-2010, 2010.
- 18
- 19 Ma, X., Bange, H. W., Eirund, G. K., and Arévalo-Martínez, D. L.: Nitrous oxide and
20 hydroxylamine measurements in the Southwest Indian Ocean. *J. Marine Syst.*,
21 <https://doi.org/10.1016/j.jmarsys.2018.03.003>, 2018.
- 22
- 23 Maas, J., Jia, Y., Quack, B., Durgadoo, J. V., Biastoch, A., and Tegtmeier, S.: Simulations of
24 anthropogenic bromoform indicate high emissions at the coast of East Asia, *Atmos. Chem.*
25 *Phys. Discuss.*, 2020, 1–31, 10.5194/acp-2019-1004, 2020.
- 26
- 27 Maas, J., Tegtmeier, S., Jia, Y., Quack, B., Durgadoo, J. V., and Biastoch, A.: Simulations of
28 anthropogenic bromoform indicate high emissions at the coast of East Asia, *Atmos. Chem.*
29 *Phys.*, 21, 4103–4121, <https://doi.org/10.5194/acp-21-4103-2021>, 2021.
- 30
- 31
- 32 Madden, R. A. and P. R. Julian: Description of Global-Scale Circulation Cells in the Tropics
33 with a 40–50 Day Period. *J. Atmos. Sci.*, 29, 1109–1123, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0469(1972)029<1109:DOGSCC>2.0.CO;2)
34 0469(1972)029<1109:DOGSCC>2.0.CO;2, 1972.
- 35
- 36 Mahajan, A. S., Plane, J. M. C., Oetjen, H., Mendes, L., Saunders, R. W., Saiz-Lopez, A., Jones,
37 C. E., Carpenter, L. J., and McFiggans, G. B.: Measurement and modelling of tropospheric
38 reactive halogen species over the tropical Atlantic Ocean, *Atmos. Chem. Phys.*, 10, 4611–4624,
39 <https://doi.org/10.5194/acp-10-4611-2010>, 2010.
- 40
- 41 Mahajan, A. S., Tinel, L., Hulswar, S., Cuevas, C. A., Wang, S., Ghude, S., et al.: Observations
42 of iodine oxide in the Indian Ocean Marine Boundary Layer: a transect from the tropics to the
43 high latitudes, *Atmos. Environ.: X*. <https://doi.org/10.1016/j.aeaoa.2019.100016>, 2019a.
- 44
- 45 Mahajan, A. S., Tinel, L., Sarkar, A., Chance, R., Carpenter, L. J., Hulswar, S., et al.:
46 Understanding Iodine Chemistry over the Northern and Equatorial Indian Ocean. *J. Geophys.*
47 *Res.*, (x), 0–3. <https://doi.org/10.1029/2018JD029063>, 2019b.
- 48
- 49 Mahajan, A.S., De Smedt, I., Biswas, M.S., Ghude, S.D., Fadnavis, S., Roy, C., van
50 Roozendaal, M.: Inter-annual variations in satellite observations of nitrogen dioxide and

formaldehyde over India. *Atmos. Environ.* 116, 194–201.
<https://doi.org/10.1016/j.atmosenv.2015.06.004>, 2015a.

Mahajan, A. S., Fadnavis, S., Thomas, M. A., Pozzoli, L., Gupta, S., Royer, S., et al.: Quantifying the impacts of an updated global dimethyl sulfide climatology on cloud microphysics and aerosol radiative forcing. *J. Geophys. Res.*, 120, 1–13.
<https://doi.org/10.1002/2014JD022687>, 2015b.

Mahowald, N. M., Hamilton, D. S., Mackey, K.R.M., Moore, J. K., Baker, A. R., Scanza, R. A., and Zhang, Y.: Aerosol trace metal leaching and impacts on marine microorganisms. *Nat. Commun.*, 9, 2614, <https://doi.org/10.1038/s41467-018-04970-7>, 2018.

Mallik, C., Lal, S., Venkataramani, S., Naja, M., and Ojha, N.: Variability in ozone and its precursors over the Bay of Bengal during post-monsoon: Transport and emission effects. *J. Geophys. Res.*, <https://doi.org/10.1002/jgrd.50764>, 2013.

Mandal, T. K., Khan, A., Ahammed, Y. N., Tanwar, R. S., Parmar, R. S., Zalpuri, K. S., et al.: Observations of trace gases and aerosols over the Indian Ocean during the monsoon transition period. *J. Earth Syst. Sci.*, 115(4), 473–484, 2006.

Manö, S., and Andreae, M. O.: Emission of Methyl Bromide from Biomass Burning, *Science*, 263, 1255–1257, [10.1126/science.263.5151.1255](https://doi.org/10.1126/science.263.5151.1255), 1994.

Marbach, T., Beirle, S., Platt, U., Hoor, P., Wittrock, F., Richter, A., Vrekoussis, M., Grzegorski, M., Burrows, J. P., and Wagner, T.: Satellite measurements of formaldehyde linked to shipping emissions, *Atmos. Chem. Phys.*, 9, 8223–8234, <https://doi.org/10.5194/acp-9-8223-2009>, 2009.

Martino, M., Mills, G. P., Woeltjen, J., and Liss, P. S.: A new source of volatile organoiodine compounds in surface seawater, *Geophys. Res. Lett.*, 36, L01609, [doi:10.1029/2008GL036334](https://doi.org/10.1029/2008GL036334), 2009.

Mason, R. P., and Sheu, G.-R.: Role of the ocean in the global mercury cycle, *Global Biogeochem. Cy.*, 16, 40–41–40–14, [10.1029/2001gb001440](https://doi.org/10.1029/2001gb001440), 2002.

Matthews, A. J., Singhruck, P. and Heywood, K. J.: Ocean temperature and salinity components of the Madden–Julian oscillation observed by Argo floats. *Clim. Dynam.*, 35, 1149–1168, <https://doi.org/10.1007/s00382-009-0631-7>, 2010.

Meehl, G. A., and J. M. Arblaster: Mechanisms for projected future changes in South Asian monsoon precipitation. *Clim. Dynam.*, 21, 659–675, <https://doi.org/10.1007/s00382-003-0343-3>, 2003.

Meenu, S., Rajeev, K., Parameswaran, K., and Suresh Raju, C., Characteristics of the double intertropical convergence zone over the tropical Indian Ocean, *J. Geophys. Res.*, 112, D11106, [doi:10.1029/2006JD007950](https://doi.org/10.1029/2006JD007950), 2007.

Mehlmann, M., Quack, B., Atlas, E., Hepach, H., and Tegtmeier, S.: Natural and anthropogenic sources of bromoform and dibromomethane in the oceanographic and biogeochemical regime of the subtropical North East Atlantic, *Environ. Sci. Process. Impacts*, 22, 679–707, [10.1039/c9em00599d](https://doi.org/10.1039/c9em00599d), 2020.

- 1
- 2 Metzl, N.: Decadal increase of oceanic carbon dioxide in Southern Indian Ocean surface waters
- 3 (1991–2007), *Deep Sea Res. Part II: Topical Studies in Oceanogr.*, 56, 607–619,
- 4 <https://doi.org/10.1016/j.dsr2.2008.12.007>, 2009.
- 5
- 6 Mihalopoulos, N., Putaud, J. P., Nguyen, B. C., and Belviso, S.: Annual variation of
- 7 atmospheric carbonyl sulfide in the marine atmosphere in the Southern Indian Ocean, *J. Atmos.*
- 8 *Chem.*, 13, 73–82, <https://doi.org/10.1007/bf00048101>, 1991.
- 9
- 10 Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., Gascon, C.: Global
- 11 biodiversity conservation: the critical role of hotspots. *Biodiversity hotspots*, Berlin,
- 12 Heidelberg: Springer. pp. 3–22. doi:10.1007/978-3-642-20992-5, 2011.
- 13
- 14 Mohan, S. and Bhaskaran, P.K.: Evaluation of CMIP5 climate model projections for surface
- 15 wind speed over the Indian Ocean region. *Clim. Dynam.*, 53(9–10), 5415–5435, 2019.
- 16
- 17 Monks, P. S., Carpenter, L. J., Penkett, S. A., Ayers, G. P., Gillett, R. W., Galbally, I. E., and
- 18 Meyer, C. P.: Fundamental ozone photochemistry in the remote marine boundary layer: The
- 19 SOAPEX experiment, measurement and theory. *Atmos. Environ.*, 32(21), 3647–3664, 1998.
- 20
- 21 Monks, PS, Archibald, AT, Colette, A, Cooper, O, Coyle, M, Derwent, R, Fowler, D, Granier,
- 22 C, Law, KS, Mills, GE, Stevenson, DS, Tarasova, O, Thouret, V, von Schneidemesser, E,
- 23 Sommariva, R, Wild, O and Williams, ML.: Tropospheric ozone and its precursors from the
- 24 urban to the global scale from air quality to short-lived climate forcer. *Atmos. Chem. Phys.* 15:
- 25 8889–8973. DOI: 10.5194/acp-15-8889-2015, 2015.
- 26
- 27 Moorthy, K. K., Satheesh, S. K., Babu, S. S., and Dutt, C. B. S.: Integrated Campaign for
- 28 Aerosols, gases and Radiation Budget (ICARB): An overview, *J. Earth Syst. Sci.*, 117, 243–
- 29 262, 10.1007/s12040-008-0029-7, 2008.
- 30
- 31 Mungall, E. L., J. P. D. Abbatt, J. J. B. Wentzell, A. K. Y. Lee, J. L. Thomas, M. Blais, M.
- 32 Gosselin, L. A. Miller, T. Papakyriakou, M. D. Willis, and J. Liggio: OVOCs in the
- 33 summertime marine Arctic atmosphere, *Proceedings of the National Academy of Sciences*, 114
- 34 (24) 6203–6208; DOI: 10.1073/pnas.1620571114, 2017.
- 35
- 36 Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestad, J. Huang, D. Koch, J.-F.
- 37 Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and H.
- 38 Zhang: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical*
- 39 *Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*
- 40 *Intergovernmental Panel on Climate Change*, 2013.
- 41
- 42 Nair, P. R., David, L. M., Girach, I. A., and George, S. K.: Ozone in the marine boundary layer
- 43 of Bay of Bengal during post-winter period: Spatial pattern and role of meteorology, *Atmos.*
- 44 *Environ.*, 45, 4671–4681, 2011.
- 45
- 46 Naja, M., Chand, D., Sahu, L., and Lal, S.: Trace gases over marine regions around India, *Indian*
- 47 *J. Mar. Sci.*, 33, 95–106, 2004.
- 48
- 49 Nalini, K., Uma, K. N., Sijikumar, S., Tiwari, Y. K., and Ramachandran, R.: Satellite- and
- 50 ground-based measurements of CO₂ over the Indian region: its seasonal dependencies, spatial

variability, and model estimates, *Int. J. Remote Sens.*, 39, 7881-7900, 10.1080/01431161.2018.1479787, 2018.

Naqvi, S. W. A., Jayakumar, D. A., Nair, M., Kumar, M. D., and George, M. D.: Nitrous oxide in the western Bay of Bengal, *Marine Chem.*, 47, 269-278, [https://doi.org/10.1016/0304-4203\(94\)90025-6](https://doi.org/10.1016/0304-4203(94)90025-6), 1994.

Naqvi, S. W. A., Bange, H. W., Gibb, S. W., Goyet, C., Hatton, A. D., and Upstill-Goddard, R. C.: Biogeochemical ocean-atmosphere transfers in the Arabian Sea, *Progress Oceanogr.*, 65, 116-144, <https://doi.org/10.1016/j.pocean.2005.03.005>, 2005.

Naqvi, S. W. A., Jayakumar, D. A., Narvekar, P. V., Naik, H., Sarma, V. V. S. S., D'Souza, W., Joseph, S., George, M. D.: Marine hypoxia/anoxia as a source of CH₄ and N₂O, *Biogeosciences* 7, 2159–2190, 2010a.

Naqvi, S. W. A., Naik, H., D'Souza, W., Narvekar, P. V., Paropkari, A. L., and Bange, H. W.: Carbon and nitrogen fluxes in the northern Indian Ocean. In: Liu, K.-K., Atkinson, L., Quiñones, R., Talaue-McManus, L. (Eds.), *Carbon and Nutrient Fluxes in Continental Margins: A Global Synthesis*. Springer-Verlag, New York, pp. 180–191, 2010b.

Nayak, R. K., Dadhwal, V. K., Majumdar, A., Patel, N. R., and Dutt, C. B. S.: Variability of atmospheric CO₂ over India and Surrounding Oceans and control by Surface Fluxes, *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, XXXVIII-8/W20, 96-101, 10.5194/isprsarchives-XXXVIII-8-W20-96-2011, 2011.

Nieves, V., J. K. Willis, and W. C. Patzert: Recent hiatus caused by decadal shift in Indo-Pacific heating. *Science*, 349, 532–535, <https://doi.org/10.1126/science.aaa4521>, 2015.

Norman, M. and Leck, C.: Distribution of marine boundary layer ammonia over the Atlantic and Indian Ocean during the Aerosols 99 cruise, *J. Geophys. Res.*, 110, D16302, doi:10.1029/2005JD005866, 2005.

Nowak, J.B., et al.: Gas-phase chemical characteristics of Asian emission plumes observed during ITCT 2K2 over the eastern North Pacific Ocean, *J. Geophys. Res.* 109, D23S19. doi:10.1029/2003JD004488, 2004.

Oetjen, H.: Measurements of halogen oxides by scattered sunlight differential optical absorption spectroscopy, University of Bremen, 2009.

Ojha, N., Naja, M., Singh, K. P., Sarangi, T., Kumar, R., Lal, S., Lawrence, M. G., Butler, T. M., and Chandola, H. C.: Variabilities in ozone at a semi-urban site in the Indo-Gangetic Plain region: Association with the meteorology and regional process, *J. Geophys. Res.*, 117, D20301, doi:10.1029/2012JD017716, 2012.

Pacyna, J. M., Pacyna, E. G.: Anthropogenic sources and global inventory of mercury emissions. In: Parsons, M.B., Percival, J.B. (Eds.), *Mercury: Sources, Measurements, Cycles, and Effects*. Short Course Series, vol. 32. Mineralogical Association of Canada, 2005.

- 1 Palastanga, V., van Leeuwen, P. J., Schouten, M. W., and de Ruijter, W. P. M.: Flow structure
2 and variability in the subtropical Indian Ocean: Instability of the South Indian Ocean
3 Countercurrent, *J. Geophys. Res.*, 112, C01001, doi:10.1029/2005JC003395, 2007.
- 4
- 5 Palmer, P. I., and Shaw, S. L.: Quantifying global marine isoprene fluxes using MODIS
6 chlorophyll observations, *Geophys. Res. Lett.*, 32, 10.1029/2005gl022592, 2005.
- 7
- 8 Pandey, P. C., Khare, N., and Sudhakar, M.: Oceanographic research: Indian efforts and
9 preliminary results from the Southern Ocean, *Current Sci.*, 90, 978-984, 2006.
- 10
- 11 Pant, V., Deshpande, C. G., and Kamra, A. K.: The concentration and number size distribution
12 measurements of the Marine Boundary Layer aerosols over the Indian Ocean, *Atmos. Res.*, 92,
13 381–393, 2009.
- 14
- 15 Pathak, H., Li, C., and Wassmann, R.: Greenhouse gas emissions from Indian rice fields:
16 calibration and upscaling using the DNDC model, *Biogeosciences*, 2, 113-123, 10.5194/bg-2-
17 113-2005, 2005.
- 18
- 19 Paytan, A., K. R. M. Mackey, Y. Chen, I. D. Lima, S. C. Doney, N. Mahowald, R. Labiosa, A.
20 F. Post: Toxicity of atmospheric aerosols on marine phytoplankton, *Proc. Natl. Acad. Sci. USA*,
21 106 (12) 4601-4605; DOI: 10.1073/pnas.0811486106, 2009.
- 22
- 23 Peters, G.P., Andrew, R.M., Canadell, J.G. et al.: Carbon dioxide emissions continue to grow
24 amidst slowly emerging climate policies. *Nat. Clim. Chang.* 10, 3–6,
25 <https://doi.org/10.1038/s41558-019-0659-6>, 2020.
- 26
- 27 Pfannerstill, E. Y., Wang, N., Edtbauer, A., Bourtsoukidis, E., Crowley, J. N., Dienhart, D.,
28 Eger, P. G., Ernle, L., Fischer, H., Hottmann, B., Paris, J.-D., Stöner, C., Tadic, I., Walter, D.,
29 Lelieveld, J., and Williams, J.: Shipborne measurements of total OH reactivity around the
30 Arabian Peninsula and its role in ozone chemistry, *Atmos. Chem. Phys.*, 19, 11501–11523,
31 <https://doi.org/10.5194/acp-19-11501-2019>, 2019.
- 32
- 33 Phillips, H. E., Tandon, A., Furue, R., Hood, R., Ummenhofer, C., Benthuyssen, J., Menezes,
34 V., Hu, S., Webber, B., Sanchez-Franks, A., Cherian, D., Shroyer, E., Feng, M., Wijesekera, H.,
35 Chatterjee, A., Yu, L., Hermes, J., Murtugudde, R., Tozuka, T., Su, D., Singh, A., Centurioni,
36 L., Prakash, S., and Wiggert, J.: Progress in understanding of Indian Ocean circulation,
37 variability, air-sea exchange and impacts on biogeochemistry, *Ocean Sci. Discuss.* [preprint],
38 <https://doi.org/10.5194/os-2021-1>, in review, 2021.
- 39
- 40 Pitts, B. J. F., and Pitts, J. N.: *Chemistry of the Upper and Lower Atmosphere*. Academic Press,
41 California, 2000
- 42 Portmann, R.W., J.S. Daniel, and A.R. Ravishankara: Stratospheric ozone depletion due to
43 nitrous oxide: influences of other gases. *Philosophical Transactions of the Royal Society*
44 *London B Biological Sciences* 367 (1593): 1256–1264. <https://doi.org/10.1098/rstb.2011.0377>,
45 2012.
- 46
- 47 Prasanna Kumar, S., Muraleedharan, P. M., Prasad, T. G., Gauns, M., Ramaiah, N., de Souza,
48 S. N., Sardesai, S., and Madhupratap, M.: Why is the Bay of Bengal less productive during
49 summer monsoon compared to the Arabian Sea? *Geophys. Res. Lett.*, 29(24), 2235,
50 doi:10.1029/2002GL016013, 2002.
- 51

- 1 Prather, M. J., Hsu, J., DeLuca, N. M., Jackman, C. H., Oman, L. D., Douglass, A. R., Fleming,
2 E. L., Strahan, S. E., Steenrod, S. D., Søvdé, O. A., Isaksen, I. S. A., Froidevaux, L., and Funke,
3 B: Measuring and modeling the lifetime of nitrous oxide including its variability. *J. Geophys.*
4 *Res.*, 120, 5693– 5705. doi: 10.1002/2015JD023267, 2015.
- 5
6 Quack, B., Atlas, E., Petrick, G., and Wallace, D.: Bromoform and dibromomethane above the
7 Mauritanian upwelling: Atmospheric distributions and oceanic emissions, *J. Geophys.*
8 *Res.*, 112, D09312, doi:10.1029/2006JD007614, 2007.
- 9
10 Quartly, G. D., J. J. H. Buck, M. A. Srokosz, and A. C. Coward: Eddies around Madagascar—
11 The retroflection reconsidered, *J. Mar. Syst.*, 63(3–4), 115–129,
12 doi:10.1016/j.jmarsys.2006.06.001, 2006.
- 13
14 Quinn, P. K., Coffman, D. J., Johnson, J. E., Upchurch, L. M., and Bates, T. S.: Small fraction
15 of marine cloud condensation nuclei made up of sea spray aerosol. *Nat. Geosci.*, 10(9), 674–
16 679, 2017.
- 17
18 Raes, E. J., Bodrossy, L., Van de Kamp, J., Holmes, B., Hardman-Mountford, N., Thompson,
19 P. A., McInnes, A. S., and Waite, A. M.: Reduction of the Powerful Greenhouse Gas N₂O in
20 the South-Eastern Indian Ocean, *PLOS ONE*, 11, e0145996, 10.1371/journal.pone.0145996,
21 2016.
- 22
23 Rahav, E., Belkin, N., Paytan, A., and Herut, B.: Phytoplankton and Bacterial Response to
24 Desert Dust Deposition in the Coastal Waters of the Southeastern Mediterranean Sea: A Four-
25 Year In Situ Survey. *Atmosphere*, 9, 305, 2018.
- 26
27 Rajak, R. and Chattopadhyay, A., Short and long-term exposure to ambient air pollution and
28 impact on health in India: a systematic review, *Int. J. Environ. Health Res.*,
29 10.1080/09603123.2019.1612042, 2019.
- 30
31 Ramanathan, V., Crutzen, P. J., Mitra, A. P., and Sikka, D.: The Indian Ocean Experiment and
32 the Asian Brown Cloud, *Curr. Sci. India*, 83, 947–955, 2002.
- 33
34 Ramanathan, V., C. Chung, D. Kim, T. Bettge, L. Buja, J. T. Kiehl, W. M. Washington, Q. Fu,
35 D. R. Sikka, M. Wild: Atmospheric brown clouds: Impacts on South Asian climate and
36 hydrological cycle, *Proc. Natl. Acad. Sci. USA*, 102 (15) 5326-5333; DOI:
37 10.1073/pnas.0500656102, 2005.
- 38
39 Randel, W. J., Park, M., Emmons, L., Kinnison, D., Bernath, P., Walker, K. A., Boone, C., and
40 Pumphrey, H.: Asian monsoon transport of pollution to the stratosphere, *Science*, 328, 611–
41 613, doi:10.1126/science.1182274, 2010.
- 42
43 Rao, R. R., and Sivakumar, R.: Seasonal variability of sea surface salinity and salt budget of
44 the mixed layer of the north Indian Ocean, *J. Geophys. Res.*, 108(C1), 3009,
45 doi:10.1029/2001JC000907, 2003.
- 46
47 Raut, N., B. K. Sitaula, L. R. Bakken, R. M. Bajracharya, and P. Dörsch.: Higher N₂O emission
48 by intensified crop production in South Asia. *Global Ecology and Conservation* 4:176–84.
49 doi:10.1016/j.gecco.2015.06.004, 2015.
- 50

- 1 Ravishankara, A., J. S. Daniel, and R. W. Portmann: Nitrous oxide (N₂O): The dominant ozone-
2 depleting substance emitted in the 21st century, *Science*, 326(5949), 123–125,
3 doi:10.1126/science.1176985, 2009.
- 4
- 5 Rayman, M. P.: The importance of selenium to human health. *Lancet*, 356, 233–241, 2000.
- 6
- 7 Read, K., A. Mahajan, L. Carpenter, M. J. Evans, B. V. E. Faria, D. E. Heard, J. R. Hopkins, J.
8 D. Lee, S. J. Moller, A. C. Lewis, L. Mendes, J. B. McQuaid, H. Oetjen, A. Saiz-Lopez, M. J.
9 Pilling, and J. M. C. Plane: Extensive halogen-mediated ozone destruction over the tropical
10 Atlantic Ocean. *Nature* 453, 1232–1235, <https://doi.org/10.1038/nature07035>, 2008.
- 11
- 12 Reppin, J., F. Schott, and J. Fischer: Equatorial currents and transports in the upper central
13 Indian Ocean: Annual cycle and interannual variability, *J. Geophys. Res.*, 104(C7), 15,495 –
14 15,514, 1999.
- 15
- 16 Richter, U. and Wallace, D. W. R.: Production of methyl iodide in the tropical Atlantic Ocean,
17 *Geophys. Res. Lett.*, 31(23), 2004.
- 18
- 19 ~~Richter, I., and S.P. Xie: Muted precipitation increase in global warming simulations: a surface~~
20 ~~evaporation perspective. *J. Geophys. Res.*, 113:D24118, doi.org/10.1029/2008J-D0105-61,~~
21 ~~2008.~~
- 22
- 23 Ridderinkhof, H., and W. P. M. de Ruijter: Moored current observations in the Mozambique
24 Channel, *Deep Sea Res. Part II*, 50, 1933–1955, 2003.
- 25
- 26 Rixen, T., Cowie, G., Gaye, B., Goes, J., do Rosário Gomes, H., Hood, R. R., Lachkar, Z.,
27 Schmidt, H., Segschneider, J., and Singh, A.: Reviews and syntheses: Present, past, and future
28 of the oxygen minimum zone in the northern Indian Ocean, *Biogeosciences*, 17, 6051–6080,
29 <https://doi.org/10.5194/bg-17-6051-2020>, 2020.
- 30
- 31 Rixen, T., Goyet, C., and Ittekkot, V.: Diatoms and their influence on the biologically mediated
32 uptake of atmospheric CO₂ in the Arabian Sea upwelling system, *Biogeosciences*, 3, 1–13,
33 <https://doi.org/10.5194/bg-3-1-2006>, 2006.
- 34
- 35 Rodríguez-Ros, P., Cortés, P., Robinson, C. M., Nunes, S., Hassler, C., Royer, S. J., Estrada,
36 M., Sala, M. M. and Simó, R.: Distribution and drivers of marine isoprene concentration across
37 the Southern Ocean, *Atmosphere (Basel)*, 11(6), 1–19, doi:10.3390/atmos11060556, 2020.
- 38
- 39 Roser, M., Ritchie H., and Ortiz-Ospina, E.: World Population Growth, published online at
40 OurWorldInData.org., 2013. Roxy, M. K., Raghu Murtugudde, A. M., Valsala, V., Panickal,
41 S., Prasanna Kumar, S., Ravichandran, M., et al.: A reduction in marine primary productivity
42 driven by rapid warming over the tropical Indian Ocean. *Geophys. Res. Lett.*, 43, 826–833. doi:
43 10.1002/2015gl066979, 2016.
- 44
- 45 Roxy M. K., Ritika K., Terray P., Murtugudde R., Ashok K., Goswami B. N.: Drying of Indian
46 subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient, *Nat.*
47 *Commun.*, 6:7423. doi.org/10.1038/ncomm s8423, 2015.
- 48
- 49 Roxy, M. K., Ritika, K., Terray, P. and Masson, S.: The curious case of Indian Ocean warming.
50 *J. Clim.* 27, 8501–8509, 2014.
- 51

- Roy, R., Pratihary, A., Narvenkar, G., Mochemadkar, S., Gauns, M., and Naqvi, S. W. A.: The relationship between volatile halocarbons and phytoplankton pigments during a Trichodesmium bloom in the coastal eastern Arabian Sea, *Estuarine, Coastal and Shelf Science*, 95, 110–118, <https://doi.org/10.1016/j.ecss.2011.08.025>, 2011.
- Saha A., Ghosh S., Sahana A. S., and E. P. Rao: Failure of CMIP5 climate models in simulating post-1950 decreasing trend of Indian monsoon, *Geophys. Res. Lett.*, 41:7323–7330. doi.org/10.1002/2014GL061573, 2014.
- Sahu, L. K., Lal, S., and Venkataramani, S.: Seasonality in the latitudinal distributions of NMHCs over the Bay of Bengal, *Atmos. Environ.*, 45, 2356–2366, <https://doi.org/10.1016/j.atmosenv.2011.02.021>, 2011.
- Sahu, L. K., Lal, S., and Venkataramani, S.: Impact of monsoon circulations on oceanic emissions of light alkenes over Bay of Bengal, *Global Biogeochem. Cycles*, 24, GB4028, [doi:10.1029/2009GB003766](https://doi.org/10.1029/2009GB003766), 2010.
- Sahu, L.K., Lal, S., and Venkataramani, S.: Distributions of O₃, CO and hydrocarbons over the Bay of Bengal: a study to assess the role of transport from southern India and marine regions during September–October 2002, *Atmos. Environ.*, 40, 4633–4645, 2006.
- Saiz-Lopez, A., Fernandez, R. P., Ordóñez, C., Kinnison, D. E., Gómez Martín, J. C., Lamarque, J.-F. and Tilmes, S.: Iodine chemistry in the troposphere and its effect on ozone, *Atmos. Chem. Phys.*, 14(23), 13119–13143, [doi:10.5194/acp-14-13119-2014](https://doi.org/10.5194/acp-14-13119-2014), 2014.
- Saiz-Lopez, A., Baidar, S., Cuevas, C. A., Koenig, T. K., Fernandez, R. P., Dix, B., Kinnison, D. E., Lamarque, J., Rodriguez-Lloveras, X., Campos, T. L. and Volkamer, R.: Injection of iodine to the stratosphere, *Geophys. Res. Lett.*, 42, 6852–6859, [doi:10.1002/2015GL064796](https://doi.org/10.1002/2015GL064796), 2015.
- Saji, N. H., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical Indian Ocean, *Nature*, volume 401, 360–363, 1999.
- Sánchez-Pérez, E. D., Marín, I., Nunes, S., Fernández-González, L., Peters, F., Pujo-Pay, M., Conan, P., Marrasé, C.: Aerosol inputs affect the optical signatures of dissolved organic matter in NW Mediterranean coastal waters, *Scientia Marina* 80(4), 437–446, <https://doi.org/10.3989/scimar.04318.20B>, 2016.
- Sardessai, S., Ramaiah, N., Prasanna Kumar, S. and de Sousa, S. N.: Influence of environmental forcings on the seasonality of dissolved oxygen and nutrients in the Bay of Bengal, *J. Marine Res.*, 65, 301–316, 2007.
- Sarma, V. V. S. S., Kumar, N. A., Prasad, V. R., Venkataramana, V., Appalanaidu, S., Sridevi, B., Kumar, B. S. K., Bharati, M. D., Subbaiah, C. V., Acharyya, T., Rao, G. D., Viswanadham, R., Gawade, L., Manjary, D. T., Kumar, P. P., Rajeev, K., Reddy, N. P. C., Sarma, V. V., Kumar, M. D., Sadhuran, Y., and Murty, T. V. R.: High CO₂ emissions from the tropical Godavari estuary (India) associated with monsoon river discharges, *Geophys. Res. Lett.*, 38, [10.1029/2011gl046928](https://doi.org/10.1029/2011gl046928), 2011.
- Sarma, V. V. S. S., Lenton, A., Law, R. M., Metzl, N., Patra, P. K., Doney, S., Lima, I. D., Dlugokencky, E., Ramonet, M., and Valsala, V., (2013), Sea–air CO₂ fluxes in the Indian Ocean

between 1990 and 2009, (2013): *Biogeosciences*, 10, 7035-7052, doi:10.5194/bg-10-7035-2013.

Sarma, V. V. S. S., Swathi, P. S., Kumar, M. D., Prasannakumar, S., Bhattathiri, P. M. A., Madhupratap, M., Ramaswamy, V., Sarin, M. M., Gauns, M., Ramaiah, N., Sardesai, S., and de Sousa, S. N.: Carbon budget in the Eastern and Central Arabian Sea: an Indian JGOFS synthesis, *Global Biogeochem. Cy.*, 17, 1102, doi:10.1029/2002GB001978, 2003.

Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J., Bergamaschi, P., Blake, D. R., Brailsford, G., Brovkin, V., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Curry, C., Frankenberg, C., Gedney, N., Höglund-Isaksson, L., Ishizawa, M., Ito, A., Joos, F., Kim, H. S., Kleinen, T., Krummel, P., Lamarque, J. F., Langenfelds, R., Locatelli, R., Machida, T., Maksyutov, S., McDonald, K. C., Marshall, J., Melton, J. R., Morino, I., Naik, V., O'Doherty, S., Parmentier, F. J. W., Patra, P. K., Peng, C., Peng, S., Peters, G. P., Pison, I., Prigent, C., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J., Spahni, R., Steele, P., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., van Weele, M., van der Werf, G. R., Weiss, R., Wiedinmyer, C., Wilton, D. J., Wiltshire, A., Worthly, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z., and Zhu, Q.: The global methane budget 2000–2012, *Earth Syst. Sci. Data*, 8, 697-751, 10.5194/essd-8-697-2016, 2016.

Savoie, D.L., and Prospero, J.M.: Comparison of oceanic and continental sources of non-sea-salt sulphate over the Pacific Ocean, *Nature*, 339, 685–687, 1989.

Sciare, J., Mihalopoulos, N., Dentener, F. J., and Sciare, L.: Interannual variability of atmospheric dimethylsulfide in the southern Indian Ocean. *J. Geophys. Res.*, 105(D21), 26369–26377, 2000.

Schott, F., and J. P. McCreary, The monsoon circulation of the Indian Ocean, *Prog. Oceanogr.*, 51, 1–123, 2001.

Schott, F. A., Xie, S.-P., and McCreary, J. P., Indian Ocean circulation and climate variability, *Rev. Geophys.*, 47, RG1002, doi:10.1029/2007RG000245, 2009.

Schouten, M. W., W. P. M. de Ruijter, P. J. van Leeuwen, and H. Ridderinkhof: Eddies and variability in the Mozambique Channel, *Deep Sea Res. Part II*, 50, 1987–2003, 2003.

Seinfeld, J. H., and S. N. Pandis: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2nd ed., John Wiley&Sons, 2006.

Selin, N. E., Jacob, D. J., Park, R. J., Yantosca, R. M., Strode, S., Jaeglé, L., and Jaffe, D.: Chemical cycling and deposition of atmospheric mercury: Global constraints from observations, *J. Geophys. Res.*, 112, D02308, doi:10.1029/2006JD007450, 2007.

Sharma, S. K., Singh, A. K., Saud, T., Mandal, T. K., Saxena, M., Singh, S., et al.: Measurement of ambient NH₃ over Bay of Bengal during W-ICARB campaign. *Annales Geophysicae*, 30(2), 371–377, <https://doi.org/10.5194/angeo-30-371-2012>, 2012.

- 1 Shaw, S. L., Chisholm, S. W., and Prinn, R. G.: Isoprene production by *Prochlorococcus*, a
2 marine cyanobacterium, and other phytoplankton, *Mar. Chem.*, 80, 227–245,
3 doi:10.1016/S0304-4203(02)00101-9, 2003.
- 4
- 5 Shechner, M. and Tas, E.: Ozone Formation Induced by the Impact of Reactive Bromine and
6 Iodine Species on Photochemistry in a Polluted Marine Environment, *Environ. Sci. Technol.*,
7 51, 24, 14030–14037, <https://doi.org/10.1021/acs.est.7b02860>, 2017.
- 8
- 9 Siedler, G., M. Rouault, and J. R. E. Lutjeharms: Structure and origin of the subtropical South
10 Indian Ocean Countercurrent, *Geophys. Res. Lett.*, 33, L24609, doi:10.1029/2006GL027399,
11 2006.
- 12
- 13 Singh, A., and Ramesh, R.: Environmental controls on new and primary production in the
14 northern Indian ocean. *Prog. Oceanogr.* 131, 138–145, 2015.
- 15
- 16 Singh D., Ghosh S., Roxy M. K., and McDermid, S.: Indian summer monsoon: extreme events,
17 historical changes, and role of anthropogenic forcings. *Wiley Interdisc. Rev. Clim. Change*,
18 doi. org/10.1002/wcc.571, 2019a.
- 19
- 20 Singh, H.B., Kanakidou, M., Crutzen, P.J., and Jacob, D.J.: High concentrations and
21 photochemical fate of oxygenated hydrocarbons in the global troposphere, *Nature*, 378, 50–54,
22 1995.
- 23
- 24 Singh, K., Panda, J., and Kant, S.: A study on variability in rainfall over India contributed by
25 cyclonic disturbances in warming climate scenario, *Int. J. Climatology*, doi: 10.1002/joc.6392,
26 2019b.
- 27
- 28 Singh, K., Panda, J., Osuri, K.K. and Vissa, N.K.: Progress in tropical cyclone predictability
29 and present status in the North Indian Ocean region. In: Lupo, A.R. (Ed.) *Tropical Cyclone*
30 *Dynamics, Prediction, and Detection*. London: Intech Open, <https://doi.org/10.5772/64333>,
31 2016.
- 32
- 33 Simpson, M. D. and Raman, S.: Role of the land plume in the transport of ozone over the ocean
34 during INDOEX (1999), *Bound. Lay. Meteorol.*, 111, 133–152, 2004.
- 35
- 36 Sprovieri, F., Pirrone, N., Bencardino, M., D'Amore, F., Carbone, F., Cinnirella, S., Mannarino,
37 V., Landis, M., Ebinghaus, R., Weigelt, A., Brunke, E.-G., Labuschagne, C., Martin, L.,
38 Munthe, J., Wängberg, I., Artaxo, P., Morais, F., Barbosa, H. D. M. J., Brito, J., Cairns, W.,
39 Barbante, C., Diéguez, M. D. C., Garcia, P. E., Dommergue, A., Angot, H., Magand, O., Skov,
40 H., Horvat, M., Kotnik, J., Read, K. A., Neves, L. M., Gawlik, B. M., Sena, F., Mashyanov, N.,
41 Obolkin, V., Wip, D., Feng, X. B., Zhang, H., Fu, X., Ramachandran, R., Cossa, D., Knoery,
42 J., Maruschak, N., Nerentorp, M., and Norstrom, C.: Atmospheric mercury concentrations
43 observed at ground-based monitoring sites globally distributed in the framework of the GMOS
44 network, *Atmos. Chem. Phys.*, 16, 11915–11935, <https://doi.org/10.5194/acp-16-11915-2016>,
45 2016.
- 46
- 47 Sreeush M. G., R. Saran, V. Valsala, S. Pentakota, K.V. S.R. Prasad, R. Murtugudde (2019):
48 Variability, trend and controlling factors of Ocean acidification over Western Arabian Sea
49 upwelling region, *Marine Chemistry*, doi.org/10.1016/j.marchem.2018.12.002.
- 50

- 1 Srivastava, S., Lal, S., Venkataramani, S., Gupta, S., and Acharya, Y. B.: Vertical distribution
2 of ozone in the lower troposphere over the Bay of Bengal and the Arabian Sea during ICARB-
3 2006: Effects of continental outflow, *J. Geophys. Res.*, 116, D13301,
4 doi:10.1029/2010JD015298, 2011.
- 5
- 6 Srivastava, S., Lal, S., Venkataramani, S., Gupta, S., and Sheel, V.: Surface distributions of O₃,
7 CO and hydrocarbons over the Bay of Bengal and the Arabian Sea during pre-monsoon season,
8 *Atmospheric Environment*, 47, 459-467, <https://doi.org/10.1016/j.atmosenv.2011.10.023>,
9 2012a.
- 10
- 11 Srivastava, S. Lal, S. Venkataramani, S., Guha, I., and Bala Subrahmanyam, D.: Airborne
12 measurements of O₃, CO, CH₄ and NMHCs over the Bay of Bengal during winter, *Atmos.*
13 *Environ.*, 59, 2012, 597-609, <https://doi.org/10.1016/j.atmosenv.2012.04.054>, 2012b.
- 14
- 15 Stemmler, I., Hense, I., and Quack, B.: Marine sources of bromoform in the global open ocean
16 – global patterns and emissions, *Biogeosciences*, 12, 1967–1981, [https://doi.org/10.5194/bg-](https://doi.org/10.5194/bg-12-1967-2015)
17 12-1967-2015, 2015.
- 18
- 19 Streets, D. G., Hao, J., Wu, Y., Jiang, J., Chan, M., Tian, H., and Feng, X.: Anthropogenic
20 mercury emissions in China, *Atmos. Environ.*, 39, 7789-7806,
21 <https://doi.org/10.1016/j.atmosenv.2005.08.029>, 2005.
- 22
- 23 Stuecker, M. F., Timmermann, A., Jin, F. F., Chikamoto, Y., Zhang, W., Wittenberg, A. T.,
24 Widiasih, E., and Zhao, S., Revisiting ENSO/Indian Ocean Dipole phase relationships.
25 *Geophys. Res. Lett.*, 44, 2481–2492. <https://doi.org/10.1002/2016GL072308>, 2017.
- 26
- 27 Sudheesh, V., Gupta, G. V. M., Sudharma, K. V., Naik, H., Shenoy, D. M., Sudhakar, M., and
28 Naqvi, S. W. A.: Upwelling intensity modulates N₂O concentrations over the western Indian
29 shelf, *J. Geophys. Res.*, 121, 8551-8565, 10.1002/2016jc012166, 2016.
- 30
- 31 Suntharalingam, P., Kettle, A. J., Montzka, S. M., and Jacob, D. J.: Global 3-D model analysis
32 of the seasonal cycle of atmospheric carbonyl sulfide: Implications for terrestrial vegetation
33 uptake, *Geophys. Res. Lett.*, 35, 10.1029/2008gl034332, 2008.
- 34
- 35 Suntharalingam, P., Zamora, L. M., Bange, H. W., Bikkina, S., Buitenhuis, E., Kanakidou, M.,
36 Lamarque, J.-F., Landolfi, A., Resplandy, L., Sarin, M. M., Seitzinger, S., and Singh, A.:
37 Anthropogenic nitrogen inputs and impacts on oceanic N₂O fluxes in the northern Indian
38 Ocean: The need for an integrated observation and modelling approach, *Deep Sea Research*
39 *Part II: Topical Studies in Oceanogr.*, 166, 104-113, <https://doi.org/10.1016/j.dsr2.2019.03.007>,
40 2019.
- 41
- 42 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey,
43 S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in
44 secondary organic aerosol formation from isoprene, *Proc. Natl. Acad. Sci.*, 107, 6640-6645,
45 10.1073/pnas.0911114107, 2010.
- 46
- 47 Tadic, I., Crowley, J. N., Dienhart, D., Eger, P., Harder, H., Hottmann, B., Martinez, M.,
48 Parchatka, U., Paris, J.-D., Pozzer, A., Rohloff, R., Schuladen, J., Shenolikar, J., Tauer, S.,
49 Lelieveld, J., and Fischer, H.: Net ozone production and its relationship to NO_x and VOCs in
50 the marine boundary layer around the Arabian Peninsula, *Atmos. Chem. Phys. Discuss.*,
51 <https://doi.org/10.5194/acp-2019-1031>, in review, 2019.

- 1
- 2 Tegtmeier, S., Krüger, K., Quack, B., Atlas, E. L., Pisso, I., Stohl, A., and Yang, X.: Emission
- 3 and transport of bromocarbons: from the West Pacific ocean into the stratosphere, *Atmos.*
- 4 *Chem. Phys.*, 12, 10633–10648, <https://doi.org/10.5194/acp-12-10633-2012>, 2012.
- 5
- 6 Tegtmeier, S., Atlas, E., Quack, B., Ziska, F., and Krüger, K.: Variability and past long-term
- 7 changes of brominated very short-lived substances at the tropical tropopause, *Atmos. Chem.*
- 8 *Phys.*, 20, 7103–7123, <https://doi.org/10.5194/acp-20-7103-2020>, 2020.
- 9
- 10 Tian, H., et al., Global methane and nitrous oxide emissions from terrestrial ecosystems due to
- 11 multiple environmental changes. *Ecosyst. Health*, 1(1):4. [http://dx.doi.org/10.1890/EHS14-](http://dx.doi.org/10.1890/EHS14-0015.1)
- 12 [0015.1](http://dx.doi.org/10.1890/EHS14-0015.1), 2015.
- 13
- 14 Toihir, Abdoulwahab M. et al. Studies on CO variation and trends over South Africa and the
- 15 Indian Ocean using TES satellite data. *S. Afr. j. sci*, 111, 9-10, 01-09,
- 16 <http://dx.doi.org/10.17159/SAJS.2015/20140174>, 2015.
- 17
- 18 Tomsche, L., Pozzer, A., Ojha, N., Parchatka, U., Lelieveld, J., and Fischer, H.: Upper
- 19 tropospheric CH₄ and CO affected by the South Asian summer monsoon during the Oxidation
- 20 Mechanism Observations mission, *Atmos. Chem. Phys.*, 19, 1915–1939,
- 21 <https://doi.org/10.5194/acp-19-1915-2019>, 2019.
- 22
- 23 Tokarczyk, R. and Moore, R. M.: Production of volatile organohalo- gens by phytoplankton
- 24 cultures, *Geophys. Res. Lett.*, 21, 285– 288, <https://doi.org/10.1029/94GL00009>, 1994.
- 25
- 26 Tournadre, J.: Anthropogenic pressure on the open ocean: The growth of ship traffic revealed
- 27 by altimeter data analysis, *Geophys. Res. Lett.*, 41, 7924–7932, [10.1002/2014gl061786](https://doi.org/10.1002/2014gl061786), 2014.
- 28
- 29 Tripathi, N., Sahu, L. K., Singh, A., Yadav, R. and Karati, K. K.: High Levels of Isoprene in
- 30 the Marine Boundary Layer of the Arabian Sea during Spring Inter-Monsoon: Role of
- 31 Phytoplankton Blooms, *ACS Earth Sp. Chem.*, 4(4), 583–590,
- 32 [doi:10.1021/acsearthspacechem.9b00325](https://doi.org/10.1021/acsearthspacechem.9b00325), 2020.
- 33
- 34 Tripathi, N., Sahu, L. K., Singh, A., Yadav, R., Patel, A., Patel, K., Patel, A., and Patel, K.: Elevated levels of biogenic nonmethane hydrocarbons in the marine boundary layer of the
- 35 Arabian Sea during the intermonsoon, *J. Geophys. Res.: Atmos.*, 124, e2020JD032869.
- 36 <https://doi.org/10.1029/2020JD032869>, 2020.
- 37
- 38
- 39 UN-Environment, 2019. Global Mercury Assessment 2018. UN-Environment Programme,
- 40 Chemicals and Health Branch, Geneva, Switzerland. 59 pp., 2019.
- 41
- 42 Valsala, V., S. Maksyutov, (2013), Interannual variability of the air–sea CO₂ flux in the north
- 43 Indian Ocean, *Ocean Dynamics*, DOI:10.1007/s10236-012-0588-7, 1-14.
- 44
- 45 Valsala, V., S. Maksyutov, and R. G. Murtugudde, (2012), A window for carbon uptake in
- 46 the southern subtropical Indian Ocean, *Geophys. Res. Lett.*, [doi:10.1029/2012GL052857](https://doi.org/10.1029/2012GL052857).
- 47
- 48 Valsala, V., and R. Murtugudde, (2015), Mesoscale and Intraseasonal Air-Sea CO₂ C₂
- 49 Exchanges in the Western Arabian Sea during Boreal Summer, *Deep Sea Research-I*,
- 50 [doi:10.1016/j.dsr.2015.06.001](https://doi.org/10.1016/j.dsr.2015.06.001).
- 51

- 1 Valsala, V., M. G. Sreeush, and K. Chakraborty, (2020), IOD impacts on Indian the Ocean
2 Carbon Cycle, *Journal of Geophysical Research*, <https://doi.org/10.1029/2020JC016485>.
3
- 4 Van Damme, M., Clarisse, L., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C., and
5 Coheur, P.-F.: Industrial and agricultural ammonia point sources exposed, *Nature*, 564, 99–103,
6 10.1038/s41586-018-0747-1, 2018.
7
- 8 Vecchi, G. A., S.-P. Xie, and A. S. Fischer: Ocean-atmosphere co-variability in the western
9 Arabian Sea, *J. Climate*, 17, 1213–1224, 2004.
- 10 Verreyken, B., Amelynck, C., Schoon, N., Müller, J.-F., Brioude, J., Kumps, N., Hermans, C.,
11 Metzger, J.-M., Colomb, A., and Stavrakou, T.: Measurement report: Source apportionment of
12 volatile organic compounds at the remote high-altitude Maïdo observatory, *Atmos. Chem.*
13 *Phys.*, 21, 12965–12988, <https://doi.org/10.5194/acp-21-12965-2021>, 2021.
14
- 15 Verver, G. H. L., Sikka, D. R., Lobert, J. M., Stossmeister, G., and Zachariasse, M.: Overview
16 of the meteorological conditions and atmospheric transport processes during INDOEX 1999, *J.*
17 *Geophys. Res.*, 106, 28399–28413, 2001.
18
- 19 Wai, K. M., Wu, S., Kumar, A., and Liao, H.: Seasonal variability and long-term evolution of
20 tropospheric composition in the tropics and Southern Hemisphere, *Atmos. Chem. Phys.*, 14,
21 4859–4874, <https://doi.org/10.5194/acp-14-4859-2014>, 2014.
22
- 23 Waliser, D. E., Intraseasonal variability. *The Asian Monsoon*, B. Wang, Ed., Springer, 203–
24 258, 2006.
25
- 26 Waliser, D. E., and C. Gautier: A Satellite-derived Climatology of the ITCZ. *J. Climate*, 6,
27 2162–2174, [https://doi.org/10.1175/1520-0442\(1993\)006<2162:ASDCOT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1993)006<2162:ASDCOT>2.0.CO;2), 1993.
28
- 29 Wang, S., Kinnison, D., Montzka, S. A., Apel, E. C., Hornbrook, R. S., Hills, A. J., et al., Ocean
30 biogeochemistry control on the marine emissions of brominated very short-lived ozone-
31 depleting substances: a machine-learning approach. *J. Geophys. Res.*, 124, 12319– 12339.
32 <https://doi.org/10.1029/2019JD031288>, 2019.
33
- 34 Wang, R., Balkanski, Y., Bopp, L., Aumont, O., Boucher, O., Ciais, P., Gehlen, M., Peñuelas,
35 J., Ethé, C., Hauglustaine, D., Li, B., Liu, J., Zhou, F., Tao, S.: Influence of anthropogenic
36 aerosol deposition on the relationship between oceanic productivity and warming, *Geophys.*
37 *Res. Lett.*, 42, 10,745– 10,754, doi:10.1002/2015GL066753, 2015.
38
- 39 Warneck, P.: The relative importance of various pathways for the oxidation of sulfur dioxide
40 and nitrogen dioxide in sunlit continental fair weather clouds, *Phys. Chem. Chem. Phys.*, 1,
41 5471–5483, 1999.
42
- 43 Watts, S. F.: The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and
44 hydrogen sulfide, *Atmos. Environ.*, 34, 761–779, 2000.
45
- 46 Webster, P. J., Moore, A. M., Loschnigg, J. P. and Leben, R. R. Coupled oceanic-atmospheric
47 dynamics in the Indian Ocean during 1997–98. *Nature*, 401, 356–360, 1999.
48
- 49 Williams, J., Custer, T., Riede, H., Sander, R., Jöckel, P., Hoor, P., Pozzer, A., Wong-
50 Zehnpfennig, S., Hosaynali-Beygi, Z., Fischer, H., Gros, V., Colomb, A., Bonsang, B., Yassaa,
51 N., Peeken, I., Atlas, E.L., Waluda, C.M., van Aardenne, J.A., Lelieveld, J.: Assessing the effect

1 of marine isoprene and ship emissions on ozone, using modeling and measurements from the
2 South Atlantic Ocean, *Environ. Chem.*, 7, 171–182, doi: [10.1071/EN09154](https://doi.org/10.1071/EN09154), 2010.

3
4 Williams, J., Fischer, H., Wong, S., Crutzen, P. J., Scheele, M. P., and Lelieveld, J.: Near
5 equatorial CO and O₃ profiles over the Indian Ocean during the winter monsoon: High O₃ levels
6 in the middle troposphere and interhemispheric exchange, *J. Geophys. Res.*, 107, 8007,
7 doi:10.1029/2001JD001126, 2002.

8 Wurl, O., Wurl, E., Miller, L., Johnson, K., and Vagle, S.: Formation and global distribution of
9 sea-surface microlayers, *Biogeosciences*, 8, 121–135, doi:10.5194/bg-8-121-2011, 2011.

10
11 Yamamoto, H., Yokouchi, Y., Otsuki, A., and Itoh, H.: Depth profiles of volatile halogenated
12 hydrocarbons in seawater in the Bay of Bengal, *Chemosphere*, 45, 371–377,
13 [https://doi.org/10.1016/S0045-6535\(00\)00541-5](https://doi.org/10.1016/S0045-6535(00)00541-5), 2001.

14
15 Yang, J., Zhao, W., Wei, L., Zhang, Q., Zhao, Y., Hu, W., Wu, L., Li, X., Pavuluri, C. M., Pan,
16 X., Sun, Y., Wang, Z., Liu, C.-Q., Kawamura, K., and Fu, P.: Molecular and spatial
17 distributions of dicarboxylic acids, oxocarboxylic acids, and α -dicarbonyls in marine aerosols
18 from the South China Sea to the eastern Indian Ocean, *Atmos. Chem. Phys.*, 20, 6841–6860,
19 <https://doi.org/10.5194/acp-20-6841-2020>, 2020.

20
21 Yao, S.L., G. Huang, R.-G. Wu, X. Qu, and D. Chen: Inhomogeneous warming of the tropical
22 Indian Ocean in the CMIP5 model simulations during 1900–2005 and associated mechanisms,
23 *Clim. Dynam.*, 2016.

24
25 Ye, H., Sheng, J., Tang, D., Morozov, E., Kalhor, M. A., Wang, S., and Xu, H.: Examining
26 the Impact of Tropical Cyclones on Air-Sea CO₂ Exchanges in the Bay of Bengal Based on
27 Satellite Data and In Situ Observations, *J. Geophys. Res.*, 124, 555–576,
28 [10.1029/2018jc014533](https://doi.org/10.1029/2018jc014533), 2019.

29
30 Yoder, J. A., McClain, C. R., Feldman, G. C., and Esaias, W. E.: Annual cycles of
31 phytoplankton chlorophyll concentrations in the global ocean: A satellite view, *Global*
32 *Biogeochem. Cy.*, 7, 181–193, [10.1029/93gb02358](https://doi.org/10.1029/93gb02358), 1993.

33
34 Yoshida, Y., Ota, Y., Eguchi, N., Kikuchi, N., Nobuta, K., Tran, H., Morino, I., and Yokota,
35 T.: Retrieval algorithm for CO₂ and CH₄ column abundances from short-wavelength infrared
36 spectral observations by the Greenhouse gases observing satellite, *Atmos. Meas. Tech.*, 4, 717–
37 734, doi:10.5194/amt-4-717-2011, 2011.

38
39 ~~Yu, L.: Global variations in oceanic evaporation (1958–2005): the role of the changing wind~~
40 ~~speed. *J. Climate*, 20:5376–5390, 2007.~~

41
42 Zavarsky, A., Goddijn-Murphy, L., Steinhoff, T., and Marandino, C. A.: Bubble-Mediated Gas
43 Transfer and Gas Transfer Suppression of DMS and CO₂, *J. Geophys. Res.*, 123, 6624–6647,
44 [10.1029/2017jd028071](https://doi.org/10.1029/2017jd028071), 2018a.

45
46 Zavarsky, A., Booge, D., Fiehn, A., Krüger, K., Atlas, E., and Marandino, C.: The Influence of
47 Air-Sea Fluxes on Atmospheric Aerosols During the Summer Monsoon Over the Tropical
48 Indian Ocean, *Geophys. Res. Lett.*, 45, 418–426, [10.1002/2017gl076410](https://doi.org/10.1002/2017gl076410), 2018b.

- 1 Zhang, N., Feng, M., Du, Y., Lan, J., Wijffels, S.E.: Seasonal and interannual variations of
2 mixed layer salinity in the southeast tropical Indian Ocean. *J. Geophys. Res.*, 121(7):4716–
3 4731, 2016.
- 4
- 5 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan,
6 L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic
7 emissions since 2010 as the consequence of clean air actions, *Atmos. Chem. Phys.*, 18, 14095–
8 14111, <https://doi.org/10.5194/acp-18-14095-2018>, 2018.
- 9
- 10 Zhou, S., Gonzalez, L., Leithead, A., Finewax, Z., Thalman, R., Vlasenko, A., Vagle, S., Miller,
11 L. A., Li, S.-M., Burekul, S., Furutani, H., Uematsu, M., Volkamer, R., and Abbatt, J.:
12 Formation of gas-phase carbonyls from heterogeneous oxidation of polyunsaturated fatty acids
13 at the air–water interface and of the sea surface microlayer, *Atmos. Chem. Phys.*, 14, 1371–
14 1384, <https://doi.org/10.5194/acp-14-1371-2014>, 2014.
- 15
- 16 Zhou, M., Langerock, B., Vigouroux, C., Sha, M. K., Ramonet, M., Delmotte, M., Mahieu, E.,
17 Bader, W., Hermans, C., Kumps, N., Metzger, J.-M., Duflot, V., Wang, Z., Palm, M., and De
18 Mazière, M.: Atmospheric CO and CH₄ time series and seasonal variations on Reunion Island
19 from ground-based in situ and FTIR (NDACC and TCCON) measurements, *Atmos. Chem.*
20 *Phys.*, 18, 13881–13901, <https://doi.org/10.5194/acp-18-13881-2018>, 2018.
- 21
- 22 Ziska, F., Quack, B., Abrahamsson, K., Archer, S. D., Atlas, E., Bell, T., Butler, J. H.,
23 Carpenter, L. J., Jones, C. E., Harris, N. R. P., Hepach, H., Heumann, K. G., Hughes, C., Kuss,
24 J., Krüger, K., Liss, P., Moore, R. M., Orlikowska, A., Raimund, S., Reeves, C. E.,
25 Reifenhäuser, W., Robinson, A. D., Schall, C., Tanhua, T., Tegtmeier, S., Turner, S., Wang, L.,
26 Wallace, D., Williams, J., Yamamoto, H., Yvon-Lewis, S., and Yokouchi, Y.: Global sea-to-
27 air flux climatology for bromoform, dibromomethane and methyl iodide, *Atmos. Chem. Phys.*,
28 13, 8915–8934, <https://doi.org/10.5194/acp-13-8915-2013>, 2013.
- 29
- 30 Zou, L.W., Zhou, T.J., Near future (2016–2040) summer precipitation changes over China as
31 projected by a regional climate model (RCM) under the RCP8.5 emissions scenario:
32 comparison between RCM downscaling and the driving GCM. *Adv. Atmos. Sci.*, 30:806–818,
33 2013.
- 34
- 35
- 36
- 37

Supplement

A) Atmospheric intraseasonal and interannual variability

Intraseasonal variability can impact atmospheric transport patterns over the Indian Ocean with the dominant mode being the eastward propagating Madden-Julian Oscillation (MJO). Equatorially trapped, baroclinic oscillations in the tropical wind field propagate slowly eastward across the Indian Ocean, Maritime Continent, and West Pacific with an intraseasonal cycle of 30–60 days (Madden and Julian, 1972). A typical MJO event exhibits large-scale convection anomalies where enhanced convection and rainfall develop over the western Indian Ocean with suppressed convection further east over the western Pacific (Zhang, 2005). The eastward propagation of the convection and circulation anomalies depends on the season and is strongest during the winter monsoon. The summer monsoon shows a north-eastward propagation of the anomalies into Southeast Asia in addition to the eastward propagation along the equator (Waliser, 2006). Among the many impacts of the MJO, strong interactions with ocean surface fluxes of mass, heat, and momentum have been observed (e.g., Matthews et al., 2010).

Similarly, modes of tropical interannual variability, such as the irregular oscillation of sea surface temperatures known as the Indian Ocean Dipole (IOD), play a role for Indian Ocean meteorology. The positive phase of the IOD is characterised by positive sea surface temperature anomalies in the western part of the Indian Ocean accompanied by negative anomalies in eastern part (Saji et al., 1999). The initial cooling off the coast of Sumatra–Java leads to a positive feedback mechanism via suppressed local convection, anomalous easterly winds, a shoaling thermocline and stronger upwelling, which in turn reinforce the initial cooling with a peak from September to November (Cai et al., 2014). Extreme positive IOD events can also impact the equatorial ocean by inducing a north-westward extension of the south-easterly trades and drying along the equatorial Indian Ocean (Webster et al., 1999).

The dominant mode of interannual climate variability of the Pacific, the El Niño–Southern Oscillation (ENSO), can also impact Indian Ocean sea surface temperatures via anomalous wind stress forcing (Latif and Barnett, 1995). In addition, ENSO modulates the depth of the Indian Ocean thermocline and contributes to changes in salinity due to shifts in rainfall and evaporation. The potential impact of ENSO on the IOD is currently under discussion (Stuecker et al., 2017 and references therein).

B.) Regional Oceanic Transport

Below we will outline the major features of the equatorial and southern Indian Ocean (Fig. S1) and describe the influence of the seasonal monsoon on the physical processes in the northern Indian Ocean (Fig. S1).

Southern Indian Ocean

The South Equatorial Current (SEC) in the Indian Ocean carries water masses entering through the Indonesian passages, with a relative salinity minimum in the Indian Ocean environment, via broad zonal inflow westward. Driven by the southeast trades, the SEC supplies the western boundary currents east of Madagascar. Part of the throughflow water in the SEC forms the northern East Madagascar Current (EMC), which partly passes in southward moving eddies through the Mozambique Channel and merges into the Agulhas (Ridderinkhof and de Ruijter, 2003; Schouten et al., 2003).

Data from satellite altimetry suggest that the eastward South Indian Countercurrent (SICC) is already present in the Mozambique Basin, southwest of Madagascar (Siedler et al., 2006). The eastward frontal SICC coincides with the thermohaline front that separates the saline

1 subtropical surface water from the fresher tropical surface water in the EMC in summer
2 (Palastanga et al., 2007). The variability of SST and salinity is high in the warm waters south
3 to south-east of Madagascar, likely due to eddy activity and upwelling. There is year-round
4 coastal upwelling along the southern stretch of the oligotrophic EMC and in the shallower
5 region just to the south of Madagascar, which leads to enhanced phytoplankton growth (Quartly
6 et al., 2006). All productivity further from the island in bands of relatively high variability along
7 25°S and along the EMC are due to a combination of local upwelling caused by eddies and,
8 more importantly, the advection of upwelled coastal waters around eddy features. The
9 anticyclones moving through the region wrap both the warm EMC waters and the nutrient-rich
10 upwelled waters into well-defined arcs. Occasionally strands of chlorophyll-rich water can
11 stretch 500 km or more eastward, which are caused by the combined effects of both cyclonic
12 and anticyclonic eddies (Quartly et al., 2006). The southward flowing EMC, as part of the
13 bifurcated SEC, and the SEC form the western and northern boundary currents of the South
14 Indian subtropical gyre, where saline surface water is formed, as there is more evaporation than
15 precipitation in this region (Schott et al., 2009). The south-eastern Indian Ocean shows the
16 strongest interannual to decadal variability of upper-ocean salinity in the Indian Ocean.
17 Seasonality of the mixed layer salinity in the south-eastern tropical Indian Ocean is influenced
18 by the annual cycles of the Indonesian Throughflow and the Leeuwin Current transports,
19 freshwater forcing, and eddy fluxes (Zhang et al., 2016).

20 Open ocean upwelling associated with the Seychelles dome, a thermocline ridge in the southern
21 tropical gyre can occur between 5 and 10°S, along the northern edge of the southeast trades,
22 where Ekman divergence occasionally appears to be strong enough to upwell subsurface waters
23 into the mixed layer (Schott et al., 2009). In this region, the South Equatorial Countercurrent
24 (SECC) flows eastward year-round, fed by the East African Coastal Current (EACC) and
25 forming the northern flank of the southern Indian Ocean tropical gyre. Low sea surface height
26 is the signature of the Indian Ocean's tropical gyre, bounded in the north by the SECC, to the
27 south by the SEC, and at the western boundary by the EACC.





Figure S1. Schematic representation of identified oceanic currents during the summer monsoon (a) and winter monsoon (b). Current branches indicated are the South Equatorial Current (SEC), South Equatorial Countercurrent (SECC), Northeast and Southeast Madagascar Current (NEMC and SEMC), East African Coastal Current (EACC), Somali Current (SC), Southern Gyre (SG) and Great Whirl (GW) and associated upwelling wedges (green shades), Southwest and Northeast Monsoon Currents (SMC and NMC), South Java Current (SJC), East Gyrar Current (EGC), and Leeuwin Current (LC). The subsurface return flow of the supergyre is shown in magenta. Depth contours shown are for 1000 m and 3000 m (grey). Updated representations are from SMC01; red vectors (Me) show directions of meridional Ekman transports. ITF indicates Indonesian Throughflow (from Schott et al., 2009, copyright 2009 by the American Geophysical Union, reproduced with permission).

Northern Indian Ocean

During the summer monsoon, the northward flowing Somali Current is supplied by the SEC and EACC. Once the Somali Current crosses the equator, a part of it turns offshore around 4°N and another part returns across the equator as a part of the southern gyre. A northern gyre occurs north of the Equator and occasionally a third gyre can be observed in many summer monsoons (Schott et al., 2009). These gyres influence the stability of the atmospheric planetary boundary layer, impacting surface wind stress and heat fluxes (Vecchi et al., 2004). Furthermore, the Southwest Monsoon Current flows towards the east, south of Sri Lanka, then turns to flow toward the north, bringing saltier Arabian Sea water into the Bay of Bengal (Jensen, 2003). In contrast, the Somali Current flows southward during the winter monsoon to meet the northward flowing EACC. They supply water for the eastward flowing SECC. The Northeast Monsoon Current flows toward the west, bringing fresher Bay of Bengal water into the Arabian Sea (Schott et al., 2009). In addition, model studies have suggested that Bay of Bengal water flows across the equator in the eastern basin in the winter monsoon (Han and McCreary, 2001; Jensen, 2003).

Unique to the Indian Ocean are strong eastward ocean surface jets during the inter-monsoon period, called Wyrtki Jets. They are produced by the semi-annual westerly equatorial winds and are important because they carry warm upper layer waters toward the east, which increases sea level and mixed layer depth in the east and decreases them in the west. These semi-annual westerlies are the reason for another unique Indian Ocean feature, namely the eastward Equatorial Undercurrent that is only present for a part of the year (i.e. February – June) when the winds have an easterly component (Schott et al., 2009; Reppin et al., 1999).