

Supplementary Information:

“Inverse modeling of fire emissions constrained by smoke plume transport using HYSPLIT dispersion model and geostationary observations” by Kim et al.

Hyun Cheol Kim^{1,2}, Tianfeng Chai^{1,2}, Ariel Stein¹ and Shobha Kondragunta³

5 ¹ Air Resources Laboratory, National Oceanic and Atmospheric Administration, College Park, MD, 20740, MD, USA

² Cooperative Institute for Satellite Earth System Studies, University of Maryland, College Park, MD, 20740, USA

³ National Environmental Satellite, Data and Information Service, National Oceanic and Atmospheric Administration, College Park, MD 20740, USA

Correspondence to: Tianfeng Chai (tianfeng.chai@noaa.gov), Hyun Cheol Kim (hyun.kim@noaa.gov)

10 **Abstract.** We provide supplementary information and descriptions of twin experiments.

List of Figures

- Figure S1 Two steps for smoke forecast with the HEIMS-fire system. Temporal coverage of observational data (oday=0,-1,2,-3) and forecast days (fdays=0,+1,+2) for operational tests. See context for the detail.2
- 15 Figure S2 Daily evolution of wildfire events over southeastern US during November 8-19, 2016.3
- Figure S3 Twin experiments setup. Constant smoke releases at 9 fire locations (left) for two days from 6Z on August 17, 2015, at 1500m or 2000m are simulated by HYSPLIT. Hourly pseudo-observations of satellite mass loadings are generated based on the HYSPLIT results (right). Regions A,B,C are used for sensitivity tests with spatial coverage in twin experiments cases 4-6.5
- 20 Figure S4 Test of fire emission vertical allocation. Comparison of the estimated smoke emission rates of Case 1 in Table 2 (left) at the nine locations (fire 1-9 as x-axis) for the two days (upper: 1st day; lower: 2nd day) and the “actual” sources (right) used in the twin experiment.5

List of tables

- 25 Table S1 List of twin experiments configurations.6
- Table S2. Smoke source term error statistics of the twin experiments. MAE: mean absolute error; NMAE: normalized MAE; RMSE: root-mean-square error; NRMSE: normalized RMSE.6




Figure S1 Two steps for smoke forecast with the HEIMS-fire system. Temporal coverage of observational data (oday=0,-1,2,-3) and forecast days (fdays=0,+1,+2) for operational tests. See context for the detail.




Figure S2 Daily evolution of wildfire events observed from MODIS truecolor images over southeastern US during November 8-19, 2016.

Twin experiments

40 A series of twin experiments have been conducted to test the range of uncertainties that come from the design of fire emission inverse modeling system. A twin experiment denotes an idealized modeling test in which we assume a modeled world mimics well the real world. In the twin experiment, we have true solution for the situation, so we can test the system's capability to reproduce the true answer. We first tested uncertainties of the system using multiple scenarios.

The HYSPLIT inverse system is further extended for the wildfire smoke problems which often have multiple sources
45 locations. The feasibility of applying the method to objectively and optimally estimate wildfire smoke sources based on satellite observations of fire plumes has been demonstrated using a set of twin experiment.



Figure S3 demonstrates the domain coverage of the twin experiments. In these experiments, hourly pseudo-observations are generated using known sources from given fire locations. In this test, we have constant smoke releases at 9 fire location for two days from 6Z on August 2017. Hourly pseudo-observations of satellite mass loadings are assumed based on the
50 HYSPLIT results. All 48 hourly non-zero mass loadings are assumed retrieved accurately.

Four types of potential uncertainties (e.g. vertical allocation, temporal coverage, spatial coverage and impact of observations errors) are test through ten cases. Table S1 summaries the list of twin experiments configurations.

Vertical allocation. We have assigned a fixed fire emissions, 10^5kg/hr , for first guess input for nine locations at five altitudes. Figure S4 shows the results of the twin experiment to test vertical allocation of the system. X-axis denote the
55 indices of nine fire locations, and y-axis denote five altitudes at which fire emissions are assigned and dispersion models are conducted. In most of cases, the system was able to reproduce the location and height of fire emission when compared with the "true" fire emission information.


Temporal coverage. Cases 4-6 were conducted to test the sensitivity from temporal availability of observational data. Case 2 assumes the observations are available for 48 hours while case 3 assumes that the observations are available for 24 hours.
60 When 48-hour observational data available, the system could reproduce the day 1 fire emission with very high accuracy (i.e. NMAE=0.77% and NRMSE=1.21%). The performance of the system drops quickly for the day 2 prediction when only less observational data (e.g. 24-hour) is available.

Spatial coverage. As clouds may block the views of satellites, the effect of missing observations due to clouds is investigated. With all 48 hourly column observations available, the smoke emissions at all 9 locations as a function of day
65 and height are recovered extremely well. Table S2 also show that missing satellite observations in key regions (Region B in Case 4) could dramatically affect the emission estimations.



70

Figure S3 Twin experiments setup. Constant smoke releases at 9 fire locations (left) for two days from 6Z on August 17, 2015, at 1500m or 2000m are simulated by HYSPLIT. Hourly pseudo-observations of satellite mass loadings are generated based on the HYSPLIT results (right). Regions A,B,C are used for sensitivity tests with spatial coverage in twin experiments cases 4-6.



75

Figure S4 Test of fire emission vertical allocation. Comparison of the estimated smoke emission rates of Case 1 in Table 2 (left) at the nine locations (fire 1-9 as x-axis) for the two days (upper: 1st day; lower: 2nd day) and the “actual” sources (right) used in the twin experiment.

80

Table S1 List of twin experiments configurations.

| Test | Case | Descriptions |
|---------------------|---------|--|
| Vertical allocation | Case 1 | Even vertical allocation |
| Temporal coverage | Case 2 | Observations available at all 48 hours |
| | Case 3 | Observations available for last 24 hours |
| Spatial coverage | Case 4 | 24-hour observations with region A blocked |
| | Case 5 | 24-hour observations with region B blocked |
| | Case 6 | 24-hour observations with region C blocked |
| Observational error | Case 7 | 10% observational error |
| | Case 8 | 20% observational error |
| | Case 9 | 50% observational error |
| | Case 10 | 100% observational error |

Table S2. Smoke source term error statistics of the twin experiments. MAE: mean absolute error; NMAE: normalized MAE; RMSE: root-mean-square error; NRMSE: normalized RMSE.

| Cases | Source term | MAE(kg/hr) | NMAE | RMSE(kg/hr) | NRMSE |
|---------------------------|-------------|------------|--------|-------------|---------|
| Case 2 (48-h obs) | Day 1 | 534.9 | 0.77% | 841.4 | 1.21% |
| | Day 2 | 1760.5 | 2.53% | 3332.5 | 4.78% |
| Case 3 (24-h obs) | Day 1 | 1985.8 | 2.85% | 3310.2 | 4.75% |
| | Day 2 | 1393.0 | 2.00% | 2943.2 | 4.22% |
| Case 4 (region A blocked) | Day 1 | 606.4 | 0.87% | 1156.3 | 1.66% |
| | Day 2 | 301.2 | 0.43% | 573.4 | 0.82% |
| Case 5 (region B blocked) | Day 1 | 23834.6 | 34.21% | 32157.9 | 46.16% |
| | Day 2 | 66177.5 | 94.99% | 78653.3 | 112.90% |
| Case 6 (region C blocked) | Day 1 | 3974.9 | 5.71% | 8803.3 | 12.64% |
| | Day 2 | 3400.6 | 4.88% | 10663.2 | 15.31% |
| Case 7 (10% error) | Day 1 | 1448.6 | 2.08% | 2259.7 | 3.24% |
| | Day 2 | 4418.7 | 6.34% | 11121.8 | 15.96% |
| Case 8 (20% error) | Day 1 | 2105.8 | 3.02% | 3954.7 | 5.68% |
| | Day 2 | 8567.9 | 12.30% | 22884.4 | 32.84% |
| Case 9 (50% error) | Day 1 | 6227.6 | 8.94% | 12047.1 | 17.29% |
| | Day 2 | 22034.5 | 31.63% | 67298.2 | 96.60% |
| Case 10 (100% error) | Day 1 | 10104.2 | 14.50% | 19759.9 | 28.36% |
| | Day 2 | 34560.9 | 49.61% | 131203.9 | 188.33% |