

~~Stratospheric pollution~~ MLS observations of biomass-burning products in the stratosphere from Canadian forest fires in August 2017

Hugh C. Pumphrey¹, Michael J. Schwartz², Michelle L. Santee², George P. Kablick III³, Michael D. Fromm³, and Nathaniel J. Livesey²

¹School of GeoSciences, The University of Edinburgh, Edinburgh, UK

²NASA Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

³US Naval Research Lab, Washington DC, USA

Correspondence: Hugh Pumphrey (Hugh.Pumphrey@ed.ac.uk)

Abstract.

Forest fires in British Columbia in August 2017 caused a pyrocumulonimbus event that injected a polluted air mass into the lower stratosphere. The Microwave Limb Sounder (MLS) on the Aura satellite first observed the polluted air mass on 14 August 2017 and continued to observe it for 60 days (100 days in water vapour). We estimate the mass of CO injected into the stratosphere to be 2–32400 TgGg. Events such as this are rare: this is the third of four such events in the 16 years since the launch of Aura, the second-largest of the four events, and the only one in the Northern Hemisphere. Unlike the preceding two events, but like the most recent event, the polluted air mass described here had an unusually high water vapour content. We describe the evolution of the polluted air mass, showing that it rose to an altitude of about 24 km (31 hPa) and divided into several identifiable parts. In addition to CO and H₂O, we observe enhanced amounts of HCN, CH₃CN, CH₃Cl and CH₃OH; these are in roughly the ratios to CO reported elsewhere. We use back-trajectories and plume-dispersion modelling to demonstrate that the pollutants observed by MLS originated in the British Columbia fires.

Copyright statement. Copyright ©statement goes here.

1 Introduction

Fires are an important natural component of many ecosystems, and of forests in particular (Bowman et al., 2009). The nature and frequency of fires can be sensitive to human intervention, even when that intervention is not the direct cause of the fires (Williams and Abatzoglou, 2016). The occurrence of fires is also sensitive to climate change; we can expect fires to become more frequent and/or more damaging in many parts of the world as the climate continues to warm (Williams et al., 2019).

The Microwave Limb Sounder (MLS) instrument (Waters et al., 2006) observes several chemical species emitted by forest fires. In tropical regions these tend to be transported to altitudes observable by this instrument by the wide-scale circulation

20 of the atmosphere, and particularly by the strong convection along the intertropical convergence zone (Pumphrey et al., 2018).
Outside of the tropics it is rare for MLS to observe the immediate effect of fires; we describe one such event in this paper.

The summer of 2017 saw the most destructive forest fires on record in British Columbia, Canada (Government of British Columbia, 2018). On 12 August 2017 a particularly intense cluster of fires gave rise to a strong convective event that lofted biomass-burning product gases to an altitude of 10–11 km (215 hPa). ~~The fires are described in some~~, as described in detail by
25 Peterson et al. (2018), who refer to them as the “Pacific Northwest Event”; ~~we~~. We adopt this name and abbreviate it to PNE. The polluted airmass was transported around the world, ascending to an altitude of 2019 km (4668 hPa) over a period of 18 days ~~. It remained observable for about 100 days, reaching a final altitude of 25 km (26 hPa)~~ (Khaykin et al., 2018). Lidar observations of extremely high levels of aerosol in the polluted airmass have been reported from Europe (Ansmann et al., 2018; Khaykin et al., 2018) (Ansmann et al., 2018; Khaykin et al., 2018; Hu et al., 2019; Baars et al., 2019) and Russia (Zuev et al., 2019). Khaykin et al.
30 (2018) describe the PNE in detail and also report space-based lidar data from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) instrument (Winker et al., 2009), showing the global evolution of the plume. Further space-based observations of aerosol from the event are described by Torres et al. (2020) and Kloss et al. (2019). Lestrelin et al. (2021) combine satellite aerosol measurements and ERA5 (Hersbach et al., 2020) reanalysis data to provide the most detailed description to date of the evolution in time and space of the polluted airmass.

35 MLS on Aura, and its predecessor on the UARS mission, have previously observed biomass burning products from intense fire events (Pumphrey et al., 2011; Livesey et al., 2004). We report here observations made by MLS of some of the gas-phase components of the polluted airmass ~~made by the Microwave Limb Sounder (MLS)~~ from the PNE. The ozone and water vapour data have been reported previously (Yu et al., 2019); here we report in more detail on the full suite of biomass-burning products observed by MLS: CO, CH₃CN, HCN, CH₃Cl and CH₃OH. We also describe any changes observed in the polluted airmass
40 for all other species observed by the instrument.

2 ~~Data from MLS~~

2 Data from MLS

The MLS instrument (Waters et al., 2006) was launched in July 2004 on NASA’s Aura satellite and has operated almost continuously since then. Aura is in a polar orbit with a 98° inclination angle. The MLS instrument views in the direction of
45 travel; as a result the measurement locations lie close to the orbit track and cover the latitude range 82° S to 82° N. Once processed, the data consist of 3495 vertical profiles per day, spaced at intervals of 167 km along the orbit track. Estimated quantities include temperature, geopotential height, and the mixing ratios of 16 chemical species, including H₂O (Lambert et al., 2007), CO (Livesey et al., 2008; Pumphrey et al., 2007) and HCN (Pumphrey et al., 2018). Data are reported on pressure levels; for most products these are spaced at 6 levels per pressure decade, giving a representation with a vertical resolution of
50 about 2.7 km. For most products the true vertical resolution is similar to this spacing, but for species with low mixing ratios and/or weak spectral lines it may be considerably poorer than this. For a few species (H₂O, O₃) the data are reported at a

resolution of 12 levels per pressure decade. ~~Full~~ We use version 4 of the MLS data; full details on use of the data may be found in Livesey et al. (2020).

3 ~~Observations~~ Results

55 3.1 ~~Carbon Monoxide~~ Observations

3.1.1 Carbon Monoxide

MLS CO data are recommended (Livesey et al., 2020) for use at pressure levels from 215 hPa (~ 11 km) up to 0.0046 hPa (~ 80 km). Data are reported at the 316 hPa level (~ 8 km), but these values are thought to mostly represent the real atmosphere at 215 hPa and are not recommended for ~~any~~ quantitative work. They can, to some extent, be a useful qualitative indicator of
60 excess CO somewhere in the ~~8-10~~ 8-10 km altitude range; we show the 316 hPa data in some of the figures in this paper, but do not use them to derive any quantitative results.




Figure 1. MLS measurements of CO (in ppbv) at 215 hPa (≈ 11 km altitude) on ~~14-15~~ 14-15 August 2017. These are the first two days after the PNE on which MLS observed the enhanced CO values.

Enhanced values of CO from the Pacific Northwest Event were first observed on 14 August 2017. Figure 1 shows MLS CO over North America at this time, at 215 hPa. The maximum values seen here are about 1.2 ppmv; this is about 24 times the typical zonal mean value of 50 ppbv and about 6 to 8 times the typical daily maximum value for this latitude band, for times
65 immediately before the PNE. Figure 2 shows a time series of the maximum CO values observed at several pressure levels. It is clear that the polluted airmass reached 215 hPa almost immediately. From there, it ascended fairly rapidly to 100 hPa and then more slowly from 100 hPa to 46 hPa.




Figure 2. Time series of maximum CO values observed polewards of $25\text{--}27.5^\circ$ N. The values shown are averages of the ten highest observed values on each day between $15\text{--}27.5^\circ$ N and 83° N. A 3-day smoothing has been applied to each curve.




Figure 3. Map showing the locations of MLS observations of enhanced CO (defined as described in the text). Colours show time in days after 12 August 2017. Symbols show pressure level in hPa. The MLS measurements lie along the satellite orbit tracks; the tracks are particularly noticeable over North America and can also be seen in Fig. 1.

To summarise the whole event, we applied the strategy used by Pumphrey et al. (2011) to detect values that exceed the usual spread expected for a given latitude band. We found the mean value and standard deviation for all measurements in a given latitude band and at a given pressure level. We then describe the CO at a point as “enhanced” if its mixing ratio was more than a chosen number, α , of standard deviations above the mean. The value chosen for α is somewhat arbitrary and was different for each level: $\alpha = 5.7$ at 215 hPa, $\alpha = 4.6$ at 147 and 100 hPa and $\alpha = 3.9$ at all other pressure levels. These choices are made in order to show the interesting events while rejecting almost all of the background points. An iterative approach is used to ensure that the mean and standard deviation are calculated from the unaffected points; there are few enough affected points that this calculation converges very quickly.

Figure 3 shows the locations where enhanced CO was detected. The plume is first observed over northwestern Canada, travelling southeastwards and then eastwards across northern Canada and then across the Atlantic to Europe. The behaviour of the plume while over Eurasia is more complex. After 30 days, one part of it reaches northwestern China, where it **divides** appears to divide into two parts, one heading east across the Pacific at about 47° N and the other returning westwards across the Atlantic at about 33° N.

As the polluted airmass moves far more rapidly in the zonal than the meridional direction, it is helpful to plot it against time and longitude, as in Fig. 4. The ascent of the polluted airmass (shown by the changing colour and shape of the points) is




Figure 4. Longitudes of enhanced CO (as defined in the text), plotted as a function of time. Colours and symbols represent pressure. Dashed lines and labels identify the various parts into which the polluted airmass is observed to divide.

clearly visible in this figure. It can also be seen in Fig. 4 that the polluted airmass is divided into two parts on three occasions: at approximately 9 days, 16 days and 41 days after 12 August 2017. We label the parts in Fig. 4 using the nomenclature introduced by Lestrelin et al. (2021), with the exception of a part which Lestrelin et al. (2021) do not identify and which we label “X”. After the first division, **one branch part X** remains at 147 to 100 hPa, travels more rapidly than the other **branch parts** and is last seen on day 16 near 80° E, 70° N. The higher **branch part** is at 100 to 68 hPa at day 9 and continues to ascend. It appears to divide into two parts (**A and B1**) somewhere around day 16; these two parts appear in Fig. 4 to cross over at around day 30, after which **one part travels east and one west. The more northern part part B1 continues eastward and part A travels westward. The two parts are at different latitudes and altitudes at this time. Part B1 lies poleward of 42°N, between 68 and 46 hPa; it travels eastward from northern China across the Pacific Ocean, while the more southern part rises to 31.6. Part A is**

equatorward of 38°N, between 46 and 31 hPa and (21 to 24 km); it moves westward across northern Africa and the Atlantic Ocean.

95 Around day 41, the eastward-travelling part part B1 appears to divide into further parts two further parts while at a longitude near 180°E. The MLS data are too sparse to determine the location where the split occurred with any accuracy. The faster-travelling part (B2) is last seen on day 60 at 90° E; the slower-part remainder (which we continue to label B1) is last seen on day 63 at 60° W. Although the MLS CO data do not make this clear, Lestrelin et al. (2021) show that B2 actually splits apart from B1 within a few days of day 25, with B1 and B2 travelling at similar longitudes until a time near day 41.




Figure 5. (a) Total mass of CO observed by MLS poleward of 27.5° N and at altitudes between 215 hPa and 31 hPa. The four times t_1, t_0, t_e and t_2 are discussed in the text. The seasonal fit includes both annual and semiannual components; the annual component is much smaller in this case. (b) As (a), but with the seasonal fit subtracted from the data, and an exponential fitted to the measurements made after the fire. The vertical dashed orange line marks 16 August 2017: Both the time of exponential fit and the earliest data used to construct it are shown in the exponential fit blue.

100 We can obtain an estimate of the total injected mass of CO by integrating the MLS data into partial column (215 hPa to 31 hPa) values and summing over all latitudes affected by the event, weighting each latitude band by its area. The seasonal behaviour of CO is fitted with a mean value and annual and semiannual sinusoids as shown in Fig. 5(a); this fit ignores the data for a period of 9993 days after the PNE. Subtracting this first fit from the data leaves a fairly featureless time series apart from

Table 1. Summary of the injected masses of CO due to the four fire events noted in the text, showing the best estimate of the total mass, with the pressure level in the “Max p ” column being the highest pressure (lowest altitude) level included. Contributions to the error in the mass estimate caused by the choice of the parameters t_0 , t_e , t_1 and t_2 are shown. Also shown is a typical error as returned by the R function `nls()`. The total error is obtained by summing the five individual errors in quadrature. This procedure is not ideal as most of the errors will be correlated with each other. However, it gives a reasonable conservative estimate. Units are Gg for all data.

Event	Max p / hPa	Mass	Error <code>nls()</code>	Error t_0	Error t_e	Error t_1	Error t_2	Total Error
ANY	215	7700	220	300	40	30	40	400
PNE	215	2450	150	90	100	56	6	210
BS	147	1350	200	200	40	20	80	300
GD	215	1520	150	230	30	13	25	280

3.2 Water vapour

a sudden increase ~~after~~ following the fire and a decay after that; we fit the decay with an exponential:

$$M = M_0 \exp\left(-\frac{t}{t_d}\right)$$

105 The exponential curve is shown in Fig. 5(b). The estimate of the injected mass, M_0 , is ~~2500 ± 150~~ 2450 ± 150 Gg, with ~~a~~ an estimated decay time t_d of ~~27 ± 2~~ 28 ± 2 days; the errors here are those given by the ~~nls()~~ nls() function in R (R Core Team, 2018). The CO data from the first three days after the event are not well fitted by the exponential used to fit the subsequent days, presumably because the plume has a small horizontal extent and lies in between the MLS orbit tracks, or because material is still being lofted into the height range where MLS can observe it. These days are therefore excluded from the exponential

110 fit. The values of M_0 and t_d obtained depend rather critically on exactly which of these early points are included. The given values are obtained using data from 16 August 2017 onward; this date is marked t_0 on Fig. 5 ~~(b) with a dashed line~~ (a). This date was chosen because to use a later date is to throw away data that are part of the exponential decay, while to use earlier dates is to include data that are not part of the exponential decay. A cutoff date earlier than 16 August 2017 results in smaller estimates of M_0 and larger estimates of t_d ; later dates give a range of similar values with larger errors. The values obtained are

115 also sensitive to the end of the window used for the exponential fit, ~~t_e~~ and to the full period used for the seasonal fit, between t_1 and t_2 . A more conservative estimate of the parameters and their errors, taking into account the spread of values obtained with reasonable ranges of start and end dates, is ~~$M_0 = 2400 ± 300$~~ $M_0 = 2450 ± 210$ Gg and $t_d = 28 ± 10$ days. We note that the decision to use a constant plus semiannual and annual cycles to fit the seasonal behaviour is also an arbitrary choice. Other reasonable choices could be made and would presumably result in similar errors.

120 To place the PNE in context we have applied this technique to three other significant fire events: the Australian new year (ANY) event of 2019-20 (Schwartz et al., 2020), the Black Saturday (BS) event of February 2009 (Pumphrey et al., 2011) and the Great Divide (GD) fire of December 2006 (Pumphrey et al., 2011). The results are shown in table 1. We use a lower




Figure 6. Longitudes of enhanced H_2O , plotted as a function of time. Colours and symbols represent pressure.

125 maximum pressure for the BS event because the 215 hPa level showed very little CO from the event after the first few days. At the time it occurred, the PNE was a record-breaking event for the period of the MLS mission; it has since been superseded by the ANY event. We repeated the ANY calculation using only the 100 hPa level and higher altitudes. We obtained a mass estimate of about 1800 Gg in reasonable agreement with the 1500 ± 900 Gg above $\theta = 380$ K obtained by Khaykin et al. (2020)

Although the method described here is useful for CO and works very well for volcanic SO_2 , it is not successful for the other species described in this paper, mainly because the signal from the event is small compared to variability due to other causes.

130 [For the larger ANY event similar techniques can be used for CH₃CN and H₂O; some results are shown in Khaykin et al. \(2020\)](#)

3.1.1 [Water vapour](#)

The event described in this paper differs from the Black Saturday event (Pumphrey et al., 2011) in that ~~a large amount of water vapour is found in the polluted airmass~~[the polluted airmass has a water vapour content which is clearly a great deal higher than in the surrounding atmosphere. For example, by the time the polluted airmass reaches 68 hPa it contains mixing ratios as high as 14 ppmv against a background of 3.5 – 5.5 ppmv.](#) Figure 6 shows the longitudes of unusually high water vapour in the same manner as Fig. 4. The values chosen for the parameter α , described above for CO, vary from $\alpha = 6$ at 147 hPa to $\alpha = 2$ between 38 and 21 hPa. Many points with enhanced water vapour are found between 215 and 147 hPa and between 60° E and 180° E. These points ~~are clearly~~[occur in all years and are presumably caused by the strong convection that occurs in this region in summer as an aspect of the Asian summer monsoon \(see e.g. Jiang et al. \(2010\); Santee et al. \(2017\)\).](#) As they are not connected with the PNE ~~so we have removed them~~[from Fig. 6 all points between 215 and 147 hPa and between 60° E and 180° E.](#) It is clear from a comparison between Figs. 4 and 6 that, where both species are enhanced, the enhanced CO and H₂O occur in the same locations. The enhanced CO can be detected against the background variability by the method described 2–3 days before the enhanced H₂O becomes apparent. Inspection of daily maps of the data reveals that H₂O is clearly enhanced on 14 August 2017 between 60°N and 70°N. The algorithm used fails to detect the enhancement at this time because of the very large background variability at lower latitudes.

After about 50 days the water vapour in the polluted airmass exceeds the background value to a greater extent than is the case for CO. After 63 days, the polluted airmass can only be identified in the water vapour data; this is presumably due to the short chemical lifetime of CO in the stratosphere [and is consistent with the CO stratospheric lifetime in the ANY plume of](#)
150 [~ 30 days estimated by Schwartz et al. \(2020\).](#) A small parcel of enhanced H₂O may be seen travelling westwards from 150° E to 80° W, between 65 and 90 days after the event, before ceasing to travel westwards ~~and~~[fading into the background. By this time, it is affecting the 31 and 26 hPa levels: an altitude of approximately 24 km.](#)

3.2 ~~Other species~~

3.1.1 [Other species](#)

155 All chemical species observed by MLS were examined for the few months after the PNE using the same thresholding technique as for CO and H₂O. For ClO, we examined the standard (640 GHz) product, a secondary product derived from the 190 GHz measurements, and a product derived from the 640 GHz data while also estimating CH₃OH. The species fall into three categories:

- Species for which no anomalous mixing ratios in the plume rise above the background variability: O₃, SO₂, BrO, HNO₃, N₂O, ClO (190 GHz). Some of these species (O₃ and N₂O are the most obvious) nevertheless ~~show~~[exhibit](#) some correlation with CO values in the plume, as we show later.

Table 2. Summary of species observed by MLS to have clearly enhanced mixing ratios in the polluted airmass. The first and last days after 12 August 2017 are listed as d_0 and d_1 ; the highest and lowest pressures at which unusual values are observed are listed as p_0 and p_1 . The “BB?” column indicates whether or not the species is a genuine biomass burning product. For some species the enhanced values are thought to be caused by a measurement problem (see main text). Species observed by the instrument but with no unusual values detected in the polluted airmass are: O₃, SO₂, BrO, HNO₃, N₂O.

Species	d_0	d_1	p_0	p_1	BB ?
CO	2	63	316	31	yes
H ₂ O	2	90	316	26	yes
HCN	8	59	100	31	yes
CH ₃ CN	5	58	147	31	yes
CH ₃ Cl	5	26	147	68	yes
CH ₃ OH	5	16	147	68	yes
CIO (640 GHz)	5	33	147	46	no
HO ₂	5	11	147	68	no
HOCl	5	8	147	100	no
HCl	5	7	147	100	no

- Genuine biomass burning products: CO, H₂O, HCN, CH₃CN, CH₃OH and CH₃Cl; elevated values of these are observed for about 60 days (100 days for H₂O).
- Species for which anomalous measurements are believed to be ~~artifacts~~ artefacts arising from spectral interference due to the presence of another molecule: HOCl, HO₂, HCl, CIO (640 GHz).

165

A summary of the ranges of time and altitude over which each species has unusual values is shown in Table 2. Pumphrey et al. (2011) demonstrate that an airmass containing biomass-burning products may appear from the MLS level 2 data to contain enhanced amounts of the 640 GHz CIO product, but that this is actually caused by enhanced amounts of methanol (CH₃OH), which has a similar spectral signature to CIO in the passband of the MLS 640 GHz radiometer.

170

As is the case for many MLS data products, CH₃OH is produced from a dedicated retrieval “phase”, in which a CIO product (not the standard one) is also retrieved (see Livesey et al. (2020) for details). The CIO data from the methanol phase show far less correlation with CO than the standard 640 GHz CIO product. This also suggests that the enhanced values seen in the standard 640 GHz CIO product are an artefact caused by enhanced methanol. HOCl, HO₂ and CH₃Cl are all detected via spectral lines that are very close to the CH₃OH lines. It seems reasonable to suppose that HOCl and HO₂ appear to be enhanced in the MLS data due to spectral contamination from CH₃OH, especially as they are both observed only for a very short period. CH₃Cl, on the other hand, is known to be a biomass-burning product (Santee et al. (2013) and references therein); its spectral signature is sufficiently different from that of CIO (or CH₃OH) that the enhancement observed is almost certainly genuine. HCl, which shows the smallest enhancement in the plume, is observed at a different frequency from the CIO, CH₃OH and CH₃Cl lines, but one very close to the main CH₃CN lines in the MLS 640 GHz radiometer. Although HCl is a minor fire

175




Figure 7. Scatter plots of the mixing ratio of CH₃CN against that of CO at five pressure levels between 14 August 2017 and 30 October 2017. Only data from north of 27.5° N are shown. The small grey dots show the entire dataset within the stated latitude and time ranges. The large coloured dots are points with enhanced CO and points within 350 km (48 s) of those points along the orbit track. The linear fits use the large points only and are performed using the `lm()` function of R (R Core Team, 2018); the R^2 and p -value are shown for each fit. The lower right panel shows the slopes of the linear fits as a function of altitude.

180 product (mainly produced from the burning of garbage (Akagi et al., 2011)), it seems likely that the enhanced values observed are spurious and are caused by the large amounts of CH₃CN in the plume.

The relationships between species in the plume were investigated further by calculating a linear fit between enhanced CO and a given species at the same retrieval level and location. Data from profiles immediately adjacent to the plume are included to provide an appropriate baseline. We show an example in Fig. 7 in which the other species is CH₃CN. Plotting the data in
 185 this manner reveals that in some cases (O₃, N₂O) there is a correlation between the two species, even when the non-CO species does not have values outside its usual range. For other species (SO₂, HNO₃, BrO), no such correlation is observed, or the correlation is small enough to suggest that it is an instrumental rather than a geophysical effect. We note values of the linear fit slope in Table 3 and show a selection of the data from Table 3 in Fig. 8. For some species (including the CH₃CN shown in Fig. 7) we have used data outside the vertical range recommended by the MLS team. This was done where the correlation
 190 with CO was clear; we note that the linear fit slopes will not be affected by a constant bias in the mixing ratio of either

Table 3. Ratios between the mixing ratio of various species and that of CO, at various pressure levels. Subscripts indicate error in the last digit. The errors are those returned by a standard fitting program (the $\text{lm}()$ function in R). The right-hand column shows values from the literature converted into ratios with CO. Superscripts in this column indicate sources: Randerson et al. (2017)¹, Rinsland et al. (2007)², Simpson et al. (2011)³, Akagi et al. (2011)⁴. Published values shown are all for boreal forests with the exception of HCl. A “—” indicates either that the species does not have good data at that level or that the linear fit is not significant at the 99% level ($p > 0.01$).

Species	215 hPa	147 hPa	100 hPa	68 hPa	46 hPa	31 hPa	Other measurements
H ₂ O	119 ₅	47 ₂	18.5 ₈	32.2 ₇	39 ₁	35 ₄	—
HCN	—	—	—	0.00097 ₄	0.00167 ₄	0.0022 ₂	0.012 ^{1,4} , 0.00242 ₃₂ ² , 0.0082 ₂₀ ³
CH ₃ CN	—	0.00140 ₇	0.00104 ₃	0.00105 ₃	0.00128 ₄	0.00145 ₁₄	0.0032 ⁴ , 0.0018 ₃ ³
CH ₃ Cl	—	0.00109 ₇	0.00141 ₅	0.00127 ₇	0.0012 ₁	0.0011 ₄	0.0056 ² , 0.00014 ₃ ³
ClO std	—	0.0048 ₂	0.0056 ₂	0.0044 ₁	0.0036 ₂	0.0014 ₅	—
HCl std	—	0.0039 ₂	0.0018 ₁	-0.0010 ₂	-0.0033 ₄	—	0.0010 ₁₀ ⁴ (chaparral)
CH ₃ OH	—	0.012 ₁	0.0107 ₅	0.0058 ₂	0.0062 ₄	0.0024 ₉	0.019 ^{1,4} ₁₃ , 0.00015 ₁₅ ³
O ₃	-0.15 ₂	-0.23 ₃	-0.39 ₈	-1.81 ₁₅	-5.2 ₃	-6 ₁	—
N ₂ O	—	—	—	0.21 ₁	0.32 ₃	0.4 ₁	0.0020 ¹




Figure 8. Slopes of scatter plots of various species against CO; the slopes are calculated as in the example shown in Fig. 7. Triangles are the data from the “other measurements” column of Table 3, with their error bars (where available). All the triangles refer to measurements made in the troposphere; their position in the vertical has no meaning.

CO or the other species. For species produced by biomass burning and ~~which that~~ have similar lifetimes to CO in the lower stratosphere, we would expect the ratios observed in Table 3 to be constant with altitude and to be in reasonable agreement with the emission ratios measured previously. The relationships with CO of CH₃CN and CH₃Cl show relatively little variation with height. The slopes we obtain are generally within the published ~~range~~ranges; for CH₃Cl ~~this the~~ range is rather large. The methanol/CO ratio is within the rather wide range of published values and decreases with height. This might appear to suggest that methanol undergoes chemical loss in the plume more rapidly than CO, but the MLS methanol data are potentially useful only at 147 hPa (and at low latitudes at 100 hPa), so any such conclusion must be regarded as speculative. It would nevertheless be consistent with the ~ 10 day lifetime for methanol estimated in Schwartz et al. (2020). The agreement between our value for the methanol/CO ratio at 147 hPa and the published values is reasonable given the large variability in the latter.

The H₂O/CO ratio varies considerably with height between 215 and 68 hPa, but is approximately constant between 68 and 31 hPa. The standard 640 GHz ClO data behave in a similar manner to the biomass-burning products. This is compatible with the fact that the standard ClO retrieval may have been affected by interference or contamination from methanol (Pumphrey et al., 2011); as noted above, the other ClO products are not enhanced so the elevated abundances of the standard product are probably due to high values of methanol.

Yu et al. (2019) showed that the negative anomaly in ozone and the positive anomaly in H₂O observed by MLS were a consequence of the transport of tropospheric air into the stratosphere (and not in situ chemical loss in the case of ozone). Such upward transport probably also gave rise to increases in N₂O and thus positive correlations between N₂O and CO. Kablick et al. (2020) discuss the low ozone and high N₂O values associated with the ~~Australian New Year (ANY)~~ANY fires of December 2019. They attribute these anomalies to rapid ascent of tropospheric air; it appears likely that the ozone and N₂O in the PNE behaved in a similar way. Khaykin et al. (2020), on the other hand, suggest that the low ozone values observed in the ANY event are at least partly due to chemical depletion.

4 ~~Trajectory and dispersion modelling~~

3.1 ~~Back Trajectories~~Trajectory and dispersion modelling

3.1.1 Back Trajectories

Back trajectories from the MLS observations in the first few days after the event were calculated using the FLEXTRA trajectory model (Stohl et al., 1995), driven by analyses from the National Centers for Environmental Prediction (NCEP) Global Forecast System (GFS) ~~analyses (National Centers for Environmental Prediction, 2000).~~(National Centers for Environmental Prediction, 2000). FLEXTRA has no mechanism for modelling the rapid rise of the polluted airmass; for this reason we restrict ourselves to trajectories launched less than four days after the fire. Trajectories launched at 316 and 215 hPa pass reasonably close to the Pacific Northwest Event; we show examples in Fig. 9. Although MLS observes enhanced CO at the 147 hPa level at this time, trajectories launched at this level pass far to the west of the PNE. ThisThe vertical resolution of the MLS grid is six levels per pressure decade, or about 2.7 km, and the true vertical resolution at the levels of concern here is 4.5 – 5 km (Livesey et al., 2020)




Figure 9. Back trajectories launched at points (non-circular symbols) where MLS observed large values of CO. Trajectory colours distinguish different pressures at the launch point; the trajectories are three-dimensional. The location of the Pacific Northwest Event (PNE) is marked in red; that of the Great Slave Lake (GSL) fires in orange. Where the trajectories pass within 400 km of the PNE, the closest point is marked with a circular dot.




Figure 10. Histogram of closest-pass times of trajectories to the PNE. The median and mean values are marked in red and orange respectively. The dotted blue lines are at 00:00 UTC.




Figure 11. FLEXPART CO mixing ratio in ppmv at altitude 12 km on 15 August 2017 at 18:00, with 00. Dots show the location of MLS CO mixing ratio (also in ppmv) profiles from an orbit at a matching time. The dots are coloured to show where MLS CO, interpolated to an altitude of 12 km, is below 0.15 ppmv, and where it exceeds 0.15, 0.3 and 0.8 ppmv.

225 . If the enhanced CO were in a thin layer at an altitude between two of the MLS reporting levels, then enhanced CO would be reported at both of those levels. The fact that trajectories launched at 147 hPa do not pass close to the fire location, and that trajectories launched at 215 hPa do pass close to the fire, suggests that over these first few days the polluted air mass is closer to 215 hPa than to 147 hPa. It is also possible that the polluted air extends to altitudes lower than 215 hPa, but we cannot infer this reliably as the MLS data at 316 hPa have a larger contribution from 215 hPa than from 316 hPa (Livesey et al., 2020). There were other forest fires burning at the same time as the PNE, notably in the region around Lake Athabasca and the Great Slave Lake (GSL); the location of these fires is marked on Fig. 9. The trajectories make it reasonably clear that the source of the polluted air mass observed by MLS is the Pacific Northwest Event and not the fires near the Great Slave Lake, a point already noted by Fromm (2019).

235 By launching trajectories at many points spread between the MLS observations of high CO, both vertically and along the measurement track, we obtain sufficient times of closest pass to the PNE to construct a histogram, as shown in Fig. 10. The mean and median closest pass times are 01:20 UTC and 00:57 UTC respectively on 2018-08-13. The standard deviation of the closest-pass times is 12 hours. The closest pass times are in good agreement (given their own standard deviation) with the times reported by Peterson et al. (2018), who state that pyrocumulonimbus clouds (pyroCb) are observed to develop between 23:00 UTC on 12 August 2017 and 00:30 UTC on 13 August 2017, and that they remain active for 1–4 hours.

3.2 Plume trajectory modelling

3.1.1 Plume trajectory modelling

240 The FLEXPART model (Stohl et al., 2005) was used to model the dispersion of the plume from the Pacific Northwest Event. FLEXPART allows the user to release a mass of a pollutant at a chosen place and time and to observe how it disperses. We released the pollutant in a small cuboid ~~region; the volume. The~~ release is defined by the location and size of this ~~region~~volume in latitude, longitude and altitude, by a start and end time, and by the mass released: nine parameters in total. We show in Fig. 11 an example of FLEXPART output for a time approximately 66 hours after the PNE. An MLS orbit that passes through
245 the polluted airmass at that time is also shown; points where the MLS CO is enhanced are marked and lie close to the modelled plume.

We adjust the properties of the mass released to best match the MLS observations using the Markov ~~Chain~~chain Monte Carlo (MCMC) inverse modelling technique (van Ravenzwaaij et al., 2018). As FLEXPART is not able to model the self-lofting of the polluted airmass, we used only the first few days of data after the fire. ~~Some experimentation; the last data used are from~~
250 18 August. We also restrict the data used to those poleward of 40°N and between 40°W and 140°W. FLEXPART output is on altitude surfaces. In order to compare it with the MLS data we ran FLEXPART so that its output was in mixing ratio units on a vertical grid with a 1 km spacing. The MLS data were interpolated onto this vertical grid from their native pressure grid using the MLS geopotential height data. The FLEXPART output was interpolated to the MLS locations in order to calculate the differences. FLEXPART is set up to assume that the atmosphere contains no CO before the time of the fire; to match this
255 the MLS data are converted to anomalies by subtracting a zonal average value for the few days before the event. The MCMC technique used is the basic Metropolis-Hastings algorithm. It starts with a reasonable guess at the parameters required to define the release of CO. It then makes a sequence of small alterations to the parameters, accepting them if they make the FLEXPART output agree better with the MLS data. Parameters that agree less well than the previous set are accepted with a probability that depends on how much worse the fit is. The resulting collection of sets of parameters should, if the process is continued
260 long enough, have a probability distribution reflecting the measurement and modelling errors. To ensure that the results are not dependent on the starting point chosen, several chains are run with different starting points. The “burn-in” period for the chains was rejected by inspection.

Initial attempts to estimate all nine release parameters by this method demonstrated that the data do not provide sufficient information to determine the horizontal extent of the release region, nor its extent in time. We therefore constrained the region to
265 be 0.5° wide in longitude, 0.375° wide in latitude and 1.5 hours in duration. (These choices were somewhat arbitrary and were intended to ensure that the box was ~~smaller than~~similar in size to the observed pyroCb clouds.) The six estimated quantities were then the ~~location~~latitude, longitude and time of the release cuboid, its vertical extent, and the total mass of CO released.

We show the horizontal release locations ~~found~~deemed acceptable by the MCMC approach in Fig. 12; each of the coloured dots represents the centre of a release. The size and shape of the release box is also shown. There were several large anvil
270 clouds in the area at the time; these are clearly visible in AVHRR images and are outlined in green in Fig. 12. The MCMC solutions have a standard deviation of about 6 km perpendicular to the wind direction and 30 km in the wind direction. The




Figure 12. Map of the source region, showing areas burned in 2017 (purple; some areas are sufficiently small that the contours enclosing them appear as dots) and anvils observed in AVHRR data (green). The anvils are numbered using the same scheme as Peterson et al. (2018); their anvil 1 is outside the area shown here. The greyscale backdrop is the AVHRR channel 5 (thermal IR) image taken at about 02:00 GMT. The orange rectangle shows the size and shape in the horizontal of the cuboid release volume. The small points are the MCMC solutions for the apparent origin of the polluted air mass and are coloured according to time; see text for details varying from 2:00 GMT on 2017-08-13 (red) to 5:00 GMT (blue).

injection altitude is determined rather precisely: 11.5 ± 0.1 km; as expected from the back trajectories, this is closer to 215 hPa than to 147 hPa. The injected mass is 620 ± 80 Gg, considerably smaller than the value of 2400 ± 300 2450 ± 210 Gg that we obtained by the simple method discussed in section 3.1.1. This is qualitatively reasonable, given that the MCMC approach only considers CO that is observable by MLS near the start of the event, while the simple method of section 3.1.1 includes could include CO that was injected at too low an altitude for MLS to observe it and which subsequently ascended to observable altitudes. We consider that the method of section 3.1.1 gives the more reliable mass estimate not only for this reason, but also because it has given mass estimates of SO₂ from volcanoes that are in broad agreement with a variety of independent measurements (Pumphrey et al., 2015). Although the MCMC method does not make use of data from as many days as the simple method of section 3.1.1, it is worth noting that it uses all the MLS data within the latitude and longitude ranges noted above. The release parameters are constrained not only by the requirement to show enhanced CO where MLS observes it, but also to show no enhancement where MLS observes none.

The average release time is about 5:00 UTC on 13 August 2017, with a spread of possible values between 3:30 UTC and 7:30 UTC. This is later than the time estimated using FLEXTRA trajectories, but is well within the 12 hour standard deviation of the FLEXTRA results. As noted above, Peterson et al. (2018) report the pyroCb events as beginning between 12 August

2017, 23:00 and 13 August 2017, 00:30 and lasting 1-4 hours. The release times we determined using FLEXPART and the MLS data are at the late end of this range.

4 Discussion

Significant injections of pyroconvective plumes into the stratosphere generally require both extremely intense fires and partic-
290 ular meteorological conditions, such as those preceding the passage of a cold front ([Peterson et al., 2018](#)). The meteorology surrounding the Pacific Northwest Event is described in some detail by Peterson et al. (2018); the fire indeed occurred while a cold front was approaching.

Most forest fires do not produce a pyroCb cloud. Moreover, most pyroCb clouds do not extend to a great enough altitude to loft the products high enough for MLS to observe them. The only events in the MLS record that are in any way comparable
295 to the PNE are the Black Saturday ([BS](#)) fire, which occurred in Australia in February 2009 (Pumphrey et al., 2011), and the ~~Australian ANY~~ fires of December 2019 to January 2020 (~~paper in preparation, also Kablick et al. (2020), who refer to this as the Australian New Year (ANY) event~~, e.g. [Khaykin et al. \(2020\)](#); [Kablick et al. \(2020\)](#); [Schwartz et al. \(2020\)](#)). The mass of CO injected into the lower stratosphere by the ~~Black Saturday BS~~ fire was, ~~when estimated by the technique of as we show in~~ section 3.1.1, ~~1.3~~[1350 Tg Gg](#) — just over half of the mass injected by the PNE. Like the PNE, the Black Saturday fire
300 occurred at a time when a cold front was approaching (Dowdy et al., 2017). The 2019/20 ANY fire injected a mass of CO considerably larger even than the PNE: initial ~~esimates, again using the method of estimates, also shown in~~ section 3.1.1, ~~place this mass in the range 8–10~~[give a mass of 7700 Tg Gg](#). MLS also observed enhanced CO from the Great Divide fires in Australia in December 2006. These fires burned for two months, but MLS only observed enhanced CO from them for a short time, beginning on ~~2006-12-12. 12 December 2006~~. The injected mass of CO from the Great Divide fires was ~~between 1.5 and~~
305 ~~1.8~~[1520 Tg Gg](#), similar to that from ~~the PNE BS~~, but the polluted airmass did not ascend in the manner observed with the PNE, ANY and ~~Black Saturday BS~~ events, and was observed by MLS for only 10 days (Pumphrey et al., 2011).

The MLS data shown in this paper are for most species the first data reported from the PNE. Measurements of aerosols emitted by the event have been reported in some detail. Zuev et al. (2019), Hu et al. (2019), Ansmann et al. (2018) and Khaykin et al. (2018) report ground-based lidar observations of the PNE plume as it passed over Europe. Khaykin et al. (2018)
310 also report aerosol data from the CALIOP lidar instrument on the CALIPSO satellite. The maps of the CALIOP data are similar to Fig. 3, indicating that the aerosol and CO travelled together; presumably the aerosol particles were too small to settle out on the timescale over which the CO was observed. Khaykin et al. (2018) suggest that the pollution is from fires near Lake Athabasca; we show clearly that it is from British Columbia and from the vicinity of the Pacific Northwest Event as described by Peterson et al. (2018). Ansmann et al. (2018) ascribe the source of the pollution to British Columbia in agreement
315 with our conclusions. [Lestrelin et al. \(2021\) track the event in detail using a combination of CALIOP lidar data and ERA5 reanalyses. Their description of the event agrees in many details with our own analysis based on the MLS H₂O and CO data. Kloss et al. \(2019\) analyse the event using aerosol data from the Stratospheric Aerosol and Gas Experiment III on the International Space Station \(SAGE III/ISS, NASA/LARC/SD/ASDC \(2017\)\) and the Ozone Mapping and Profiler Suite Limb](#)

320 Profiler (OMPS-LP, Loughman et al. (2018)) instruments, claiming that the aerosol is transported around the eastern edge of the Asian monsoon anticyclone (AMA) into tropical latitudes. We observe no such behaviour in the MLS CO data. One reason for this is that by the time parts A and B1 (as labelled in Fig. 4) of the polluted airmass are in the vicinity of the AMA, they are at altitudes higher than the AMA and also higher than the aerosols discussed by Kloss et al. (2019).

In addition to the MCMC results, Fig. 12 shows the areas burned in 2017 marked in purple (British Columbia Wildfire Service, 2018). Figure 12 also shows the AVHRR thermal image; three large anvils of high cloud are clearly visible. Examination
325 of the visible bands of the same image suggests that the anvils marked “Anvil 3 & 4” and “Anvil 5” have an active pyroCb cloud at the southwest edge. It is not clear from the visible images that an active pyroCb is associated with Anvil 2; images taken during the previous two hours (Peterson et al., 2018) show that Anvil 2 formed near 00:00 UTC. Anvil 3 also formed around 00:00 UTC; Anvil 4 formed half an hour later and had merged with Anvil 3 by 01:45 UTC.

The possible solutions for the source of the polluted airmass that emerge from the MCMC inverse method are shown in
330 Fig. 12 as a cloud of small points. The colours of these dots indicate the time, varying from 2:00 GMT (red) to 5:00 GMT (blue). The location of these dots suggests that the polluted airmass observed by MLS is probably associated with Anvils 2, 3 and 4 and is less likely to be associated with Anvil 5. This in turn suggests that the emissions come from the ~~plateau~~ Plateau complex of fires (west of Nazko) and not from the Elephant Hill fires (north of Ashcroft). It could be argued that the polluted airmass did not come from the region shown in Fig. 12, but rather was injected at some point along the trajectory between
335 this region and the first MLS observation of enhanced CO. We consider such a possibility unlikely as the injection would have to have occurred in a manner that matched the shape of the polluted region as modelled by FLEXPART at the time of the hypothesised injection.

The polluted airmasses from the PNE and ANY contained large amounts of water — this distinguishes them from the Black Saturday event, in which the water content of the polluted airmass was similar to that of the surrounding air, except at 100 hPa
340 and lower altitudes during the first few days of the event.

The rapid ascent of the polluted airmass from the PNE ~~is due to its containing a large amount of~~ after its initial injection results from solar heating of its extraordinarily dense black carbon aerosol (Yu et al., 2019; Kablick et al., 2020). The ~~aerosol absorbs incident sunlight, heating the polluted airmass and causing it to rise. The~~ event can in some ways be considered a small-scale demonstration of what would happen in the aftermath of a nuclear war: such a war would cause widespread fires
345 that would put a great deal more black carbon into the middle atmosphere than the PNE did. Self-lofting of this carbon (as demonstrated by the PNE) would raise it to an altitude where it would remain for a number of years, giving rise to the so-called *nuclear winter* [Robock et al. (2007) and references therein]. Simulations (Reisner et al., 2018; Robock et al., 2007) of the response of the climate to a regional-scale nuclear war disagree on the severity of this effect. However, much of the disagreement is caused by widely different estimates of how much black carbon would reach the stratosphere from a given
350 nuclear explosion. Once it did reach the stratosphere, the black carbon aerosol from a nuclear war would ascend in the manner that we observe in the PNE.

5 Conclusions

The Pacific Northwest Event was a forest fire accompanied by unusually strong ~~convection~~pyroconvection, injecting both aerosol and gas-phase pollutants into the stratosphere. Observations of ~~gas-phase pollutants~~CO from the MLS instrument
355 show that the polluted airmass reached all longitudes on a timescale of 60 days and ascended to ~~an altitude of 24~~a pressure of 46 hPa (approximately 22 km altitude). The polluted airmass was much wetter than the surrounding air in the stratosphere; in this respect it was different from the otherwise similar Black Saturday event and similar to the recent ANY event. In water vapour, the plume was observable for 100 days, ascending to 31 hPa (approximately 24 km altitude).

The polluted airmass contained biomass-burning products HCN, CH₃CN, CH₃Cl and CH₃OH in broadly similar proportions to CO as observed by other techniques in other biomass-burning events. The evolution of the gas-phase components of the polluted airmass as observed by MLS was similar to that of its aerosol components reported elsewhere (Lestrelin et al., 2021; Khaykin et al.
360

Back trajectories link the observed excess CO clearly to the PNE as described by Peterson et al. (2018), and more detailed plume dispersion modelling shows that the pollution observed by MLS probably came from the Plateau complex of fires to the west of Nazko. The time of injection given by this method is in agreement with the description of the event given by Peterson et al. (2018).
365

A simple approach to estimating the mass of CO injected above 215 hPa leads to a value of 2450±210 Gg. We consider this figure more reliable than the lower number of 620±80 Gg obtained by plume dispersion modelling. However, we also consider that the discrepancy bears further investigation.

370 Events of the type we describe here are rare; the PNE, ANY, Black Saturday and Great Divide fires are the only such events in the MLS dataset at the time of writing. Whether they will become ~~rarer or more~~more or less frequent as the climate changes over the coming decades remains to be seen.

Code and data availability. The FLEXTRA and FLEXPART models are available at <https://www.flexpart.eu/>. The MLS data used in this paper are available at <https://disc.gsfc.nasa.gov/>. The data for each species has a separate DOI, e.g. 10.5067/Aura/MLS/DATA2005 (CO),
375 10.5067/Aura/MLS/DATA2009 (H₂O). The AVHRR L1B data can be obtained from <https://www.bou.class.noaa.gov/saa/products/welcome>. The GFS analyses (National Centers for Environmental Prediction, 2000) are available from <https://doi.org/10.1038/s41612-018-0039-3>.

Author contributions. HCP wrote most of the paper and carried out a large part of the data analysis and modelling. MJS provided some of the data analysis and suggested the use of figures 4, 6 and 7. MLS provided advice on the CH₃OH, ClO, CH₃CN and CH₃Cl products, was responsible for the use of Fig. 8 and suggested many improvements to Fig. 7. MDF assisted with the trajectory analysis. All co-authors
380 commented on multiple drafts prior to submission, helped to shape the analysis, provided HCP with a great deal of assistance in responding to the referees' comments and suggested many improvements to the text.

Competing interests. The authors declare that they have no competing interests.

Acknowledgements. [The authors thank Albert Ansmann for drawing our attention to two important references.](#) Work at the Jet Propulsion Laboratory, California Institute of Technology, was done under contract with the National Aeronautics and Space Administration. MLS
385 research at Edinburgh was funded by NERC under grant NE/E003990/1 and previous grants.

References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- 390 Ansmann, A., Baars, H., Chudnovsky, A., Mattis, I., Veselovskii, I., Haarig, M., Seifert, P., Engelmann, R., and Wandinger, U.: Extreme levels of Canadian wildfire smoke in the stratosphere over central Europe on 21–22 August 2017, *Atmospheric Chemistry and Physics*, 18, 11 831–11 845, <https://doi.org/10.5194/acp-18-11831-2018>, 2018.
- Baars, H., Ansmann, A., Ohneiser, K., Haarig, M., Engelmann, R., Althausen, D., Hanssen, I., Gausa, M., Pietruczuk, A., Szkop, A., Stachlewska, I. S., Wang, D., Reichardt, J., Skupin, A., Mattis, I., Trickl, T., Vogelmann, H., Navas-Guzmán, F., Haeffele, A., Acheson, K., Ruth, A. A., Tatarov, B., Müller, D., Hu, Q., Podvin, T., Goloub, P., Veselovskii, I., Pietras, C., Haeffelin, M., Fréville, P., Sicard, M., Comerón, A., Fernández García, A. J., Molero Menéndez, F., Córdoba-Jabonero, C., Guerrero-Rascado, J. L., Alados-Arboledas, L., Bortoli, D., Costa, M. J., Dionisi, D., Liberti, G. L., Wang, X., Sannino, A., Papagiannopoulos, N., Boselli, A., Mona, L., D’Amico, G., Romano, S., Perrone, M. R., Belegante, L., Nicolae, D., Grigorov, I., Gialitaki, A., Amiridis, V., Soupiona, O., Papayannis, A., Mamouri, R.-E., Nisantzi, A., Heese, B., Hofer, J., Schechner, Y. Y., Wandinger, U., and Pappalardo, G.: The unprecedented 2017–2018 stratospheric smoke event: decay phase and aerosol properties observed with the EARLINET, *Atmospheric Chemistry and Physics*, 19, 15 183–15 198, <https://doi.org/10.5194/acp-19-15183-2019>, 2019.
- 400 Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D’Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth System, *Science*, 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.
- 405 British Columbia Wildfire Service: Fire Perimeters — Historical, <https://catalogue.data.gov.bc.ca/dataset/22c7cb44-1463-48f7-8e47-88857f207702>, accessed: 2019-06-29, 2018.
- Dowdy, A. J., Fromm, M. D., and McCarthy, N.: Pyrocumulonimbus lightning and fire ignition on Black Saturday in southeast Australia, *J. Geophys. Res. Atmos.*, 122, 7342–7354, <https://doi.org/10.1002/2017JD026577>, 2017.
- 410 Fromm, M.: Interactive comment on “Transport of the 2017 Canadian wildfire plume to the tropics and global stratosphere via the Asian monsoon circulation” by Corinna Kloss et al., *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2019-204-SC1>, 2019.
- Government of British Columbia: 2017 Wildfire Season Summary, <https://www2.gov.bc.ca/gov/content/safety/wildfire-status/about-bcws/wildfire-history/wildfire-season-summary>, accessed: 2017-06-17, 2018.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 420 Hu, Q., Goloub, P., Veselovskii, I., Bravo-Aranda, J.-A., Popovici, I. E., Podvin, T., Haeffelin, M., Lopatin, A., Dubovik, O., Pietras, C., Huang, X., Torres, B., and Chen, C.: Long-range-transported Canadian smoke plumes in the lower stratosphere over northern France, *Atmospheric Chemistry and Physics*, 19, 1173–1193, <https://doi.org/10.5194/acp-19-1173-2019>, 2019.

- Jiang, J. H., Su, H., Pawson, S., Liu, H.-C., Read, W. G., Waters, J. W., Santee, M. L., Wu, D. L., Schwartz, M. J., Livesey, N. J., Lambert, A., Fuller, R. A., and Lee, J. N.: Five year (2004–2009) observations of upper tropospheric water vapor and cloud ice from MLS and comparisons with GEOS-5 analyses, *Journal of Geophysical Research: Atmospheres*, 115, D15 103, <https://doi.org/10.1029/2009JD013256>, 2010.
- Kablick, III, G. P., Allen, D. R., Fromm, M. D., and Nedoluha, G. E.: Australian pyroCb smoke generates synoptic-scale stratospheric anticyclones, *Geophysical Research Letters*, 47, e2020GL088 101, <https://doi.org/10.1029/2020GL088101>, 2020.
- Khaykin, S., Legras, B., Bucci, S., Sellitto, P., Isaksen, L., Tencé, F., Bekki, S., Bourassa, A., Rieger, L., Zawada, D., Jumelet, J., and Godin-Beekmann, S.: The 2019/20 Australian wildfires generated a persistent smoke-charged vortex rising up to 35 km altitude, *Commun. Earth Environ.*, 1, 22, <https://doi.org/10.1038/s43247-020-00022-5>, 2020.
- Khaykin, S. M., Godin-Beekmann, S., Hauchecorne, A., Pelon, J., Ravetta, F., and Keckhut, P.: Stratospheric smoke with unprecedentedly high backscatter observed by lidars above southern France., *Geophys. Res. Lett.*, 45, 1639–1646, <https://doi.org/10.1002/2017GL076763>, 2018.
- Kloss, C., Berthet, G., Sellitto, P., Ploeger, F., Bucci, S., Khaykin, S., Jégou, F., Taha, G., Thomason, L. W., Barret, B., Le Flochmoen, E., von Hobe, M., Bossolasco, A., Bègue, N., and Legras, B.: Transport of the 2017 Canadian wildfire plume to the tropics and global stratosphere via the Asian monsoon circulation, *Atmospheric Chemistry and Physics Discussions*, 2019, 1–22, <https://doi.org/10.5194/acp-2019-204>, 2019.
- Lambert, A., Read, W., Livesey, N., Santee, M., Manney, G., Froidevaux, L., Wu, D., Schwartz, M., Pumphrey, H., Jimenez, C., Nedoluha, G., Cofield, R., Cuddy, D., Daffer, W., Drouin, B., Fuller, R., Jarnot, R., Knosp, B., Pickett, H., Perun, V., Snyder, W., Stek, P., Thurstans, R., Wagner, P., Waters, J., Jucks, K., Toon, G., Stachnik, R., Bernath, P., Boone, C., Walker, K., Urban, J., Murtagh, D., Elkins, J., and Atlas, E.: Validation of the Aura Microwave Limb Sounder middle atmosphere water vapor and nitrous oxide measurements, *J. Geophys. Res.*, 112, D24S35, <https://doi.org/10.1029/2007JD008752>, 2007.
- Lestrelin, H., Legras, B., Podglajen, A., and Salihoglu, M.: Smoke-charged vortices in the stratosphere generated by wildfires and their behaviour in both hemispheres: comparing Australia 2020 to Canada 2017, *Atmospheric Chemistry and Physics*, 21, 7113–7134, <https://doi.org/10.5194/acp-21-7113-2021>, 2021.
- Livesey, N. J., Fromm, M. D., Waters, J. W., Manney, G. L., Santee, M. L., and Read, W. G.: Enhancements in lower stratospheric CH₃CN observed by UARS MLS following boreal forest fires., *J. Geophys. Res.*, 109, D06308, <https://doi.org/10.1029/2003JD004055>, 2004.
- Livesey, N. J., Filipiak, M., Froidevaux, L., Read, W., Lambert, A., Santee, M., Jiang, J., Pumphrey, H., Waters, J., Cofield, R., Cuddy, D., Daffer, W., Drouin, B., Fuller, R., Jarnot, R., Jiang, Y., Knosp, B., Li, Q., Perun, V., Schwartz, M., Snyder, W., Stek, P., Thurstans, R., Wagner, P., Avery, M., Browell, E., Cammas, J.-P., Christensen, L., Diskin, G., Gao, R.-S., Jost, H.-J., Loewenstein, M., Lopez, J., Nedelec, P., Osterman, G., Sachse, G., and Webster, C.: Validation of Aura Microwave Limb Sounder O₃ and CO observations in the upper troposphere and lower stratosphere, *J. Geophys. Res.*, 113, D15S02, <https://doi.org/10.1029/2007JD008805>, 2008.
- Livesey, N. J., Read, W. G., Wagner, P. A., Froidevaux, L., Lambert, A., Manney, G. L., Valle, L. M., Pumphrey, H. C., Santee, M. L., Schwartz, M. J., Wang, S., Fuller, R. A., Jarnot, R. F., Knosp, B. W., Martinez, E., and Lay, R. R.: Earth Observing System (EOS) Aura Microwave Limb Sounder (MLS) Version 4.2x Level 2 and 3 data quality and description document., Tech. Rep. JPL D-33509 Rev E, NASA Jet Propulsion Laboratory California Institute of Technology, Pasadena, California, 91109-8099, <http://mls.jpl.nasa.gov>, 2020.
- Loughman, R., Bhartia, P. K., Chen, Z., Xu, P., Nyaku, E., and Taha, G.: The Ozone Mapping and Profiler Suite (OMPS) Limb Profiler (LP) Version 1 aerosol extinction retrieval algorithm: theoretical basis, *Atmospheric Measurement Techniques*, 11, 2633–2651, <https://doi.org/10.5194/amt-11-2633-2018>, 2018.

- NASA/LARC/SD/ASDC: SAGE III/ISS L2 Solar Event Species Profiles (Native) V051, https://doi.org/10.5067/ISS/SAGEIII/SOLAR_BINARY_L2-V5.1, 2017.
- National Centers for Environmental Prediction: NCEP FNL Operational Model Global Tropospheric Analyses, continuing from July 1999, <https://doi.org/10.5065/D6M043C6>, 2000.
- 465 Peterson, D. A., Campbell, J. R., Hyer, E. J., Fromm, M. D., Kablick III, G. P., Cossuth, J. H., and DeLand, M. T.: Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke, *Nature Partner Journal: Climate and Atmospheric Science*, 1, 30, <https://doi.org/10.1038/s41612-018-0039-3>, 2018.
- Pumphrey, H. C., Filipiak, M. J., Livesey, N. J., Schwartz, M. J., Boone, C., Walker, K. A., Bernath, P., Ricaud, P., Barret, B., Clerbaux, C., Jarnot, R. F., Kovalenko, L. J., Manney, G. L., and Waters, J. W.: Validation of middle-atmosphere carbon monoxide retrievals from MLS
470 on Aura, *Journal of Geophysical Research*, 112, D24S38, <https://doi.org/10.1029/2007JD008723>, 2007.
- Pumphrey, H. C., Santee, M. L., Livesey, N. J., Schwartz, M. J., and Read, W. G.: Microwave Limb Sounder observations of biomass-burning products from the Australian bush fires of February 2009, *Atmos. Chem. Phys.*, 11, 6285–6296, <https://doi.org/10.5194/acp-11-6285-2011>, 2011.
- Pumphrey, H. C., Read, W. G., Livesey, N. J., and Yang, K.: Observations of volcanic SO₂ from MLS on Aura, *Atmos. Meas. Tech.*, 8,
475 195–209, <https://doi.org/10.5194/amt-8-195-2015>, 2015.
- Pumphrey, H. C., Glatthor, N., Bernath, P. F., Boone, C. D., Hannigan, J., Ortega, I., Livesey, N. J., and Read, W. G.: MLS measurements of stratospheric hydrogen cyanide during the 2015–16 El Niño event, *Atmospheric Chemistry and Physics*, 2018, 691–703, <https://doi.org/10.5194/acp-18-691-2018>, 2018.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2018.
- 480 //www.R-project.org/, 2018.
- Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire Emissions Database, Version 4.1 (GFEDv4), <https://doi.org/10.3334/ORN LDAAC/1293>, <https://doi.org/10.3334/ORN LDAAC/1293>, last access 15 January 2018, 2017.
- Reisner, J., D’Angelo, G., Koo, E., Even, W., Hecht, M., Hunke, E., Comeau, D., Bos, R., and Cooley, J.: Climate Impact of a Regional Nuclear Weapons Exchange: An Improved Assessment Based On Detailed Source Calculations, *J. Geophys. Res. Atmospheres*, 123,
485 2752–2772, <https://doi.org/10.1002/2017JD027331>, 2018.
- Rinsland, C. P., Dufour, G., Boone, C. D., Bernath, P. F., Chiou, L., Pierre-Fran c. C., Turquety, S., and Clerbaux, C.: Satellite boreal measurements over Alaska and Canada during June–July 2004: Simultaneous measurements of upper tropospheric CO, C₂H₆, HCN, CH₃Cl, CH₄, C₂H₂, CH₃OH, HCOOH, OCS, and SF₆ mixing ratios, *Global Biogeochemical Cycles*, 21, GB3008, <https://doi.org/10.1029/2006GB002795>, 2007.
- 490 Robock, A., Oman, L., Stenchikov, G. L., Toon, O. B., Bardeen, C., and Turco, R. P.: Climatic consequences of regional nuclear conflicts, *Atmospheric Chemistry and Physics*, 7, 2003–2012, <https://doi.org/10.5194/acp-7-2003-2007>, 2007.
- Santee, M., Livesey, N., Manney, G., Lambert, A., and Read, W.: Methyl chloride from the Aura Microwave Limb Sounder: First global climatology and assessment of variability in the upper troposphere and stratosphere, *J. Geophys. Res. Atmospheres*, 118, 13 532 – 13 560, <https://doi.org/10.1002/2013JD020235>, 2013.
- 495 Santee, M. L., Manney, G. L., Livesey, N. J., Schwartz, M. J., Neu, J. L., and Read, W. G.: A comprehensive overview of the climatological composition of the Asian summer monsoon anticyclone based on 10 years of Aura Microwave Limb Sounder measurements, *Journal of Geophysical Research: Atmospheres*, 122, 5491–5514, <https://doi.org/10.1002/2016JD026408>, 2017.

- Schwartz, M. J., Pumphrey, H. C., Santee, M. L., Manney, G. L., Lambert, A., Livesey, N. J., Millán, L., Neu, J. L., Read, W. J., and Werner, F.: Australian New Year's PyroCb impact on Stratospheric Composition, *Geophys. Res. Lett.*, 47, e2020GL090831, <https://doi.org/10.1029/2020GL090831>, 2020.
- 500
- Simpson, I. J., Akagi, S. K., Barletta, B., Blake, N. J., Choi, Y., Diskin, G. S., Fried, A., Fuelberg, H. E., Meinardi, S., Rowland, F. S., Vay, S. A., Weinheimer, A. J., Wennberg, P. O., Wiebring, P., Wisthaler, A., Yang, M., Yokelson, R. J., and Blake, D. R.: Boreal forest fire emissions in fresh Canadian smoke plumes: C₁-C₁₀ volatile organic compounds (VOCs), CO₂, CO, NO₂, NO, HCN and CH₃CN, *Atmospheric Chemistry and Physics*, 11, 6445–6463, <https://doi.org/10.5194/acp-11-6445-2011>, 2011.
- 505
- Stohl, A., Wotawa, G., Siebert, G., and Kromp-Kolb, H.: Interpolation errors in wind fields as a function of spatial and temporal resolution and their impact on different types of kinematic trajectories, *J. Appl. Meteor.*, 34, 2149–2165, 1995.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, 2005.
- Torres, O., Bhartia, P. K., Taha, G., Jethva, H., Das, S., Colarco, P., Krotkov, N., Omar, A., and Ahn, C.: Stratospheric Injection of Massive Smoke Plume From Canadian Boreal Fires in 2017 as Seen by DSCOVR-EPIC, CALIOP, and OMPS-LP Observations, *J. Geophys. Res. Atmospheres*, 125, e2020JD032579, <https://doi.org/10.1029/2020JD032579>, 2020.
- 510
- van Ravenzwaaij, D., Cassey, P., and Brown, S. D.: A simple introduction to Markov Chain Monte-Carlo sampling, *Psychon. Bull. Rev.*, 25, 143–154, <https://doi.org/10.3758/s13423-016-1015-8>, 2018.
- Waters, J. W., Froidevaux, L., Harwood, R., Jarnot, R., Pickett, H., Read, W., Siegel, P., Cofield, R., Filipiak, M., Flower, D., Holden, J., Lau, G., Livesey, N., Manney, G., Pumphrey, H., Santee, M., Wu, D., Cuddy, D., Lay, R., Loo, M., Perun, V., Schwartz, M., Stek, P., Thurstans, R., Boyles, M., Chandra, S., Chavez, M., Chen, G.-S., Chudasama, B., Dodge, R., Fuller, R., Girard, M., Jiang, J., Jiang, Y., Knosp, B., LaBelle, R., Lam, J., Lee, K., Miller, D., Oswald, J., Patel, N., Pukala, D., Quintero, O., Scaff, D., Snyder, W., Tope, M., Wagner, P., and Walch, M.: The Earth Observing System Microwave Limb Sounder (EOS MLS) on the Aura satellite, *IEEE Trans. Geoscience and Remote Sensing*, 44, 1106–1121, 2006.
- 515
- Williams, A. P. and Abatzoglou, J. T.: Recent Advances and Remaining Uncertainties in Resolving Past and Future Climate Effects on Global Fire Activity, *Curr Clim Change Rep.*, 2, 1–14, <https://doi.org/10.1007/s40641-016-0031-0>, 2016.
- 520
- Williams, A. P., Abatzoglou, J. T., Gershunov, A., Guzman-Morales, J., Bishop, D. A., Balch, J. K., and Lettenmaier, D. P.: Observed Impacts of Anthropogenic Climate Change on Wildfire in California, *Earth's Future*, 7, 892–910, <https://doi.org/10.1029/2019EF001210>, 2019.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the CALIPSO Mission and CALIOP Data Processing Algorithms, *Journal of Atmospheric and Oceanic Technology*, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009.
- 525
- Yu, P., Toon, O. B., Bardeen, C. G., Zhu, Y., Rosenlof, K. H., Portmann, R. W., Thornberry, T. D., Gao, R.-S., Davis, S. M., Wolf, E. T., de Gouw, J., Peterson, D. A., Fromm, M. D., and Robock, A.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume, *Science*, 365, 587–590, <https://doi.org/10.1126/science.aax1748>, 2019.
- 530
- Zuev, V. V., Gerasimov, V. V., Nevzorov, A. V., and Savelieva, E. S.: Lidar observations of pyrocumulonimbus smoke plumes in the UTLS over Tomsk (Western Siberia, Russia) from 2000 to 2017, *Atmospheric Chemistry and Physics*, 19, 3341–3356, <https://doi.org/10.5194/acp-19-3341-2019>, 2019.