

Responses to Anonymous Referee #1

General comments:

This manuscript investigated the aerosol hygroscopic properties for relatively large particles with a diameter larger than 300nm. As previous studies focus on fine particles, this work presented the significant hygroscopic variation of large particles.

- 5 This unique measurement can enhance our understanding of aerosol hygroscopic properties. Therefore, I recommend the manuscript for publication in Atmos. Chem. Phys., after properly addressing following technical but important problems.

General response: Thank you for your thorough and detail review of our manuscript. Your comments are very helpful in improving this work. Here we respond to your comments one by one. Your comments are in italics, and my responses are in plain text.

10

Specific Comments:

1. *Line 14: Please define the GF.*

Response: The definition of GF was added in the text.

15

2. *Line 28-29: Please check the citation format again.*

Response: The citation and reference was updated to be consistent with ACP format.

3. *Line 48: The citation is wrong, should be (Swietlicki et al. 2008).*

Response: Thank you for your comment. The correction was made in the new manuscript.

20

4. *Section 2.1: I did not understand why the authors introduced the AERONET station here, since it has not been used in this study.*

Response: Thank you for your suggestion. The description concerning the AERONET station was deleted in the new manuscript.

25

5. *Line 131: The GF should be defined before.*

Response: It was defined in the abstract.

6. *Line 142: Why the third-moment mean values? Please specify.*

30

Response: The third-moment mean value is the volume-weighted mean value of growth factor. It can be directly used to simplify the calculation of total liquid water content for the whole particle population.

7. *Line 143: "... can be calculated from GF_{mean} using the equation above." Please specify which equation. The author should number the formula.*

35

Response: Thank you for your comment. The formula was numbered and the equation was also updated in the new manuscript.

8. Line 152: It seems that the BC information is not used in this study. Is there a need to mention this?

Response: The BC information was used in the Mie model to calculate the aerosol extinction.

40


9. Section 4.2: The Mie theory was used to describe the aerosol effects on visibility degradation. Please describe to what extent this theory can be used in the estimation.

Response: The Mie model is widely employed to estimate the aerosol optical properties given specific physical and chemical characteristics. To evaluate its applicability in aerosol light scattering calculation, an optical closure study was done.

45

In the closure study, we have measured aerosol light scattering coefficients at (635, 525, 450) nm from Aurora 3000 nephelometer. We can also calculate the aerosol light scattering at 525 nm from particle number size distribution (PNSD) using Mie scattering model. The comparison between the measured and calculated aerosol light scattering is shown in the figure below. It can be seen that good consistency is achieved between measured and calculated values, which shows both the data reliability and model applicability.

50



10. In Fig.7(b): Please accurately replot this panel with the wind direction. I do not understand the color bar, which only shows south wind and north wind.

55

Response: Thank you for your suggestion. This figure was updated with both wind speeds and directions in the new manuscript.

11. Line 290: It's better to specify the Koschmieder relation here.

Response: The Koschmieder relation was added in the text.

60

12. *Line 293: Please at least give some descriptions about the calculation method instead of just putting a citation here. Some necessary but not key information can also be offered in the supplement.*

Response: Thank you for your comment. We added some brief description of the calculation process in the new manuscript.

Responses to Anonymous Referee #2

General Response: Thank you for your positive review of our manuscript. We sincerely appreciate the efforts you have put in the review process, and we improve this work based on your comments and suggestions. Below we will respond to your comments one by one. Your comments are in italics, and my responses are in plain text. All the changes have been included in the newest version of our manuscript.

Specific comments:

The manuscript by Shen et al. report size-resolved aerosol hygroscopicity measurements over an extended size range of 50-600 nm. This size range covers the mode diameter of ambient aerosol particles, and can thus provide more useful information about the optical properties and the climate impact of aerosol hygroscopic growth. They show that on average the number fraction of more hygroscopic mode particles decreases with increasing particle size for 400 nm or larger. However, the more hygroscopic mode in the larger size range is dominant during the polluted events, consistent with the general consensus that aqueous production of secondary species plays an important role in the formation of winter haze in Beijing. I think the measurement data and analysis are solid. The paper is overall clearly written but could be further improved in English. I, therefore, recommend publication in ACP once the authors address a few issues:

1. *In Line 16, the authors mention “unexpected low hygroscopicity”; however, in line 18 they say “this result is supported by previous chemical composition analysis”. These two statements seem to be contradictory. Also, the authors should briefly discuss why the hygroscopicity decreases with particle size for large particles, based on previous chemical analysis (inorganic/organic fraction, dust, etc).*

response: Thank you for your comment. These two statements are not contradictory. The “unexpected low hygroscopicity” means that the relatively low average hygroscopicity, especially during clean conditions, is unexpected because the accumulation mode particles used to be thought as more hygroscopic. However, when the pollution evolves, the growth and expansion of MH (more hygroscopic) group particles is consistent with general consensus that secondary species play a dominant role in the pollution evolution.

For the second comment, we have some discussions in the section 3.1 (line 175-183). In the discussion, we compare our results with previous chemical analysis and find that the decreased hygroscopicity may be related to the increased mass ratio of EC, dust, or other undefined mass in larger sizes. Since we didn't have simultaneous chemical constituent measurements, we cannot give direct chemical evidence.

2. *It might be helpful to explain why the TDMA used in this study can extend the measurements to larger particle sizes. Is this because of a lower sheath flow rate or different geometry of the DMA?*

Response: The TDMA can extend the upper size limit because of its unique geometry design. For the DMA, the relationship between electrical mobility size and voltage follows the equation:

$$V = \frac{Q_{sheath} \cdot \ln\left(\frac{R_{outer}}{R_{inner}}\right)}{2\pi L_{DMA} Z_p}$$

For the same electrical mobility diameter, a lower voltage is needed if the $\frac{R_{outer}}{R_{inner}}$ is smaller. That's to say, a lower ratio of R_{outer} to R_{inner} will extend the upper size limit within the same voltage limit.

We compare the commonly used TSI 3080 DMA with the BMI 2100 DMA in the table below. It can be seen that the special geometry of the BMI 2100 DMA ensures that it can have a larger size limit.

105

Model	V	L _{DMA} (cm)	$\frac{R_{outer}}{R_{inner}}$
TSI 3080	10000	44.37	$\frac{1.961}{0.937} = 2.09$
BMI 2100	5600	34.02	$\frac{3.613 \text{ cm}}{3.120 \text{ cm}} = 1.16$

3. *What does the red line in Fig. 1c mean?*

110

Response: the red line in Fig.1c represents the mean values of calculated size-resolved light extinction coefficients during the whole measurement period. Please check the text in red in the following:

Figure 1: Frequency distribution of (a) aerosol number size distribution during the whole measurement period, (b) corresponding volume size distribution and (c) size-resolved light extinction coefficient contribution calculated from Mie theory. The red lines in (a-c) represent the mean values. The lines with marks in Fig. 1(c) denote the hygroscopicity measurement by HTDMA in previous and our studies.

115

4. *The citation and bibliography styles should be consistent with the ACP format.*

Response: Thank you for your suggestion. We updated the citation and bibliography format in the new manuscript.

120 **Measurement report: Aerosol hygroscopic properties extended to 600 nm in the urban environment**

Chuanyang Shen¹, Gang Zhao^{1,2}, Weilun Zhao¹, Ping Tian³, Chunsheng Zhao^{1*}

¹ Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing 100871, China

² College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

125 ³ Beijing Key Laboratory of Cloud, Precipitation and Atmospheric Water Resources, Beijing 100089, China

Correspondence to: Chunsheng Zhao (zcs@pku.edu.cn)

Abstract. Submicron particles larger than 300 nm dominate the aerosol light extinction and mass concentration in the atmosphere. The water uptake ability of this size range greatly influences the particle mass, visibility degradation, and particle chemistry. However, most of previous field measurements on aerosol hygroscopicity are limited within 350 nm. In this study, the size-resolved aerosol hygroscopic properties over extended size range (50-600 nm) at 85% relative humidity were investigated in Beijing winter from 27 November 2019 to 14 January 2020 using a Humidity Tandem Differential Mobility Analyser (HTDMA) instrument. The corresponding aerosol optical properties were also analyzed using the Mie scattering theory. Results show that the averaged probability distribution of growth factor GF (GF-PDF) is generally a constitute of a more-hygroscopic (MH) group and a less-hygroscopic (LHHeyder et al.) group (including hydrophobic). For the particles larger than 300 nm, there exist a large fraction of LH group particles, resulting in an unexpected low hygroscopicity. During the development of pollution when particles are gradually aged and accumulated, the bulk hygroscopicity above 300 nm is enhanced significantly by the growth and expansion of the MH group. This result is supported by previous chemical composition analysis, and we give a more direct and detailed evidence from the growth factor and mixing state aspects. Our calculations indicate that the optical contribution of particles larger than 300 nm constitutes about 2/3 of the total aerosol extinction. The large hygroscopic variation of aerosols above 300 nm will influence the light degradation comparably with the increase of aerosol loading in the low visibility haze events. Our studies highlight that the hygroscopic properties above 300 nm are complex and vary greatly with different pollution stages. Therefore more field measurements and investigations need to be done in the future.

1 Introduction

145 Aerosol particles can scatter and absorb solar radiation, and act as cloud condensation nuclei in the atmosphere. Through these effects, they can influence the radiative balance of the Earth-Atmosphere system (Twomey, 1974;ALBRECHT, 1989;CHARLSON et al., 1992;Penner et al., 2001;Lohmann and Feichter, 2005;Haywood and Boucher, 2000). They can also impact human life by degrading environmental visibility and despairing respiratory health (Ibald-Mulli et al., 2002;Peters et al., 1997;Chang et al., 2009). All these effects can be largely enhanced by aerosol's hygroscopic property (Kreidenweis and Asa-Awuku, 2014), which is the ability to absorb water under an elevated relative humidity (RH). The particle's water uptake

property, which is mainly related to the water-soluble materials contained, determines aerosol liquid water content, affects the multiphase chemistry and local photochemistry, and facilitates particle formation and aging processes (Wu et al., 2018). Thus, a thorough understanding of aerosol hygroscopicity is crucial to quantify all these effects.

155 So far, while many field measurements on aerosol hygroscopicity have been carried out worldwide using Hygroscopicity
Tandem Differential Mobility Analyzer (HTDMA) (Massling et al., 2009; Park et al., 2009; Hersey et al., 2011; Kitamori et al.,
2009; Martin et al., 2011), Cloud Condensation Nuclei Counter (CCNC) (Deng et al., 2013; Ma et al., 2016; Tao et al., 2018),
and optical humidified measurements ($f(\text{RH})$) (Carrico et al., 2000; Zieger et al., 2010; Yan et al., 2009; Fitzgerald et al.,
1982; Kuang et al., 2015), detailed hygroscopic properties over large sizes ($> 300 \text{ nm}$) are still very scarce. CCNC can give an
160 estimate of bulk aerosol hygroscopicity under different supersaturations. Because of the supersaturation limits, it mostly
reveals hygroscopicity property within 200 nm and cannot provide direct size-resolved or mixing state information (Roberts
and Nenes, 2005). $F(\text{RH})$ measures the humidity-dependent scattering coefficient over the whole size range. It focuses more
on the hygroscopicity of sizes that contribute more to the bulk optical properties (Kuang et al., 2018), and cannot give size
information or mixing states, either. HTDMA is a widely used instrument to obtain the size-resolved growth factor of aerosol
165 particles under different RH, and has been applied to many field measurements to obtain aerosol bulk hygroscopicity as well
as mixing state information. However, most of these field studies were limited within the size range of 300 nm (Swietlicki et
al., 2008) ~~(Swietlicki et al., 2017, 2008).~~

For the submicron aerosols in the urban environment, especially areas suffering from severe haze pollution and affected by
170 intensive anthropogenic activities, the particles larger than 300 nm contribute significantly to the particle mass and surface
concentration (Lundgren and Paulus, 2012; Sverdrup, 1977), liquid water content (Bian et al., 2014), and optical properties
(Ouimette and Flagan, 1982; Kuang et al., 2018). For the submicron particles, although the number distribution is dominated
by particles smaller than 0.1 μm , most of the surface area is in the 0.1-0.5 μm size range (Seinfeld and Pandis, 2006). The
aerosol mass distribution peaks at an even higher size range of about 0.2-1 μm . Moreover, light scattering is usually
175 approximately proportional to aerosol volume or mass (Pinnick et al., 1980), which means that the light degradation by aerosol
particles is also concentrated at a relatively ~~larger~~ higher accumulation size. Ma et al. (2012) demonstrated that the aerosol
particles between 200 nm and 1 μm usually contribute more than 80% to light extinction at 550 nm during summer on the
North China Plain (NCP). In the pollution period, the enhanced particle growth by coagulation and condensation of vapors,
aerosol size distribution will shift to larger diameters. The aerosol mass and optical extinction contributed by the larger
180 accumulation mode particles will also be increased. When exposed to a high RH, these factors can be further enhanced with
the addition of water. Considering the dominant contribution of this part to aerosol optical properties, it's necessary to study
the detailed hygroscopic properties over extended size range and investigate its variation during different pollution conditions.

Some previous studies have tried to derive the hygroscopicity in a larger accumulation size above 300 nm. However, there is
185 no direct evidence and measurements supporting their assumptions or parameterization schemes. Kuang et al. (2017) calculated
the equivalent aerosol hygroscopicity parameter κ from $f(\text{RH})$ curves ($\kappa_{f(\text{RH})}$) measured at Wangdu on the NCP and the results
lied between 0.06 and 0.51, presenting a very large variation range. Because the light scattering is mostly contributed by
particles at a larger accumulation size, so $\kappa_{f(\text{RH})}$ can generally represent the hygroscopicity in this size range. Liu et al. (2014)
190 also derived the hygroscopicity over a larger size range of 30 nm- 4 μm from chemical composition and found that κ values
vary relatively less for particles between 250 nm and 1 μm than particles smaller than 250 nm. Chen et al. (2012) assumed that
particles in each aerosol mode had the same hygroscopicity, and then estimated size-resolved κ for aerosols with diameters of
3 nm ~10 μm based on the contribution of each mode to a specific particle size. However, all these estimations are indirect
methods and have some unproven assumptions.

195 Beijing, one of the biggest cities in China and the world, is a densely populated area with severe particle pollution.
Representative of the urban environment, the consumption of fossil fuels is quite considerable, and a lot of pollutants are
emitted into the ~~atmosphere-environment~~ every day. Exacerbated by meteorological conditions, these pollutants are readily
trapped in ~~the~~ local region, go through a series of physical and chemical processes, and finally evolved into severe haze events
(Guo et al., 2014;Sun et al., 2014;Jiang et al., 2015;Ye et al., 2016). Influenced by the particle pollution and strong hygroscopic
200 growth of aerosol particles, visibility degradation is also a frequent and urgent environmental problem. Based on this condition,
many field measurements focusing on the particles' microphysical properties, including hygroscopicity, were conducted in
Beijing and the surrounding region. Massling et al. (2009) used a custom-made HTDMA to measure the hygroscopic
properties of ambient particles in Beijing and reported the corresponding measurement results. Haze in China (HaChi)
campaign also investigate aerosol particles' various microphysical properties, including CCN activity, optical enhancement
205 factor and hygroscopic growth properties under high humidity (Deng et al., 2011;Liu et al., 2014;Liu et al., 2011b;Chen et al.,
2014;Shen et al., 2018). Wang et al. (2018) made statistical analysis and proposed parameterization of the hygroscopic growth
of urban aerosol (50-350 nm) in Beijing from ~~the~~ long-term measurement. However, there are no direct measurements focusing
on the size range larger than 350 nm.

210 In this study, we deployed an HTDMA instrument to our laboratory in Beijing to conduct aerosol hygroscopicity measurements
from 27 November 2019 to 14 January 2020. Other particle microphysical properties, optical properties, and meteorology
parameters were also monitored simultaneously. Three questions need to be answered here: 1) what are the basic characteristics
of aerosol hygroscopicity over ~~an~~ extended size range in the urban environment? 2) how the hygroscopic properties vary with
pollution conditions, and 3) how the variation influences the visibility degradation? In the following, section 2 will ~~give a~~
215 ~~description of~~ describe the data and methods used in this study; section 3 will present the measurement results, including overall

aerosol hygroscopicity and its daily variations; the hygroscopic variation under different pollution conditions and potential impacts will be shown in section 4, and the last section comes the conclusions.

2. Data and methods

2.1 measurement site

220 The measurement was conducted in the aerosol laboratory, which is located on the rooftop of a six-floor building ~~on~~ the
campus of Peking University. ~~It shares the same location with the AERONET station of BEIJING_PKU~~ (39° 59' N, 116° 18'
E). The sampling site is in the northwest of Beijing, surrounded by schools, residential buildings and shopping centers.
Moreover, two main streets: Zhongguancun North Street is located to the west and Chengfu Road is to the south. Except for
the road traffic, there are no large emission sources in the neighboring. Aerosol particles here are representative of the urban
225 environments. More details can be found in Zhao et al. (2018).

In the measurement, a PM10 impactor was used to remove aerosol particles with aerodynamic diameters larger than 10 μm .
Then a dryer was used to decrease the RH to less than 30%. Next, the dried poly-disperse particles were guided into a splitter
with different instruments located ~~on the~~ downstream. From 27 November 2019 to 14 January 2020, aerosol number size
230 distributions, size-resolved hygroscopicity, and optical properties including scattering and absorption coefficients were
measured. Meteorological parameters like temperature, wind speed, wind direction, and RH were also monitored ~~at~~ this site.

2.2 Instrumentation and data

A BMI Humidified Tandem Differential Mobility Analyser (BMI HTDMA, Model 3100) was mainly used in this study. The
detailed instrument description and performance evaluation ~~has~~ been introduced in the previous publication (Lopez-Yglesias
235 et al., 2014). Similar to other conventional HTDMA systems, this instrument mainly incorporates an upstream Differential
Mobility Analyzer (DMA1), a humidification system that offers humidified sheath air to the humidifier and downstream DMA
to expose the particles to a prescribed RH, and a downstream DMA (DMA2) in series with a Mixing Condensation Particle
Counter (BMI MCPC, Model 1720) to measure the particles' size distribution after water uptake. Unique features of this BMI
HTDMA include a diffusion-based particle humidifier, a DMA design allowing selection of particles up to 2 μm diameter at
240 only 5600 volts, and the ability to study the complete deliquescence and efflorescence cycle. It also offers ~~a~~ processing software
to do the corrections and data inversion, which transforms the raw measured counts to the particle's growth factor distribution.

During the measurement, the RH of the second DMA was set to be 85%. The dry diameters selected by DMA1 were 50 nm,
100 nm, 200 nm, 300 nm, 400 nm, 500 nm, 600 nm. ~~In the calibration of~~ ~~To calibrate~~ the measurement system, ammonium

245 sulfate particles were also tested to compare with the theoretical values. The calibration includes both the dry test and the RH
test. The temperature and RH in the humidifier and DMA2 sheath were recorded in real-time for later check of system stability.
For the best working performance, the room was air-conditioned at 25 °C and circulated all the time.

The hygroscopic diameter growth factor (GF) is the ratio of the particle wet diameter at a given RH to the dry diameter, D_0 :
250
$$GF = \frac{Dp(RH)}{D_0} \quad (1)$$

Growth factor probability distribution function (GF-PDF, $c(GF)$) is normalized to unity. In order to simplify parameterization
and better compare the aerosol hygroscopic properties among different measurements, the hygroscopicity parameter κ was
also calculated in this study (Petters and Kreidenweis, 2007). It can be calculated from the following equation:

255
$$\kappa = (GF^3 - 1) \left(\frac{1}{RH} \exp\left(\frac{4\sigma_{s/a}M_w}{RT\rho_w D_0 GF}\right) - 1 \right) \quad (2)$$

where $\sigma_{s/a}$ is the droplet surface tension, M_w is the molecular weight of water, ρ_w is the density of liquid water, R is the
universal gas constant, and T is the absolute temperature.

The κ -PDF can also be derived from the GF-PDF. The volume-weighted mean growth factor is defined as the 3rd-moment
260 mean values of the GF-PDF: $GF_{mean} = (GF^3 \cdot c(GF)dGF)^{1/3}$, and it can be directly used to calculate the total liquid water
content of all particles at a specific size. The corresponding mean hygroscopic parameter κ_{mean} can be calculated from GF_{mean}
using the equation (2) above. To better understand the mixing state of particles and group them in terms of hygroscopic
properties, we classify the particles into a less-hygroscopic group with GF lower than 1.2, and a more-hygroscopic group with
GF larger than 1.2. For each group, the number fraction, GF_{mean} , and standard deviation σ can be determined from the GF-
265 PDF.

Apart from HTDMA, the aerosol light scattering coefficients at three wavelengths (450 nm, ~~52550~~ nm, and ~~635700~~ nm) were
measured using an Aurora 3000 nephelometer-TSI 3563 nephelometer (Anderson and Ogren, 1998). The particle number size
distribution (PNSD) was measured simultaneously by a BMI scanning electrical mobility sizer (BMI SEMS, Model 2100) and
270 a TSI scanning mobility particle sizer (SMPS). The BMI SEMS measures PNSD with a time resolution of 10 min over the size
range of 10-1000 nm. The SMPS has a time resolution of 5 min and covers the size from 10 to 600 nm. The black carbon (BC)
mass concentrations were measured using an MA200 (Aethlabs, serial number 0083) at 1 min temporal resolution.

3. Results and Discussions

3.1 Overview of GF and κ distribution

275 The aerosol number size distributions during the whole measurement period are summarized in Fig. 1(a). We also summarize
some previous studies on urban aerosol hygroscopicity from HTDMA measurements and present statistical results in Fig. 1(c).
The reference can be found in Table 1. We can clearly see that in Beijing winter, the peak contribution to the surface area is
200-300 nm and to the mass is 300-400 nm. For optical light extinction coefficients at 550 nm, the peak diameter is around
400-500 nm. However, the focuses of previous HTDMA measurements are only around 50-300 nm. It's this dislocation that
280 prompts us to conduct hygroscopicity measurements over an extended size range and investigate the potential impacts on
aerosol optical properties.

Figure 2 gives an overview of the average GF-PDF and the corresponding κ -PDF during the whole measurement period. It
can be seen that the mean growth factor distributions for all sizes show a distinct bi-modal pattern, consisting of a less-
285 hygroscopic mode and a more-hygroscopic mode. For particles of 50 nm, the LH and MH modes cannot be clearly
distinguished from each other. In general, particles of this size are not readily hygroscopic. For the size range between 100 nm
and 400 nm, there exists a dominant MH group and a minor but distinct LH group. The peak of the MH group shifts
significantly from about 1.2-1.4 to 1.4-1.6 as the size increases, indicating that the water-soluble or water-uptake materials in
larger sizes have a relatively stronger hygroscopicity than the smaller sizes. In contrast, the peak of the LH group barely shifts
290 slightly. The number fraction of the LH group decreases with size when particles are smaller than 200 nm, whereas it increases
with size when the size exceeds 200 nm. The spread (stand deviation) of the LH group has the same trend. For the particles
larger than 400 nm, the LH group dominates the number fraction, resulting in the decrease of bulk hygroscopicity of large
particles. Similar to the GF-PDF, κ -PDF also presents two modes corresponding to the LH and MH groups.

295 In general, our results are consistent with Wang et al. (2018) for the overlap region of 50-350 nm. We can also see a stronger
fraction of the nearly-hydrophobic group with size in lower and moderate pollution conditions in their study. This trend is
captured by our results and further extended to a larger size above 350 nm. Massling et al. (2009) also reported at the similar
trend of increasing hydrophobic particles with size in winter. They also gave the size-resolved chemical composition across
the submicron size range. For particles larger than 300 nm, there is an increased mass ratio of Elemental Carbon and undefined
300 mass, which may correspond to this increased group of particles. The size-resolved κ in Liu et al. (2014) is are different from
our results in the overlap region because their measurements were done in summer, in which the emissions and particle
photochemistry are quite different from those in winter. An apparent difference is the missing of the nearly-hydrophobic
group in accumulation mode particles, which is the main cause of the decreased hygroscopicity above 300 nm in our study.

305 The statistics of the HTDMA measurement are summarized in Table 2. The number fraction, mean GF and mean κ , spread of
GF and κ for the LH and MH groups during the whole measurement period are calculated. Corresponding standard deviations
are also presented in the table. The mean GF values for 50 nm, 100 nm, 200 nm, 300 nm, 400 nm, 500 nm, 600 nm are
1.13 \pm 0.07, 1.25 \pm 0.06, 1.31 \pm 0.07, 1.30 \pm 0.08, 1.24 \pm 0.12, 1.16 \pm 0.14, 1.12 \pm 0.16, respectively. The number fraction of LH group
310 particles reaches a trough at 200 nm and then increases with size. The ensemble mean κ peaks at 300 nm and then decreases
with size, which is related to the increase of LH group number fraction.

3.2. Time series and diurnal variation

The time series of aerosol size distribution and growth factor distribution over different sizes are shown in Fig. 3. The black
line in panel (a) is the time-dependent volume concentrations calculated from particle number size distribution. This value can
generally represent the particle pollution level. As elucidated by Guo et al. (2014), the PM episodes in Beijing present a
315 periodic cycle governed by meteorological conditions. This kind of cycle lasts from 3 days to 1 week, causing different degrees
of pollution. Our measurement captured about ~~six~~6 pollution cycles. Because the sampling was done at a fixed site, so the
complete cycle may go interrupted by transport or local physical or chemical processes. Nevertheless, the measured aerosol
properties from 27 November 2019 to 14 January 2020 can still be representative of the typical winter conditions in Beijing.
The blank area in the figure denotes missing data, which is caused by the instrument shutdown or measurement calibration.

320

From Fig. 3(a), we can see that the particle number concentration usually has an abrupt increase at the beginning of a pollution
cycle. Then the volume or mass concentration begins to develop and increase to a high level. During the clean period, the wind
is often dominated by strong northwest or northeast winds because it can bring unpolluted air masses from the northern
mountainous area. Polluted days are often accompanied by a weak southerly wind. Furthermore, high relative humidity tends
325 to appear at the end of each pollution cycle, which may help to push the pollution level to a peak (Fig. 8).

For each pollution cycle, we can see from Fig. 3 (b-c) that the GF-PDF of 50 nm and 100 nm particles does not have large
variations. But for other sizes, an increase of growth factor along with the pollution development can be clearly seen from the
GF-PDF contour. Different sizes have different lag time to respond to the evolution of pollution: the larger the size, the more
330 time needed to change the hygroscopic properties.

The average diurnal variations of GF-PDF for all the sizes are presented in Fig. 4. For better comparison, we mark the line
between daytime and nighttime. During the measurement period, the time of sunrise was about 7.00 a.m. and the sunset was
around 5.00 p.m. For particles of 50 nm, their hygroscopicity starts to decrease around sunrise and begins to increase around
335 sunset. So the hygroscopicity parameter κ peaks at 6.00 a.m. and reaches a minimum at 19.00 p.m. For particles larger than
200 nm, the hygroscopicity at daytime ~~isare~~ about 10-50% larger than that of nighttime.

For accumulation sizes, the number fraction of the LH group ($n_{f_{LH}}$) in the daytime is smaller than that in the nighttime. On one hand, the cause could be that more hydrophobic particles are generated or emitted in the night, and these particles include fresh BC or organic aerosols (Ma et al., 2011); on the other hand, more MH group particles could be formed in the daytime, or those aged particles in the night residual layer are mixed in the planetary boundary layer again (Liu et al., 2014). No diurnal variation is found in the peak of the LH group, indicating ~~the~~ similar chemical compositions in the LH group. In contrast, the peak hygroscopic parameter κ for the MH group varied significantly with time of a day, indicating that the chemical compositions in the MH group varied greatly at different day time. It usually reaches a maximum around noon and starts to decrease in the afternoon. The black lines in Fig. 4(a-g) are the ensemble mean κ for each size. For particles larger than 50 nm, this value also peaks around noon, which may be the result of decreased $n_{f_{LH}}$ and increased MH hygroscopicity. More statistical results about daytime and nighttime can be found in Table 3.

4. Evolution of Large Particle Hygroscopicity and Effects

4.1 Overview of GF and κ distribution

Guo et al. (2014) reported that particle pollution in Beijing is characterized by two distinct aerosol formation processes of nucleation and growth. During different aerosol formation processes, aerosol chemical compositions and corresponding hygroscopic properties may vary greatly. In order to investigate the evolution of particle hygroscopicity with different pollution levels, we group the data into three periods: 1) clean period in which the volume concentration is lower than $30 \mu\text{m}^3/\text{cm}^3$; (2) transition period in which the particle volume concentration is larger than 30 but smaller than $40 \mu\text{m}^3/\text{cm}^3$; (Fierce et al.) polluted period in which the volume concentration exceeds $40 \mu\text{m}^3/\text{cm}^3$.

Fig.5 shows the GF-PDF and average size distribution under different pollution levels. From the clean to the transition period, the most apparent change is in the number concentration. The number concentrations of both the LH and MH mode particles have increased significantly. At the same time, the number fraction of the MH group has also been enhanced. When the air quality develops from transition to polluted level, the number concentration remains similar, but the MH group is greatly enhanced, not only in the number fraction but also in ~~the~~ terms of the size range. The MH mode grows to the size of 600 nm, greatly squeezing the fraction of nearly-hydrophobic mode particles. From the statistical results in Table 4, the number fraction of less hygroscopic mode for 600 nm is 0.86 ± 0.22 , 0.75 ± 0.29 , 0.39 ± 0.27 for the clean, transition and pollution stage, respectively. For smaller particles, the variation of this value is not so distinct. Taking particles of 100 nm as an example, the $n_{f_{LH}}$ for clean, transition and pollution stage is 0.36 ± 0.14 , 0.33 ± 0.15 , 0.30 ± 0.13 , respectively. The difference of $n_{f_{LH}}$ among different stages increases with size.

It needs to be noted that the great decrease of nf_{LH} for large particles in the pollution period doesn't necessarily mean the decrease of absolute number concentration of LH mode particles. From Fig .5(f), the number concentration of LH mode particles remains similar to the transition stage. However, the total number concentration and corresponding mass concentration of large particles increases. These particles are grown from those smaller particles and are more aged in the environment. These processes are also called secondary aerosol formation. This part of particles contributes to the increase of the MH mode in a larger size and correspondingly total large particle number concentration. With the evolution of pollution levels, this process will continue. Zheng et al. (2016) showed that the fraction of the secondary components in PM_{2.5} are enhanced with increasing PM_{2.5} mass concentration. Our measurement results give ~~an~~ another direct evidence supporting this conclusion.

For each size, we summarize the average hygroscopicity under different pollution conditions (Fig. 6). The results show that for particles smaller than 300 nm, the hygroscopicity variation is not very large. But for particles larger than 300 nm, particle hygroscopicity can vary significantly with different pollution levels, which correspond to different aging stages of urban aerosol particles. In conclusion, when urban aerosols in Beijing go through a series of aging processes, their hygroscopicity is enhanced, especially for larger accumulation particles.

4.2 The effects on visibility degradation

To better understand the meteorological, microphysical and hygroscopic evolutions during the aging processes, and the effects of hygroscopicity variation on visibility degradation, a 3-day pollution period from 27 November to 1 December 2019 is selected for the analysis. Fig. 7 shows the overall situation of this pollution event. From the noon of 27 November, the wind speed began to decrease to below 3 m/s. The synoptic situation of Beijing remained stagnant until the end of this pollution period, with an average wind speed of 1.04 m/s. This meteorological situation indicated that the local emissions were the primary contribution of the surface air pollution and the aging processes were not influenced too much ~~by~~~~from~~ transport from other areas. In this respect, our measurement can represent the temporal evolution of aerosols in this urban region.

We can see from Fig. 7(b-c) that in the first 24 hours, there was a considerable increase of aerosol number concentration for particles smaller than 100 nm, which corresponded to the aerosol nucleation. The visibility dropped down from ~~the~~ upper limit to around 30 km. In this stage, aerosol hygroscopicity was relatively low. In the transition stage, the mean particle size grew continuously and the aerosol volume concentration also had a stable increase. At the same time, the aerosol became more hygroscopic in the accumulation size range between 100-300 nm. During this period, the visibility decreased slightly to 20 km. In the polluted stage, under the combined influence ~~of~~~~from~~ the high relative humidity and large hygroscopicity enhancement, the visibility decreased considerably to below 10 km, causing the most severe haze event, even though the aerosol volume concentration remained similar to that in ~~the~~ transition stage.

In this haze formation process, RH, aerosol loading (AL) and hygroscopicity all contribute to ~~the~~ visibility degradation. In order to quantify how much these factors took effect, we summarize the mean values of these factors in different stages, as shown in Fig.8(a-c). We can see that as the pollution developed, the number concentration of particles larger than 200 nm increased while the small particles (<100 nm) decreased. The aerosol hygroscopicity was enhanced greatly, from a value below 0.2 to above 0.4 for particles larger than 300 nm. At the same time, the mean relative humidity also increased from 25% to 77%.

The visibility is closely related to aerosol light extinction (σ) through Koschmieder relation (Griffing, 1980; Husar et al., 2000; Carrico, 2003):

$$\sigma_{ex}(km^{-1}) = \frac{3}{VIS(km)} \quad \text{(Fierce et al.)}$$

In this study, the particles are assumed to be spherical and their size-dependent extinction behavior is computed from the Mie model using particle dry size distribution, refractive indices, size-resolved hygroscopicity, and ambient RH. For each particle, we can determine its dry diameter and its wet diameter after hygroscopic growth. The refractive index (m) for the wet particle is classified between the BC core and coating materials, and m(coating) is computed as a volume-weighted average of water and dry non-absorbing materials. Then the light extinction for one particle can be calculated. The total aerosol light extinction can be integrated over all sizes given a measured PNSD. The detailed calculation description can be found in Chen et al. (2012). From this method, we can not only quantify the contribution from each size bin, but can also determine the role each factor (AL, RH and κ) played in the low visibility haze event. Fig. 8(d) compares the size-resolved aerosol light extinction coefficient during the clean period (~~blue~~-solid blue line) and the pollution period (~~black~~-solid black line). The blue line is considered as the base-line, which corresponds to low aerosol loading, low hygroscopicity and low RH values. When each of these factors is considered, we can obtain the following size-resolved light extinction, as shown in dotted lines. The magenta dotted line represents the condition with AL effect considered, which is calculated from the PNSD in pollution stage but with the κ and RH in the clean stage. The red dotted line represents the condition when both AL and RH are considered. The shading between lines is the difference, indicating the contribution to the aerosol light extinction from each considered factor.

From Fig. 8, we can see that particles larger than 300 nm contribute more than half of the total light extinction. Under this condition, if the hygroscopicity properties and variations in this size range are not known, it would be hard to fully quantify and predict the visibility impairment caused by aerosol particles in the urban environment. As we have seen from Fig.6 and Fig. 8(b), the hygroscopicity variation is very large in this size range. From our calculation (pie chart in Fig. 8(d)), it would bring about 40% of the total difference between the clean and pollution period, which is similar to the contribution of added aerosol loading. It's to be noted that all these three factors are coupled together to influence the haze formation, and the hygroscopicity variation would amplify the effect of added aerosol loading and increased ambient RH.

5. Conclusions

435 Submicron particles larger than 300 nm dominate the aerosol light extinction and mass concentration, and constitute a large part of the aerosol surface concentration. Under the exposure to high RH, all these parameters will be enhanced with the addition of water, and then greatly influence the particle mass, visibility degradation, particle chemistry and haze formation. However, field measurements on aerosol hygroscopicity in previous studies are focused below the size of 350 nm. For larger particles, detailed descriptions of hygroscopicity characteristics and variations are scarce. Some studies tried to derive the
440 hygroscopicity of a larger size range from other indirect methods like chemical composition or based on assumptions that accumulation mode particles share the same hygroscopicity. Till now, no direct evidence has been provided to support these assumptions or parameterization schemes.

In this study, a comprehensive aerosol field measurement focusing on hygroscopicity properties and variations over an
445 extended size range (50-600 nm) was conducted at a Beijing urban site in winter 2019. During the measurement, an HTDMA instrument was employed to measure hygroscopic growth factors of particles with dry diameters of 50, 100, 200, 300, 400, 500, and 600 nm at 85% RH from 27 November 2019 to 14 January 2020. This in-situ field measurement of atmospheric aerosols is representative of urban environment and of great importance to better understand the frequent low visibility events over North China Plain.

450

The measurement results show that the mean growth factor (GF) values for the sizes above were 1.13 ± 0.07 , 1.25 ± 0.06 , 1.31 ± 0.07 , 1.30 ± 0.09 , 1.24 ± 0.13 , 1.16 ± 0.15 , 1.12 ± 0.15 , respectively. The average GF-PDF presents a bi-modal structure including a less hygroscopic group and a more hygroscopic group. For the particles larger than 300 nm, there is a large fraction of nearly-hydrophobic or less hygroscopic particles, which decreases the bulk hygroscopicity in this size range. However,
455 during the polluted episode when the aerosol particles are fully aged, the ensemble mean hygroscopicity of larger particles will be enhanced significantly because of the growth and expansion of the MH group to this size range. Our measurements give a direct evidence supporting that the variation of the MH group brings the biggest change and uncertainty to the bulk hygroscopicity of particles larger than 300 nm. Our results are consistent with previous HTDMA studies in the overlap region (Massling et al., 2009; Wang et al., 2018; Liu et al., 2011a) and the chemical composition studies during the pollution
460 development (Guo et al., 2014; Zheng et al., 2016; Wu et al., 2018)

The significant variation of larger particle hygroscopicity will contribute much to the variation of particle optical properties like light extinction coefficient. Size-resolved light extinction coefficients under different pollution conditions are calculated using Mie scattering theory. From our results, when coupled with increased RH, the influence of κ variation on light
465 degradation could compete with the added aerosol loading in the haze development. Based on these results, we strongly

recommend the simultaneous measurement of hygroscopic properties over a size range larger than 300 nm in the future urban aerosol field studies.

Competing interests. The authors declare that they have no conflict of interest.

470 **Data availability.** The data used in this study can be obtained from this link:

<https://pan.baidu.com/s/1Sv9Zi3SjBf0vhMRH2NNRA>. It is also available when requesting the authors.

Author contributions. Chuanyang Shen, Gang Zhao, Weilun, Zhao and Chunsheng Zhao discussed the results; Ping Tian offered his help in the experiment; Chuanyang Shen wrote the manuscript.


475 **Acknowledgements.** This work is supported by the National Key Research and Development Program of China (2016YFC020000: Task 5) and the National Natural Science Foundation of China (41590872).

References


- 480 ALBRECHT, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227-1230, 10.1126/science.245.4923.1227, 1989.
- Anderson, T. L., and Ogren, J. A.: Determining aerosol radiative properties using the TSI 3563 integrating nephelometer, *Aerosol Science and Technology*, 29, 57-69, 1998.
- Bian, Y. X., Zhao, C. S., Ma, N., Chen, J., and Xu, W. Y.: A study of aerosol liquid water content based on hygroscopicity measurements at high relative humidity in the North China Plain, *Atmospheric Chemistry and Physics*, 14, 6417-6426, 10.5194/acp-14-6417-2014, 2014.
- 485 Carrico, C. M., Rood, M. J., Ogren, J. A., Neus üß C., Wiedensohler, A., and Heintzenberg, J.: Aerosol Optical properties at Sagres, Portugal during ACE-2, *Tellus B*, 52, 694-715, 10.1034/j.1600-0889.2000.00049.x, 2000.
- Carrico, C. M.: Mixtures of pollution, dust, sea salt, and volcanic aerosol during ACE-Asia: Radiative properties as a function of relative humidity, *Journal of Geophysical Research*, 108, 10.1029/2003jd003405, 2003.
- 490 Chang, D., Song, Y., and Liu, B.: Visibility trends in six megacities in China 1973–2007, *Atmos Res*, 94, 161-167, 10.1016/j.atmosres.2009.05.006, 2009.
- CHARLSON, R. J., SCHWARTZ, S. E., HALES, J. M., CESS, R. D., COAKLEY, J. A., HANSEN, J. E., and HOFMANN, D. J.: Climate Forcing by Anthropogenic Aerosols, *Science*, 255, 423-430, 10.1126/science.255.5043.423, 1992.
- Chen, J., Zhao, C. S., Ma, N., Liu, P. F., Göbel, T., Hallbauer, E., Deng, Z. Z., Ran, L., Xu, W. Y., Liang, Z., Liu, H. J., Yan, P., Zhou, X. J., and Wiedensohler, A.: A parameterization of low visibilities for hazy days in the North China Plain, *Atmospheric Chemistry and Physics*, 495 12, 4935-4950, 10.5194/acp-12-4935-2012, 2012.
- Chen, J., Zhao, C. S., Ma, N., and Yan, P.: Aerosol hygroscopicity parameter derived from the light scattering enhancement factor measurements in the North China Plain, *Atmospheric Chemistry and Physics*, 14, 8105-8118, 10.5194/acp-14-8105-2014, 2014.
- Deng, Z. Z., Zhao, C. S., Ma, N., Liu, P. F., Ran, L., Xu, W. Y., Chen, J., Liang, Z., Liang, S., Huang, M. Y., Ma, X. C., Zhang, Q., Quan, J. N., Yan, P., Henning, S., Mildnerberger, K., Sommerhage, E., Schafer, M., Stratmann, F., and Wiedensohler, A.: Size-resolved and bulk activation properties of aerosols in the North China Plain, *Atmospheric Chemistry and Physics*, 500 11, 3835-3846, 10.5194/acp-11-3835-2011, 2011.
- Deng, Z. Z., Zhao, C. S., Ma, N., Ran, L., Zhou, G. Q., Lu, D. R., and Zhou, X. J.: An examination of parameterizations for the CCN number concentration based on in situ measurements of aerosol activation properties in the North China Plain, *Atmospheric Chemistry and Physics*, 13, 6227-6237, 10.5194/acp-13-6227-2013, 2013.
- 505 Fierce, L., Onasch, T. B., Cappa, C. D., Mazzoleni, C., China, S., Bhandari, J., Davidovits, P., Fischer, D. A., Helgestad, T., Lambe, A. T., Sedlacek, A. J., 3rd, Smith, G. D., and Wolff, L.: Radiative absorption enhancements by black carbon controlled by particle-to-particle heterogeneity in composition, *Proc Natl Acad Sci U S A*, 117, 5196-5203, 10.1073/pnas.1919723117, 2020.
- Fitzgerald, J. W., Hoppel, W. A., and Vietti, M. A.: The Size and Scattering Coefficient of Urban Aerosol Particles at Washington, DC as a Function of Relative Humidity, *J Atmos Sci*, 39, 1838-1852, 10.1175/1520-0469(1982)039<1838:TSASCO>2.0.CO;2, 1982.

- 510 Griffing, G. W.: Relations between the prevailing visibility, nephelometer scattering coefficient and sunphotometer turbidity coefficient, *Atmospheric Environment* (1967), 14, 577-584, [https://doi.org/10.1016/0004-6981\(80\)90089-X](https://doi.org/10.1016/0004-6981(80)90089-X), 1980.
- Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., Du, Z., Wu, Z., Shao, M., Zeng, L., Molina, M. J., and Zhang, R.: Elucidating severe urban haze formation in China, *Proc Natl Acad Sci U S A*, 111, 17373-17378, 10.1073/pnas.1419604111, 2014.
- Haywood, J., and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review, *Reviews of Geophysics*, 38, 513-543, 10.1029/1999rg000078, 2000.
- 515 Hersey, S. P., Craven, J. S., Schilling, K. A., Metcalf, A. R., Sorooshian, A., Chan, M. N., Flagan, R. C., and Seinfeld, J. H.: The Pasadena Aerosol Characterization Observatory (PACO): chemical and physical analysis of the Western Los Angeles basin aerosol, *Atmospheric Chemistry and Physics*, 11, 7417-7443, 10.5194/acp-11-7417-2011, 2011.
- ~~Heyder, J., Gebhart, J., Rudolf, G., Schiller, C. F., and Stahlhofen, W.: Deposition of particles in the human respiratory tract in the size range 0.005–15 µm, *J Aerosol Sci*, 17, 811–825, [https://doi.org/10.1016/0021-8502\(86\)90035-2](https://doi.org/10.1016/0021-8502(86)90035-2), 1986.~~
- 520 Husar, R. B., Husar, J. D., and Martin, L.: Distribution of continental surface aerosol extinction based on visual range data, *Atmospheric Environment*, 34, 5067-5078, [https://doi.org/10.1016/S1352-2310\(00\)00324-1](https://doi.org/10.1016/S1352-2310(00)00324-1), 2000.
- Ibald-Mulli, A., Wichmann, H. E., Kreyling, W., and Peters, A.: Epidemiological Evidence on Health Effects of Ultrafine Particles, *Journal of Aerosol Medicine*, 15, 189-201, 10.1089/089426802320282310, 2002.
- 525 Jiang, C., Wang, H., Zhao, T., Li, T., and Che, H.: Modeling study of PM_{2.5} pollutant transport across cities in China's Jing-Jin-Ji region during a severe haze episode in December 2013, *Atmospheric Chemistry and Physics*, 15, 5803-5814, 10.5194/acp-15-5803-2015, 2015.
- Kitamori, Y., Mochida, M., and Kawamura, K.: Assessment of the aerosol water content in urban atmospheric particles by the hygroscopic growth measurements in Sapporo, Japan, *Atmospheric Environment*, 43, 3416-3423, 10.1016/j.atmosenv.2009.03.037, 2009.
- Kreidenweis, S. M., and Asa-Awuku, A.: Aerosol Hygroscopicity: Particle Water Content and Its Role in Atmospheric Processes, 331-361, 10.1016/b978-0-08-095975-7.00418-6, 2014.
- 530 Kuang, Y., Zhao, C. S., Tao, J. C., and Ma, N.: Diurnal variations of aerosol optical properties in the North China Plain and their influences on the estimates of direct aerosol radiative effect, *Atmos. Chem. Phys.*, 15, 5761-5772, 10.5194/acp-15-5761-2015, 2015.
- Kuang, Y., Zhao, C., Tao, J., Bian, Y., Ma, N., and Zhao, G.: A novel method for deriving the aerosol hygroscopicity parameter based only on measurements from a humidified nephelometer system, *Atmospheric Chemistry and Physics*, 17, 6651-6662, 10.5194/acp-17-6651-2017, 2017.
- 535 Kuang, Y., Zhao, C. S., Zhao, G., Tao, J. C., Xu, W., Ma, N., and Bian, Y. X.: A novel method for calculating ambient aerosol liquid water content based on measurements of a humidified nephelometer system, *Atmos. Meas. Tech.*, 11, 2967-2982, 10.5194/amt-11-2967-2018, 2018.
- Liu, H. J., Zhao, C. S., Nekat, B., Ma, N., Wiedensohler, A., van Pinxteren, D., Spindler, G., Müller, K., and Herrmann, H.: Aerosol hygroscopicity derived from size-segregated chemical composition and its parameterization in the North China Plain, *Atmospheric Chemistry and Physics*, 14, 2525-2539, 10.5194/acp-14-2525-2014, 2014.
- 540 Liu, P. F., Zhao, C. S., Gobel, T., Hallbauer, E., Nowak, A., Ran, L., Xu, W. Y., Deng, Z. Z., Ma, N., Mildnerberger, K., Henning, S., Stratmann, F., and Wiedensohler, A.: Hygroscopic properties of aerosol particles at high relative humidity and their diurnal variations in the North China Plain, *Atmospheric Chemistry and Physics*, 11, 3479-3494, 10.5194/acp-11-3479-2011, 2011a.
- 545 Liu, P. F., Zhao, C. S., Göbel, T., Hallbauer, E., Nowak, A., Ran, L., Xu, W. Y., Deng, Z. Z., Ma, N., Mildnerberger, K., Henning, S., Stratmann, F., and Wiedensohler, A.: Hygroscopic properties of aerosol particles at high relative humidity and their diurnal variations in the North China Plain, *Atmospheric Chemistry and Physics*, 11, 3479-3494, 10.5194/acp-11-3479-2011, 2011b.
- Lohmann, U., and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715-737, 10.5194/acp-5-715-2005, 2005.
- 550 Lopez-Yglesias, X. F., Yeung, M. C., Dey, S. E., Brechtel, F. J., and Chan, C. K.: Performance Evaluation of the Brechtel Mfg. Humidified Tandem Differential Mobility Analyzer (BMI HTDMA) for Studying Hygroscopic Properties of Aerosol Particles, *Aerosol Science and Technology*, 48, 969-980, 10.1080/02786826.2014.952366, 2014.
- Lundgren, D. A., and Paulus, H. J.: The Mass Distribution of Large Atmospheric Particles, *Journal of the Air Pollution Control Association*, 25, 1227-1231, 10.1080/00022470.1975.10470200, 2012.
- 555 Ma, N., Zhao, C. S., Nowak, A., Muller, T., Pfeifer, S., Cheng, Y. F., Deng, Z. Z., Liu, P. F., Xu, W. Y., Ran, L., Yan, P., Gobel, T., Hallbauer, E., Mildnerberger, K., Henning, S., Yu, J., Chen, L. L., Zhou, X. J., Stratmann, F., and Wiedensohler, A.: Aerosol optical properties in the North China Plain during HaChi campaign: an in-situ optical closure study, *Atmospheric Chemistry and Physics*, 11, 5959-5973, 10.5194/acp-11-5959-2011, 2011.
- Ma, N., Zhao, C. S., Müller, T., Cheng, Y. F., Liu, P. F., Deng, Z. Z., Xu, W. Y., Ran, L., Nekat, B., van Pinxteren, D., Gnauk, T., Müller, K., Herrmann, H., Yan, P., Zhou, X. J., and Wiedensohler, A.: A new method to determine the mixing state of light absorbing carbonaceous using the measured aerosol optical properties and number size distributions, *Atmospheric Chemistry and Physics*, 12, 2381-2397, 10.5194/acp-12-2381-2012, 2012.
- 560 Ma, N., Zhao, C., Tao, J., Wu, Z., Kecorius, S., Wang, Z., Groß J., Liu, H., Bian, Y., Kuang, Y., Teich, M., Spindler, G., Müller, K., van Pinxteren, D., Herrmann, H., Hu, M., and Wiedensohler, A.: Variation of CCN activity during new particle formation events in the North China Plain, *Atmospheric Chemistry and Physics*, 16, 8593-8607, 10.5194/acp-16-8593-2016, 2016.

- 565 Martin, M., Chang, R. Y. W., Sierau, B., Sjogren, S., Swietlicki, E., Abbatt, J. P. D., Leck, C., and Lohmann, U.: Cloud condensation nuclei closure study on summer arctic aerosol, *Atmospheric Chemistry and Physics*, 11, 11335-11350, 10.5194/acp-11-11335-2011, 2011.
- Massling, A., Stock, M., Wehner, B., Wu, Z. J., Hu, M., Brüggemann, E., Gnauk, T., Herrmann, H., and Wiedensohler, A.: Size segregated water uptake of the urban submicrometer aerosol in Beijing, *Atmospheric Environment*, 43, 1578-1589, 10.1016/j.atmosenv.2008.06.003, 2009.
- 570 Ouimette, J. R., and Flagan, R. C.: The extinction coefficient of multicomponent aerosols, *Atmospheric Environment* (1967), 16, 2405-2419, [https://doi.org/10.1016/0004-6981\(82\)90131-7](https://doi.org/10.1016/0004-6981(82)90131-7), 1982.
- Park, K., Kim, J.-S., and Park, S. H.: Measurements of Hygroscopicity and Volatility of Atmospheric Ultrafine Particles during Ultrafine Particle Formation Events at Urban, Industrial, and Coastal Sites, *Environ. Sci. Technol.*, 43, 6710-6716, 10.1021/es900398q, 2009.
- Penner, J. E., Hegg, D., and Leaitch, R.: Unraveling the role of aerosols in climate change, *Environ. Sci. Technol.*, 35, 332A-340A, 10.1021/es0124414, 2001.
- 575 Peters, A., Wichmann, H. E., Tuch, T., Heinrich, J., and Heyder, J.: Respiratory effects are associated with the number of ultrafine particles, *American Journal of Respiratory and Critical Care Medicine*, 155, 1376-1383, 10.1164/ajrccm.155.4.9105082, 1997.
- Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmospheric Chemistry and Physics*, 7, 1961-1971, 2007.
- 580 Pinnick, R. G., Jennings, S. G., and Chylek, P.: Relationships between extinction, absorption, backscattering, and mass content of sulfuric acid aerosols, *Journal of Geophysical Research*, 85, 10.1029/JC085iC07p04059, 1980.
- Roberts, G. C., and Nenes, A.: A Continuous-Flow Streamwise Thermal-Gradient CCN Chamber for Atmospheric Measurements, *Aerosol Science and Technology*, 39, 206-221, 10.1080/027868290913988, 2005.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics-From Air Pollution to Climate Change*, 2nd ed.; , John Wiley & Sons: Hoboken, NJ, 2006.
- 585 Shen, C., Zhao, C., Ma, N., Tao, J., Zhao, G., Yu, Y., and Kuang, Y.: Method to Estimate Water Vapor Supersaturation in the Ambient Activation Process Using Aerosol and Droplet Measurement Data, *Journal of Geophysical Research: Atmospheres*, 123, 10.1029/2018jd028315, 2018.
- Sun, Y. L., Jiang, Q., Wang, Z. F., Fu, P. Q., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013, *J Geophys Res-Atmos*, 119, 4380-4398, 10.1002/2014jd021641, 2014.
- 590 Sverdrup, G. M.: Parametric measurement of submicron atmospheric aerosol size distributions, University of Minnesota., 1977.
- Swietlicki, E., Hansson, H. C., Hämeri, K., Svenningsson, B., Massling, A., McFiggans, G., McMurry, P. H., Pet ä ä T., Tunved, P., Gysel, M., Topping, D., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments—a review, *Tellus B: Chemical and Physical Meteorology*, 60, 432-469, 10.1111/j.1600-0889.2008.00350.x, 2008.
- 595 Swietlicki, E., Hansson, H. C., Hämeri, K., Svenningsson, B., Massling, A., McFiggans, G., McMurry, P. H., Pet ä ä T., Tunved, P., Gysel, M., Topping, D., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments—a review, *Tellus B: Chemical and Physical Meteorology*, 60, 432-469, 10.1111/j.1600-0889.2008.00350.x, 2017.
- 600 Tao, J., Zhao, C., Ma, N., and Kuang, Y.: Consistency and applicability of parameterization schemes for the size-resolved aerosol activation ratio based on field measurements in the North China Plain, *Atmospheric Environment*, 173, 316-324, <https://doi.org/10.1016/j.atmosenv.2017.11.021>, 2018.
- Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment* (1967), 8, 1251-1256, [http://dx.doi.org/10.1016/0004-6981\(74\)90004-3](http://dx.doi.org/10.1016/0004-6981(74)90004-3), 1974.
- 605 Wang, Y., Wu, Z., Ma, N., Wu, Y., Zeng, L., Zhao, C., and Wiedensohler, A.: Statistical analysis and parameterization of the hygroscopic growth of the sub-micrometer urban background aerosol in Beijing, *Atmospheric Environment*, 175, 184-191, 10.1016/j.atmosenv.2017.12.003, 2018.
- Wu, Z., Wang, Y., Tan, T., Zhu, Y., Li, M., Shang, D., Wang, H., Lu, K., Guo, S., Zeng, L., and Zhang, Y.: Aerosol Liquid Water Driven by Anthropogenic Inorganic Salts: Implying Its Key Role in Haze Formation over the North China Plain, *Environmental Science & Technology Letters*, 5, 160-166, 10.1021/acs.estlett.8b00021, 2018.
- 610 Yan, P., Pan, X., Tang, J., Zhou, X., Zhang, R., and Zeng, L.: Hygroscopic growth of aerosol scattering coefficient: A comparative analysis between urban and suburban sites at winter in Beijing, *Particology*, 7, 52-60, <https://doi.org/10.1016/j.partic.2008.11.009>, 2009.
- Ye, X. X., Song, Y., Cai, X. H., and Zhang, H. S.: Study on the synoptic flow patterns and boundary layer process of the severe haze events over the North China Plain in January 2013, *Atmospheric Environment*, 124, 129-145, 10.1016/j.atmosenv.2015.06.011, 2016.
- 615 Zhao, G., Zhao, C., Kuang, Y., Bian, Y., Tao, J., Shen, C., and Yu, Y.: Calculating the aerosol asymmetry factor based on measurements from the humidified nephelometer system, *Atmospheric Chemistry and Physics*, 18, 9049-9060, 10.5194/acp-18-9049-2018, 2018.
- Zheng, J., Hu, M., Peng, J., Wu, Z., Kumar, P., Li, M., Wang, Y., and Guo, S.: Spatial distributions and chemical properties of PM_{2.5} based on 21 field campaigns at 17 sites in China, *Chemosphere*, 159, 480-487, <https://doi.org/10.1016/j.chemosphere.2016.06.032>, 2016.
- Zieger, P., Fierz-Schmidhauser, R., Gysel, M., Ström, J., Henne, S., Yttri, K. E., Baltensperger, U., and Weingartner, E.: Effects of relative humidity on aerosol light scattering in the Arctic, *Atmos. Chem. Phys.*, 10, 3875-3890, 10.5194/acp-10-3875-2010, 2010.
- 620



625 **Figure 24:** Frequency distribution of (a) aerosol number size distribution during the whole measurement period, (b) corresponding volume size distribution and (c) size-resolved light extinction coefficient contribution calculated from Mie theory. The red lines in (a-c) represent the mean values. The lines with marks in Fig. 1(c) denote the hygroscopicity measurement by HTDMA in previous and our studies.



630

Figure 2: Averaged probability distribution functions of (a) growth factor and (b) hygroscopicity parameter κ . The black lines in these two figures represent the ensemble mean values.

635






Figure 3: Temporal evolutions of (a) particle number size distribution and integrated volume concentration (black line), (b-h) growth factor probability distributions over different sizes from 50 nm to 600 nm. The colour contour in (a) denotes the particle number


640 concentration ($dn/d\log D_p$ (cm^{-3})), in (b-h) denotes the growth factor probability distribution function. Y-axis of (a) is particle diameter in log-scale.



645 **Figure 4: Diurnal variations of aerosol hygroscopicity distribution function over different sizes from 50 nm to 600 nm. Panel (h) shows the mean diurnal temperature and relative humidity. The black lines in (a-g) are the mean hygroscopicity calculated from the average κ -PDF. The vertical dotted lines in all the figures denote the time of sunrise (~7.00 a.m.) and sunset (~5.00 p.m.).**





650 **Figure 5:** (a-c) are the growth factor probability distribution under different pollution conditions. (d-f) are the corresponding mean number size distribution for LH and MH mode. For comparison, we also mark the mean number size distribution under transition stage in dashed lines in panel (f).



655

Figure 6: Mean size-resolved κ under different pollution conditions. The upper and lower limit of semi-transparent patch is the 25% and 75% percentile.






660

Figure 7: A pollution cycle from 11/27/2019 to 12/01/2019. (a) is the temporal evolution of temperature (blue line) and RH (orange line); (b) is the time series of visibility (black line) and wind speed. The length of the arrow indicates the wind speed, and the direction of the arrow indicates the wind direction. colour of the points denotes the direction of wind. (c) stands for the time series of particle number size distribution and the black line is the integrated volume concentration. Left axis is the log scale of particle diameters and right axis is for the volume concentration. (d) is the size-resolved hygroscopicity parameter κ .

665



670 **Figure 8:** (a) is the mean PNSD under different pollution conditions. (b) is the mean size-resolved κ under different pollution conditions. (c) is the mean RH under different conditions. (d) shows the effects of different factors (aerosol loading, RH and κ variation) to the size-resolved light extinction. The areas of different colors are shown in the pie chart on the left.

675 **Table 1:** Summary of previous HTDMA measurement of urban aerosol particles.

<i>Reference</i>	<i>Site</i>	<i>time</i>	<i>Size range (nm)</i>
<i>Zhang et al. (1993)</i>	Claremont, California, USA	Summer 1987	50-500
<i>Tschiersch et al. (1997)</i>	Neuherberg, Germany	Jan-Feb 1997	50-300
<i>Baltensperger et al. (2002)</i>	Milano (IT)	June-July 1998	20-200
<i>Ferron et al.(2005)</i>	Bavaria, Germany	Feb-Mar 1998	50-250
<i>Cocker et al. (2001)</i>	Pasadena, California (Husar et al.)	Aug–Sept 1999	50-150
<i>Massling et al. (2005)</i>	Leipzig, Germany	May–Aug 2000	50-150
<i>Mochida et al. (2006)</i>	Tokyo, Japan	Nov 2000	80-150

<i>Chen et al. (2003)</i>	Taipei, Taiwan	Oct–Dec 2001	53-202
<i>Zieger et al. (2014)</i>	Central Europe	Feb-Mar 2009	50-265
<i>Liu et al, (2011)</i>	Wuqing, China	7.17-8.12 2009	50-250
<i>Fors et al. (2011)</i>	South Sweden	May 2008-July 2010	35-265
<i>Holmgren et al. (2014)</i>	Central France	Sep 2008-Dec 2012	25-165
<i>Y Wang et al. (2018)</i>	Beijing, China	May 2014-Jan 2015	50-350

680

Table 2: Summary of the HTDMA measurements of hygroscopic growth at 85% RH.

<i>Size (nm)</i>	<i>50</i>	<i>100</i>	<i>200</i>	<i>300</i>	<i>400</i>	<i>500</i>	<i>600</i>
<i>Number of scans</i>	1654	1680	1673	1699	1710	1738	1732
<i>Less-Hygroscopic(GF<1.2)</i>							
<i>Number fraction</i>	0.70±0.19	0.35±0.14	0.29±0.16	0.36±0.20	0.50±0.27	0.66±0.30	0.78±0.29
<i>Mean GF</i>	1.07±0.03	1.08±0.03	1.09±0.04	1.09±0.05	1.07±0.05	1.05±0.05	1.05±0.05
<i>Spread of GF</i>	0.08±0.01	0.07±0.01	0.07±0.02	0.08±0.02	0.08±0.02	0.08±0.02	0.09±0.02
<i>Mean κ</i>	0.05±0.02	0.06±0.02	0.06±0.03	0.06±0.03	0.05±0.03	0.04±0.03	0.03±0.03
<i>Spread of κ</i>	0.059±0.0	0.050±0.0	0.046±0.0	0.049±0.0	0.051±0.0	0.051±0.0	0.051±0.0
	07	08	12	12	12	12	12
<i>More-Hygroscopic(GF>1.2)</i>							
<i>Number fraction</i>	0.30±0.19	0.65±0.14	0.71±0.16	0.64±0.20	0.50±0.27	0.34±0.30	0.22±0.29
<i>Mean GF</i>	1.29±0.04	1.37±0.04	1.44±0.05	1.46±0.05	1.45±0.07	1.43±0.10	1.43±0.12
<i>Spread of GF</i>	0.05±0.02	0.09±0.02	0.10±0.01	0.11±0.02	0.10±0.03	0.08±0.04	0.07±0.04
<i>Mean κ</i>	0.25±0.05	0.31±0.05	0.38±0.05	0.40±0.06	0.38±0.09	0.35±0.12	0.36±0.15
<i>Spread of κ</i>	0.06±0.03	0.10±0.02	0.12±0.02	0.13±0.03	0.11±0.04	0.10±0.05	0.09±0.06
<i>Ensemble of all groups</i>							
<i>Mean GF</i>	1.13±0.07	1.25±0.06	1.31±0.07	1.30±0.09	1.24±0.13	1.16±0.15	1.12±0.15
<i>Mean κ</i>	0.12±0.06	0.23±0.06	0.31±0.07	0.31±0.10	0.25±0.13	0.18±0.15	0.15±0.15

Table 3. Diurnal variations of aerosol hygroscopic properties for different sizes.

<i>Size (nm)</i>	<i>50</i>	<i>100</i>	<i>200</i>	<i>300</i>	<i>400</i>	<i>500</i>	<i>600</i>
<i>Peak (h)</i>	6	10	13	12	13	13	13

<i>Bottom (h)</i>	19	18	19	18	2	0	20
Winter Daytime							
κ_{mean}	0.11±0.07	0.24±0.06	0.33±0.06	0.34±0.09	0.27±0.14	0.20±0.15	0.18±0.17
$nf_{LH-group}$	0.72±0.20	0.31±0.13	0.24±0.13	0.31±0.18	0.45±0.27	0.62±0.31	0.73±0.32
$\kappa_{LH-group}$	0.05±0.02	0.06±0.02	0.07±0.03	0.06±0.03	0.05±0.03	0.04±0.04	0.04±0.04
$\kappa_{MH-group}$	0.24±0.05	0.31±0.05	0.38±0.05	0.41±0.06	0.39±0.10	0.37±0.14	0.37±0.15
Winter Nighttime							
κ_{mean}	0.12±0.06	0.22±0.06	0.30±0.08	0.30±0.10	0.23±0.13	0.17±0.14	0.14±0.14
$nf_{LH-group}$	0.68±0.18	0.37±0.15	0.32±0.17	0.39±0.21	0.53±0.28	0.69±0.29	0.80±0.27
$\kappa_{LH-group}$	0.05±0.02	0.05±0.02	0.06±0.03	0.06±0.03	0.04±0.03	0.04±0.03	0.03±0.03
$\kappa_{MH-group}$	0.26±0.05	0.31±0.05	0.37±0.05	0.39±0.07	0.37±0.08	0.35±0.11	0.35±0.15

Table 4. Statistical results of aerosol hygroscopic properties under different pollution conditions.

<i>Size (nm)</i>	50	100	200	300	400	500	600
Clean							
nf_{LH}	0.68±0.20	0.36±0.14	0.33±0.16	0.41±0.20	0.58±0.26	0.75±0.26	0.86±0.22
<i>Mean GF</i>	1.13±0.07	1.25±0.06	1.30±0.07	1.28±0.09	1.21±0.13	1.23±0.13	1.08±0.12
<i>Mean κ</i>	0.12±0.07	0.23±0.06	0.29±0.07	0.29±0.10	0.22±0.13	0.15±0.14	0.12±0.14
Transition							
nf_{LH}	0.73±0.17	0.33±0.15	0.21±0.14	0.29±0.17	0.43±0.25	0.61±0.28	0.75±0.29
<i>Mean GF</i>	1.12±0.07	1.26±0.06	1.34±0.06	1.33±0.07	1.27±0.11	1.19±0.13	1.13±0.14
<i>Mean κ</i>	0.11±0.06	0.24±0.06	0.34±0.07	0.34±0.09	0.28±0.11	0.20±0.13	0.16±0.14
Pollution							
nf_{LH}	0.74±0.16	0.30±0.13	0.14±0.11	0.16±0.12	0.21±0.16	0.31±0.22	0.39±0.27
<i>Mean GF</i>	1.11±0.06	1.27±0.05	1.36±0.05	1.38±0.06	1.37±0.07	1.33±0.10	1.30±0.13
<i>Mean κ</i>	0.10±0.05	0.25±0.05	0.36±0.06	0.40±0.07	0.38±0.08	0.34±0.10	0.32±0.14