

Supplement to “Emissions of non-methane volatile organic compounds from combustion of domestic fuels in Delhi, India”

Gareth J. Stewart¹, W. Joe F. Acton^{2,a}, Beth S. Nelson¹, Adam R. Vaughan¹, James R. Hopkins^{1,3}, Rahul Arya^{4,5}, Arnab Mondal^{4,5}, Ritu Jangirh^{4,5}, Sakshi Ahlawat^{4,5}, Lokesh Yadav^{4,5}, Sudhir K. Sharma^{4,5}, Rachel E. Dunmore¹, Siti S. M. Yunus⁶, C. Nicholas Hewitt², Eiko Nemitz⁷, Neil Mullinger⁷, Ranu Gadi⁸, Lokesh. K. Sahu⁹, Nidhi Tripathi⁹, Andrew R. Rickard^{1,3}, James D Lee^{1,3}, Tuhin K. Mandal^{4,5} and Jacqueline F. Hamilton¹.

¹ Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, YO10 5DD, UK

² Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK

³ National Centre for Atmospheric Science, University of York, York, YO10 5DD, UK

⁴ CSIR-National Physical Laboratory, Dr. K.S. Krishnan Marg, New Delhi, Delhi 110012, India

⁵ Academy of Scientific & Innovative Research, Ghaziabad, Uttar Pradesh- 201 002, India

⁶ School of Water, Environment and Energy, Cranfield University, Cranfield, MK43 0AL, UK

⁷ UK Centre for Ecology and Hydrology, Penicuik, EH26 0QB, UK


⁸ Indira Gandhi Delhi Technical University for Women, Kashmiri Gate, New Delhi, Delhi 110006, India

⁹ Physical Research Laboratory (PRL), Ahmedabad 380009, India

^a Now at: School of Geography, Earth and Environmental Sciences, University of Birmingham, B15 2TT, Birmingham, UK

20 **S1 - Breakthrough test**

21



22 Figure S1. Results of the GCxGC-FID breakthrough test. Left: Benzene. Right: *n*-octane.

23

24 **S2 – Emission factors**

25 Please see the attached excel file for the emission factors measured in this study.

26

S3 – Calculation of total volume of air sampled

NMVOC mass in $\mu\text{g m}^{-3}$ was determined from:

$$\text{NMVOC}_{\text{mass}} = \frac{\text{NMVOC}_{\text{ppbv}} \times 12.187 \times \text{MW}}{273.15 + T}$$

where $\text{NMVOC}_{\text{mass}}$ = NMVOC concentration in $\mu\text{g m}^{-3}$, $\text{NMVOC}_{\text{ppbv}}$ = NMVOC concentration in ppbv, MW = molecular mass in g mol^{-1} and T = temperature in $^{\circ}\text{C}$ which was measured for each experiment.

The total volume of air convectively moving up the stack was determined from:

$$V_d = \sqrt{\frac{2gP_s}{D_s}}$$

where V_d = vertical displacement up the flue (ms^{-1}), $g = 9.81 \text{ m}^{-2}$, P_s = average stack pressure (mmH_2O) and D_s is determined by:

$$D_s = \frac{TD_a}{T_s}$$

where T = ambient temperature (k), D_a = density of air (1.1455 kg m^{-3}), T_s = average stack temperature ($^{\circ}\text{K}$). The emission factor (EF) was calculated by:

$$\text{EF} = \frac{tCV_dA_d}{M}$$

where t = time burned (s), C = concentration (g m^{-3}), A_d = area of flue, M = mass of fuel burnt (kg).

S4 – Calculation of total emission factor

Table S1 shows a summary of the instruments used to calculate the total emission factor from this study. The total signal from the PTR-ToF-MS had isotope peaks removed as well as reagent ion peaks (m/z 31 Th), water cluster peaks (m/z 37 Th), ammonia (m/z = 18.034 Th), acetylene (m/z = 26.015 Th), hydrogen cyanide (m/z = 28.019 Th), ethene (m/z = 28.03 Th), propene (m/z = 43.054 Th), nitrous acid (m/z = 48.008 Th), butadienes (m/z = 55.053 Th), butenes, other hydrocarbons (m/z = 57.069 Th), DMS (m/z = 63.032 Th), isoprene (m/z = 69.069 Th), pentenes, methylbutenes (m/z = 71.084 Th), benzene (m/z = 79.053 Th), toluene (m/z = 93.069 Th), C₂ substituted monoaromatics (m/z = 107.084 Th) and C₃ substituted monoaromatics (m/z = 121.099 Th). The emission factors for removed species have been incorporated from the DC-GC-FID and the GCxGC-FID.

Table S1. Summary of instruments used for calculation of total emission factor.

Species	Instrument
Aromatics	
Benzene-C ₃ substituted monoaromatics	GCxGC-FID
C ₄ -C ₅ substituted monoaromatics	PTR-ToF-MS
Phenolics	
All species	PTR-ToF-MS
Oxygenated aromatics	
All species	PTR-ToF-MS
Oxygenated aliphatics	
All species	PTR-ToF-MS
NVOCs	
All species	PTR-ToF-MS
Furanics	
All species	PTR-ToF-MS
Monoterpenes	
All species	PTR-ToF-MS ^a
PAHs	
All species	PTR-ToF-MS ^a
Alkenes	
All DC-GC-FID alkenes to C ₅	DC-GC-FID
Remaining alkenes ^b	PTR-ToF-MS
Alkanes ^c	

C ₂ -C ₆ alkanes	DC-GC-FID
C ₇ -C ₁₂ alkanes	GCxGC-FID
C ₁₃ -C ₂₀ alkanes	SPE-GCxGC-ToF-MS

58

59 ^a PTR-ToF-MS signal used as may include isomers not measured by GCxGC-FID.

60 ^b Only alkenes not measured by DC-GC-FID.

61 ^c No peaks for > C₆ alkanes in PTR-ToF-MS spectra.

62

63

64

65

66

67



69

70