

PM₁ composition and source apportionment at two sites in Delhi, India across multiple seasons

Ernesto Reyes-Villegas^{1,*}, Upasana Panda², Eoghan Darbyshire^{1,△}, James M. Cash^{3,4}, Rutambhara Joshi¹, Ben Langford³, Chiara F. Di Marco³, Neil J. Mullinger³, Mohammed S. Alam⁵, Leigh R. Crilley^{5,a} Daniel J. Rooney⁵, W. Joe F. Acton⁶, Will Drysdale⁷, Eiko Nemitz³, Michael Flynn¹, Aristeidis Voliotis¹, Gordon McFiggans^{1,*}, Hugh Coe¹, James Lee⁷, C. Nicholas Hewitt⁶, Mathew R. Heal⁴, Sachin S Gunthe^{2,8}, Tuhin K. Mandal⁹, Bhola R. Gurjar¹⁰, Shivani¹¹, Ranu Gadi¹¹, Siddhartha Singh¹², Vijay Soni¹², James D. Allan^{1,13,*}

¹Department of Earth and Environmental Sciences, University of Manchester, Manchester, M13 9PL, UK

¹⁰EWRE Division, Dept of Civil Engineering, Indian Institute of Technology Madras, Chennai, 600 036, India

³UK Centre for Ecology & Hydrology, Edinburgh Research Station, Penicuik, EH26 0QB, UK

⁴School of Chemistry, University of Edinburgh, Edinburgh, EH9 3FJ, UK

⁵School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK

⁶Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

¹⁵Wolfson Atmospheric Chemistry Laboratory, University of York, York, YO10 5DD, UK

⁸Laboratory for Atmospheric and Climate Sciences, Indian Institute of Technology Madras, Chennai-36, India

⁹CSIR-National Physical Laboratory, Dr. K.S. Krishnan Marg, New Delhi, Delhi 110012, India

¹⁰Dpt. of Civil Engineering, Indian Institute of Technology, Roorkee, India

¹¹Department of Applied Sciences and Humanities, Indira Gandhi Delhi Technical University for Women, Delhi, 110006, India

¹²India Meteorology Department, New Delhi, 110003, India

¹³National Centre for Atmospheric Science, University of Manchester, Manchester, M13 9PL, UK

[△] Now at: The conflict and Environment Observatory, Hebden Bridge, HX7 5HZ, UK

^a Now at: Department of Chemistry, York University, Toronto, ON, Canada

* Correspondence to: James Allan (James.Allan@manchester.ac.uk), Gordon McFiggans (g.mcfiggans@manchester.ac.uk) and Ernesto Reyes-Villegas (ernesto.reyesvillegas@manchester.ac.uk)

Abstract.

Air pollution in urban environments has been shown to have a negative impact on air quality and human health, particularly in megacities. Over recent decades, Delhi, India has suffered high atmospheric pollution, with significant particulate matter (PM) concentrations as result of anthropogenic activities. Organic aerosols (OA) are composed of thousands of different chemical species and are one of the main constituents of submicron particles. However, quantitative knowledge of OA composition, their sources and processes in urban environments is still limited. This is important particularly in India, as Delhi is a massive, inhomogeneous conurbation, which we would expect that the apportionment and concentrations will vary depending on where in Delhi the measurements/source apportionment is performed, indicating the need of multi-site measurements. This study presents the first multisite analysis carried out in India over different seasons, with a focus on identifying OA sources. The measurements were taken during 2018 at two sites in Delhi, India. One site was located at the

India Meteorological Department, New Delhi (ND). The other site was located at the Indira Gandhi Delhi Technical University for Women, Old Delhi (OD). Non-refractory submicron aerosol (NR-PM₁) concentrations (ammonium, nitrate, sulphate, 40 chloride and organic aerosols) of four aerosol mass spectrometers were analysed. Collocated measurements of VOC, black carbon, NO_x and CO were performed. Positive matrix factorization (PMF) analysis was performed to separate the organic fraction, identifying a number of conventional factors: hydrocarbon-like OA (HOA) related to traffic emissions, biomass burning OA (BBOA), cooking OA (COA) and secondary OA (SOA).

A composition-based estimate of PM₁ is defined by combining BC and NR-PM₁ (C-PM₁ = BC + NR-PM₁). No significant 45 difference was observed on C-PM₁ concentrations between sites; OD (142 +/- 117 $\mu\text{g.m}^{-3}$) compared to ND (123 +/- 71 $\mu\text{g.m}^{-3}$), from post-monsoon measurements. A wider variability was observed between seasons, where pre-monsoon and monsoon showed C-PM₁ concentrations lower than 60 $\mu\text{g.m}^{-3}$. A seasonal variation in C-PM₁ composition was observed; SO₄²⁻ showed a high contribution over pre-monsoon and monsoon seasons while NO₃⁻ and Cl⁻ had a higher contribution in winter and post-monsoon. The main primary aerosol source was from traffic, which is consistent with the PMF analysis and aethalometer 50 model analysis. Thus, in order to reduce PM₁ concentrations in Delhi through local emission controls traffic emissions control offers the greatest opportunity. PMF-AMS mass spectra will help to improve future aerosol source apportionment studies. The information generated in this study increases our understanding of PM₁ composition and OA sources in Delhi, India. Furthermore, the scientific findings provide significant information to strengthen legislation that aims to improve air quality in India.

55

Keywords: AMS, PMF, on-line measurements, air quality, aerosol sources, Urban air pollution.

60

65

70

1 Introduction

Air pollution in urban environments has been shown to have a negative impact on human health, particularly in megacities. The World Health Organization (WHO) stated that in 2016 about 90% of the global population living in urban environments was exposed to particulate matter concentrations exceeding the WHO air quality guidelines 80 (https://www.who.int/gho/phe/air_pollution_pm25_concentrations/en/, accessed 20/06/2019). According to the WHO global air quality database 2018, Delhi was ranked in 11th position of the cities with high particulate mass, measured as the mass of particles whose size is equal to or lower than 2.5 μm (PM_{2.5}), having an annual average concentration of 143 $\mu\text{g.m}^{-3}$. Other Indian cities are in the top 20 with Varanasi being in first place with a PM_{2.5} concentration of 217 $\mu\text{g.m}^{-3}$ 85 (<https://www.who.int/airpollution/data/cities/en/>, accessed 17/06/2019). Previous studies have identified ambient fine particle concentrations to have detrimental effects on health, particularly the submicron fraction lower than or equal to 1 μm (PM₁) (Pope et al., 2002; Ramgolam et al., 2009).

Organic aerosol (OA) constitute a large proportion of PM₁ mass, with fractions between 40-60% in urban environments (Zhang et al., 2007). OA is composed of thousands of different chemical species and is difficult to study due to a variety of sources 90 and processes. Black carbon (BC), another important component of PM₁, is a product of incomplete combustion and is recognised as one of the main climate-forcing components (Bond and Bergstrom, 2006; Lack et al., 2014). Moreover, the WHO, in the health effects of BC report, states that while there was no clear evidence of BC directly affecting human health, it may however work as a medium to transport a wide variety of chemical species of varying toxicities (Janssen et al., 2012). BC sources and concentrations have been studied in India (Thamban et al., 2017; Jain et al., 2018a). In 2014, Gupta et al. (2017) 95 studied the seasonal variation of BC in Agra, identifying biomass burning to be the major BC source, especially during winter. The BC mixing state was measured in a clean site (Nainital) and a polluted site (Gurgaon), with average BC concentrations of 1.0 and 11.0 $\mu\text{g.m}^{-3}$, respectively (Raatikainen et al., 2017). In 2016, a BC network in India was initiated with 16 Aethalometers (LASKAR et al., 2016); the availability of these instruments allowed the study of BC sources in different environments (Rajesh and Ramachandran, 2017; Kolhe et al., 2018; Nazeer Hussain et al., 2018).

Aerosol mass spectrometer (AMS) instruments (Aerodyne Research, Billerica, MA) have been widely used in different locations around the world to quantify online real-time non-refractory PM₁ (NR-PM₁): OA, nitrate (NO₃⁻), sulphate (SO₄²⁻), ammonium (NH₄⁺) and chloride (Cl⁻). In India, the first NR-PM₁ measurements were taken on campus at the Indian Institute

of Technology (IIT) in Kanpur in winter 2011, where OA was found to contribute 70% to NR-PM₁ concentrations.
105 (Chakraborty et al., 2015). Subsequently, a number of studies have been conducted at the same site showing dominance of the OA fraction with substantial contributions from secondary sources (Kumar et al., 2016) and looking at organo-nitrates and OA sources (Chakraborty et al., 2016;Chakraborty et al., 2018) December 2015 to January 2016, where NR-PM₁ measurements helped to analyse the temporal characteristics of brown carbon (Satish et al., 2017). Aircraft measurements during the pre-monsoon and monsoon periods in 2016, covering the NE Bay of Bengal and IGP regions, allowed the vertical and horizontal aerosol chemical composition to be studied (Brooks et al., 2019). Recent long-term measurements in Mahabaleshwar (Mukherjee et al., 2018), Bhubaneswar (Kompalli et al., 2020) and New Delhi (Gani et al., 2019) have allowed the study of NR-PM₁ seasonal variability.

Receptor modelling tools have been used in aerosol source apportionment studies in India, such as chemical mass balance,
115 principal component analysis and positive matrix factorization (PMF) (Tiwari et al., 2013;Bhuyan et al., 2018;Jain et al., 2018b;Gadi et al., 2019;Shivani et al., 2019;Jain et al., 2020). PMF has been shown to be a useful tool to determine OA sources from aerosol mass spectrometer measurements across the northern hemisphere (Ng et al., 2010). In India, source apportionment studies using PMF have identified different OA sources, for instance, seasonal analysis in 2016-2017 in Mahabaleshwar, a high altitude site, identified hydrocarbon-like OA (HOA), biomass burning OA (BBOA) and two different types of OOA as
120 the OA sources (Mukherjee et al., 2018;Singla et al., 2019). Similar OA factors were identified in a recent study performed in New Delhi, with the main OA sources being HOA, BBOA, and OOA from a 1 year analysis (Bhandari et al., 2019).

This work is part of the Atmospheric Pollution and Human Health in an Indian Megacity (APHH-India) programme
125 (<https://www.urbanair-india.org/>). The main objective of the programme is to support research on the sources and emissions of urban air pollution in New Delhi, India, the processes underlying and impacting on these, and how air pollution then impacts on health. In agreement with two of the working packages of the programme (DelhiFlux and PROMOTE), the objectives in this work are:

- 130 • To study the aerosol concentrations in high-time resolution in Old and New Delhi through the year in two different locations.
- To identify main OA sources and to determine the contribution of primary and secondary OA.
- To study the interaction of aerosol concentrations with local meteorology
- To provide information to be used by the modelling community for improvements in the emission inventories, air pollution forecast and source apportionment.

135

2 Methodology

140 2.1 Site description and meteorology

The measurements were taken during 2018 at two sites in Delhi, India. One site was located at the Indian Meteorological Department (IMD), Mausam Bhawan, Lodhi Road, New Delhi. Lat 28.588, Lon 77.217, hereafter ND. The other site was located at the Indira Gandhi Delhi Technical University for Women. (IGDTUW) at New Church Rd, Kashmere Gate, Old Delhi. Lat 28.664, Lon 77.232, hereafter OD. Section S1 in the Supplementary Material shows the site locations.

145

The meteorology in India is highly influenced by the monsoon season, which takes places between June and September and is characterised by an increase in precipitation (Turner et al., 2019). Recent long-term measurements in Mahabaleshwar (Mukherjee et al., 2018) and New Delhi (Gani et al., 2019) have identified a clear pattern over the pre monsoon, monsoon and post monsoon seasons. Airborne measurements have studied the monsoon progression across the Indo-Gangetic Plain region 150 and its impacts on aerosol characteristics (Brooks et al., 2019). In the present study, temperature showed a wide range of variability; winter was cool with an average temperature of 20 °C, whereas pre-monsoon, monsoon and post-monsoon seasons were characterised by average temperatures of 36, 31 and 25 °C, respectively (Figure 1). As expected, relative humidity (RH) showed the highest average over monsoon (76%) compared to the other seasons with RH lower than 60 %. Similar temperatures have been previously observed by Brooks et al. (2019) with 31 °C in pre-monsoon and 22° C in monsoon as well 155 as by Gani et al. (2019) with temperatures between 10-20 °C in winter, 25-40 °C in pre-monsoon and 25-35 °C in monsoon. Figure S3 in supplement shows box plots of temperature, wind speed and RH for the different seasons.

2.2 Instrumentation

Four Aerodyne Aerosol Mass Spectrometers were deployed across the different seasons in 2018 (Table 1). One Aerosol Chemical Speciation Monitor (ACSM) (Ng et al., 2011), one compact time-of-flight Aerosol Mass Spectrometer (cToF-AMS) 160 (Drewnick et al., 2005) and two High-Resolution Time-of-Flight AMS (HR-ToF-AMS) (DeCarlo et al., 2006). The principle of operation of aerosol mass spectrometer instruments has been widely explained in previous publications; in general, these instruments quantify real time concentrations of NR-PM₁ (OA, NO₃⁻, SO₄²⁻, NH₄⁺ and Cl⁻) by vaporising the aerosols at 600 °C with electron impact ionisation at 70 eV and final detection in the mass spectrometer. The main difference between the instruments used in this study is the mass spectrometer, with differences in the sensitivity and the mass to charge (m/z) 165 resolution. However, all are reliable instruments and successful intercomparisons with parallel measurements have been performed in previous studies (Ng et al., 2011; Crenn et al., 2015). In this study, the cToF-AMS and the HR-AMS agree to within 21% from parallel ambient measurements in OD during the pre-monsoon season (Figure S7). Previous comparisons of AMS instruments have shown agreement between instruments of between 19 and 50% and 35% is widely recommended as

the absolute accuracy of AMS instruments (DeCarlo et al., 2008; Dunlea et al., 2009; Bahreini et al., 2009; Crenn et al., 170 2015; Shinozuka et al., 2020). Section S2 shows the calibrations and quality assurance analysis performed to the AMS measurements. The measurements in PostM_OD_T and PostM_ND_T, were taken from a height of 32 m from a tower (T). The tower was used to perform eddy-covariance flux measurements, which will be reported elsewhere. The rest were ground-based measurements (~3 m high). All set ups included a PM_{2.5} cyclone to cut particle size and a drier to reduce humidity. The 175 AMS measurements during the Diwali festival (7th November) for the period PostM_ND_T_C were lost due to a power cut, thus this special episode was removed from the NR-PM₁ analysis.

Table 1. Measurement dates and instrumentation. Winter (Win), pre-monsoon (PreM), monsoon (Mon) and post-monsoon (PostM). New Delhi (ND) and Old Delhi (OD) sites. ACSM (A), two HR-ToF-AMS instruments were used (H1 and H2), cToF-AMS (C). Eddy-covariance flux measurements tower (T).


| Instrument | Event | start | end | Days | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|------------|---------------|-------|-------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| ACSM | Win_ND_A | 05/02 | 03/03 | 26 | | | | | | | | | | | |
| HR-AMS_1 | PreM_ND_H1 | 24/04 | 30/05 | 36 | | | | | | | | | | | |
| cToF-AMS | PreM_OD_C | 28/05 | 09/06 | 12 | | | | | | | | | | | |
| HR-AMS_2 | PreM_OD_H2 | 26/05 | 28/06 | 33 | | | | | | | | | | | |
| HR-AMS_2 | Mon_OD_H2 | 03/08 | 19/08 | 16 | | | | | | | | | | | |
| HR-AMS_2 | PostM_OD_H2 | 09/10 | 04/11 | 26 | | | | | | | | | | | |
| cToF-AMS | PostM_OD_C | 11/10 | 15/10 | 4 | | | | | | | | | | | |
| HR-AMS_2 | PostM_OD_T_H2 | 06/11 | 20/11 | 18 | | | | | | | | | | | |
| cToF-AMS | PostM_ND_T_C | 06/11 | 20/11 | 32 | | | | | | | | | | | |

180

Other instruments used along with the mass spectrometers include: two Aethalometers (Magee Scientific; one model AE31 and one model AE33) and one Multi-Angle Absorption Photometer (MAAP, Thermo Fisher Scientific) (Petzold et al., 2002). Table 1 shows details about the instrument locations and sampling periods and table S1 presents the collocated instruments 185 with the mass spectrometers. Meteorology data, apart from Planetary Boundary Layer Height (PBLH), was downloaded from https://ncdc.noaa.gov/ (last access: 05/01/2019) at 1-hourly resolution. PBLH data sourced from ECMWF ERA5 0.25 deg. results in 1-hour resolution.

PMF (Paatero et al., 2002) was used to identify potential sources of OA, based on the Source Finder tool, SoFi 4.8 (Canonaco et al., 2013). Section S3 in supplement provides detailed information about the criteria to select the OA factors. The PMF 190 analysis of the OD_H2 measurements presented in this work was performed, by Cash et al. (2020) for the complete dataset (26/05 – 20/11) using high-resolution organic mass spectra. The PMF analysis to the PreM_ND_H1 dataset was performed to unit mass resolution data. BC concentrations measured with the Aethalometer AE-31 were corrected following the Weingartner method (Weingartner et al., 2003) and using the SP-2 as a reference BC measurement. The aethalometer model was applied, following the Sandradewi method (Sandradewi et al., 2008), to aethalometer measurements, both AE-31 and AE-33, to identify

195 the contribution of biomass burning and fossil fuel to BC concentrations. Section S4 in the Supplement shows the BC data processing.



200 **Figure 1. Box plots with (a) planetary boundary layer height (PBLH), (b) temperature, (c) relative humidity and (d) wind speed for the different seasons. The marker represents the mean.**

3. Results

3.1 Aerosol concentrations and composition

205 Aerosol measurements were taken in OD and ND over different seasons in 2018. The highest concentrations were observed in the post-monsoon season, where OA concentrations of $400 \mu\text{g.m}^{-3}$ were found (Fig. S8). Lower aerosol concentrations were observed in the monsoon season, where OA reached between $80 - 200 \mu\text{g.m}^{-3}$. Similar to Gani et al. (2019), we combine BC and NR-PM₁ concentrations to develop a composition based estimate of PM₁ ($\text{C-PM}_1 = \text{BC} + \text{NR-PM}_1$). The season with the highest averaged C-PM₁ concentrations were observed in PostM-OD, with average concentrations ($\pm \text{SD}$) of $142 \pm 117 \mu\text{g.m}^{-3}$ in PostM_OD_T_H2 (Fig. 2.a). Lower aerosol concentrations were observed in pre-monsoon and seasons with C-PM₁ average concentrations lower than $40 \mu\text{g.m}^{-3}$ in PreM_ND_H1, and Mon_OD_H2, with OA being the component with the leading contributor to PM₁.

Measurements during post-monsoon were taken in OD (PostM_OD_T_H2) and ND (PostM_ND_T_C) over the same time (6-20 Nov). This comparison shows average C-PM₁ concentrations in OD were slightly higher (142 +/- 117 $\mu\text{g.m}^{-3}$) compared to ND (123 +/- 71 $\mu\text{g.m}^{-3}$). It is worth mentioning that, while removing the Diwali Festival as a special event, the measurements shown in this work still include high concentrations from festivities that took place before and after the Diwali Festival on November 7th. The ND site is located in affluent residential green area, where the transit of heavy good vehicles is controlled; thus, it could be expected that the use of fireworks is also restricted here, whereas in OD, there is less control of the firework activities. While no significant differences were observed when comparing between sites, a clear difference between seasons was observed; pre-monsoon and monsoon with similar C-PM₁ concentrations lower than 60 $\mu\text{g.m}^{-3}$, while post-monsoon with C-PM₁ concentrations higher than 120 $\mu\text{g.m}^{-3}$. Detailed statistical analysis for NR-PM₁ and BC is presented in Table S3.




Figure 2. Average aerosol concentrations (a) and relative contribution to C-PM1 (b).

225 OA makes the main relative contribution to PM₁ with values of 0.45-0.62 followed by BC with values of 0.13-0.24 (Fig. 2.b). This high relative contribution of OA has been observed in previous studies in India (Bhandari et al., 2019;Singla et al., 2019). The high BC contributions of around 18%, suggest an important contribution from primary sources. Sulphate made a higher contribution in pre-monsoon and monsoon while NO_3^- and Cl^- had a higher contribution in winter and post-monsoon, similar to a previous study performed in Delhi (Gani et al., 2019).

230

This wide range of concentrations is related to the meteorological conditions (Fig. 1), mainly with boundary layer height as a result of changes in temperature; winter and PostM-T had a temperature of 20-25° C, resulting in wind speeds of 2.5-3.0 m.s^{-1} , which may have resulted in a more stable mixing layer, accumulating aerosol concentrations. In early pre-monsoon, an average temperature of 35° C with wind speeds of 3.5 m.s^{-1} facilitated greater aerosol dispersion. From the diurnal cycles (Fig. 235 3), it can be seen that NO_3^- and Cl^- showed high concentrations during the morning while BC and OA showed high concentrations at night. SO_4^{2-} showed less diurnal variability, suggesting a regional origin with large sources at midday when

the photochemistry takes place compensating for the boundary layer effect. BC is an aerosol constituent related to primary emissions, fossil fuel (mainly traffic) and biomass burning (Sandradewi et al., 2008), hence the fact that OA had a diurnal pattern similar to BC during the post-monsoon season suggests OA was potentially dominated by primary sources; while in the pre-monsoon season, OA showed an increase in concentrations during the day, suggesting a larger contribution from secondary OA (Shrivastava et al., 2017). This primary-secondary origin of OA will be analysed in the next section.




Figure 3. Median diurnal concentrations of the aerosol chemical components

245 3.2 Source apportionment of the organic aerosol

Organic aerosol is composed of thousands of chemical species, making source attribution challenging. Methods using receptor modelling tools such as PMF are often employed to deconvolve OA into various OA factors that, depending on their composition and temporal variations, can be attributed to potential OA sources. In this work a number of factors were identified with PMF analysis (Fig. 4); three common factors were found in all the seasons, hydrocarbon-like OA (HOA), biomass burning 250 OA (BBOA) and more oxidized oxygenated OA (MO-OOA). Other identified sources were cooking OA (COA), less oxidized oxygenated OA (LO-OOA) and oxygenated primary OA (oPOA). HOA is related to fossil fuel emissions, mainly attributed to traffic, whose mass spectrum is characterised by peaks at mass to charge ratio (m/z) 55 and 57, and is dominated by the $C_xH_y^+$ family (Zhang et al., 2005). BBOA is characterised by m/z 60 ($C_2H_4O^+$) which is related to anhydrosugar fragments, such as levoglucosan (Alfarra et al., 2007). The mass spectrum of COA presents peaks at m/z 55 and m/z 57, similar to the HOA 255 spectrum but with a lower peak at m/z 57 (Allan et al., 2010; Mohr et al., 2012). The two oxygenated OA (OOA) factors

identified MO-OOA and LO-OOA. MO-OOA, with a dominant peak at m/z 44 (CO_2^+), and LO-OOA, typically with a peak at m/z 44 and a larger peak at m/z 43 (mostly C_2H_3^+) compared to MO-OOA (Zhang et al., 2011).

The HOA mass spectrum was consistent across the collection periods and between the different instruments deployed, with
260 peaks at m/z 41, 43, 55 and 57, and its diurnal cycle showed high concentrations in the morning and at night resulting from
| the traffic rush hour. Moreover, HOA showed its highest concentrations during periods of low wind speeds (Fig. S197) and
shallow boundary-layers, highlighting the importance of HOA as a primary local source: even a small local source emitting
265 into a shallow nocturnal boundary-layer can result in significant accumulation of concentrations. The MO-OOA mass spectrum
is also consistent over all the seasons with peaks at m/z 18 and m/z 44, its diurnal cycle shows the highest concentrations to
have been around 10:00 to 15:00 hrs, suggesting being related to the stronger radiation occurring at that time, resulting in
secondary aerosol production from photochemistry. We can observe larger variability in the BBOA mass spectra, with changes
270 to the peak at m/z 60, which has a higher relative intensity over the post-monsoon and winter seasons. oPOA, which shows
peaks at m/z at 43 and m/z 44, has a characteristic diurnal pattern with high concentrations around 09:00 hrs, while the mass
spectrum resembles that of OOA, the diurnal cycle behaved different to those of LO-OOA and MO-OOA. A more detailed
275 analysis about its possible origin will be provided in the next section. COA was identified over per-monsoon and post-
monsoon, but not in PostM_ND_T or during winter. Moreover, previous studies in urban environments have identified a lunch
peak in the diurnal plots (Allan et al., 2010; Young et al., 2015). However, this peak was not clearly observed in any of the
seasons where COA was identified. Perhaps, the high OA concentrations resulting from the other anthropogenic emissions,
i.e. traffic and biomass burning along with the boundary layer effect, would favour the mixing of OA sources and PMF will
struggle to completely separate the OA sources (Lanz et al., 2008).





Figure 4. Mass spectra and diurnal plots of OA factors from PMF analysis to the different periods. Here the period all_OD_H2 is displayed to compare with the other periods. The period all_OD_H2, which shows the high resolution PMF analysis performed to the complete dataset of AMS (H2) 26th of May – 23rd November 2018, is analysed in detail by Cash et al. (2020) . Detail analysis of the all_OD_H2 dataset is presented in Table S4, where the PMF analysis identified 7 factors, we are adding HOA = HOA + nHOA and BBOA = SFOA + SVBBOA to compare with our 5-factor solutions.

4 Discussion

4.1 The role of meteorology on aerosol concentrations.

The meteorology in India has been summarised in Section 2.1, stating the impact of the monsoon season and the difference between pre-monsoon and post-monsoon. This work allowed the study of NR-PM₁ seasonal variability, which showed strong variations through the seasons (Figure 2), where a seasonal cycle was observed with high aerosol concentrations in post-monsoon and winter with low temperatures. From the previous diurnal plots in winter_ND (Fig. 3), high Cl⁻ concentrations were observed in the morning. PMF analysis performed on this dataset also identified a factor referred to as oPOA with a morning peak (Fig. 4.a). Figure 5 shows the diurnal trends of Cl⁻ and oPOA along with meteorology parameters. Both Cl⁻ and oPOA peak around 09:00 hrs, at the time when RH shows its highest value and their concentrations start decreasing when the PBLH starts increasing around 10:00 hrs. This decrease in Cl⁻ and oPOA concentrations is perhaps due to mixing with fresh air masses from the break-up of the boundary layer, moreover, it is likely that the Cl⁻ reduction is due to some repartitioning into the gas-phase as temperature increases. NH₄Cl has a higher vapour pressure than NH₄NO₃ and will dissociate first. This is also consistent with the decrease on Cl⁻ concentrations before NO³⁻ concentrations increase in the morning. The same high Cl⁻ concentrations were observed by Chakraborty et al. (2018) in NR-PM₁ measurements taken during Sep-Oct 2014 in Kanpur,

India where similar meteorological conditions were present. This suggests the impact RH, PBLH and temperature have on Cl^- and oPOA concentrations. Figure S2048 shows Cl^- and oPOA have different sources, it can be observed that Cl^- follows the trend of NH_4^+ , which is expected as the AMS is sensitive mainly to NH_4Cl but not to other Cl^- compounds, and oPOA follows the same pattern as SO_4^{2-} . It is possible that the properties of the condensed material in oPOA may be similar to the assumed NH_4Cl , even though the sources are different, i.e. both have a similar volatility. A more detailed analysis on the Cl^- processes is performed by Gunthe et al. (2021), where the ACSM_Winter data is part of the analysed datasets.



305 **Figure 5. Diurnal median values of meteorological parameters, Cl^- and oPOA concentrations in Winter_ND**

4.2 Organic aerosol sources and their contribution to NR- PM_1 .

Seasonal dependencies of the PMF factors can be investigated by looking at their average contributions (Fig. 6). HOA is the factor with the highest concentration with $6.0 - 55.0 \mu\text{g.m}^{-3}$ for Pre-monsoon and Post-monsoon, respectively, in the PostM_OD_H2 dataset (Table S4), HOA concentrations of $30 \mu\text{g.m}^{-3}$ were identified. These HOA concentrations represent a 310 relative contribution to OA sources of 20 to 50%. Traditionally, HOA has been related mainly to traffic emissions of vehicles using petrol or diesel (Zhang et al., 2005;Platt et al., 2017). However, in Delhi, compressed natural gas vehicles and generators might also contribute to HOA concentrations (Prakash et al., 2020). It is interesting to see that BBOA has low concentrations (2.0 – $8.0 \mu\text{g.m}^{-3}$) in the ND and pre-monsoon seasons with a low contribution to OA sources (0.10 – 0.15). However, Cash et al. (2020), in the PostM_OD_T_H2 analysis, identified a BBOA average concentration of $48.22 \mu\text{g.m}^{-3}$, representing a relative 315 contribution of 25%. COA is an important primary source in OD, with average concentrations of $2.0 - 12.0 \mu\text{g.m}^{-3}$. It is worth mentioning that COA concentrations might be overestimated due to a potential higher relative ionisation efficiency, which has been observed previously in a laboratory experiment (Reyes-Villegas et al., 2018). Considering HOA, BBOA and COA as primary OA (POA) and the rest of the sources as OOA and looking at the POA relative contribution to OA concentrations (Fig. 6.d), there is a more constant concentration in ND with values between 54 – 56%, while POA relative contributions in 320 OD range between 46 – 66%. These POA relative contributions close to 50% over different times of the year have been previously observed in a study performed in 2017 at IIT Delhi, New Delhi (Bhandari et al., 2019), with similar average

concentrations for POA and OOA (52 and 56 $\mu\text{g.m}^{-3}$ for winter as well as 30 and 31 $\mu\text{g.m}^{-3}$ for pre-monsoon, respectively). Cash et al. (2020) identified POA relative contributions of 45% and 52% for pre-monsoon and monsoon and 64% for post-monsoon, showing the importance of POA concentrations in OD during this season. OOA concentrations are produced by both POA and precursor gas. In other urban environments, it is possible to see a more variable POA-OOA ratio contribution to OA concentrations (Jimenez et al., 2009). For instance, in London, UK, in 2012 POA contributions ranged from 80% in winter to 45% in summer (Young et al., 2015), perhaps as a result of the lower temperatures during winter compared to summer along with less photochemistry in winter at higher altitudes. These findings demonstrate the impact primary PM_1 emissions in Delhi have on OA concentrations over the year.

330





Figure 6. OA average concentrations (6.a) and relative contribution (6.b). The whiskers represent the standard deviation of POA/OOA contribution after running PMF with different Fpeaks between -1 to 1 with steps of 0.2 to prove the stability of the separation.

335 4.3 fossil fuel and biomass burning sources of BC.

The high contribution of POA has been explained in Section 4.2 and more specifically, traffic was identified as an important OA source. This finding is supported by the aethalometer model analysis. This model based on the Sandradewi method (Sandradewi et al., 2008) uses the absorption coefficient (b_{abs}) and the absorption angstrom exponent to apportion the contribution of fossil fuel (b_{abs_950ff}) and biomass burning (b_{abs_470bb}) to Aethalometer measurements (refer to Section S4 in the 340 Supplement for details). Figure 7 shows the mean absorption coefficients for fossil fuel and biomass burning, where high b_{abs_950ff} values compared to b_{abs_470bb} are observed. Pre-monsoon and monsoon are the seasons with low biomass burning influence, with average values between $8 - 20 \text{ Mm}^{-1}$; during these seasons, fossil fuel showed b_{abs} values between 35 and 50 Mm^{-1} . In post-monsoon the influence of biomass burning starts to increase with average values around 50 Mm^{-1} . However, at 345 this time, traffic showed higher average values between 100 and 150 Mm^{-1} . In winter, biomass burning has an important influence, with an average of 80 Mm^{-1} , while fossil fuel still has a major influence with a value of 95 Mm^{-1} . It is only during the Diwali festival, from OD-PostMon (Fig. S1⁶⁴ where biomass burning has a significant contribution to BC (700 Mm^{-1}). However, during this period, there is an important contribution of BC from fossil fuel (500 Mm^{-1}), showing the importance of fossil fuel as a BC source.



350

Figure 7. Average absorption coefficients for fossil fuel (b_{abs_950ff}) and biomass burning (b_{abs_470bb}). The whiskers represent ± 1 standard deviation.

5. Conclusions

Aerosol composition was studied to compare Old Delhi (OD) and New Delhi (ND) over different seasons in 2018 with a focus on source apportionment. No significant difference was observed on C-PM₁ concentrations (defined as total AMS mass plus black carbon, BC) from post-monsoon measurements between sites; OD (142 +/- 117 $\mu\text{g.m}^{-3}$) compared to ND (123 +/- 71 $\mu\text{g.m}^{-3}$). A wider variability was observed between seasons, where pre-monsoon and monsoon showed average C-PM₁ concentrations lower than 60 $\mu\text{g.m}^{-3}$. The fact that there is no remarkable difference on C-PM₁ concentrations in the two sites, suggests that, when aiming to control C-PM₁ high concentrations, the actions should be region orientated, for example as Delhi region, rather than considering controlling air pollution in OD or ND only. OD presented high SO₄²⁻ concentrations (7.0 $\mu\text{g.m}^{-3}$ in OD and 3.4 $\mu\text{g.m}^{-3}$ in ND). In the post-monsoon and winter seasons, the C-PM₁ mass was dominated by organic matter (OA) rather than inorganic and the largest single contributor was hydrocarbon-like organic aerosol (HOA). NH₄⁺, NO₃⁻ and Cl⁻ peaked in the morning while OA peaked at night. A seasonal variation of PM₁ composition was observed; SO₄²⁻ showed a high contribution over pre-monsoon and monsoon seasons while (more volatile) NO₃⁻ and Cl⁻ made a larger contribution in the cooler seasons, winter and post-monsoon. The fact that no remarkable difference between sites was found suggests the air pollution in Delhi is a regional problem that is influenced by meteorological conditions over the year, being the post-monsoon and winter the seasons with highest pollutant concentrations.

Analysis of the organic mass spectra by positive matrix factorisation (PMF) identified a number of conventional factors: HOA, biomass burning OA (BBOA), cooking-like OA (COA), less oxidized oxygenated OA (LO-OOA) and more oxidized oxygenated OA (MO-OOA). One additional factor (oPOA) had a particular diurnal trend, similar to Cl⁻, and a mass spectral signature similar to OOA. However, from examination of the polar plots (Fig. S20), oPOA appears to have similar source sectors to SO₄²⁻. This suggests oPOA may be semi-volatile and driven by changes in T and RH, like Cl⁻, whilst having different sources, undetermined at this time. Traffic showed to be the main primary aerosol source for both OA and BC, as seen with the PMF analysis (HOA) and aethalometer model analysis (b_{abs_950ff}). The measurements indicate that in Delhi further control of primary traffic exhaust emissions would make a significant contribution to reducing PM₁ concentrations. Moreover, by controlling gas precursor emissions, OOA concentrations may also be reduced.

Primary organic aerosol (POA) made a relatively constant contribution at the different sites and seasons with a relative contribution around 50 % to OA. This shows both the importance of primary sources and the fact that daily average temperatures remain above 20 °C throughout the year; 20 °C in winter and 43 °C in pre-monsoon, compared, for example, with European urban environments where average temperatures can drop to 5 °C or lower in winter and summer temperatures are around 25-30 °C. This temperature might increase the regional pollutant transport allowing an even POA – SOA distribution over the Delhi region. When comparing these results with Cash et al. (2020) , it shows the importance POA has in OD during the post-monsoon season, where a POA relative contribution of 63% was observed. Meteorology played an important role in

aerosol concentrations. High concentrations of Cl^- and oPOA were observed in winter_ND which was characterised by high RH, with a sudden drop in concentrations when the boundary layer broke up in the morning and temperatures rose. This Cl^- peak has been observed in a previous post-monsoon study in Kanpur India (Chakraborty et al., 2018; Gadi et al., 2019). More detailed research should be done in order to identify the potential sources/processes that caused the peak in OA concentrations 390 around 9-10 am in urban Indian environments, which has been observed taking place during post-monsoon and winter.

The information generated in this study increases our understanding of PM_1 composition and OA sources in Delhi, India. The PMF-AMS mass spectra can be used in future source apportionment studies in India and perhaps other urban environments to improve the identification of sources and provides a unique dataset for the assessment of atmospheric chemistry and transport 395 models. Furthermore, the scientific findings provide significant information to target legislation that aim to improve air quality in India.

6. Data availability

AMS data used in this paper have been archived at

<https://catalogue.ceda.ac.uk/uuid/b7c9aeb6aee54698aef82f20365fc441>

400 Acknowledgements

This work was supported by the UK NERC through the DelhiFlux and PROMOTE projects under the Newton-Bhabha Fund Programme “Air Pollution and Human Health in a Developing Megacity (APHH-India)”. NERC reference numbers: NE/P016502/1, NE/P01643X/1, NE/P016472/1 and NE/P016480/1. The monsoon measurements were supported by the NERC 405 National Capability award SUNRISE (NE/R000131/1). James Cash is recipient of a NERC E³ DTP studentship (NE/L002558/1).

References

Alfarra, M. R., Prevot, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M., and Baltensperger, U.: Identification of the mass spectral signature of organic aerosols from wood burning emissions, Environmental Science & Technology, 41, 5770-5777, Doi 10.1021/Es062289b, 2007.

410 Allan, J. D., Williams, P. I., Morgan, W. T., Martin, C. L., Flynn, M. J., Lee, J., Nemitz, E., Phillips, G. J., Gallagher, M. W., and Coe, H.: Contributions from transport, solid fuel burning and cooking to primary organic aerosols in two UK cities, Atmos Chem Phys, 10, 647-668, 2010.

Bahreini, R., Ervens, B., Middlebrook, A. M., Warneke, C., de Gouw, J. A., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., Neuman, J. A., Ryerson, T. B., Stark, H., Atlas, E., Brioude, J., Fried, A., Holloway, J. S., Peischl, J., Richter, D., Walega, J., Weibring, P., Wollny, A. G., and Fehsenfeld, F. C.: Organic aerosol formation in urban and industrial plumes near Houston 415 and Dallas, Texas, Journal of Geophysical Research: Atmospheres, 114, <https://doi.org/10.1029/2008JD011493>, 2009.

Bhandari, S., Gani, S., Patel, K., Wang, D. S., Soni, P., Arub, Z., Habib, G., Apte, J. S., and Hildebrandt Ruiz, L.: Sources and atmospheric dynamics of organic aerosol in New Delhi, India: Insights from receptor modeling, *Atmos. Chem. Phys. Discuss.*, 2019, 1-33, 10.5194/acp-2019-403, 2019.

420 Bhuyan, P., Deka, P., Prakash, A., Balachandran, S., and Hoque, R. R.: Chemical characterization and source apportionment of aerosol over mid Brahmaputra Valley, India, *Environmental Pollution*, 234, 997-1010, 10.1016/j.envpol.2017.12.009, 2018.

Bond, T. C., and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Science and Technology*, 40, 27-67, 10.1080/02786820500421521, 2006.

425 Brooks, J., Allan, J. D., Williams, P. I., Liu, D., Fox, C., Haywood, J., Langridge, J. M., Highwood, E. J., Kompalli, S. K., O'Sullivan, D., Babu, S. S., Satheesh, S. K., Turner, A. G., and Coe, H.: Vertical and horizontal distribution of submicron aerosol chemical composition and physical characteristics across northern India during pre-monsoon and monsoon seasons, *Atmos. Chem. Phys.*, 19, 5615-5634, 10.5194/acp-19-5615-2019, 2019.

430 Canonaco, F., Crippa, M., Slowik, J. G., Baltensperger, U., and Prevot, A. S. H.: SoFi, an IGOR-based interface for the efficient use of the generalized multilinear engine (ME-2) for the source apportionment: ME-2 application to aerosol mass spectrometer data, *Atmos Meas Tech*, 6, 3649-3661, DOI 10.5194/amt-6-3649-2013, 2013.

Cash, J. M., Langford, B., Di Marco, C., Mullinger, N., Allan, J., Reyes-Villegas, E., Joshi, R., Heal, M. R., Acton, W. J. F., Hewitt, N., Misztal, P., Drysdale, W., Mandal, T. K., Shivani, Gadi, R., and Nemitz, E.: Seasonal analysis of submicron aerosol in Old Delhi using high resolution aerosol mass spectrometry: Chemical characterisation, source apportionment and new marker identification, *Atmos. Chem. Phys. Discuss.*, 2020, 1-42, 10.5194/acp-2020-1009, 2020.

435 Chakraborty, A., Bhattu, D., Gupta, T., Tripathi, S. N., and Canagaratna, M. R.: Real-time measurements of ambient aerosols in a polluted Indian city: Sources, characteristics, and processing of organic aerosols during foggy and nonfoggy periods, *Journal of Geophysical Research: Atmospheres*, 120, 9006-9019, 10.1002/2015jd023419, 2015.

Chakraborty, A., Gupta, T., and Tripathi, S. N.: Chemical composition and characteristics of ambient aerosols and rainwater residues during Indian summer monsoon: Insight from aerosol mass spectrometry, *Atmos Environ*, 136, 144-155, 10.1016/j.atmosenv.2016.04.024, 2016.

440 Chakraborty, A., Mandariya, A. K., Chakraborti, R., Gupta, T., and Tripathi, S. N.: Realtime chemical characterization of post monsoon organic aerosols in a polluted urban city: Sources, composition, and comparison with other seasons, *Environmental Pollution*, 232, 310-321, 10.1016/j.envpol.2017.09.079, 2018.

445 Crenn, V., Sciare, J., Croteau, P. L., Verlhac, S., Fröhlich, R., Belis, C. A., Aas, W., Äijälä, M., Alastuey, A., Artiñano, B., Baisnée, D., Bonnaire, N., Bressi, M., Canagaratna, M., Canonaco, F., Carbone, C., Cavalli, F., Coz, E., Cubison, M. J., Esser-Gietl, J. K., Green, D. C., Gros, V., Heikkinen, L., Herrmann, H., Lunder, C., Minguillón, M. C., Močnik, G., O'Dowd, C. D., Ovadnevaite, J., Petit, J. E., Petralia, E., Poulain, L., Priestman, M., Riffault, V., Ripoll, A., Sarda-Estève, R., Slowik, J. G., Setyan, A., Wiedensohler, A., Baltensperger, U., Prévôt, A. S. H., Jayne, J. T., and Favez, O.: ACTRIS ACSM intercomparison – Part 1: Reproducibility of concentration and fragment results from 13 individual Quadrupole Aerosol Chemical Speciation Monitors (Q-ACSM) and consistency with co-located instruments, *Atmos. Meas. Tech.*, 8, 5063-5087, 10.5194/amt-8-5063-2015, 2015.

450 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal Chem*, 78, 8281-8289, Doi 10.1021/Ac061249n, 2006.

455 DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crounse, J., Wennberg, P. O., Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D., Weinheimer, A. J., Montzka, D. D., Campos, T., and Jimenez, J. L.: Fast airborne aerosol size and chemistry measurements above Mexico City and Central Mexico during the MILAGRO campaign, *Atmos Chem Phys*, 8, 4027-4048, 2008.

Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., 460 Borrmann, S., and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) - Instrument description and first field deployment, *Aerosol Science and Technology*, 39, 637-658, Doi 10.1080/02786820500182040, 2005.

Dunlea, E. J., DeCarlo, P. F., Aiken, A. C., Kimmel, J. R., Peltier, R. E., Weber, R. J., Tomlinson, J., Collins, D. R., Shinozuka, Y., McNaughton, C. S., Howell, S. G., Clarke, A. D., Emmons, L. K., Apel, E. C., Pfister, G. G., van Donkelaar, A., Martin, R. V., Millet, D. B., Heald, C. L., and Jimenez, J. L.: Evolution of Asian aerosols during transpacific transport in INTEX-B, *Atmos. Chem. Phys.*, 9, 7257-7287, 10.5194/acp-9-7257-2009, 2009.

Gadi, R., Shivani, Sharma, S. K., and Mandal, T. K.: Source apportionment and health risk assessment of organic constituents in fine ambient aerosols (PM2.5): A complete year study over National Capital Region of India, *Chemosphere*, 221, 583-596, <https://doi.org/10.1016/j.chemosphere.2019.01.067>, 2019.

470 Gani, S., Bhandari, S., Seraj, S., Wang, D. S., Patel, K., Soni, P., Arub, Z., Habib, G., Hildebrandt Ruiz, L., and Apte, J. S.: Submicron aerosol composition in the world's most polluted megacity: the Delhi Aerosol Supersite study, *Atmos. Chem. Phys.*, 19, 6843-6859, 10.5194/acp-19-6843-2019, 2019.

475 Gunthe, S. S., Liu, P., Panda, U., Raj, S. S., Sharma, A., Darbyshire, E., Reyes-Villegas, E., Allan, J., Chen, Y., Wang, X., Song, S., Pöhlker, M. L., Shi, L., Wang, Y., Kommula, S. M., Liu, T., Ravikrishna, R., McFiggans, G., Mickley, L. J., Martin, S. T., Pöschl, U., Andreae, M. O., and Coe, H.: Enhanced aerosol particle growth sustained by high continental chlorine emission in India, *Nat Geosci.*, 14, 77-84, 10.1038/s41561-020-00677-x, 2021.

480 Gupta, P., Singh, S. P., Jangid, A., and Kumar, R.: Characterization of black carbon in the ambient air of Agra, India: Seasonal variation and meteorological influence, *Advances in Atmospheric Sciences*, 34, 1082-1094, 10.1007/s00376-017-6234-z, 2017.

Jain, C. D., Gadhavi, H. S., Wankhede, T., Kallelapu, K., Sudhesh, S., Das, L. N., Pai, R. U., and Jayaraman, A.: Spectral properties of black carbon produced during biomass burning, *Aerosol Air Qual Res.*, 18, 671-679, 10.4209/aaqr.2017.03.0102, 2018a.

Jain, S., Sharma, S. K., Mandal, T. K., and Saxena, M.: Source apportionment of PM10 in Delhi, India using PCA/APCS, UNMIX and PMF, *Particuology*, 37, 107-118, 10.1016/j.partic.2017.05.009, 2018b.

485 Jain, S., Sharma, S. K., Vijayan, N., and Mandal, T. K.: Seasonal characteristics of aerosols (PM2.5 and PM10) and their source apportionment using PMF: A four year study over Delhi, India, *Environmental Pollution*, 262, 114337, <https://doi.org/10.1016/j.envpol.2020.114337>, 2020.

Jimenez, J., Canagaratna, M., Donahue, N., Prevot, A., Zhang, Q., Kroll, J., DeCarlo, P., Allan, J., Coe, H., and Ng, N.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525-1529, 2009.

490 Kolhe, A. R., Aher, G. R., Ralegankar, S. D., and Safai, P. D.: Investigation of aerosol black carbon over semi-urban and urban locations in south-western India, *Atmospheric Pollution Research*, 10.1016/j.apr.2018.04.010, 2018.

Kompalli, S. K., Suresh Babu, S. N., Satheesh, S. K., Krishna Moorthy, K., Das, T., Boopathy, R., Liu, D., Darbyshire, E., Allan, J. D., Brooks, J., Flynn, M. J., and Coe, H.: Seasonal contrast in size distributions and mixing state of black carbon and its association with PM1.0 chemical composition from the eastern coast of India, *Atmos. Chem. Phys.*, 20, 3965-3985, 10.5194/acp-20-3965-2020, 2020.

495 Kumar, B., Chakraborty, A., Tripathi, S. N., and Bhattu, D.: Highly time resolved chemical characterization of submicron organic aerosols at a polluted urban location, *Environmental Science: Processes and Impacts*, 18, 1285-1296, 10.1039/c6em00392c, 2016.

Lack, D. A., Moosmüller, H., McMeeking, G. R., Chakrabarty, R. K., and Baumgardner, D.: Characterizing elemental, equivalent black, and refractory black carbon aerosol particles: a review of techniques, their limitations and uncertainties, *500 Analytical and bioanalytical chemistry*, 406, 99-122, 10.1007/s00216-013-7402-3, 2014.

Lanz, V. A., Alfara, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., Szidat, S., Wehrli, M. N., Wacker, L., Weimer, S., Caseiro, A., Puxbaum, H., and Prevot, A. S. H.: Source attribution of submicron organic aerosols during wintertime inversions by advanced factor analysis of aerosol mass spectra, *Environmental Science & Technology*, 42, 214-220, Doi 10.1021/Es0707207, 2008.

505 LASKAR, S. I., JASWAL, K., BHATNAGAR, M. K., and RATHORE, L. S.: India Meteorological Department, India, 1021-1037, 2016.

Mohr, C., DeCarlo, P. F., Heringa, M. F., Chirico, R., Slowik, J. G., Richter, R., Reche, C., Alastuey, A., Querol, X., Seco, R., Penuelas, J., Jimenez, J. L., Crippa, M., Zimmermann, R., Baltensperger, U., and Prevot, A. S. H.: Identification and quantification of organic aerosol from cooking and other sources in Barcelona using aerosol mass spectrometer data, *Atmos. Chem. Phys.*, 12, 1649-1665, DOI 10.5194/acp-12-1649-2012, 2012.

510 Mukherjee, S., Singla, V., Pandithurai, G., Safai, P. D., Meena, G. S., Dani, K. K., and Anil Kumar, V.: Seasonal variability in chemical composition and source apportionment of sub-micron aerosol over a high altitude site in Western Ghats, India, *Atmos. Environ.*, 180, 79-92, 10.1016/j.atmosenv.2018.02.048, 2018.

Nazeer Hussain, S., Chakradhar Rao, T., Balakrishnaiah, G., Rama Gopal, K., Raja Obul Reddy, K., Siva Kumar Reddy, N., Lokeswara Reddy, T., Pavan Kumari, S., Ramanjaneya Reddy, P., and Ramakrishna Reddy, R.: Investigation of black carbon

aerosols and their characteristics over tropical urban and semi-arid rural environments in peninsular India, *Journal of Atmospheric and Solar-Terrestrial Physics*, 167, 48-57, 10.1016/j.jastp.2017.10.010, 2018.

520 Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prevot, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos Chem Phys*, 10, 4625-4641, DOI 10.5194/acp-10-4625-2010, 2010.

Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., and Sun, Y.: An Aerosol Chemical Speciation Monitor (ACSM) for routine monitoring of the composition and mass concentrations of ambient aerosol, *Aerosol Science and Technology*, 45, 780-794, 2011.

525 Paatero, P., Hopke, P. K., Song, X. H., and Ramadan, Z.: Understanding and controlling rotations in factor analytic models, *Chemometr Intell Lab*, 60, 253-264, Doi 10.1016/S0169-7439(01)00200-3, 2002.

Petzold, A., Kramer, H., and Schonlinner, M.: Continuous measurement of atmospheric black carbon using a multi-angle absorption photometer, *Environ Sci Pollut R*, 9, 78-82, 2002.

530 Platt, S. M., El Haddad, I., Pieber, S. M., Zardini, A. A., Suarez-Bertoa, R., Clairotte, M., Daellenbach, K. R., Huang, R. J., Slowik, J. G., Hellebust, S., Temime-Roussel, B., Marchand, N., de Gouw, J., Jimenez, J. L., Hayes, P. L., Robinson, A. L., Baltensperger, U., Astorga, C., and Prévôt, A. S. H.: Gasoline cars produce more carbonaceous particulate matter than modern filter-equipped diesel cars, *Sci Rep-Uk*, 7, 4926, 10.1038/s41598-017-03714-9, 2017.

Pope, I. C., Burnett, R. T., Thun, M. J., and et al.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *JAMA*, 287, 1132-1141, 10.1001/jama.287.9.1132, 2002.

535 Prakash, J., Vats, P., Sharma, A. K., Ganguly, D., and Habib, G.: New Emission Inventory of Carbonaceous Aerosols from the On-road Transport Sector in India and its Implications for Direct Radiative Forcing over the Region, *Aerosol Air Qual Res*, 20, 741-761, 10.4209/aaqr.2019.08.0393, 2020.

Raatikainen, T., Brus, D., Hooza, R. K., Hyvarinen, A. P., Asmi, E., Sharma, V. P., Arola, A., and Lihavainen, H.: Size-selected black carbon mass distributions and mixing state in polluted and clean environments of northern India, *Atmos Chem Phys*, 17, 371-383, 10.5194/acp-17-371-2017, 2017.

540 Rajesh, T. A., and Ramachandran, S.: Characteristics and source apportionment of black carbon aerosols over an urban site, *Environ Sci Pollut R*, 24, 8411-8424, 10.1007/s11356-017-8453-3, 2017.

Ramgolam, K., Favez, O., Cachier, H., Gaudichet, A., Marano, F., Martinon, L., and Baeza-Squiban, A.: Size-partitioning of an urban aerosol to identify particle determinants involved in the proinflammatory response induced in airway epithelial cells, *Part Fibre Toxicol*, 6, 6-9, 2009.

545 Reyes-Villegas, E., Bannan, T., Le Breton, M., Mehra, A., Priestley, M., Percival, C., Coe, H., and Allan, J. D.: Online Chemical Characterization of Food-Cooking Organic Aerosols: Implications for Source Apportionment, *Environmental Science & Technology*, 52, 5308-5318, 10.1021/acs.est.7b06278, 2018.

Sandradewi, J., Prévôt, A. S. H., Szidat, S., Perron, N., Alfarra, M. R., Lanz, V. A., Weingartner, E., and Baltensperger, U. R. 550 S.: Using aerosol light absorption measurements for the quantitative determination of wood burning and traffic emission contribution to particulate matter, *Environmental Science and Technology*, 42, 3316-3323, 10.1021/es702253m, 2008.

Satish, R., Shamjad, P., Thamban, N., Tripathi, S., and Rastogi, N.: Temporal Characteristics of Brown Carbon over the Central Indo-Gangetic Plain, *Environmental Science and Technology*, 51, 6765-6772, 10.1021/acs.est.7b00734, 2017.

555 Shinozuka, Y., Saide, P. E., Ferrada, G. A., Burton, S. P., Ferrare, R., Doherty, S. J., Gordon, H., Longo, K., Mallet, M., Feng, Y., Wang, Q., Cheng, Y., Dobrucki, A., Freitag, S., Howell, S. G., LeBlanc, S., Flynn, C., Segal-Rosenheimer, M., Pistone, K., Podolske, J. R., Stith, E. J., Bennett, J. R., Carmichael, G. R., da Silva, A., Govindaraju, R., Leung, R., Zhang, Y., Pfister, L., Ryoo, J. M., Redemann, J., Wood, R., and Zuidema, P.: Modeling the smoky troposphere of the southeast Atlantic: a comparison to ORACLES airborne observations from September of 2016, *Atmos. Chem. Phys.*, 20, 11491-11526, 10.5194/acp-20-11491-2020, 2020.

560 Shivani, Gadi, R., Sharma, S. K., and Mandal, T. K.: Seasonal variation, source apportionment and source attributed health risk of fine carbonaceous aerosols over National Capital Region, India, *Chemosphere*, 237, 124500, <https://doi.org/10.1016/j.chemosphere.2019.124500>, 2019.

Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer,

565 R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509-559, 10.1002/2016rg000540, 2017.

570 Singla, V., Mukherjee, S., Pandithurai, G., Dani, K. K., and Safai, P. D.: Evidence of Organonitrate Formation at a High Altitude Site, Mahabaleshwar, during the Pre-monsoon Season, *Aerosol Air Qual Res*, 19, 1241-1251, 10.4209/aaqr.2018.03.0110, 2019.

575 Thamban, N. M., Tripathi, S. N., Moosakutty, S. P., Kuntamukkala, P., and Kanawade, V. P.: Internally mixed black carbon in the Indo-Gangetic Plain and its effect on absorption enhancement, *Atmos Res*, 197, 211-223, <https://doi.org/10.1016/j.atmosres.2017.07.007>, 2017.

Tiwari, S., Pervez, S., Cinzia, P., Bisht, D. S., Kumar, A., and Chate, D.: Chemical characterization of atmospheric particulate matter in Delhi, India, part II: Source apportionment studies using PMF 3.0, *Sustainable Environment Research*, 23, 295-306, 2013.

580 Turner, A. G., Bhat, G. S., Martin, G. M., Parker, D. J., Taylor, C. M., Mitra, A. K., Tripathi, S. N., Milton, S., Rajagopal, E. N., Evans, J. G., Morrison, R., Pattnaik, S., Sekhar, M., Bhattacharya, B. K., Madan, R., Govindankutty, M., Fletcher, J. K., Willetts, P. D., Menon, A., Marsham, J. H., team, a. t. I., Hunt, K. M. R., Chakraborty, T., George, G., Krishnan, M., Sarangi, C., Belušić, D., Garcia-Carreras, L., Brooks, M., Webster, S., Brooke, J. K., Fox, C., Harlow, R. C., Langridge, J. M., Jayakumar, A., Böing, S. J., Halliday, O., Bowles, J., Kent, J., O'Sullivan, D., Wilson, A., Woods, C., Rogers, S., Smout-Day, R., Tideman, D., Desai, D., Nigam, R., Paleri, S., Sattar, A., Smith, M., Anderson, D., Bauguitte, S., Carling, R., Chan, C., Devereau, S., Gratton, G., MacLeod, D., Nott, G., Pickering, M., Price, H., Rastall, S., Reed, C., Trembath, J., Woolley, A., Volonté, A., and New, B.: Interaction of convective organization with monsoon precipitation, atmosphere, surface and sea: The 2016 INCOMPASS field campaign in India, *Quarterly Journal of the Royal Meteorological Society*, n/a, 10.1002/qj.3633, 2019.

585 Weingartner, E., Saathoff, H., Schnaiter, M., Streit, N., Bitnar, B., and Baltensperger, U.: Absorption of light by soot particles: determination of the absorption coefficient by means of aethalometers, *J Aerosol Sci*, 34, 1445-1463, 10.1016/S0021-8502(03)00359-8, 2003.

590 Young, D. E., Allan, J. D., Williams, P. I., Green, D. C., Flynn, M. J., Harrison, R. M., Yin, J., Gallagher, M. W., and Coe, H.: Investigating the annual behaviour of submicron secondary inorganic and organic aerosols in London, *Atmos. Chem. Phys.*, 15, 6351-6366, 10.5194/acp-15-6351-2015, 2015.

Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, *Environmental Science & Technology*, 39, 4938-4952, 10.1021/es048568l, 2005.

595 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys Res Lett*, 34, L13801, Artn L13801
600 Doi 10.1029/2007gl029979, 2007.

Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Analytical and bioanalytical chemistry*, 401, 3045-3067, 2011.