



Tracking Separate Contributions of Diesel and Gasoline Vehicles to Roadside PM_{2.5} Through Online Monitoring of Volatile Organic Compounds, PM_{2.5} Organic and Elemental Carbon: A Six-Year Study in Hong Kong

5 Yee Ka Wong¹, X.H. Hilda Huang¹, Peter K.K. Louie², Alfred L.C. Yu², Damgy H.L. Chan², Jian Zhen Yu^{1,3}

¹Division of Environment & Sustainability, Hong Kong University of Science & Technology, Clear Water Bay, Kowloon, Hong Kong

²Hong Kong Environmental Protection Department, 47/F, Revenue Tower, 5 Gloucester Road, Wan Chai, Hong Kong

10 ³Department of Chemistry, Hong Kong University of Science & Technology, Clear Water Bay, Kowloon, Hong Kong

Correspondence to: Jian Zhen Yu (jian.yu@ust.hk)

Abstract. Vehicular emissions contribute a significant portion to fine particulate matter (PM_{2.5}) air pollution in urban areas. Knowledge of the relative contribution of gasoline versus diesel powered vehicles is highly policy relevant and yet there lacks an effective observation-based method to determine this quantity, especially for its robust tracking over a period of

15 years. In this work, we present an approach to track separate contributions by gasoline and diesel vehicles through positive matrix factorization (PMF) analysis of online monitoring data measurable by relatively inexpensive analytical instruments. They are PM_{2.5} organic and elemental carbon (OC and EC), C₂–C₉ volatile organic compounds (VOCs) (e.g., pentanes, benzene, xylenes, etc.) and nitrogen oxides concentrations. The method was applied to monitoring data spanning over six

20 years between 2011 and 2017 in a roadside environment in Hong Kong. We found that diesel vehicles accounted for ~ 70–90 % of the vehicular PM_{2.5} (PM_{vehicle}) over the years and the remaining from gasoline vehicles. The diesel PM_{vehicle} during a truck- and a bus-dominated periods showed declining trends, in coincidence with control efforts targeting at diesel commercial vehicles and franchised buses in the intervening period. The combined PM_{vehicle} from diesel and gasoline vehicles by PMF agrees well with an independent estimate by the EC-tracer method, both confirming PM_{vehicle} contributed significantly to the PM_{2.5} in this urban environment (~ 4–8 µg m⁻³, representing 30–60 % in summer and 10–20 % in winter).

25 Our work shows that long-term monitoring of roadside VOCs and PM_{2.5} OC and EC is effective for tracking gaseous and PM pollutants from different vehicle categories. This work also demonstrates the value of evidence-based approach in support of effective control policy formulation.

1 Introduction

Vehicular emissions (VE) are among the major sources of air pollution in the urban environment. Major constituents in VE

30 include nitrogen oxides (NO_x), carbon monoxide (CO), volatile organic compounds (VOCs) and fine particulate matter



(PM_{2.5}). Two primary components in vehicular PM_{2.5} (PM_{vehicle}) are elemental carbon (EC) and organic matter (OM) (Kleeman et al., 2000; Chow et al., 2011). Growing evidence has shown exposure to VE affects human health (Peters et al., 2004; Beelen et al., 2008; Benbrahim-Tallaa et al., 2012; Rice et al., 2015). With an increasing number of the global population residing in urban areas, VE have become the major target for source control in many parts of the world. But
35 uncertainty over the relative importance of diesel and gasoline vehicles to PM_{vehicle} often poses challenge in effective policymaking (Gertler, 2005).

The Chemical Mass Balance (CMB) Model and Positive Matrix Factorization (PMF) Model are two prevalent receptor models for quantifying contributing sources to PM_{2.5} including VE. However, their capability to resolve separate contributions for diesel and gasoline vehicles is often severely constrained when relying on chemical constituents residing in
40 the PM (particulate matter) fraction alone. In CMB, EC and certain organic compounds (e.g. hopanes, benzo[ghi]perylene and coronene) have been specifically used as tracers for deriving diesel and gasoline PM_{vehicle} contributions, respectively (Schauer et al., 1996; Subramanian et al., 2006; Chow et al., 2007). However, the contributions of the two are often subjected to large uncertainty due to the substantial variability in source profiles and oxidation degradation of organic tracers (Subramanian et al., 2006; Weitzkamp et al., 2008). On the other hand, PMF analysis of PM compositions often yields one
45 overall VE factor due to the lack of tracers specific to individual VE types and the difficulty is compounded by the often similar temporal pattern of traffic activity among different vehicle types (Dallmann et al., 2014; Lee et al., 2015; Wang et al., 2017). Some other chemical characteristics such as sub-fractions of OC and EC obtained from thermal analysis and metals (e.g. Mn and Fe) have been used in PMF to differentiate diesel and gasoline contributions (Kelly et al., 2013). These characteristics, however, are relatively less specific and thus they are often not applicable to typical urban areas where a
50 complex mix of contributing sources exists.

In Hong Kong (HK), PM_{vehicle} contribution from diesel vehicles has reduced significantly over the last two decades, resulting from a series of ambitious control efforts. The success has been verified by results from an ad hoc roadside study and a study comparing the emissions in a local tunnel between 2003 and 2015 (Lee et al., 2017; Wang et al., 2018). Most HK studies in the past only reported overall PM_{vehicle} contribution due to the lack of separate local source profiles for diesel and gasoline
55 vehicles and constraints in PMF model posed by a lack of vehicle type specific tracers (Li et al., 2012; Huang et al., 2014; Cheng et al., 2015; Sun et al., 2016). Some studies achieved the separation either by using non-local source profiles in CMB, or by coupling vehicle type specific traffic data collected in a short period with aerosol mass spectrometry-based PMF (Zheng et al., 2006; Lee et al., 2017). The lack of a robust means to differentiate diesel and gasoline contributions to PM_{vehicle} calls for a need to develop a more effective source apportionment strategy, especially considering the long-term need in
60 monitoring their impact on air quality.

In VOC source apportionment studies, certain VOC species have been used to track the contributions of specific vehicle types, such as propane and butanes for vehicles fueled by liquefied petroleum gas (LPG) and pentanes and toluene for gasoline vehicles (Lyu et al., 2016; Yao et al., 2019). These gaseous species, however, are rarely considered for deriving vehicle type specific PM_{vehicle}. Lambe et al. (2009) added a few VOC species into their 2 h resolution organic tracer-based



65 PMF to explicitly apportion black carbon in Pittsburgh to diesel and gasoline vehicles. Thornhill et al. (2010) also used PMF
with real-time gaseous species (including VOCs) and $PM_{2.5}$ concentration data captured by a suit of mobile equipment to
quantify $PM_{2.5}$ contributions from diesel and gasoline vehicles in Mexico City. Our group reported limited exploration of the
combined use of hourly VOCs and $PM_{2.5}$ organic carbon (OC) and EC data in PMF to estimate the total $PM_{vehicle}$ at a
roadside environment in HK (Huang et al., 2014). Here we present a detailed investigation on the feasibility of such an
70 approach for separating $PM_{vehicle}$ contributions by diesel and gasoline vehicles using a comprehensive dataset covering six
years between 2011 and 2017.

The objective of this work is to establish an approach to obtain vehicle type specific $PM_{vehicle}$, through integrating online
routine monitoring data, such as NO_x , hourly VOCs and OC-EC measurement data, into PMF analysis. The study features a
six-year-long monitoring (2011–2017) in a roadside environment in HK. We evaluate this new method through comparing
75 the total $PM_{vehicle}$ derived from an independent EC-tracer method developed previously, and for the first time report the long-
term trends in $PM_{vehicle}$ for diesel and gasoline vehicles in HK (Huang et al., 2014; Wong et al., 2019). Several policies
targeting at diesel vehicles fell within the timeline of the study period, providing a valuable opportunity to examine their
effectiveness. The methodology presented in this study for instrument deployment, data collection and analysis are highly
recommended to the air quality management authorities as the research outcome has demonstrated how the routine
80 monitoring dataset could be turned into measurement-based evidence for evaluating effectiveness of control policies
targeting vehicles.

2 Methods

2.1 Roadside Measurements

The study time window spans over a six-year period from May 2011 to August 2017. Sampling was conducted at Mong Kok
85 Air Quality Monitoring Station (MK AQMS), which is a roadside site in the Air Quality Monitoring Network operated by
the HK Environmental Protection Department (HKEPD). The station is located at the junction of two trunk roads, with an
annual average daily traffic count of $\sim 45,000$ (Transport Department, 2018). Previous vehicle counting exercise showed that
private cars fueled by gasoline, goods vehicles and buses fueled by diesel and taxis running on LPG made up 32 %, 22 %, 16
% and 29 % of the vehicle fleet in the sampling area (Lee et al., 2017). In addition to the busy traffic, there are small shops,
90 restaurants, and tall residential and commercial buildings in the immediate vicinity of the station. A previous study has
shown that on-road vehicles are the most important sources of submicron carbonaceous aerosols at the site, followed by
cooking activities (Lee et al., 2015).

A semi-continuous OC-EC field analyzer system (RT-4, Sunset Laboratory, OR, USA) was operated to obtain hourly OC-
EC concentrations data in $PM_{2.5}$. Details for the on-site operation, instrument conditions and quality control work during the
95 entire sampling period are provided in the supplementary material. Hourly concentration data for $PM_{2.5}$, C_2 – C_9 VOCs, and
trace gases including NO_x and CO measured at the MK AQMS were provided by HKEPD. Details of the monitoring



equipment can be found in HKEPD's annual air quality report (Environmental Protection Department, 2018). In particular, the VOCs were measured with a GC955 series 611/811 VOC analyzer (Syntech Spectras, The Netherlands) which quantifies 30 species including 11 C₂–C₈ alkanes, 9 C₂–C₅ alkenes, 1 C₂ alkyne and 9 C₆–C₉ single-ring aromatics.

100 2.2 Data Treatment

To avoid biased interpretation of the relationships between the measured species, we exclude hourly samples with one or more missing species from the subsequent data analyses. Sampling days with data cover rate (i.e., number of valid data/total number of hours during the study period) < 75 % are also excluded to maximize the representativeness of the concentration data of a sampling day. The trends presented throughout this study are constructed from the monthly averages. Only months
105 with data cover rate > 33 % are considered. The monthly data cover rates are summarized in Fig. S1.

2.3 Estimation of Vehicular PM_{2.5} by Positive Matrix Factorization

Vehicular contributions to PM_{2.5} are quantified by PMF analysis using the EPA PMF 5.0 software (Paatero and Tapper, 1994; Paatero, 1997; Norris et al., 2014). PMF is a receptor model that solves the chemical mass balance of a speciated sample data matrix by decomposing it into factor profiles and factor contributions with non-negative constraints, with the
110 objective to minimize the uncertainty weighted deviation between observed and modeled species concentrations. In this work, concentrations of OC, EC, NO_x, CO and twelve VOC species that were consistently detected above detection limit (> 80 %) in each calendar year are included in the input dataset. The twelve VOCs include ethene, ethane, propane, propene, *i*-butane, *n*-butane, *i*-pentane, *n*-pentane, benzene, toluene, ethylbenzene and *m*-&*p*-xylene. Some examples of the excluded VOCs are butadiene, *n*-hexane and *n*-heptane. Preliminary PMF analysis showed that including these species had no
115 advantage in identifying more sources. Meanwhile, the considerable fraction of below detection limit data for these species would affect the quality of the PMF solutions. Details regarding other modeling inputs are provided in the supplementary material. Vehicle-related VOCs in the roadside environment are freshly emitted and thus should be mostly conserved, rendering them suitable for receptor modeling. PM_{vehicle} for individual vehicle types are calculated as the sum of OM and EC in the corresponding factor profiles, considering it is primarily composed of OM and EC. OM is estimated to be OC
120 multiplying by a factor of 1.2 based on organic aerosol mass spectra measured for both diesel and gasoline VE (Dallmann et al., 2014; Lee et al., 2015).

We note that cooking emissions, a known OC source in MK, is not considered in the current PMF analysis, as the relevant tracer compounds for cooking was not supported with the instrumentation deployed in this study. We therefore relax the modeling of OC (i.e., allow the modeled OC to have a relatively larger deviation from measurement) by tripling its
125 uncertainty. This approach allows the apportioning of OC a larger degree of freedom, which in effect allows the model to only capture the OC that is associated with the fitting species, whilst leaving the unrelated fraction as unapportioned OC. A sensitivity test showed that further doubling the OC uncertainty would not cause a discernible impact on the PMF solution. The robustness of the PMF solutions is examined through executing the displacement (DISP) and Fpeak (strength values of –



5 and 5) functions. In bootstrap (BS) analysis, the input dataset was split into three groups of equal sample size for execution because of the limited computing capability of the software (total sample size = 24,586). Nevertheless, this practice allows us to assess the model uncertainty associated with using different subsets of samples, which will be discussed in the result section.

2.4 Estimation of Vehicular PM_{2.5} by EC-tracer Method

To evaluate the PMF estimation, an EC-tracer method specifically designed for estimating PM_{vehicle} contribution in roadside environments is applied. Details regarding the principle and application of this method are documented in our previous work and a brief account is given here (Huang et al., 2014; Wong et al., 2019). In this method, VE is assumed to have a characteristic OC-to-EC ratio ([OC/EC]_{vehicle}) and be responsible for all ambient EC. The latter represents a reasonable approximation given that the EC at MK AQMS was dominated by traffic exhaust over the entire study period, which will be discussed further in Sect. 3.1.2. With these assumptions, OC_{vehicle} can be estimated as the product of the ambient EC concentration and the [OC/EC]_{vehicle}, while PM_{vehicle} can be estimated using the approach similar to that introduced in Sect. 2.3 (i.e., $PM_{vehicle} = EC_{ambient} + OC_{vehicle} \times 1.2$).

The [OC/EC]_{vehicle} is determined using the minimum OC/EC ratio approach, in which the OC in a certain lowest percentage range by OC/EC ratio are regressed on EC, and the slope obtained represents the target ratio (Lim and Turpin, 2002). These lowest OC/EC ratios typically represent the composition of primary emissions from fossil fuel combustion sources, which are clearly dominated by VE in roadside environment. In this study, the optimal Deming regression evaluated previously was applied on the lowest 5 % data by OC/EC ratio on a monthly basis (Huang et al., 2014). The analysis was performed using the Igor (WaveMetrics, Inc. Lake Oswego, OR, USA) based computer program developed by Wu (2017).

3 Results and Discussion

3.1 Ambient Trends

3.1.1 OC Trends

The monthly OC concentration showed a decreasing trend over the six-year period from May 2011 to August 2017 as shown in Fig. 1a. A consistent seasonal cycle with fall–winter (mid-September to mid-March of next year) high, summer (mid-May to mid-September) low and spring (mid-March to mid-May) in between is observed over the years. The study-wide OC in the three seasonal periods were 6.9 ± 3.4 , 3.9 ± 2.6 and $5.9 \pm 2.8 \mu\text{g C m}^{-3}$, respectively (Fig. S2). The main cause for the seasonal variations of OC is related to the geographical location of HK, which is in the coastal area facing the South China Sea to the south and mainland China to the north. During the fall/winter monsoon season, the prevailing northeasterly wind transports pollutants from the continental area to HK (Hagler et al., 2006). Other reasons for the elevated OC observed in wintertime include reduced mixing height and enhanced partitioning of semivolatile organic compounds (SVOCs) into the



particle phase due to lower temperature and higher organic aerosol loading. It is noted that the decreasing trend of OC was
 160 mainly driven by the reduction in winter OC. The winter average OC concentration dropped significantly from $10.7 \mu\text{g C m}^{-3}$
 3 in 2011 to $4.3 \mu\text{g C m}^{-3}$ in 2017, while the decrease in summer was from 5.1 to $2.8 \mu\text{g C m}^{-3}$ during the same period.

3.1.2 EC Trends

The six-year trend of EC concentration is plotted in Fig. 1b, which shows a different temporal characteristic compared to
 OC. A main feature of the EC trend is the lack of seasonality throughout the years. The study-wide seasonal concentrations
 165 remained at $\sim 5 \mu\text{g C m}^{-3}$ in all seasons (Fig. S2). We previously demonstrated that EC at MK AQMS was mainly influenced
 by vehicular traffic by showing similarities in their diurnal and weekday–holiday variation patterns (Huang et al., 2014).
 Such correlations persisted over the years (Fig. S3). These indicate that the EC concentration at the site was mainly affected
 by local VE sources and less impacted by regional sources regardless of season. In the first three years, the monthly EC
 concentrations fluctuated near the $5\text{--}6 \mu\text{g C m}^{-3}$ range. Starting from mid-2014, it declined significantly to the level of ~ 3
 170 $\mu\text{g C m}^{-3}$ toward the end of the measurement period. Similar variation trend was also observed for NO_x (Fig. 1c), which is
 mainly generated by on-road diesel vehicles in the roadside environment. Notably, these decreasing trends coincided with
 the launch of a Phasing Out Pre-Euro IV Diesel Commercial Vehicles Program in March 2014 in HK. The results here imply
 diesel vehicles were the major EC contributor at the sampling site.

3.1.3 Carbonaceous Aerosols and $\text{PM}_{2.5}$ Trends

175 The relative contributions of carbonaceous aerosols to $\text{PM}_{2.5}$ at MK AQMS over the study time window is shown in Fig. 1d.
 OM was approximated as $\text{OC} \times 1.4$ for typical urban aerosols with primary and secondary origins. The $\text{PM}_{2.5}$ concentration
 is overlaid on the same plot. As shown in the figure, $\text{PM}_{2.5}$ concentration displayed a seasonal variation (winter high and
 summer low) similar to that of OC over the years, which was a result of combined effect of regional air pollutant transport
 and meteorological conditions, as discussed in Sect. 3.1.1. In the middle of the year with warmer weather and lower $\text{PM}_{2.5}$
 180 mass ($\sim 20 \mu\text{g m}^{-3}$), EC showed an elevated relative contribution to $\sim 30\%$, and EC and OM had comparable contributions
 to $\text{PM}_{2.5}$. The opposite was shown in colder season with higher $\text{PM}_{2.5}$ mass ($30\text{--}60 \mu\text{g m}^{-3}$). EC only made up $\sim 15\%$ of
 aerosol mass while OM accounted for $\sim 30\%$. About two-thirds of $\text{PM}_{2.5}$ in these periods were composed of non-
 carbonaceous materials. Based on HKEPD's continuous filter-based $\text{PM}_{2.5}$ speciation effort, these materials mainly consist
 of secondary inorganics (sulfate, nitrate and ammonium), followed by crustal material, and trace elements (Yu and Zhang,
 185 2018). The secondary inorganic components have long been attributed to regional air pollution. The persistently large
 contributions from these components across the years indicates that controlling $\text{PM}_{2.5}$ including its gaseous precursors (SO_2 ,
 NO_x and NH_3) on a regional scale is still important to reduce the overall $\text{PM}_{2.5}$ at this roadside location.

3.1.4 *n*-Butane and *i*-Pentane Trends

Figure 1 also shows multi-year trends in VOCs that are associated with specific vehicle types (LPG and gasoline). *n*-Butane



190 has been used to track LPG-fueled vehicles at MK AQMS (Lyu et al., 2016; Yao et al., 2019). As shown in Fig. 1e, no seasonality in *n*-butane level was observed over the years, supporting this species was predominantly emitted by local LPG vehicles. It remained at ~ 10 – 15 ppbv level in mid-2011–mid-2013, dropped precipitously to ~ 7 ppbv in the second half of 2013, followed by a steadily declining trend until the end of the study period. Yao et al. (2019) reported similar trend characteristics for the same site with a more continuous dataset (September 2012–April 2017). The drop in the second half of 195 2013 was a response to a catalytic converter replacement scheme for LPG-fueled vehicles implemented by the Government (Lyu et al., 2016).

Similar to *n*-butane, *i*-pentane was also dominated by a local source as reflected by the absence of seasonality over the years (Fig. 1f). This gasoline exhaust/evaporation tracer remained fairly stable at ~ 1 ppbv level over the entire study period. Such an invariability is in line with a previous study showing the VOCs contributions from gasoline-powered vehicles in the same 200 study area were relatively stable over the similar period (Yao et al., 2019).

3.2 $[\text{OC}/\text{EC}]_{\text{vehicle}}$ for EC-tracer Method

$[\text{OC}/\text{EC}]_{\text{vehicle}}$ determined using summer data (June, July and August) are very similar over the years and do not exhibit obvious trend over the years, as shown in Fig. 2a. The ratios range from 0.30 to 0.47, with R^2 between 0.56–0.96 (sample size $n = 18$ –33). Figure 2b plots the frequency of occurrence of the lowest 5 % OC/EC ratios in 24 hours of a day for the 205 summer months. It shows that the lowest ratios occurred most frequently near the morning rush hour (7:00–10:00 a.m.) with minimal contribution from other primary sources (e.g. cooking emissions), supporting that these ratios were dominated by VE. The ratios were a factor of 2–3 higher in winter (Fig. S4), likely as a result of enhanced OC contributions from aged air mass and biomass burning from regional transport as discussed in Sect. 3.1.1. Thus, these values were likely biased high. Another complicating factor is that the reduction in ambient temperature and elevation in organic aerosol concentration in 210 colder season would favor the partitioning of SVOCs into particle phase, thereby inflating the $[\text{OC}/\text{EC}]_{\text{vehicle}}$ (Robinson et al., 2007). We did not account for this effect in this study. Instead, the summer values were adopted for deriving the $[\text{OC}/\text{EC}]_{\text{vehicle}}$ and this ratio is considered a lower-estimate for colder season samples. Given that similar $[\text{OC}/\text{EC}]_{\text{vehicle}}$ values were obtained in the summer months across the years, the mean value of 0.35 (standard deviation = 0.05) is considered to be the best estimate of $[\text{OC}/\text{EC}]_{\text{vehicle}}$ for subsequent analysis.

215 3.3 Vehicular Contributions by PMF Analysis

3.3.1 Source Identification

Among various PMF runs, the result of five-factor run is the most interpretable for source identification and quantification. The contributing sources are identified through studying the presence of marker species and temporal variations in normalized contributions as shown in Fig. 3. Three vehicular factors are identified, corresponding to the vehicle types 220 observed near the sampling site. The first factor represents diesel exhaust as it contains the majority of NO_x and EC, which



are largely attributed to diesel vehicles. The low OC/EC ratio of 0.5 and diurnal profile, in line with traffic flow of diesel vehicles next to the site, further confirm its source identity.

Factor 2 is associated with gasoline vehicles due to the dominant presence of *i*-/*n*-pentane. This factor also has a diurnal profile consistent with gasoline vehicle flow near the site, with evening peak occurring two hours later than that of diesel vehicles. The *i*-/*n*-pentane could be emitted through fuel evaporation and as unburnt gasoline in tailpipe exhaust, whereas the carbonaceous particulates with OC/EC ratio of 2.5 and CO signal tailpipe exhaust.

Contribution from LPG-fueled vehicles is identified in the third factor by propane and *i*-/*n*-butane originating mainly from fuel evaporation. Its diurnal variation pattern is consistent with the activity pattern (busy in small hours) of local taxis running on LPG (Yu et al., 2016). Note this vehicle type has negligible contribution to PM, in agreement with the highly volatile nature of LPG.

The fourth factor contains notable amount of toluene, ethylbenzene and *m*-&*p*-xylene that commonly exist in consumer and industrial products as solvent or in gasoline as additives (Bolden et al., 2015). Previous studies attributed this factor to solvent usage (Lyu et al., 2016; Yao et al., 2019). However, upon closer examination to its diurnal pattern, we found that this factor shows regular peaks around 11:00 and 17:00. Given the MK AQMS is surrounded by sixteen 24/7 gas stations within 1.5 km (Fig. S5), and the peak business hours of some of these stations show similar diurnal variation pattern (based on popular times information from Google Maps, Fig. S5), we classify this factor as fuel-filling process instead of solvent usage.

The last factor has abundant chemically stable ethane and benzene. Particulates in this factor are also enriched substantially in OC (OC/EC ratio ~ 9). In contrast to the three vehicle-related factors, absence of diurnal variation and presence of winter-high and summer-low contribution were noted (Fig. S6). These characteristics collectively indicate this factor is aged air mass.

3.3.2 Model Evaluation

Modeling uncertainty estimation from DISP, Fpeak and BS (grouped samples) show that the solutions are rotationally and statistically robust, with details provided in the supplementary material. In particular, the factor profiles from the three grouped PMF analyses are very similar, as shown in Fig. S7. The [OC/EC]_{vehicle} ratios for diesel vehicles are 0.5 in the all grouped runs, while those for gasoline vehicles are 1.8–2.2, implying the chemical characteristics of PM_{vehicle} of the two remained similar over the study time period. The PMF solution (base run) is also evaluated from the modeling performance of the fitting species, which are summarized in Table S3. Most gaseous species are well reproduced ($R^2 = 0.73$ – 0.95), except ethene and ethane ($R^2 = 0.32$ – 0.40) due to their higher measurement uncertainties. Modeled EC is also consistent with the measurement ($R^2 = 0.90$), but OC in comparison shows larger discrepancy ($R^2 = 0.64$).

To unveil the cause of the OC discrepancy, we compare the modeled and measured OC concentrations on a diurnal basis. The result is presented in Fig. 4 using box diagram. The plot reveals that the discrepancy mainly occurred during 12:00–14:00 and 18:00–21:00. More specifically, the modeled OC were considerably lower than the measurement during these two



periods. This feature remained across all seasons as depicted in Fig. S8. Previous studies have attributed the two organic peaks during these two mealtime periods to cooking emissions (Lee et al., 2015; Sun et al., 2016). As mentioned in Sect. 2.3, we were unable to account for this source in PMF due to the lack of suitable tracer species for cooking emissions. Nevertheless, Fig. 4 shows that the modeled and measured OC had a good agreement during the non-mealtime hours (i.e., 0:00–11:00 and 15:00–17:00). This indicates that the aged air mass factor and VE factors resolved by the PMF were able to explain the non-cooking OC, lending support to the PMF-derived OC_{vehicle} . It should be noted that although PMF without down-weighting OC could bring the modeled OC into better agreement with measurement, the resolved factor profiles were less consistent among the three grouped PMF analyses, causing discontinuity in factor contributions between different periods. The improved interpretability of the PMF results indicates that down-weighting OC is necessary in our situation.

3.3.3 PMF-derived OC_{vehicle} Trends

The separate diesel and gasoline OC_{vehicle} concentrations over the entire study period are shown in Fig. 5a. A decreasing trend in the overall OC_{vehicle} started to emerge in mid-2014, which was driven by the reduction in diesel OC_{vehicle} . Between mid-2011 and mid-2014, diesel OC_{vehicle} hovered at $\sim 2 \mu\text{g C m}^{-3}$ level. It then started decreasing until 2017, upon which the concentration had dropped by half. Gasoline contribution, on the other hand, remained at $\sim 1 \mu\text{g C m}^{-3}$ level over the entire study time period. As a result of the different pace of reductions, gasoline OC_{vehicle} has grown in relative importance over the years as shown in Fig. 5d. In the first three years, gasoline vehicles were a smaller contributor compared to diesel vehicles, accounting for $\sim 30\%$ of the OC_{vehicle} , but afterwards, its contribution has become comparable to diesel vehicles, growing to 40–50 %. Diesel and gasoline OC_{vehicle} derived from the grouped PMF analyses and the associated uncertainties derived from the 5th and 95th percentile of the BS results are plotted in Fig. S9a, while the results obtained from the base run are also shown as solid lines in the graph. It appears that the division of samples into three time periods does not exert discernible influence on the results, though the gasoline OC_{vehicle} occasionally deviated from the base result noticeably in the first time period. But in general, OC_{vehicle} from the two vehicle categories have converged over the whole study time period.

The overall OC_{vehicle} derived from the EC-tracer method is plotted in Fig. 5a for comparison. It appears that this independent method can only account for the diesel fraction of PMF-derived OC_{vehicle} . A plausible reason is that the $[OC/EC]_{\text{vehicle}}$ determined through the minimum OC/EC ratio approach (0.35) is biased toward diesel exhaust with low OC/EC ratio (0.47 from PMF). The comparison here suggests the OC_{vehicle} derived from the EC-tracer method reflects diesel influence better.

The relative contribution of total OC_{vehicle} to ambient OC exhibited a large seasonal dependence (Fig. S10a). In winter months when ambient OC concentration was high, VE made up 30–40 % of OC. The percentage share increased sharply to 70–100 % in summer months when ambient OC was low. However, it should be noted that the PMF-modeled OC occasionally exceeded the measurement, and the exceedance increased with decreasing ambient OC level (Fig. S11). This could be attributed to the uncertainties arising from PMF modeling and measuring low level of OC, and thus the relative importance in summer months is likely overestimated.



3.3.4 PMF-derived EC_{vehicle} Trends

Trends in vehicular EC (EC_{vehicle}) were very similar to those for OC_{vehicle} , as shown in Fig. 5b. Different from OC_{vehicle} , the EC_{vehicle} derived from the PMF and EC-tracer method, which is essentially the ambient EC, agree well with each other. This is because PMF attributed the majority of the ambient EC to VE sources, more than 70 % as shown in Fig. S10b. As a result, the change in EC_{vehicle} was very similar to that of ambient EC and thus is not repeated here. A key finding is that diesel vehicles dominated the EC_{vehicle} over the entire study period, constituting more than 80 % of EC_{vehicle} as shown in Fig. 5e. Such dominance remains valid after consideration of PMF modeling uncertainties (Fig. S9b). The above findings emphasize the reduction in EC over the years was mostly attributed to the control of diesel vehicles, and this vehicle category should deserve closer attention for further EC abatement.

3.3.5 PMF-derived PM_{vehicle} Trends

The positive impact on air quality of vehicle control policies, if any, is more obvious if EC and OM from VE are considered together (i.e., PM_{vehicle}). The monthly average PM_{vehicle} by vehicle category are given in Fig. 5c. As of mid-2014, PM_{vehicle} fluctuated slightly around the $8 \mu\text{g m}^{-3}$ level, followed by a considerable reduction to $\sim 4 \mu\text{g m}^{-3}$ level in the beginning of 2017. This amount of reduction represents one-fifth of typical $PM_{2.5}$ concentration at MK AQMS during summer ($\sim 20 \mu\text{g m}^{-3}$). As noted from Fig. 5c, the reduction in PM_{vehicle} was mainly driven by diesel vehicles, which was also the dominant PM_{vehicle} contributor over the whole study period as given in Fig. 5f. Another finding from Fig. 5f is that the relative importance of gasoline vehicles has only grown slightly from ~ 20 % before mid-2014 to ~ 30 % afterwards despite the drastic decrease in diesel PM_{vehicle} . After considering the PMF modeling uncertainties as depicted in Fig. S9c, it is still clear that diesel vehicles have dominated PM_{vehicle} at MK AQMS over the entire study period, and this vehicle class should remain the focal point for further control of PM_{vehicle} .

PM_{vehicle} estimated by the EC-tracer method are also compared with the PMF method in Fig. 5c. As shown, the disparity between the two approaches has narrowed considerably compared to OC_{vehicle} . The improvement mainly involves the consideration of EC, which is similarly perceived as tracer for VE in the two estimation methods. The similarity between the PM_{vehicle} obtained from the two methods lends support to our PMF model in producing reasonable estimates for separate diesel and gasoline PM_{vehicle} contributions. Furthermore, the consistent results obtained from both methods highlight that PM_{vehicle} was an important contributor to the $PM_{2.5}$ at MK AQMS, with noticeable seasonal variation similar to that of OC_{vehicle} , as given in Fig. S10c. Based on PMF results, during summer and under the dominance of local sources, VE was responsible for ~ 30 – 60 % of ambient $PM_{2.5}$ during the study period. When $PM_{2.5}$ concentration increased in winter due to regional influence, VE contribution dropped to roughly 10–20 %.



3.4 Policy Evaluation

Tackling tailpipe emissions from diesel commercial vehicles (DCVs) and franchised bus is a long-term need in HK (Environment Bureau, 2013). A three-day detailed traffic counting exercise was conducted by the Government at the MK AQMS in May 2013. The details were reported in Lee et al. (2017). From that we are able to identify a truck-dominated period around midday (11:00–13:00) during which the trucks (i.e., DCVs) number is a factor of 2–3 higher than buses (~ 100 vs. ~ 40 vehicle h^{-1}), as well as a mid-night period (22:00–0:00) with buses count being 3 times higher than trucks (~ 60 vs. ~ 20 vehicle h^{-1}). Hence, the annual trends in diesel $\text{PM}_{\text{vehicle}}$ extracted from these two periods could provide indication on how diesel $\text{PM}_{\text{vehicle}}$ has been impacted by different policies. The results are presented in Fig. 6. Also shown in the figure is a background condition represented by the small hours (2:00–4:00) when diesel vehicle number reaches the minimum (~ 10 vehicle h^{-1}). A clear declining trend in diesel $\text{PM}_{\text{vehicle}}$ is noted for both vehicle-dominated periods, with both trends approaching the levels in the background period.

For the truck-dominated trend, the $\text{PM}_{\text{vehicle}}$ levels started to drop after March 2014, which marks the commencement of the Phasing Out Pre-Euro IV DCVs Program implemented by the Government. This program aims at progressively replacing all Pre-Euro IV DCVs ($\sim 82,000$) in HK by the end of 2019 (Environment Bureau, 2013). The diesel $\text{PM}_{\text{vehicle}}$ concentrations in the pre-DCV Program period and the start of 2017 were respectively ~ 9 and $\sim 5 \mu\text{g m}^{-3}$, representing almost a 50 % reduction. The reduction also appeared to respond reasonably with the progress of the program shown in Fig. 6. Prior to this DCV program, another scheme that replaces $\sim 7,400$ Euro II DCVs was launched during July 2010–June 2013. That program, however, did not produce obvious impact on diesel $\text{PM}_{\text{vehicle}}$, possibly because of the smaller scale of the implementation compared to the more recent DCV program.

The declining trend in diesel $\text{PM}_{\text{vehicle}}$ during the bus-dominated period could be attributed to a host of control measures for franchised buses. In the intervening years, local franchised bus companies have been continuously scrapping and replacing old model buses with newer model buses complying with higher Euro Standards (Environment Bureau, 2013). At the level of transportation management, the Government has been pursuing reduction of bus trips in congested corridors through rationalization of bus routes. Setting up of low emission zones in densely populated spots including the MK area, where only cleaner model franchised buses are allowed, might also contribute to the decrease in bus-related $\text{PM}_{\text{vehicle}}$ contribution (Environment Bureau, 2013).

Conclusions

We present a holistic analysis on the long-term monitoring data of hourly $\text{PM}_{2.5}$ OC and EC, vehicle-specific VOCs (e.g. *n*-butane and *i*-pentane) and NO_x concentrations in an urban roadside environment in HK. The dataset covers a six-year period from May 2011 to August 2017. Both OC and EC concentrations were observed to decrease notably over the entire study period, plausibly due to the efficient control of pollution sources on both regional and local context. By integrating OC, EC and VOCs (e.g. *n*-butane, *i*-pentane, benzene and xylene) datasets into PMF analysis, we successfully differentiate $\text{PM}_{\text{vehicle}}$



contributions from diesel and gasoline vehicles, and for the first time report their individual long-term trends. The overall PM_{vehicle} is also estimated by the EC-tracer method, which shows good agreement with that from the PMF analysis, supporting the PM_{vehicle} estimate from the PMF. Our work identifies diesel vehicles as the dominant vehicle type in contributing PM_{vehicle} ($\sim 70\text{--}80\%$) over the entire study period. Thus, further VE control effort for mitigating roadside PM_{vehicle} in HK should focus on diesel vehicles. The technique developed in this work could be extrapolated to other roadside environments with mixed vehicular contributions, considering both continuous OC and EC analyzers and online VOC instruments are increasingly incorporated in governments' air quality monitoring programs. We note that the OC_{vehicle} estimated by this approach serves as a lower limit for the vehicle-contributed OC since the fitting species considered here are all tracers for primary emissions. The primary emissions from on-road vehicles also have high potential to form secondary organic aerosols (Gentner et al., 2017), and this secondary PM derived from vehicles is not captured in PM_{vehicle} estimates by our method. Future work should attempt to quantify this missing fraction of vehicular $PM_{2.5}$ for more insightful policy implications.

Data availability. The data presented in this article are available from the authors upon request. Please contact Jian Zhen Yu (jian.yu@ust.hk).

Author Contribution. JZY, XHHH, and YKW designed the field experiments with assistance from PKKL, ALCY and DHL. YKW and XHHH carried out the ECOC measurements. YKW performed the data analysis and prepared the manuscript with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. The content of this paper does not necessarily reflect the views and policies of the HKSAR Government, nor does mention of trade names or commercial products constitute an endorsement or recommendation of their use.

Acknowledgements. This work is supported by the Hong Kong Environmental Protection Department (HKEPD) (tender refs. AS10-336, 12-00255, and 14-05593) and Hong Kong Research Grants Council (16305418). We thank HKEPD for provision of the hourly VOC dataset.

References

- Beelen, R., Hoek, G., van den Brandt, P. A., Goldbohm, R. A., Fischer, P., Schouten, L. J., Jerrett, M., Hughes, E., Armstrong, B., and Brunekreef, B.: Long-term effects of traffic-related air pollution on mortality in a Dutch cohort (NLCS-AIR Study), *Environ. Health Persp.*, 116, 196–202, 2008.
- Benbrahim-Tallaa, L., Baan, R. A., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Guha, N., Loomis, D., and Straif, K.: Carcinogenicity of diesel-engine and gasoline-engine exhausts and some nitroarenes, *Lancet Oncol.*, 13, 663–664, 2012.
- Bolden, A. L., Kwiatkowski, C. F., and Colborn, T.: New look at BTEX: Are ambient levels a problem? *Environ. Sci. Technol.*, 49, 5261–5276, 2015.



- 380 Cheng, Y., Lee, S. C., Gu, Z. L., Ho, K. F., Zhang, Y. W., Huang, Y., Chow, J. C., Watson, J. G., Cao, J. J., and Zhang, R. J.: PM_{2.5} and PM_{10-2.5} chemical composition and source apportionment near a Hong Kong roadway, *Particuology*, 18, 96–104, 2015.
- Chow, J. C., Watson, J. G., Lowenthal, D. H., Chen, L. W. A., Zielinska, B., Mazzoleni, L. R., and Magliano, K. L.: Evaluation of organic markers for chemical mass balance source apportionment at the Fresno Supersite, *Atmos. Chem. Phys.*, 7, 1741–1754, 2007.
- 385 Chow, J. C., Watson, J. G., Lowenthal, D. H., Chen, L. W. A., and Motallebi, N.: PM_{2.5} source profiles for black and organic carbon emission inventories, *Atmos. Environ.*, 45, 5407–5414, 2011.
- Dallmann, T. R., Onasch, T. B., Kirchstetter, T. W., Worton, D. R., Fortner, E. C., Herndon, S. C., Wood, E. C., Franklin, J. P., Worsno, D. R., Goldstein, A. H., and Harley, R. A.: Characterization of particulate matter emissions from on-road gasoline and diesel vehicles using a soot particle aerosol mass spectrometer, *Atmos. Chem. Phys.*, 14, 7585–7599, 2014.
- 390 Environment Bureau: A Clean Air Plan for Hong Kong, available at: https://www.enb.gov.hk/en/files/New_Air_Plan_en.pdf, 2013.
- Environmental Protection Department: Air quality in Hong Kong 2017, The Government of the Hong Kong Special Administrative Region, Hong Kong, 2018.
- 395 Gentner, D. R., Jathar, S. H., Gordon, T. D., Bahreini, R., Day, D. A., El Haddad, I., Hayes, P. L., Pieber, S. M., Platt, S. M., de Gouw, J., Goldstein, A. H., Harley, R. A., Jimenez, J. L., Prévôt, A. S. H., and Robinson, A. L.: Review of urban secondary organic aerosol formation from gasoline and diesel motor vehicle emissions, *Environ. Sci. Technol.*, 51, 1074–1093, 2017.
- Gertler, A. W.: Diesel vs. gasoline emissions: Does PM from diesel or gasoline vehicles dominate in the US? *Atmos. Environ.*, 39, 2349–2355, 2005.
- 400 Hagler, G. S. W., Bergin, M. H., Salmon, L. G., Yu, J. Z., Wan, E. C. H., Zheng, M., Zeng, L. M., Kiang, C. S., Zhang, Y. H., Lau, A. K. H., and Schauer, J. J.: Source areas and chemical composition of fine particulate matter in the Pearl River Delta region of China, *Atmos. Environ.*, 40, 3802–3815, 2006.
- Huang, X. H. H., Bian, Q. J., Louie, P. K. K., and Yu, J. Z.: Contribution of vehicular carbonaceous aerosols to PM_{2.5} in a roadside environment in Hong Kong, *Atmos. Chem. Phys.*, 14, 9279–9293, 2014.
- 405 Kelly, K. E., Kotchenruther, R., Kuprov, R., and Silcox, G. D.: Receptor model source attributions for Utah's Salt Lake City airshed and the impacts of wintertime secondary ammonium nitrate and ammonium chloride aerosol, *J. Air Waste Manage.*, 63, 575–590, 2013.
- Kleeman, M. J., Schauer, J. J., and Cass, G. R.: Size and composition distribution of fine particulate matter emitted from motor vehicles, *Environ. Sci. Technol.*, 34, 1132–1142, 2000.
- 410 Lambe, A. T., Logue, J. M., Kreisberg, N. M., Hering, S. V., Worton, D. R., Goldstein, A. H., Donahue, N. M., and Robinson, A. L.: Apportioning black carbon to sources using highly time-resolved ambient measurements of organic molecular markers in Pittsburgh, *Atmos. Environ.*, 43, 3941–3950, 2009.



- Lee, B. P., Li, Y. J., Yu, J. Z., Louie, P. K. K., and Chan, C. K.: Characteristics of submicron particulate matter at the urban
 415 roadside in downtown Hong Kong—Overview of 4 months of continuous high-resolution aerosol mass spectrometer
 measurements, *J. Geophys. Res.-Atmos.*, 120, 7040–7058, 2015.
- Lee, B. P., Louie, P. K. K., Luk, C., and Chan, C. K.: Evaluation of traffic exhaust contributions to ambient carbonaceous
 submicron particulate matter in an urban roadside environment in Hong Kong, *Atmos. Chem. Phys.*, 17, 15121–15135, 2017.
- Li, Y. C., Yu, J. Z., Ho, S. S. H., Yuan, Z. B., Lau, A. K. H., and Huang, X. F.: Chemical characteristics of PM_{2.5} and
 420 organic aerosol source analysis during cold front episodes in Hong Kong, China, *Atmos. Res.*, 118, 41–51, 2012.
- Lim, H. J. and Turpin, B. J.: Origins of primary and secondary organic aerosol in Atlanta: results of time resolved
 measurements during the Atlanta supersite experiment, *Environ. Sci. Technol.*, 36, 4489–4496, 2002.
- Lyu, X. P., Guo, H., Simpson, I. J., Meinardi, S., Louie, P. K. K., Ling, Z. H., Wang, Y., Liu, M., Luk, C. W. Y., Wang, N.,
 and Blake, D. R.: Effectiveness of replacing catalytic converters in LPG-fueled vehicles in Hong Kong, *Atmos. Chem.*
 425 *Phys.*, 16, 6609–6626, 2016.
- Norris, G., Duvall, R., Brown, S., and Bai, S.: EPA Positive Matrix Factorization (PMF) 5.0 Fundamentals and User Guide,
 U.S. Environmental Protection Agency, 2014.
- Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometr. Intell. Lab.*, 37, 23–35, 1997.
- Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error
 430 estimates of data values, *Environmetrics*, 5, 111–126, 1994.
- Peters, A., von Klot, S., Heier, M., Trentinaglia, I., Hörmann, A., Wichmann, H. E., and Löwel, H.: Exposure to traffic and
 the onset of myocardial infarction, *New Engl. J. Med.*, 351, 1721–1730, 2004.
- Rice, M. B., Ljungman, P. L., Wilker, E. H., Dorans, K. S., Gold, D. R., Schwartz, J., Koutrakis, P., Washko, G. R.,
 O'Connor, G. T., and Mittleman, M. A.: Long-term exposure to traffic emissions and fine particulate matter and lung
 435 function decline in the Framingham Heart Study, *Am. J. Resp. Crit. Care.*, 191, 656–664, 2015.
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J.
 R., and Pandis, S. N.: Rethinking organic aerosols: Semivolatile emissions and photochemical aging, *Science*, 315, 1259–
 1262, 2007.
- Schauer, J. J., Rogge, W. F., Hildemann, L., M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Source apportionment
 440 of airborne particulate matter using organic compounds as tracers, *Atmos. Environ.*, 30, 3837–3855, 1996.
- Subramanian, R., Donahue, N. M., Bernardo-Bricker, A., Rogge, W. F., and Robinson, A. L.: Contribution of motor vehicle
 emissions to organic carbon and fine particle mass in Pittsburgh, Pennsylvania: Effects of varying source profiles and
 seasonal trends in ambient marker concentrations, *Atmos. Environ.*, 40, 8002–8019, 2006.
- Sun, C., Lee, B. P., Huang, D., Li, Y. J., Schurman, M. I., Louie, P. K. K., Luk, C., and Chan, C. K.: Continuous
 445 measurements at the urban roadside in an Asian megacity by Aerosol Chemical Speciation Monitor (ACSM): particulate
 matter characteristics during fall and winter seasons in Hong Kong, *Atmos. Chem. Phys.*, 16, 1713–1728, 2016.



- Thornhill, D. A., Williams, A. E., Onasch, T. B., Wood, E., Herndon, S. C., Kolb, C. E., Knighton, W. B., Zavala, M.,
Molina, L. T., and Marr, L. C.: Application of positive matrix factorization to on-road measurements for source
apportionment of diesel- and gasoline-powered vehicle emissions in Mexico City, *Atmos. Chem. Phys.*, 10, 3629–3644,
450 2010.
- Transport Department: The Annual Traffic Census 2017, The Government of the Hong Kong Special Administrative
Region, Hong Kong, 2018.
- Wang, Q. Q., He, X., Huang, X. H. H., Griffith, S. M., Feng, Y. M., Zhang, T., Zhang, Q. Y., Wu, D., and Yu, J. Z.: Impact
of secondary organic aerosol tracers on tracer-based source apportionment of organic carbon and $\text{PM}_{2.5}$: A case study in the
455 Pearl River Delta, China, *ACS Earth Space Chem.*, 1, 562–571, 2017.
- Wang, X. L., Ho, K. F., Chow, J. C., Kohl, S. D., Chan, C. S., Cui, L., Lee, S. C. F., Chen, L. W. A., Ho, S. S. H., Cheng, Y.,
and Watson, J. G.: Hong Kong vehicle emission changes from 2003 to 2015 in the Shing Mun Tunnel, *Aerosol Sci.*
Technol., 52, 1085–1098, 2018.
- Weitkamp, E. A., Lambe, A. T., Donahue, N. M., and Robinson, A. L.: Laboratory measurements of the heterogeneous
460 oxidation of condensed-phase organic molecular markers for motor vehicle exhaust, *Environ. Sci. Technol.*, 42, 7950–7956,
2008.
- Wong, Y. K., Huang, X. H. H., Cheng, Y. Y., Louie, P. K. K., Yu, A. L. C., Tang, A. W. Y., Chan, D. H. L., and Yu, J. Z.:
Estimating contributions of vehicular emissions to $\text{PM}_{2.5}$ in a roadside environment: A multiple approach study, *Sci. Total*
Environ., 672, 776–788, 2019.
- 465 Wu, C.: Scatter Plot, Zenodo, <https://zenodo.org/record/832417>, 2017.
- Yao, D. W., Lyu, X. P., Murray, F., Morawska, L., Yu, W., Wang, J. Y., and Guo, H.: Continuous effectiveness of replacing
catalytic converters on liquefied petroleum gas-fueled vehicles in Hong Kong, *Sci. Total Environ.*, 648, 830–838, 2019.
- Yu, J. Z. and Zhang, T.: Chemical speciation of $\text{PM}_{2.5}$ filter samples – January 1 through December 31, 2017, Final report
submitted to the Hong Kong Environmental Protection Department, The Government of the Hong Kong Special
470 Administrative Region (Ref. 16-02709), Hong Kong, 2018.
- Yu, J. Z., Huang, X. H. H., Griffith, S. M., Yuan, Z. B., and Ng, W. M.: Integrated assessment on roadside air quality, Final
report submitted to the Hong Kong Environmental Protection Department, The Government of the Hong Kong Special
Administrative Region (Ref. 12-00255), Hong Kong, 2016.
- Zheng, M., Hagler, G. S. W., Ke, L., Bergin, M. H., Wang, F., Louie, P. K. K., Salmon, L., Sin, D. W. M., Yu, J. Z., and
475 Schauer, J. J.: Composition and sources of carbonaceous aerosols at three contrasting sites in Hong Kong, *J. Geophys. Res.*,
111, D20313, 2006.




Figure 1. Trends in concentrations of (a) OC, (b) EC, (c) NO_x , (d) $\text{PM}_{2.5}$, (e) *n*-butane and (f) *i*-pentane at MK AQMS. Each data point represents the monthly average of the hourly concentrations. Shaded areas represent one standard deviation for the hourly data. Commencements of the Phasing Out Pre-Euro IV Diesel Commercial Vehicles (DCV) Program and Catalytic Converters Replacement Program for LPG taxis are marked in figure (b), (c) and (e). In figure (d), relative contributions to $\text{PM}_{2.5}$ from EC, OM and other components are shown in secondary axis. OM is approximated as $\text{OC} \times 1.4$. Other components are the difference between measured $\text{PM}_{2.5}$ and EC + OM.




Figure 2. Determination of $[\text{OC}/\text{EC}]_{\text{vehicle}}$ for MK AQMS using optimal Deming regression of the lowest 5 % data by OC/EC ratio from summer months. Figure (a) presents the month-by-month $[\text{OC}/\text{EC}]_{\text{vehicle}}$ (grey columns), R^2 of OC and EC (red markers) and number of data point considered (blue markers). Figure (b) shows the frequency of occurrence of the lowest 5 % OC/EC ratios at different hours of the day considering all of the summer data.




Figure 3. Factor profiles resolved by PMF analysis of hourly $\text{PM}_{2.5}$ OC-EC, C_2 - C_8 VOCs and other trace gases data at MK AQMS. Panels on left present the species relative contributions to each factor. Panels on right show the diurnal profile of normalized contribution from each factor in box-plots. For each box, solid square marker, horizontal line, lower and upper bound are mean, median, 25th and 75th percentile, respectively.




Figure 4. Diurnal variations of OC concentrations in box-plots derived from ambient measurement (green boxes) and PMF modeling (grey boxes). For each box, solid square marker, horizontal line, lower and upper bound are mean, median, 25th and 75th percentile, respectively. The afternoon and evening mealtime periods are indicated in the orange frames.




Figure 5. Trends in contributions of diesel and gasoline vehicles at MK AQMS derived from PMF analysis (stacked areas) in terms of (a) $OC_{vehicle}$, (b) $EC_{vehicle}$ and (c) $PM_{vehicle}$ concentrations. Relative contributions of the two vehicle types to the corresponding pollutants are shown in figure (d)–(f). The stacked areas are constructed by interpolation of the monthly data points. In all plots, each data point represents the monthly average of the hourly concentrations. Only months with data cover rate > 33 % are considered. Overall vehicular contributions derived from the EC-tracer method are shown in blue markers in figure (a)–(c).




Figure 6. Trends in PMF-resolved diesel $PM_{vehicle}$ concentrations (left axis) at the MK AQMS during truck-dominated hours (11:00–13:00, blue line), bus-dominated hours (22:00–0:00, brown line), and background hours (2:00–4:00, grey line). Vertical



510 **dashed line represents commencement of the Phasing Out Pre-Euro IV Diesel Commercial Vehicles Program (in March 2014).
Green-shaded area represents the percent completion of the program (right axis).**