

Radiative and chemical implications of the size and composition of aerosol particles in the existing or modified global stratosphere

Daniel M. Murphy¹, Karl D. Froyd^{1,2}, Ilann Bourgeois^{1,2}, Charles A. Brock¹, Agnieszka Kupc^{1,2,3}, Jeff Peischl^{1,2}, Gregory P. Schill^{1,2}, Chelsea R. Thompson^{1,2}, Christina J. Williamson^{1,2}, Pengfei Yu⁴

¹NOAA Chemical Sciences Laboratory, Boulder, CO 80305, USA

²Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309, USA

³Faculty of Physics, Aerosol Physics and Environmental Physics, University of Vienna, 1090 Vienna, Austria

⁴Institute for Environment and Climate Research, Jinan University, Guangzhou, China

Correspondence to: Daniel Murphy (daniel.m.murphy@noaa.gov)

Abstract. The size of aerosol particles has fundamental effects on their chemistry and radiative effects. We explore those effects using aerosol size and composition data in the lowermost stratosphere along with calculations of light scattering. In the size range between about 0.1 and 1.0 μm diameter (accumulation mode), there are at least two modes of particles in the lowermost stratosphere. The larger mode consists mostly of particles produced in the stratosphere and the smaller mode consists mostly of particles transported from the troposphere. The stratospheric mode is similar in the Northern and Southern hemispheres whereas the tropospheric mode is much more abundant in the Northern Hemisphere. The purity of sulfuric acid particles in the stratospheric mode shows that there is limited production of secondary organic aerosol in the stratosphere, especially in the Southern Hemisphere. Out of eight sets of flights sampling the lowermost stratosphere (four seasons and two hemispheres) there were three with large injections of specific materials: volcanic, biomass burning, or dust. The stratospheric and tropospheric modes have very different roles for radiative effects on climate and for heterogeneous chemistry. Because the larger particles are more efficient at scattering light, most of the radiative effect in the lowermost stratosphere is due to stratospheric particles. In contrast, the tropospheric particles can have more surface area, at least in the Northern Hemisphere. The surface area of tropospheric particles could have significant implications for heterogeneous chemistry because these particles, which are partially neutralized and contain organics, do not correspond to the surfaces used for laboratory studies of stratospheric heterogeneous chemistry. We then extend the analysis of size-dependent properties to particles ~~injected into the stratosphere, either intentionally or from volcanoes~~. There is no single size that will simultaneously ~~maximize~~ the climate impact relative to the injected mass, infrared heating, potential for heterogeneous chemistry, and undesired changes in direct sunlight. In addition, light absorption in the far ultraviolet is identified as an issue requiring more study for both the existing and potentially modified stratosphere.

Deleted: considered for intentional climate modification

Deleted: optimize

1 Introduction

Stratospheric particles have been studied for over 50 years (Junge and Manson, 1961; reviewed by Kremser et al., 2016). The stratospheric aerosol layer has a maximum in mixing ratio between about 20 and 25 km altitude. However, the larger air density

35 at lower altitudes means that the majority of the mass of stratospheric aerosol is in the lowermost stratosphere, below the maximum in mixing ratio (Yu et al., 2016).

Various trends have been reported for the background stratospheric aerosol at times not influenced by major volcanic eruptions. Deshler et al. (2006) concluded there was little long-term change in background stratospheric aerosol from 1971 to 2004. 40 Hofmann et al. (2009) found an increasing trend after 2000, and Friberg et al. (2014) found an increasing trend from 1999 to 2008. There has been recognition that moderate volcanic eruptions frequently influence the stratospheric aerosol and true nonvolcanic “background” concentrations are not necessarily present just because there has been no Pinatubo-scale eruption (Solomon et al., 1996, Vernier et al. 2011). Different altitudes may exhibit different trends (Khaykin et al., 2017). Moderate volcanic eruptions tend to mask trends in the non-volcanic background (Kremser et al., 2016). The data shown here reinforce 45 the notion of modest but frequent perturbations to the lower stratosphere.

The overall circulation of air in the stratosphere, with rising air in the tropics and descending air in the extratropics, is mostly fed by air entering at the tropical tropopause. The lowermost stratosphere is a region at middle and high latitudes between the local tropopause and slightly above the altitude of the tropical tropopause. Air in this lowermost stratosphere is affected by 50 both downward motion in the extratropical stratosphere and adiabatic mixing with the troposphere (Holton et al., 1995). The tropospheric influence can extend as high as about 450 K potential temperature (Rosenlof et al., 1997). All of the data described here are in the lowermost stratosphere and show the influence of both air from higher in the stratosphere and air from the troposphere.

55 The chemical composition of particles in the lower stratosphere has been measured by several techniques. Impactor samples collected from the CARIBIC platform have been quantitatively analyzed for N, O, S, K, Fe, and other elements (Nguyen and Martinsson, 2007; Friberg et al, 2014; Martinsson et al., 2019). The moles of oxygen were approximately four times the moles of sulfur, plus about 0.2 times the moles of carbon, indicating SO₄ in sulfate and sulfuric acid plus some contribution from oxygenated organics. Much of the detailed information on the chemical composition of aerosols in the lower stratosphere has 60 come from the Particle Analysis by Laser Mass Spectrometry (PALMS) instrument (Murphy et al., 1998). These data show that most particles larger than about 110 nm fall into three distinct types: sulfuric acid with or without meteoric metals and mixed organic-sulfate particles from the troposphere (Murphy et al., 2014). Because the pure sulfuric acid particles do not contain biomass burning residues, they are not simply tropospheric particles that have lost organics after entering the stratosphere (Murphy et al., 2007). Recent data from another single particle mass spectrometer have found comparable 65 abundances of sulfuric acid particles with meteoric metals (Schneider et al., in discussion).

We extend the previous results to show that the mixed organic-sulfate particles from the troposphere are generally smaller than both types of sulfuric acid particles. We then show how this size difference has significant implications for light scattering and heterogeneous chemistry.

70

2 Methods

This paper includes data in the lowermost stratosphere from the Atmospheric Tomography (ATom) mission with deployments in four seasons during 2016 to 2018. Although the ATom mission was not specifically designed to sample the stratosphere, it encountered stratospheric air in both the Northern and Southern Hemispheres during its regular vertical profiles. Stratospheric air was encountered periodically at altitudes greater than about 7 km and latitudes poleward of about 30 degrees north or 35 degrees south (Figure S1 and Table S1). Because the NASA DC8 aircraft has a ceiling of about 12 km, stratospheric air was always associated with low tropopauses, sometimes in tropopause folds. We therefore use ozone rather than altitude as the primary definition of how far into the stratosphere measurements were taken. If there were mixing with a great deal of tropospheric air it would reduce the ozone concentration below the thresholds we set for stratospheric measurements. Data below 7 km altitude were excluded just in case there were unusually high ozone concentrations at low altitude.

75

80

Size distributions for accumulation-mode particles were measured using two modified commercial laser optical particle spectrometers, an ultra-high sensitivity aerosol spectrometer (UHSAS; Droplet Measurement Technologies, Longmont, USA) from 0.07 to 0.6 μm diameter and a laser aerosol spectrometer (LAS, TSI Inc., St. Paul, USA) from 0.6 to \sim 4.8 μm diameter (Kupc et al., 2018; Brock et al., 2019). The diameters are based on calibration by ammonium sulfate particles. The size resolution of the reported data is 20 bins per decade of particle size. Data are recorded at 1s intervals although averaging is needed in the stratosphere to improve counting statistics for particles in the LAS size range (Brock et al., 2019). Ozone measurements are described by Bourgeois et al., 2020.

85

90

Particle composition was measured with the Particle Analysis by Laser Mass Spectrometry (PALMS) instrument (Thomson et al., 2000; Froyd et al. 2019). A pressure-controlled aerodynamic focusing inlet brings particles into a vacuum where they cross two continuous laser beams. The transit time between the beams measures the aerodynamic diameter of each particle. The aerodynamic diameters are under vacuum conditions with most particles much smaller than the mean free path at the inlet exhaust. Transition flow corrections are considered. Transit times were calibrated to known particle sizes before and after every field deployment. A 193 nm pulse from an excimer laser is triggered when a particle arrives at the second laser beam. Either positive or negative ions are analyzed with a time-of-flight mass spectrometer. The polarity was switched every few minutes. Most of the data shown here are from positive ion spectra. Negative ion spectra do not distinguish sulfuric acid with and without meteoric metals because the metal ions only appear in the positive ion spectra.

95

100

The optical size distributions are combined with the PALMS single-particle composition data for particles larger than 100 nm to create size distributions that are resolved by composition. In each size range, the number of particles is obtained from the optical size distributions and the fraction of particles with different compositions is obtained from PALMS. This combination requires converting the PALMS aerodynamic diameters to correspond to the optical diameters (Froyd et al., 2019). The composition-resolved size distributions presented here use wider size bins than the native optical particle counter resolution but narrower size bins than the standard ATom products (Froyd et al., 2019). The narrower bins are possible because of improved statistics after averaging over all of the data within a specific band of ozone and latitude, even if those data were not contiguous in time.

For the purpose of this study particles are classified into four basic categories: sulfuric acid with and without meteoric metals, mixed organic-sulfate particles, and other particles including dust. Example spectra are shown in Murphy et al. (2014). When appropriate further distinctions can be made, such as separating biomass burning particles from other mixed organic-sulfate particles.

Sources of uncertainty include particle statistics, identifying particle types, the particle volume from the optical particle counters, and combining the PALMS spectra with the optical size distributions. Overall statistics are excellent, with approximately 800,000 single particle mass spectra acquired above 7 km altitude during ATom. Of those, approximately 78,000 were positive ion spectra at ozone concentrations larger than 250 ppbv (Table S1). However, low ambient number densities for large particles result in fewer mass spectra of particles larger than about 1 μm , leading to statistical noise for large particles visible in the figures in this paper. Except for ATom4 Northern Hemisphere, particles larger than 1 μm contributed only a very small fraction of the aerosol volume in the lower stratosphere. Classification uncertainty depends on the type of particle. The meteoric-sulfuric particles are very distinctive. From manually examining mass spectra we estimate that only a few percent are incorrectly classified. Sulfuric acid particles are more difficult to classify because some tropospheric particles with low organic content are similar to stratospheric sulfuric acid particles. The criteria used here probably err on the side of underestimating tropospheric particles in the lower stratosphere, especially in the Southern Hemisphere. Using various criteria for separating organic-sulfate and sulfuric acid particles suggests a classification uncertainty of up to 25% for those particle types in the lower stratosphere. With sufficient averaging (minutes), the volume derived from optical size distributions has an uncertainty propagated from size and flow uncertainties of about +13/-28% in the accumulation mode and up to +/- 50% above 1 μm (Kupc et al., 2018; Brock et al., 2019). Excellent agreement between extinctions calculated from the size distributions and independent extinction measurements indicates that systematic errors may actually be less than this (Brock et al., 2019). Because PALMS measures aerodynamic diameters, mapping the PALMS spectra onto the optical size distributions requires particle density and shape (Froyd et al., 2019). These are well known for sulfuric acid particles, less so for some particles such as dust. How uncertainty in density affects the combination of PALMS and optical data depends on the shape of the size distribution and the number of types of particles. In cases with flat size distributions or where only one dominant type of

Deleted: uncertainty in

Deleted: mean that there are few

Deleted: perhaps

Deleted: if there

particle is present, uncertainty in density introduces very little additional uncertainty when combining PALMS data with size
140 distributions. The uncertainty is larger if there are several types of particles present in a size range with rapidly changing
particle volume versus size. Statistical uncertainty in the fractional organic content of mixed particles is less than 10% if at
least dozens of mass spectra are averaged (Froyd et al., 2019).

Calculations of light scattered back to outer space are made for an optically thin layer uniformly spread over a sunlit hemisphere
145 as described by Murphy (2009), except that these calculations use an atmospheric transmittance appropriate for approximately
11 km altitude (Arvesen, 1969). Changing the solar spectrum over the entire range from top-of-atmosphere to surface gives
qualitatively similar results. At the low relative humidities in the stratosphere, water uptake is less important for optical
properties than it is in the troposphere. The mean relative humidity for the ATom data at ozone > 250 ppbv was less than 10%.

150 A sectional aerosol model (CARMA) coupled with the NSF/DOE Community Earth System Model (CESM) is used in the
study to simulate the composition and size distributions of stratospheric aerosols (Yu et al., 2015; Toon et al., 1988). CESM-
CARMA tracks two external-mixed groups of aerosols. The first group consists of pure sulfate particles (formed through
nucleation and condensation of water vapor and sulfuric acid) with 20 size bins ranging from 0.4 nm to 2.6 μm in diameter;
the second group consists of internal mixed aerosols (containing condensed sulfate, organics, black carbon, salt and dust) with
155 20 size bins from 0.1 μm to 17 μm . The model does not explicitly separate meteoric-sulfuric particles from other sulfuric acid
particles. The model is run at a horizontal resolution of 1.9° (latitude) x 2.5° (longitude). It has 56 vertical layers from the
surface to 1.8 hPa with a vertical resolution of ~ 1 km near the tropopause.

3 Composition-resolved size distributions in the stratosphere

160 Figure 1 shows the composition-resolved size distributions measured in the lowermost stratosphere for the four ATom
deployments, separated by the Northern and Southern Hemisphere. The data are for ozone between 250 and 400 ppbv. This
range of ozone is chosen to be definitely in stratospheric air and to include data from both hemispheres on all four deployments.
For comparison, the median ozone concentration at the lapse rate thermal tropopause was slightly less than 100 ppbv. On each
165 panel the thick black line is the size distribution from the optical particle counters. At each size the fraction of particle types
from PALMS is shaded. A number of features in Figure 1 are worthy of comment.

The volume distributions show a peak near 400 nm diameter and another peak, or at least a shoulder, near 180 or 200 nm.
These sizes are both within what is considered the accumulation mode. The composition reveals why there are two modes.
170 Indeed, without the composition it would be difficult to be sure that there were two separate modes. For example, the Wyoming
particle counters used on stratospheric balloon flights (Deshler et al., 2003) do not clearly resolve the modes.

Deleted: in

175 The 400-nm mode is from sulfuric acid particles produced in the stratosphere, especially those with meteoric metals. The size
of these meteoric-sulfuric particles is extremely consistent through both hemisphere and the four deployments (Table 1). The
primary source of sulfuric acid in the stratosphere, oxidation of carbonyl sulfide, is similar in the two hemispheres. The
meteoric-sulfuric particles also have a narrow size distribution, with a typical geometric standard deviation of about 1.4 when
fit with a log-normal distribution. This is consistent with condensational growth, which tends to lead to narrow size
distributions because smaller particles have relatively more surface area for condensation. Sulfuric acid particles without
180 meteoric material have more diverse sizes except for the volcanically influenced ATom1 Southern Hemisphere, when the
sulfuric acid particles had a narrow size distribution very similar to the meteoric-sulfuric particles.

The smaller mode near 200 nm is from mixed organic-sulfate particles that have mixed into the stratosphere. The mass spectra
of particles in the smaller mode are essentially identical to those of particles in the upper troposphere (Figure S2). Unlike
185 studies that relied solely on bulk composition (e.g. Martinsson et al., 2019), we identify the tropospheric contribution based
on the mass spectra of individual particles (Murphy et al., 2007). Comparing the Northern and Southern Hemispheres in Figure
1, the concentration of the smaller mode is larger in the Northern Hemisphere. The upper troposphere in the Southern
Hemisphere has generally lower aerosol concentrations, so mixing in a given amount of tropospheric air will bring in fewer
particles than the same amount of mixing in the Northern Hemisphere.

190 Of the eight cases in Figure 1, three have much higher aerosol concentrations than the others, for three very different reasons.
The ATom1 [wintertime](#) Southern Hemisphere had a mode of pure sulfuric acid particles. These were most likely produced
from SO₂ injected into the stratosphere by the Calbuco eruption in April 2015, about 16 months before the measurements
(Bègue et al., 2017). The sulfuric acid particles are remarkably pure, except for associated water. This is consistent with
195 previous data on aerosol volatility and infrared spectra after the eruption of Mt. Pinatubo (Deshler et al., 1992; Grainger et al.,
1993). The size distribution is also quite narrow, with a geometric standard deviation σ_g of 1.29 for just the sulfuric acid
particles (Table 1). Satellite aerosol retrievals may use wider size distributions. For example, Bauman et al. (2003) use a lookup
range for σ_g of 1.3 to 2.3 and show that retrievals may not find solutions if $\sigma_g \approx 1.1$. The size distribution of these volcanic
200 particles suggests caution in satellite and lidar retrievals of stratospheric aerosols dominated by one source of particles grown
by condensation of sulfuric acid. As can be seen in Figure 1, overall size distributions typically are not this narrow because
they are broadened with particles from multiple sources.

The ATom3 Northern Hemisphere had a large tropospheric organic-sulfate contribution. More detailed composition shows
that these included a large fraction of biomass burning particles. The ATom3 flights were about two months after Canadian
205 fires produced a massive injection of smoke into the stratosphere, [with additional injection](#) rising through the tropopause due
to diabatic heating (Torres et al., 2020). A separate paper is in preparation about the in-situ data from this wildfire event. The
ATom4 [springtime](#) Northern Hemisphere had both a large contribution from organic-sulfate particles and a remarkable amount

Deleted: ~

Deleted: and also

210 of dust at and above the tropopause. Concentrations at the tropopause often reached several micrograms per standard cubic
meter. The dust was very widespread: it was measured over both the north Atlantic and Pacific Oceans over more than 40
degrees of latitude. This may be Asian dust and pollution carried to high altitude in an event similar to that described by Huang
et al. (2008). The ATom4 and Huang et al. events were both in May. A separate paper is also planned about this dust event.

215 Figure 2 shows the CESM/CARMA model results for the ATom2 flights for the same 250 – 400 ppbv range of ozone as the
data in Figure 1. ATom2 is chosen because neither hemisphere was perturbed by volcanic sulfate, biomass burning, or dust.
The model reproduces the tropospheric mode well in the Northern Hemisphere but overestimates it in the Southern
Hemisphere. The model reproduces the total volume of stratospheric particles well in both hemispheres but the modeled
diameter of these particles is too small. A possible reason is that the model does not include meteoric smoke particles on which
220 sulfuric acid can condense. That is, the model treats both the meteoric-sulfuric and sulfuric acid particles observed by PALMS
as a single type. Figure S3 compares the Northern Hemisphere ATom data to Wilson et al. (2008). Consistent with the ATom
observations, the tropospheric particle mode at about 200 nm is in some cases distinct from the larger stratospheric mode.

Figure 3 shows composition-resolved size distributions further into the stratosphere with ozone between 500 and 850 ppbv.
225 Only the Northern Hemisphere during ATom2 and ATom4 had significant amounts of PALMS data in this ozone range. For
ATom2, the primary difference at higher ozone was more meteoric-sulfuric particles, a result also found by Schneider et al.
(in discussion). The large mixing event of dust and other tropospheric particles during ATom2 barely affected “altitudes” of
more than 500 ppbv ozone. Figure 4 has some of the data in Figures 1 and 3 replotted to emphasize the size distributions of
each component. The size distributions of the meteoric-sulfuric particles are extremely consistent. The size distributions of the
230 sulfuric acid particles without meteoric content are usually broader, possibly indicating more diverse sources. One exception
is the very narrow size distribution of the particles after the Calbuco eruption. The ATom3 Northern Hemisphere has a mode
of larger organic particles from the pyrocumulus injection event. The dust event in the Northern Hemisphere during ATom4
also brought up organic-sulfate and sulfuric particles, or their precursors, from near the surface as well as dust.

235 Figure 4 can also be used to illustrate what features of the size distributions are more certain or uncertain. The positions and
widths of the modes are robust, as are broad features such as the presence of a bimodal organic-sulfate distribution for ATom3.
Narrow variations in the distributions, such as the minimum near 0.55 μm in the ATom3 distribution, could be due to gain
stitching or Mie scattering effects in the optical particle counters. In the bottom panel, the increased concentration of particles
larger than 1 μm during ATom4 compared to the other deployments is robust but the sharpness of the increase at 2 μm could
240 easily be due to limited statistics for large particles.

4 Vertical profiles

245 The shaded regions in Figure 1 can be integrated over all sizes to determine the volume associated with each type of particle,
then multiplied by a density to determine the mass. Figure 5 shows vertical profiles of the volume concentrations for the
meteoric sulfuric particles and organic-sulfate particles. As expected for a high-altitude source, the concentration of meteoric-
sulfuric particles increases with ozone concentration. Except for the ATom3 wildfire biomass burning injection, the
concentration of organic-sulfate particles decreased with increasing ozone or, in one case, stayed roughly constant. The
concentration of the meteoric-sulfuric particles is fairly consistent between hemispheres and deployments. More measurements
250 are needed to see if the small seasonal differences are persistent in other years. In contrast, the concentration of organic-sulfate
particles was larger in the Northern Hemisphere than in the Southern Hemisphere and varied considerably between
deployments. It is worth noting that the highest concentration of tropospheric particles in each hemisphere, ATom3 for the
Southern Hemisphere and ATom4 for the Northern Hemisphere, were both local springtime.

255 Figure 6 shows the ratio of the C⁺ peak, an indicator of organic content, to two peaks indicative of sulfate or sulfuric acid. The
top axis gives an approximate mass fraction of organics adapted from calibrations described by Froyd et al. (2019). The vertical
axis of ozone serves as a measure of distance into the stratosphere. The organic content is separated by particle type, something
not possible with bulk analysis. That the stratospheric and tropospheric particle compositions remain distinct implies that there
is very limited redistribution of semi-volatile organics between particles. Like most upper tropospheric particles, the organic-
260 sulfate particles are internally mixed with on average about 40 to 80 percent organic material by mass. There is little variation
with ozone, indicating a long lifetime for the organic material as well as little uptake of sulfuric acid. The latter is consistent
with most of the sulfuric acid coming from carbonyl sulfide above 20 km rather than SO₂ near the tropopause (Kremser et al.,
2016; Rollins et al., 2017). That there are two types of particles with different compositions present in the same air also means
that any semi-volatile gas-phase organics and ammonia cannot be in equilibrium with both types of particles.

265 There is some limited uptake of organics onto the stratospheric particles, although the maximum organic concentration is still
much less than for tropospheric particles. Meteoric-sulfuric particles definitely formed in the stratosphere, so any significant
organic content indicates net uptake of organics. Their organic content grows as the particles descend to the lowermost
stratosphere and the upper troposphere. The meteoric-sulfuric particles contain much less than 1% organic mass at altitudes
270 with ozone greater than 500 ppbv and as much as 2 to 4% near the tropopause in the Northern Hemisphere. Such limited
formation of secondary organic mass in the lowermost stratosphere is consistent with previous PALMS measurements
(Murphy et al., 2007).

A new finding from ATom is that there is a very distinct and consistent difference between the hemispheres in the small amount
275 of organic content that does form in the meteoric-sulfuric acid particles. Since the particles start from similar formation
processes much higher in the stratosphere, we conclude that there is more condensable or reactive organic vapor in the Northern

Deleted: give

Deleted: give

280 Hemisphere lower stratosphere. This could be either gas phase species mixed from the troposphere or semi-volatile organics transferring from organic-sulfate particles.

285 Adding a few percent mass to the meteoric-sulfuric particles represents a very small amount of organic vapor. Without knowing uptake coefficients the amount of vapor cannot be determined uniquely, but a representative calculation is that one or two pptv of an organic gas-phase species with molecular weight of about 100 daltons that reacts with sulfuric acid on every collision would add few percent mass to a 450 nm particle in a few months. The same order of magnitude can be obtained by noting from Figure 5 that at 200 ppbv ozone there is about 100 ng standard m^{-3} of meteoric sulfuric particles. 1% by mass of these particles corresponds to about a part per trillion by mass of air. We conclude that an order of magnitude for highly condensable organic vapor in the lowermost stratosphere is a few parts per trillion in the Northern Hemisphere and less in the Southern Hemisphere. A less reactive or condensable organic molecule could be present at a correspondingly higher concentration.

290

5 Radiative and chemical implications

The different sizes of the sulfuric acid and organic-sulfate particles lead to substantial differences in their radiative and chemical effects. Important properties are the amount of infrared heating, the amount of light scattered, implications for 295 photolysis, the surface area available for heterogeneous chemistry, and the mass sedimentation rate.

A key part of the radiative implications is the efficiency of light scattering as a function of particle size. Figure 7 shows the mass scattering efficiency as a function of particle size averaged over the solar spectrum and a sunlit Earth. Calculations are for a real refractive index of 1.45 and minimal absorption. Atmospheric extinction is determined by the solid total scattering 300 curve. Much of the light scattered by particles continues downward to the Earth and so does not directly affect climate. Separating out the light scattered to outer space (dashed curve) gives a maximum that is slightly broader and shifted to smaller sizes than light extinction. Over much of the size range of particles that scatter light efficiently, only about 1/5 of the light that is scattered goes to outer space; the remainder becomes diffuse light. This is a reason for the large increases in diffuse light (with decreases in direct sunlight) after volcanic eruptions (Murphy, 2009).

305

The top panel of Figure 8 replots the chemically resolved size distribution for ATom2 in the [wintertime](#) Northern Hemisphere. This was chosen as an example because it is similar to several other locations and seasons such as ATom1 [summertime](#) Northern Hemisphere and ATom3 [springtime](#) Southern Hemisphere. There was more sampling time in the ATom2 Northern Hemisphere stratosphere so the particle statistics are better, with [mass spectra of](#) about 10,000 particles. In Figure 8 the two 310 sulfuric acid particle categories have been combined. Of the particles larger than 0.1 μm diameter, about 39% of the volume was organic-sulfate particles from the troposphere and 61% sulfuric acid particles from the stratosphere (including both those with and without meteoric metals). The percentage contribution to each parameter in Figure 8 by stratospheric aerosol will be

somewhat larger at ambient conditions because sulfuric acid has some water uptake even at <10% relative humidity. Ambient sulfuric acid particles may have roughly 5 to 15% larger diameters than measured in the warm aircraft cabin.

315

5.1 Infrared heating

An important property for stratospheric particles is their absorption and emission of infrared radiation. Infrared properties are more important for stratospheric aerosol than very low-altitude aerosol because the latter are close to the surface temperature and so absorb and emit similar amounts of energy.

320

Infrared absorption by stratospheric particles is important for two reasons. First, it heats the stratosphere around the particles. Changes in circulation due to infrared heating were responsible for significant changes in ozone after the Pinatubo eruption (Labitze and McCormick, 1992; Pitari and Rizi, 1993). The heating-induced changes in ozone were as large or larger than those due to heterogeneous chemistry. There are additional feedbacks on the circulation after changes in infrared heating due to changes in ozone and water vapor (Visoni et al., 2017). Infrared heating is largest in the lower stratosphere where the temperature contrast with the surface is greatest (Lacis, 2015).

325

Second, infrared absorption by stratospheric particles offsets some or even all of the shortwave cooling of the Earth. For sulfuric acid particles similar to those after the Pinatubo eruption, longwave heating offset roughly 25% of the shortwave cooling (Hansen et al., 2005). This increases to about 50% for large injection rates because larger particles (>0.6 μm) become increasingly less efficient at scattering sunlight to outer space (Figure 7) compared to their volume (Niemeier and Timmreck, 2015). Large particles can even cause net warming: alumina with the size distribution from rocket emissions was calculated to cause net warming (Ross and Sheaffer, 2014).

330

335

For wavelengths much larger than the particles, absorption and emission are approximately proportional to total particle volume (and the material) and do not depend on particle size (van de Hulst, 1981). Therefore, net thermal infrared heating is insensitive to the size of particles and will approximately follow the volume distribution in the top panel of Figure 8. For ATom2, most of the volume distribution was composed of particles of stratospheric origin, but that was not always the case (Figure 1). The infrared effects of the tropospheric particles found in the lower stratosphere are hard to assess. Longwave radiative heating depends not only on the strength of the absorption bands but also on their overlap with atmospheric windows in the infrared spectrum. The presence of a significant fraction of organic material has unknown implications for infrared heating. Like sulfate, infrared heating will be proportional to the volume of organic aerosol, but the absolute amount of heating depends on how aerosol absorption features correlate with gas-phase absorption or window regions in the infrared spectrum.

340

345

We are not aware of any infrared spectra of organic material in aerosols in the stratosphere or upper troposphere.

5.2 Scattering of visible light

350 The middle panel in Figure 8 shows the size distribution weighted by the amount of sunlight scattered to outer space, which is
relevant for shortwave climate effects. Weighting the size distribution by the light extinction would shift the peak just slightly
further to larger particles. Because of their size, the sulfuric acid particles contribute a greater fraction of the light scattering
than their mass fraction. In fact, comparing Figures 1 and 7, the most abundant size of the sulfuric acid particles in the lower
stratosphere was close to the maximum in light scattering to outer space per unit mass. The ATom mission [took place](#) during
355 a time with small volcanic influence. In contrast, particles shortly after the Pinatubo eruption had volume mean diameters
greater than $0.7\ \mu\text{m}$ (Wilson et al., 2008; Figure S3), large enough that their mass scattering efficiency decreased. Figure S4
shows the relative extinction due to various particle types for cases other than the ATom2 Northern Hemisphere used as an
example in Figure 8.

5.3 Heterogeneous chemistry

360 Particle size can affect heterogeneous chemistry. Reactions with sulfuric acid particles that are important to stratospheric
chemistry span the range from reactions that occur in the interior of liquid particles and hence are proportional to volume to
reactions that occur on the surface and hence are proportional to surface area (Hanson et al., 1994). Heterogeneous chemistry
can be especially important within or at the edge of the polar vortex (Solomon et al., 2015; Stone et al., 2017).

365 The bottom panel of Figure 8 shows the size distribution weighted by surface area rather than volume. For these conditions
surface reactions are closely proportional to surface area; gas-phase diffusion is a minor correction. For the case shown in
Figure 8, the organic-sulfate particles from the troposphere are about half of the surface area in the lowermost stratosphere.
This is significant because whereas stratospheric heterogeneous chemistry on sulfuric acid has been extensively studied, little
370 is known about the same reactions on organic-sulfate particles. Figure 8 shows ATom2 Northern Hemisphere as an example.
Surface area fractions for other cases are shown in Figure S4.

The organic-sulfate particles differ from the sulfuric acid particles in important ways. Most obviously, they contain a high
proportion of organics that may participate in new chemistry with halogen radicals. Iodine in particular may react with organic
375 aerosols (Murphy and Thomson, 2000; Pechtl et al., 2007). Although not fully neutralized, the organic-sulfate particles are not
nearly as acidic as the relatively pure sulfuric acid particles. This can be determined from acid cluster peaks in the PALMS
mass spectra. Some chlorine activation reactions that lead to ozone destruction are acid-catalyzed (Burley and Johnston, 1992)
and therefore may be slower on partially neutralized particles. The organic-sulfate particles also contain less water – sulfuric
acid is extremely hygroscopic compared to other species at the low relative humidities in the stratosphere. The availability of

380 condensed water for heterogeneous reactions could be further reduced if the organic-sulfate particles are glassy at the low
temperatures and humidity in the stratosphere (Krieger et al., 2012; Price et al., 2014).

5.4 Sedimentation

385 Sedimentation is a key process in the stratospheric aerosol budget (Wilson et al., 2008). It is more important than in the
stratosphere than it is near ground level partly because particles fall faster at lower air density. A bigger reason sedimentation
is important in the stratosphere is the relevant time scale: a fall speed of a kilometer per month would be unimportant in the
lower troposphere but can control the residence time of a particle in the stratosphere. Particles larger than about 1 μm diameter
have sedimentation rates greater than 10 km yr^{-1} in the lower stratosphere. The sedimentation flux as a function of size, shown
390 in Figure S5, is similar to the light scattering panel in Figure 8). For the ATom2 example used in Figure 8, the sulfuric acid
particles have roughly twice the volume of the organic-sulfate particles. Their source strength must be somewhere between
about twice as large as the tropospheric particles (if loss is controlled by bulk air motion) and three times as large (if loss is
controlled by sedimentation).

395 5.5 Ultraviolet scattering and absorption

Absorption and scattering of ultraviolet light are distinct from that of visible light because of the impact on photolysis rates.
Light scattering by stratospheric aerosol changes the path length of light in the stratosphere, which in turn changes photolysis
rates (Huang and Massie, 1997; Pitari et al., 2015). The calculations are complex because Rayleigh scattering in the ultraviolet
400 leads to strong effects from multiple scattering (Bian and Prather, 2002), especially at twilight or if a scattering aerosol layer
is located above gas-phase absorption (Davies et al., 1993; Anderson et al., 1995). In addition, the long path lengths magnify
the importance of any absorption of ultraviolet light by aerosols.

The scattering of ultraviolet light peaks at smaller particle sizes than for sunlight. This means that the smaller tropospheric-
405 organic particles contribute substantially to the scattering of ultraviolet light. The relative contributions to scattering of light
at < 240 nm are shown in Figure S5 and are similar to the surface area panel in Figure 8 except that sizes smaller about 80 nm
and larger about 600 nm contribute less to UV scattering than they do to surface area.

One important wavelength band is 200 to 242 nm, where photolysis of O_2 is responsible for formation of ozone and photolysis
410 of N_2O produces odd nitrogen (NO_x). (Brasseur and Solomon, 1986). For purely scattering particles, changes in photolysis in
this wavelength range are reduced by large cancellations in direct and diffuse light (Michelangeli et al., 1989). Light scattering
by the El Chichon volcanic cloud was estimated to reduce O_2 photolysis by about 10% (Michelangeli et al., 1989). The overall
effect of scattering seems to be to reduce ozone formation (Pitari et al., 2015).

415 Unlike pure scattering particles, absorbing particles would not have a similar partial cancellation between direct and diffuse
sunlight. Huang and Massie (1997) examined the effect of substituting ash with visible and UV absorption for non-absorbing
sulfuric acid in a simple model of photolysis after a volcanic eruption. There are competing effects on the ozone column
because both J_{O_2} and J_{O_3} are reduced by UV absorption, with one reducing and the other increasing ozone (Pitari and Rizzi,
1993). The individual effects were several percent of the ozone column with the net impact difficult to assess because their
420 simple model did not include NO_x or halogen chemistry. The imaginary refractive index of the organic-sulfate particles at
wavelengths below 242 nm is not known but could easily be large enough to lead to significant absorption compared to
scattering.

6 Relevance to volcanic or intentional aerosol injection

425 Seeing the different radiative efficiency and surface area per unit mass of the native stratospheric particles compared to the
particles transported from the troposphere motivates more general calculations about the radiative and chemical effects of
particles as a function of size. Because we are concerned with the climate implications of stratospheric particles we will
compare the sunlight scattered to outer space with various parameters such as surface area and sedimentation rate. The presence
430 of more than one type of particle in the stratosphere also motivates investigating why different types of particles have different
sizes.

Sulfur dioxide is periodically injected into the stratosphere by major volcanic eruptions, leading to an increase in sulfuric acid
aerosol. One such example is visible in the ATom1 Southern Hemisphere panel in Figure 1. There have also been numerous
435 proposals for, and studies of, injecting material into the stratosphere for the purpose of solar radiation management (National
Research Council, 2015). Regardless of the desirability of such actions, some of the calculations presented above on the optical
properties and potential for heterogeneous chemistry have implications for added material whether it be from volcanoes or
deliberate intervention. Even if various types of particles have similar optical properties, the chemically resolved size
distributions are important for attributing radiative effects to different sources. Without information about particle composition
440 we could not say, for example, what percentage of extinction is due to particles formed in the stratosphere and what percentage
is due to particles formed in the troposphere.

6.1 Complex controls on particle size

445 The previous sections demonstrate that particle size is important for many of the properties relevant to climate and chemical
effects. In the real world, one cannot instantly fill a box with monodisperse particles of a chosen size, the way one might in a
model. It is important to understand what controls the size distribution of particles in the stratosphere and how it will change

with additional aerosol or precursors such as sulfur dioxide. The mean **particle** diameter in the stratosphere is not a **constant** but varies with the aerosol loading, altitude, and latitude (English et al., 2012).

Deleted: single number

450

The size of particles in the unperturbed or perturbed stratosphere can be understood in two complementary ways. The first way, a top-down approach, says that for a given mass of stratospheric aerosol, the more particles there are, the smaller they must be. The second way, a bottom-up approach, considers how the size of each particle is set by a balance of growth and removal processes in the stratosphere.

455

For the top-down approach, one must consider at least three sources of particle number in the stratosphere. Particles come down from the mesosphere, up from the troposphere, and new particles can form in the stratosphere (Murphy et al., 2014). The meteoric source of particles to the stratosphere is mostly “smoke” consisting of material that evaporated from ablating meteoroids and condensed into new particles high in the atmosphere. Much of this material descends near the winter poles (Bardeen et al., 2008). Second, tropospheric particles provide an important source of stratospheric particles below 20 km altitude (Yu et al., 2016). The fate of tropospheric particles entering the stratosphere is poorly represented in most models. New particle formation is also important for the stratosphere. The pure sulfuric acid category in Figure 1 is probably from growth of particles formed in (or at the edge of) the stratosphere. One formation region is near the tropical tropopause with upward transport into the stratosphere (Brock et al., 1995; English et al., 2011). There is probably also formation of new sulfuric acid particles higher in the stratosphere over the winter poles (Wilson et al., 1989), although this must be distinguished from meteoric smoke descending in the same regions (Curtius et al., 2005).

460

465

Adding sulfuric acid or its precursors will have complex effects on new particle formation, with more vapor to condense but also more surface area sink. In contrast, injected solid particles would provide a surface sink for background sulfuric acid from oxidation of carbonyl sulfide, likely reducing new particle formation. This implies that injected solid particles would probably change the size of the natural sulfuric acid particles in the stratosphere. Sufficiently small injected solid particles might reach high altitudes where existing sulfuric acid particles have evaporated (Weisenstein et al., 2015; Jones et al., 2016). There could be unknown effects if they were later entrained in descending air in the winter polar regions.

470

475

The bottom-up approach considers how the size of stratospheric particles is determined by a balance of growth and removal processes. Particles grow by coagulation and by condensation of sulfuric acid and other species. Coagulation in the unperturbed stratosphere is slow except for special situations such as shortly after new particle formation (Brock et al., 1995; Hamill et al., 1997). Coagulation increases non-linearly with aerosol concentration so it becomes more significant after volcanic eruptions (Pinto et al., 1989) or large injection scenarios (Weisenstein et al., 2015). In these cases, coagulation helps drive the extra mass primarily to larger particles rather than more numerous particles (Heckendorn et al., 2009; Niemeier and Timmreck, 2015). Both sedimentation and downward motion are important removal processes (Wilson et al., 2008).

480

One implication of having multiple sources of particles in the stratosphere is that there is no single response to injected material.

485 It is only in the last few years that stratospheric models have incorporated multiple sources of particles along with detailed microphysics (Pitari et al. 2014; Yu et al., 2016; Mills et al., 2017). There is still considerable uncertainty in quantitatively understanding the size of particles in the current stratosphere, let alone after a perturbation. Figure 2 demonstrates that a detailed microphysics model of the stratosphere did not grow the sulfuric acid particles to large enough sizes.

490 6.2 Impacts on the stratosphere as a function of size

The preceding section suggests that the ultimate size of particles is not something that can simply be chosen when injecting aerosol particles or their precursors; instead it is also strongly influenced by subsequent particle formation and growth processes in the stratosphere. Even solid particles may coagulate or grow by condensation of sulfuric acid and not retain their
495 original size. Despite this, it is of interest to examine the impact of particle size on the relative magnitude of effects. Figure 9 shows an estimate of net cooling compared to particle volume, surface area, and sedimentation rate.

The shortwave effect is modified to include longwave effects by adapting a calculation ~~from~~ Lacis et al. (1992), who showed that heating exceeded shortwave cooling for several different types of particles when their diameters were more than about 4
500 μm . Since infrared absorption is almost independent of size, an approximate net cooling was estimated by subtracting a constant from the calculated shortwave scattering to space such that the net was zero at 4 μm diameter. Figure 7 shows the infrared effect on net cooling as a function of size. The vertical scale of the net cooling curve is consistent with the scattering curves. For example, at peak scattering efficiency near 0.5 μm , infrared effects reduce the cooling of the Earth by stratospheric aerosol by 10 to 15%. The relative infrared effect is larger for either larger or smaller particles. This calculation is only
505 approximate but the patterns in Figure 9 are not sensitive to the exact amount subtracted from cooling by pure scattering. The relative sedimentation curve in Figure 9 also varies slightly with altitude but again the patterns are similar.

The size that has the least infrared absorption for a given amount of shortwave climate impact is about 0.5 μm diameter (filled circles). This is true regardless of details of the infrared spectrum of the particles. Larger or smaller particles will be less
510 effective at cooling the Earth and will cause more stratospheric circulation changes for a given amount of cooling. Sufficiently large ($> 4 \mu\text{m}$) or small ($< 0.1 \mu\text{m}$) particles cause net heating of the Earth (Lacis et al., 1992). The crossing point between net cooling and net heating will depend on particle composition. Even for the optimal size, the infrared heating due to deliberate injections of sulfuric acid or its precursors into the stratosphere would cause significant changes in circulation (Aquila et al.,
515 2014).

Deleted: in

The potential for increased heterogeneous chemistry would be reduced by using larger particles with less surface area. Particles with a diameter of about 1 μm have the largest cooling effect for a given surface area (open circles). These larger particles, however, have high sedimentation rates (downward triangles) and limited stratospheric lifetimes. For a given climate impact, the mass flux due to sedimentation is minimized by particles with a diameter of about 300 nm. The least diffuse light is created by the smallest particles (Figure 7). The exact sizes of the various maxima depend on the refractive index and density. However, the patterns of which variables are maximized at larger or smaller sizes will be similar.

7 Discussion

The title of Solomon et al. (2011) includes the phrase “the persistently variable ‘background’ stratospheric aerosol”. The ATom data presented here add new meaning to that phrase. Out of eight samplings of the lowermost stratosphere, three exhibited much higher aerosol concentrations for three different reasons: a volcanic eruption, biomass burning aerosol, and transport of dust and other near-surface particles. None of these events were targeted during the flights. Such variations in the stratospheric aerosol layer are important for both heterogeneous chemistry (Solomon et al., 1996) and climate (Solomon et al., 2011).

There are important differences in the aerosol in the lower stratosphere between the Northern and Southern Hemispheres. A smaller amount of tropospheric aerosol in the Southern Hemisphere stratosphere indicates that the tropospheric particles are mixing into the lower stratosphere within each hemisphere rather than entering in rising air in the tropics and splitting into the two hemispheres. Sulfuric acid particles in the Southern Hemisphere also acquire less organic content. This suggests that there are lower concentrations of gas-phase organics in the Southern Hemisphere. One of several possible formation routes is that small organic compounds such as acetone and formaldehyde can react with concentrated sulfuric acid to form polymers that stay in the aerosol (Iraci and Tolbert, 1997; Williams et al., 2010). Other routes would be if low-volatility organic molecules were formed in the gas phase or evaporated from the tropospheric particles and recondensed on the sulfuric acid particles. Even in the Northern Hemisphere, only low part-per-trillion range concentrations of gas-phase organics are required to explain the very small amounts of organics taken up by the sulfuric acid particles.

The data here add support to the concept of Yu et al. (2016) that tropospheric particles comprise a significant fraction of the aerosol in the lowermost stratosphere. Such tropospheric particles offer a route for anthropogenic influence on the stratosphere. The Yu et al. model also correctly predicts that tropospheric particles are smaller than sulfuric acid particles formed in the stratosphere (Figure 2).

The data here have several implications for satellite retrievals. First, the reason lower stratospheric size distributions are often broad is that they are really two or more overlapping distributions. As the tropospheric/stratospheric mix shifts the mean diameter will shift. Second, unlike sulfuric acid, the smaller, organic-rich particles coming up from the troposphere may be

effloresced and/or glassy. That could explain a very small amount of depolarization. Finally, when only the stratospheric mode is dominant the size distribution can be very narrow.

555 Absorption of ultraviolet light means that impurities should be considered when assessing deliberately added materials. For example, absorption appropriate for optical-quality sapphire should probably not be used when evaluating proposals to add industrial quantities of alumina to the stratosphere. Even part-per-million impurities in alumina increase absorption in the ultraviolet (Innocenzi et al., 1990). Compared to many materials, sulfuric acid has extremely low absorption in the ultraviolet (Noziere and Esteve, 2005; Dykema et al., 2016).

560 The broad distribution of particle sizes in the unperturbed stratosphere is the superposition of several narrower distributions. Single particle types, particularly meteoric-sulfuric acid particles, can have narrow size distributions (lognormal standard deviation ~ 1.4).

565 Multiple formation mechanisms for stratospheric particles imply that the size of particles after a volcanic or intentional injection may be difficult to predict. Yet an accurate prediction of size is important: The diameter must be known to perhaps 25% to accurately estimate tradeoffs between climate impact and side effects (Figure 9). A state-of-the-art microphysical bin model underestimates the size of stratospheric sulfuric acid particles, indicating that we do not fully understand what controls the size of particles in the stratosphere. The size difference has significant impacts on properties: the modeled particles have about 65% of the climate impact per unit mass as calculated from observed sizes of stratospheric sulfuric acid particles, 160% of the surface area, and sediment about 60% as fast.

575 There is no single diameter that produces the largest shortwave climate impact with the fewest side effects (Figure 9). To the extent that one could control the size of particles after an intentional injection, any chosen size involves tradeoffs. Particles smaller than about $0.6 \mu\text{m}$ diameter have more surface area for possible heterogeneous chemistry. Particles larger than about $0.4 \mu\text{m}$ require more injected mass and produce more diffuse light. For a given amount of scattered sunlight, either sufficiently large or small particles have more infrared absorption and hence more impacts on stratospheric circulation. Most of the mass of particles after the Mt. Pinatubo eruption was larger than $0.6 \mu\text{m}$ diameter (Brock et al., 1993; Wilson et al. 2008), a size range with relatively little surface area compared to their climate impact. The heterogeneous chemistry observed after Mt. Pinatubo may therefore underestimate what might happen with intentionally added material.

580

Data availability and author contributions

Data are available at <https://daac.ornl.gov/ATOM/campaign/>, <https://espo.nasa.gov/atom/content/ATom>, and <https://www.esrl.noaa.gov/csl/projects/atom/data/MurphyACP2020/>. DMM wrote the manuscript with assistance from all

other authors. DMM, KDF, CAB, AK, GPS, and CJW made aerosol measurements and analysed the data. IB and CRT made
585 ozone measurements and analysed the data. PY performed model calculations. Authors declare no competing interests.

Acknowledgements

T. Ryerson helped provide ozone measurements. J. C. Wilson provided data from his 2008 paper. These measurements were
590 supported by NOAA internal climate funding and also in part by NASA award NNH15AB12I. The ATom mission as a whole
was supported by NASA's Earth System Science Pathfinder Program EVS-2 funding. A. Kupc was supported by the Austrian
Science Fund's Erwin Schrodinger Fellowship J-3613.

References

- 595 Anderson, D. E., DeMajistre, R. and Lloyd, S. A.: Impact of aerosols and clouds on the troposphere and stratosphere
radiation field with application to twilight photochemistry at 20 km, *J. Geophys. Res.*, 100, 7135-7145,
<https://doi.org/10.1029/95JD00024>, 1995.
- Aquila, V., Garfinkel, C. I., Newman, P. A., Oman, L. D., and Waugh, D. W.: Modifications of the quasi-biennial oscillation
600 by a geoengineering perturbation of the stratospheric aerosol layer, *Geophys. Res. Lett.*, 41, 1738-1744,
[doi:10.1002/2013GL058818](https://doi.org/10.1002/2013GL058818), 2014.
- Arvesen, J. C., Griffin, Jr., R. N., and Pearson, Jr., B. D.: Determination of extraterrestrial solar spectral irradiance from a
research aircraft, *Appl. Optics*, 8, 2215-2232, <https://doi.org/10.1364/AO.8.002215>, 1969.
- 605 Bardeen, C. G., Toon, O. B., Jensen, E. J., Marsh, D. R., and Harvey, V. L.: Numerical simulations of the three-dimensional
distribution of meteoric dust in the mesosphere and upper stratosphere, *J. Geophys. Res.*, 113, D17202,
[doi:10.1029/2007JD009515](https://doi.org/10.1029/2007JD009515), 2008.
- 610 Bauman, J. J., Russell, P. B., Geller, M. A., and Hamill, P.: A stratospheric aerosol climatology from SAGE II and CLAES
measurements: 1. Methodology, *J. Geophys. Res.*, 108, 4382, [doi:10.1029/2002JD002992](https://doi.org/10.1029/2002JD002992), 2003.
- Bègue, N., Damien Vignelles, D., Berthet, G., Portafaix, T., Payen, G., Jégou, F., Bencherif, H., Jumelet, J., Vernier, J.-P.,
Lurton, T., Renard, J.-B., Clarisse, L., Duverger, V., Posny, F., Metzger, J.-M., and Godin-Beekmann, S.: Long-range

- 615 transport of stratospheric aerosols in the Southern Hemisphere following the 2015 Calbuco eruption, *Atmos. Chem. Phys.*,
17, 15019–15036, <https://doi.org/10.5194/acp-17-15019-2017>, 2017.
- Bian, H., and Prather, M. J.: Fast-J2: Accurate simulation of stratospheric photolysis in global chemical models, *J. Atmos.*
Chem., 41, 281-296, <https://doi.org/10.1023/A:1014980619462>, 2002.
- 620 Bourgeois, I., Peischl, J., Thompson, C. R., Aikin, K. C., Campos, T., Clark, H., Commane, R., Daube, B., Diskin, G. W.,
Elkins, J. W., Gao, R.-S., Gaudel, A., Hints, E. J., Johnson, B. J., Kivi, R., McKain, K., Moore, F. L., Parrish, D. D.,
Querel, R., Ray, E., Sánchez, R., Sweeney, C., Tarasick, D. W., Thompson, A. M., Thouret, V., Witte, J. C., Wofsy, S. C.,
and Ryerson, T. B.: Global-scale distribution of ozone in the remote troposphere from the ATom and HIPPO airborne field
625 missions, *Atmos. Chem. Phys.*, 20, 10611–10635, [10.5194/acp-20-10611-2020](https://doi.org/10.5194/acp-20-10611-2020), 2020.
- Brasseur, G., and Solomon, S.: *Aeronomy of the Middle Atmosphere*, Kluwer Academic Publishers, Dordrecht, Holland,
2nd Edition, ISBN 9781402038241, 1986.
- 630 Brock, C. A., Jonsson, H. H., Wilson, J. C., Dye, J. E., Baumgardner, D., Borrmann, S., Pitts, M. C., Osborn, M. T.,
DeCoursey, R. J., and Woods, D.C.: Relationships between optical extinction, backscatter, and aerosol surface area and
volume in the stratosphere following the eruption of Mt. Pinatubo, *Geophys. Res. Lett.*, 20, 2555-2558,
<https://doi.org/10.1029/93GL01691>, 1993.
- 635 Brock, C. A., Hamill, P., Wilson, J. C., Jonsson, H. H., and Chan, K. R.: Particle formation in the upper tropical troposphere
– A source of nuclei for the stratospheric aerosol, *Science*, 270, 1650-1653, [10.1126/science.270.5242.1650](https://doi.org/10.1126/science.270.5242.1650), 1995.
- Brock, C. A., Williamson, C., Kupc, A., Froyd, K. D., Erdesz, F., Wagner, N., Richardson, M., Schwarz, J. P., Gao, R.-S.,
Katch, J. M., Campuzano-Jost, P., Nault, B. A., Schroder, J. C., Jimenez, J. L., Weinzierl, B., Dollner, M., Bui, T., and
640 Murphy, D. M.: Aerosol size distributions during the Atmospheric Tomography Mission (ATom): methods, uncertainties,
and data products, *Atmos. Meas. Tech.*, 12, 3081-3099, <https://doi.org/10.5194/amt-12-3081-2019>, 2019.
- Burley, J. D., and Johnston, H. S.: Ionic mechanisms for heterogeneous stratospheric reactions and ultraviolet
photoabsorption cross sections for NO₂⁺, HNO₃, AND NO₃⁻ in sulfuric acid, *Geophys. Res. Lett.*, 19, 1359-1362, doi
645 [10.1029/92GL01115](https://doi.org/10.1029/92GL01115), 1992.
- Curtius, J., Weigel, R., Vössing, H.-J., Wernli, H., Werner, A., Volk, C.-M., Konopka, P., Krebsbach, M., Schiller, C., Roiger,
A., Schlager, H., Dreiling, V., and Borrmann, S.: Observations of meteoric material and implications for aerosol nucleation

- in the winter Arctic lower stratosphere derived from in situ particle measurements, *Atmos. Chem. Phys.*, 5, 3053-3069,
650 <https://doi.org/10.5194/acp-5-3053-2005>, 2005.
- Cziczo, D. J., Thomson, T.S., Murphy, D. M.: Ablation, flux, and atmospheric implications of meteors Inferred from
stratospheric aerosol, *Science*, 291, 1772-1775, 10.1126/science.1057737 2001.
- 655 Davies, R.: Increased transmission of ultraviolet radiation to the surface due to stratospheric scattering, *J. Geophys. Res.*, 98,
7251-7253, <https://doi.org/10.1029/93JD00408>, 1993.
- Deshler, T.: A review of global stratospheric aerosol: measurements, importance, life cycle, and local stratospheric aerosol,
Atmos. Res., 90, 223-232, doi.org/10.1016/j.atmosres.2008.03.016, 2008.
- 660 Deshler, T., Hofmann, D. J., Johnson, B. J., and Wozier, W. R.: Balloonborne measurements of the Pinatubo aerosol size
distribution and volatility at Laramie, Wyoming during the summer of 1991, *Geophys. Res. Lett.*, 19, 199-202, 1992.
- Deshler T., Hervig, M. E., Hofmann, D. J., Rosen, J. M., and Liley, J. B.: Thirty years of in situ stratospheric aerosol size
665 distribution measurements from Laramie, Wyoming (41°N), using balloon-borne instruments, *J. Geophys. Res.*, 108: article
4167, [doi:10.1029/2002JD002514](https://doi.org/10.1029/2002JD002514), 2003.
- Deshler, T., Anderson-Sprecher, R., Jäger, H., Barnes, J., Hofmann, D. J., Clemesha, B., Simonich, D., Osborn, M.,
Grainger, R. G., and Godin-Beekmann, S.: Trends in the nonvolcanic component of stratospheric aerosol over the period
670 1971 – 2004, *J. Geophys. Res.*, 111, D01201, [doi:10.1029/2005JD006089](https://doi.org/10.1029/2005JD006089), 2006.
- Ditas, J., Ma, N., Zhang, Y., Assmann, D., Neumaier, M., Riede, H., Karu, E., Williams, J., Scharffe, D., Wang, Q., Saturno,
J., Schwarz, J. P., Katic, J. M., McMeeking, G. R., Zahn, A., Hermann, M., Brenninkmeijer, C. A. M., Andreaeg, M. O.,
Pöschl, U., Sua, H., and Cheng, Y.: Strong impact of wildfires on the abundance and aging of black carbon in the lowermost
stratosphere, *P. Natl. Acad. Sci. USA*, 115, E11595-E11603, <https://doi.org/10.1073/pnas.1806868115>, 2018.
- 675 Dykema, J. A., Keith, D. W., and Keutsch, F. N.: Improved aerosol radiative properties as a foundation for solar
geoengineering risk assessment, *Geophys. Res. Lett.*, 43, 7758-7766, [doi:10.1002/2016GL069258](https://doi.org/10.1002/2016GL069258), 2006.
- English, J. M., Toon, O. B., Mills, M. J., and Yu, F.: Microphysical simulations of new particle formation in the upper
troposphere and lower stratosphere, *Atmos. Chem. Phys.*, 11, 9303-9322, [doi:10.5194/acpd-11-12441-2011](https://doi.org/10.5194/acpd-11-12441-2011), 2011.
- 680

- English, J. M., Toon, O. B., and Mills, M. J.: Microphysical simulations of sulfur burdens from stratospheric sulfur geoengineering, *Atmos. Chem. Phys.*, 12, 4775-4793, doi:10.5194/acp-12-4775-2012, 2012.
- 685 Friberg, J., Martinsson, B. G., Andersson, S. M., Brenninkmeijer, C. A. M., Hermann, M., Van Velthoven, P. F. J., and Zahn, A.: Sources of increase in lowermost stratospheric sulphurous and carbonaceous aerosol background concentrations during 1999–2008 derived from CARIBIC flights, *Tellus B*, 66:1, 23428, DOI:10.3402/tellusb.v66.23428, 2014.
- 690 Froyd, K. D., Murphy, D. M., Brock, C. A., Campuzano-Jost, P., Dibb, J. E., Jimenez, J-L., Kupc, A., Middlebrook, A. M., Schill, G. P., Thornhill, K. L., Williamson, C. J., Wilson, J. C., and Ziemba, L. D.: A new method to quantify mineral dust and other aerosol species from aircraft platforms using single-particle mass spectrometry, *Atmos. Meas. Tech.*, 12, 6209-8239, DOI:10.5194/amt-12-6209-2019, 2019.
- 695 Grainger, R. G., Lambert, A., Taylor, F. W., Remedios, J. J., Rodgers, C. D., and Corney, M.: Infrared absorption by volcanic stratospheric aerosols observed by ISAMS, *Geophys. Res. Lett.*, 20, 1283-1286, 1993.
- Hansen, J., et al.: Efficacy of climate forcings, *J. Geophys. Res.*, 110, D18104, doi:10.1029/2005JD005776, 2005.
- 700 Hanson, D. R., Ravishankara, A. R., and Solomon, S.: Heterogeneous reactions in sulfuric acid aerosols: A framework for model calculations, *J. Geophys. Res.*, 99, 3615-3629, <https://doi.org/10.1029/93JD02932>, 1994.
- Heckendorn, P., Weisenstein, D., Fueglistaler, S., Luo, B. P., Rozanov, E., Schraner, M., Thomason, L. W., and Peter, T.: The impact of geoengineering aerosols on stratospheric temperature and ozone, *Environ. Res. Lett.*, 4, 045108, <https://doi.org/10.1088/1748-9326/4/4/045108>, 2009.
- 705 Hofmann, D., Barnes, J., O'Neill, M., Trudeau, M., and Neely, R.: Increase in background stratospheric aerosol observed with lidar at Mauna Loa Observatory and Boulder, Colorado, *Geophys. Res. Lett.*, 36, L15808, doi:10.1029/2009GL039008, 2009.
- 710 Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, *Rev. Geophys.*, 33, 403-439, <https://doi.org/10.1029/95RG02097>, 1995.
- Huang, T. Y. W., and Massie, S. T.: Effect of volcanic particles on the O₂ and O₃ photolysis rates and their impact on ozone in the tropical stratosphere, *J. Geophys. Res.*, 102, 1239-1249, <https://doi.org/10.1029/96JD02967>, 1997.

- 715 Huang, J., Minnis, P., Chen, B., Huang, Z., Liu, Z., Zhao, Q., Yi, Y., and Ayers, J. K.: Long-range transport and vertical structure of Asian dust from CALIPSO and surface measurements during PACDEX, *J. Geophys. Res.*, 113, D23212, doi:10.1029/2008JD010620, 2008.
- 720 Innocenzi, M. E., Swimm, R. T., Bass, M., French, R. H., Villaverde, A. B., and Kokta, M. R.: Room-temperature optical absorption in undoped α -Al₂O₃, *J. Appl. Phys.*, 67, 7542-7546, <https://doi.org/10.1063/1.345817>, 1990.
- Iraci, L. T., and Tolbert, M. A.: Heterogeneous interaction of formaldehyde with cold sulfuric acid: Implications for the upper troposphere and lower stratosphere, *J. Geophys. Res.*, 102, 16,099-16,107, <https://doi.org/10.1029/97JD01259>, 1997.
- 725 Jones, A. C., Haywood, J. M., and Jones, A.: Climatic impacts of stratospheric geoengineering with sulfate, black carbon and titania injection, *Atmos. Chem. Phys.*, 16, 2843-2862, doi:10.5194/acp-16-2843-2016, 2016.
- Junge C. E., and Manson J. E.: Stratospheric aerosol studies, *J. Geophys. Res.*, 66, 2163-2182, <https://doi.org/10.1029/JZ066i007p02163>, 1961.
- 730 Khaykin, S. M., Godin-Beckmann, S., Keckhut, P., Hauchecorne, A., Jumelet, J., Vernier, J.-P., Bourassa, A., Degenstein, D. A., Rieger, L. A., Bingen, C., Vanhellemont, F., Robert, C., DeLand, M., and Bhartia, P. K.: Variability and evolution of the midlatitude stratospheric aerosol budget from 22 years of ground-based lidar and satellite observations, *Atmos. Chem. Phys.*, 17, 1829-1845, doi:10.5194/acp-17-1829-2017, 2017.
- Kremser, S., et al.: Stratospheric aerosol – Observations, processes, and impact on climate, *Rev. Geophys.*, 54, 278-335, doi:10.1002/2015RG000511, 2016.
- 740 Krieger, U. K., Marcolli, C., and Reid, J. P.: Exploring the complexity of aerosol particle properties and processes using single particle techniques, *Chem. Soc. Rev.*, 41, 6631-6662, <https://doi.org/10.1039/C2CS35082C>, 2012.
- Kupc, A., Williamson, C., Wagner, N. L., Richardson, M. and Brock, C. A.: Modification, calibration, and performance of the Ultra-High Sensitivity Aerosol Spectrometer for particle size distribution and volatility measurements during the Atmospheric Tomography Mission (ATom) airborne campaign, *Atmos. Meas. Tech.*, 11, 369-383, <https://doi.org/10.5194/amt-11-369-2018>, 2018.
- 745

- Labitzke, K., and McCormick, M. P.: Stratospheric temperature increases due to Pinatubo aerosols, *Geophys. Res. Lett.*, 19, 207-210, <https://doi.org/10.1029/91GL02940>, 1992.
- 750
- Lacis, A., Hansen, J., and Sato, M.: Climate forcing by stratospheric aerosols, *Geophys. Res. Lett.*, 19, 1607-1610, <https://doi.org/10.1029/92GL01620>, 1992.
- Lacis, A.: Volcanic aerosol radiative properties, *Pages magazine*, 23, 51-51, 2015.
- 755
- Martinsson, B. G., Friberg, J., Sandvik, O. S., Hermann, M., van Velthoven, P. F. J., and A. Zahn, A., Formation and composition of the UTLS aerosol, *npj Climate Atmos. Sci.*, 40, <https://doi.org/10.1038/s41612-019-0097-1>, 2019.
- Michelangeli, D. V., Allen, M., and Yung, Y. L.: El Chichon volcanic aerosols: Impact of radiative, thermal, and chemical perturbations, *J. Geophys. Res.*, 94, 18429-08443, <https://doi.org/10.1029/JD094iD15p18429>, 1989.
- 760
- Mills, M. J., Richter, J. H., Tilmes, S., Kravitz, B., MacMartin, D. G., Glanville, A. A., Tibbia, J. J., Lamarque, J-F., Vitt, F., Schmidt, A., Gettelman, A., Hannay, C., Bacmeister, D., and Kinnison, D. E.: Radiative and chemical response to interactive stratospheric sulfate aerosols in fully coupled CESM1(WACCM), *Journal of Geophysical Research: Atmospheres*, 122, 13,061-13,078, <https://doi.org/10.1002/2017JD027006>, 2017.
- 765
- Murphy D. M., Thomson D. S., and Mahoney M. J.: In situ measurements of organics, meteoric material, mercury, and other elements in aerosols at 5 to 19 km, *Science*, 282: 1664-1669, DOI: 10.1126/science.282.5394.1664, 1998.
- 770
- Murphy, D. M., and Thomson, D. S.: Halogen ions and NO⁺ in the mass spectra of aerosols in the upper troposphere and lower stratosphere, *Geophys. Res. Lett.*, 27, 3217-3220, <https://doi.org/10.1029/1999GL011267>, 2000.
- Murphy, D. M., Cziczo, D. J., Hudson, P. K., and Thomson, D. S.: Carbonaceous material in aerosol particles in the lower stratosphere and tropopause region, *J. Geophys. Res.*, 112, D04203, doi:10.1029/2006JD007297, 2007.
- 775
- Murphy, D. M.: Effect of Stratospheric aerosols on direct sunlight and implications for concentrating solar power, *Environ. Sci. Technol.*, 43, 2784-2786, <https://doi.org/10.1021/es802206b>, 2009.
- Murphy, D. M., Froyd, K. D., Schwarz, J. P., and Wilson, J. C.: Observations of the chemical composition of stratospheric aerosol particles, *Q. J. Roy. Meteor. Soc.*, 140, 1269-1278, <https://doi.org/10.1002/qj.2213>, 2014.
- 780

- National Research Council: Climate Intervention: Reflecting Sunlight to Cool Earth. Washington, DC: The National Academies Press. <https://doi.org/10.17226/18988>, 2015.
- 785 Nguyen, H. N., and Martinsson, B. G.: Analysis of C, N and O in aerosol collected on an organic backing using internal blank measurements and variable beam size, *Nucl. Instrum. Meth. B.*, 264, 96-102, <https://doi.org/10.1016/j.nimb.2007.08.001>, 2007.
- Niemeier, U., and Timmreck, C.: What is the limit of climate engineering by stratospheric injection of SO₂?, *Atmos. Chem. Phys.*, 15, 9129-9141, doi:10.5194/acp-15-9129-2015, 2015.
- 790 Noziere, B., and Esteve, W.: Organic reactions increasing the absorption index of atmospheric sulfuric acid aerosols, *Geophys. Res. Lett.*, 32, L03812, doi:10.1029/2004GL021942, 2005.
- 795 Pechtl, S., Schmitz, G., and Von Glasow, R.: Modelling iodide – iodate speciation in atmospheric aerosol: Contributions of inorganic and organic iodine chemistry, *Atmos. Chem. Phys.*, 7, 1381-1393, 2007.
- Pinto, J. P., Turco, R. P., and Toon, O. B.: Self-limiting physical and chemical effects in volcanic eruption clouds, *J. Geophys. Res.*, 94, 11165-11174, <https://doi.org/10.1029/JD094iD08p11165>, 1989.
- 800 Pitari, G., and Rizi, V.: An estimate of the chemical and radiative perturbation of stratospheric ozone following the eruption of Mt. Pinatubo, *J. Atmos. Sci.*, 50, 3260-3276, [https://doi.org/10.1175/1520-0469\(1993\)050%3C3260:AEOTCA%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050%3C3260:AEOTCA%3E2.0.CO;2), 1993.
- 805 Pitari, G., Aquila, V., Kravitz, B., Robock, A., Watanabe, S., Cionni, I., De Luca, N., Di Genova, G., Mancini, E., and Tilmes, S.: Stratospheric ozone response to sulfate geoengineering: Results from the Geoengineering Model Intercomparison Project (GeoMIP), *J. Geophys. Res. Atmos.*, 119, 2629-2653, doi:10.1002/2013JD020566, 2014.
- Price, H. C., Murray, B. J., Mattsson, J., O'Sullivan, D., Wilson, T. W., Baustian, K. J., and Benning, L. G.: Quantifying water diffusion in high-viscosity and glassy aqueous solutions using a Raman isotope tracer method, *Atmos. Chem. Phys.*, 14, 3817-3830, doi:10.5194/acp-14-3817-2014, 2014.
- 815 Rollins, A. W., et al.: The role of sulfur dioxide in stratospheric aerosol formation evaluated by using in situ measurements in the tropical lower stratosphere, *Geophys. Res. Lett.*, 44, 4280-4286, doi:10.1002/2017GL072754, 2017.

- Rosenlof, K. H., Tuck, A. F., Kelly, K. K., Russell III, J. M., and McCormick, M. P.: Hemispheric asymmetries in water vapor and inferences about transport in the lower stratosphere, *J. Geophys. Res.*, 102, 13,213-13,234, <https://doi.org/10.1029/97JD00873>, 1997.
- 820 Ross, M. N., and Sheaffer, P. M.: Radiative forcing caused by rocket engine emissions, *Earth's Future*, 2, 177-196, doi:10.1002/2013EF000160, 2014.
- Schneider, J., Weigel, R., Klimach, T., Dragoneas, A., Appel, O., Hünig, A., Molleker, S., Köllner, F., Clemen, H.-C., Eppers, O., Hoppe, P., Hoor, P., Mahnke, C., Krämer, M., Rolf, C., Groß, J.-U., Zahn, A., Obersteiner, F., Ravegnani, F.,
825 Ulanovsky, A., Schlager, H., Scheibe, M., Diskin, G. S., DiGangi, J. P., Nowak, J. B., Zöger, M., and Borrmann, S.: Aircraft-based observation of meteoric material in lower stratospheric aerosol particles between 15 and 68° N, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2020-660>, in review, 2020.
- Solomon, S., Portmann R. W., Garcia R. R., Thomason L. W., Poole L. R., and McCormick M. P.: The role of aerosol
830 variations in anthropogenic ozone depletion at northern midlatitudes, *J. Geophys. Res.* 101: 6713-6727, <https://doi.org/10.1029/95JD03353>, 1996.
- Solomon, S., Daniel, J. S., Neely III, R-R., Vernier, J-P., Dutton, E. G., Thomason, L. W.: The persistently variable “background” stratospheric aerosol layer and global climate change, *Science*, 333, 866-870, DOI: 10.1126/science.1206027,
835 2011.
- Solomon, S., Kinnison, D., Bandoro, J., and Garcia, R.: Simulation of polar ozone depletion: An update, *J. Geophys. Res. Atmos.*, 120, 7958-7974, doi:10.1002/2015JD023365, 2015.
- 840 Stone, K. A., Solomon, S., Kinnison, D. E., Pitts, M. C., Poole, L. R., Mills, M. J., and Hagiya, S.: Observing the impact of Calbuco volcanic aerosols on South Polar ozone depletion in 2015, *J. Geophys. Res.*, 122, 11,862-11,879, <https://doi.org/10.1002/2017JD026987>, 2017.
- Thomson, D. S., Schein, M. E., and Murphy, D. M.: Particle analysis by laser mass spectrometry WB-57F instrument
845 overview, *Aerosol Sci. Tech.*, 33, 153-169, <https://doi.org/10.1080/027868200410903>, 2000.
- Toon, O. B., Turco, R. P., Westphal, D., Malone, R., and Liu, M.: A multidimensional model for aerosols—Description of computational analogs, *J. Atmos. Sci.*, 45, 2123-2144, [https://doi.org/10.1175/1520-0469\(1988\)](https://doi.org/10.1175/1520-0469(1988)), 1988.

850 Torres, O., Bhartia, P. K., Taha, G., Jethva, H., Das, S., Colarco, P., Krotkov, N., Omar, A., and Ahn, A.: Stratospheric injection of massive smoke plume from Canadian boreal fires in 2017 as seen by DSCOVR-EPIC, CALIOP, and OMPS-LP observations. *J. Geophys. Res.*, 125, e2020JD032579, [https://doi.org/ 10.1029/2020JD032579](https://doi.org/10.1029/2020JD032579), 2020.

Van de Hulst, H. C.: *Light scattering by small particles*, Dover, ISBN 9780486642284, 1981.

855 Vernier, J.-P., et al.: Major influence of tropical volcanic eruptions on the stratospheric aerosol layer during the last decade, *Geophys. Res. Lett.*, 38, L12807, doi:10.1029/2011GL047563, 2011.

860 Visioni, D., Pitari, G., and Aquila, V.: Sulfate geoengineering: a review of the factors controlling the needed injection of sulfur dioxide, *Atmos. Chem. Phys.*, 17, 3879-3889, doi:10.5194/acp-17-3879-2017, 2017.

Weisenstein, D. K., Keith, D. W., and Dykema, J. A.: Solar geoengineering using solid aerosol in the stratosphere, *Atmos. Chem. Phys.*, 15, 11835-11859, doi:10.5194/acp-15-11835-2015, 2015.

865 Williams, M. B., Michelsen, R. H., Axson, J. L., and Iraci, L. T.: Uptake of acetone, acetaldehyde and ethanol in cold sulfuric acid solutions containing organic material: Carbon accretion mechanisms, *Atmos. Environ.*, 44, 1145-1151, <https://doi.org/10.1016/j.atmosenv.2009.12.036>, 2010.

870 Wilson, J. C., Loewenstein, M., Fahey, D. W., Gary, B., Smith, S. D., Kelly, K. K., Ferry, G. V., and Chan, K. R.: Observations of condensation nuclei in the airborne Antarctic ozone experiment: Implications for new particle formation and polar stratospheric cloud formation, *J. Geophys. Res.*, 94, 16 437-16 448, <https://doi.org/10.1029/JD094iD14p16437>, 1989.

875 Wilson J. C., Lee S-H., Reeves J. M., Brock C. A., Jonsson H. H., Lafleur B. G., Loewenstein M., Podolske J., Atlas E., Boering K., Toon G., Fahey D. W., Bui T. P., Diskin G., and Moore F.: Steady-state aerosol distributions in the extra-tropical, lower stratosphere and the processes that maintain them, *Atmos. Chem. Phys.*, 8: 6617-6626, <https://doi.org/10.5194/acp-8-6617-2008>, 2008.

880 Yu, P., Toon, O. B., Bardeen, C. G., Mills, M. J., Fan, T., English, J. M., and Neely, R. R.: Evaluations of tropospheric aerosol properties simulated by the Community Earth System Model with a sectional aerosol microphysics scheme, *J Adv. Model Earth Sys.*, 7, 865-914, <https://doi.org/10.1002/2014MS000421>, 2015.

Yu, P., Murphy, D. M., Portmann, R. W., Toon, O. B., Froyd, K. D., Rollins, A. W., Gao, R-S., and Rosenlof, K. H.: Radiative forcing from anthropogenic sulfur and organic emissions reaching the stratosphere, *Geophys. Res. Lett.*, 43, 9361-9367, doi:10.1002/2016GL070153, 2016.




Figure 1. Composition resolved size distributions from the lower stratosphere during the Atmospheric Tomography (ATom) mission. The size distributions from optical particle counter measurements are apportioned at each size into classes of particles based on the PALMS single particle composition data. The smallest few size bins sometimes had too few particles with PALMS mass spectra to apportion the composition. Sizes in all figures are geometric diameter at low relative humidity and concentrations are per standard cm^3 . Fine structure in total volume distributions at $0.6 \mu\text{m}$ and larger may be artifacts due to Mie resonances in the optical particle counter; minima near 0.2 or $0.3 \mu\text{m}$ are robust.




Figure 2. A comparison of the CARMA bin microphysics aerosol model in the CESM climate model with observed size distributions during ATom2. Total particle volumes are proportional to areas under each curve.






Figure 3. Composition-resolved size distributions for the times when data are available for ozone greater than 500 ppbv, all in the Northern Hemisphere. The dashed curves are to facilitate comparison with Figure 1.



905

Figure 4. Size distributions of several types of particles in the lower stratosphere. Northern Hemisphere data are solid and Southern Hemisphere data are dashed. Data are from 250 to 400 ppbv ozone except the curves with circles are data at more than 500 ppbv ozone. The three major events are noted where they may have influenced the size distributions. Lognormal fits are given in Table 1.


910




915

Figure 5. Profiles of the volume concentration of two types of particles in the lower stratosphere, using ozone as a surrogate for a vertical coordinate. This does not include particles smaller than 0.1 μm . Sulfuric acid particles with meteoric metals have a high-altitude source whereas organic-sulfate particles, as selected by their mass spectra, originate in the troposphere. The organic-sulfate particles are more abundant in the Northern Hemisphere than they are in the Southern Hemisphere.

920




925 **Figure 6.** The organic content of two types of stratospheric particles using ozone as a surrogate for altitude. Shown are the relative areas of the C^+ peak in the mass spectra relative to two peaks indicative of sulfate. The top axis is an approximate mass fraction. The organic-sulfate particles are about 40 to 70% organic by mass, consistent with a tropospheric source. The sulfuric acid particles have near-zero organic content well into the stratosphere, increasing up to a few percent by mass near the tropopause. There was a strong difference in the organic content of the sulfuric acid particles between the hemispheres. Each point is the average of about 200 mass spectra.



930

Figure 7. The mass scattering efficiency of particles with a refractive index similar to sulfuric acid. Shown are the total scattered light, the light scattered away from the Earth to space, and their ratio. Larger particles have more forward scattering and therefore a smaller fraction scattered to space. The lowest curve shows the climate impact after an approximate correction for warming due to infrared absorption. Particles smaller than about 0.1 μm or larger than about 4 μm warm the Earth (Lacis et al., 1992). Tropospheric particles containing ammonium sulfate or organic material with higher refractive indices would shift the curves to slightly larger diameters.

935



940

Figure 8. The composition-resolved size distribution from Figure 1 for the ATom2 Northern Hemisphere weighted by volume, light scattered to outer space, and surface area. The two sulfuric acid categories (with and without meteoric metals) have been combined. Percentages refer to sizes above 0.1 μm only. ATom2 is chosen as an example; Figure S4 shows the percentage contributions of various particle types to these processes for other deployments and for the Southern Hemisphere. Scaling to net thermal infrared heating gives nearly identical relative contributions as volume in the top panel. Scaling to sedimentation rate gives a similar shape to the middle panel.

945




Figure 9. Calculated net cooling per unit volume, surface area, and sedimentation velocity. Calculations of sunlight reflected to outer space are for a single size aerosol averaged over the solar spectrum and zenith angles characteristic of the sunlit Earth. The wavelength averaging eliminates the oscillations from Mie scattering. Particles of about 0.3 μm diameter give the most cooling per unit sedimentation rate and particles of about 0.5 μm diameter give the most cooling per unit mass. Each curve is normalized so the maximum point is unity. For example, 0.2 μm particles have about one tenth the net cooling per unit surface area as 1 μm diameter particles.

950

955

	Total Volume		Meteoric-sulfuric			Sulfuric			Organic-sulfate			Other
	6 nm-1.7 μm $\mu\text{m}^3 \text{cm}^{-3}$	0.1-1.7 μm $\mu\text{m}^3 \text{cm}^{-3}$	Volume $\mu\text{m}^3 \text{cm}^{-3}$	Fit diam. μm	Fit σ	Volume $\mu\text{m}^3 \text{cm}^{-3}$	Fit diam. μm	Fit σ	Volume $\mu\text{m}^3 \text{cm}^{-3}$	Fit diam. μm	Fit σ	Volume $\mu\text{m}^3 \text{cm}^{-3}$
ATom1 NH	0.35	0.30	0.046	0.37	1.35	0.103	0.29	1.55	0.150	0.13	1.89	0.004
ATom2 NH	0.35	0.33	0.083	0.46	1.44	0.115	0.35	2.14	0.124	0.17	1.45	0.005
ATom3 NH	0.78	0.76	0.150	0.54	1.51	0.160	0.50	1.74	0.436	0.39	2.89	0.024
Atom4 NH	0.68	0.64	0.066	0.44	1.49	0.175	0.22	2.15	0.292	0.30	2.04	0.107
ATom1 SH	0.49	0.48	0.101	0.39	1.27	0.330	0.40	1.29	0.050	0.27	1.43	0.011
ATom2 SH	0.32	0.31	0.110	0.40	1.39	0.147	0.39	1.64	0.052	0.12	1.80	0.001
ATom3 SH	0.28	0.26	0.056	0.40	1.29	0.088	0.35	1.73	0.120	0.19	1.93	0.003
ATom4 SH	0.19	0.17	0.040	0.36	1.47	0.090	0.16	2.30	0.033	0.20	2.94	0.004
ATom2 NH high	0.40	0.39	0.156	0.42	1.36	0.122	0.40	1.64	0.112	0.18	1.36	0.005
ATom4 NH high	0.37	0.35	0.132	0.41	1.59	0.104	0.31	2.57	0.093	0.38	3.64	0.021

960

Table 1. Total volumes and volumes and log-normal fit parameters for individual types of particles. Lognormal fits were performed for particle volume so the fit diameter is for the volume distribution. The volumes and fits are for the size range 0.1 to 1.7 μm diameter except as noted (first column). Data are for 250-400 ppbv ozone except the last two rows are for 500-850 ppbv. Volumes are per standard cm^3 . Fits are to the equation $V = a^x \exp(-(\ln D - \ln D_{fit})^2 / (2\sigma^2))$ where D is the diameter and a is a scaling factor. Each fit is for a specific component. Fits were not attempted for “other” particles (mostly dust) because they did not generally show a defined mode. Italics indicate a poor fit to a single lognormal shape.

965