1	Changes in biomass burning, wetland extent, or agriculture
2	drive atmospheric NH3 trends in <u>key</u> African regions
3	
4	
5	
6 7 8 9	Jonathan E. Hickman ^{1*,} Niels Andela ^{2†} , Enrico Dammers ³ , Lieven Clarisse ⁴ , Pierre-François Coheur ⁴ , Martin Van Damme ⁴ , Courtney Di Vittorio ⁵ , Money Ossohou ⁶ , Corinne Galy- Lacaux ⁷ , Kostas Tsigirdis ^{1,8} , Susanne Bauer ¹
10	¹ NASA Goddard Institute for Space Studies, New York, USA
11	² NASA Goddard Space Flight Center, Beltsville, USA
12	³ Air Quality Research Division, Environment and Climate Change Canada, Toronto,
13	Canada
14	⁴ Université libre de Bruxelles (ULB), Service de Chimie Quantique et Photophysique,
15	Atmospheric Spectroscopy, Brussels, Belgium
16	⁵ Wake Forest University, Winston-Salem, USA
17	⁶ Laboratoire des Sciences de la Matière, de l'Environnement et de l'Energie Solaire,
18	Université Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire
19	⁷ Laboratoire d'Aérologie, Université Toulouse III Paul Sabatier / CNRS, France
20	⁸ Columbia University, New York, USA
21 22	$^{\dagger}\text{Now}$ at School of Earth and Ocean Sciences, Cardiff University, Cardiff, UK
23	*Correspondence to: jonathan.e.hickman@nasa.gov
24	
25	Abstract
26	Atmospheric ammonia (NH_3) is a precursor to fine particulate matter and a source
27	of nitrogen (N) deposition that can adversely affect ecosystem health. The main sources of
28	NH_3 —agriculture and biomass burning—are undergoing or expected to undergo
29	substantial changes in Africa. Although evidence of increasing NH_3 over parts of Africa has
30	been observed, the mechanisms behind these trends are not well understood. Here we use

Deleted: several

32 observations of atmospheric NH₃ vertical column densities (VCDs) from the Infrared 33 Atmospheric Sounding Interferometer (IASI) along with other satellite observations of the 34 land surface and atmosphere to evaluate how NH₃ concentrations have changed over Africa 35 from 2008 through 2018, and what has caused those changes. In West Africa NH₃ VCDs are 36 observed to increase during the late dry season, with increases of over 6% yr⁻¹ in Nigeria 37 during February and March (p<0.01). These positive trends are associated with increasing 38 burned area and CO trends during these months, likely related to agricultural preparation. 39 Increases are also observed in the Lake Victoria Basin, where they are associated with 40 expanding agricultural area. In contrast, NH₃ VCDs declined over the Sudd wetlands in 41 South Sudan by over <u>1.5</u>% yr⁻¹, though not significantly (p=0.28). Annual maxima in NH₃ 42 VCDs in South Sudan occur during February through May and are associated with drying of 43 temporarily flooded wetland soils, which favor emissions of NH₃. The change in mean NH₃ 44 VCDs over the Sudd is strongly correlated with variation in wetland extent in the Sudd: in 45 years when more area remained flooded during the dry season, NH₃ concentrations were 46 higher (r=0.64, p≤0.05). Relationships between biomass burning and NH₃ may be observed 47 when evaluating national-scale statistics: countries with the highest rates of increasing NH₃ VCDs also had high rates of growth in CO VCDs; burned area displayed a similar pattern, 48 49 though not significantly. Livestock numbers were also higher in countries with 50 intermediate or high rates of NH₃ VCD growth. Fertilizer use in Africa is currently low but 51 growing; implementing practices that can limit NH₃ losses from fertilizer as agriculture is 52 intensified may help mitigate impacts on health and ecosystems. 53 1. Introduction: 54 55 56 Ammonia (NH₃), a reactive nitrogen (N) trace gas, plays a number of important roles 57 in the atmosphere, with implications for human health, climate, and ecosystems. Once in 58 the atmosphere, NH₃ contributes to the production of inorganic aerosols, the primary constituents of fine particulate matter and a serious health hazard (Bauer et al., 2016; 59 60 Lelieveld et al., 2015; Pope et al., 2002). NH₃ can also be deposited to downwind 61 ecosystems, contributing to eutrophication, soil acidification, vegetation damage, 62 productivity declines, reductions in biodiversity, and indirect greenhouse gas emissions

Deleted: 7

Deleted: 2 Deleted: Deleted: 0

Deleted: 5	
Deleted: =)
Deleted: 4)

70	(Denier Van Der Gon and Bleeker, 2005; Krupa, 2003; Matson et al., 1999; Stevens et al.,
71	2018; Tian and Niu, 2015).
72	Although NH $_3$ is emitted from natural soils, agriculture is by far the largest source of
73	NH_3 globally (Behera et al., 2013; Bouwman et al., 1997). Urea fertilizer and livestock
74	excreta are particularly important substrates for NH_3 formation, and can be volatilized
75	quickly under favorable environmental conditions (Bouwman et al., 1997). In all soils, NH_3
76	is formed in solution following the dissociation of ammonium (NH $_4$ +; Eq. 1).
77	
78	$NH_{4^{+}} + OH^{-} \leftrightarrow H_2O + NH_3$ (Eq. 1)
79	
80	Soil NH $_3$ production is temperature-dependent, doubling with every 5°C temperature
81	increase, though the actual soil $\rm NH_3$ flux is determined in part by plant and soil physiological
82	and physical factors (Sutton et al., 2013). On average, fertilizer use has been extremely low
83	in sub-Saharan Africa—often an order of magnitude or more lower than typical in Europe,
84	the United States, or China (Hazell and Wood, 2008; Vitousek et al., 2009). Livestock manure
85	N content also tends to be very low in sub-Saharan Africa (Rufino et al., 2006), The low
86	fertilizer use suggests that natural soils (as opposed to agricultural soils) may be a more
87	important source in the region than elsewhere in the world. However, agricultural
88	intensification and increasing fertilizer use has been a central policy focus for many African
89	countries, with national and regional efforts to increase N inputs by an order of magnitude
90	or more (AGRA, 2009).
91	After agriculture, biomass burning is the most important source of NH_3 globally
92	(Bouwman et al., 1997), with roughly 60 to 70% of global $\rm NH_3$ emissions from fires occurring
93	in Africa (Cahoon et al., 1992; Whitburn et al., 2015). The amount of $\rm NH_3$ emitted from
94	biomass fires is controlled primarily by the type of burning that occurs. N in fuel is present
95	predominantly in a chemically reduced state, and \ensuremath{NH}_3 is emitted in greater quantities from
96	low temperature smoldering combustion in which fuel N is incompletely oxidized (Goode et
97	al., 1999; Yokelson et al., 2008). Fuel moisture content, which can help determine whether

- 98 combustion is smoldering or flaming, is thus an important determinant of biomass burning
- 99 NH_3 emissions (Chen et al., 2010).

Deleted: ; Deleted: t

Deleted: ,

103	In contrast to other reactive N gases such as NO_x (nitric oxide + nitrogen dioxide),	
104	NH_3 emissions are typically unregulated outside of Europe (Anker et al., 2018; Kanter,	
105	2018; USDA Agricultural Air Quality Task Force, 2014), and substantial increasing trends	
106	have been observed by the NASA Atmospheric InfraRed Sounder (AIRS) and the Infrared	
107	Atmospheric Sounding Interferometer (IASI) over many of the world's major agricultural	
108	and biomass burning regions during the $21^{ m st}$ century (Van Damme et al., 2021; Warner et	
109	al., 2017). West Africa has been identified as an important NH_3 source region (Van Damme	
110	et al., 2018), where a trend of increasing NH_3 concentrations <u>in recent decades</u> has been	Deleted: over 2002 to 2013
111	attributed at least in part to increased fertilizer use (Van Damme et al., 2021; Warner et al.,	
112	2017). Increasing trends have also been observed over central Africa, and attributed to	
113	higher rates of biomass burning (Van Damme et al., 2021; Warner et al., 2017). However,	
114	the studies by Warner et al. (2017), and Van Damme et al. (2021) were global in nature, and	Deleted:
115	as such could not include detailed explorations of the drivers of trends such as	
116	consideration of emission seasonality or the geographic distribution of emission drivers.	
117	Consideration of these factors is particularly important across large parts of Africa where	Deleted: , which are
118	both biomass burning and soils are potentially important sources <u>of NH_3</u> (van der A et al.,	
119	2008).	
120	Here we use a <u>n eleven</u> -year satellite record to evaluate trends in atmospheric NH_3	Deleted: ten
121	concentrations over Africa from 2008 through 201 <mark>8, including detailed examination of</mark>	Deleted: 7
122	three regions where changes are pronounced: West Africa, the Lake Victoria Region, and	
123	South Sudan.	
124		
125	2. Data and Methods	
126	2.1 Global gridded data	
127	Multiple data products were used, including satellite observations and spatial	
128	datasets:	
129	-IASI-A, launched aboard the European Space Agency's MetOp-A in 2006, provides	
130	measurements of atmospheric NH_3 and carbon monoxide (CO) twice a day (9:30 in the	
131	morning and evening, Local Solar Time at the equator). Here we use morning observations,	
132	when the thermal contrast is more favorable for retrievals (Clarisse et al., 2009; Van	
133	Damme et al., 2014a). The NH $_3$ retrieval product used (ANNI-NH $_3$ -v3R) follows a neural	

139 network retrieval approach. We refer to Van Damme et al. (2017) and Van Damme et al. 140 (2021) for a detailed description of the algorithm. For CO, we used the product obtained 141 with the FORLI v20140922 retrieval algorithm (Hurtmans et al., 2012). Given the absence 142 of hourly or even daily observations of NH3 concentrations in sub-Saharan Africa, the 143 detection limit of IASI is difficult to determine with certainty. However, the region 144 experiences high thermal contrast, and IASI seems to be able to reliably observe NH₃ down 145 to 1 to 2 ppb at the surface (Clarisse et al., 2009; Van Damme et al., 2014b). We gridded the 146 Level-2 IASI NH₃ and CO products to $0.5^{\circ} \times 0.5^{\circ}$ resolution. We used a conventional binning 147 approach based on the center of each satellite footprint. We did not apply an averaging 148 weight. Quality control procedures were followed as detailed in van Damme et al. 2017 and 149 Van Damme et al., 2021. Specifically, the screening of retrievals included filtering of 150 retrievals where cloud cover is over 10%, where the total column density is below zero and 151 the absolute value of the hyperspectral range index (HRI) is above 1.5, and where the ratio 152 of the total column density to HRI is larger than 1.5 x 10¹⁶ molecules cm⁻². 153 The IASI products have been validated using ground-based Fourier transform 154 infrared (FTIR) observations of NH₂ total columns, with robust correlations at sites with 155 high NH₃ concentrations, but lower at sites where atmospheric concentrations approach IASI's detection limits (Dammers et al., 2016; Guo et al., 2021). Compared to the FTIR 156 observations, total columns from previous IASI NH₃ products (IASI-LUT and IASI-NNv1) 157

- are biased low by \sim 30% which varies per region depending on the local concentrations.
- 159 Although FTIR observations are absent from Africa, earlier work has shown fair agreement
- 160 between previous versions of IASI total column densities and surface observations of NH₃
- 161 using passive samplers across the International Network to study Deposition and
- 162 Atmospheric chemistry in AFrica (INDAAF) network in West Africa (Van Damme et al.,
- 163 2015), including in observations of seasonal variation (Hickman et al., 2018; Ossohou et al.,
- 164 2019). Validation of the IASI CO product using surface, aircraft, and satellite observations
- 165 have found total columns to have an error that is generally below 10-15% in the tropics
- and mid-latitudes (George et al., 2009; Kerzenmacher et al., 2012; Pommier et al., 2010; De
- 167 Wachter et al., 2012). The IASI NH_3 and CO products were used for the years 2008—the
- 168 first full year of data available—to the end of 2018. Random errors in observations can be

Deleted: 2

Deleted: 2 Deleted: to match the resolution of other data used in the analysis. ... 173 assumed to cancel out in the annual mean, which is what we used in our analysis. With the

174 assumption that random errors cancel out, only systematic errors related to tropospheric

175 vertical column contents remain; these systematic errors do not contribute to uncertainty

176 in trend analyses. In addition, we first take monthly averages based on all daily

177 observations within a given month before calculating seasonal means to minimize any

178 potential effects of temporal variability in cloud cover.

179 -The Tropical Rainfall Measuring Mission (TRMM) daily precipitation product (3B42)
180 is based on a combination of TRMM observations, geo-synchronous infrared observations,
181 and rain gauge observations (Huffman et al., 2007). Independent rain gauge observations
182 from West Africa have been used to validate the product, with no indication of bias in the
183 product (Nicholson et al., 2003).

NOAA Global Surface Temperature Dataset, a 0.5° gridded 2m monthly land surface
 temperature product (Fan and van den Dool, 2008). The data set is based on a combination
 of station observations from the Global Historical Climatology Network version 2 and the
 Climate Anomaly Monitoring System (GHCN_CAMS), and uses an anomaly interpolation
 approach which relies on observation-based reanalysis data to derive spatio-temporal
 variation in temperature lapse rates for topographic temperature adjustment.

- 500m MCD64A1 collection 6 Moderate Resolution Imaging Spectroradiometer
(MODIS) burned area product for the period 2008-2018 (Giglio et al., 2018). The burned
area data are aggregated by month and gridded to 0.25° resolution, and do not include
burned area from small fires.

MODIS MCD12C1 (collection 5) land cover product, which provides the percentage
 of cropped area in each 0.25° grid cell (Friedl et al., 2002). In Africa, agriculture is often
 practiced in complex mosaics of agricultural and natural land cover, so we used both the crop
 and crop/natural area mosaic MODIS classifications as agricultural area in our analysis.

-We also used data on the spatio-temporal distribution of armed conflict events from
the Armed Conflict Location & Event Data Project (ACLED; Raleigh et al., 2010). We included
data for both violent and non-violent conflict events over the period 2008-2018,

202 2.2 Sudd wetland extent

203 Monthly flooded area extents of the Sudd Wetland, South Sudan from 2000 to 2017 204 were derived from 8-day composite MODIS land surface reflectance imagery (MOD09A1); data from 2005 through 2017 were used in the analyses. We refer to Di Vittorio and 205 206 Georgakakos (2018) for a detailed description of the classification procedure designed to 207 retrieve these data. In summary, monthly flood maps were obtained through a two-stage classification procedure. The first stage used the full 18-year data set to produce a wetland 208 209 land cover map that distinguishes between wetland vegetation classes and their flooding 210 regimes (permanently flooded, seasonally flooded, or non-flooded). The second stage 211 compares seasonally flooded pixels from each vegetation class to their non-flooded 212 counterparts on a monthly basis to identify the timing and duration of flooding for each pixel. 213 These data were originally derived to calibrate a hydrologic model of the Sudd that is 214 dependent on Nile flows; therefore, a connectivity algorithm was applied to ensure that all 215 flooded pixels were physically connected to the Nile River. A few adjustments have been 216 made to the previously published dataset for the application of this study. The classification 217 algorithm has been improved to more accurately capture the inter-annual fluctuations in the 218 permanently flooded areas. The dataset was also extended through 2017, and the total 219 flooded area was quantified prior to applying the connectivity algorithm. The magnitudes of 220 the monthly flooded area estimates are now substantially larger because they include areas 221 flooded from local runoff in addition to areas flooded by the Nile River.

222

223

2.3 Spatial and national analyses

We evaluated spatial relationships between mean annual tropospheric NH₃ concentration and several independent variables at 0.25° resolution: population density, livestock density, and cropped area. Population density and livestock density data are not available as time series suitable for trend analysis, so we use single year values in our analyses. We calculated population density based on the 2017 version of the US Department of Energy's Gridded Landscan population dataset (Dobson et al., 2000; 230 available at https://landscan.ornl.gov). Livestock density was based on the FAO global 231 gridded livestock dataset for the year 2007 (Robinson et al., 2014). Before analysis, we 232 converted the livestock densities of chickens, goats, pigs, and sheep to tropical livestock 233 units (TLU), using values of 0.01, 0.1, 0.2, and 0.1 TLU, respectively; North African cattle 234 were converted using a factor of 0.7, whereas sub-Saharan cattle were converted using a factor of 0.5 (Chilonda and Otte, 2006). For cropped area, we used the MODIS MCD12C1 235 236 (collection 5) land cover product as described above. We conducted spatial analyses by 237 establishing a map of 3° grid cells and calculating the correlation between the value of each 238 independent variable and NH₃ for all 0.5° grid cells within the larger grid cells (N = <u>36</u> 239 including water grid cells, though these were excluded from the analysis). 240

241 National data on annual livestock numbers, crop production, and fertilizer N use were 242 obtained from the UN Food and Agriculture Organization FAOSTAT for 51 African countries 243 (FAO, 2020). Livestock data consisting of sheep, goats, cattle, and pigs were converted to 244 tropical livestock units as described above, and buffaloes were converted using a conversion 245 factors of 0.7 (Chilonda and Otte, 2006). National emissions of CO2 were obtained from 246 World Bank Open Data (World Bank, 2019). National-level mean annual cropland area, 247 burned area, and atmospheric NH3 and CO concentrations were also calculated for each of 248 the 51 countries from the spatial datasets described above. Countries were sorted into three 249 bins based on whether their relative change in mean annual NH₃ concentration was low, 250 medium or high, and means and standard errors were calculated for each of the three 17-251 country bins.

Linear trend analyses were conducted using linregress from the scipy.stats package
in Python v3.6.3. Statistical analyses of national scale data were conducted using ANOVA in
R. Data were log or rank transformed when necessary to meet the assumptions of ANOVA.
Values of α for treatment comparisons following significant ANOVA results were corrected
for multiple testing using Benjamini-Hochberg corrections.

- 257
- 258 259

3. Results & Discussion

Deleted: 1.5

263 **3.1 Continental distributions and trends**

Mean annual NH₃ concentrations for 2008-2018 are highest across the savannas 264 265 and forest-savanna mosaics in North equatorial Africa, and especially in West Africa; there 266 are smaller regional hotspots in the Lake Victoria basin, South Sudanese wetlands, and 267 along the Nile delta and river (Fig. 1a). Parts of these regions experience substantial 268 biomass burning (Fig. 1e), high livestock densities (Fig. 1g), and/or high cropland cover 269 (Fig. 1h), all of which can contribute to NH₃ emissions. The high concentrations in West 270 Africa, which is one of the major global NH₃ hotspots (Van Damme et al., 2018), is likely the 271 result of biomass burning emissions. Biomass burning emissions tend to drive seasonal 272 variation in NH₃ VCDs in West Africa, with the largest emissions occurring late in the dry 273 season and early rainy season (Hickman et al., 2021b). In addition to local emissions, 274 biomass burning emissions and their reactive products are transported to the coast of West 275 Africa during both the northern hemisphere rainy season, when it is transported from 276 central and southern Africa, and during the dry season, when it is transported from 277 biomass burning regions to the east (Sauvage et al., 2007). Most areas with trends are 278 significant at P=0.2 or higher (Fig. S1). 279 In addition to being hotspots of mean NH₃ concentrations, some of these regions 280 have also experienced increases in NH₃ concentrations from 2008 to 2018 (Fig. 1b). Like 281 Warner et al. (2017) and Van Damme et al. (2021), we observed some increases in the 282 northern grasslands, central African forests, and the Nile region, but we also observe trends 283 in the Lake Victoria Basin, which Warner et al. (2017) did not, but Van Damme et al. (2021) 284 did. Also in contrast to Warner et al. (2017) but in line with Van Damme et al. (2021), we 285 observe a prominent decline in NH₃ VCDs over South Sudan (Fig. 1b, S1). 286 The Nile region exhibits elevated NH₃ concentrations and a modest positive trend 287 over the observation period (Fig. 1a, 1b). This trend appears largely to be related to 288 agriculture and livestock: in a spatial analysis, snapshots of livestock densities and of 289 population densities are both positively related to changes in NH₃ VCDs (Fig. 2). Although 290 there is not a positive relationship between agricultural area and NH₃ VCDs over the Nile 291 region from 2008 to 2018, Egypt's population increased by roughly 25% over that period 292 (World Bank, 2019), and fertilizer N use increased by roughly 8% after a decline in use



area, as well as spatial distribution of livestock density and cropped area across seven sub-

Saharan African ecoregions. Mean annual (a) and trend (b) in atmospheric NH₃ VCDs from

297

298



Deleted: several of these

- 303 IASI for the period 2008 through 2018. Mean annual (c) and trend (d) in annual
- atmospheric CO VCDs from IASI for the same period. Mean annual (e) and trend (f) in
- annual burned area from MODIS for 2008-2018. Livestock densities for 2007 from the FAO
- 306 (f), and mean cropped area from MODIS for 2008-2018 (g). The border of South Sudan is
- 307 highlighted in black, and several regions boxed: the Nile region at 30°N, the Sudd wetland
- 308 in South Sudan, the Lake Victoria region at the equator, and West Africa centered around
- 309 10°N.



311 Figure 2. Relationships between NH₃ trends and livestock density, population density, and

312 cropland area, as well as changes in cropland area. Spatial correlations between changes in

- 313 annual atmospheric NH₃ VCDs and livestock density (a) and population density (b).
- 314 Correlation between cropland area and NH₃ VCDs for 2008 through 2018 (c). Change in

315 crop area for 2008 through 2018 (d). The NH₃ and crop area trends are based on data for

316 2008 through 2018, livestock density data are for the year 2007, population density data

are for the year 2017.





320

324

Figure 3. Change in mean monthly atmospheric NH₃ VCDs for the period 2008 through

322 2018. Grid cells where mean annual $\rm NH_3$ VCDs for the entire period are under $5 x 10^{15}$

molecules cm⁻² are not displayed. Results significant at P=0.05 are presented in Figure S 5_{2}



330	Figure 4. Correlation coefficient for the relationship between mean annual CO and
331	$\rm NH_3$ VCDs (a), changes in $\rm NH_3$ VCDs (b) and changes in CO VCDs (c) over 2008 through
332	2018 in West Africa. Grid cells where mean annual NH_3 VCDs for the entire period are
333	under $5x10^{15}$ molecules cm ⁻² are not displayed. Results significant at P=0.05 for the entire
334	continent are presented in Figure S <mark>6,</mark>

3.2 West Africa

337	The increasing trend in NH_3 VCDs over West Africa are centered over Nigeria and
338	the southern coast, and to a lesser extent across parts of the wet savanna (Fig. 1b).
339	Increases in NH_3 VCDs tend to be higher in grid cells with higher population densities in
340	Nigeria <u>and other parts of West Africa</u> (Fig. 2b), suggesting a possible anthropogenic
341	influence. The spatial distribution of the mean annual NH_3 trend is overlapped by a
342	substantial increase in mean annual CO VCDs (Fig. 1b, 1d), pointing to a biomass burning
343	source, as is also the case in central Africa. Earlier studies have found substantial declines
344	in annual burned area across the north equatorial African biomass burning region as
345	detected by MODIS (Andela et al., 2017; Andela and van der Werf, 2014) and related
346	declines in NO_2 VCDs across the region (Hickman et al., 2021a), which would seem to stand
347	in contrast to the increasing CO and NH_3 trends observed here.
348	However, the annual decline in burned area and NO_2 VCDs is characterized by
349	heterogeneity when considering individual months. In West Africa, the dry season is
350	typically November to February or March. During the transition from the dry to rainy
351	season in February and March, $NO_2 \ VCDs$ exhibit increasing rather than decreasing trends
352	in West Africa, though burned area patterns are not as clear when 2018 is included
353	(Hickman et al., 2021a; Fig. S2, S3). Although these increases in NO_2 VCDs are small in the
354	annual context, they occur at a time of year when biomass burning combustion is less
355	complete, potentially due to greater fuel moisture and declining fire radiative power
356	(Hickman et al., 2021a; Zheng et al., 2018). These conditions would lead to greater

359 species such as CO₂ and NO₂ that dominate emissions during the peak of the biomass 360 burning season (Fig. S2, S4). Indeed, our observations suggest that much of the increasing 361 NH₃ trend occurs during this transitional period, with NH₃ VCDs increasing by roughly <u>6</u>% 362 yr⁻¹ for all of Nigeria during February and March (Fig. 3, <u>S5</u>; p<0.01). Variation in NH₃ VCDs are positively correlated with CO VCDs (Fig. 4a, S6), which are also increasing during this 363 364 period (Fig. 4c, S4). 365 These correlations imply a biomass burning source for the increasing NH₃ VCDs in 366 West Africa; although the burned area trends are not as clear, it is important to remember 367 that MODIS undercounts burned area during this time of year by a factor of 3 to 6, and so 368 would be less sensitive to trends (Ramo et al., 2021; Roteta et al., 2019). Although there is 369 considerable gas flaring in Nigeria, gas flaring emissions have exhibited long-term negative 370 trends (Doumbia et al., 2019). In addition, although NO2 VCDs were found to decrease 371 across the productive savannas of West Africa, regions of increasing NO₂ VCDs were 372 observed over large parts of Nigeria, further suggesting that there may be increases—or at 373 least smaller decreases—in biomass burning in the country (Hickman et al., 2021a). It is 374 unlikely that changes in chemical sinks—specifically, the formation of nitrate aerosols in 375 reactions with NO_x or sulfate—are responsible for the increasing trend: the observed 376 increase in NO2 VCDs observed during February and March would be expected to lead to a 377 shorter NH_3 lifetime and decreasing VCDs. In addition, emissions of SO_2 are relatively low 378 in West Africa, with moderate emissions occurring in Nigeria, but neither emissions nor 379 lifetime exhibit clear seasonal variation (Lee et al., 2011). 380 Small agricultural fires are likely an important contributor to the increasing NH₃ 381 VCDs during the dry-to-rainy season transitional period—a period when agricultural fires 382 are common in the region (Korontzi et al., 2006). There are large numbers of small fires 383 that are not detected by MODIS during these months: as noted above, estimates of burned 384 area during February, March, and April are revised upwards by roughly a factor of 3 to 6 385 over MODIS when small fires are included (Ramo et al., 2021; Roteta et al., 2019). Many of 386 these small fires are likely related to agricultural field preparation prior to planting 387 (Gbadegesin and Olusesi, 1994), which typically takes place in March or April (Vrieling et 388 al., 2011; Yegbemey et al., 2014). An increase in fires during this transitional period is also

emissions of less oxidized species such as CO and NH₃, rather than the more fully oxidized

358

Deleted: 7

391 consistent with one of the primary mechanisms behind the overall decline in burned area: 392 roughly half of the decline is attributed to increased population density and the expansion 393 of agricultural area, which contributes to the anthropogenic suppression of larger fires 394 (Andela et al., 2017; Andela and van der Werf, 2014). This agricultural expansion, 395 however, can be expected to be accompanied by increases in small fires used for the 396 removal of stubble or harvest byproduct (Gbadegesin and Olusesi, 1994), leading to the 397 increased emissions during the rainy-to-dry season transition observed here. 398 Globally, agricultural emissions from fertilized soils and livestock excreta are the 399 largest source of NH₃ (Bauer et al., 2016), and Warner et al. (Warner et al., 2017) suggest 400 that national-scale changes in fertilizer use could explain the NH₃ trend over Nigeria. 401 However, as noted above, much of the increase in West Africa occurs prior to the start of 402 the planting season—before fertilizer is applied—and appears likely to be due to biomass 403 burning emissions instead, potentially related to field preparation. Fertilizer or manure 404 may make a contribution to the increasing trend later in the year, as NH₃ VCDs increase in 405 the wet savanna during May, June, and July (Fig. 3), though there are also significant 406 correlations between NH₃ and CO VCDs (Fig. 4), suggesting that biomass burning may 407 continue to play an important role. However, average N fertilizer use in West Africa is universally under 40 kg N ha⁻¹ yr⁻¹, typically under 20 kg N ha⁻¹ yr⁻¹, and is under 10 kg N 408 409 ha⁻¹ yr⁻¹ in Nigeria—over an order of magnitude lower than rates in Europe, the United 410 States, and China (FAO, 2020). Although percentage changes in fertilizer use are 411 substantial, in absolute terms they represent increases of less than 2 kg N ha⁻¹ yr⁻¹, and 412 frequently less than 1 kg N ha⁻¹ yr⁻¹, a relatively small but perhaps not entirely trivial 413 perturbation to the N cycle; Between 2000 and 2007, total N deposition averaged 8.38 kg N Deleted: . 414 ha-1 yr-1 in wet savanna and 14.75 kg N ha-1 yr-1 in forest ecosystems based on surface 415 sampling sites (Galy-Lacaux and Delon, 2014), and biological N fixation in tropical and wet 416 savannas has been estimated as ranging from 16 to 44 kg N ha⁻¹ yr⁻¹ (Bustamante et al., 417 2006). These estimates suggest, that fertilizer increases may represent a 1 to 2% annual Deleted: , suggesting 418 increase in N inputs. But given the small magnitude of fertilizer applications, it appears 419 unlikely that changes in fertilizer use can explain the entirety of NH₃ increases during the 420 growing season. Our analyses do suggest that livestock may contribute to increasing NH₃ 421 VCDs over the Sahel, from roughly 15 to 18N (Fig. 2a). However, many of these pixels are

424 also those where population density appears to be playing a role (Figure 2b) and where

425 <u>correlations between NH₃ and CO VCDs are present during the transition from the dry to</u>

426 rainy season (Fig. S7), which may reflect a contribution from agricultural fires.

427

428 **3.3. South Sudan**

429 The most notable declining trend in NH₃ VCDs occurs in South Sudan over the Sudd wetlands at a rate of over 2% yr⁻¹ (Fig. 1b; p=0.20). It appears that this decline is related to 430 431 interannual variation in the flooded extent of the Sudd, a vast wetland that connects the 432 White and Blue Nile tributaries. Seasonal variation of inflow to the Sudd leads to variation 433 in flooded extent: an area of roughly 15,000 km² is permanently flooded, and another 434 roughly 15,000 km² is temporarily flooded each year, with considerable interannual 435 variation in the total flooded area (Di Vittorio and Georgakakos, 2018). Among other 436 factors, drying soils should increase production and emissions of NH3 from soils, as Eq. 1 is 437 shifted to the right (Clarisse et al., 2019). Earlier work evaluating an NH₃ hotspot over Lake 438 Natron in Tanzania found that the drying of seasonally flooded soils leads to large 439 emissions of NH₃: As the waters of Lake Natron recede during the dry season each year 440 and the surrounding mud flats dry out, NH₃ VCDs increase rapidly, with hotspots appearing 441 over the mudflats (Clarisse et al., 2019). These elevated VCDs are attributed to multiple 442 possible factors, including the effects of drying on concentrations of NH₃ in solution (which 443 increases the concentration gradient with the atmosphere), reduced biological uptake of 444 NH₃, convective transport of dissolved NH₃ from depth to the soil surface, and increased mineralization of labile organic matter (Clarisse et al., 2019). 445 446 We find the same clear seasonal relationship between wetland flooded extent and 447 NH₃ concentrations over the Sudd—VCDs increase as waters recede from the temporarily 448 flooded area, leading to annual maxima from February through May (Fig. 5a; bounding box 449 of 29E to 31.5E and 6N to 9.9N). Like the entire country, seasonal variation in NH₃ VCDs 450 over the Sudd follow variation in surface temperature, but NH3 concentrations over the

451 Sudd are substantially elevated compared to surrounding regions during this time of year

but not others, suggesting that a mechanism in addition to temperature is contributing to

Deleted: n











471 and the entirety of South Sudan for the period 2008 through 2017.

480	It is possible that conflict in South Sudan could contribute to the decline in NH_3	
481	VCDs. In 2013, a civil conflict emerged in South Sudan that was ultimately responsible for	
482	the displacement of millions of people (Global Internal Displacement Monitoring Centre,	
483	2020; World Bank, 2019) and the disruption of livestock migration patterns (Idris, 2018).	
484	However, these disruptions appeared only after the onset of the long-term change in NH_3 ,	
485	and appear unlikely to make an important contribution to the observed interannual	
486	variation (SI Text, Fig. S <mark>9, S10)</mark> .	Deleted: 7
		Deleted: 8
487	It is unlikely that changes in chemical sinks are responsible for the decline in NH_3	
488	VCDs. VCDs of tropospheric NO_2 are also decreasing in the region (Fig. S11), which is	Deleted: 9
489	suggestive of less formation of particulate-phase ammonium rather than more.	
490	Anthropogenic SO_2 emissions in Africa in general and South Sudan in particular are very	
491	low (European Commission Joint Research Centre (JRC)/Netherlands Environmental	
492	Assessment Agency (PBL), 2016), and would not be expected to be emitted from the Sudd;	
493	more generally, the clear spatial association between the NH_3 trend and the Sudd (Fig. 1,	
494	Fig. S12) is strongly suggestive of changes in emissions rather than atmospheric processes	Deleted: 0
495	being responsible for the trend.	
496		
407		
497	3.4 Lake victoria Basin region	
498	The Lake Victoria Basin and its surroundings—an area including elevated mean NH_3	
499	VCDs—exhibit an increasing NH3 trend (Fig. 1b, Fig. 7, Fig. S13), which appears to be the	Deleted: 1
500	result of increasing agricultural activity in the area. The region includes a high and	
501	increasing density of agricultural land (Fig. 1h, Fig. 2d, Fig. S1 <u>4</u>), and these increases in	Deleted: 2
502	cropped area are positively correlated with increases in NH_3 VCDs across much of the	
503	region (Fig. 2c). The northern and southern halves of the Lake Victoria region <u>—which</u>	
504	straddles the equator—have distinct growing seasons: in the north, the season generally	
I		

511	starts in April, whereas in the south, it starts in November or December (Vrieling et al.,	
512	2011). <u>Some of the long-term trend reflects this seasonality, with increases in the north</u>	Deleted: The
513	and south occurring during their respective growing seasons (Fig. 3 <u>, Fig. S15</u>). Fertilizer	
514	use in the Lake Victoria region is low: national averages range from about 1 to 3 kg	
515	nutrients ha $^{\cdot 1}$ yr $^{\cdot 1}$ in Uganda to about 35 to 40 kg nutrients ha $^{\cdot 1}$ yr $^{\cdot 1}$ in Kenya (Elrys et al.,	
516	2019; World Bank, 2019); to put these numbers in context, Organization for Economic	
517	Cooperation and Development (OECD) countries use about 135-140 kg nutrients ha $^{-1}$ yr $^{-1}$	
518	(World Bank, 2019). Although rates of fertilizer use have increased by substantial	
519	proportions, the absolute amount of increase is relatively small, typically roughly 1 to $10~\mathrm{kg}$	
520	nutrients decade ⁻¹ . Unlike in West Africa, however, interannual variation in burned area	
521	(Fig. 7. S16) does not exhibit a clear relationship with changes in NH ₃ VCDs. Consequently,	Deleted: 3
522	we expect that both the expansion and intensification of agriculture in the region	
523	contribute to the increasing NH_3 VCDs.	
524	We note that there is a negative correlation between cropland area and NH ₃ VCDs in	
525	<u>Uganda, north of Lake Victoria (Figure 7b). We expect this is a consequence of the</u>	
526	<u>extremely low fertilizer use in Uganda (Masso et al., 2017) , which leads to depletion of soil</u>	
527	<u>N—and thus substrate for ammonia volatilization—over time (</u> Cobo et al., 2010) <u>.</u>	
528	We also note that there is an apparent increase in NH3 VCDs over the lake itself. It is	
529	important to note that differences in conditions over the lake and adjacent land cover—e.g.,	
530	emissivity, thermal contrast, etc.—contribute to substantial differences in mean retrieved	
531	$\rm NH_3$ VCDs over the lake relative to the surrounding land surface. Both monthly and	
532	interannual variation in NH ₃ VCDs over Lake Victoria correspond closely to variation in	
533	<u>NH₃ VCDs over the surrounding land surface (Figure <mark>\$</mark>17, S18), suggesting that the trend</u>	Formatted: Highlight
534	over the lake results from transport of NH_3 emitted from the surrounding land surface.	





Figure 7. Changes in NH₃ VCDs and their relationship with burned area and 538 539 cropped area over the Lake Victoria region for the 2008 through 2018 period. (a) 540 Correlation coefficients for the relationship between NH₃ VCDs and burned area. (b) 541 Correlation coefficients for the relationship between NH₃ VCDs and cropped area, including 542 mosaics of crops and natural vegetation cover. (c) Changes in NH₃ VCDs. 543 544 3.5 National-scale relationships 545 Examining relationships at a national scale can provide insight into relationships 546 between changes in agricultural or biomass burning and changes in atmospheric NH₃ VCDs 547 at larger scales. When grouping countries into three bins based on their annual percentage 548 changes in NH₃ VCDs, there is some evidence for a broad relationship between livestock 549 and NH₃ VCDs at the national scale (Fig. 8). The rate of change in national-scale NH₃ VCDs 550 varies significantly among bins (p<0.001; rank transformed, though note that residuals 551 may still deviate from normality). The annual percentage changes in livestock in TLUs vary 552 significantly by bin (p=0.042; rank transformed), with the middle bin higher than the 553 bottom bin (p=0.1) and the high bin higher than the bottom bin (p=0.06). Annual 554 percentage changes in fertilizer N (p=0.58) and crop production (p=0.62; rank

555 transformed) did not vary by bin.

-(Deleted: 56
-(Deleted: 03
-(Deleted: , though not significantly so
•(Deleted: 21
Y	Deleted: ,
ľ	Deleted: 23





Figure 8. Annual percentage changes in national mean annual NH₃ VCDs, burned area, CO
VCDs, livestock, crop yield, and fertilizer N use for African countries with low, medium, or
high rates of NH₃ VCD change. Error bars represent the standard error of the mean. See
Table S1 for the list of countries in each bin and Fig. S19, for an expanded set of variables.

569

570 571 Instead of a direct agricultural relationship with changes in NH₃ VCDs, there is the 572 possibility that changes in biomass burning are associated with changes in NH₃ VCDs. 573 Although the differences in the annual percentage change in burned area were not 574 significant among bins (p=0.54; rank transformed), the overall pattern is consistent with 575 earlier results finding that a reduction in burned area across the northern biomass burning 576 region was associated in part with the expansion of agriculture and presumed

577 anthropogenic suppression of fire (Andela et al., 2017; Andela and van der Werf, 2014).

 $578 \qquad \text{However, burned area as measured by MODIS is likely an imperfect predictor for NH_3}$

Deleted: 48

Deleted: 4

582	emissions—as noted previously, MODIS underestimates burned area by a factor of 3 to 6	
583	during shoulder seasons (Roteta et al., 2019), which is when fires are expected to emit	
584	more reduced species such as NH_3 (Zheng et al., 2018). In contrast to burned area, the	
585	annual change in column densities of CO—which tends to be co-emitted with $\rm NH_3$ from	
586	fires—differed significantly among bins (p<0.001; rank transformed) and was significantly	
587	higher in the high bin than in the low or medium bins, (p<0.001, post-hoc tests). The higher	
588	annual CO changes in the high bin could related to larger anthropogenic fossil fuel	
589	emissions, but we see no difference among bins in growth rates of CO_2 emissions (p=0.48;	
590	Figure S1 $\underline{9}$; such a difference would be expected if differences in economic development	
591	were responsible for the CO differences. These results leave open the possibility that	
592	changes in either biofuel emissions or biomass burning emissions—perhaps from smaller	
593	fires not observed in the MODIS burned area product—may be primarily responsible for	
594	the difference in CO between bins, and may be contributing to the differences in \ensuremath{NH}_3	
595	between bins. Changes in NO_2 VCDs and SO_2 concentrations can affect the lifetime of NH_3	
596	(the latter by changing SO $_4$ concentrations), but do not appear to make an important	
597	contribution to the observed trends in NH_3 VCDs among bins (Fig. S12, SI text).	
598	Temperature, likewise, does not appear to play an important role (SI text).	
599		
600	4. Conclusion	
601	Using IASI, we have observed both increases and decreases in atmospheric $\rm NH_3$	
602	VCDs in different regions in Africa between 2008 and 2018, with different factors affecting	
603	trends in different regions.	
604	We observed increases in NH_3 VCDs in West Africa, which earlier work had	
605	concluded was likely related to increased fertilizer use. Fertilizer is not typically applied in	
606	West Africa until the start of the growing season—often April—but we find that much of	
607	the $\rm NH_3$ increase occurs during February and March, suggesting that increasing fertilizer	
608	use is unlikely to provide a complete explanation for the $\rm NH_3$ trend. Agriculture may	
609	nevertheless play a role, with enhanced burned area and especially CO concentrations in	
610	February suggestive of increased burning of crop stubble in preparation for planting during	
611	this time of year. Fires in this region tend to emit a greater proportion of less oxidized	

Deleted:

Deleted: Deleted: 45 Deleted: 4

617 species such as NH₃ at the end of the dry season, consistent with a biomass burning source 618 for the increasing NH₃ VCDs. 619 Decreases in NH₃ VCDs were largest in South Sudan, especially over the Sudd 620 wetland, where NH₃ VCDs vary seasonally with the extent of area flooded. As the 621 temporarily flooded areas of the Sudd dry out each year, NH₃ VCDs increase as reduction in 622 soil moisture drives increased production and volatilization of NH₃. The area of the Sudd 623 that is flooded each year varies, and from 2008 until 2015, the area that remains flooded 624 during the dry season generally increased, producing a positive overall trend for the period 625 of 2008 through 2017. This increase in the dry season flooded area drove a decrease in 626 NH₃ VCDs: with less soil drying out, the seasonal maxima in NH₃ VCDs were lower. 627 Although it is possible that conflict in South Sudan could contribute to changes in NH₃ 628 VCDs, the timing and distribution of conflict events and human displacement suggest that 629 other factors are likely more important. 630 Modest increases in NH₃ VCDs were observed in the Lake Victoria region. This 631 region has experienced increases in agricultural area during the IASI observation period, 632 and these changes explained a large proportion of the variation in NH₃ VCDs across large 633 patches of the region, where biomass burning could not. We expect that both expansion 634 and intensification of agriculture in this region could contribute to the positive NH_3 trend. 635 Considering national-scale statistics, comparisons between equally sized bins of 17 countries each suggested that changes in biomass burning emissions and livestock 636 637 emissions could contribute to differences in NH₃ VCDs among countries, but variables 638 related to cropped agriculture such as cropped area or fertilizer N use did not appear to be 639 important factors at this scale. This may be because although fertilizer use has been 640 increasing in sub-Saharan Africa, it remains extremely low relative to other continents, and 641 relative to the levels needed to attain food security. Average fertilizer use in most 642 countries in the region is under 20 kg N ha⁻¹ yr⁻¹, and sometimes less than 5 kg N ha⁻¹ yr⁻¹. 643 Although recommended fertilizer rates are lower in most African countries than in the U.S. 644 or Europe, increasing N inputs to 50 or, 100 kg N ha⁻¹ yr⁻¹ would represent a major 645 perturbation to the regional N cycle, and potentially a large new source of NH₃ to the 646 atmosphere. West Africa is already a global NH₃ hotspot (Van Damme et al., 2018), 647 suggesting that encouraging policies that can help to limit NH₃ emissions during the early

Deleted: ,

Deleted: , or 150

- 650 stages of agricultural intensification in Africa may help mitigate potential impacts on the atmosphere. Fortunately, agricultural practices such as sub-surface application of 651 652 fertilizer, which is already being promoted to smallholder farmers, can serve to both limit 653 NH₃ emissions also help to increase crop yields. 654 These past and anticipated future trends also make the case for expanding capacity 655 for atmospheric monitoring in sub-Saharan Africa. Although long-term monitoring 656 networks have been established in West Africa (Adon et al., 2010; Ossohou et al., 2019) and 657 South Africa (Conradie et al., 2016) as part of the INDAAF network, it is mainly focused on 658 deposition and the spatio-temporal resolution of surface measurements is very coarse 659 when compared to the data available in other parts of the world, which will limit our ability 660 to understand how agricultural and socio-economic development in Africa affect the 661 atmosphere. Satellite observations can help to bridge some of these data gaps, but have 662 their own spatio-temporal limitations, and would further benefit from additional high-663 quality surface observations for evaluation of retrieval products.
- 664

665 Data availability: All data used in this study are available from public sources, with the 666 exception of Sudd wetland extent, which is available by request from Courtney Di Vittorio. The 667 IASI NH3 and CO data are available from The IASI https://iasi.aeris-data.fr. The NOAA Global 668 Surface Dataset available https://data.nodc.noaa.gov/cgi-Temperature is at bin/iso?id=gov.noaa.ncdc:C01585. MODIS 669 burned area data are available from https://www.globalfiredata.org/data.html. MODIS agricultural area are available at 670 https://lpdaac.usgs.gov/products/mcd12c1v006/. TRMM 3B42 precipitation data are available 671 672 from https://pmm.nasa.gov/data-access/downloads/trmm. The Gridded Livestock of the World 673 data are available from https://livestock.geo-wiki.org/home-2/. Population density data for 2017 674 are available at https://landscan.ornl.gov/downloads/2017. FAO national crop production and 675 fertilizer N data are available at http://www.fao.org/faostat/en/. Data on conflict events from 676 ACLED are available at https://acleddata.com/#/dashboard. World Bank national statistics 677 on refugees and internally displaced people are available at https://data.worldbank.org.

678

Deleted: ,

Deleted: and

681	Author Contribution: J.E.H. designed the study, conducted the analysis, and wrote the paper.
682	NA, ED, CD, MO, CG-L, KT, and SEB contributed to study design and edited the paper. LC, P-
683	FC, and MVD developed the original IASI trace gas retrievals and edited the paper.
(04	
084 koz	The authors declare that they have no conflict of interest.
685	
686	Acknowledgements: J.E.H.'s research was supported by an appointment to the NASA
687	Postdoctoral Program at the NASA Goddard Institute for Space Studies administered by
688	Universities Space Research Association under contract with NASA.
689	
690	
691	References
692	van der A, R. J., Eskes, H. J., Boersma, K. F., van Noije, T. P. C., Van Roozendael, M., De Smedt,
693	I., Peters, D. H. M. U. and Meijer, E. W.: Trends, seasonal variability and dominant NOx
694	source derived from a ten year record of NO2 measured from space, J. Geophys. Res.
695	Atmos., 113(4), D04302, doi:10.1029/2007JD009021, 2008.
696	Adon, M., Galy-Lacaux, C., Yoboué, V., Delon, C., Lacaux, J. P., Castera, P., Gardrat, E., Pienaar,
697	J., Al Ourabi, H., Laouali, D., Diop, B., Sigha-Nkamdjou, L., Akpo, a., Tathy, J. P., Lavenu, F. and
698	Mougin, E.: Long term measurements of sulfur dioxide, nitrogen dioxide, ammonia, nitric
699	acid and ozone in Africa using passive samplers, Atmos. Chem. Phys., 10(15), 7467–7487,
700	doi:10.5194/acp-10-7467-2010, 2010.
701	AGRA: AGRA in 2008: Building on the New Momentum in African Agriculture., 2009.
702	Andela, N. and van der Werf, G. R.: Recent trends in African fires driven by cropland
703	expansion and El Niño to la Niña transition, Nat. Clim. Chang., 4(9), 791–795,
704	doi:10.1038/nclimate2313, 2014.
705	Andela, N., Morton, D. C., Giglio, L., Chen, Y., Van Der Werf, G. R., Kasibhatla, P. S., DeFries, R.
706	S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon,
707	S., Melton, J. R., Yue, C. and Randerson, J. T.: A human-driven decline in global burned area,
708	Science (80)., 356(6345), 1356–1362, doi:10.1126/science.aal4108, 2017.

- 709 Anker, H. T., Baaner, L., Backes, C., Keessen, A. and Möckel, S.: Comparison of ammonia
- 710 regulation in Germany , the Netherlands and Denmark legal framework, Copenhagen.,

- 711 2018.
- 712 Bauer, S. E., Tsigaridis, K. and Miller, R.: Significant atmospheric aerosol pollution caused by
- vorld food cultivation, Geophys. Res. Lett., 43(10), 5394–5400,
- 714 doi:10.1002/2016GL068354, 2016.
- 715 Behera, S. N., Sharma, M., Aneja, V. P. and Balasubramanian, R.: Ammonia in the
- 716 atmosphere: A review on emission sources, atmospheric chemistry and deposition on
- 717 terrestrial bodies, Environ. Sci. Pollut. Res., 20(11), 8092–8131, doi:10.1007/s11356-013-
- 718 2051-9, 2013.
- 719 Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van Der Hoek, K. W. and Olivier, J.
- 720 G. J.: A global high-resolution emission inventory for ammonia, Global Biogeochem. Cycles,
- 721 11(4), 561–587, 1997.
- 722 Bustamante, M. M. C., Medina, E., Asner, G. P., Nardoto, G. B. and Garcia-Montiel, D. C.:
- Nitrogen cycling in tropical and temperate savannas, Biogeochemistry, 79(1-2), 209–237,
- 724 doi:10.1007/s10533-006-9006-x, 2006.
- 725 Cahoon, D. R., Stocks, B. J., Levine, J. S., Cofer, W. R. and O'Neill, K. P.: Seasonal distribution of
- 726 African savanna fires, Nature, 359(6398), 812–815, doi:10.1038/359812a0, 1992.
- 727 Chen, L.-W. A., Verburg, P., Shackelford, A., Zhu, D., Susfalk, R., Chow, J. C. and Watson, J. G.:
- 728 Moisture effects on carbon and nitrogen emission from burning of wildland biomass,
- 729 Atmos. Chem. Phys., 10(14), 6617–6625, doi:10.5194/acp-10-6617-2010, 2010.
- 730 Chilonda, P. and Otte, J.: Indicators to monitor trends in livestock production at national,
- regional and international levels, Livest. Res. Rural Dev., 18(8), 1–12, 2006.
- 732 Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D. and Coheur, P. F.: Global ammonia
- distribution derived from infrared satellite observations, Nat. Geosci., 2(7), 479-483,
- 734 doi:10.1038/ngeo551, 2009.
- 735 Clarisse, L., Van Damme, M., Gardner, W., Coheur, P.-F., Clerbaux, C., Whitburn, S., Hadji-
- 736 Lazaro, J. and Hurtmans, D.: Atmospheric ammonia (NH3) emanations from Lake Natron's
- 737 saline mudflats, Sci. Rep., 9, 4441, doi:10.1038/s41598-019-39935-3, 2019.
- 738 Cobo, J. G., Dercon, G. and Cadisch, G.: Nutrient balances in African land use systems across
- 739 different spatial scales: A review of approaches, challenges and progress, Agric. Ecosyst.
- 740 Environ., 136(1-2), 1-15, doi:10.1016/j.agee.2009.11.006, 2010.
- 741 Conradie, E. H., Van Zyl, P. G., Pienaar, J. J., Beukes, J. P., Galy-Lacaux, C., Venter, A. D. and

- 742 Mkhatshwa, G. V.: The chemical composition and fluxes of atmospheric wet deposition at
- four sites in South Africa, Atmos. Environ., 146, 113–131,
- 744 doi:10.1016/j.atmosenv.2016.07.033, 2016.
- 745 Van Damme, M., Wichink Kruit, R. J., Schaap, M., Clarisse, L., Clerbaux, C., Coheur, P. F.,
- 746 Dammers, E., Dolman, A. J. and Erisman, J. W.: Evaluating 4 years of atmospheric ammonia
- 747 (NH3) over Europe using IASI satellite observations and LOTOS-EUROS model results, J.
- 748 Geophys. Res., 119(15), 9549–9566, doi:10.1002/2014JD021911, 2014a.
- 749 Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J.,
- 750 Erisman, J. W. and Coheur, P. F.: Global distributions, time series and error characterization
- of atmospheric ammonia NH3 from IASI satellite observations, Atmos. Chem. Phys., 14(6),
- 752 2905–2922, doi:10.5194/acp-14-2905-2014, 2014b.
- 753 Van Damme, M., Clarisse, L., Dammers, E., Liu, X., Nowak, J. B., Clerbaux, C., Flechard, C. R.,
- Galy-Lacaux, C., Xu, W., Neuman, J. A., Tang, Y. S., Sutton, M. A., Erisman, J. W. and Coheur, P.
- 755 F.: Towards validation of ammonia (NH3) measurements from the IASI satellite, Atmos.
- 756 Meas. Tech., 8(3), 1575–1591, doi:10.5194/amt-8-1575-2015, 2015.
- 757 Van Damme, M., Whitburn, S., Clarisse, L., Clerbaux, C., Hurtmans, D. and Coheur, P. F.:
- 758 Version 2 of the IASI NH3 neural network retrieval algorithm: Near-real-time and
- reanalysed datasets, Atmos. Meas. Tech., 10(12), 4905-4914, doi:10.5194/amt-10-49052017, 2017.
- 761 Van Damme, M., Clarisse, L., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C. and
- Coheur, P.: Industrial and agricultural ammonia point sources exposed, Nature, 564, 99–103, 2018.
- 764 Van Damme, M., Clarisse, L., Franco, B., Sutton, M. A., Erisman, J. W., Wichink Kruit, R., van
- 765 Zanten, M., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C. and Coheur, P.-F.:
- 766 Global, regional and national trends of atmospheric ammonia derived from a decadal
- 767 (2008-2018) satellite record, Environ. Res. Lett., 2021.
- 768 Dammers, E., Palm, M., Van Damme, M., Vigouroux, C., Smale, D., Conway, S., Toon, G. C.,
- 769 Jones, N., Nussbaumer, E., Warneke, T., Petri, C., Clarisse, L., Clerbaux, C., Hermans, C.,
- 770 Lutsch, E., Strong, K., Hannigan, J. W., Nakajima, H., Morino, I., Herrera, B., Stremme, W.,
- 771 Grutter, M., Schaap, M., Kruit, R. J. W., Notholt, J., Coheur, P. F. and Erisman, J. W.: An
- 772 evaluation of IASI-NH3 with ground-based Fourier transform infrared spectroscopy

- 773 measurements, Atmos. Chem. Phys., 16(16), 10351-10368, doi:10.5194/acp-16-10351-
- 2016, 2016.
- 775 Denier Van Der Gon, H. and Bleeker, A.: Indirect N2O emission due to atmospheric N
- deposition for the Netherlands, Atmos. Environ., 39(32), 5827–5838,
- 777 doi:10.1016/j.atmosenv.2005.06.019, 2005.
- 778 Dobson, J. E., Bright, E. A., Coleman, P. R., Durfee, R. C. and Worley, B. A.: A global population
- database for estimating populations at risk, Photogramm. Eng. Remote Sens., 66(7), 849-
- 780 857 [online] Available from: In Book..., 2000.
- 781 Doumbia, E. H. T., Liousse, C., Keita, S., Granier, L., Granier, C., Elvidge, C. D., Elguindi, N. and
- 782 Law, K.: Flaring emissions in Africa: Distribution, evolution and comparison with current
- 783 inventories, Atmos. Environ., 199(November 2018), 423-434,
- 784 doi:10.1016/j.atmosenv.2018.11.006, 2019.
- 785 Elrys, A. S., Abdel-Fattah, M. K., Raza, S., Chen, Z. and Zhou, J.: Spatial trends in the nitrogen
- 786 budget of the African agro-food system over the past five decades, Environ. Res. Lett.,
- 787 14(12), doi:10.1088/1748-9326/ab5d9e, 2019.
- 788 European Commission Joint Research Centre (JRC)/Netherlands Environmental
- 789 Assessment Agency (PBL): Emission Database for Global Atmospheric Research (EDGAR),
- 790 release version 4.3.1, 2016.
- 791 Fan, Y. and van den Dool, H.: A global monthly land surface air temperature analysis for
- 792 1948-present, J. Geophys. Res. Atmos., 113(1), D01103, doi:10.1029/2007JD008470, 2008.
- 793 FAO: FAO Statistics Database, FAOSTAT Stat. Database [online] Available from:
- http://www.fao.org/faostat/en/ (Accessed 1 January 2019), 2020.
- 795 Friedl, M. A., McIver, D. K., Hodges, J. C. F., Zhang, X. Y., Muchoney, D., Strahler, A. H.,
- 796 Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F. and Schaaf, C.: Global
- 797 land cover mapping from MODIS: Algorithms and early results, Remote Sens. Environ.,
- 798 83(1–2), 287–302, doi:10.1016/S0034-4257(02)00078-0, 2002.
- 799 Galy-Lacaux, C. and Delon, C.: Nitrogen emission and deposition budget in West and Central
- 800 Africa, Environ. Res. Lett., 9(12), doi:10.1088/1748-9326/9/12/125002, 2014.
- 801 Gbadegesin, A. and Olusesi, B. B.: Effects of land clearing methods on soil physical and
- 802 hydrological properties in southwestern Nigeria, Environmentalist, 14(4), 297–303
- 803 [online] Available from: http://www.scopus.com/inward/record.url?eid=2-s2.0-

- 804 0022858519&partnerID=40&md5=3051b1dd91ff2990e37fe7466872923e, 1994.
- 805 George, M., Clerbaux, C., Hurtmans, D., Turquety, S., Coheur, P. F., Pommier, M., Hadji-
- 806 Lazaro, J., Edwards, D. P., Worden, H., Luo, M., Rinsland, C. and McMillan, W.: Carbon
- 807 monoxide distributions from the IASI/METOP mission: Evaluation with other space-borne
- remote sensors, Atmos. Chem. Phys., 9(21), 8317–8330, doi:10.5194/acp-9-8317-2009,
 2009.
- 810 Giglio, L., Boschetti, L., Roy, D. P., Humber, M. L. and Justice, C. O.: The Collection 6 MODIS
- 811 burned area mapping algorithm and product, Remote Sens. Environ., 217, 72–85,
- 812 doi:10.1016/j.rse.2018.08.005, 2018.
- 813 Global Internal Displacement Monitoring Centre: Global Internal Displacement Database,
- 814 [online] Available from: https://www.internal-displacement.org/database/displacement-815 data, 2020.
- 816 Goode, J. G., Yokelson, R. J., Susott, R. A. and Ward, D. E.: Trace gas emissions from
- 817 laboratory biomass fires measured by open-path Fourier transform infrared spectroscopy;,
- 818 J. Chem. Inf. Model., 104(D17), 21237–21245, 1999.
- 819 Guo, X., Clarisse, L., Wang, R., Van Damme, M., Whitburn, S., Coheur, P., Clerbaux, C., Franco,
- 820 B., Pan, D., Golston, L. M., Wendt, L., Sun, K., Tao, L., Miller, D., Mikoviny, T., Müller, M.,
- 821 Wisthaler, A., Tevlin, A. G., Murphy, J. G., Nowak, J. B., Roscioli, J. R., Volkamer, R., Kille, N.,
- 822 Neuman, J. A., Eilerman, S. J., Crawford, J. H., Yacovitch, T. I., Barrick, J. D., Scarino, A. J. and
- 823 Zondlo, M. A.: Validation of IASI satellite ammonia observations at the pixel scale using in-
- situ vertical profiles, J. Geophys. Res. Atmos., 126, e2020JD033475,
- 825 doi:10.1029/2020jd033475, 2021.
- 826 Hazell, P. and Wood, S.: Drivers of change in global agriculture., Philos. Trans. R. Soc. Lond.
- 827 B. Biol. Sci., 363(1491), 495–515, doi:10.1098/rstb.2007.2166, 2008.
- 828 Hickman, J. E., Dammers, E., Galy-Lacaux, C. and Van Der Werf, G. R.: Satellite evidence of
- substantial rain-induced soil emissions of ammonia across the Sahel, Atmos. Chem. Phys.,
 18(22), 16713–16727, doi:10.5194/acp-18-16713-2018, 2018.
- 831 Hickman, J. E., Andela, N., Tsigaridis, K., Galy-Lacaux, C., Ossohou, M. and Bauer, S. E.:
- 832 Reductions in NO 2 burden over north equatorial Africa from decline in biomass burning in
- spite of growing fossil fuel use, 2005 to 2017, Proc. Natl. Acad. Sci., 118(7), e2002579118,
- 834 doi:10.1073/pnas.2002579118, 2021a.

- 835 Hickman, J. E., Dammers, E., Galy-Lacaux, C., Ossohou, M. and Bauer, S. E.: Continental and
- 836 ecoregion-specific drivers of atmospheric NO2 and NH3 seasonality over Africa revealed by
- 837 satellite observations, Global Biogeochem. Cycles, 35, e2020GB006916,
- 838 doi:10.1029/2020GB006916, 2021b.
- 839 Huffman, G. J., Adler, R. F., Bolvin, D. T., Gu, G., Nelkin, E. J., Bowman, K. P., Hong, Y., Stocker,
- 840 E. F. and Wolff, D. B.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global,
- 841 Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales, J. Hydrometeorol., 8(1),
- 842 38–55, doi:10.1175/jhm560.1, 2007.
- 843 Hurtmans, D., Coheur, P. F., Wespes, C., Clarisse, L., Scharf, O., Clerbaux, C., Hadji-Lazaro, J.,
- 844 George, M. and Turquety, S.: FORLI radiative transfer and retrieval code for IASI, J. Quant.
- 845 Spectrosc. Radiat. Transf., 113(11), 1391–1408, doi:10.1016/j.jqsrt.2012.02.036, 2012.
- 846 Idris, I.: Livestock and conflict in South Sudan K4D Helpdesk Report 484, Brighton.
- 847 [online] Available from:
- 848 https://assets.publishing.service.gov.uk/media/5c6abdec40f0b61a22792fd5/484_Livest
- 849 ock_and_Conflict_in_South_Sudan.pdf, 2018.
- 850 Kanter, D. R.: Nitrogen pollution: a key building block for addressing climate change, Clim.
- 851 Change, 147(1–2), 11–21, doi:10.1007/s10584-017-2126-6, 2018.
- 852 Kerzenmacher, T., Dils, B., Kumps, N., Blumenstock, T., Clerbaux, C., Coheur, P. F., Demoulin,
- 853 P., García, O., George, M., Griffith, D. W. T., Hase, F., Hadji-Lazaro, J., Hurtmans, D., Jones, N.,
- 854 Mahieu, E., Notholt, J., Paton-Walsh, C., Raffalski, U., Ridder, T., Schneider, M., Servais, C. and
- 855 De Mazière, M.: Validation of IASI FORLI carbon monoxide retrievals using FTIR data from
- 856 NDACC, Atmos. Meas. Tech., 5(11), 2751–2761, doi:10.5194/amt-5-2751-2012, 2012.
- 857 Korontzi, S., McCarty, J., Loboda, T., Kumar, S. and Justice, C.: Global distribution of
- 858 agricultural fires in croplands from 3 years of Moderate Resolution Imaging
- 859 Spectroradiometer (MODIS) data, Global Biogeochem. Cycles, 20(2), GB2021,
- 860 doi:10.1029/2005GB002529, 2006.
- 861 Krupa, S. V.: Effects of atmospheric ammonia (NH3) on terrestrial vegetation: A review,
- 862 Environ. Pollut., 124(2), 179–221, doi:10.1016/S0269-7491(02)00434-7, 2003.
- 863 Lee, C., Martin, R. V., Van Donkelaar, A., Lee, H., Dickerson, R. R., Hains, J. C., Krotkov, N.,
- 864 Richter, A., Vinnikov, K. and Schwab, J. J.: SO2 emissions and lifetimes: Estimates from
- 865 inverse modeling using in situ and global, space-based (SCIAMACHY and OMI)

- 866 observations, J. Geophys. Res. Atmos., 116(6), 1–13, doi:10.1029/2010JD014758, 2011.
- 867 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D. and Pozzer, A.: The contribution of outdoor
- air pollution sources to premature mortality on a global scale, Nature, 525(7569), 367–371,
- 869 doi:10.1038/nature15371, 2015.
- 870 Masso, C., Nziguheba, G., Mtutegi, J., Galy-Lacaux, C., Wendt, K., Butterbach-Bahl, K., Wairegi,
- 871 L. and Datta, A.: Soil fertility Managment in Sub-Saharan Africa, in Sustainable Agriculture
- 872 Reviews, edited by E. Lichtfouse, p. 304, Springer International Publishing., 2017.
- 873 Nicholson, S., Some, B., McCollum, J., Nelkin, E., Klotter, D., Berte, Y., Diallo, B., Gaye, I.,
- 874 Kpabeba, G., Ndiaye, O., Noukpozounkou, J., Tanu, M., Thiam, A., Toure, A. and Traore, A.:
- 875 Validation of TRMM and other rainfall estimates with a high-density gauge dataset for West
- 876 Africa. Part II: Validation of TRMM rainfall products, J. Appl. Meteorol., 42, 1355–1368,
- 877 doi:Article, 2003.
- 878 Ossohou, M., Galy-Lacaux, C., Yoboué, V., Hickman, J. E., Gardrat, E., Adon, M., Darras, S.,
- 879 Laouali, D., Akpo, A., Ouafo, M., Diop, B. and Opepa, C.: Trends and seasonal variability of
- 880 atmospheric NO2 and HNO3 concentrations across three major African biomes inferred
- from long-term series of ground-based and satellite measurements, Atmos. Environ., 207,
 148–166, 2019.
- 883 Pamela A. Matson, McDowell, W. H., Townsend, A. R. and Vitousek, P. M.: The globalization
- 884 of N deposition: ecosystem consequences in tropical environments, Biogeochemistry, 46,
- 885 67-83 [online] Available from: http://academic.engr.arizona.edu/HWR/Brooks/GC572-
- 886 2004/readings/matson.pdf, 1999.
- 887 Pommier, M., Law, K. S., Clerbaux, C., Turquety, S., Hurtmans, D., Hadji-Lazaro, J., Coheur, P.
- 888 F., Schlager, H., Ancellet, G., Paris, J. D., Nédlec, P., Diskin, G. S., Podolske, J. R., Holloway, J. S.
- and Bernath, P.: IASI carbon monoxide validation over the Arctic during POLARCAT spring
- and summer campaigns, Atmos. Chem. Phys., 10(21), 10655–10678, doi:10.5194/acp-10-
- 891 10655-2010, 2010.
- 892 Pope, A., Burnett, R., Thun, M., EE, C., D, K., I, K. and GD, T.: Long-term Exposure to Fine
- 893 Particulate Air Pollution, J. Am. Med. Assoc., 287(9), 1132–1141,
- 894 doi:10.1001/jama.287.9.1132, 2002.
- 895 Raleigh, C., Linke, A., Hegre, H. and Karlsen, J.: Introducing ACLED: An Armed Conflict
- 896 Location and Event Dataset: Special Data Feature, J. Peace Res., 47(5), 651–660,

- 897 doi:https://doi.org/10.1177/0022343310378914, 2010.
- 898 Ramo, R., Roteta, E., Bistinas, I., van Wees, D., Bastarrika, A., Chuvieco, E. and van der Werf,
- 899 G. R.: African burned area and fire carbon emissions are strongly impacted by small fires
- 900 undetected by coarse resolution satellite data, Proc. Natl. Acad. Sci. U. S. A., 118(9), 1–7,
- 901 doi:10.1073/pnas.2011160118, 2021.
- 902 Robinson, T. P., Wint, G. R. W., Conchedda, G., Van Boeckel, T. P., Ercoli, V., Palamara, E.,
- 903 Cinardi, G., D'Aietti, L., Hay, S. I. and Gilbert, M.: Mapping the Global Distribution of
- 904 Livestock, PLoS One, 9(5), e96084, doi:10.1371/journal.pone.0096084, 2014.
- 905 Roteta, E., Bastarrika, A., Padilla, M., Storm, T. and Chuvieco, E.: Development of a Sentinel-2
- 906 burned area algorithm: Generation of a small fire database for sub-Saharan Africa, Remote
- 907 Sens. Environ., 222(November 2018), 1–17, doi:10.1016/j.rse.2018.12.011, 2019.
- 908 Rufino, M. C., Rowe, E. C., Delve, R. J. and Giller, K. E.: Nitrogen cycling efficiencies through
- 909 resource-poor African crop-livestock systems, Agric. Ecosyst. Environ., 112(4), 261–282,
- 910 doi:10.1016/j.agee.2005.08.028, 2006.
- 911 Sauvage, B., Gheusi, F., Thouret, V., Cammas, J. P., Duron, J., Escobar, J., Mari, C., Mascart, P.
- 912 and Pont, V.: Medium-range mid-tropospheric transport of ozone and precursors over
- 913 Africa: Two numerical case studies in dry and wet seasons, Atmos. Chem. Phys., 7(20),
- 914 5357-5370, doi:10.5194/acp-7-5357-2007, 2007.
- 915 Stevens, C. J., David, T. I. and Storkey, J.: Atmospheric nitrogen deposition in terrestrial
- 916 ecosystems: Its impact on plant communities and consequences across trophic levels,
- 917 Funct. Ecol., 32(7), 1757–1769, doi:10.1111/1365-2435.13063, 2018.
- 918 Sutton, M. A., Reis, S., Riddick, S. N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S.,
- 919 Braban, C. F., Vieno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall,
- 920 T. D., Milford, C., Flechard, C. R., Loubet, B., Massad, R., Cellier, P., Personne, E., Coheur, P. F.,
- 921 Clarisse, L., Van Damme, M., Ngadi, Y., Clerbaux, C., Skøth, C. A., Geels, C., Hertel, O., Wichink
- 922 Kruit, R. J., Pinder, R. W., Bash, J. O., Walker, J. T., Simpson, D., Horváth, L., Misselbrook, T. H.,
- 923 Bleeker, A., Dentener, F. and de Vries, W.: Towards a climate-dependent paradigm of
- 924 ammonia emission and deposition, Philos. Trans. R. Soc. B Biol. Sci., 368(1621), 20130166-
- 925 20130166, doi:10.1098/rstb.2013.0166, 2013.
- 926 Tian, D. and Niu, S.: A global analysis of soil acidification caused by nitrogen addition,
- 927 Environ. Res. Lett., 10(2), 024019, doi:10.1088/1748-9326/10/2/024019, 2015.

- 928 USDA Agricultural Air Quality Task Force: Ammonia Emissions : What To Know Before You
- 929 Regulate, Washington, DC. [online] Available from:
- 930 http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/air/taskforce/?cid=stelprdb1
- 931 268645, 2014.
- 932 Vitousek, P., Naylor, R., Crews, T., David, M., Drinkwater, L., Holland, E., Johnes, P.,
- 933 Katzenberger, J., Martinelli, L. A., Matson, P. A., Nziguheba, G., Ojima, D., Palm, C. A.,
- 934 Robertson, G., Sanchez, P., Townsend, A. and Zhang, F.: Nutrient Imbalances in Agricultural
- 935 Development, Science (80-.)., 324, 1519–1520, 2009.
- 936 Di Vittorio, C. A. and Georgakakos, A. P.: Land cover classification and wetland inundation
- 937 mapping using MODIS, Remote Sens. Environ., 204(May 2017), 1–17,
- 938 doi:10.1016/j.rse.2017.11.001, 2018.
- 939 Vrieling, A., de Beurs, K. M. and Brown, M. E.: Variability of African farming systems from
- 940 phenological analysis of NDVI time series, Clim. Change, 109(3-4), 455-477,
- 941 doi:10.1007/s10584-011-0049-1, 2011.
- 942 De Wachter, E., Barret, B., Le Flochmoën, E., Pavelin, E., Matricardi, M., Clerbaux, C., Hadji-
- 943 Lazaro, J., George, M., Hurtmans, D., Coheur, P. F., Nedelec, P. and Cammas, J. P.: Retrieval of
- 944 MetOp-A/IASI CO profiles and validation with MOZAIC data, Atmos. Meas. Tech., 5(11),
- 945 2843-2857, doi:10.5194/amt-5-2843-2012, 2012.
- 946 Warner, J. X., Dickerson, R. R., Wei, Z., Strow, L. L., Wang, Y. and Liang, Q.: Increased
- 947 atmospheric ammonia over the world's major agricultural areas detected from space,
- 948 Geophys. Res. Lett., 44(6), 2875–2884, doi:10.1002/2016GL072305, 2017.
- 949 Whitburn, S., Van Damme, M., Kaiser, J. W., Van Der Werf, G. R., Turquety, S., Hurtmans, D.,
- 950 Clarisse, L., Clerbaux, C. and Coheur, P. F.: Ammonia emissions in tropical biomass burning
- 951 regions: Comparison between satellite-derived emissions and bottom-up fire inventories,
- 952 Atmos. Environ., 121, 42–54, doi:10.1016/j.atmosenv.2015.03.015, 2015.
- 953 World Bank: World Bank Open Data, World Bank Open Data [online] Available from:
- 954 https://www.data.worldbank.org (Accessed 2 February 2019), n.d.
- 955 Yegbemey, R. N., Kabir, H., Awoye, O. H. R., Yabi, J. A. and Paraïso, A. A.: Managing the
- 956 agricultural calendar as coping mechanism to climate variability: A case study of maize
- 957 farming in northern Benin, West Africa, Clim. Risk Manag., 3, 13–23,
- 958 doi:10.1016/j.crm.2014.04.001, 2014.

- 959 Yokelson, R. J., Christian, T. J., Karl, T. G. and Guenther, A.: The tropical forest and fire
- 960 emissions experiment: laboratory fire measurements and synthesis of campaign, Rev. Int.
- 961 Acupunt., 8, 3509–3527, doi:10.1016/s1887-8369(09)71579-0, 2008.
- 962 Zheng, B., Chevallier, F., Ciais, P., Yin, Y. and Wang, Y.: On the Role of the Flaming to
- 963 Smoldering Transition in the Seasonal Cycle of African Fire Emissions, Geophys. Res. Lett.,
- 964 45(21), 11,998-12,007, doi:10.1029/2018GL079092, 2018.