



Changes in source contributions of particulate matter during COVID-19 pandemic in the Yangtze River Delta, China

Jinlong Ma¹, Juanyong Shen², Peng Wang³, Shengqiang Zhu¹, Yu Wang¹, Pengfei Wang⁴, Gehui Wang^{5,6}, Jianmin Chen^{1,6*}, Hongliang Zhang^{1,6*}

⁵ ¹Department of Environmental Science and Engineering, Fudan University, Shanghai 200438, China ²School of Environmental Science and Engineering, Shanghai Jiao Tong University, Shanghai 200240, China ³Department of Civil and Environmental Engineering, Hong Kong Polytechnic University, Hong Kong 99907, China ⁴Department of Civil and Environmental Engineering, Louisiana State University, Baton Rouge, LA 70803, USA ⁵Key Lab of Geographic Information Science of the Ministry of Education, School of Geographic Sciences, East China Normal

University, Shanghai 200241, China
 ⁶Institute of Eco-Chongming (IEC), Shanghai 200062, China

Correspondence to: Hongliang Zhang (zhanghl@fudan.edu.cn); Jianmin Chen (jmchen@fudan.edu.cn)

Abstract. Within a short time after the outbreak of coronavirus disease 2019 (COVID-19) in Wuhan, Hubei, the Chinese government took the nationwide lockdown to prevent the spread of the pandemic. The quarantine measures have significantly

- 15 decreased the anthropogenic activities, and thus improving the air quality. To study the impacts caused by the lockdown on specific source sectors and regions in the Yangtze River Delta (YRD), the Community Multiscale Air Quality (CMAQ) model was used to investigate the changes in source contributions to fine particulate matter (PM_{2.5}) from January 23 to February 28, 2020, based on different emission control cases. Compared to Case 1 (without emission reductions), the total PM_{2.5} mass for Case 2 (with emission reductions) decreased by larger than 20% over the entire YRD and the reduction ratios of its components
- 20 were 15%, 16%, 20%, 43%, 34%, and 35% in primary organic aerosol (POA), elemental carbon (EC), sulfate, nitrate, ammonium, and secondary organic aerosol (SOA), respectively. The source appointment results showed that PM_{2.5} concentrations from transportation decreased by 40% while from residential and power decreased by less than 10% due to the lockdown. Although all sources decreased, the relative contribution changed differently. Contributions of residential increased by more than 10% to 35%, while that of industry decreased to 33%. Considering the regional transport, the total PM_{2.5} mass
- 25 of all regions decreased 20-30% in the YRD with the largest decreased value of 5.0 μg m⁻³ in Henan, Hebei, Beijing and Tianjin (Ha-BTH). In Shanghai, the lower contributions from local emissions and regional transmission (mainly Shandong and Ha-BTH) led to the reduced PM_{2.5}. This study suggested adjustments of control measures for various sources and regions. Also, it was necessary to strengthen the regional collaboration at a large scale to improve the air quality in the YRD.

1 Introduction

30 Fine particulate matter (PM_{2.5}, an aerodynamic diameter of fewer than 2.5 μm) has been a great concern in China since 2013 due to its high levels and related health risks (Lelieveld et al., 2015;Huang et al., 2014;He and Christakos, 2018;Shang et al.,





2018;Song et al., 2017;Song et al., 2016;Yan et al., 2018;Du and Li, 2016;Liu et al., 2016;Shen et al., 2020a). To improve air quality, China has promulgated stringent emission control plans such as the Air Pollution Prevention and Control Action Plan and PM_{2.5} concentrations have been reduced significantly in different regions (Zheng et al., 2018;Cai et al., 2017;Zhang et al.,

- 2016;Zheng et al., 2017). The Yangtze River Delta (YRD), one of the largest economic centers, PM_{2.5} concentrations were reduced by 34.3% from 2013 to 2017 due to significant efforts (China, 2018). However, PM_{2.5} concentrations are still much higher than the recommended annual mean criteria of 10 μg m⁻³ by the World Health Organization (WHO). The significant reductions in emissions lead to changes in the local and regional transport contributions of key pollutants. Consequently, the air quality strategies need further improvement according to the source apportionment results.
- 40 PM_{2.5} is a complete mixture of primary PM components (PPM) and secondary formed components, and its source apportionment based on quantifying the contributions of different sources to all the components. Statistical methods using observed PM_{2.5} composition information can only resolve the contributions of source sectors to PPM. Source-oriented chemical transport models (CTMs) are capable of investigating the contributions of both source sectors and regional transports to both PPM and secondary components (Wang et al., 2014; Ying et al., 2014; Wang et al., 2013; Yang et al., 2020). For instance, Hu
- 45 et al. (2015) reported that local emissions account for the highest fraction of PPM compared to the regional transport in Shanghai. Zhang et al. (2012) showed that the power sector (~30%) was the predominant contributor to sulfate, a component of secondary inorganic aerosol (SIA), and remaining contributions are from industrial and residential sectors in Shanghai. Liu et al. (2020) reported that the industry sector was the major secondary organic aerosol (SOA) emissions source, and additionally both regional transport and local emissions were critical to Shanghai. With source contributions changed, the
- 50 information provided by these studies is not suitable for the further reduction of PM_{2.5} in the YRD. Therefore, the updated source appointment information is needed to support the further reduction policy. To prevent the spread of the COVID-19 pandemic, the unprecedented nationwide lockdown has been implemented to limit

anthropogenic activities from January 2020. As a result, anthropogenic emissions decreased drastically, especially in the transportation and industry sectors (Wang et al., 2020a). As a natural experiment with high research values, this provides a

- valuable opportunity to understand pollution changes with extremely strict measures. Studies have reported significant decreases of $PM_{2.5}$ in the YRD based on absolute concentrations (Chen et al., 2020;Li et al., 2020;Chauhan and Singh, 2020;Yuan et al., 2020). However, it is not clear how the contributions of local sources and regional transport changed, and cannot be used to design control strategies. Thus, it is critical to investigate changes in source sectors and regions during the COVID-19 pandemic.
- 60 In this study, a source-oriented version of the Community Multiscale Air Quality (CMAQ) model was used to determine the contributions of source sectors and regional transport to $PM_{2.5}$ in the YRD from January 23 to February 28. The impacts of quarantine measures were estimated by comparing the contributions before and after January 23, the start point of the lockdown. The results offer a deep insight into $PM_{2.5}$ source changes and help develop suitable emission control measures.





2 Methodology

65 2.1 Model description

The SAPRC-11 photochemical mechanism and AERO6 aerosol module were applied in the CMAQ v5.0.2 to separately quantify source contributions to PPM and SIA (Carter and Heo, 2013;Zhang et al., 2015). The CMAQ model used in this study was modified to add additional non-reactive tracers of PPM from various source sectors and regions (Hu et al., 2015). The emission rates of these tracers only account for 0.001% of total PPM emission rates in each grid cell so that they will not have

- an impact on the atmospheric process. The concentrations of tracers from a given source or region were then estimated by multiplying 10^5 to represent the concentrations of PPM from that source or region. The concentrations of components in PPM were calculated based on the ratio of each component to total PPM from sources or regions. Details are discussed in Hu et al. (2015).
- The contributions of source sectors and regions to SIA were quantified by tagging reactive tracers. Precisely, both the components of SIA and their precursors from diverse source types and regions were tracked separately by adding labels on
- NO_x, SO₂, and NH₃ through the atmospheric process (Shi et al., 2017). In this study, contributions from different emission sectors including residential, industry, transportation, power, and agriculture, and those from source regions including Jiangsu, Shanghai, Zhejiang, Anhui, Ha-BTH (Henan, Hebei, Beijing, and Tianjin), Shandong, HnHb (Hunan and Hubei) and other provinces were tracked (Fig. S1 and Table S1). It should be noted that considerable uncertainties due to emission of precursors
- and lack of formation pathways, contributions of SOA to total $PM_{2.5}$ was small (less than 10%) and SOA sources were not tracked in this study. More information of SOA source apportionment was discussed in Wang et al. (2018).

2.2 Model application

Two nested domains were used to simulate pollution changes during the COVID-19 pandemic from January 5 to February 28, 2020. As shown in Fig. S1, China and its surrounding areas were covered in the outer 36 km domain (197×127 grid cells),

- 85 and the YRD was covered by the inner 12km domain (97 × 88 grid cells). The first five-day simulation was removed to minimize the effect of initial conditions. The boundary conditions used in the 12 km domain were offered by the 36 km simulations. Meteorology inputs were generated by the Weather Research and Forecasting (WRF) model v3.6.1. The boundary and initial conditions for WRF were from the National Centers for Environmental Prediction (NCEP) Final (FNL) Operational Model Global Tropospheric Analyses dataset (available at http://rda.ucar.edu/datasets/ds083.2/). The anthropogenic emissions
- 90 in China, based on the Multi-resolution Emission Inventory for China (MEIC) (<u>http://www.meicmodel.org</u>), include industry, power, agriculture, residential and transportation. The emissions from other countries were obtained from the Emissions Database for Global Atmospheric Research (EDGAR) v4.3 (<u>http://edgar.jrc.ec.europa.eu/overview.php?v=431</u>). Biogenic emissions were generated using the Model for Emissions of Gases and Aerosols from Nature (MEGAN) v2.1 (Guenther et al., 2012).
- 95 Two cases were simulated in this study (Table 1). The base case (Case 1) used the original inventory. In Case 2, the emissions





of carbon monoxide (CO), nitric oxide (NO_x), sulfur dioxide (SO₂), volatile organic compounds (VOC), and PM were decreased during the COVID-19 period since January 23 with provincial specific factors from Huang et al. (2020). The differences between the cases represent the changes in sources and regions.

3 Results and discussion

3.1 Model performance 100

105

3.1.1 WRF evaluation

Since air quality simulations would be influenced by meteorology difference, it is critical to validate WRF performance before simulating source appointment (Zhang et al., 2015). The model performance of meteorological parameters temperature at 2 m above the ground surface (T2), wind speed (WSPD), wind direction (WD) and relative humidity (RH) in the COVID-19 period could be found in Table S2. The statistical values of mean prediction (PRE); mean observation (OBS); mean bias (MB); gross error (GE); and root mean square error (RMSE) have been calculated and the calculation formulas were listed in Table S4. The WRF model predicted slightly higher T2 than observations in the two periods. The MB values of T2 before and after the lockdown were both 1.6, while the GE value of T2 before the lockdown period was slightly larger than the recommended criterion based on Emery et al. (2001). Except for the MB values of WSPD, both GE (1.3 and 1.6) and RMSE (1.7 and 2.0) were met the benchmarks during the two periods. The MB (1.8) and GE (29.2) values of WD were all within the benchmarks 110

- after the lockdown, but the GE value of WD before the lockdown was slightly higher than the benchmark. The simulated RH was underestimated with the MB values of -2.4 and -5.6 during the two periods. The hourly comparisons of T2, WSPD and RH were shown in Fig. S9 based on Wang et al. (2020b, under review) also indicated good model performance. Compared to previous studies (Chen et al., 2019;Liu et al., 2020), the meteorology predictions in this study were robust to drive air quality
- 115 simulation. Generally, the simulation of meteorological conditions in this study was acceptable.

3.1.2 CMAQ evaluation

The model performance of O₃, NO₂, SO₂, PM_{2.5} and PM₁₀ mass in the YRD during the COVID-19 pandemic has been described in Table S2 of a previous study (Wang et al., 2020b, under review). During the whole simulated period, the predicted PM_{2.5} and O_3 were slightly higher than observations, but the model performance was within the criteria for PM_{2.5} (mean fractional

- 120 bias (MFB) $\leq \pm 60\%$ and mean fractional error (MFE) $\leq 75\%$, suggested by Boylan and Russell (2006)) and for O₃ (MFB $\leq \pm 15\%$ and MFE≤30%, suggested by U.S.EPA (2007)). Figure 1 showed predicted and observed daily PM2.5 and its components averaged over the YRD and at three major cities based on Case 2. Generally, the temporal trends of PM_{2.5} mass before and during the lockdown were successfully captured by the model simulations. The MFB and MFE values of PM_{2.5} mass were 0.14-0.41 and 0.38-0.57, which were all within the criteria. In Shanghai, the simulations missed the PM_{2.5} episodes from
- 125 January 11 to 13, but the overall performance was good. In addition, observed SIA (including sulfate, nitrate and ammonium)





from January 08 to February 10 2020 in Shanghai reported by Chen et al. (2020) were used to evaluate the model performance, as shown in Fig. S3. The daily simulated trends of SIA generally agreed with the observations, although the model slightly overpredicted SIA concentrations with the MFB values of 0.19-0.37 and the MFE values of 0.41-0.68 (Table S3). The overestimation of nitrate has been reported in the previous studies (Chang et al., 2018;Shen et al., 2020b;Choi et al., 2019) and

- 130 the possible reason was the lack of chlorine heterogeneous chemistry in the model (Qiu et al., 2019). Compared to the predictions before the lockdown, the MFB and MFE values of $PM_{2.5}$ and its components during the lockdown were slightly lower than the values before the lockdown (Table S4), indicating the model performance during the lockdown was better. This discrepancy could be caused by the uncertainties in the emissions and meteorology (Ying et al., 2014). The model performance of the WRF model was similar during the two periods (Table S2). The 2016 MEIC emission was used for the year 2020, which
- 135 may overestimate the anthropogenic emissions and thus the $PM_{2.5}$ concentration in the before lockdown period. However, the emission adjustments based on Huang et al. (2020) during lockdown may be closer to the real condition, leading to better model performance. Despite these uncertainties, the model results were acceptable for source apportionment studies.

3.2 Changes of PM_{2.5} and components during the lockdown

- Figure 2 shows the predicted total PM_{2.5} and its components in the YRD during the COVID-19 lockdown. In both cases, PM_{2.5} and its components showed similar spatial distributions with the highest concentrations in the northwest and lower concentrations in the southeast. Substantial PM_{2.5} was observed in north Anhui, similar patterns were found in elemental carbon (EC) and primary organic aerosol (POA), indicating similar sources and large contributions. For Case 2, averaged PM_{2.5} concentrations mainly decreased in south and northeast YRD due to the lockdown and all major components decreased in varying degrees. For EC and POA, similar decreases of 15% were observed in Anhui compared to Case 1. More important
- 145 decreases were found in other regions especially in Zhejiang (up to 25%). SIA had the maximum decrease in Anhui (30-40%), which was related to sharp drops of concentrations in nitrate and ammonium with decreasing ratios of 40-50% and 30-40% (Fig. S4), respectively. On the contrary, the reductions of sulfate in Shanghai were higher than other regions in the YRD, mainly due to reduced more SO₂ from industries during the lockdown based on Huang et al. (2020). Except for central and northwest YRD, SOA decreased significantly (35-40%) also due to the reductions of industrial activities, which was an
- 150 important contributor to SOA (Liu et al., 2020). Figure 3 shows the contributions of components to PM_{2.5} in the YRD and three major cities during the lockdown. For Case 2, over the entire YRD, the reductions in POA, EC, sulfate, nitrate, ammonium, and SOA was 2.4, 0.8, 2.1, 7.8, 2.9, and 0.9 μg m⁻³ with a total of 17.0 μg m⁻³ decrease in PM_{2.5}. The most significant decrease was found in nitrate with the highest decrease rate of over 40%. In selected cities, PM_{2.5} was decreased by 15.1, 14.8 and 16.8 μg m⁻³ in Shanghai, Hangzhou and Nanjing,
- 155 respectively, with the largest decrease ratio of 27% in Hangzhou. Secondary components (SIA + SOA) dropped more significantly than primary components, especially for nitrate (35-45%) due to the severe decrease of NO_x from transportation. This also indicates that atmospheric reactions and meteorological conditions were important during the pandemic period. In addition to nitrate, a sharp decrease was observed in ammonium due to the decrease of both nitrate and sulfate (Erisman and Context).





Schaap, 2004). SIA concentrations contributed the most to PM_{2.5} in selected cities with the highest values of 26.5 μg m⁻³ in Nanjing. Furthermore, the largest contributor to SIA was nitrate in the YRD, Hangzhou, and Nanjing during the lockdown, while sulfate became the dominant contributor in Shanghai and accounted for 22% of total PM_{2.5}, similar to the result in Chen et al. (2020). This shows the changes in PM_{2.5}, but sector and region contributions are unclear to help design regional control policies. Below shows the source appointment and regional transport of PM_{2.5} and changes caused by lockdown measures.

3.3 Source sector contributions to PM_{2.5}

- 165 Figure 4 shows the contributions of different source sectors to $PM_{2.5}$ in the YRD during the lockdown. Source appointments of SIA and PPM in two cases are illustrated in Fig. S5 and Fig. S7, respectively. The agricultural source of PPM is not shown due to minor contributions. Generally, residential activities were the most significant contributor to $PM_{2.5}$ with the highest value of 45.0 µg m⁻³, mainly due to the large contribution to PPM (Fig. S7). The contribution in Shanghai was ~20.0 µg m⁻³ and decreased to 15.0 µg m⁻³ during the lockdown. The overall decrease was less than 10% in the middle YRD and less than
- 170 15% in the rest regions. Contributions from transportation decreased the most due to the lockdown from larger than 10.0 μ g m⁻³ in Case 1 to less than 7.5 μ g m⁻³ in most areas. This was shown in SIA as well (Fig. S5), over 40% decreases were found in the YRD except for southeast, with the maximum decrease value of ~7.0 μ g m⁻³. The industry contributed the most to PM_{2.5} values in industrial cities such as Suzhou and Hefei, positions as shown in Fig. S1, which decreased significantly by ~10.0 μ g m⁻³ from >30.0 μ g m⁻³ to ~20.0 μ g m⁻³ in Case 2. PM_{2.5} from the power sector was decreased by less than 5% to less than 6 μ g
- m⁻³ in most areas due to reduced emissions of SO₂ and associated sulfate (Fig. S6). PM_{2.5} from agriculture also decreased by the lockdown with the largest decrease of 5.0 μg m⁻³ in northwest YRD.
 Figure 5 shows the changes in contributions of sources to PM_{2.5} in the YRD, Shanghai, Hangzhou, and Nanjing caused by the lockdown. Overall in the YRD, residential and industrial sources were major sources with contributions of 35% and 33% with decreases of less than 20%. Transportation, power and agriculture sources contributed similar to PM_{2.5} but with different
- 180 changing ratios of 40%, 6%, and 17%, respectively. Although all sources decreased, the relative contribution did not remain unchanged. The contribution ratio of transportation decreased by 27% due to the decrease in both primary emission and secondary formation, as shown in Fig. S8 and Fig. S9. The contribution ratios of residential and power increased by more than 10% while industry and agriculture showed slight changes. In large cities, industrial sources were leading with 5.0-10.0 µg m⁻ ³ higher contribution than residential sources, while other sources were similar to the YRD averages. In Shanghai, the
- 185 contributions of power and agriculture showed insignificant changes, that of the industry changed by ~20% and transportation decreased by more than 30%. The relative contribution of transportation decreased by more than 15%, while that of power and agriculture increased by 14% and 9%, respectively. In Hangzhou and Nanjing, the trends were similar except contributions and changes of all sources were larger in Nanjing. Due to the lockdown measures, contributions of different sources decreased but their relative contribution changed differently, implying that adjustment of control measures for various sources is needed.

190 3.4 Regional contributions to PM_{2.5}





Figure 6 illustrates the distribution of PM_{2.5} contributed by emissions from different regions for two cases in the YRD during the lockdown. Regional transmission of SIA and PPM was shown in Fig. S10 and Fig. S11, respectively. It is clear that the regional distributions of each source were the same in both cases and Case 2 had lower values and narrower distributions. Contributions of local emissions from Jiangsu, Shanghai, Zhejiang, and Anhui generally peaked near the source regions with less than 5.0 µg m⁻³ transported to other areas. Emissions from HnHb were barely transported to the central YRD area. Shandong and Ha-BTH emissions could be transported further due to north winds, as shown in Fig. S2, with ~10.0 µg m⁻³ and ~5.0 µg m⁻³ contributions to north YRD, respectively. It indicates that the regional transports among provinces were notable, which is consistent with Du et al. (2017). Other regions also had small contributions to YRD, but the contributions decreased significantly during the lockdown. The limitation of commercial activities and traffics caused by the pandemic lockdown significantly suppressed the emission and dispersion of PM_{2.5}. Compared to Case 1, contributions from Jiangsu, Anhui, Shandong, and Ha-BTH in Case 2 decreased by 20-30%. More significant decreases of 30-40% were found in Shanghai, Zhejiang, and HnHb. The largest decrease of ~18.0 µg m⁻³ was observed in Hubei, the center of the COVID-19 pandemic in

China due to more strict lockdown measures. Figure S9 shows that after the implementation of quarantine measures, the SIA contributions decreased by more than 30% among each region and HnHb decreased by 51% to less than 10.0 µg m⁻³. Figure
S10 shows the narrower distributions and smaller decreases of PPM in Case 2 compared with SIA, with a decrease of less than 30% in all selected regions.

Figure 7 illustrates $PM_{2.5}$ contributed by eight regions averagely in the YRD and Shanghai. In the YRD, averaged contributions due to local emissions from Jiangsu, Shanghai, Zhejiang and Anhui were 6.8, 0.8, 1.5, and 6.3 µg m⁻³ during the lockdown period, while the contribution of outside YRD areas from HnHb, Shandong, Ha-BTH and Other were 5.0, 9.1, 14.4, and 8.2

- 210 μg m⁻³, respectively. The contributions of all regions decreased due to the COVID-19 lockdown with the averaged decreasing rate of 20-30% with the largest decrease rate of 33% in HnHb and the least decrease of 21% in Jiangsu. In addition to the absolute contributions, Figure 7(b) also shows the relative contribution of different regions. Ha-BTH had the largest contribution of ~30%, followed by Shandong and Other. Jiangsu and Anhui were the largest local contributors with ~12% each. It is clear that long-range transport played an important role in PM_{2.5} pollution in the YRD with more than 70%
- 215 contribution. Due to the COVID-19, although the absolute contributions decreased universally, their relative contributions did not. The importance of Jiangsu and Shandong increased by ~5%, while that of Shanghai, Zhejiang and HnHb decreased with the largest rate of 12% in HnHb. The results show that although all regions reduced their concentrations to the YRD, the relative contribution changed. In the future, regional cooperative control is needed for the YRD and strategies should be adjusted according to changes in contributions.
- 220 At the city level, local emission was the major contributor with contributions of 10.0 μg m⁻³ within the YRD to Shanghai, the largest city in the YRD (Fig. 7(c)). Jiangsu contributed 16% to Shanghai while Zhejiang and Anhui had few effects. Outside the YRD, Shandong had the largest contribution (11.5 μg m⁻³), followed by Ha-BTH and Other areas. In total, contributions from neighbor provinces (<10.0 μg m⁻³) were much smaller than long-range transport from outside YRD (23.7 μg m⁻³). Prevailing northerly winds were a key factor in this (Fig. S2). The lockdown decreased the contributions from all regions by





225 20-45%, with the largest decrease from HnHb. The contribution order of different regions was unchanged but their relative contributions changed. The relative contributions of local emissions from Shanghai decreased by ~10%, while that of Shandong and Jiangsu increased by ~10%. The relative contribution of HnHb decreased by more than 20%, although the absolute changes were small.

The quarantine measures during COVID-19 lockdown reduced emissions from transportation and industrials, and the total

230 emissions for different areas changed differently. Although $PM_{2.5}$ concentrations decreased in the whole YRD, the contributions of source sectors and regions changed differently. It highlights the need for regional cooperative emission reduction and adjusting control strategies when significant reductions are achieved.

4 Conclusion

- A source-oriented CMAQ model investigated the changes in contributions of source sectors and regions to $PM_{2.5}$ during the 235 COVID-19 lockdown in the YRD. Total $PM_{2.5}$ mass decreased by larger than 20% across the YRD due to decreases of 30-40% and 10-20% in secondary and primary components, respectively. The results of the source appointment showed that the residential and industrial sources were the major sources with contributions of 35% (18.0 µg m⁻³) and 33% (17.1 µg m⁻³) and decreased by less than 20% due to the lockdown. Contributions from transportation decreased by 40%, which was the most significant decrease, while the decrease in power was less than 10%. The relative contribution of sources changed due to
- 240 differences in source decreases. The relative contribution of transportation decreased by more than 25%, while that of residential and power increased by more than 10%, suggesting that further abatement policy should adjust control measures for various sources. Contributions from regional transport of emission outside YRD were the dominant contributor (more than 70%) to the YRD, and contributions from all regions decreased due to the lockdown. The relative contribution of each region also changed with increases in Jiangsu and Shandong (~10%) but decreases in all other regions. This implies that strengthening
- the regional joint preventions and control of transported pollution from heavily polluted regions could effectively mitigate PM_{2.5} pollution in the YRD.

Author contributions. JM conducted the modelling and led the writing. JS assisted in writing papers. PW, SZ, YW and PW collected data and provided technical supports. GW assisted with data analysis. JC and ZH designed the study, discussed the results, and edited the paper.

250 Competing interests. The authors declare that they have no conflict of interest.

Financial support. This project was funded by Institute of Eco-Chongming (ECNU-IEC-202001).

References





Boylan, J. W., and Russell, A. G.: PM and light extinction model performance metrics, goals, and criteria for three-dimensional air quality models, Atmospheric Environment, 40, 4946-4959, <u>https://doi.org/10.1016/j.atmosenv.2005.09.087</u>, 2006.

255 Cai, S., Wang, Y., Zhao, B., Wang, S., Chang, X., and Hao, J.: The impact of the "Air Pollution Prevention and Control Action Plan" on PM2.5 concentrations in Jing-Jin-Ji region during 2012-2020, Science of the Total Environment, 580, 197-209, 10.1016/j.scitotenv.2016.11.188, 2017.

Carter, W. P. L., and Heo, G.: Development of revised SAPRC aromatics mechanisms, Atmospheric Environment, 77, 404-414, https://doi.org/10.1016/j.atmosenv.2013.05.021, 2013.

- 260 Chang, X., Wang, S., Zhao, B., Cai, S., and Hao, J.: Assessment of inter-city transport of particulate matter in the Beijing–Tianjin–Hebei region, Atmospheric Chemistry and Physics, 18, 4843-4858, 10.5194/acp-18-4843-2018, 2018. Chauhan, A., and Singh, R. P.: Decline in PM2.5 concentrations over major cities around the world associated with COVID-19, Environmental Research, 187, 10.1016/j.envres.2020.109634, 2020.
- Chen, D., Tian, X., Lang, J., Zhou, Y., Li, Y., Guo, X., Wang, W., and Liu, B.: The impact of ship emissions on PM2.5 and the deposition
 of nitrogen and sulfur in Yangtze River Delta, China, Science of The Total Environment, 649, 1609-1619, https://doi.org/10.1016/j.scitotenv.2018.08.313, 2019.

Chen, H., Huo, J., Fu, Q., Duan, Y., Xiao, H., and Chen, J.: Impact of quarantine measures on chemical compositions of PM2.5 during the COVID-19 epidemic in Shanghai, China, Sci Total Environ, 743, 140758, 10.1016/j.scitotenv.2020.140758, 2020.

Air quality targets set by the Action Plan have been fully realized: <u>http://www.gov.cn/xinwen/2018-02/01/content_5262720.htm/</u>, access: 1st Febuary 2018.

Choi, Lee, Woo, Kim, and Lee: Comparison of PM2.5 Chemical Components over East Asia Simulated by the WRF-Chem and WRF/CMAQ Models: On the Models' Prediction Inconsistency, Atmosphere, 10, 10.3390/atmos10100618, 2019.

Du, W., Hong, Y., Xiao, H., Zhang, Y., Chen, Y., Xu, L., Chen, J., and Deng, J.: Chemical Characterization and Source Apportionment of PM2.5 during Spring and Winter in the Yangtze River Delta, China, Aerosol and Air Quality Research, 17, 2165-2180, 10.4209/aaqr.2017.03.0108, 2017.

Du, Y., and Li, T.: Assessment of health-based economic costs linked to fine particulate (PM2.5) pollution: a case study of haze during January 2013 in Beijing, China, Air Quality Atmosphere and Health, 9, 439-445, 10.1007/s11869-015-0387-7, 2016.

Emery, C., Tai, E., and Yarwood, G.: Enhanced meteorological modeling and performance evaluation for two Texas episodes. Report to the Texas Natural Resources Conservation Commission. p.b.E., International Corp, Novato, CA, 2001.

280 Erisman, J. W., and Schaap, M.: The need for ammonia abatement with respect to secondary PM reductions in Europe, Environ Pollut, 129, 159-163, 10.1016/j.envpol.2003.08.042, 2004.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, Geoscientific Model Development, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.

- He, J., and Christakos, G.: Space-time PM2.5 mapping in the severe haze region of Jing-Jin-Ji (China) using a synthetic approach, Environmental Pollution, 240, 319-329, 10.1016/j.envpol.2018.04.092, 2018.
 Hu, J., Wu, L., Zheng, B., Zhang, Q., He, K., Chang, Q., Li, X., Yang, F., Ying, Q., and Zhang, H.: Source contributions and regional transport of primary particulate matter in China, Environ Pollut, 207, 31-42, 10.1016/j.envpol.2015.08.037, 2015.
- Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P.,
 Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J.,
 Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prévôt, A. S. H.: High secondary aerosol contribution to particulate
 pollution during haze events in China, Nature, 514, 218-222, 10.1038/nature13774, 2014.
 Huang, X., Ding, A., Gao, J., Zheng, B., Zhou, D., Qi, X., Tang, R., Wang, J., Ren, C., Nie, W., Chi, X., Xu, Z., Chen, L., Li, Y., Che, F.,
- Pang, N., Wang, H., Tong, D., Qin, W., Cheng, W., Liu, W., Fu, Q., Liu, B., Chai, F., Davis, S. J., Zhang, Q., and He, K.: Enhanced secondary
 pollution offset reduction of primary emissions during COVID-19 lockdown in China, National Science Review, 10.1093/nsr/nwaa137, 2020

Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, Nature, 525, 367-+, 10.1038/nature15371, 2015.

Li, L., Li, Q., Huang, L., Wang, Q., Zhu, A., Xu, J., Liu, Z., Li, H., Shi, L., Li, R., Azari, M., Wang, Y., Zhang, X., Liu, Z., Zhu, Y., Zhang, 300
K., Xue, S., Ooi, M. C. G., Zhang, D., and Chan, A.: Air quality changes during the COVID-19 lockdown over the Yangtze River Delta Region: An insight into the impact of human activity pattern changes on air pollution variation, Sci Total Environ, 732, 139282, 10.1016/j.scitotenv.2020.139282, 2020.

Liu, J., Shen, J., Cheng, Z., Wang, P., Ying, Q., Zhao, Q., Zhang, Y., Zhao, Y., and Fu, Q.: Source apportionment and regional transport of anthropogenic secondary organic aerosol during winter pollution periods in the Yangtze River Delta, China, Sci Total Environ, 710, 135620, 10.1016/j.scitotenv.2019.135620, 2020.

Liu, Q., Baumgartner, J., Zhang, Y., and Schauer, J. J.: Source apportionment of Beijing air pollution during a severe winter haze event and associated pro-inflammatory responses in lung epithelial cells, Atmospheric Environment, 126, 28-35, 10.1016/j.atmosenv.2015.11.031, 2016.





- Qiu, X., Ying, Q., Wang, S., Duan, L., Zhao, J., Xing, J., Ding, D., Sun, Y., Liu, B., Shi, A., Yan, X., Xu, Q., and Hao, J.: Modeling the impact of heterogeneous reactions of chlorine on summertime nitrate formation in Beijing, China, Atmospheric Chemistry and Physics, 19, 6737-6747, 10.5194/acp-19-6737-2019, 2019.
 Shang, X., Zhang, K., Mang, F., Wang, S., Lee, M., Suh, L. Kim, D., Jeon, K., Park, H., Wang, Y., and Zhao, Y.: Characteristics and source.
 - Shang, X., Zhang, K., Meng, F., Wang, S., Lee, M., Suh, I., Kim, D., Jeon, K., Park, H., Wang, X., and Zhao, Y.: Characteristics and source apportionment of fine haze aerosol in Beijing during the winter of 2013, Atmospheric Chemistry and Physics, 18, 2573-2584, 10.5194/acp-18-2573-2018, 2018.
- 315 Shen, J., Zhao, Q., Cheng, Z., Huo, J., Zhu, W., Zhang, Y., Duan, Y., Wang, X., Antony Chen, L. W., and Fu, Q.: Evolution of source contributions during heavy fine particulate matter (PM2.5) pollution episodes in eastern China through online measurements, Atmospheric Environment, 232, 10.1016/j.atmosenv.2020.117569, 2020a.
 Shen, L. Zhao, O., Cheng, Z., Wang, P., Ving, O., Lin, L. Duen, Y., and Fu, O.: Insights into source origins and formation mechanisms of

Shen, J., Zhao, Q., Cheng, Z., Wang, P., Ying, Q., Liu, J., Duan, Y., and Fu, Q.: Insights into source origins and formation mechanisms of nitrate during winter haze episodes in the Yangtze River Delta, Sci Total Environ, 741, 140187, 10.1016/j.scitotenv.2020.140187, 2020b.

320 Shi, Z., Li, J., Huang, L., Wang, P., Wu, L., Ying, Q., Zhang, H., Lu, L., Liu, X., Liao, H., and Hu, J.: Source apportionment of fine particulate matter in China in 2013 using a source-oriented chemical transport model, Sci Total Environ, 601-602, 1476-1487, 10.1016/j.scitotenv.2017.06.019, 2017.
Sing G. Win, L. Xia, Y. Liu, Y. Liu, Y. Liu, Y. Liu, Y. Dai, O. Liu, P. Wang, Y. a. and Mag. Hu Air.

Song, C., Wu, L., Xie, Y., He, J., Chen, X., Wang, T., Lin, Y., Jin, T., Wang, A., Liu, Y., Dai, Q., Liu, B., Wang, Y.-n., and Mao, H.: Air pollution in China: Status and spatiotemporal variations, Environmental Pollution, 227, 334-347, 10.1016/j.envpol.2017.04.075, 2017.

- 325 Song, Y., Wang, X., Maher, B. A., Li, F., Xu, C., Liu, X., Sun, X., and Zhang, Z.: The spatial-temporal characteristics and health impacts of ambient fine particulate matter in China, Journal of Cleaner Production, 112, 1312-1318, 10.1016/j.jclepro.2015.05.006, 2016. U.S.EPA: Agency, U.S.E.P. (Ed.), Guidance on the Use of Models and Other Analyses for Demonstrating Attainment of Air Quality Goals for Ozone, PM2.5 and Regional Haze. Research Triangle Park, North Carolina., 2007.
- Wang, L. T., Wei, Z., Yang, J., Zhang, Y., Zhang, F. F., Su, J., Meng, C. C., and Zhang, Q.: The 2013 severe haze over the southern Hebei,
 China: model evaluation, source apportionment, and policy implications, Atmospheric Chemistry and Physics Discussions, 13, 28395-28451,
 10.5194/acpd-13-28395-2013, 2013.

Wang, P., Ying, Q., Zhang, H., Hu, J., Lin, Y., and Mao, H.: Source apportionment of secondary organic aerosol in China using a regional source-oriented chemical transport model and two emission inventories, Environmental Pollution, 237, 756-766, https://doi.org/10.1016/j.envpol.2017.10.122, 2018.

- Wang, P., Chen, K., Zhu, S., Wang, P., and Zhang, H.: Severe air pollution events not avoided by reduced anthropogenic activities during COVID-19 outbreak, Resour Conserv Recycl, 158, 104814, 10.1016/j.resconrec.2020.104814, 2020a.
 Wang, Y., Li, L., Chen, C., Huang, C., Huang, H., Feng, J., Wang, S., Wang, H., Zhang, G., Zhou, M., Cheng, P., Wu, M., Sheng, G., Fu, J., Hu, Y., Russell, A. G., and Wumaer, A.: Source apportionment of fine particulate matter during autumn haze episodes in Shanghai, China, Journal of Geophysical Research: Atmospheres, 119, 1903-1914, 10.1002/2013JD019630, 2014.
- 340 Wang, Y., Zhu, S., Ma, J., Wang, P., Wang, P., and Zhang, H.: Enhanced atmospheric oxidation capacity and associated ozone increases during COVID-19 lockdown in the Yangtze River Delta, submitted to the Science of the Total Environment, 2020b. Yan, D., Lei, Y., Shi, Y., Zhu, Q., Li, L., and Zhang, Z.: Evolution of the spatiotemporal pattern of PM2.5 concentrations in China - A case study from the Beijing-Tianjin-Hebei region, Atmospheric Environment, 183, 225-233, 10.1016/j.atmosenv.2018.03.041, 2018.
- Yang, X., Xiao, H., Wu, Q., Wang, L., Guo, Q., Cheng, H., Wang, R., and Tang, Z.: Numerical study of air pollution over a typical basin topography: Source appointment of fine particulate matter during one severe haze in the megacity Xi'an, Sci Total Environ, 708, 135213, 10.1016/j.scitotenv.2019.135213, 2020.

Ying, Q., Wu, L., and Zhang, H.: Local and inter-regional contributions to PM2.5 nitrate and sulfate in China, Atmospheric Environment, 94, 582-592, 10.1016/j.atmosenv.2014.05.078, 2014.

Yuan, Q., Qi, B., Hu, D., Wang, J., Zhang, J., Yang, H., Zhang, S., Liu, L., Xu, L., and Li, W.: Spatiotemporal variations and reduction of
 air pollutants during the COVID-19 pandemic in a megacity of Yangtze River Delta in China, The Science of the total environment, 751,
 141820-141820, 10.1016/j.scitotenv.2020.141820, 2020.

Zhang, H., Li, J., Ying, Q., Yu, J. Z., Wu, D., Cheng, Y., He, K., and Jiang, J.: Source apportionment of PM2.5 nitrate and sulfate in China using a source-oriented chemical transport model, Atmospheric Environment, 62, 228-242, 10.1016/j.atmosenv.2012.08.014, 2012.

- Zhang, H., Wang, Y., Hu, J., Ying, Q., and Hu, X. M.: Relationships between meteorological parameters and criteria air pollutants in three megacities in China, Environ Res, 140, 242-254, 10.1016/j.envres.2015.04.004, 2015.
- Zhang, H., Wang, S., Hao, J., Wang, X., Wang, S., Chai, F., and Li, M.: Air pollution and control action in Beijing, Journal of Cleaner Production, 112, 1519-1527, 10.1016/j.jclepro.2015.04.092, 2016.

Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., and Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, Atmospheric Chemistry and Physics, 18, 14095-14111, 10.5194/acp-18-14095-2018, 2018.

Zheng, Y., Xue, T., Zhang, Q., Geng, G., Tong, D., Li, X., and He, K.: Air quality improvements and health benefits from China's clean air action since 2013, Environmental Research Letters, 12, 10.1088/1748-9326/aa8a32, 2017.





| | Province | CO | NO _x | SO_2 | VOC | PM _{2.5} | BC | OC |
|--------|----------------|-----|-----------------|--------|----------|-------------------|-----|-----|
| Case 1 | All | | | | No chang | es | | |
| Case 2 | Beijing | 22% | 45% | 26% | 45% | 18% | 46% | 8% |
| | Tianjin | 21% | 38% | 20% | 41% | 14% | 22% | 6% |
| | Hebei | 15% | 45% | 16% | 36% | 12% | 17% | 5% |
| | Shanxi | 18% | 40% | 20% | 33% | 16% | 19% | 10% |
| | Inner Mongolia | 14% | 29% | 15% | 34% | 13% | 16% | 6% |
| | Liaoning | 21% | 40% | 28% | 36% | 16% | 28% | 8% |
| | Jilin | 16% | 39% | 23% | 34% | 13% | 18% | 5% |
| | Heilongjiang | 17% | 37% | 27% | 28% | 13% | 15% | 7% |
| | Shanghai | 35% | 48% | 42% | 45% | 34% | 54% | 42% |
| | Jiangsu | 23% | 50% | 26% | 41% | 16% | 35% | 7% |
| | Zhejiang | 41% | 50% | 29% | 45% | 30% | 49% | 20% |
| | Anhui | 14% | 56% | 22% | 31% | 11% | 22% | 4% |
| | Fujian | 29% | 51% | 30% | 42% | 19% | 31% | 7% |
| | Jiangxi | 24% | 53% | 21% | 43% | 19% | 30% | 9% |
| | Shandong | 23% | 50% | 25% | 39% | 19% | 35% | 9% |
| | Henan | 23% | 57% | 22% | 41% | 18% | 35% | 8% |
| | Hubei | 19% | 55% | 23% | 35% | 16% | 23% | 10% |
| | Hunan | 22% | 51% | 25% | 36% | 20% | 24% | 15% |
| | Guangdong | 38% | 50% | 33% | 46% | 27% | 42% | 13% |
| | Guangxi | 24% | 50% | 28% | 39% | 17% | 27% | 5% |
| | Hainan | 24% | 44% | 25% | 36% | 14% | 25% | 4% |
| | Chongqing | 18% | 53% | 32% | 37% | 14% | 20% | 4% |
| | Sichuan | 16% | 50% | 27% | 33% | 9% | 15% | 3% |
| | Guizhou | 24% | 39% | 25% | 30% | 22% | 25% | 20% |
| | Yunnan | 24% | 51% | 25% | 41% | 18% | 21% | 8% |
| | Tibet | 16% | 35% | 15% | 35% | 14% | 14% | 5% |
| | Shaanxi | 19% | 45% | 18% | 34% | 13% | 22% | 5% |
| | Gansu | 13% | 47% | 16% | 29% | 9% | 13% | 3% |
| | Qinghai | 23% | 46% | 22% | 39% | 20% | 20% | 7% |
| | Ningxia | 24% | 36% | 24% | 39% | 20% | 23% | 8% |
| | Xinjiang | 16% | 35% | 15% | 35% | 14% | 14% | 5% |
| | | | | | | | | |

| Table 1. Simulation scenario | s during the COVID-19 | period in this study | based on Huang et al. (2020). |
|------------------------------|-----------------------|----------------------|-------------------------------|
|------------------------------|-----------------------|----------------------|-------------------------------|







Figure 1: Predicted daily PM_{2.5} and its components with observed daily PM_{2.5} in the YRD and three major cities in Case 2 before (shaded area) and during the lockdown period (white area). Units are µg m⁻³.







Figure 2: Spatial distribution of predicted PM_{2.5} total and major components and changes caused by the lockdown measures in the 370 YRD from January 23 to February 28, 2020. EC is elemental carbon, POA is primary organic aerosol. The relative difference is calculated by (Case 2 – Case 1) / Case 1, using the concentration. Note color ranges are different among panels.







Figure 3: Predicted PM2.5 and its major components of Case 2 (red histogram corresponding to left Y-axis) and relative concentration (circle corresponding to right Y-axis) from January 23 to February 28, 2020 in the YRD and Shanghai, Hangzhou, and Nanjing. Relative concentration means the relative change of concentration between Case 1 and Case 2, which is calculated by (Case 2 - Case 1) / Case 1.









Figure 4: Predicted PM_{2.5} from different source sectors of two cases and the relative difference in the YRD from January 23 to February 28, 2020. Note color ranges are different among panels.



380





Figure 5. Concentrations and contributions of different emission sectors to PM2.5 in the YRD and three major cities of Case 2 from January 23 to February 28, 2020. The values of histograms correspond to the left Y-axis and the values of relative changes correspond to the right Y-axis. The relative contribution means the relative change of contribution between Case 1 and Case 2, calculated by (Case 2 - Case 1) / Case 1. The meaning of relative concentration is the same as in Figure 3.







385

Figure 6: Averaged regional contributions of predicted $PM_{2.5}$ in the YRD from 23 to February 28, 2020. Note color ranges are different among panels.







Figure 7: Concentrations and contributions of predicted PM_{2.5} from different regions in the YRD ((a) and (b)) and Shanghai ((c) 390 and (d)) of Case 2 corresponding to left Y-axis and the relative change (corresponding to right Y-axis) from January 23 to February 28, 2020. The meanings of relative concentration and relative contribution are the same as in Figure 3 and Figure 5.