



Long-range transport patterns into the tropical northwest Pacific during the CAMP²Ex aircraft campaign: chemical composition, size distributions, and the impact of convection

Miguel Ricardo A. Hilario^{1,+}, Ewan Crosbie^{2,3}, Michael Shook², Jeffrey S. Reid⁴, Maria Obiminda L. Cambaliza^{1,5}, James Bernard B. Simpas^{1,5}, Luke Ziemba², Joshua P. DiGangi², Glenn S. Diskin², Phu Nguyen⁶, Joseph Turk⁷, Edward Winstead^{2,3}, Claire E. Robinson^{2,3}, Jian Wang⁸, Jiaoshi Zhang⁸, Yang Wang⁹, Subin Yoon¹⁰, James Flynn¹⁰, Sergio L. Alvarez¹⁰, Ali Behrangi^{11,12}, Armin Sorooshian^{11,13,*}

10 ¹ Manila Observatory, Quezon City 1108, Philippines

² NASA Langley Research Center, Hampton, VA, USA

³ Science Systems and Applications, Inc., Hampton, VA, USA

⁴ Marine Meteorology Division, Naval Research Laboratory, Monterey, CA, USA

⁵ Department of Physics, Ateneo de Manila University, Quezon City 1108, Philippines

15 ⁶ Department of Civil & Environmental Engineering, University of California Irvine, Irvine, CA 92697, USA

⁷ NASA Jet Propulsion Laboratory, Pasadena, CA, USA

⁸ Center for Aerosol Science and Engineering, Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, St. Louis, MO 63130, USA

20 ⁹ Department of Civil, Architectural and Environmental Engineering, Missouri University of Science and Technology, Rolla, MO 65409, USA

¹⁰ Department of Earth and Atmospheric Science, University of Houston, Texas, 77204, USA

¹¹ Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ 85721, USA

¹² Department of Geosciences, University of Arizona, Tucson, AZ 85721, USA

¹³ Department of Chemical and Environmental Engineering, University of Arizona, Tucson, AZ 85721, USA

25

*Correspondence to: Armin Sorooshian (armin@email.arizona.edu)

⁺Now at: Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ 85721, USA



Abstract. The tropical Western North Pacific (TWNP) is a receptor for pollution sources throughout Asia and is highly susceptible to climate change, making it imperative to understand long-range transport in this complex aerosol-
30 meteorological environment. Measurements from the NASA Cloud, Aerosol, and Monsoon Processes Philippines Experiment (CAMP²Ex; 24 Aug to 5 Oct 2019) and back trajectories from the National Oceanic and Atmospheric Administration Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPLIT) were used to examine transport into the TWNP from the Maritime Continent (MC), Peninsular Southeast Asia (PSEA), East Asia (EA), and West Pacific (WP). A mid-campaign monsoon shift on 20 Sep 2019 led to distinct transport patterns between the
35 southwest monsoon (before 20 Sep) and monsoon transition (after 20 Sep). During the southwest monsoon, long-range transport was a function of southwesterly winds and cyclones over the South China Sea. Low (high) altitude air generally came from MC (PSEA), implying distinct aerosol processing related to convection and perhaps wind shear. The monsoon transition saw transport from EA and WP, driven by Pacific northeasterly winds, continental anticyclones, and cyclones over the East China Sea. Composition of transported air differed by emission source and accumulated precipitation along trajectories (APT) as an indicator of convection. MC air was characterized by biomass
40 burning tracers while major components of EA air pointed to Asian outflow and secondary formation. Convective scavenging of PSEA air was evidenced by considerable vertical differences between aerosol species but not trace gases, as well as notably higher APT and smaller particles than other regions. Finally, we observed a possible wet scavenging mechanism acting on MC air aloft that was not strictly linked to precipitation. These results are important
45 for understanding the transport and processing of air masses with further implications for modeling aerosol lifecycles and guiding international policymaking on public health and climate.



1. Introduction

50 As pollution emissions from Asian countries have surpassed those of countries in Europe and North America (Akimoto, 2003; Smith et al., 2011), Asia becomes increasingly important from a global climate and health perspective. The tropical Western North Pacific (TWNP), situated adjacent to Southeast Asia (Fig. 1), is a receptor for multiple sources of aerosol particles throughout the region (Bagtasa et al., 2018; Hilario et al., 2020a; Huang et al., 2019; Reid et al., 2015) and is one of the most susceptible regions to global climate change (IPCC, 2014; Reid et al., 2013; Yusuf and Francisco, 2009). Amidst several multi-scale meteorological phenomena ranging from the Asian monsoon system (Akasaka et al., 2007; Chang et al., 2005), the El Niño Southern Oscillation (Cruz et al., 2013; Jose et al., 1996), the Madden-Julian Oscillation (Maloney & Hartmann, 2001; Pullen et al., 2015), and intermittent typhoons (Bagtasa, 2017; Maloney and Dickinson, 2003), the TWNP hosts arguably one of the most complex meteorological environments in the world with likewise intricate relationships to aerosol lifecycle and climate impacts (Reid et al., 2012; Ross et al., 2018).

60 Owing to atmospheric residence times ranging from days to weeks (Balkanski et al., 1993; Kritz and Rancher, 1980) and enabled by the surrounding meteorology, aerosol particles from multiple sources can undergo long-range transport into the TWNP (Lin et al., 2007; Xian et al., 2013). These sources include biomass burning from the Maritime Continent (MC) (Hilario et al., 2020a, 2020b; Reid et al., 2015), anthropogenic and dust outflow from East Asia (EA) (Bagtasa et al., 2019; Braun et al., 2020; Geng et al., 2019; Miyazaki, 2003; Oshima et al., 2012; Tan et al., 2012), 65 emissions from Peninsular Southeast Asia (PSEA) (Bagtasa et al., 2019; Geng et al., 2019; Huang et al., 2020; Lin et al., 2009; Nguyen et al., 2020), and marine aerosol particles from the western Pacific (WP). Such transport is controlled by the interplay of several factors such as topography, sea breeze, monsoon flows, and typhoons (Reid et al., 2012; Wang et al., 2013b). Aside from the risk posed by transported anthropogenic aerosol on public health (Lelieveld et al., 2015; Zhang et al., 2017), such a diverse set of aerosol sources and types can result in variable aerosol-cloud-climate interactions (Hamid et al., 2001; Heald et al., 2014; Rosenfeld, 1999; Ross et al., 2018; Sorooshian et al., 2009; Yu et al., 2006; Yuan et al., 2011), which are complicated further by the spatial inhomogeneity of transported aerosol particles (Akimoto, 2003). As the influence of aerosol particles on climate remains one of the largest uncertainties in our understanding of the atmosphere (IPCC, 2014), investigating the composition and transport mechanisms of air masses from different source regions will aid in the future development of transport models and lead to a better understanding of the transport pathways that modulate aerosol particles in this part of the world.

75 Previous aircraft campaigns in Asia and the Pacific include the Transport and Chemical Evolution Over the Pacific (TRACE-P) campaign (Jacob et al., 2003), the Aerosol Radiative Forcing in East Asia (A-FORCE) campaign (Oshima et al., 2012), the Pacific Exploratory Mission – West A and B (PEM-West) (Hoell et al., 1996, 1997), and the Oxidant and Particulate Photochemical Processes Above a South East Asian Rainforest (OP3) project (Hewitt et al., 2010). 80 These campaigns examined springtime outflow from the Asian continent (e.g., Koike et al., 2003; Kondo et al., 2004; Park, 2005) and early-summertime characteristics of local and transported aerosol over Borneo (e.g., Robinson et al., 2011, 2012); however, no study to our knowledge has utilized aircraft data to characterize long-range transport patterns over the TWNP coinciding with the peak agricultural burning period for Indonesia and Malaysia. Limited ship observations in association with the 7 Southeast Asian Studies (7SEAS) program (e.g., Reid et al., 2015, 2016a, 2016b) 85 found a highly dynamic aerosol environment (Atwood et al., 2017; Hilario et al., 2020c; Reid et al., 2015).

The NASA Cloud, Aerosol, and Monsoon Processes-Philippines Experiment (CAMP²Ex) aircraft campaign examined the influence of meteorology, convection, and radiative effects on gas and aerosol species in the TNWP. Based at Clark, Luzon, Philippines, from 24 August to 5 October 2019, CAMP²Ex obtained a wide array of measurements between 0 – 9 km above ground level (AGL) across 19 research flights (RF). Some RFs were conducted in 90 coordination with the seaborne research vessel R/V Sally Ride as part of the Office of Naval Research Propagation of InterSeasonal Tropical Oscillation (PISTON) project (<https://onrpiston.colostate.edu/>). The CAMP²Ex campaign is unique in that it began during the peak of the Asian southwest monsoon (SWM) and coincided with an early monsoon transition (MT), which occurred in late-September instead of the more common time in October (Chang et al., 2005; Matsumoto et al., 2020). The early MT led to diverse transport patterns (Fig. 2) that offered an opportunity to examine 95 long-range transport into the TWNP.



100 Aircraft campaigns allow for vertically-resolved measurements of air mass properties, which are essential to better understand the atmosphere, as aerosol-cloud-climate interactions vary by altitude (Dahutia et al., 2019; Dong et al., 2017; Hansen, 2005; Mishra et al., 2015) and vertical transport can influence air mass composition (Matsui et al., 2011; Moteki et al., 2012; Oshima et al., 2012, 2013). Furthermore, as the main route of aerosol removal from the atmosphere, wet scavenging is a crucially important aspect of aerosol vertical profiles and are linked to significant uncertainties in climate models (Neu & Prather, 2012; Wang et al., 2013). Vertically-resolved in situ observations in field campaigns targeting aerosol-cloud-meteorology interactions are necessary to advance understanding of scavenging processes to inform the spectrum of models ranging from smaller-scale process models to larger-scale climate models (MacDonald et al., 2018; Sorooshian et al., 2019).

105 As Asian emissions continue to increase as a consequence of rapid economic development, it is imperative to understand the influence of long-range transport on air quality and aerosol-cloud-climate feedbacks in this region. In this study, we focus on characterizing transported air masses from four key regions: the Maritime Continent (MC: 5° S – 6.8° N, 94.9° E – 119.5° E), Peninsular Southeast Asia (PSEA: 10° N – 23° N, 95° E – 109.5° E), East Asia (EA: 22° N – 44° N, 100° E – 122° E and 30° N – 44° N, 122° E – 145° E), and the West Pacific Ocean (WP: 3° N – 25° N, 130° E – 145° E). Using air mass back trajectories to complement the CAMP²Ex data, this study aims to (1) identify regional transport pathways into the TWNP and their associated synoptic conditions, (2) characterize air masses coming from different source regions in terms of composition and aerosol size distribution, and (3) estimate the influence of convection and precipitation on transported air masses. By examining how transport and scavenging mechanisms impact air mass composition, our results have implications for improving modeling of aerosol lifecycles in this meteorologically complex region of the world and guiding policymaking related to public health and climate.

2. Methods

2.1. CAMP²Ex observations

120 A major goal of the 2019 CAMP²Ex aircraft campaign was to understand aerosol-cloud-climate feedbacks in the TWNP (Di Girolamo et al., 2018). Although multiple aircraft were deployed, this study focused on measurements made onboard the NASA P-3B Orion (N426NA) aircraft. Aircraft altitude (m AGL hereafter) was calculated as the difference between GPS altitude and ground elevation provided by the Google Maps API, with an uncertainty of ± 5 m. Dry optical size distribution data were collected by the Laser Aerosol Spectrometer (LAS; TSI Model 3340) and are presented as an integrated particle number concentration for diameters between 100 and 1000 nm ($N_{100-1000\text{nm}}$; cm^{-3}). Uncertainty of $N_{100-1000\text{nm}}$ is estimated at 20%. Carbon monoxide (CO; ppm) and methane (CH_4 ; ppm) mixing ratios were measured by a dried-airstream near-infrared cavity ringdown absorption spectrometer (G2401-m; PICARRO, Inc.), with uncertainties of 2% and 1% and precisions of 0.005 ppm and 0.001 ppm, respectively. Ozone (O_3 ; ppbv) measurements were conducted with a dual beam UV absorption sensor (Model 205; 2B Technologies) with an uncertainty of 6 ppbv. Non-refractory aerosol composition in the submicrometer range was measured with a High-Resolution Time-of-Flight Aerosol Mass Spectrometer (AMS; Aerodyne) (DeCarlo et al., 2008). The species of relevance to this study include sulfate (SO_4^{2-}), ammonium (NH_4^+), nitrate (NO_3^-), and organic aerosol (OA), all of which are reported in units of $\mu\text{g m}^{-3}$ with uncertainties up to 50%. The AMS was operated in 1 Hz Fast-MS mode and averaged to 30-second time resolution for this study, with campaign-averaged 1-sigma detection limits (in $\mu\text{g m}^{-3}$) as follows: 0.169 (OA), 0.039 (SO_4^{2-}), 0.035 (NO_3^-), 0.169 (NH_4^+). Mass concentrations below these detection limit values, which are sometimes negative due to the AMS difference method, are statistically equal to zero. Black carbon (BC; ng m^{-3}) was measured with a Single-Particle Soot Photometer (SP2) (Moteki & Kondo, 2007, 2010), including an uncertainty of 10%. We note that BC data were unavailable during one flight covering a major Borneo smoke event (RF9); thus, the BC value presented in Table 1 is likely under-represented compared to the AMS data. A Fast Integrated Mobility Spectrometer (FIMS) measured aerosol size distribution between 10 nm and ~ 600 nm with a concentration uncertainty of $\sim 15\%$ and size uncertainty of $\sim 3\%$ (Wang et al., 2017a, 2017b, 2018a).

140 All aerosol data are reported at standard temperature and pressure (STP; 273 K, 1013 hPa). Only data collected from outside of clouds via isokinetic sampling (McNaughton et al., 2007) were used to eliminate sampling artifacts from the shattering of large water and ice particles (Murphy et al., 2004). Background concentrations of each species were defined as the lowest 10th percentile of all CAMP²Ex data along vertical profiles for every 5 K range of potential temperature (Koike et al., 2003; Matsui et al., 2011a). Enhancements above these background concentrations are



145 denoted by Δ . For species ratios, only data with $\Delta\text{CO} > 0.02$ ppm were included similar to past work (Kleinman et al., 2007; Kondo et al., 2011; Matsui et al., 2011b).

150 Only data along profiles extending vertically more than 2 km were considered for analysis as they provide a “snapshot” of the atmosphere with which we can compare more directly air mass characteristics across different altitudes. Data collected when the P-3B aircraft sampled directly over urbanized Luzon ($13^\circ\text{N} - 15.8^\circ\text{N}$, $120^\circ\text{E} - 122^\circ\text{E}$) were excluded from analysis to minimize the impact of local emissions. Flight tracks and identified vertical profiles are shown in Fig. S1 of the Supplementary Information (SI).

2.2. Back trajectory classification

155 The National Oceanic and Atmospheric Administration (NOAA) Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPPLIT) (Rolph et al., 2017; Stein et al., 2015) was used to generate 120-hour back trajectories along vertical profiles with one minute temporal resolution. Input meteorological data was from the National Centers for Environmental Prediction (NCEP) Global Forecast System (GFS) at a horizontal resolution of $0.25^\circ \times 0.25^\circ$.

160 Our analysis focused on transport from key source regions (MC, PSEA, EA, WP). We note here that “source region” refers to the attributed origin of an air mass as identified by our trajectory classification scheme and does not preclude the possibility of entrainment from emission sources during transport (e.g., shipping). Our classification scheme considered two important environmental factors: (1) the synoptic shift that occurred around 20 Sep 2019, dividing the CAMP²Ex period into the SWM (before 20 Sep) and MT (after 20 Sep); and (2) the vertical wind shear across the region (Fig. 2). To better capture the pronounced effect of the monsoon shift, air masses were only classified as MC or PSEA (EA or WP) if sampled during the SWM (MT). For example, instances of EA air sampled during the SWM were classified as “Other” while air from EA sampled during MT were classified as EA. The inclusion of a monsoon phase filter more explicitly highlights the temporal aspect of meteorology in the TWNP; however, without this monsoon phase criterion, resulting air mass classifications remain largely unchanged (Section 3.2). Furthermore, to account for regional vertical inhomogeneity (Atwood et al., 2013; Sarkar et al., 2018), our analysis of air mass characteristics differentiates between boundary layer (BL; < 2 km) and free troposphere (FT; > 2 km) (Section 3.3).

170 For an air mass to be classified into a source region, its back trajectory must pass within a source region’s bounding box for more than 6 h at an altitude below 2 km, which is the typical summertime BL height in the region (Chien et al., 2019). In addition to excluding data collected over urbanized Luzon (Section 2.1), trajectories passing through the Philippines ($12^\circ\text{N} - 18.25^\circ\text{N}$, $120.5^\circ\text{E} - 122.5^\circ\text{E}$ and $5.1^\circ\text{N} - 14.5^\circ\text{N}$, $122.5^\circ\text{E} - 126.7^\circ\text{E}$) under our trajectory classification scheme were excluded to further focus our analysis on long-range transport and associated processes.

175 Most air masses came from only one of the four source regions: WP (117 occurrences), MC (174 occurrences), PSEA (88 occurrences), EA (130 occurrences). Air masses that passed through both EA and WP (12 occurrences) were considered as EA air due to the considerable influence of EA outflow on air mass composition (Talbot et al., 1997). Other transport permutations (e.g., air that passed through both MC and PSEA) did not meet the requirements of our classification scheme and were omitted. Thus, we focus on the four major transport pathways (MC, PSEA, WP, EA). Focusing on these major pathways adds robustness to the analysis by partly compensating for the limits of the trajectory model in capturing more complex meteorological phenomena such as wind shear (Freitag et al., 2014), which have been shown to contribute to trajectory uncertainty (Siems et al., 2000; Stohl et al., 2002). In addition to requiring that the back trajectories pass through the source regions, the additional criteria imposed (e.g., altitude < 2 km over the source region) increase our confidence that the remaining cases represent instances of long-range transport. Resulting source region contributions per RF are shown in Fig. S2. We emphasize that these source region contributions represent frequencies of observation rather than frequencies of occurrence, as the sampling location of the aircraft introduces a bias inherent in aircraft campaigns (Section 3.2).

185
190 As a consequence of our filtering scheme, a large portion of trajectories were tagged as “Other” (66.8%). This is attributable to several scenarios: (1) air masses that passed over source regions, but above our BL height threshold of 2 km; (2) air masses influenced by the Philippines (i.e., air masses that stayed over the Philippines at < 2 km for more than 6 hours); (3) stagnant air masses that did not reach any source region; and (4) other transport permutations that occurred too infrequently to provide robust statistics.



195 Trajectory clustering was performed using two well-established methods: k-means and Ward linkage (Govender and
Sivakumar, 2020) in order to confirm the robustness of our predefined source regions. K-means clustering classifies
data into k clusters such that the sum of squares per cluster is minimized (Hartigan and Wong, 1979), with the
drawback that k must be specified beforehand. Ward linkage is a hierarchical clustering method that merges clusters
such that the increase in intra-cluster Ward's distance is minimized (Ward, 1963) and has been described as the method
that most closely accomplishes the goals of clustering (Tufféry, 2011). More comprehensive descriptions of these
clustering methods can be found elsewhere (Govender and Sivakumar, 2020; Pérez et al., 2017). Prior to clustering, a
200 weighted distance matrix was calculated, similar to Taubman et al. (2006): (1) normalized trajectory coordinates to
give equal weighting to both horizontal and vertical transport; (2) weighted time steps linearly back in time; and (3)
assigned nearby points (time step < 6 h) zero weighting on the clustering to remove the influence of aircraft position
on the clustering.

2.3. Accumulated precipitation along individual trajectories

205 Accumulated precipitation along individual trajectories (APT) was calculated using data from satellite precipitation
products (SPPs): (1) the Precipitation Estimation from Remotely Sensed Information using Artificial Neural Networks
– Climate Data Record (PERSIANN-CDR) (Ashouri et al., 2015; Nguyen et al., 2018); (2) the Integrated Multi-
satellitE Retrievals for the Global Precipitation Measurement (GPM) mission (IMERG) (Huffman et al., 2019); and
210 (3) the Tropical Rainfall Measuring Mission (TRMM) Multi-satellite Precipitation Analysis (TMPA) 3B42-V7
(Huffman et al., 2007). The purpose of utilizing multiple SPPs is to account for the uncertainties inherent in satellite
retrievals, particularly during very light or heavy rainfall conditions, providing us with an ensemble of estimates rather
than relying on a single SPP (Chen et al., 2020; Liu, 2016; Maggioni et al., 2016; Mahmud et al., 2017; Tan & Santo,
2018). Furthermore, although these SPPs measure surface precipitation and do not fully capture scavenging effects
aloft, we use APT as an indicator of whether an air mass passed through a convectively active region.

215 PERSIANN-CDR ($0.25^\circ \times 0.25^\circ$, daily resolution) uses a modified PERSIANN algorithm utilizing NCEP Stage IV
hourly precipitation and monthly precipitation from the Global Precipitation Climatology Project (GPCP) to maintain
monthly amounts consistent with GPCP (Ashouri et al., 2015). PERSIANN-CDR data are available at the Center for
Hydrometeorology and Remote Sensing (CHRS) Data Portal (<http://chrsdata.eng.uci.edu>) (Nguyen et al., 2019).

220 IMERG ($0.1^\circ \times 0.1^\circ$, 30-min resolution) integrates multiple satellite retrievals from the passive microwave (MW)
precipitation retrievals provided by the suite of GPM constellation passive microwave radiometers (Kummerow et al.,
2015), the Climate Prediction Center MORPHing technique (CMORPH) from NOAA, and PERSIANN-Cloud
Classification System (PERSIANN-CCS; Hong et al., 2004) from the University of California, Irvine. These data are
available from the NASA Precipitation Processing System (Skofronick-Jackson et al., 2018). For inter-calibrating
various MW precipitation products, IMERG uses the GPM_2BCMB product (Olson et al., 2018) that utilizes the
225 GPM Microwave Imager (GMI) and Dual-frequency Precipitation Radar (DPR) instruments on the GPM core satellite
IMERG (Hou et al., 2014; Kidd et al., 2020). For this study, we use IMERG Final Run data, available at
<https://pmm.nasa.gov/data-access/downloads/gpm>.

230 TMPA ($0.25^\circ \times 0.25^\circ$, 3-h resolution) similarly combines data from multiple satellites such as TRMM (pre-2015),
NASA's *Aqua*, and the NOAA satellite series, involving calibration with gauge data when feasible (Huffman et al.,
2007). Though TRMM ended its service in 2015, the TMPA 3B42 algorithm was continued in parallel with IMERG
through December 2019 and had been based on a climatological calibration since 2014. As TMPA is climatologically
calibrated, TMPA may be less sensitive to interannual variability in precipitation; thus, including TMPA in this study
may provide a better idea of the spread among SPPs. TMPA data are accessible through [https://pmm.nasa.gov/data-
access/downloads/trmm](https://pmm.nasa.gov/data-access/downloads/trmm).

235 Precipitation along each trajectory was integrated from the trajectory spawn point (i.e., P3-B sampling location) to the
point when it reaches the boundary of a source region. An additional 24 h along the trajectory after reaching a source
region was included in the APT integration to account for precipitation effects within the source region (Matsui et al.,
2011a, b). No significant APT differences were found between using 0, 24, or 48 h for the APT calculation, suggesting
that our results are robust with regard to the added duration. Furthermore, APT comparisons demonstrate that our
results are independent of chosen SPP in terms of APT ranking (i.e., all SPPs agreed on which source regions are
240 associated with the highest or lowest APTs).



2.4. Other Data

Elevation data (Fig. 1a) were acquired from the United States Geological Survey (USGS) Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010) (Danielson and Gesch, 2011). Population density data for 2020 (Fig. 1b) were retrieved from the Gridded Population of the World (GPW), v4 (Center for International Earth Science
245 Information Network, 2018) (<https://sedac.ciesin.columbia.edu/data/set/gpw-v4-population-density-rev11/data-download>). Depicted in Fig. 1c, active fires tagged with high confidence ($> 80\%$; Bhardwaj et al., 2016) were obtained from the Moderate Resolution Imaging Spectroradiometer (MODIS) Collection 6 algorithm (<https://firms.modaps.eosdis.nasa.gov/>) (Levy et al., 2013) and converted into average fire density at $0.5^\circ \times 0.5^\circ$ resolution. Planetary BL (PBL) sulfur dioxide (SO_2) was retrieved by the Ozone Monitoring Instrument (OMI) and
250 obtained from NASA Goddard Earth Sciences Data and Information Services Center (GES DISC) (Krotkov et al., 2015). The OMI SO_2 data were then resampled to $1^\circ \times 1^\circ$ resolution and averaged between 1 August and 15 October 2019 (Fig. 1d). Reanalysis data from NCEP/NCAR ($2.5^\circ \times 2.5^\circ$) were used to examine synoptic conditions (Kalnay et al., 1996).

3. Results and Discussion

255 3.1. Observed transport patterns during CAMP²Ex

Figure 1 provides an overview of the general source regions impacting the TWNP. The TWNP is surrounded by areas of high population density in EA, MC, and PSEA (Fig. 1b). Burning was prevalent mainly in the MC (Fig. 1c); though, fires were also detected along the eastern PSEA coast. Satellite retrievals of PBL SO_2 reveal possible point sources (Fig. 1d), perhaps from volcanoes, shipping, burning, and industry (Fioletov et al., 2016; Guttikunda et al., 2001;
260 Zhang et al., 2019); however, we caution that cloud contamination may influence the SO_2 retrievals and are used here to demonstrate the variety of sources in Asia.

Trajectories from each source region show distinct pathways (Fig. 3; left column), indicative of differences in accompanying synoptic-scale circulation. These pathways are generally corroborated by both k-means and Ward linkage clustering methods (Figs. S3 and S4), confirming the robustness of our predefined source regions (Fig. 1a).
265 Prior to further discussion, we emphasize the temporal aspect of these observed transport patterns (Figs. 2 and 3), in particular, their dependence on both synoptic (e.g., SWM) and mesoscale meteorology (e.g., typhoons), which varied during CAMP²Ex in terms of phase and frequency, respectively. Consequently, a specific transport pattern may be more dominant in one monsoon phase and less so in another while being enhanced (or suppressed) by intermittent mesoscale phenomena.

270 3.1.1. Southwest monsoon

Beginning with transport during the SWM prior to 20 Sep 2019, PSEA air is advected by uniform westerlies (Fig. 3a) associated with cyclonic activity over the northern South China Sea (SCS) (Fig. 3b) (Cheng et al., 2013; Huang et al., 2020; Lin et al., 2009). In comparison, although MC transport also occurs during the SWM, the mechanism behind
275 MC transport is driven instead by southwesterlies originating across the MC (Fig. 3d) (Ge et al., 2014; Wang et al., 2013; Xian et al., 2013). Transport from the MC is further promoted by well-developed cyclones entering PSEA (Fig. 3e), as previously highlighted by observational (Hilario et al., 2020; Reid et al., 2015) and modeling studies (Wang et al., 2013). The similar cyclonic activity over northern SCS/PSEA may explain the confluence of air masses from PSEA and MC (e.g., RF6), indicated by the frequent sampling of MC and PSEA air in the same RF (Fig. S2).

A key difference between PSEA and MC air is that PSEA air passed through convective areas over the PSEA (Takahashi et al., 2010), the SCS (Fig. 3a, c) (Chen et al., 2017), and along the western coast of the Philippines (Akasaka et al., 2007; Chen et al., 2017; Cruz et al., 2013; Hilario et al., 2020b) while MC air passed through areas with less precipitation (Figs. 3d, f). As a result, PSEA air showed much higher APT than MC air (Table 1) and was more likely to have been processed by clouds. The transport pathway of PSEA through convective regions may lead to wet scavenging and aqueous-phase processing (MacDonald et al., 2018; Moteki et al., 2012; Sorooshian et al., 2006,
285 2007; Wonaschuetz et al., 2012), affecting both air mass composition and particle size distributions (Section 3.3).

In terms of sampled air masses, PSEA and MC showed contributions of 5.7% and 11.3%, respectively (Fig. 4a), and differ in terms of vertical distribution (Fig. 4b). PSEA air was sampled across a wide range of altitudes with the



majority of observations over 900 hPa, similar to Kondo et al. (2004), explainable by convection-related lofting (Fig. 5a). Very few observations of PSEA air were made near the surface. The lofting of PSEA air can occur over the PSEA itself (Fig. 3c) (Kondo et al., 2004), through mechanisms like orographic effects (Lin et al., 2009). Convection over the SCS trough likely also contributes to lofting (Fig. 3c) (Chen et al., 2017). Lofting of PSEA air into the FT has important downstream ramifications as it modulates both aerosol composition and size distribution (Matsui et al., 2011a; Moteki et al., 2012; Oshima et al., 2013). However, we note that vertical motion is an important source of uncertainty in trajectory models (Harris et al., 2005) and should be interpreted with caution.

In contrast to PSEA air, sampled MC air was well-mixed within the BL (Fig. 5b), but the observation frequency of MC air dropped sharply above 750 hPa, consistent with previous modeling studies (e.g., Xian et al., 2013). Such distinct vertical distributions between MC and PSEA air are perhaps due to highly sheared environment during the SWM, generally contributing to distinct air mass sources across different altitudes (Atwood et al., 2013; Sarkar et al., 2018) and varying degrees to which these air masses are processed (Section 3.3).

3.1.2. Monsoon transition

The arrival of the MT period after 20 Sep 2019 led to a synoptic-scale shift (Fig. 2), allowing the sampling of air from EA and WP (Fig. 3g, j). Transport from EA was observed across several MT flights (Fig. S2) and originated mainly from southeastern China, Korea, and Japan (Fig. 3g), suggesting the entrainment of anthropogenic emissions (Section 3.3) (Cheng et al., 2013; Hatakeyama et al., 2001, 2004; Kim et al., 2009; Wang et al., 2016). Depicted in Fig. 3h, Asian outflow was promoted by the pairing of a well-developed cyclone passing over the East China Sea (Hatakeyama et al., 2001, 2004; Uno et al., 1998) and an anticyclone over the Asian continent (Honomichl and Pan, 2020). In comparison, WP transport was observed mainly towards the end of the campaign (Fig. S2), likely a consequence of sampling location, and was driven by Pacific northeasterlies (Figs. 3j – k). Transport from the WP, similar to that of EA, coincided with an anticyclone over the Asian continent (Fig. 3k); however, WP transport is marked by the absence of the East China Sea cyclone that promoted southward transport of EA air (Fig. 3h). This difference may explain why EA and WP air were usually sampled in separate RFs (Fig. S2), in contrast to PSEA and MC air, which tended to be sampled together.

Air from EA and WP show similarly low APT (Table 1), explainable by the generally lower precipitation in MT (Figs. 3i, l) compared to SWM (Figs. 3c, f) (Matsumoto et al., 2020), as well as the lower number of cyclone occurrences after 20 Sep 2019. Although EA transport was driven by a well-developed cyclone (Fig. 3h), trajectories suggest that EA air traveled through the outer bands of the cyclone (Fig. 3g), largely avoiding high precipitation areas (Fig. 3i). This suggests that anthropogenic emissions entrained in EA air experienced low levels of scavenging and were more likely to be sampled, in contrast with high APT urban source regions like PSEA (Section 3.3).

Transport from EA and WP were quite similar in terms of relative contribution (8.5% and 7.6%, respectively; Fig. 4a) and vertical sampling distribution (Fig. 4b). Sampling of EA and WP air were largely constrained to the BL, though sampled EA air was unimodal while WP air was more evenly sampled. In terms of vertical motion during transport, some EA trajectories exhibited downward motion (Fig. 5c), likely due to subsidence from the continental anticyclone (Fig. 3h), contrasting the vertical motion of PSEA air, which generally experienced upward motion associated with convection (Fig. 5a).

In summary, important transport features over the TWNP include the following: (1) SWM transport from the MC and PSEA was driven by southwesterlies and cyclonic activity over northern SCS/PSEA while MT transport from EA and the WP was driven by Pacific northeasterlies, anticyclones over the Asian continent, and well-developed cyclones over the East China Sea; (2) EA and MC air were sampled largely within the BL, did not exhibit significant upward motion, and experienced low APT, suggesting that they likely carry urban/continental or biomass burning emissions; in contrast, (3) PSEA air may have undergone a high degree of aerosol scavenging over convective regions (e.g., SCS), indicated by high APT and upward motion of trajectories.

3.2. Sensitivity analysis

In order to assess the uncertainty associated with our trajectory classification, we evaluated the effect of the following variables on source classification: (1) trajectory height threshold over source regions; (2) back trajectory run time; (3) vertical profile filtering; (4) monsoon phase; and (5) aircraft sampling location. Results are provided in Table S1 and



340

summarized below. Except for aircraft sampling location, independently changing any of these variables had little effect on the resulting source-region distribution. The relative contributions of source regions did vary significantly with sampling location, though areas surrounding Luzon (e.g., East of Luzon, North of Luzon) showed some degree of similarity. Thus, we emphasize that, as with any aircraft campaign, observed transport is to some degree dependent on aircraft location.

345

In order to reduce the effect of local emissions, we excluded trajectories classified as influenced by the Philippines (PH). To evaluate our filter for Philippine-influenced trajectories (hereafter, PH filter), air mass characteristics were compared between transported air unaffected by PH (No-PH air; e.g., MC) and transported air mixed with PH air (With PH; e.g., MC-PH). A local signal was observed for $N_{100-1000\text{nm}}$, suggested by differences in the histograms of $N_{100-1000\text{nm}}$ between non-PH and PH-mixed air (Fig. S5), particularly for MC and PSEA air. Differences in the species concentration histograms of non-PH and PH-mixed air were also observed for other anthropogenic species (BC, OA, SO_4^{2-} ; not shown), confirming the effectiveness of the PH filter.

3.3. Chemical composition of transported air masses

350

A convenient opportunity afforded by having conducted the air mass classification presented above was to examine how gas and aerosol properties vary for each source region based on vertically-resolved in situ aircraft measurements. To account for regional vertical wind shear (Fig. 2) while considering the generally lower classification frequency at higher altitudes (Fig. 4b), air mass characterization was resolved into BL and FT subsets and composited by source region (Table 2). The delineation between BL and FT composition is demonstrated by selected species (Fig. 6), which generally dropped in concentration above the BL (< 850 hPa). Prior to further discussion, we note here that shipping is a major regional source (Streets et al., 1997, 2000) and may contribute appreciably to all air masses.

355

Significant differences in composition were observed in the same monsoon season (e.g., SWM) depending on air mass origin. Air from PSEA had much lower species concentrations than MC (Table 2) due to decreased emissions and increased potential for wet scavenging. Sampled MC air showed statistically significant differences between BL and FT concentrations for both gas and aerosol species (Table S2), indicative of emissions constrained to the BL, and exhibited strong biomass burning signals in its composition profile (e.g., $N_{100-1000\text{nm}}$, CO, NO_3^- , OA, and BC) (Maudlin et al., 2015; Pósfai et al., 2003; Reid et al., 1998, 2005; Theodoritsi et al., 2020; Yadav et al., 2017). We note that peaks of CO (Fig. 6b) and CO_2 (not shown) were observed in MC samples at around 650 hPa, suggestive of MC burning emissions lofted into the FT; however, this feature consisted of few samples and did not appear in other gases (e.g., SO_2) and thus warrants caution in its interpretation.

360

365

In contrast, PSEA air was generally characterized by concentration magnitudes between those of MC and WP. Aerosol concentrations of PSEA air in the FT were lower by at least an order of magnitude than those in the BL (SO_4^{2-} , NH_4^+ , OA, BC) while trace gases (CO, CH_4) showed more similar BL and FT concentrations (Tables 2 and S2; Fig. S6). These aerosol-gas differences point to: (1) the lofting of PSEA air into the FT, as suggested by the similarity of trace gas concentrations between BL and FT, and (2) the consequent scavenging of aerosol particles, explaining the much lower aerosol concentrations in the FT (Oshima et al., 2012, 2013; Sievering et al., 1984). For comparison, MC air showed large BL and FT differences in both aerosol and gas species, the latter of which indicates the infrequent lofting of MC air (Figs. 4b and 5b). Since PSEA air came from a populated region (Fig. 1b) and likely originally contained anthropogenic aerosol particles, these unique characteristics of PSEA air compared to MC and EA air support the likelihood of aerosol scavenging in PSEA air. These observations are robust due to the relatively even sampling frequency of PSEA across altitudes (Fig. 4b).

370

375

380

Transport during the MT season showed similarly distinct composition profiles depending on air mass origin. Air from EA exhibited higher concentrations of SO_4^{2-} , O_3 , CH_4 , and NH_4^+ , owing to urban emissions in continental outflow (Chuang et al., 2014; Talbot et al., 1997; Thornton et al., 1999; Umezawa et al., 2014; Wang et al., 2007) and extensive secondary aerosol formation (Hatakeyama et al., 2001, 2004, 2011; Krupa and Manning, 1988; Matsui et al., 2014). In contrast, WP air is characterized as pure marine due to composition similar to those previously reported in Pacific marine environments (Davis et al., 1996; Matsumoto et al., 1998; Talbot et al., 1997).



3.3.1. Species ratios

Composition profiles between regions (Table 2) reveal clear differences as a function of (1) emission source and (2) passage through convective regions indicated by APT (Table 1). The role of emission source was most evident when comparing air masses of low APT (EA, MC). Though EA and MC had similar BL values for $N_{100-1000\text{nm}}$ (Table 2), they showed distinct chemical differences: MC air was characterized by higher concentrations of biomass burning tracers (e.g., CO, CH₄, NO₃⁻, OA) while EA air showed influence from urban/continental sources and secondary formation (e.g., O₃, CH₄, NH₄⁺, SO₄²⁻). Such differences in composition are corroborated by species ratios derived with linear regression (Fig. 7). Prior to a discussion on the species ratios, we note that the reported species ratios in this study are difficult to compare directly to at-source measurements of the same quantity because the composition of air masses was likely influenced by sources and sinks during transport (e.g., Choi et al., 2019; Conte et al., 2019; Gruber et al., 2019; Yang et al., 2009); however, differences in species ratios can still aid in air mass characterization and point to possible emission sources.

In Fig. 7a, the lower $\Delta\text{CO}/\Delta\text{CO}_2$ ratio of EA air versus MC air (Fig. 7a) is indicative of an inefficient combustion signature in MC air (Halliday et al., 2019), attributable to the predominantly smoldering phase of MC fires (Gras et al., 1999; Reid et al., 2013). We note that the poor $\Delta\text{CO}-\Delta\text{CO}_2$ correlation for MC air indicates that our reported ratio does not reflect expected emission ratios (Andreae, 2019; Hurst et al., 1994). This further suggests (1) our source region classification (i.e., MC, EA) may not perfectly capture air mass differences, and (2) additional sources of CO or CO₂ during transport. Thus, it is necessary to use multiple species ratios to supplement air mass chemical characterization.

The strong relationship between CH₄ and CO in MC air is a good indication of biomass burning influence (Andreae, 2019; Hecobian et al., 2011). The ratio of $\Delta\text{CH}_4/\Delta\text{CO}$ (Fig. 7b) was much higher in EA air compared to MC air, indicating the dominance of CH₄ from residential and industrial activity (Geng et al., 2019; He et al., 2019; Tohjima et al., 2014) as well as from rice cultivation in EA (Xia et al., 2020).

As an indicator of aerosol hygroscopicity (Kreidenweis and Asa-Awuku, 2014; Malm et al., 2005; Svenningsson et al., 2006), the $\Delta\text{OA}/\Delta\text{SO}_4^{2-}$ ratio pointed to higher hygroscopicity in EA air than MC air (Cheung et al., 2020; Saxena et al., 1995; Wang et al., 2017, 2018, 2019). Interestingly, though peat burning in the MC has been associated with SO₄²⁻ (Ikegami et al., 2001; Reid et al., 2013), the elevated $\Delta\text{OA}/\Delta\text{SO}_4^{2-}$ ratio in MC air (4.85 ± 0.24) is indicative of lower hygroscopicity, explainable by the high levels of OA emitted during biomass burning (Radzi bin Abas et al., 2004; Theodoritsi et al., 2020) or produced through gas-to-particle conversion during transport (Cappa et al., 2020; Mardi et al., 2018; Zhou et al., 2012). The lower $\Delta\text{OA}/\Delta\text{SO}_4^{2-}$ ratio of EA air (0.75 ± 0.09) signaling high hygroscopicity is attributable to the high levels of SO₂ in EA (Fig. 1d), leading to the secondary formation of SO₄²⁻ (e.g., Hatakeyama et al., 2011).

Although MC was calculated to have low APT (Table 1), a comparison of BL and FT air from MC (Figs. S6 and S7) allows for speculation on a possible scavenging mechanism acting on FT air. Linear regressions of $\Delta\text{SO}_4^{2-}/\Delta\text{CO}$ (Fig. S7a) suggested the removal of SO₄²⁻ from FT air while $\Delta\text{OA}/\Delta\text{CO}$ (Fig. S7b) indicated no such effect on OA. Considering the higher hygroscopicity and therefore scavenging susceptibility of SO₄²⁻ compared to OA (e.g., Kreidenweis and Asa-Awuku, 2014), we speculate the removal of SO₄²⁻ to be related to wet scavenging. Comparisons of BL and FT species concentrations (Fig. S6) further supports the possibility of this hypothesized scavenging, as aerosol species have significantly lower concentrations in the FT compared to BL, a difference not observed for trace gases. The potential for scavenging was also supported by number size distributions (Section 3.4).

We note that this wet scavenging mechanism is not apparent from APT, which suggested dry conditions for MC air (Table 1). This disagreement with APT stems from the usage of SPPs which typically describe surface precipitation and, consequently, our APT may not fully capture potential wet scavenging effects aloft. Thus, the speculated scavenging mechanism of MC air in the FT may occur at higher levels and may not be strictly linked to surface precipitation. Perhaps, this mechanism is related to processes such as in-cloud scavenging (Sievering et al., 1984; Yang et al., 2020; Yu et al., 2020) but, indeed, this requires a deeper investigation in future work.

Due to BC's lack of secondary sources, the ratio of $\Delta\text{BC}/\Delta\text{CO}$ has been used to gauge transport efficiency as affected by physical removal processes on air masses (Matsui et al., 2011b; Moteki et al., 2012; Oshima et al., 2012) and as an



430 indicator of combustion efficiency, which increases with $\Delta BC/\Delta CO$ (Kondo et al., 2011). The ratio of $\Delta BC/\Delta CO$ was much higher in EA air than in MC air (Fig. 7f), similar to observations by Pani et al. (2019) in Taiwan. This difference is explainable by burning in industrial and residential areas in EA (Bond et al., 2004; Geng et al., 2019) and the predominance of smoldering fires in the MC (Gras et al., 1999; Reid et al., 2013), which yield a lower $\Delta BC/\Delta CO$ than flaming fires (Kondo et al., 2011).

435 3.4. Size distributions of transported air masses

To more deeply characterize the air masses from different source regions, we examine the differences in normalized FIMS number (Fig. 8) and volume (Fig. S8) size distributions between BL and FT, which can also offer insights into the convection-related removal of PSEA air. In-cloud processing during transport may influence particle sizes in these air masses, whereby a combination of the following processes can occur (e.g., Ervens, 2015; Sorooshian et al., 2007; Wonaschuetz et al., 2012), followed by detrainment from the cloud or wet removal: (1) collisions between interstitial aerosol and droplets; (2) coalescence among droplets; and (3) aqueous-phase processing in droplets. However, comparisons between size distributions between regions and between BL and FT may still lend valuable insights into transport-related processes (e.g., Moteki et al., 2012).

445 Firstly, comparing source regions of low APT and high aerosol concentration, EA air in the BL (Fig. 8a) showed a wider peak in its size distribution (40 – 200 nm) than that of MC (Fig. 8b), which showed a clear unimodal peak (100 nm). This was perhaps an effect of multiple sources contributing to EA air masses, ranging from industrial activities to rice cultivation (Chen et al., 2020b; Geng et al., 2019; Wang et al., 2016; Xia et al., 2020). In comparison, biomass burning emissions from the MC have been shown to greatly influence air mass composition (Engling et al., 2014; Fujii et al., 2015; Santoso et al., 2011) and, by extension, such a dominant emission source can explain the large
450 unimodal peak in MC's BL size distribution (Figs. 8b and S8b).

A comparison of the FT and BL size distributions of MC air may point to a potential scavenging mechanism of MC air lofted into the FT, signaled by significant BL and FT differences above 50 nm (Fig. 8b). This potential scavenging mechanism of MC air was previously proposed using species ratios (Section 3.3.1), and the size distribution here provides further evidence for the hypothesized removal process; however, we emphasize that the hypothesized
455 scavenging mechanism is for now speculative and warrants future examination.

In contrast, the size distribution of PSEA air in the BL shows two peaks at around 50 nm and 200 nm. Comparing PSEA and MC air in the BL reveals much smaller particle sizes in PSEA air (Fig. 8c), explainable by differences in source emissions as well as in APT. A comparison of BL and FT air from PSEA pointed to scavenging during lofting into the FT: the FT size distribution of PSEA air showed a sharp drop in particle number concentrations above 50 nm
460 while the BL size distribution of PSEA air was much broader. In fact, the size distribution of FT air from the PSEA is more similar in shape to that of WP air (Fig. 8d), which is representative of background FT air. Due to this similarity, the peaks at 30 nm in FT air from PSEA and WP may originate from new particle formation (Williamson et al., 2019) that has been shown to be connected to APT in marine environments (Ueda et al., 2016), such as the convectively active SCS.

465 3.5. Influence of convection on transported air masses

The relationship between composition and convection was further investigated through scatterplots of $\Delta BC/\Delta CO$ ratio, an indicator of physical removal processes (Moteki et al., 2012), as a function of APT, an indicator of convection. The decrease in $\Delta BC/\Delta CO$ ratio with APT (Fig. 9a) indicates that convection during transport is one of the main scavenging mechanisms in the TWNP. Both EA and MC air showed very high $\Delta BC/\Delta CO$ ratios compared to PSEA,
470 indicative of more efficient transport, while having low APT, which allowed for a clear transport signal as shown by the high concentrations of anthropogenic or burning species in these air masses (Section 3.3). In contrast, PSEA air is characterized by a lower $\Delta BC/\Delta CO$ ratio coinciding with high APT (Fig. 3c; Table 1), pointing to particle scavenging over convective areas. To further demonstrate the impact of convection on transport efficiency, Fig. 9b reveals $\Delta BC/\Delta CO$ distributions resolved by low and high APT. The shift towards higher $\Delta BC/\Delta CO$ ratios under low APT
475 implies that dry and non-convective conditions are conducive to transport, suggesting the higher BL concentrations of anthropogenic species in MC, EA air were likely enabled by lower levels of wet removal.



4. Summary and Conclusions

Utilizing airborne CAMP²Ex measurements between 24 Aug and 5 Oct 2019 and HYSPLIT back trajectories, we examined transport patterns into TWNP from key source regions (PSEA, MC, EA, WP). Key conclusions from this study include the following:

1. Meteorological phenomena driving transport as well as the origins of transported air masses shifted significantly with the monsoon phase. During the SWM, MC and PSEA transport were associated with monsoon-driven southwesterly winds and cyclonic activity over the northern SCS. Wind shear was associated with predominantly BL (FT) sampling of MC (PSEA) air, implying distinct aerosol processing between these two source regions. In comparison, transport during the MT period from EA and WP was driven by northeasterly winds from the Pacific, anticyclones over the Asian continent, and well-developed cyclones passing through the East China Sea. These transport differences led to varying degrees of convection experienced by transported air masses. PSEA air generally passed through convective regions (SCS, west of Luzon, and over the PSEA itself) and was lofted into the FT, which led to scavenging of aerosols. In contrast, air masses from the MC and EA underwent relatively little convection, indicated by low APT, and mainly were confined to the BL, enabling the transport of anthropogenic emissions.
2. Characteristics of transported air masses differed primarily by emission source and passage through convective regions. Due to low APT and high $\Delta BC/\Delta CO$, transported air from MC and EA exhibited characteristic emissions: MC air from biomass burning (CO, well-correlated CO and CH₄, NO₃⁻, OA) and EA air from anthropogenic outflow and secondary formation (O₃, CH₄, NH₄⁺, SO₄²⁻). Key species ratios corroborated distinct sources between MC and EA air. Aerosol size distributions in EA air suggested multiple primary sources (industry, residential emissions, rice cultivation) as well as secondary formation, indicated by its relatively broad peak; in contrast, the narrower peak in the size distribution of MC air pointed to the predominance of biomass burning emissions.
3. Air from the PSEA showed strong evidence of particle scavenging: passage over high precipitation areas, convective lofting, high APT, low $\Delta BC/\Delta CO$, relatively low levels of anthropogenic species, and a size distribution shifted towards smaller particle sizes. Aerosol concentrations of PSEA air in the FT were lower by at least an order of magnitude than those in the BL, a difference that was not observed for trace gases, which pointed to scavenging of aerosol particles in the FT. Furthermore, PSEA air in the FT lacked the larger peak ($D_p = 200$ nm) observed in BL air and instead peaked at much smaller sizes ($D_p = 30$ nm), suggesting large particle removal during convective lofting. The fine mode peak ($D_p = 30$ nm) for PSEA FT air also resembled that of WP air, suggestive of new particle formation during transport from the PSEA, perhaps occurring over the convective SCS.
4. A possible wet scavenging mechanism for MC FT air was inferred from $\Delta SO_4^{2-}/\Delta CO$ and $\Delta OA/\Delta CO$ ratios between BL and FT, and corroborated by size distributions, which showed significant BL and FT differences for larger particles (> 50 nm). The disagreement with APT was attributed to SPP limitations in capturing scavenging effects aloft, hinting that the scavenging mechanism acts at higher layers and may not be linked to surface precipitation. However, we emphasize that the exact scavenging mechanism is for now speculative and warrants its own investigation in the future.

Recommendations for future work include: (1) investigating the hypothesized scavenging mechanism of MC air aloft using vertically-resolved moisture and convection retrievals; (2) examining deep convection periods to further evaluate wet scavenging effects on transported air; (3) characterizing aerosol hygroscopicity as a function of air mass source region and transport processes; and (4) comparing different sampling areas over the Philippines as impacted by transported air masses.

Data Availability. The CAMP²Ex dataset can be found at <https://doi.org/10.5067/Suborbital/CAMP2EX2018/DATA001> (last access: 11 September 2020). HYSPLIT data are accessible through the NOAA READY website (<http://www.ready.noaa.gov>; last accessed: 13 July 2020). Global elevation data from GMTED2010 are available at <http://temis.nl/data/gmted2010/> (last access: 12 March 2020). Population density data are provided by the Center for International Earth Science Information Network, available at <https://sedac.ciesin.columbia.edu/data/set/gpw-v4-population-density-rev11/data-download>; last access: 3 July 2020). MODIS active fire data can be downloaded through the Fire Information for Resource Management System



530 (<https://firms.modaps.eosdis.nasa.gov/>; last access: 29 June 2020). OMI data were retrieved from the NASA GES DISC website (<https://doi.org/10.5067/Aura/OMI/DATA3008>; last access: 27 June 2020). NCEP/NCAR Reanalysis data were provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their website (<https://www.esrl.noaa.gov/psd/>; last access: 13 June 2020).

Author contributions. MRAH performed the analysis and prepared the manuscript. All authors provided input for the manuscript and/or participated in data collection and processing.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This research was funded by NASA grant 80NSSC18K0148 in support of CAMP²Ex.

535 References

- Akasaka, I., Morishima, W. and Mikami, T.: Seasonal march and its spatial difference of rainfall in the Philippines, *Int. J. Climatol.*, 27(6), 715–725, doi:10.1002/joc.1428, 2007.
- Akimoto, H.: Global Air Quality and Pollution, *Science*, 302(5651), 1716–1719, doi:10.1126/science.1092666, 2003.
- 540 Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmospheric Chem. Phys.*, 19(13), 8523–8546, doi:10.5194/acp-19-8523-2019, 2019.
- Ashouri, H., Hsu, K.-L., Sorooshian, S., Braithwaite, D. K., Knapp, K. R., Cecil, L. D., Nelson, B. R. and Prat, O. P.: PERSIANN-CDR: Daily Precipitation Climate Data Record from
- 545 Multisatellite Observations for Hydrological and Climate Studies, *Bull. Am. Meteorol. Soc.*, 96(1), 69–83, doi:10.1175/BAMS-D-13-00068.1, 2015.
- Atwood, S. A., Reid, J. S., Kreidenweis, S. M., Cliff, S. S., Zhao, Y., Lin, N.-H., Tsay, S.-C., Chu, Y.-C. and Westphal, D. L.: Size resolved measurements of springtime aerosol particles over the northern South China Sea, *Atmos. Environ.*, 78, 134–143,
- 550 doi:10.1016/j.atmosenv.2012.11.024, 2013.
- Atwood, S. A., Reid, J. S., Kreidenweis, S. M., Blake, D. R., Jonsson, H. H., Lagrosas, N. D., Xian, P., Reid, E. A., Sessions, W. R. and Simpas, J. B.: Size-resolved aerosol and cloud condensation nuclei (CCN) properties in the remote marine South China Sea – Part 1: Observations and source classification, *Atmospheric Chem. Phys.*, 17(2), 1105–1123,
- 555 doi:10.5194/acp-17-1105-2017, 2017.
- Bagtasa, G.: Contribution of Tropical Cyclones to Rainfall in the Philippines, *J. Clim.*, 30(10), 3621–3633, doi:10.1175/JCLI-D-16-0150.1, 2017.
- Bagtasa, G., Cayetano, M. G. and Yuan, C.-S.: Seasonal variation and chemical characterization of PM_{2.5} in northwestern Philippines, *Atmospheric Chem. Phys.*, 18(7), 4965–4980,
- 560 doi:10.5194/acp-18-4965-2018, 2018.
- Bagtasa, G., Cayetano, M. G., Yuan, C.-S., Uchino, O., Sakai, T., Izumi, T., Morino, I., Nagai, T., Macatangay, R. C. and Velazco, V. A.: Long-range transport of aerosols from East and



- Southeast Asia to northern Philippines and its direct radiative forcing effect, *Atmos. Environ.*, 218, 117007, doi:10.1016/j.atmosenv.2019.117007, 2019.
- 565 Balkanski, Y. J., Jacob, D. J., Gardner, G. M., Graustein, W. C. and Turekian, K. K.: Transport and residence times of tropospheric aerosols inferred from a global three-dimensional simulation of ^{210}Pb , *J. Geophys. Res.*, 98(D11), 20573, doi:10.1029/93JD02456, 1993.
- Bhardwaj, P., Naja, M., Kumar, R. and Chandola, H. C.: Seasonal, interannual, and long-term variabilities in biomass burning activity over South Asia, *Environ. Sci. Pollut. Res.*, 23(5), 4397–4410, doi:10.1007/s11356-015-5629-6, 2016.
- 570 Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H. and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109(D14), D14203, doi:10.1029/2003JD003697, 2004.
- Braun, R. A., Aghdam, M. A., Bañaga, P. A., Betito, G., Cambaliza, M. O., Cruz, M. T.,
575 Lorenzo, G. R., MacDonald, A. B., Simpas, J. B., Stahl, C. and Sorooshian, A.: Long-range aerosol transport and impacts on size-resolved aerosol composition in Metro Manila, Philippines, *Atmospheric Chem. Phys.*, 20(4), 2387–2405, doi:10.5194/acp-20-2387-2020, 2020.
- Cappa, C. D., Lim, C. Y., Hagan, D. H., Coggon, M., Koss, A., Sekimoto, K., de Gouw, J.,
580 Onasch, T. B., Warneke, C. and Kroll, J. H.: Biomass-burning-derived particles from a wide variety of fuels – Part 2: Effects of photochemical aging on particle optical and chemical properties, *Atmospheric Chem. Phys.*, 20(14), 8511–8532, doi:https://doi.org/10.5194/acp-20-8511-2020, 2020.
- Center for International Earth Science Information Network: Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11, [online] Available from:
585 https://doi.org/10.7927/H49C6VHW (Accessed 12 March 2020), 2018.
- Chang, C.-P., Wang, Z., McBride, J. and Liu, C.-H.: Annual Cycle of Southeast Asia—Maritime Continent Rainfall and the Asymmetric Monsoon Transition, *J. Clim.*, 18(2), 287–301, doi:10.1175/JCLI-3257.1, 2005.
- Chen, S., Liu, B., Tan, X. and Wu, Y.: Inter-comparison of spatiotemporal features of
590 precipitation extremes within six daily precipitation products, *Clim. Dyn.*, 54(1–2), 1057–1076, doi:10.1007/s00382-019-05045-z, 2020a.
- Chen, T.-C., Tsay, J.-D. and Matsumoto, J.: Interannual Variation of the Summer Rainfall Center in the South China Sea, *J. Clim.*, 30(19), 7909–7931, doi:10.1175/JCLI-D-16-0889.1, 2017.
- Chen, Y., Cai, J., Wang, Z., Peng, C., Yao, X., Tian, M., Han, Y., Shi, G., Shi, Z., Liu, Y., Yang,
595 X., Zheng, M., Zhu, T., He, K., Zhang, Q. and Yang, F.: Simultaneous measurements of urban and rural particles in Beijing – Part 1: Chemical composition and mixing state, *Atmospheric Chem. Phys.*, 20(15), 9231–9247, doi:https://doi.org/10.5194/acp-20-9231-2020, 2020b.
- Cheng, F.-Y., Yang, Z.-M., Ou-Yang, C.-F. and Ngan, F.: A numerical study of the dependence of long-range transport of CO to a mountain station in Taiwan on synoptic weather patterns



- 600 during the Southeast Asia biomass-burning season, *Atmos. Environ.*, 78, 277–290,
doi:10.1016/j.atmosenv.2013.03.020, 2013.
- Cheung, H. C., Chou, C. C.-K., Lee, C. S. L., Kuo, W.-C. and Chang, S.-C.: Hygroscopic
properties and cloud condensation nuclei activity of atmospheric aerosols under the influences of
Asian continental outflow and new particle formation at a coastal site in eastern Asia,
605 *Atmospheric Chem. Phys.*, 20(10), 5911–5922, doi:10.5194/acp-20-5911-2020, 2020.
- Chien, F.-C., Hong, J.-S. and Kuo, Y.-H.: The Marine Boundary Layer Height over the Western
North Pacific Based on GPS Radio Occultation, Island Soundings, and Numerical Models,
Sensors, 19(1), 155, doi:10.3390/s19010155, 2019.
- Choi, Y., Kim, D., Cho, S. and Kim, T.-W.: Southeastern Yellow Sea as a sink for atmospheric
610 carbon dioxide, *Mar. Pollut. Bull.*, 149, 110550, doi:10.1016/j.marpolbul.2019.110550, 2019.
- Chuang, M.-T., Lee, C.-T., Chou, C. C.-K., Lin, N.-H., Sheu, G.-R., Wang, J.-L., Chang, S.-C.,
Wang, S.-H., Chi, K. H., Young, C.-Y., Huang, H., Chen, H.-W., Weng, G.-H., Lai, S.-Y., Hsu,
S.-P., Chang, Y.-J., Chang, J.-H. and Wu, X.-C.: Carbonaceous aerosols in the air masses
transported from Indochina to Taiwan: Long-term observation at Mt. Lulin, *Atmos. Environ.*, 89,
615 507–516, doi:10.1016/j.atmosenv.2013.11.066, 2014.
- Conte, L., Szopa, S., Séférian, R. and Bopp, L.: The oceanic cycle of carbon monoxide and its
emissions to the atmosphere, *Biogeosciences*, 16(4), 881–902, doi:10.5194/bg-16-881-2019,
2019.
- Cruz, F. T., Narisma, G. T., Villafuerte, M. Q., Cheng Chua, K. U. and Olaguera, L. M.: A
620 climatological analysis of the southwest monsoon rainfall in the Philippines, *Atmospheric Res.*,
122, 609–616, doi:10.1016/j.atmosres.2012.06.010, 2013.
- Dahutia, P., Pathak, B. and Bhuyan, P. K.: Vertical distribution of aerosols and clouds over
north-eastern South Asia: Aerosol-cloud interactions, *Atmos. Environ.*, 215, 116882,
doi:10.1016/j.atmosenv.2019.116882, 2019.
- 625 Danielson, J. J. and Gesch, D. B.: Global Multi-resolution Terrain Elevation Data 2010
(GMTED2010): U.S. Geological Survey Open-File Report 2011-1073, , 34, 2011.
- Davis, D. D., Crawford, J., Chen, G., Chameides, W., Liu, S., Bradshaw, J., Sandholm, S.,
Sachse, G., Gregory, G., Anderson, B., Barrick, J., Bachmeier, A., Collins, J., Browell, E., Blake,
D., Rowland, S., Kondo, Y., Singh, H., Talbot, R., Heikes, B., Merrill, J., Rodriguez, J. and
630 Newell, R. E.: Assessment of ozone photochemistry in the western North Pacific as inferred
from PEM-West A observations during the fall 1991, *J. Geophys. Res. Atmospheres*, 101(D1),
2111–2134, doi:10.1029/95JD02755, 1996.
- DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crouse, J., Wennberg, P.
O., Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D.,
635 Weinheimer, A. J., Montzka, D. D., Campos, T. and Jimenez, J. L.: Fast airborne aerosol size
and chemistry measurements above Mexico City and Central Mexico during the MILAGRO
campaign, *Atmospheric Chem. Phys.*, 8(14), 4027–4048, doi:10.5194/acp-8-4027-2008, 2008.



- 640 Di Girolamo, L., Reid, J. S., Holz, R., Tanelli, S., van den Heever, S. C., Narisma, G. T. T. and
Simpas, J. B. B.: Cloud and Aerosol Monsoonal Processes-Philippines Experiment (CAMP2Ex),
[online] Available from: https://espo.nasa.gov/CAMP2Ex_White_Paper (Accessed 31 July
2019), 2018.
- Dong, Z., Li, Z., Yu, X., Cribb, M., Li, X. and Dai, J.: Opposite long-term trends in aerosols
between low and high altitudes: a testimony to the aerosol–PBL feedback, *Atmospheric Chem.
Phys.*, 17(12), 7997–8009, doi:10.5194/acp-17-7997-2017, 2017.
- 645 Engling, G., He, J., Betha, R. and Balasubramanian, R.: Assessing the regional impact of
indonesian biomass burning emissions based on organic molecular tracers and chemical mass
balance modeling, *Atmospheric Chem. Phys.*, 14(15), 8043–8054, doi:10.5194/acp-14-8043-
2014, 2014.
- 650 Ervens, B.: Modeling the Processing of Aerosol and Trace Gases in Clouds and Fogs, *Chem.
Rev.*, 115(10), 4157–4198, doi:10.1021/cr5005887, 2015.
- Fioletov, V. E., McLinden, C. A., Krotkov, N., Li, C., Joiner, J., Theys, N., Carn, S. and Moran,
M. D.: A global catalogue of large SO₂ sources and emissions derived from the Ozone
Monitoring Instrument, *Atmospheric Chem. Phys.*, 16(18), 11497–11519, doi:10.5194/acp-16-
11497-2016, 2016.
- 655 Freitag, S., Clarke, A. D., Howell, S. G., Kapustin, V. N., Campos, T., Brekhovskikh, V. L. and
Zhou, J.: Combining airborne gas and aerosol measurements with HYSPLIT: a visualization tool
for simultaneous evaluation of air mass history and back trajectory consistency, *Atmospheric
Meas. Tech.*, 7(1), 107–128, doi:10.5194/amt-7-107-2014, 2014.
- 660 Fujii, Y., Tohno, S., Amil, N., Latif, M. T., Oda, M., Matsumoto, J. and Mizohata, A.: Annual
variations of carbonaceous PM_{2.5} in Malaysia: influence by Indonesian peatland fires,
Atmospheric Chem. Phys., 15(23), 13319–13329, doi:10.5194/acp-15-13319-2015, 2015.
- Ge, C., Wang, J. and Reid, J. S.: Mesoscale modeling of smoke transport over the Southeast
Asian Maritime Continent: coupling of smoke direct radiative effect below and above the low-
level clouds, *Atmospheric Chem. Phys.*, 14(1), 159–174, doi:10.5194/acp-14-159-2014, 2014.
- 665 Geng, X., Zhong, G., Li, J., Cheng, Z., Mo, Y., Mao, S., Su, T., Jiang, H., Ni, K. and Zhang, G.:
Molecular marker study of aerosols in the northern South China Sea: Impact of atmospheric
outflow from the Indo-China Peninsula and South China, *Atmos. Environ.*, 206, 225–236,
doi:10.1016/j.atmosenv.2019.02.033, 2019.
- 670 Govender, P. and Sivakumar, V.: Application of k-means and hierarchical clustering techniques
for analysis of air pollution: A review (1980–2019), *Atmospheric Pollut. Res.*, 11(1), 40–56,
doi:10.1016/j.apr.2019.09.009, 2020.
- Gras, J. L., Jensen, J. B., Okada, K., Ikegami, M., Zaizen, Y. and Makino, Y.: Some optical
properties of smoke aerosol in Indonesia and tropical Australia, *Geophys. Res. Lett.*, 26(10),
1393–1396, doi:10.1029/1999GL900275, 1999.



- 675 Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T. and Wanninkhof, R.: The oceanic sink for anthropogenic CO₂ from 1994 to 2007, *Science*, 363(6432), 1193–1199, doi:10.1126/science.aau5153, 2019.
- 680 Guttikunda, S. K., Thongboonchoo, N., Calori, G., Carmichael, G. R. and Streets, D. G.: Sulfur Deposition in Asia: Seasonal Behavior and Contributions from Various Energy Sectors, *Water, Air, Soil Pollut.*, 24, doi:10.1023/A:1011912902825, 2001.
- 685 Halliday, H. S., DiGangi, J. P., Choi, Y., Diskin, G. S., Pusede, S. E., Rana, M., Nowak, J. B., Knote, C., Ren, X., He, H., Dickerson, R. R. and Li, Z.: Using Short-Term CO/CO₂ Ratios to Assess Air Mass Differences Over the Korean Peninsula During KORUS-AQ, *J. Geophys. Res. Atmospheres*, 124(20), 10951–10972, doi:10.1029/2018JD029697, 2019.
- Hamid, E. Y., Kawasaki, Z.-I. and Mardiana, R.: Impact of the 1997-98 El Niño Event on lightning activity over Indonesia, *Geophys. Res. Lett.*, 28(1), 147–150, doi:10.1029/2000GL011374, 2001.
- 690 Hansen, J.: Efficacy of climate forcings, *J. Geophys. Res.*, 110(D18), D18104, doi:10.1029/2005JD005776, 2005.
- Harris, J. M., Draxler, R. R. and Oltmans, S. J.: Trajectory model sensitivity to differences in input data and vertical transport method: TRAJECTORY MODEL SENSITIVITY, *J. Geophys. Res. Atmospheres*, 110(D14), n/a-n/a, doi:10.1029/2004JD005750, 2005.
- 695 Hartigan, J. A. and Wong, M. A.: Algorithm AS 136: A K-Means Clustering Algorithm, *J. R. Stat. Soc. Ser. C Appl. Stat.*, 28(1), 100–108, doi:10.2307/2346830, 1979.
- Hatakeyama, S., Murano, K., Sakamaki, F., Mukai, H., Bandow, H. and Komazaki, Y.: Transport of Atmospheric Pollutants from East Asia, *Water, Air, Soil Pollut.*, 130(1/4), 373–378, doi:10.1023/A:1013877000169, 2001.
- 700 Hatakeyama, S., Takami, A., Sakamaki, F., Mukai, H., Sugimoto, N., Shimizu, A. and Bandow, H.: Aerial measurement of air pollutants and aerosols during 20-22 March 2001 over the East China Sea, *J. Geophys. Res. Atmospheres*, 109(D13), n/a-n/a, doi:10.1029/2003JD004271, 2004.
- 705 Hatakeyama, S., Hanaoka, S., Ikeda, K., Watanabe, I., Arakaki, T., Sadanaga, Y., Bandow, H., Kato, S., Kajii, Y., Sato, K., Shimizu, A. and Takami, A.: Aerial Observation of Aerosols Transported from East Asia — Chemical Composition of Aerosols and Layered Structure of an Air Mass over the East China Sea, *Aerosol Air Qual. Res.*, 11(5), 497–507, doi:10.4209/aaqr.2011.06.0076, 2011.
- 710 He, L., Zeng, Z., Pongetti, T. J., Wong, C., Liang, J., Gurney, K. R., Newman, S., Yadav, V., Verhulst, K., Miller, C. E., Duren, R., Frankenberg, C., Wennberg, P. O., Shia, R., Yung, Y. L. and Sander, S. P.: Atmospheric Methane Emissions Correlate With Natural Gas Consumption From Residential and Commercial Sectors in Los Angeles, *Geophys. Res. Lett.*, 46(14), 8563–8571, doi:10.1029/2019GL083400, 2019.



- Heald, C. L., Ridley, D. A., Kroll, J. H., Barrett, S. R. H., Cady-Pereira, K. E., Alvarado, M. J. and Holmes, C. D.: Contrasting the direct radiative effect and direct radiative forcing of aerosols, *Atmospheric Chem. Phys.*, 14(11), 5513–5527, doi:10.5194/acp-14-5513-2014, 2014.
- 715 Hecobian, A., Liu, Z., Hennigan, C. J., Huey, L. G., Jimenez, J. L., Cubison, M. J., Vay, S., Diskin, G. S., Sachse, G. W., Wisthaler, A., Mikoviny, T., Weinheimer, A. J., Liao, J., Knapp, D. J., Wennberg, P. O., Kürten, A., Crouse, J. D., Clair, J. St., Wang, Y. and Weber, R. J.: Comparison of chemical characteristics of 495 biomass burning plumes intercepted by the NASA DC-8 aircraft during the ARCTAS/CARB-2008 field campaign, *Atmospheric Chem. Phys.*, 11(24), 13325–13337, doi:10.5194/acp-11-13325-2011, 2011.
- 720 Hewitt, C. N., Lee, J. D., MacKenzie, A. R., Barkley, M. P., Carslaw, N., Carver, G. D., Chappell, N. A., Coe, H., Collier, C., Commane, R., Davies, F., Davison, B., DiCarlo, P., Di Marco, C. F., Dorsey, J. R., Edwards, P. M., Evans, M. J., Fowler, D., Furneaux, K. L., Gallagher, M., Guenther, A., Heard, D. E., Helfter, C., Hopkins, J., Ingham, T., Irwin, M., Jones, C., Karunaharan, A., Langford, B., Lewis, A. C., Lim, S. F., MacDonald, S. M., Mahajan, A. S., Malpass, S., McFiggans, G., Mills, G., Misztal, P., Moller, S., Monks, P. S., Nemitz, E., Nicolas-Perea, V., Oetjen, H., Oram, D. E., Palmer, P. I., Phillips, G. J., Pike, R., Plane, J. M. C., Pugh, T., Pyle, J. A., Reeves, C. E., Robinson, N. H., Stewart, D., Stone, D., Whalley, L. K. and Yin, X.: Overview: oxidant and particle photochemical processes above a south-east Asian tropical rainforest (the OP3 project): introduction, rationale, location characteristics and tools, *Atmospheric Chem. Phys.*, 10(1), 169–199, doi:10.5194/acp-10-169-2010, 2010.
- 725 Hilario, M. R. A., Cruz, M. T., Bañaga, P. A., Betito, G., Braun, R. A., Stahl, C., Cambaliza, M. O., Lorenzo, G. R., MacDonald, A. B., AzadiAghdam, M., Pabroa, P. C., Yee, J. R., Simpas, J. B. and Sorooshian, A.: Characterizing weekly cycles of particulate matter in a coastal megacity: The importance of a seasonal, size-resolved, and chemically-speciated analysis, *J. Geophys. Res. Atmospheres*, doi:10.1029/2020JD032614, 2020a.
- 735 Hilario, M. R. A., Olaguera, L. M., Narisma, G. T. and Matsumoto, J.: Diurnal characteristics of summer precipitation over Luzon Island, Philippines, *Asia-Pac. J. Atmospheric Sci.*, doi:10.1007/s13143-020-00214-1, 2020b.
- 740 Hilario, M. R. A., Cruz, M. T., Cambaliza, M. O. L., Reid, J. S., Xian, P., Simpas, J. B., Lagrosas, N. D., Uy, S. N. Y., Cliff, S. and Zhao, Y.: Investigating size-segregated sources of elemental composition of particulate matter in the South China Sea during the 2011 Vasco cruise, *Atmospheric Chem. Phys.*, 20(3), 1255–1276, doi:10.5194/acp-20-1255-2020, 2020c.
- 745 Hoell, J. M., Davis, D. D., Liu, S. C., Newell, R., Shipham, M., Akimoto, H., McNeal, R. J., Bendura, R. J. and Drewry, J. W.: Pacific Exploratory Mission-West A (PEM-West A): September–October 1991, *J. Geophys. Res. Atmospheres*, 101(D1), 1641–1653, doi:10.1029/95JD00622, 1996.
- 750 Hoell, J. M., Davis, D. D., Liu, S. C., Newell, R. E., Akimoto, H., McNeal, R. J. and Bendura, R. J.: The Pacific Exploratory Mission-West Phase B: February–March, 1994, *J. Geophys. Res. Atmospheres*, 102(D23), 28223–28239, doi:10.1029/97JD02581, 1997.



- Hong, Y., Hsu, K.-L., Sorooshian, S. and Gao, X.: Precipitation Estimation from Remotely Sensed Imagery Using an Artificial Neural Network Cloud Classification System, *J. Appl. Meteorol.*, 43(12), 1834–1853, doi:10.1175/JAM2173.1, 2004.
- 755 Honomichl, S. B. and Pan, L. L.: Transport from the Asian Summer Monsoon Anticyclone over the Western Pacific, *J. Geophys. Res. Atmospheres*, doi:10.1029/2019JD032094, 2020.
- Hou, A. Y., Kakar, R. K., Neeck, S., Azarbarzin, A. A., Kummerow, C. D., Kojima, M., Oki, R., Nakamura, K. and Iguchi, T.: The Global Precipitation Measurement Mission, *Bull. Am. Meteorol. Soc.*, 95(5), 701–722, doi:10.1175/BAMS-D-13-00164.1, 2014.
- 760 Huang, H.-Y., Wang, S.-H., Huang, W.-X., Lin, N.-H., Chuang, M.-T., Silva, A. M. da and Peng, C.-M.: Influence of Synoptic-Dynamic Meteorology on the Long-Range Transport of Indochina Biomass Burning Aerosols, *J. Geophys. Res. Atmospheres*, 125(3), e2019JD031260, doi:10.1029/2019JD031260, 2020.
- Huang, L., Lin, W., Li, F., Wang, Y. and Jiang, B.: Climate Impacts of the Biomass Burning in Indochina on Atmospheric Conditions over Southern China, *Aerosol Air Qual. Res.*, 9(12), 2707–2720, doi:10.4209/aaqr.2019.01.0028, 2019.
- 765 Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P. and Stocker, E. F.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales, *J. Hydrometeorol.*, 8(1), 38–55, doi:10.1175/JHM560.1, 2007.
- 770 Huffman, G. J., Bolvin, D. T., Nelkin, E. J. and Tan, J.: NASA Global Precipitation Measurement (GPM) Integrated Multi-satellite Retrievals for GPM (IMERG). Algorithm Theoretical Basis Document, version 6, [online] Available from: https://docsserver.gesdisc.eosdis.nasa.gov/public/project/GPM/IMERG_doc.06.pdf (Accessed 20 March 2020), 2019.
- 775 Hurst, D. F., Griffith, D. W. T. and Cook, G. D.: Trace gas emissions from biomass burning in tropical Australian savannas, *J. Geophys. Res.*, 99(D8), 16441, doi:10.1029/94JD00670, 1994.
- Ikegami, M., Okada, K., Zaizen, Y., Makino, Y., Jensen, J. B., Gras, J. L. and Harjanto, H.: Very high weight ratios of S/K in individual haze particles over Kalimantan during the 1997 Indonesian forest fires, *Atmos. Environ.*, 35(25), 4237–4243, doi:10.1016/S1352-2310(01)00247-3, 2001.
- 780 IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, [online] Available from: <https://www.ipcc.ch/report/ar5/syr/> (Accessed 2 January 2020), 2014.
- 785 Jacob, D. J., Crawford, J. H., Kleb, M. M., Connors, V. S., Bendura, R. J., Raper, J. L., Sachse, G. W., Gille, J. C., Emmons, L. and Heald, C. L.: Transport and Chemical Evolution over the Pacific (TRACE-P) aircraft mission: Design, execution, and first results, *J. Geophys. Res.*, 108(D20), 9000, doi:10.1029/2002JD003276, 2003.



- 790 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, B., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R. and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project., *Bull. Am. Meteorol. Soc.*, 77, 437–472, doi:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.
- 795 Kidd, C., Takayabu, Y. N., Skofronick-Jackson, G. M., Huffman, G. J., Braun, S. A., Kubota, T. and Turk, F. J.: The Global Precipitation Measurement (GPM) Mission, in *Satellite Precipitation Measurement: Volume 1*, edited by V. Levizzani, C. Kidd, D. B. Kirschbaum, C. D. Kummerow, K. Nakamura, and F. J. Turk, pp. 3–23, Springer International Publishing, Cham., 2020.
- Kim, J. H., Yum, S. S., Lee, Y.-G. and Choi, B.-C.: Ship measurements of submicron aerosol size distributions over the Yellow Sea and the East China Sea, *Atmospheric Res.*, 93(4), 700–714, doi:10.1016/j.atmosres.2009.02.011, 2009.
- 800 Kleinman, L. I., Daum, P. H., Lee, Y.-N., Senum, G. I., Springston, S. R., Wang, J., Berkowitz, C., Hubbe, J., Zaveri, R. A., Brechtel, F. J., Jayne, J., Onasch, T. B. and Worsnop, D.: Aircraft observations of aerosol composition and ageing in New England and Mid-Atlantic States during the summer 2002 New England Air Quality Study field campaign, *J. Geophys. Res.*, 112(D9), D09310, doi:10.1029/2006JD007786, 2007.
- 805 Koike, M., Kondo, Y., Kita, K., Takegawa, N., Masui, Y., Miyazaki, Y., Ko, M. W., Weinheimer, A. J., Flocke, F., Weber, R. J., Thornton, D. C., Sachse, G. W., Vay, S. A., Blake, D. R., Streets, D. G., Eisele, F. L., Sandholm, S. T., Singh, H. B. and Talbot, R. W.: Export of anthropogenic reactive nitrogen and sulfur compounds from the East Asia region in spring, *J. Geophys. Res.*, 108(D20), 8789, doi:10.1029/2002JD003284, 2003.
- 810 Kondo, Y., Morino, Y., Takegawa, N., Koike, M., Kita, K., Miyazaki, Y., Sachse, G. W., Vay, S. A., Avery, M. A., Flocke, F., Weinheimer, A. J., Eisele, F. L., Zondlo, M. A., Weber, R. J., Singh, H. B., Crawford, J. H., Blake, D. R., Fuelberg, H. E., Clarke, A. D., Talbot, R. W., Sandholm, S. T., Browell, E. V., Streets, D. G. and Liley, B.: Impacts of biomass burning in Southeast Asia on ozone and reactive nitrogen over the western Pacific in spring, *J. Geophys. Res.*, 109(D15), D15S12, doi:10.1029/2003JD004203, 2004.
- 815 Kondo, Y., Matsui, H., Moteki, N., Sahu, L., Takegawa, N., Kajino, M., Zhao, Y., Cubison, M. J., Jimenez, J. L., Vay, S., Diskin, G. S., Anderson, B., Wisthaler, A., Mikoviny, T., Fuelberg, H. E., Blake, D. R., Huey, G., Weinheimer, A. J., Knapp, D. J. and Brune, W. H.: Emissions of black carbon, organic, and inorganic aerosols from biomass burning in North America and Asia in 2008, *J. Geophys. Res.*, 116(D8), D08204, doi:10.1029/2010JD015152, 2011.
- Kreidenweis, S. M. and Asa-Awuku, A.: Aerosol Hygroscopicity: Particle Water Content and Its Role in Atmospheric Processes, in *Treatise on Geochemistry*, pp. 331–361, Elsevier., 2014.
- Kritz, M. A. and Rancher, J.: Circulation of Na, Cl, and Br in the tropical marine atmosphere, *J. Geophys. Res.*, 85(C3), 1633, doi:10.1029/JC085iC03p01633, 1980.



- 825 Krotkov, N. A., Li, C. and Leonard, P.: OMI/Aura Sulfur Dioxide (SO₂) Total Column L3 1 day Best Pixel in 0.25 degree x 0.25 degree V3, [online] Available from: <https://doi.org/10.5067/Aura/OMI/DATA3008> (Accessed 3 March 2020), 2015.
- Krupa, S. V. and Manning, W. J.: Atmospheric ozone: Formation and effects on vegetation, *Environ. Pollut.*, 50(1–2), 101–137, doi:10.1016/0269-7491(88)90187-X, 1988.
- 830 Kummerow, C. D., Randel, D. L., Kulie, M., Wang, N.-Y., Ferraro, R., Joseph Munchak, S. and Petkovic, V.: The Evolution of the Goddard Profiling Algorithm to a Fully Parametric Scheme, *J. Atmospheric Ocean. Technol.*, 32(12), 2265–2280, doi:10.1175/JTECH-D-15-0039.1, 2015.
- Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D. and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525(7569), 367–371, doi:10.1038/nature15371, 2015.
- 835 Levy, R. C., Mattoo, S., Munchak, L. A., Remer, L. A., Sayer, A. M., Patadia, F. and Hsu, N. C.: The Collection 6 MODIS aerosol products over land and ocean, *Atmospheric Meas. Tech.*, 6(11), 2989–3034, doi:10.5194/amt-6-2989-2013, 2013.
- Lin, C.-Y., Hsu, H. -m, Lee, Y. H., Kuo, C. H., Sheng, Y.-F. and Chu, D. A.: A new transport mechanism of biomass burning from Indochina as identified by modeling studies, *Atmospheric Chem. Phys.*, 9(20), 7901–7911, doi:https://doi.org/10.5194/acp-9-7901-2009, 2009.
- 840 Lin, I.-I., Chen, J.-P., Wong, G. T. F., Huang, C.-W. and Lien, C.-C.: Aerosol input to the South China Sea: Results from the MODerate Resolution Imaging Spectro-radiometer, the Quick Scatterometer, and the Measurements of Pollution in the Troposphere Sensor, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 54(14–15), 1589–1601, doi:10.1016/j.dsr2.2007.05.013, 2007.
- 845 Liu, Z.: Comparison of Integrated Multisatellite Retrievals for GPM (IMERG) and TRMM Multisatellite Precipitation Analysis (TMPA) Monthly Precipitation Products: Initial Results, *J. Hydrometeorol.*, 17(3), 777–790, doi:10.1175/JHM-D-15-0068.1, 2016.
- MacDonald, A. B., Dadashazar, H., Chuang, P. Y., Crosbie, E., Wang, H., Wang, Z., Jonsson, H., H., Flagan, R. C., Seinfeld, J. H. and Sorooshian, A.: Characteristic Vertical Profiles of Cloud Water Composition in Marine Stratocumulus Clouds and Relationships With Precipitation, *J. Geophys. Res. Atmospheres*, 123(7), 3704–3723, doi:10.1002/2017JD027900, 2018.
- 850 Maggioni, V., Meyers, P. C. and Robinson, M. D.: A Review of Merged High-Resolution Satellite Precipitation Product Accuracy during the Tropical Rainfall Measuring Mission (TRMM) Era, *J. Hydrometeorol.*, 17(4), 1101–1117, doi:10.1175/JHM-D-15-0190.1, 2016.
- 855 Mahmud, M. R., Hashim, M. and Reba, M. N. M.: How effective is the new generation of GPM satellite precipitation in characterizing the rainfall variability over Malaysia?, *Asia-Pac. J. Atmospheric Sci.*, 53(3), 375–384, doi:10.1007/s13143-017-0042-3, 2017.
- 860 Malm, W., Day, D., Kreidenweis, S., Collettjr, J., Carrico, C., Mcmeeking, G. and Lee, T.: Hygroscopic properties of an organic-laden aerosol, *Atmos. Environ.*, 39(27), 4969–4982, doi:10.1016/j.atmosenv.2005.05.014, 2005.



- Maloney, E. D. and Dickinson, M. J.: The Intraseasonal Oscillation and the Energetics of Summertime Tropical Western North Pacific Synoptic-Scale Disturbances, *J. ATMOSPHERIC Sci.*, 60, 16, 2003.
- 865 Mardi, A. H., Dadashazar, H., MacDonald, A. B., Braun, R. A., Crosbie, E., Xian, P., Thorsen, T. J., Coggon, M. M., Fenn, M. A., Ferrare, R. A., Hair, J. W., Woods, R. K., Jonsson, H. H., Flagan, R. C., Seinfeld, J. H. and Sorooshian, A.: Biomass Burning Plumes in the Vicinity of the California Coast: Airborne Characterization of Physicochemical Properties, Heating Rates, and Spatiotemporal Features, *J. Geophys. Res. Atmospheres*, 123(23), doi:10.1029/2018JD029134, 870 2018.
- Matsui, H., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Koike, M., Zhao, Y., Fuelberg, H. E., Sessions, W. R., Diskin, G., Anderson, B. E., Blake, D. R., Wisthaler, A., Cubison, M. J. and Jimenez, J. L.: Accumulation-mode aerosol number concentrations in the Arctic during the ARCTAS aircraft campaign: Long-range transport of polluted and clean air from the Asian 875 continent, *J. Geophys. Res.*, 116(D20), D20217, doi:10.1029/2011JD016189, 2011a.
- Matsui, H., Kondo, Y., Moteki, N., Takegawa, N., Sahu, L. K., Zhao, Y., Fuelberg, H. E., Sessions, W. R., Diskin, G., Blake, D. R., Wisthaler, A. and Koike, M.: Seasonal variation of the transport of black carbon aerosol from the Asian continent to the Arctic during the ARCTAS aircraft campaign, *J. Geophys. Res.*, 116(D5), D05202, doi:10.1029/2010JD015067, 2011b.
- 880 Matsui, H., Koike, M., Kondo, Y., Takami, A., Fast, J. D., Kanaya, Y. and Takigawa, M.: Volatility basis-set approach simulation of organic aerosol formation in East Asia: implications for anthropogenic–biogenic interaction and controllable amounts, *Atmospheric Chem. Phys.*, 14(18), 9513–9535, doi:10.5194/acp-14-9513-2014, 2014.
- 885 Matsumoto, J., Olaguera, L. M. P., Nguyen-Le, D., Kubota, H. and Villafuerte, M. Q.: Climatological seasonal changes of wind and rainfall in the Philippines, *Int. J. Climatol.*, joc.6492, doi:10.1002/joc.6492, 2020.
- 890 Matsumoto, K., Nagao, I., Tanaka, H., Miyaji, H., Iida, T. and Ikebe, Y.: Seasonal characteristics of organic and inorganic species and their size distributions in atmospheric aerosols over the northwest pacific ocean, *Atmos. Environ.*, 32(11), 1931–1946, doi:10.1016/S1352-2310(97)00499-8, 1998.
- Maudlin, L. C., Wang, Z., Jonsson, H. H. and Sorooshian, A.: Impact of wildfires on size-resolved aerosol composition at a coastal California site, *Atmos. Environ.*, 119, 59–68, doi:10.1016/j.atmosenv.2015.08.039, 2015.
- 895 McNaughton, C. S., Clarke, A. D., Howell, S. G., Pinkerton, M., Anderson, B., Thornhill, L., Hudgins, C., Winstead, E., Dibb, J. E., Scheuer, E. and Maring, H.: Results from the DC-8 Inlet Characterization Experiment (DICE): Airborne Versus Surface Sampling of Mineral Dust and Sea Salt Aerosols, *Aerosol Sci. Technol.*, 41(2), 136–159, doi:10.1080/02786820601118406, 2007.



- 900 Mishra, A. K., Koren, I. and Rudich, Y.: Effect of aerosol vertical distribution on aerosol-radiation interaction: A theoretical prospect, *Heliyon*, 1(2), e00036, doi:10.1016/j.heliyon.2015.e00036, 2015.
- Miyazaki, Y.: Synoptic-scale transport of reactive nitrogen over the western Pacific in spring, *J. Geophys. Res.*, 108(D20), 8788, doi:10.1029/2002JD003248, 2003.
- 905 Moteki, N. and Kondo, Y.: Effects of Mixing State on Black Carbon Measurements by Laser-Induced Incandescence, *Aerosol Sci. Technol.*, 41(4), 398–417, doi:10.1080/02786820701199728, 2007.
- Moteki, N. and Kondo, Y.: Dependence of Laser-Induced Incandescence on Physical Properties of Black Carbon Aerosols: Measurements and Theoretical Interpretation, *Aerosol Sci. Technol.*, 44(8), 663–675, doi:10.1080/02786826.2010.484450, 2010.
- 910 Moteki, N., Kondo, Y., Oshima, N., Takegawa, N., Koike, M., Kita, K., Matsui, H. and Kajino, M.: Size dependence of wet removal of black carbon aerosols during transport from the boundary layer to the free troposphere, *Geophys. Res. Lett.*, 39(13), n/a-n/a, doi:10.1029/2012GL052034, 2012.
- 915 Murphy, D. M., Cziczo, D. J., Hudson, P. K., Thomson, D. S., Wilson, J. C., Kojima, T. and Buseck, P. R.: Particle Generation and Resuspension in Aircraft Inlets when Flying in Clouds, *Aerosol Sci. Technol.*, 38(4), 401–409, doi:10.1080/02786820490443094, 2004.
- Neu, J. L. and Prather, M. J.: Toward a more physical representation of precipitation scavenging in global chemistry models: cloud overlap and ice physics and their impact on tropospheric ozone, *Atmospheric Chem. Phys.*, 12(7), 3289–3310, doi:10.5194/acp-12-3289-2012, 2012.
- 920 Nguyen, L. S. P., Huang, H.-Y., Lei, T. L., Bui, T. T., Wang, S.-H., Chi, K. H., Sheu, G.-R., Lee, C.-T., Ou-Yang, C.-F. and Lin, N.-H.: Characterizing a landmark biomass-burning event and its implication for aging processes during long-range transport, *Atmos. Environ.*, 241, 117766, doi:10.1016/j.atmosenv.2020.117766, 2020.
- 925 Nguyen, P., Ombadi, M., Sorooshian, S., Hsu, K., AghaKouchak, A., Braithwaite, D., Ashouri, H. and Thorstensen, A. R.: The PERSIANN family of global satellite precipitation data: a review and evaluation of products, *Hydrol. Earth Syst. Sci.*, 22(11), 5801–5816, doi:10.5194/hess-22-5801-2018, 2018.
- 930 Nguyen, P., Shearer, E. J., Tran, H., Ombadi, M., Hayatbini, N., Palacios, T., Huynh, P., Braithwaite, D., Updegraff, G., Hsu, K., Kuligowski, B., Logan, W. S. and Sorooshian, S.: The CHRS Data Portal, an easily accessible public repository for PERSIANN global satellite precipitation data, *Sci. Data*, 6(1), 180296, doi:10.1038/sdata.2018.296, 2019.
- Olson, W. S.: GPM Combined Radar-Radiometer Precipitation Algorithm Theoretical Basis Document (Version 5), , 68, 2018.
- 935 Oshima, N., Kondo, Y., Moteki, N., Takegawa, N., Koike, M., Kita, K., Matsui, H., Kajino, M., Nakamura, H., Jung, J. S. and Kim, Y. J.: Wet removal of black carbon in Asian outflow:



- Aerosol Radiative Forcing in East Asia (A-FORCE) aircraft campaign, *J. Geophys. Res. Atmospheres*, 117(D3), n/a-n/a, doi:10.1029/2011JD016552, 2012.
- 940 Oshima, N., Koike, M., Kondo, Y., Nakamura, H., Moteki, N., Matsui, H., Takegawa, N. and Kita, K.: Vertical transport mechanisms of black carbon over East Asia in spring during the A-FORCE aircraft campaign, *J. Geophys. Res. Atmospheres*, 118(23), 13,175-13,198, doi:10.1002/2013JD020262, 2013.
- 945 Pani, S. K., Ou-Yang, C.-F., Wang, S.-H., Ogren, J. A., Sheridan, P. J., Sheu, G.-R. and Lin, N.-H.: Relationship between long-range transported atmospheric black carbon and carbon monoxide at a high-altitude background station in East Asia, *Atmos. Environ.*, 210, 86–99, doi:10.1016/j.atmosenv.2019.04.053, 2019.
- Park, R. J.: Export efficiency of black carbon aerosol in continental outflow: Global implications, *J. Geophys. Res.*, 110(D11), D11205, doi:10.1029/2004JD005432, 2005.
- 950 Pérez, I. A., Sánchez, M. L., García, M. A. and Pardo, N.: Boundaries of air mass trajectory clustering: key points and applications, *Int. J. Environ. Sci. Technol.*, 14(3), 653–662, doi:10.1007/s13762-016-1140-y, 2017.
- Pósfai, M., Simonics, R., Li, J., Hobbs, P. V. and Buseck, P. R.: Individual aerosol particles from biomass burning in southern Africa: 1. Compositions and size distributions of carbonaceous particles, *J. Geophys. Res. Atmospheres*, 108(D13), n/a-n/a, doi:10.1029/2002JD002291, 2003.
- 955 Radzi bin Abas, M., Oros, D. R. and Simoneit, B. R. T.: Biomass burning as the main source of organic aerosol particulate matter in Malaysia during haze episodes, *Chemosphere*, 55(8), 1089–1095, doi:10.1016/j.chemosphere.2004.02.002, 2004.
- Reid, J. S., Hobbs, P. V., Ferek, R. J., Blake, D. R., Martins, J. V., Dunlap, M. R. and Liou, S. C.: Physical, chemical, and optical properties of regional hazes dominated by smoke in Brazil, *J. Geophys. Res. Atmospheres*, 103(D24), 32059–32080, doi:10.1029/98JD00458, 1998.
- 960 Reid, J. S., Koppmann, R., Eck, T. F. and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmos Chem Phys*, 27, 2005.
- 965 Reid, J. S., Xian, P., Hyer, E. J., Flatau, M. K., Ramirez, E. M., Turk, F. J., Sampson, C. R., Zhang, C., Fukada, E. M. and Maloney, E. D.: Multi-scale meteorological conceptual analysis of observed active fire hotspot activity and smoke optical depth in the Maritime Continent, *Atmospheric Chem. Phys.*, 12(4), 2117–2147, doi:10.5194/acp-12-2117-2012, 2012.
- 970 Reid, J. S., Hyer, E. J., Johnson, R. S., Holben, B. N., Yokelson, R. J., Zhang, J., Campbell, J. R., Christopher, S. A., Di Girolamo, L., Giglio, L., Holz, R. E., Kearney, C., Miettinen, J., Reid, E. A., Turk, F. J., Wang, J., Xian, P., Zhao, G., Balasubramanian, R., Chew, B. N., Janjai, S., Lagrosas, N., Lestari, P., Lin, N.-H., Mahmud, M., Nguyen, A. X., Norris, B., Oanh, N. T. K., Oo, M., Salinas, S. V., Welton, E. J. and Liew, S. C.: Observing and understanding the Southeast Asian aerosol system by remote sensing: An initial review and analysis for the Seven Southeast



Asian Studies (7SEAS) program, *Atmospheric Res.*, 122, 403–468,
doi:10.1016/j.atmosres.2012.06.005, 2013.

975 Reid, J. S., Lagrosas, N. D., Jonsson, H. H., Reid, E. A., Sessions, W. R., Simpas, J. B., Uy, S.
N., Boyd, T. J., Atwood, S. A., Blake, D. R., Campbell, J. R., Cliff, S. S., Holben, B. N., Holz,
R. E., Hyer, E. J., Lynch, P., Meinardi, S., Posselt, D. J., Richardson, K. A., Salinas, S. V.,
Smirnov, A., Wang, Q., Yu, L. and Zhang, J.: Observations of the temporal variability in aerosol
980 properties and their relationships to meteorology in the summer monsoonal South China Sea/East
Sea: the scale-dependent role of monsoonal flows, the Madden–Julian Oscillation, tropical
cyclones, squall lines and cold pools, *Atmospheric Chem. Phys.*, 15(4), 1745–1768,
doi:10.5194/acp-15-1745-2015, 2015.

Reid, J. S., Lagrosas, N. D., Jonsson, H. H., Reid, E. A., Atwood, S. A., Boyd, T. J., Ghate, V.
P., Xian, P., Posselt, D. J., Simpas, J. B., Uy, S. N., Zaiger, K., Blake, D. R., Bucholtz, A.,
985 Campbell, J. R., Chew, B. N., Cliff, S. S., Holben, B. N., Holz, R. E., Hyer, E. J., Kreidenweis,
S. M., Kuciauskas, A. P., Lolli, S., Oo, M., Perry, K. D., Salinas, S. V., Sessions, W. R.,
Smirnov, A., Walker, A. L., Wang, Q., Yu, L., Zhang, J. and Zhao, Y.: Aerosol meteorology of
Maritime Continent for the 2012 7SEAS southwest monsoon intensive study – Part 2: Philippine
receptor observations of fine-scale aerosol behavior, *Atmospheric Chem. Phys.*, 16(22), 14057–
990 14078, doi:10.5194/acp-16-14057-2016, 2016a.

Reid, J. S., Xian, P., Holben, B. N., Hyer, E. J., Reid, E. A., Salinas, S. V., Zhang, J., Campbell,
J. R., Chew, B. N., Holz, R. E., Kuciauskas, A. P., Lagrosas, N., Posselt, D. J., Sampson, C. R.,
Walker, A. L., Welton, E. J. and Zhang, C.: Aerosol meteorology of the Maritime Continent for
995 the 2012 7SEAS southwest monsoon intensive study – Part 1: regional-scale phenomena,
Atmospheric Chem. Phys., 16(22), 14041–14056, doi:10.5194/acp-16-14041-2016, 2016b.

Robinson, N. H., Newton, H. M., Allan, J. D., Irwin, M., Hamilton, J. F., Flynn, M., Bower, K.
N., Williams, P. I., Mills, G., Reeves, C. E., McFiggans, G. and Coe, H.: Source attribution of
Bornean air masses by back trajectory analysis during the OP3 project, *Atmospheric Chem.*
1000 *Phys.*, 11(18), 9605–9630, doi:10.5194/acp-11-9605-2011, 2011.

Robinson, N. H., Allan, J. D., Trembath, J. A., Rosenberg, P. D., Allen, G. and Coe, H.: The
lofting of Western Pacific regional aerosol by island thermodynamics as observed around
Borneo, *Atmospheric Chem. Phys.*, 12(13), 5963–5983, doi:10.5194/acp-12-5963-2012, 2012.

Rolph, G., Stein, A. and Stunder, B.: Real-time Environmental Applications and Display
sYstem: READY, *Environ. Model. Softw.*, 95, 210–228, doi:10.1016/j.envsoft.2017.06.025,
1005 2017.

Rosenfeld, D.: TRMM observed first direct evidence of smoke from forest fires inhibiting
rainfall, *Geophys. Res. Lett.*, 26(20), 3105–3108, doi:10.1029/1999GL006066, 1999.

Ross, A. D., Holz, R. E., Quinn, G., Reid, J. S., Xian, P., Turk, F. J. and Posselt, D. J.: Exploring
the First Aerosol Indirect Effect over the MaritimeContinent Using a Ten-Year Collocated
1010 MODIS, CALIOP, andModel Dataset, preprint, *Aerosols/Remote Sensing/Troposphere/Physics*
(physical properties and processes)., 2018.



- Santoso, M., Lestiani, D. D., Mukhtar, R., Hamonangan, E., Syafrul, H., Markwitz, A. and Hopke, P. K.: Preliminary study of the sources of ambient air pollution in Serpong, Indonesia, *Atmospheric Pollut. Res.*, 2(2), 190–196, doi:10.5094/APR.2011.024, 2011.
- 1015 Sarkar, S., Fan, W. H., Jia, S., Blake, D. R., Reid, J. S., Lestari, P. and Yu, L. E.: A quantitative assessment of distributions and sources of tropospheric halocarbons measured in Singapore, *Sci. Total Environ.*, 619–620, 528–544, doi:10.1016/j.scitotenv.2017.11.087, 2018.
- Saxena, P., Hildemann, L. M., McMurry, P. H. and Seinfeld, J. H.: Organics alter hygroscopic behavior of atmospheric particles, *J. Geophys. Res.*, 100(D9), 18755, doi:10.1029/95JD01835, 1995.
- 1020 Siems, S. T., Hess, G., Suhre, K., Businger, S. and Draxler, R. R.: The impact of wind shear on observed and simulated trajectories during the ACE-1 Lagrangian experiments., 2000.
- Sievering, H., Van Valin, C. C., Barrett, E. W. and Pueschel, R. F.: Cloud scavenging of aerosol sulfur: Two case studies, *Atmospheric Environ.* 1967, 18(12), 2685–2690, doi:10.1016/0004-6981(84)90333-0, 1984.
- 1025 Skofronick-Jackson, G., Kirschbaum, D., Petersen, W., Huffman, G., Kidd, C., Stocker, E. and Kakar, R.: The Global Precipitation Measurement (GPM) mission’s scientific achievements and societal contributions: reviewing four years of advanced rain and snow observations, *Q. J. R. Meteorol. Soc.*, 144(S1), 27–48, doi:10.1002/qj.3313, 2018.
- 1030 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A. and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmospheric Chem. Phys.*, 11(3), 1101–1116, doi:10.5194/acp-11-1101-2011, 2011.
- 1035 Sorooshian, A., Varutbangkul, V., Brechtel, F. J., Ervens, B., Feingold, G., Bahreini, R., Murphy, S. M., Holloway, J. S., Atlas, E. L., Buzorius, G., Jonsson, H., Flagan, R. C. and Seinfeld, J. H.: Oxalic acid in clear and cloudy atmospheres: Analysis of data from International Consortium for Atmospheric Research on Transport and Transformation 2004: OXALIC ACID DATA ANALYSIS FROM ICARTT, *J. Geophys. Res. Atmospheres*, 111(D23), doi:10.1029/2005JD006880, 2006.
- 1040 Sorooshian, A., Lu, M.-L., Brechtel, F. J., Jonsson, H., Feingold, G., Flagan, R. C. and Seinfeld, J. H.: On the Source of Organic Acid Aerosol Layers above Clouds, *Environ. Sci. Technol.*, 41(13), 4647–4654, doi:10.1021/es0630442, 2007.
- 1045 Sorooshian, A., Padró, L. T., Nenes, A., Feingold, G., McComiskey, A., Hersey, S. P., Gates, H., Jonsson, H. H., Miller, S. D., Stephens, G. L., Flagan, R. C. and Seinfeld, J. H.: On the link between ocean biota emissions, aerosol, and maritime clouds: Airborne, ground, and satellite measurements off the coast of California: OCEAN EMISSIONS, AEROSOL, AND CLOUDS, *Glob. Biogeochem. Cycles*, 23(4), n/a-n/a, doi:10.1029/2009GB003464, 2009.
- Sorooshian, A., Anderson, B., Bauer, S. E., Braun, R. A., Cairns, B., Crosbie, E., Dadashazar, H., Diskin, G., Ferrare, R., Flagan, R. C., Hair, J., Hostetler, C., Jonsson, H. H., Kleb, M. M., Liu, H., MacDonald, A. B., McComiskey, A., Moore, R., Painemal, D., Russell, L. M., Seinfeld,



- 1050 J. H., Shook, M., Smith, W. L., Thornhill, K., Tselioudis, G., Wang, H., Zeng, X., Zhang, B., Ziemba, L. and Zuidema, P.: Aerosol–Cloud–Meteorology Interaction Airborne Field Investigations: Using Lessons Learned from the U.S. West Coast in the Design of ACTIVATE off the U.S. East Coast, *Bull. Am. Meteorol. Soc.*, 100(8), 1511–1528, doi:10.1175/BAMS-D-18-0100.1, 2019.
- 1055 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: NOAA’s HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bull. Am. Meteorol. Soc.*, 96(12), 2059–2077, doi:10.1175/BAMS-D-14-00110.1, 2015.
- 1060 Stohl, A., Eckhardt, S., Forster, C., James, P., Spichtinger, N. and Seibert, P.: A replacement for simple back trajectory calculations in the interpretation of atmospheric trace substance measurements, *Atmos. Environ.*, 36(29), 4635–4648, doi:10.1016/S1352-2310(02)00416-8, 2002.
- Streets, D. G., Carmichael, G. R. and Arndt, R. L.: Sulfur dioxide emissions and sulfur deposition from international shipping in Asian waters, *Atmos. Environ.*, 31(10), 1573–1582, doi:10.1016/S1352-2310(96)00204-X, 1997.
- 1065 Streets, D. G., Guttikunda, S. K. and Carmichael, G. R.: The growing contribution of sulfur emissions from ships in Asian waters, 1988–1995, *Atmos. Environ.*, 34(26), 4425–4439, doi:10.1016/S1352-2310(00)00175-8, 2000.
- 1070 Svenningsson, B., Rissler, J., Swietlicki, E., Mircea, M., Bilde, M., Facchini, M. C., Decesari, S., Fuzzi, S., Zhou, J., Mønster, J. and Rosenørn, T.: Hygroscopic growth and critical supersaturations for mixed aerosol particles of inorganic and organic compounds of atmospheric relevance, *Atmospheric Chem. Phys.*, 6(7), 1937–1952, doi:10.5194/acp-6-1937-2006, 2006.
- Takahashi, H. G., Fujinami, H., Yasunari, T. and Matsumoto, J.: Diurnal rainfall pattern observed by Tropical Rainfall Measuring Mission Precipitation Radar (TRMM-PR) around the Indochina peninsula, *J. Geophys. Res.*, 115(D7), D07109, doi:10.1029/2009JD012155, 2010.
- 1075 Talbot, R. W., Dibb, J. E., Lefer, B. L., Bradshaw, J. D., Sandholm, S. T., Blake, D. R., Blake, N. J., Sachse, G. W., Collins, J. E., Heikes, B. G., Merrill, J. T., Gregory, G. L., Anderson, B. E., Singh, H. B., Thornton, D. C., Bandy, A. R. and Pueschel, R. F.: Chemical characteristics of continental outflow from Asia to the troposphere over the western Pacific Ocean during February–March 1994: Results from PEM-West B, *J. Geophys. Res. Atmospheres*, 102(D23), 28255–28274, doi:10.1029/96JD02340, 1997.
- 1080 Tan, M. L. and Santo, H.: Comparison of GPM IMERG, TMPA 3B42 and PERSIANN-CDR satellite precipitation products over Malaysia, *Atmospheric Res.*, 202, 63–76, doi:10.1016/j.atmosres.2017.11.006, 2018.
- 1085 Tan, S.-C., Shi, G.-Y. and Wang, H.: Long-range transport of spring dust storms in Inner Mongolia and impact on the China seas, *Atmos. Environ.*, 46, 299–308, doi:10.1016/j.atmosenv.2011.09.058, 2012.



- 1090 Taubman, B. F., Hains, J. C., Thompson, A. M., Marufu, L. T., Doddridge, B. G., Stehr, J. W., Piety, C. A. and Dickerson, R. R.: Aircraft vertical profiles of trace gas and aerosol pollution over the mid-Atlantic United States: Statistics and meteorological cluster analysis, *J. Geophys. Res. Atmospheres*, 111(D10), n/a-n/a, doi:10.1029/2005JD006196, 2006.
- Theodoritsi, G. N., Posner, L. N., Robinson, A. L., Yarwood, G., Koo, B., Morris, R., Mavko, M., Moore, T. and Pandis, S. N.: Biomass burning organic aerosol from prescribed burning and other activities in the United States, *Atmos. Environ.*, 241, 117753, doi:10.1016/j.atmosenv.2020.117753, 2020.
- 1095 Thornton, D. C., Bandy, A. R., Blomquist, B. W., Driedger, A. R. and Wade, T. P.: Sulfur dioxide distribution over the Pacific Ocean 1991-1996, *J. Geophys. Res. Atmospheres*, 104(D5), 5845–5854, doi:10.1029/1998JD100048, 1999.
- 1100 Tohjima, Y., Kubo, M., Minejima, C., Mukai, H., Tanimoto, H., Ganshin, A., Maksyutov, S., Katsumata, K., Machida, T. and Kita, K.: Temporal changes in the emissions of CH₄ and CO from China estimated from CH₄ / CO₂ and CO / CO₂ correlations observed at Hateruma Island, *Atmospheric Chem. Phys.*, 14(3), 1663–1677, doi:10.5194/acp-14-1663-2014, 2014.
- Tufféry, S.: *Data Mining and Statistics for Decision Making: Tufféry/Data Mining and Statistics for Decision Making*, John Wiley & Sons, Ltd, Chichester, UK., 2011.
- 1105 Ueda, S., Miura, K., Kawata, R., Furutani, H., Uematsu, M., Omori, Y. and Tanimoto, H.: Number–size distribution of aerosol particles and new particle formation events in tropical and subtropical Pacific Oceans, *Atmos. Environ.*, 142, 324–339, doi:10.1016/j.atmosenv.2016.07.055, 2016.
- 1110 Umezawa, T., Goto, D., Aoki, S., Ishijima, K., Patra, P. K., Sugawara, S., Morimoto, S. and Nakazawa, T.: Variations of tropospheric methane over Japan during 1988–2010, *Tellus B Chem. Phys. Meteorol.*, 66(1), 23837, doi:10.3402/tellusb.v66.23837, 2014.
- Uno, I., Ohara, T. and Murano, K.: Simulated Acidic Aerosol Long-Range Transport and Deposition over East Asia - Role of Synoptic Scale Weather Systems, in *Air Pollution Modeling and Its Application XII*, Springer US. [online] Available from: https://doi.org/10.1007/978-1-4757-9128-0_20 (Accessed 16 April 2020), 1998.
- 1115 Wang, F., Chen, Y., Meng, X., Fu, J. and Wang, B.: The contribution of anthropogenic sources to the aerosols over East China Sea, *Atmos. Environ.*, 127, 22–33, doi:10.1016/j.atmosenv.2015.12.002, 2016.
- 1120 Wang, H., Easter, R. C., Rasch, P. J., Wang, M., Liu, X., Ghan, S. J., Qian, Y., Yoon, J.-H., Ma, P.-L. and Vinoj, V.: Sensitivity of remote aerosol distributions to representation of cloud–aerosol interactions in a global climate model, *Geosci. Model Dev.*, 6(3), 765–782, doi:10.5194/gmd-6-765-2013, 2013a.
- Wang, J., Ge, C., Yang, Z., Hyer, E. J., Reid, J. S., Chew, B.-N., Mahmud, M., Zhang, Y. and Zhang, M.: Mesoscale modeling of smoke transport over the Southeast Asian Maritime



1125 Continent: Interplay of sea breeze, trade wind, typhoon, and topography, *Atmospheric Res.*, 122, 486–503, doi:10.1016/j.atmosres.2012.05.009, 2013b.

Wang, J., Pikridas, M., Spielman, S. R. and Pinterich, T.: A fast integrated mobility spectrometer for rapid measurement of sub-micrometer aerosol size distribution, Part I: Design and model evaluation, *J. Aerosol Sci.*, 108, 44–55, doi:10.1016/j.jaerosci.2017.02.012, 2017a.

1130 Wang, J., Pikridas, M., Pinterich, T., Spielman, S. R., Tsang, T., McMahon, A. and Smith, S.: A Fast Integrated Mobility Spectrometer for rapid measurement of sub-micrometer aerosol size distribution, Part II: Experimental characterization, *J. Aerosol Sci.*, 113, 119–129, doi:10.1016/j.jaerosci.2017.05.001, 2017b.

1135 Wang, N., Jing, B., Wang, P., Wang, Z., Li, J., Pang, S., Zhang, Y. and Ge, M.: Hygroscopicity and Compositional Evolution of Atmospheric Aerosols Containing Water-Soluble Carboxylic Acid Salts and Ammonium Sulfate: Influence of Ammonium Depletion, *Environ. Sci. Technol.*, 53(11), 6225–6234, doi:10.1021/acs.est.8b07052, 2019.

1140 Wang, X., Jing, B., Tan, F., Ma, J., Zhang, Y. and Ge, M.: Hygroscopic behavior and chemical composition evolution of internally mixed aerosols composed of oxalic acid and ammonium sulfate, *Atmospheric Chem. Phys.*, 17(20), 12797–12812, doi:10.5194/acp-17-12797-2017, 2017c.

Wang, Y., Zhuang, G., Chen, S., An, Z. and Zheng, A.: Characteristics and sources of formic, acetic and oxalic acids in PM_{2.5} and PM₁₀ aerosols in Beijing, China, *Atmospheric Res.*, 84(2), 169–181, doi:10.1016/j.atmosres.2006.07.001, 2007.

1145 Wang, Y., Pinterich, T. and Wang, J.: Rapid measurement of sub-micrometer aerosol size distribution using a fast integrated mobility spectrometer, *J. Aerosol Sci.*, 121, 12–20, doi:10.1016/j.jaerosci.2018.03.006, 2018a.

Wang, Z., Jing, B., Shi, X., Tong, S., Wang, W. and Ge, M.: Importance of water-soluble organic acid on the hygroscopicity of nitrate, *Atmos. Environ.*, 190, 65–73, doi:10.1016/j.atmosenv.2018.07.010, 2018b.

1150 Ward, J. H.: Hierarchical Grouping to Optimize an Objective Function, *J. Am. Stat. Assoc.*, 58(301), 236–244, doi:10.1080/01621459.1963.10500845, 1963.

1155 Williamson, C. J., Kupc, A., Axisa, D., Bilsback, K. R., Bui, T., Campuzano-Jost, P., Dollner, M., Froyd, K. D., Hodshire, A. L., Jimenez, J. L., Kodros, J. K., Luo, G., Murphy, D. M., Nault, B. A., Ray, E. A., Weinzierl, B., Wilson, J. C., Yu, F., Yu, P., Pierce, J. R. and Brock, C. A.: A large source of cloud condensation nuclei from new particle formation in the tropics, *Nature*, 574(7778), 399–403, doi:10.1038/s41586-019-1638-9, 2019.

1160 Wonaschuetz, A., Sorooshian, A., Ervens, B., Chuang, P. Y., Feingold, G., Murphy, S. M., de Gouw, J., Warneke, C. and Jonsson, H. H.: Aerosol and gas re-distribution by shallow cumulus clouds: An investigation using airborne measurements, *J. Geophys. Res. Atmospheres*, 117(D17), n/a-n/a, doi:10.1029/2012JD018089, 2012.



- Xia, L., Zhang, G., Zhan, M., Li, B. and Kong, P.: Seasonal variations of atmospheric CH₄ at Jingdezhen station in Central China: Understanding the regional transport and its correlation with CO₂ and CO, *Atmospheric Res.*, 241, 104982, doi:10.1016/j.atmosres.2020.104982, 2020.
- 1165 Xian, P., Reid, J. S., Atwood, S. A., Johnson, R. S., Hyer, E. J., Westphal, D. L. and Sessions, W.: Smoke aerosol transport patterns over the Maritime Continent, *Atmospheric Res.*, 122, 469–485, doi:10.1016/j.atmosres.2012.05.006, 2013.
- Yadav, I. C., Linthoingambi Devi, N., Li, J., Syed, J. H., Zhang, G. and Watanabe, H.: Biomass burning in Indo-China peninsula and its impacts on regional air quality and global climate change—a review, *Environ. Pollut.*, 227, 414–427, doi:10.1016/j.envpol.2017.04.085, 2017.
- 1170 Yang, G.-P., Zhang, H.-H., Su, L.-P. and Zhou, L.-M.: Biogenic emission of dimethylsulfide (DMS) from the North Yellow Sea, China and its contribution to sulfate in aerosol during summer, *Atmos. Environ.*, 43(13), 2196–2203, doi:10.1016/j.atmosenv.2009.01.011, 2009.
- 1175 Yang, Y., Lin, Q., Fu, Y., Lian, X., Jiang, F., Peng, L., Zhang, G., Li, L., Chen, D., Li, M., Ou, J., Bi, X., Wang, X. and Sheng, G.: Stage-resolved in-cloud scavenging of submicron and BC-containing particles: A case study, *Atmos. Environ.*, 117883, doi:10.1016/j.atmosenv.2020.117883, 2020.
- 1180 Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. A., Anderson, T. L., Balkanski, Y., Bellouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T. and Zhou, M.: A review of measurement-based assessments of the aerosol direct radiative effect and forcing, *Atmospheric Chem. Phys.*, 6(3), 613–666, doi:10.5194/acp-6-613-2006, 2006.
- 1185 Yu, X., Li, D., Li, D., Zhang, G., Zhou, H., Li, S., Song, W., Zhang, Y., Bi, X., Yu, J. and Wang, X.: Enhanced wet deposition of water-soluble organic nitrogen during the harvest season: influence of biomass burning and in-cloud scavenging, *J. Geophys. Res. Atmospheres*, doi:10.1029/2020JD032699, 2020.
- Yuan, T., Remer, L. A., Pickering, K. E. and Yu, H.: Observational evidence of aerosol enhancement of lightning activity and convective invigoration: aerosol enhancement of lightning, *Geophys. Res. Lett.*, 38(4), doi:10.1029/2010GL046052, 2011.
- 1190 Yusuf, A. A. and Francisco, H.: Climate change vulnerability mapping for Southeast Asia, Economy and Environment Program for Southeast Asia (EEPSEA) report. [online] Available from: <http://www.eepsea.net/> (Accessed 1 January 2019), 2009.
- 1195 Zhang, H., Di, B., Liu, D., Li, J. and Zhan, Y.: Spatiotemporal distributions of ambient SO₂ across China based on satellite retrievals and ground observations: Substantial decrease in human exposure during 2013–2016, *Environ. Res.*, 179, 108795, doi:10.1016/j.envres.2019.108795, 2019.
- Zhang, Q., Jiang, X., Tong, D., Davis, S. J., Zhao, H., Geng, G., Feng, T., Zheng, B., Lu, Z., Streets, D. G., Ni, R., Brauer, M., van Donkelaar, A., Martin, R. V., Huo, H., Liu, Z., Pan, D., Kan, H., Yan, Y., Lin, J., He, K. and Guan, D.: Transboundary health impacts of transported



1200 global air pollution and international trade, *Nature*, 543(7647), 705–709,
doi:10.1038/nature21712, 2017.

Zhou, S., Wang, Z., Gao, R., Xue, L., Yuan, C., Wang, T., Gao, X., Wang, X., Nie, W., Xu, Z.,
Zhang, Q. and Wang, W.: Formation of secondary organic carbon and long-range transport of
carbonaceous aerosols at Mount Heng in South China, *Atmos. Environ.*, 63, 203–212,
doi:10.1016/j.atmosenv.2012.09.021, 2012.

1205



1206 **Table 1: Statistics for accumulated precipitation along trajectories (APT, mm) for Peninsular Southeast Asia (PSEA), Maritime Continent (MC), East Asia (EA), and West**
1207 **Pacific (WP), calculated with IMERG, TMPA, and PERSIANN-CDR. Median values are presented along with the 25th and 75th percentiles provided in parentheses.**

	IMERG	TMPA	PERSIANN-CDR
PSEA	34.11 (19.47 – 49.13)	39.83 (22.25 – 60.08)	34.74 (28.89 – 43.84)
MC	1.70 (0.34 – 11.78)	2.15 (0.00 – 12.64)	6.10 (3.11 – 17.30)
EA	0.54 (0.04 – 1.48)	1.23 (0.14 – 2.94)	5.14 (2.67 – 12.13)
WP	1.31 (0.12 – 7.45)	3.17 (0.00 – 13.17)	14.30 (4.78 – 20.05)

1208

1209




1210 **Table 2: Comparisons of boundary layer (BL; < 2 km) and free troposphere (FT; > 2 km) mean values across source regions. Uncertainty values are presented as 1**
 1211 **standard deviation. Bold values denote when significant statistical differences are found ($p < 0.05$). Corresponding p-values are provided in Table S2 of the Supplementary**
 1212 **Information. The EA-FT column was left blank due to the infrequent sampling of EA air in the FT.**

	EA		MC		PSEA		WP	
	BL	FT	BL	FT	BL	FT	BL	FT
$N_{100-1000nm}$ (cm^{-3})	839.11 ± 507.34	-	818.43 ± 571.90	223.87 ± 316.26	272.55 ± 125.29	35.88 ± 41.23	71.18 ± 66.13	14.72 ± 5.99
CO (ppm)	0.16 ± 0.07	-	0.18 ± 0.12	0.11 ± 0.07	0.10 ± 0.02	0.10 ± 0.02	0.08 ± 0.00	0.08 ± 0.01
O ₃ (ppbv)	45.29 ± 28.89	-	31.22 ± 10.62	23.69 ± 7.62	24.03 ± 3.44	29.69 ± 5.52	12.73 ± 3.27	18.81 ± 7.79
CH ₄ (ppm)	1.96 ± 0.06	-	1.86 ± 0.01	1.85 ± 0.01	1.88 ± 0.03	1.87 ± 0.01	1.88 ± 0.00	1.88 ± 0.01
SO ₄ ²⁻ ($\mu g m^{-3}$)	5.14 ± 2.40	-	2.43 ± 1.33	0.69 ± 0.75	1.03 ± 0.54	0.17 ± 0.15	0.79 ± 0.98	0.23 ± 0.09
NO ₃ ⁻ ($\mu g m^{-3}$)	0.19 ± 0.45	-	0.24 ± 0.32	0.08 ± 0.20	0.04 ± 0.04	0.00 ± 0.03	0.01 ± 0.04	-0.01 ± 0.05
NH ₄ ⁺ ($\mu g m^{-3}$)	1.32 ± 1.19	-	0.86 ± 0.75	0.28 ± 0.43	0.23 ± 0.24	-0.01 ± 0.18	0.10 ± 0.30	0.00 ± 0.20
OA ($\mu g m^{-3}$)	2.68 ± 3.54	-	7.23 ± 8.80	2.10 ± 4.52	0.96 ± 1.00	0.07 ± 0.24	0.13 ± 0.32	0.01 ± 0.21
BC ($ng m^{-3}$)	87.29 ± 98.53	-	71.81 ± 41.79	16.06 ± 17.40	24.90 ± 15.83	2.79 ± 5.59	1.03 ± 2.92	0.20 ± 0.91

1213




1214




1215

1216 **Figure 1:** Maps of (a) ground elevation from the Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010), flight
1217 tracks in red, and approximate source regions in labeled boxes: Peninsular Southeast Asia (PSEA), Maritime Continent
1218 (MC), East Asia (EA), and West Pacific (WP), (b) 2020 population density from the Center for International Earth Science
1219 Information Network (CIESIN) Gridded Population of the World (GPW) v4, (c) MODIS active fire hotspot density (only
1220 fires tagged with > 80% confidence) averaged at 0.5° × 0.5° resolution from 1 Aug to 15 Oct 2019, and (d) OMI-retrieved
1221 PBL SO₂ averaged over the same period.



1222


1223 Figure 2: Trajectory densities resolved by monsoon phase and sampling vertical level. Monsoon phases are southwest
1224 monsoon (SWM; before 20 Sep 2019) and the monsoon transition (MT; after 20 Sep 2019). Vertical sampling levels are
1225 divided into boundary layer (BL; < 2 km) and free troposphere (FT; > 2 km). Red star denotes Clark International Airport,
1226 Philippines. Number of trajectories (n) is shown in the upper right of each panel.




1227



1228 **Figure 3: Classified trajectories and synoptic conditions associated with transport from (a – c) Peninsular Southeast Asia**
 1229 **(PSEA), (d – f) Maritime Continent (MC), (g – i) East Asia (EA), and (j – l) West Pacific (WP). Left: Trajectory density**
 1230 **normalized to the mean per source region, with the number of trajectories classified into each source region**
 1231 **annotated on the lower left of each panel. Middle: NCEP 850 mb geopotential height anomaly from the mean for 2019, overlaid with**
 1232 **horizontal winds ($\geq 2 \text{ m s}^{-1}$). Right: PERSIANN-CDR average precipitation overlaid with NCEP 850 mb ω where red (blue)**
 1233 **contour lines represent ω values above 0.05 Pa s^{-1} (below -0.05 Pa s^{-1}).**



1234 **Figure 4: Relative contributions (in percentages) of air masses arriving from study regions (a) averaged over all altitudes**
 1235 **and (b) pressure levels (hPa). Total number of trajectories per pressure bin is provided on the right end of (b). Source**
 1236 **regions are Peninsular Southeast Asia (PSEA), Maritime Continent (MC), East Asia (EA), and West Pacific (WP).**
 1237
 1238




1239

1240

1241

1242

Figure 5: Vertical motion during transport from (a) Peninsular Southeast Asia (PSEA), (b) Maritime Continent (MC), (c) East Asia (EA), and (d) West Pacific (WP). Color corresponds to density as a function of trajectory altitude (pressure) and time step. Example trajectories are plotted in black to show actual vertical motion.



1243

1244


1245

1246

Figure 6: Vertical median profiles of composition for Peninsular Southeast Asia (PSEA), Maritime Continent (MC), East Asia (EA), and West Pacific (WP). (a) $N_{100-1000nm}$ (cm^{-3}), (b) CO (ppm), (c) SO_4^{2-} ($\mu\text{g m}^{-3}$), (d) OA ($\mu\text{g m}^{-3}$). Left and right sides of shaded areas refer to 25th and 75th percentiles, respectively.


1247

1248



1249

1250 **Figure 7: Linear regressions of (a) $\Delta\text{CO}/\Delta\text{CO}_2$, (b) $\Delta\text{CH}_4/\Delta\text{CO}_2$, (c) $\Delta\text{OA}/\Delta\text{SO}_4^{2-}$, and (d) $\Delta\text{ABC}/\Delta\text{CO}$ for the Maritime**
1251 **Continent (MC) and East Asia (EA). Annotated are species ratios (slope) per region and standard error (SE) as a measure**
1252 **of uncertainty (slope \pm SE). Pearson's R values are provided in parentheses. Only data with $\Delta\text{CO} > 0.02$ ppm were included**
1253 **to better identify combustion-related ratios. Peninsular Southeast Asia and West Pacific data were not plotted due to low**
1254 **correlations.**



1255

1256


1257

1258

1259

1260

Figure 8: Median FIMS number size distributions ($dN/d\log D_p$; cm^{-3}) as a function of geometric mean particle diameter (D_p ; nm), resolved by source region and sampling altitude in the boundary layer (BL; < 2 km) and free troposphere (FT; > 2 km). Source regions are Peninsular Southeast Asia (PSEA), Maritime Continent (MC), East Asia (EA), and West Pacific (WP). Upper and lower bounds of the shaded areas refer to 25th and 75th percentiles, respectively. The size distribution of EA air in the FT was not plotted due to the infrequent sampling of EA air in the FT.



1261

1262
1263
1264
1265
1266
1267

Figure 9: (a) $\Delta BC/\Delta CO$ ratio ($\Delta CO > 0.02$ ppm) as a function of accumulated precipitation along individual trajectories (APT) calculated with PERSIANN-CDR, and (b) histograms of $\Delta BC/\Delta CO$ resolved by wet (APT > 10 mm) and dry (APT < 10 mm) air masses. Medians and 25th/75th percentiles (error bars) are shown for (a) each source region: Peninsular Southeast Asia (PSEA), Maritime Continent (MC), and East Asia (EA). West Pacific (WP) was not plotted because of few data points where $\Delta CO > 0.02$ ppm. Black line in (a) represents the trend for all source-classified data (EA, MC, PSEA, WP) grouped into APT bins. Number of observations plotted per source region are provided in parentheses in (a).