

Global Dust Optical Depth Climatology Derived from CALIOP and MODIS
Aerosol Retrievals on Decadal Time Scales: Regional and Interannual Variability

5 Qianqian Song^{1,2}, Zhibo Zhang^{1,2,*}, Hongbin Yu³, Paul Ginoux⁴, Jerry Shen^{3,#}

1. Physics Department, UMBC, Baltimore, Maryland, USA

2. Joint Center of Earth Systems Technology, UMBC, Baltimore, Maryland,
USA

10 3. Earth Sciences Division, NASA Goddard Space Flight Center, Greenbelt,
Maryland, USA

4. NOAA Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey,
USA.

15

*Correspondence to: Zhibo Zhang, Zhibo.Zhang@umbc.edu

who worked as summer intern at NASA Goddard Space Flight Center during
June–August 2020.

Abstract

We derived two observation-based global monthly mean dust aerosol optical depth (DAOD) climatological datasets from 2007 to 2019 with a 2° (latitude) \times 5° (longitude) spatial resolution, one based on CALIOP and the other on MODIS observations. In addition, the CALIOP climatological dataset also includes dust vertical extinction profiles. Dust is distinguished from non-dust aerosols based on particle shape information (e.g., lidar depolarization ratio) for CALIOP, and on dust size and absorption information (e.g., fine-mode fraction, Ångström exponent, and single-scattering albedo) for MODIS, respectively. The two datasets compare reasonably well with the results reported in previous studies and the collocated AERONET coarse mode AOD. Based on these two datasets, we carried out a comprehensive comparative study of the spatial and temporal climatology of dust. On multi-year average basis, the global (60°S - 60°N) annual mean DAOD is 0.032 and 0.067 according to CALIOP and MODIS retrievals, respectively. In most dust active regions, CALIOP DAOD generally correlates well (correlation coefficient $R > 0.6$) with the MODIS DAOD, although CALIOP value is significantly smaller. CALIOP DAOD is 18%, 34%, 54% and 31% smaller than MODIS DAOD over Sahara Desert, the tropical Atlantic Ocean, the Caribbean Sea, and the Arabian Sea, respectively. Applying a regional specific lidar ratio (LR) of 58sr instead of the 44sr used in the CALIOP operational retrieval reduces the difference from 18% to 8% over the Sahara and from 34% to 12% over Tropical Atlantic Ocean. However, over Eastern Asia and the Northwestern Pacific Ocean (NWP), the two datasets show weak correlation. Despite these discrepancies, CALIOP and MODIS show similar seasonal and interannual variations in regional DAOD. For dust aerosol over NWP, both CALIOP and MODIS show a declining trend of DAOD at a rate of about $2\% \text{ yr}^{-1}$. This decreasing trend is consistent with the observed declining trend of DAOD

in the southern Gobi Desert at a rate of $3\% \text{ yr}^{-1}$ and $5\% \text{ yr}^{-1}$ according to CALIOP and MODIS, respectively. The decreasing trend of DAOD in the southern Gobi Desert is in turn
45 found to be significantly correlated with increasing vegetation and decreasing surface wind speed
in the area.

1 Introduction

Mineral dust, referred to as dust for short, is one of the most abundant type of atmospheric aerosol in terms of dry mass (Textor et al. 2006; Yu et al. 2012; Kok et al. 2017). Dust aerosol
50 directly interacts with both solar and thermal infrared radiation, known as the direct radiative
effect, and thereby influences the Earth's radiative energy budget (Kok et al, 2017; Song et al.,
2018; Di Biagio et al. 2020). Dust also influences the life cycle and properties of clouds by
altering the thermal structure of the atmosphere (known as semi-direct effects) (Hansen et al.,
1997) and acting as cloud condensation nuclei (CCN) and ice nuclei (IN) (known as indirect
55 effects) (Albrecht 1989; Rosenfeld and Lensky 1998; Twomey 1977). Dust storms and plumes
can degrade air quality affecting human health (Griffin, 2007; Querol et al., 2019). Dust
deposition provides essential nutrients to marine and terrestrial ecosystems (Jickells et al. 2005;
Yu et al., 2015b) but reduces the snow albedo increasing snow melt (Painter et al., 2007). All
these impacts manifest the important role of mineral dust in the Earth systems (e.g. Evan et al.,
60 2006; Lau & Kim, 2007; Miller & Tegen, 1998; Shao et al., 2011)

Dust production is sporadic in nature. Dust aerosol can be transported on intercontinental,
hemispherical, and even global scales (Grousset et al. 2003; Uno et al. 2009; Yu et al. 2012,
2013). Thus, global and routine measurements of dust spanning over years or even decades are
65 vital for studying dust transport and deposition, estimating the dust radiative effects, and

evaluating and constraining dust simulations in numerical weather and climate models. Satellite remote sensing is the only means to observe dust on regional to global scales. Satellite remote sensing techniques usually retrieve the optical depth or extinction profile for total aerosol in the atmosphere with additional retrievals of particle size, shape, or absorption properties that are sensor specific. Passive sensors have been used to detect dust sources and track dust plumes at global scales. A few examples are the Total Ozone Mapping Spectrometer (TOMS) (Prospero et al., 2002), Ozone Monitoring Instrument (OMI) (Chimot et al. 2017), Multi-angle Imaging SpectroRadiometer (MISR) (Ge et al., 2014 and Y. Yu et al. 2019), Moderate Resolution Imaging Spectroradiometer (MODIS) (Ginoux et al., 2010; Remer et al., 2005; Yu et al., 2009), multi-angular and polarimetric POLarization of Directionality of the Earth's Reflectances / Polarization and Anisotropy of Reflectances for Atmospheric science coupled with Observations from a Lidar (POLDER/PARASOL) measurements (Chen et al. 2018) and the International Association of Structural Integrators (IASI) (Klüser et al., 2011; Clarisse et al. 2019). On one hand, these passive sensors provide global or quasi global coverage of column integrated properties of aerosol with satisfactory temporal resolution. On the other hand, they do not provide the vertical structure of aerosol that is critical for studying aerosol-cloud interactions and aerosol influences on the thermal structure of the atmosphere. Space-borne lidar systems, such as the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) onboard the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) spacecraft (Winker et al., 2010) and the Cloud-Aerosol Transport System (CATS) onboard the International Space Station (Yorks et al. 2015) are able to provide the vertical structure of aerosol and clouds, albeit with limited spatial coverage. All these passive and active remote sensing observations have been

used extensively in studies of the spatial and temporal evolution of aerosol over the past decade (e.g., Proestakis et al. 2018).

90

A significant hurdle of applying satellite remote sensing measurements for dust studies is how to distinguish dust from other aerosol types in a quantitative way. While many studies have used total aerosol retrievals by focusing on the regions and seasons where dust dominates, some studies have developed sensor-specific methods of partitioning total aerosol into dust and non-
95 dust components with varying assumptions (Kaufman et al., 2005; Kalashnikova et al. 2005; Dubovik et al. 2006; Ginoux et al., 2010; Yu et al., 2009, 2013, 2015a, 2019). In general, the dust separation methods are based on dust physical and optical properties such as their large size, their irregular or non-spherical shape, and absorption characteristics. For example, CALIOP dust classification is mainly based on the fact that dust aerosols are non-spherical in shape and their
100 lidar depolarization ratio is significantly larger than those spherical aerosols. In contrast, the wide spectral coverage of MODIS measurements enables the retrieval of aerosol particle size information, such as effective radius, fine-mode fraction (FMF), and aerosol extinction Angstrom exponent, as well as spectral gradient of absorption (decreasing of absorption from UV to red) (Remer et al., 2005). The combinations of these retrievals provide the basis for dust
105 separation and DAOD retrievals from MODIS. Some recent studies have also characterized dust distribution through integrating satellite measurements with other data sources and model simulations. For example, Voss and Evan (2020) (referred to as VE20 hereafter) developed a dust optical depth record from MODIS retrievals, similar to Kaufman et al. (2005) over ocean and Ginoux et al. (2012) over land. Unlike Kaufman et al. (2005) and Yu et al. (2020) that
110 derived characteristic FMF values for combustion, dust, and marine aerosol from MODIS

retrievals, VE20 determined these characteristic FMFs from AERONET measurements. VE20 also extended the MODIS-based method to AVHRR over-ocean retrievals with some assumptions and produced the long-term (1981-2018) record of dust optical depth. Gkikas et al. 2021 developed a global fine resolution ($0.1^\circ \times 0.1^\circ$) DAOD dataset for the period 2006-2017 by scaling MODIS retrieved Collection 6.1 Aerosol Optical Depth (AOD) with the DAOD-to-AOD ratios provided by MERRA-2 (Modern-Era Retrospective analysis for Research and Applications, Version 2) reanalysis (Gelaro et al. 2017). Given that MODIS and other remote sensing measurements (e.g., MISR and AERONET) have been assimilated in the MERRA-2 reanalysis to constrain the aerosol optical depth, the DAOD-to-AOD ratio reported by MERRA-2 is the same as that from the underlying GOCART aerosol transport model in the MERRA-2 reanalysis system.

In this study, we focus on the DAOD derived from CALIOP and MODIS with two major objectives. First, we produce a decadal (2007-2019) record of global DAOD and dust vertical extinction coefficient profile climatology from the CALIOP observations, which represents an extension of the trans-Atlantic dust transport and deposition studies by Yu et al. (2015a, 2015b, 2019), both in terms of spatial and temporal coverages. Second, we compare the CALIOP DAOD climatology with the MODIS DAOD over both land and ocean (Yu et al. 2020; Pu and Ginoux, 2018) to identify and understand their differences in terms of global dust distribution and interannual variabilities including interannual trend in key dust regions. Our analysis goes beyond broad dust-laden regions by zooming into potential dust source areas, which provides important insights into local dust activities. A systematic comparison and better understanding of DAOD from the two sensors based on distinct retrieval algorithms is critical for applying

satellite measurements to evaluate global dust modeling (Kim et al. 2019). In comparison to
135 some most recent studies (Voss and Evan, 2020; Gkikas et al. 2021), our dust climatology is
derived using the satellite observations in a self-consistent way without blending in other
measurements (e.g., AERONET) or models (e.g., MERRA-2). As discussed in Yu et al. (2009),
the self-consistent use of MODIS data could minimize the introduction of additional biases due
to discrepancies in FMF between MODIS and AERONET. Furthermore, we use the latest
140 version 4.2 CALIOP products and version 6.1 MODIS products to characterize the spatial and
temporal distributions of dust. The rest of the paper is organized as follows. Section 2 provides a
description of the methodology of deriving dust climatology from CALIOP and MODIS. In
Section 3, we compare our DAOD datasets with previous studies and collocated AERONET
retrievals. In Section 4, we compare and study the DAOD climatology from CALIOP and
145 MODIS. Section 5 provides a summary of the study along with the main conclusions.

2 Dust Detection and AOD Partition Schemes

2.1 CALIOP Dust Detection and AOD Partition

CALIPSO is in a sun-synchronous polar orbit with an equator crossing time of around 13:30
150 local time and 98° orbit inclination. CALIOP is a two-wavelength (532nm and 1064nm)
polarization-sensitive lidar onboard CALIPSO. CALIPSO orbit track repeats every 16 days,
CALIOP sensor never provides global coverage due to its small footprint. At Earth's surface, the
diameter of CALIOP footprint is around 70m, with spacing distance of 333m between two
adjacent footprints along the orbit track. CALIOP utilizes three receiver channels (one measuring
155 the 1064nm backscatter intensity and two measuring orthogonally polarized components of the

532nm backscatter) to provide high vertical resolution 30-60m of aerosol and cloud structure profiles (Winker et al., 2009).

Aerosol subtype classification and a priori assumption of LR (extinction to backscatter ratio) for specific aerosol type are critical for CALIOP aerosol retrievals. CALIOP Level 2 product has been validated by comparing with ground-based measurements. The comparison between aerosol subtypes in CALIOP level 2 V2.01 and NASA Aerosol Robotic Network (AERONET) aerosol types shows that 70% of the CALIOP and AERONET aerosol types are in agreement and best agreement is achieved for dust and polluted dust (Mielonen et al. 2009). Schuster et al. (2012) compared CALIOP AOD to the collocated AERONET AOD measurements and found a CALIPSO bias of -13%, corresponding to an absolute bias of -0.029 relative to AERONET AOD on global average. Further comparison between CALIPSO AOD measurements and the collocated AERONET AOD measurements for the columns that contain the dust subtype exclusively showed a larger bias (i.e., -29% and corresponding absolute bias of -0.1), although they show a relatively high correlation of $R=0.58$; this indicates that the assumed LR of 40 sr for the CALIPSO dust retrievals is too low. Omar et al. 2013 showed that CALIOP AOD are lower than AERONET AOD especially for low AOD. Furthermore, they found that the median of relative AOD difference between CALIOP and AERONET (500nm) is 25% of AERONET AOD for $AOD > 0.1$.

CALIOP observations have been used widely in previous studies of the spatial and temporal evolution of dust aerosols over the past decade (Huang et al. 2007, 2008; Yang et al. 2012; Xu et al. 2016; Kim et al., 2019). It is important to note that these studies are regional in scope and

they use the standard CALIPSO product and aerosol subtype classification algorithm (Omar et al. 2009). In the standard CALIPSO product, each detected aerosol layer is classified as one of the six subtypes: dust, polluted dust, polluted continental, smoke, clean marine and clean continental. In the latest CALIOP version, another sub-type “marine-dust” is introduced (Kim et al. 2018). In these studies, the “dust” subtype or a combination of “dust” and “polluted dust” subtypes is categorized as dust. While the former assumption leads to an underestimate of dust due to neglecting dust component in the “polluted-dust” subtype, the latter assumption results in an overestimate of dust because of accounting for non-dust component in the “polluted-dust” subtype. In order to better distinguish dust component from each CALIOP detected aerosol layers, Yu et al. (2015a) developed an algorithm independent of the standard aerosol subtype classification to distinguish dust from non-dust aerosol by using their respective thresholds of particulate depolarization ratio (Table 1). The depolarization-based dust separation algorithm is based on the method developed by Shimizu et al. 2004, Hayasaka et al. 2007 and Tesche et al. 2009. The algorithm has been implemented in the framework of surface lidar network such as European Aerosol Research Lidar Network (EARLINET) (Ansmann et al. 2011) and also applied to CALIOP observations (Yu et al., 2012; Amiridis et al. 2013; Yu et al., 2015a). They further used the derived three-dimensional distribution of dust extinction to quantify the trans-Atlantic dust transport and deposition and its implications for Amazon rainforest (Yu et al., 2015b, 2019).

In this study, we use the methodology in Yu et al. (2015a) to derive the monthly mean dust extinction profile under clear-sky conditions from the latest V4.20 CALIOP products on a global scale from 2007 to 2019. First, we select the cloud-free columns based on the CALIOP cloud

layer product. In order to increase the sampling, we define clear-sky cases in this study either as columns that are completely cloud-free or with the presence of optically thin (cloud optical depth < 0.2) and high-level (cloud base $> 7\text{km}$) clouds. This is justified that the presence of high-level optically thin clouds does not significantly affect the retrieval of aerosol layers below the clouds (Yu et al. 2015a). After clear-sky screening, we use the operational 5 km level 2 CALIOP aerosol profile product that contains aerosol depolarization, backscatter and extinction profiles over a global scale (Young et al. 2018) to derive the dust extinction profile. The depolarization ratio from CALIOP is a key variable for detecting and distinguishing dust from non-dust aerosol. Backscatter by spherical particle largely retains the polarization of the incident light, resulting in a depolarization ratio of nearly zero. In contrast, dust particles are generally non-spherical in shape and large in size, which gives them non-zero depolarization ratio that is significantly larger than other types of aerosol. The cloud-aerosol discrimination (CAD) score in the products gauges the level of confidence for a feature being classified as aerosol or cloud. In this study, in order to screen out low-confidence aerosol and cloud discrimination, we select layers with CAD scores between -90 and -100 (high level of confidence for aerosol feature) by following Yu et al. (2019). Aerosol profile product also provides extinction quality control flag (Ext_QC) to indicate problematic retrievals. This study only uses layers with Ext_QC values of 0, 1, 18, and 16 (Winker et al., 2013). Only nighttime data are used to avoid sunlight interference in aerosol signals.

For each aerosol backscatter coefficient profile, we derive the fraction of dust backscatter to total backscatter (f_d) at each altitude from the following equation

$$f_d = \frac{(\delta - \delta_{nd})(1 + \delta_d)}{(\delta_d - \delta_{nd})(1 + \delta)} , \quad (1)$$

where δ is CALIOP observed particulate depolarization ratio, δ_d and δ_{nd} is a priori knowledge of depolarization ratios of dust and non-dust aerosols respectively. Clearly, the calculations of f_d in Eq. (1) rely on the a priori depolarization ratios of dust and non-dust aerosols (i.e., δ_d and δ_{nd}). To account for various types of non-dust aerosols with different depolarization ratio, we follow Yu et al. 2015a and assume 0.02 and 0.07 as lower and upper bounds for δ_{nd} (Burton et al., 2012; Fiebig et al., 2002; Sakai et al., 2010). Dust aerosols have significantly larger depolarization ratio compared to non-dust aerosols. To account for the variability of dust shape and size, we use 0.2 and 0.3 as lower and upper bounds for δ_d (Ansmann et al., 2012; Esselborn et al., 2009; Sakai et al., 2010). Given an observed dust depolarization ratio δ , the f_d based on Eq. (1) has the minimum value when $\delta_d = 0.30$ and $\delta_{nd} = 0.07$ and the maximum value when $\delta_d = 0.20$ and $\delta_{nd} = 0.02$. To account for this variability, the final f_d is based on the mean of the lowest (i.e., $\delta_d = 0.30$ and $\delta_{nd} = 0.07$) and the highest (i.e., $\delta_d = 0.20$ and $\delta_{nd} = 0.02$) dust scenario.

In each 2° (latitude) $\times 5^\circ$ (longitude) grid, at each altitude, dust backscatter coefficient for per clear-sky overpass is derived by multiplying CALIOP total backscatter coefficient with the calculated f_d from Eq. (1). To derive dust extinction coefficient from dust backscatter coefficient, we assume dust LR, i.e., extinction to backscatter ratio, of $44 \pm 9 \text{ sr}$ at 532nm, consistent with CALIOP Version 4.20 operational retrieval (Kim et al., 2018). The monthly mean dust extinction coefficient is calculated at each altitude when overpass samples within the month is larger than 5. Then DAOD is calculated by integrating the monthly mean extinction coefficient profile for each grid. The use of globally uniform LR and the selection of δ_d and δ_{nd} could induce uncertainty to the derived DAOD. This is discussed in section 3.

It is important to note that in this study we use only nighttime CALIOP observations for DAOD retrievals. This is because the daytime CALIOP observations are often contaminated by background solar noise (Getzewich et al. 2018). As shown in Figure S1 in the supplementary material, when the above DAOD retrieval method is applied to daytime CALIOP observation, there is a widespread non-zero DAOD retrieval over remote ocean regions where dust should be scarce. This is apparently an artifact caused by solar contamination on CALIOP daytime observations, which motivates and justifies our use of nighttime CALIOP observations. On the other hand, however, this leads to an inconsistency with the MODIS DAOD retrieval which is based on daytime observations (see section 2.2). Although the diurnal cycle of dust has been investigated using model simulations (e.g., Yue et al. 2009), it is extremely difficult to assess dust diurnal variation from polar orbiting remote sensing observations, especially using elastic lidar in visible region like CALIOP, due to the inherent instrument limitation. For example, a recent study by Yu et al. 2021 attempted to use the retrievals from the Cloud-Aerosol Transport System (CATS) lidar to study the diurnal cycle of dust. The 51.6-degree inclination orbit allows CATS to sample the tropical and midlatitude regions multiple times a day, which make it more advantageous than CALIOP for diurnal variability studies. Unfortunately, after a validation comparison with AERONET observations (i.e., solar-based during daytime and lunar-based during nighttime), they found a significant day–night inconsistency in their retrieval quality. Because of this inconsistency, they concluded that diurnal variability in dust and dust mixture characteristics have to be examined separately for daytime and nighttime periods. Nevertheless, Yu et al. 2021 plotted the daytime and nighttime DAOD together for several dust active regions (see their Figures 3 and 10-13). The contrast between daytime and nighttime DAOD based on these plots is roughly between 10-15%, which is smaller than other uncertainties in CALIOP

retrievals as analyzed in section 3. Again, it has to be emphasized that this contrast is partly due
270 to the day–night inconsistency in CATS data quality.

2.2 MODIS Dust Detection and AOD Partition

As described above, the CALIOP-based DAOD derivation mainly makes use of dust non-sphericity in shape to separate dust aerosol from others. Another important difference of dust aerosol from other types of aerosols is their relatively large size. This difference provides the
275 basis for the dust separation. DAOD derivation scheme based on the Moderate Resolution Imaging Spectroradiometer (MODIS) retrievals is introduced in this section.

MODIS sensors onboard of the Aqua and Terra satellites measure radiances at 36 spectral bands ranging from 0.41 to 14 μm , with a 2330 km swath that provides near-global coverage
280 every day. MODIS aerosol retrievals employ two complementary algorithms to achieve the global coverage. The Dark Target (DT) algorithm is applicable for the retrieval of aerosol loading and properties over dark surfaces, including ocean-water and vegetated land. The MODIS aerosol AOD retrievals over ocean are found within the retrieval errors of $\Delta\tau_a = \pm 0.03 \pm 0.05\tau_a$ relative to AERONET AOD measurements (Remer et al. 2005). An approach
285 was developed in previous studies to separate DAOD from other types of aerosol by using aerosol optical depth (τ) and fine mode fraction (f) retrieved from MODIS DT retrieval over ocean. Both τ and f refer to properties at 550nm hereafter, unless specified otherwise. In this approach, both τ and fine-mode AOD ($f\tau$) are assumed to be composed of marine aerosol, dust and combustion aerosols, i.e.,

$$\tau = \tau_m + \tau_d + \tau_c , \quad (2)$$

$$f\tau = f_m\tau_m + f_d\tau_d + f_c\tau_c , \quad (3)$$

290 Where the subscripts m, d, and c represent marine aerosol, dust and combustion aerosol, respectively. Based on Eq. (2) and (3), τ_d can be calculated from MODIS-retrieved τ and f , with appropriate parameterizations for f_m, f_d, f_c and τ_m . More specifically, f_m, f_d, f_c were determined from retrieved f in selected regions and seasons for which a specific aerosol type dominates, τ_m was parameterized as a function of wind speed (details can be found in Kaufman et al. 2005; Yu
295 et al., 2009, 2020).

Over land, MODIS aerosol properties including AOD, Ångström exponent, SSA are retrieved from the Deep Blue (DB) algorithm (Hsu et al. 2004, 2013). The MODIS aerosol AOD retrievals over land are found within the retrieval errors of $\Delta\tau_a = \pm 0.05 \pm 0.15\tau_a$ relative to
300 AERONET AOD measurements (Remer et al. 2005). DAOD over land is derived from the AOD using one criterion based on size distribution (to distinguish fine and coarse modes) and the other criterion based on absorption (to distinguish between scattering sea salt and absorbing dust). To apply first criterion, we use the following formula established by Anderson et al. 2005 using in-situ data:

$$COD_M = AOD \times (0.98 - 0.5089\alpha + 0.051\alpha^2) , \quad (4)$$

305 Where α is the Ångström exponent (a measure of the wavelength dependence of optical depth) which has been shown to be highly sensitive to particle size (Eck et al. 1999), COD_M is the coarse mode fraction (aerodynamic diameters larger than $1\mu m$) of AOD retrieved from MODIS, with a contribution from absorbing (DAOD) and scattering aerosols (sea salt aerosol optical depth). The second criterion requires the single-scattering albedo at 470nm to be less than 0.99
310 for the retrieval of DAOD (more details can be found in Pu and Ginoux, 2018).

Overall, multi-wavelength observations from MODIS contains aerosol size information such as fine-mode fraction and Ångström exponent in the observed reflectance spectral pattern, which was used to separate dust aerosol from others in MODIS dust retrieval over ocean and land (Table 1). In this study, the latest retrieved aerosol properties from MODIS Collection 6.1 are used. We use data from Aqua MODIS only, because Terra MODIS retrievals may generate spurious dust trend (Yu et al. 2020). In order to minimize cloud contamination and avoid the infrequent sampling to bias DAOD in MODIS dust retrieval over ocean, we screen the data by requiring a minimum of 10 DAOD retrievals in a month.

The relevant variables and the quality assurance procedures used in CALIOP- and MODIS-based DAOD retrievals are summarized in Table 1 and Table S1, respectively.

3 Comparison with previous studies and Uncertainty Analysis

Based on the dust detection and separation schemes of two sensors described in section 2, we derived the following three datasets:

1. The monthly mean CALIOP-based total aerosol optical depth (TAOD) and DAOD, as well as the vertical extinction profile on a 2° (latitude) $\times 5^\circ$ (longitude) spatial resolution grids for the period of 2007 – 2019. This relatively coarse resolution is limited by CALIOP's sampling.
2. We combine the monthly mean Aqua MODIS over-ocean (Yu et al., 2020) and over-land (Pu and Ginoux, 2018) TAOD and DAOD on a $1^\circ \times 1^\circ$ spatial resolution grids to get the monthly mean MODIS-based TAOD and DAOD from 2003 to 2019. In order to compare with CALIOP-based dust climatology data, we aggregate the $1^\circ \times 1^\circ$ MODIS-based data to $2^\circ \times 5^\circ$ resolution grids.

3. For evaluation and comparison purpose (see section 4.1), we also produce a seasonal global distribution of conditionally sampled DAOD from CALIOP (Marinou et al. 2017, Proestakis et al. 2018). While the standard climatological DAOD includes all cloud-free cases in the average of dust extinction and DAOD regardless of the presence of dust, the conditionally sampled DAOD calculation only averages those cases where dust is detected (i.e., DAOD and dust extinction are non-zero). Therefore, the conditionally sampled DAOD is directly related to the intensity of the detected dust events, whereas the climatological DAOD is determined by a number of factors including not only the intensity of the detected dust events but also the frequency of the dust events as well as the capability of the instrument to sample the dust events.

3.1 Comparison with previous studies

Before we compare and study the DAOD climatology from MODIS and CALIOP in detail in the next section, we first evaluate our retrievals through comparisons with the regional and global DAOD values reported in the previous studies and explore the potential reasons for the differences.

Table 2 summarizes a comprehensive comparison of our DAOD datasets with previous studies. In Ridley et al. 2016, DAOD is first estimated in 14 dust-laden regions from the combination of AERONET measurements, MODIS and MISR retrievals. Then the observation-based, regional DAOD estimates are estimated to the global scale based on the model-estimated regional-to-global DAOD ratio. Using this method, they estimated that the global ($90^{\circ}\text{S}\sim 90^{\circ}\text{N}$) DAOD@550nm is 0.03 ± 0.005 . Using the DAOD-to-AOD ratio from MERRA-2, Gkikas et al. 2021 converted the MODIS AOD retrievals to DAOD and found a similar global ($90^{\circ}\text{S}\sim 90^{\circ}\text{N}$)

DAOD@550nm around 0.033. In contrast, as shown in Table 2 our MODIS-based global (90°S~90°N) DAOD is 0.057. However, it is important to note that the global mean DAOD values from these studies are not directly comparable to our global mean results because of the methodology differences. In particular, both of aforementioned studies used model simulations to aid their global DAOD estimate, while our estimates are completely based on observations (More precisely, DAOD of the scope 60°S~60°N are completely based on observations, while outside of the scope, DAOD is assumed to be zero). Nevertheless, to gain a more insightful understanding of the differences, we select the same 14 dust-laden regions as in the Ridley et al. 2016 (see Figure S2 in the supplementary material) and derive the corresponding regional DAOD (see Figure S3 and Table S2 in the supplementary material). As aforementioned, in Ridley et al. 2016 the DAOD in these dust-laden regions is based on AERONET measurements and satellite retrievals, and therefore more comparable with our results. As shown in the supplementary material (Figure S3), our regional MODIS-based DAOD values are in excellent agreement with those reported in Ridley et al. 2016 (relative bias $Br = -5.8\%$ in DJF, -0.2% in MAM, -2.5% in JJA and -10.4% in SON). This regional comparison suggests that the difference in global DAOD between our study and Ridley et al. 2016 is probably because we used different methods to derive the DAOD in the regions with less frequent dust activities (i.e., observation-based vs. model-based).

Recently, VE20 used a method similar to our MODIS-based DAOD estimate methodology to derive the global DAOD. Because of the use of similar methodology and data, VE20 is more comparable to our study than Ridley et al. 2016 or Gkikas et al. 2021. They estimated that the long-term mean DAOD to be 0.1 over land between 50°S and 60°N, which is almost identical to

our estimate of 0.103 (60°S ~ 60°N) as shown in Table 2. However, when averaged over the ocean, their DAOD estimate (0.03 ± 0.01) is significantly smaller than our result (0.055). As explained in the supplementary material, this difference is probably because different parameterizations of f_m, f_d, f_c and τ_m in Eq. (3) used in the two studies (see Table S4 and discussions in supplementary material).

A recent study by Proestakis et al. 2018 used a method similar to ours as described in section 2.1 to derive CALIOP-based regional DAOD in five dust-laden regions in Asia. We compared our CALIOP-based regional DAOD for the same regions (Figure S4) and compare the results with the values reported in Proestakis et al. 2018. As shown in Figure S5 of the supplementary material, the two studies are in excellent agreement with relative difference Br = 5.5% in DJF, -6.0% in MAM, -6.9% in JJA and 0.8% in SON, respectively.

Overall, the above comparisons indicate that our DAOD retrievals are in reasonable agreement with previous studies (where directly comparable). However, none of the aforementioned previous studies performed a systematic comparison between MODIS- and CALIOP-based DAOD, which is one of the motivations for this study and will be addressed in the Section 4.

3.2 Uncertainty Analysis

In order to understand the differences between the MODIS- and CALIOP-based DAOD, it is important to identify and quantify the uncertainties in each retrieval. The uncertainty of CALIOP DAOD retrieval come from several sources: An important source is the inherent uncertainty associated with CALIOP observations and its retrieval algorithm, such as instrument

calibration errors, errors in discriminating cloud from aerosol and failure to detect aerosol layers (including tenuous aerosol layer and the lower part of heavy dust layer. For example, Thorsen and Fu (2015) estimated that CALIOP may have underestimated 30%-50% in the magnitude of aerosol direct radiative effect due to its low sensitivity to tenuous layer), which is likely to translate into low bias in DAOD. In heavy aerosol conditions (e.g., strong dust storms in source regions and outflow regions), CALIOP laser cannot penetrate to the bottom of aerosol layer due to the laser attenuation (Chamara et al., 2017), which could also lead to a low bias in CALIOP DAOD.

CALIOP-based DAOD is also subject to the uncertainty associated with the assumed dust LR. Different deserts produce dust with different minerology, size and shape, and thus different LRs. Voss et al., (2001) measures LR for African dust as 41 ± 8 sr using a micropulse lidar system and Liu et al. (2002) measures LR for Asian dust as 42-55 sr. Globally observed LRs are summarized in Müller et al., (2007) and Baars et al., (2016). In this study, we assume dust LR to be 44 ± 9 sr at 532nm to be consistent with the value used in the CALIOP V4 product (Kim et al. 2018). This LR range is also comparable to previous studies and basically covers the range of typical dust LRs from 35 sr to 55 sr (Muller et al. 2007, Baars et al. 2016). The ± 9 sr induces $\pm 20\%$ DAOD uncertainties. When separating dust from non-dust aerosol, the choice of depolarization ratio (DPR) for dust aerosols and non-dust aerosols also introduces uncertainty in DAOD. To quantify the uncertainty caused by DPR selection, we also calculated DAOD in the lowest ($\delta_d = 0.30$ and $\delta_{nd} = 0.07$) and the highest ($\delta_d = 0.20$ and $\delta_{nd} = 0.02$) dust fraction scenarios. The uncertainty induced by DPR is region dependent (Figure S6). The uncertainty is

much lower in dust dominant regions than other regions. The averaged uncertainty for regions with DAOD>0.05 is 20%, while the averaged uncertainty for other regions is 38%.

MODIS dust detection is also subject to a number of uncertainties. Over ocean, the persistent presence of clouds, especially broken clouds, poses a great challenge to the MODIS aerosol retrievals (Martins et al. 2002). If a cloud is mistaken as aerosol, it would lead to a high AOD and low FMF bias, and thereby a high DAOD bias. In addition, DAOD was calculated from the MODIS-retrieved AOD (τ) and FMF (f) with appropriate parameterizations of marine aerosol AOD (τ_m), FMF of dust (f_{dust}), combustion (f_c) and marine (f_m) aerosols (see Table 2 in Yu et al. 2020 for the parameterization values). All the parameterizations could also introduce uncertainty in the derived DAOD, in particular on a regional basis (see details in Yu et al. 2020). Over land, the derived MODIS DAOD represents the coarse-mode fraction (aerodynamic diameters larger than $1\mu m$) of dust only. The exclusion of submicron dust aerosol could induce around 3% underestimation of the global atmospheric dust mass load and around 15% underestimation of the global DAOD (see Figure S1 in Kok et al. 2017).

One way to evaluate these uncertainties and validate the two dust detection methods is to compare with an independent measurement of DAOD. AERONET measurements have been considered as ground truth and often used to evaluate satellite aerosol optical depth retrievals. However, so far there is not a valid method to derive DAOD from AERONET AOD measurements to compare our results with. Therefore, we use coarse-mode AOD (COD) from AERONET measurements as a proxy for DAOD (Pu and Ginoux, 2018) to compare with our DAOD datasets and further estimate the absolute expected errors (EE) associated with our

450 DAOD datasets. The fine mode and coarse mode AOD in AERONET product are defined optically, rather than in terms of a microphysical cutoff of the associated particle size distribution at some specific radius (see details in O'Neill et al. 2003). Over land especially dust source regions, dust aerosols are predominantly in coarse mode, therefore, AERONET COD could be considered as a good proxy of DAOD over land. Over ocean, the exclusion of fine mode DAOD
455 could be partially cancelled by the inclusion of coarse sea salt AOD in AERONET COD retrievals. Therefore, AERONET COD is considered as a proxy of DAOD over ocean as well.

We use AERONET monthly mean COD retrieved at 500nm from the level 2 (cloud screened and quality assured) Spectral Deconvolution Algorithm (SDA) version 4.1 in this study.
460 The AERONET COD is converted to 550nm and 532nm using Angstrom Exponent to compare with MODIS and CALIOP DAOD retrievals, respectively. In addition, we produce a finer resolution ($1^\circ \times 1^\circ$) CALIOP-based DAOD retrieval to compare with AERONET COD.

For overland dust retrievals, between 2007 and 2019, there are 16653 MODIS, CALIOP
465 monthly mean DAOD retrievals collocated with 761 AERONET sites located within a 1-degree MODIS and CALIOP grid cell (Figure 1). MODIS DAOD ($DAOD_M$) overall bias high compared to AERONET COD with absolute bias $B_a = 0.01$, and relative bias $B_r = 26.7\%$. While CALIOP DAOD ($DAOD_C$) generally bias low with $B_a = -0.02$ and $B_r = -27.9\%$. Using a methodology suggested in Sayer et al. 2013, the estimated EE (take 68th percentiles referring to
470 Sayer et al. 2013) for all collocated MODIS DAOD over land is approximately $0.65 \times DAOD_M + 0$, and for CALIOP DAOD over land is approximately $0.52 \times DAOD_C + 0.02$ (Figure 2).

For over-ocean dust retrievals, between 2007 and 2019, there are 7755 MODIS, CALIOP monthly mean DAOD retrievals collocated with 311 AERONET sites located within a 1-degree MODIS and CALIOP grid cell (Figure 3). MODIS DAOD overall bias high compared with AERONET COD with absolute bias $B_a = 0.01$, and relative bias $B_r = 18.1\%$. While CALIOP DAOD generally bias low with $B_a = -0.02$ and $B_r = -35\%$. The estimated EE for all collocated MODIS DAOD over land is approximately $0.50 \times \text{DAOD}_M + 0$, and for CALIOP DAOD over land is approximately $0.54 \times \text{DAOD}_C + 0.02$ (Figure 4).

We further analyze the statistical parameters and EE by continents for MODIS and CALIOP DAOD (Table 3). The lowest EE, B_r and highest correlation (R) are estimated over Africa, followed by Asia, Europe, Americas and Australia. This implies that our DAOD retrievals are subject to higher bias under high AOD in polluted regions. Overall, MODIS-based monthly mean DAOD retrievals are larger than AERONET COD measurements, while CALIOP-based DAOD retrievals are smaller than AERONET COD, which seems to suggest that the true DAOD fall between the MODIS and CALIOP DAOD products.

4 Global Dust Climatology

In this section, we compare CALIOP global dust retrieval against MODIS dust retrieval, more specifically MODIS ocean dust retrieval from Yu et al. (2009, 2020) and land dust retrieval from Pu and Ginoux (2018), we analyze the similarities and differences between two dust climatological datasets and furthermore study the seasonal cycle and decadal trend of dust aerosols based on these datasets.

4.1 Comparison between CALIOP and MODIS DAOD Climatology

The DAOD climatology datasets derived from the CALIOP and MODIS observations, as described in Section 3, have two major sources of uncertainty:

1) The uncertainty associated with the TAOD retrieval. The primary uncertainty sources in MODIS TAOD retrieval include instrument calibration errors, cloud-masking errors, inappropriate assumption of surface reflectance and aerosol model selection (Remer et al. 2005; (Levy et al. 2013, 2018). Uncertainty sources in CALIOP aerosol retrieval include instrument calibration errors, errors in discriminating cloud from aerosol, uncertainties associated with the a priori assumption of LRs, under detection of tenuous aerosol layers, and overestimation of the elevation height of heavy aerosol plume base (Winker et al. 2009; Yu et al., 2010; Schuster et al., 2012; Thorsen and Fu, 2015; Rajapakshe et al. 2017).

2) The uncertainty associated with dust detection and separation. As explained in section 2, CALIOP- and MODIS-based dust detection and separation methods are based on different characteristics of dust aerosols in comparison with other types of aerosols, as summarized in Table 1. The CALIOP-based method makes use of the fact that depolarization ratio of dust aerosols is much higher than other types of aerosols, primarily because of irregular non-spherical shape and also to a lesser extent because of coarse size of dust particles (Gasteiger et al. 2011, Järvinen et al. 2016). MODIS-based method is largely based on the characteristics of coarse particle size. Over ocean, DAOD is derived from MODIS-retrieved TAOD and Fine Mode Fraction (FMF) with a priori characteristic FMF for individual aerosol types. Over land, DAOD is derived using spectral dependence of aerosol extinction (i.e., Angstrom exponent) and single scattering albedo. In other words, MODIS retrieves overland DAOD based on dust size supplemented by absorption characteristics.

Given these retrieval uncertainties and methodological differences, some discrepancies
520 between the two DAOD climatological datasets are expected. In this section, we will compare
the two datasets to identify and understand their similarities and differences. Since the
mechanisms of dust generation, dust transport and dust removal processes all have a seasonal
cycle (Mbourou et al. 1997; Parrington et al. 1983), we first present and discuss dust spatial
distributions for each season in this section. Table 4 summarizes the seasonal and annual mean
525 DAOD and TAOD values averaged over ocean, land and the globe (all limited to 60°S-60°N),
respectively, based on MODIS and CALIOP dust retrievals from 2007 to 2019. On multi-year
average basis, the global DAOD was found to be 0.055 over the ocean and 0.103 over land based
on MODIS, and 0.020 over ocean and 0.068 over land based on CALIOP. The global, annual
mean DAOD (TAOD) is 0.032 (0.121) and 0.067 (0.171) according to CALIOP and MODIS
530 retrieval, respectively.

As a comparison of two DAOD retrievals in this study, generally, DAOD from two
retrievals differ by a factor of about 3 over ocean and less than 2 over land, while TAOD differ
by a factor of less than 2 over both ocean and land. The ratio of DAOD over land to that over
ocean is about 2 and 3 for MODIS and CALIOP, respectively. For TAOD, the land to ocean ratio
535 is about 2 for both products. Overall, the difference in TAOD between two retrievals is less than
their difference in DAOD. On a global average, both MODIS and CALIOP-based DAOD peaks
in boreal summer (June-July-August). DAOD reaches minimum in boreal Fall (September-
October-November) for MODIS but in boreal Winter (December-January-February) for
CALIOP. The MODIS and CALIOP differences are region dependent, which is discussed as
540 follows.

Figure 5 shows the spatial distribution of seasonal mean DAOD and the percentage of DAOD to the TAOD based on 13-year (2007-2019) CALIOP and MODIS observations. Note that this period is chosen because both datasets are available. Generally, MODIS-based DAOD is larger than CALIOP-based DAOD. As expected, high values are seen from both CALIOP-based and MODIS-based DAOD over the ‘dust belt’ regions extending from the west coast of North Africa to the Middle East, Central Asia, and China, where large-scale dust activities occur persistently throughout the year. However, the CALIOP-based DAOD is rather low in some other regions that are known to be dusty in certain seasons, such as the southwestern United States, South America (Patagonian Desert), Australia, and South Africa (i.e., Kalahari Desert). These regions do stand out in MODIS DAOD maps (i.e., the second column in Figure 5). Then we plot DAOD-to-TAOD ratio based on DAOD and TAOD retrievals from two sensors (the last two columns in Figure 5). These regions indeed show up in the DAOD-to-TAOD ratio plot based on both sensors (i.e., the last two columns in Figure 5). This means that in those regions both sensor-specific methodologies are able to distinguish dust aerosol from sensor-detected total aerosol to some extent so that the DAOD-to-TAOD ratio stands out in those regions for both sensors.

The climatological dust product shown in Figure 5 is a measure of the average dust loading over a geographical domain and time interval. It contains information of both the intensity and frequency of dust activities. The seasonal conditionally sampled DAOD shown in the first column of Figure 6 eliminates the impacts from dust frequency by excluding dust-free cases in the average. It is mainly related to the intensity of observed dust events. Therefore, the comparison between climatological and conditionally sampled DAOD sheds a light on the

frequency and intensity of dust events detected by CALIOP. Therefore, we further compare the
565 seasonal climatological DAOD and conditional DAOD product. The second column of Figure 6
shows the seasonal climatological DAOD. The third column in Figure 6 shows the relative
difference between conditionally sampled DAOD and climatological DAOD with respect to the
climatological DAOD. In ‘dust belt’ regions, especially in Sahara Desert and Middle East where
dust activities are persistent, climatological DAOD is very close to conditional DAOD. In
570 Australia, Southwest United State, South America and South Africa, however, the conditional
DAOD (column 1 in Figure 6) and the difference (column 3 in Figure 6) are relatively high. This
suggests that dust activities in those regions are highly episodic and/or occur in relatively small
scales. The difference also is very large in open oceans, suggesting that dust aerosols are present
at a very low frequency.

575
Having analyzed the conditionally sampled DAOD from CALIOP, we now return to
climatological DAOD and comparison between CALIOP and MODIS. Hereafter, all AOD
values are climatological without otherwise explicit statement. Figure 7 shows the difference in
seasonal mean TAOD, DAOD, and the percentage of DAOD in TAOD between MODIS
580 retrievals and CALIOP retrievals. We first focus on ‘dust belt’ and its ocean out-flow regions
extending from Northeast Atlantic, North Africa to the Middle East, Central Asia, China and
Northwest Pacific. We note that in Figure 7 CALIOP-based TAOD and DAOD is generally
smaller than MODIS-based ones over North Africa and Saharan dust out-flow region over the
tropical Atlantic Ocean. One of the reasons of this large discrepancy is the choose of LR in
585 CALIOP aerosol retrieval in these regions. CALIOP V4 products retrieve dust extinction
coefficients with two steps. First, apply a globally uniform LR of 44sr for the identified dust

aerosol layers to retrieve backscatter coefficients. Second, use the same LR of $44sr$ value to convert backscatter coefficients to extinction coefficients. Amiridis et al. 2013 shows that in the second step applying LR of $58sr$ to CALIOP dust backscatter coefficients in North Africa improves the resulting aerosol extinction in terms of optical depth comparison with synchronous and collocated AERONET and MODIS measurements. Similarly, over Sahara Desert and the tropical Atlantic Ocean (see Figure 8 (a) and (d)), we apply LR of $58sr$ to the derived backscatter coefficient of dust component to get extinction coefficient of dust component. The resulting DAOD for LR of $58sr$ shows an improvement in comparison with MODIS DAOD relative to LR of $44sr$ (Figure 9 (a) and (d)). Therefore, the choose of LR can largely explain the difference between MODIS and CALIOP DAOD over North Africa and tropical Atlantic Ocean. For other regions, typical values of LR of desert dust aerosols vary between 35 and 55 sr , which is basically covered by the range of $44 \pm 9sr$ used in this study. The DAOD uncertainty induced by $\pm 9sr$ is estimated to be around 20% as shown in the shaded area in Figure 9.

In Middle East (the region indicated by Figure 8 (b)), the second column in Figure 7 shows that MODIS-DAOD is generally larger than CALIOP-DAOD in Arabian Peninsula, while opposite in India.

In Arabian Sea (the region indicated by Figure 8 (h)), comparing column 2 and column 4 in Figure 7, we could see that MODIS-DAOD is significantly larger than CALIOP-DAOD during JJA, during which cloud fraction is very high in the region. MODIS aerosol retrieval is more susceptible to cloud contamination. Specifically, the cloud contamination can lead to an overestimation of TAOD but underestimation of FMF. Although the MODIS retrieval algorithm neither assume coarse particles are exclusively from dust aerosols nor assume dust particles are

all coarse particles (Yu et al., 2020), coarse mode aerosols are primarily dust. Thus, the overestimation of TAOD and underestimation of FMF will lead to an overestimation in DAOD.

Over Eastern Asia and Asian dust outflow region (Northwest Pacific-NWP), CALIOP-based DAOD is generally smaller than MODIS-based DAOD. There could be several reasons for this. First, this region is a major outflow region of Asian pollution (Yu et al., 2020). It is possible that the internal mixing of dust aerosols with industrial pollution in this region changes the dust morphology making it less non-spherical (Li and Shao 2009, Huang et al. 2020) but larger in size, which leads to smaller depolarization ratio and smaller fine-mode fraction. As a result, CALIOP shape-based DAOD derivation method could not capture the dust particles contained in the mixture, while those dust particles can be captured by MODIS size-based method. Another potential reason could be associated with that dust plumes in this region are vertically dispersed (Yu et al., 2010; Su and Toon, 2011). These tenuous dust layers are likely to go undetected by CALIOP because of its relatively low sensitivity. However, MODIS retrieves aerosol from the columnal integrated reflectance which is not dependent on the vertical distribution of aerosol. The difference may also be caused by uncertainties in MODIS aerosol retrievals. The West Pacific Ocean is cloudy almost all year long (see the last column in Figure 7), which makes MODIS aerosol retrievals bias high due to its more susceptibility to cloud contamination. An exception occurs during winter when cloud fraction is large in NWP. The MODIS-based DAOD is smaller than CALIOP-based DAOD, even though MODIS TAOD is larger than CALIOP TAOD. Similarly, over the southeastern Atlantic Ocean, CALIOP-based DAOD is also generally smaller than MODIS-based DAOD. On one hand, cloud contamination may have biased the MODIS dust retrieval high. On the other hand, CALIOP clear-sky sampling is not large enough to capture some dust events in this region.

We further compare DAOD (Figure 9) and TAOD (Figure S7 in the supplementary) retrievals from CALIOP and MODIS over major dust laden regions (as shown in Figure 8), including three source regions on land (i.e., Sahara Desert, Middle East and Eastern Asia) and six oceanic outflow regions (i.e., the Tropical Atlantic Ocean - TAT, the Caribbean Basin - CRB, the Mediterranean Sea - MED, the Northwest Pacific Ocean - NWP, the Arabian Sea - ARB as well as the tropical Indian Ocean and the Bay of Bengal - IND). Each data point in the scatter plot represents a monthly mean DAOD (or TAOD) in a $2^\circ \times 5^\circ$ grid. The density of data is represented by different color. To avoid our analysis being biased by some extreme and rare cases, we exclude those data points within the lowest 5% of data density (grey points in Figure 9 and Figure S7). Overall, the DAOD from the two instruments correlate well ($R > 0.75$) and on average CALIOP-based DAOD is 18%, 34%, 54% and 31% lower than MODIS-based DAOD over the Sahara (Figure 9(a)), TAT (Figure 9(d)), CRB(Figure 9(e)) and ARB(Figure 9(h)) regions, respectively. Applying LR of 58sr to Sahara dust reduces the difference from 18% to 8% over the Sahara and from 34% to 12% over TAT. Over the Sahara Desert, the good agreement in DAOD between the two sensors (bias of 8% and $R = 0.78$) suggests that over the Sahara Desert dust particles can be adequately characterized by both irregular non-spherical shape and coarse size. As a result, both CALIOP- and MODIS-based methods are able to detect and separate the dust from other types of aerosols. In TAT and ARB regions, two instruments correlate well ($R > 0.8$) in both DAOD and TAOD. For TAOD, CALIOP is smaller than MODIS by 2% in TAT and larger than MODIS by 15% in ARB. Differences in DAOD are larger, with CALIOP DAOD lower than the MODIS DAOD by 12% and 31% in TAT and ARB, respectively. This suggests that the differences in DAOD from the two instruments are mainly resulted from differences in the dust separation method. In East Asia and NWP, on contrast, both TAOD and DAOD show

poor correlation between the two methods (Figure 9(c), 9(g), S7(c) and S7(g)). As discussed earlier, the poor correlation between the two methods may be contributed by many factors. For example, the total TAOD retrievals from MODIS are subject to larger uncertainties due to cloud contamination, or the DAOD retrieval from CALIOP may miss spherical dust particles that are coated by large combustion emissions from East Asia.

4.2 Comparison between CALIOP and MODIS DAOD Seasonality

Figure 10 compares annual cycle of MODIS and CALIOP DAOD based on the 13-year (2007-2019) average over the nine dust laden regions. Each data point represents domain-averaged 13-year mean DAOD for a month while the error bar indicates $\pm 1\sigma$ (one standard deviation of DAOD). The seasonal cycles of dust activities and dust transport are consistent with results in literature. For example, Prospero et al. 2002 shows that dust activity peaks in May-July in North Africa and Middle East, while peaks in spring in China. These seasonal cycles are consistent with our results shown in the first row of Figure 10. Yu et al. 2015a shows that DAOD peaks in June-July-August in La Parguera, which is consistent with the seasonal cycle in CRB in this study. Generally, CALIOP and MODIS show very similar seasonality over those dust laden regions. DAOD peaks in summer June-July-August (JJA) over Sahara Desert, Middle East, TAT, CRB, ARB and IND, but in spring March-April-May (MAM) over Eastern Asia, MED and NWP. Over NWP, the seasonal cycle of MODIS DAOD is somewhat different from that of CALIOP DAOD. While CALIOP DAOD peaks in spring, MODIS DAOD shows a peak in late spring or even summer months for some years. This could have resulted from cloud contamination in MODIS retrievals due to the large cloud fraction in summer [Yu et al., 2020].

Compared to the MODIS dust retrieval, CALIOP has a unique capability of detecting dust aerosol vertical distribution. Figure 11 shows seasonal mean dust extinction vertical profile

from CALIOP for the nine dust-laden regions. The values on each plot represent the seasonal
680 mean DAOD. Both DAOD and dust vertical structure have a seasonal dependence. In Sahara (a),
Middle East (b) and their dust outflow regions the Tropical Atlantic (d) and the Arabian Sea (h),
summertime dust aerosol has the highest DAOD and reaches to the highest altitude extending
from surface up to 6km in altitude.

The analysis above has been performed over the broad dust-laden regions. Here we focus
685 on MODIS and CALIOP comparison in major potential source areas (PSAs) for dust in North
Africa, namely NAF-1 to NAF-6 as illustrated in Figure 12 (adapted from Fig. 1 in Formenti et
al., 2011). Among all dust source regions around the globe, the Sahara Desert and its margins in
North Africa are the largest dust emitter. Within this region, prominent dust sources are often
associated with topographical lows and foothills of mountains (Prospero et al. 2002). Seasonal
690 variations of DAOD in the six PSAs are shown in Figure 13. Two B values are shown in the
upper left of each panel in Figure 13, where B is defined as the average of CALIOP DAOD /
MODIS DAOD ratios of all data pairs. $B=1$, >1 , <1 indicates no bias, high bias and low bias.
They are calculated based on CALIOP DAOD using dust LR of 44sr and 58sr respectively. The
CALIOP DAOD derived using larger LR (58sr) achieve a better agreement (B values are closer
695 to 1) with MODIS DAOD. Striking CALIOP and MODIS differences in DAOD exist in NAF-5
where the mean bias (B) deviate far from 1. NAF-5 (14N-20N, 15E-20E) is located in Bodélé
Depression, Western Chad. This region is reported as the most intense dust source in the world
(Prospero et al. 2002), and dust activity in the region occurs with a high frequency during all
seasons except fall (Mbourou et al., 1997). However, CALIOP DAOD are much smaller than
700 MODIS retrievals in this region. In terms of dust seasonality (Figure 13), the MODIS DAOD
indicates intense dust aerosol loading all year long with a lower DAOD in Fall, while CALIOP

shows a more distinct seasonality with the highest DAOD of about 0.3 in May-July and the lowest DAOD of <0.1 in winter. Over other PSAs in North Africa, MODIS and CALIOP DAOD show similar seasonality with B closing to 1 (Figure 13).

705 In summary, MODIS and CALIOP DAOD show largest differences under the following conditions: (1) highly cloudy oceanic regions and (2) dust-pollution internal mixtures with high relative humidity. Their differences can be explained as follows.

1. Over cloudy ocean, effective cloud screening is critical to the quality of aerosol retrievals.

710 As an active sensor, CALIOP is more reliable in discriminating clouds and aerosols than passive imager MODIS. In addition, active sensor is able to avoid impact from cloud side scattering. Therefore, MODIS is subject to more cloud contamination than CALIOP. Large cloud contamination in MODIS results in overestimation in TAOD and underestimation in FMF, introducing a high bias in DAOD over ocean cloudy regions (e.g., NWP).

- 715 2. Pure dust particles are hydrophobic and will not absorb water vapor. However, for dust aerosols coated by other types of aerosols (such as the deliquescent dust-nitrate $\text{Ca}(\text{NO}_3)_2$) and saline mineral dust particles emitted from saline topsoil in arid and semiarid areas (Tang et al. 2019), those types of dust particles will take up water vapor and grow to be larger in size and more spherical in shape (Wu et al. 2020). This phenomenon is most prominent for dust aerosols in polluted region (e.g., EAS) as well as with relatively high relative humidity. While such coarse spherical dust particles will not be accounted as dust in CALIOP shape-based method, they are categorized as dust in the MODIS size-based method.
- 720

4.3 DAOD Inter-annual variation from CALIOP and MODIS observations

In this section we examine the inter-annual variation of DAOD captured by two sensors over several major dust source and outflow regions. Figure 14 shows a global map of DAOD interannual trend derived based on the 13-year (2007-2019) time series of annual mean DAOD from CALIOP and MODIS. DAOD trend are calculated for each $2^{\circ} \times 5^{\circ}$ grid. Red color indicates positive trend and blue negative trend. Regions where the trend is statistically significant ($p < 0.05$) are marked with symbol '+'. The similar trend map for total aerosol optical depth is shown in Figure S8 in the supplementary. Overall, DAOD global pattern of interannual trend is similar to TAOD in major dust-laden regions. For example, Over Sahara Desert and tropical Atlantic Ocean region, both CALIOP and MODIS do not show statistically significant trend in DAOD and TAOD. In East Asia and the northwest Pacific Ocean, both sensors show negative trend in DAOD and TAOD.

Figure 15 displays interannual variability of annual-mean DAOD for the major dust-laden regions as defined in Figure 8. Seasonal and annual DAOD trends in the nine regions are listed in Table 5. Both MODIS and CALIOP show a clear DAOD trend in certain seasons over the Eastern Asia, ARB and NWP regions. In Eastern Asia, MODIS and CALIOP show a consistent DAOD decreasing trend at a rate of $-1.7\% \text{ yr}^{-1}$ annually. The two sensors show a DAOD decreasing trend of $-3.5\% \text{ yr}^{-1}$ and $-2.5\% \text{ yr}^{-1}$ respectively in Eastern Asia during spring and show a consistent trend of DAOD in ARB during the fall, though with a factor of 2 difference in magnitude. In NWP, both MODIS- and CALIOP-based DAOD shows a decreasing trend of $-1.7\% \text{ yr}^{-1}$ and $-1.6\% \text{ yr}^{-1}$, respectively. The annual DAOD decreasing trend in NWP is mainly attributed to the DAOD decline in spring at a rate of $-2.3\% \text{ yr}^{-1}$ and $-3.0\% \text{ yr}^{-1}$ for MODIS and CALIOP, respectively. For comparison, Shimizu et al. (2017) detect the decreasing

DAOD trends of $-4.3\% \text{ yr}^{-1}$ in spring and $-2.5\% \text{ yr}^{-1}$ on annual mean basis from the Asian Dust Network (AD-Net) lidar observations over Japan (2007-2016). These trends are greater
750 than our results based on MODIS and CALIOP data records.

Dust over NWP comes mainly from East Asian dust sources. The broad East Asian region (ESA defined in Figure 12) shows statistically significant DAOD decreasing trends (Figure 15c) which is consistent with the DAOD decreasing trend in NWP. It is also imperative to further examine which of six major PSAs in East Asia (ESA-1 to ESA-6 in Figure 12) contribute to the
755 decreasing trend of DAOD. As shown in Figure 16, among the six PSAs, the satellite data show consistent interannual declining trend of DAOD in EAS-5 (Southern Gobi Desert) at a rate of $-4.8\% \text{ yr}^{-1}$ and $-2.8\% \text{ yr}^{-1}$ for MODIS and CALIOP, respectively. In spring, DAOD in EAS-5 shows a significantly declining trend at a rate of $-5.6\% \text{ yr}^{-1}$ and $-3.3\% \text{ yr}^{-1}$ for MODIS and CALIOP (Figure S9). Figure 17 assesses the correlation between DAOD in EAS-5 and
760 DAOD in NWP based on MODIS and CALIOP, respectively. For annual mean DAOD from 2007 to 2019, both sensors show a good correlation between EAS-5 and NWP with $R \approx 0.6$ ($p = 0.02$). In spring, the correlation of DAOD from two regions is good based on CALIOP ($R = 0.6, p = 0.03$), while a weaker correlation ($R = 0.53, p = 0.07$) was found based on MODIS. We further examine potential factors contribute to the declining trend of DAOD in
765 ESA-5. The first row in Figure 18 shows the springtime trend of MODIS enhance vegetation index (EVI), MERRA2 near-surface (at 10 m) wind speed and precipitation in EAS-5 region (Qian et al. 2002; Kurosaki and Mikami 2003; Lee and Sohn 2011). EVI and precipitation show a statistically significant ($p < 0.05$) increasing trend with $R = 0.82$ and $R = 0.58$, respectively. Surface wind speed shows a statistically significant ($p < 0.05$) decreasing trend with $R = -0.66$.

770 The second row in Figure 18 shows the correlations of the three factors with MODIS DAOD and

CALIOP DAOD, respectively. Clearly, EVI is anti-correlated with both MODIS and CALIOP DAOD with $|R| > 0.7$ and $p < 0.05$. Surface wind speed is correlated with MODIS DAOD and CALIOP DAOD with $|R| > 0.6$ and $p < 0.05$. While the correlation with precipitation is not statistically significant ($p > 0.05$). Note that EVI and surface wind speed are not independent variables that affect dust emissions. An increase of EVI or vegetation cover could reduce the surface wind speed. However, given the relatively coarse resolution of MERRA2, the surface wind speed trend may largely reflect the change in atmospheric circulations other than local wind decrease induced by more vegetation. The precipitation shows no statistically significant correlation with MODIS and CALIOP DAOD.

As discussed earlier, our results suggest that the decrease of NWP DAOD is likely a result of the decreasing dust events in Asian deserts (i.e., EAS-5 Gobi) in turn likely due to change of vegetation. This is also reported in several recent studies. Sternberg et al. (2015) found that Gobi Desert contracted from 2000 to 2012 due to increased moisture availability. Song et al. (2016) used an Integrated Wind Erosion Modeling System to simulate the spring dust emissions in northern China over the period of 1982 to 2011. They found a significant decrease of the magnitude of spring dust event in China which is attributed to both climate change and local mitigation strategies. Similarly, An et al., (2018) also noted a significant decrease of dust storm event in East Asian after analyzing observational data from ground stations, numerical modeling, and vegetation indices obtained from both satellite and reanalysis data. Over the last few decades, The Chinese government has been taking actions to restore overgrazed land in Inner Mongolia, the enlarged vegetation coverage and the expected earlier vegetation green-up due to global warming could have mitigated dust activity in this region (Fan et al. 2014). Together the results

from our analysis, along with the aforementioned recent studies, suggest that the decreasing
795 springtime DAOD trend in the NWP region is a result of the decline of dust activities in the Inner
Mongolia (i.e., EAS-5) which is likely linked to vegetation coverage changes in recent years as a
result of China's mitigation projects to hold back desertification.

Some caveats must be mentioned, however, when interpreting the trend analysis here. First
of all, due to the limitation of satellite data record, we have only 13 years' CALIOP data and 17
800 years' MODIS data available. Other climate variabilities, such as the El Nino-Southern
Oscillation (ENSO), could confound the trend analysis. For example, Abish and Mohanakumar
(2013) shows that La Nina (El Nino) weakens (strengthens) the zonal circulation over the Indian
subcontinent, which result in low (high) aerosol concentration over Indian subcontinent
transported from Arabian Desert over the period. Gong et al. (2005) also shows the impact of
805 ENSO on the interannual variability of Asian dust loading and deposition. According to the
NOAA Oceanic Nino Index (ONI), the climate switched from a strong La Niña phase in 2010-
2011 to a strong El Niño phase in 2015-2016. However, the potential impact of ENSO on the
dust inter-annual variability is beyond the scope of this study and will be left for the future
research.

810 5 Summary and Conclusion

We derive two observation-based global monthly mean dust aerosol optical depth
(DAOD) climatological datasets from 2007 to 2019 with a 2° (latitude) ×5° (longitude) spatial
resolution, one based on CALIOP and the other on MODIS observations. Our product captures
815 very well as much hot spots along the 'dust belt' region well, as weaker signals in other dust
active regions such as Southwestern United States, Patagonian Desert in South America, Central

Australia, and South Africa (Figure 5). Since DAOD climatology product contains and mixes the information of the intensity and frequency of dust activities, we introduce the conditional DAOD product, which diminishes impacts from dust frequency by excluding dust-free cases in the average. The comparison between DAOD climatology data and conditional DAOD data suggests that dust activities in those regions are highly episodic. The two data records compare reasonably well with the results reported in previous studies and the collocated AERONET coarse model AOD. The comparison of our MODIS-based and CALIOP-based DAOD with AERONET COD indicates that MODIS overestimates DAOD, while CALIOP underestimates DAOD. It is highly probably that the true DAOD fall between MODIS and CALIOP DAOD.

CALIOP distinguishes dust aerosols based on its non-spherical shape, whereas MODIS separates dust aerosols from others based on its large size characteristics. The discrepancy in dust retrieval based on two instruments are expected due to the uncertainty associated with their TAOD retrieval and the uncertainty associated with their different mechanism in dust detection and separation. The comparison between CALIOP dust retrieval and MODIS dust retrieval facilitate a better understanding of advantages and limitations of each dust product and also provide some insights on dust morphology and dust size. Through the comparison, we found generally CALIOP-based DAOD correlates well with MODIS-based DAOD over dust-laden regions such as Sahara ($R=0.78$), TAT ($R=0.84$), CRB ($R=0.75$) and ARB ($R=0.85$), but with CALIOP-based DAOD 18%, 34%, 54% and 31% lower than MODIS-based DAOD over those regions respectively. This result is consistent with the different treatment of the dust-pollution mixtures in the dust separation approaches of two instruments. Applying LR of 58sr to Sahara dust reduce the difference from 18% to 8% over the Sahara and from 34% to 12% over TAT.

840 Over the Sahara Desert, the good agreement in DAOD between the two sensors (bias of 8% and
R = 0.78) suggests that dust aerosols are irregular non-spherical and at the same time large in size
in this region. In some regions such as NWP, the DAOD correlation between two sensors is quite
low. There could be many reasons for this, for example, the total TAOD retrievals from MODIS
have larger uncertainty due to cloud contamination, or the DAOD retrieval from CALIOP may
845 miss coarse spherical dust-pollution mixtures.

The interannual variability of DAOD over dust-laden regions show a clear trend in
Eastern Asia a rate of $-1.7\% \text{ yr}^{-1}$ based on two sensors. Over the outflow region of Easter Asia,
DAOD in NWP region shows a clear trend at a rate of $-1.6\% \text{ yr}^{-1}$ and $-1.7\% \text{ yr}^{-1}$ based on
CALIOP and MODIS respectively, this trend is mainly attributed to the decreasing trend in
850 spring with a rate of $-3.0\% \text{ yr}^{-1}$ based on CALIOP and $-2.3\% \text{ yr}^{-1}$ based on MODIS.
Further investigation of DAOD trend in six dust source areas in Eastern Asia where NWP dust
aerosols come from shows that there is an obvious decreasing trend in DAOD during 2007 -
2019 over Southern Gobi Desert based on both CALIOP and MODIS dust retrievals. The
decreasing trend of DAOD is correlated significantly with the vegetation index and surface wind
855 speed in the area, whereas there is almost no correlation with the precipitation.

Data availability. The global DAOD and dust vertical extinction coefficient climatology data derived from CALIOP in this study and the MODIS DAOD retrieval data over land and ocean are available at

[‘https://drive.google.com/drive/folders/1aQVupe7govPwR6qmsqUbR4fJQsp1DBCX?usp=sharing’](https://drive.google.com/drive/folders/1aQVupe7govPwR6qmsqUbR4fJQsp1DBCX?usp=sharing). The MODIS Enhanced Vegetation Index (EVI) data could be downloaded from [‘https://lpdaac.usgs.gov/products/myd13c2v006/#tools’](https://lpdaac.usgs.gov/products/myd13c2v006/#tools). The MERRA2 surface wind speed and precipitation data are available at

[‘https://disc.sci.gsfc.nasa.gov/datasets/M2T1NXFLX_5.12.4/summary?keywords=%22MERRA-2%22’](https://disc.sci.gsfc.nasa.gov/datasets/M2T1NXFLX_5.12.4/summary?keywords=%22MERRA-2%22).

Acknowledgement. Qianqian Song and Zhibo Zhang cordially acknowledge the funding support from the Future Investigators in NASA Earth and Space Science and Technology (FINESST).

Zhibo Zhang’s research is supported by NASA grant (80NSSC20K0130) from the CALIPSO and CloudSat program managed by Dr. David Considine. HY was supported by NASA’s the Science of Terra, Aqua, and Suomi-NPP and the CALIPSO/CloudSat Science Team programs administered by Dr. Hal Maring and Dr. David Considine, respectively. The computations in this study were performed at the UMBC High Performance Computing Facility (HPCF). The facility is supported by the US National Science Foundation through the MRI program (grant nos. CNS-0821258 and CNS-1228778) and the SCREMS program (grant no. DMS-0821311), with substantial support from UMBC. The MODIS aerosol data were obtained from the NASA Level-1 and Atmosphere Archive and Distribution System (LAADS) webpage (<https://ladsweb.nascom.nasa.gov/>). The CALIOP aerosol products were obtained from NASA Langley Research Center Atmospheric Science Data Center (<https://eosweb.larc.nasa.gov/>).

Table 1. Summary of DAOD retrievals from MODIS and CALIOP

Sensors	Retrieve Scope	Relevant variables used to derive DAOD	References
MODIS	Ocean	AOD, fine-mode AOD	Yu et al. (2009, 2020)
MODIS	Land	AOD, SSA at 470nm, Angstrom exponent	Pu and Ginoux et al. (2018)
CALIOP	Globe	Profiles of backscatter, extinction, depolarization ratio	Yu et al. (2015a)

885 Table 2. Compare global mean DAOD retrievals in this study with some relevant studies (Note the definition of global scope is different for different studies).

Region		DAOD@550nm	Reference
90°S~90°N	Global	0.03±0.005	Ridley et al. 2016 Use multiple satellite platforms, in-situ AOD observations and four global models
90°S~90°N	Global	0.033	Gkikas et al 2021 Use AOD from Aqua MODIS and DOD-to-AOD ratio from MERRA2
50°S~60°N	Over Ocean	0.03±0.06	Voss and Evan 2020 Over Ocean: use method in Kaufman et al 2005 Over Land: use method in Ginoux et al. 2012
	Over Land	0.1	
60°S~60°N	Over Ocean	0.055, 0.020	This Study MODIS-based, CALIOP-based DAOD (To calculate global mean DAOD for scope 90°S~90°N, we assume zero DAOD outside of region 60°S~60°N. We weight each grid-cell surface area into ocean, land and global DAOD average)
	Over Land	0.103, 0.068	
90°S~90°N	Global	0.057, 0.028	

890

Table 3 Statistical parameters and absolute error by continents using the method indicated in Figure 1. Sites is the number of AERONET sites involved; N is the number of MODIS, CALIOP and AERONET matchups. R is correlation coefficient; C is the intercept of the linear fit; K is the slope of the linear fit; RMSE is root mean square error of the linear fit; Ba is the absolute bias; Br is the relative bias. For cells with two rows of values, the upper row is for MODIS, the lower row is for CALIOP.

Region	Sites	N	R	C	K	RMSE	Ba	Br (%)	Absolute Error
Global	761	16653	0.72	0.01	1.05	0.08	0.01	26.7	$0.65 \times \text{DAOD}_M$
			0.70	-0.01	0.90	0.07	-0.02	-27.9	$0.52 \times \text{DAOD}_C + 0.02$
Africa	44	706	0.79	0.04	0.72	0.10	0.01	4.5	$0.37 \times \text{DAOD}_M + 0.01$
			0.72	0.01	0.75	0.12	-0.02	-19.8	$0.51 \times \text{DAOD}_C + 0.02$
Asia	143	2507	0.64	0.04	0.88	0.10	0.03	34.2	$0.61 \times \text{DAOD}_M$
			0.57	0.00	0.84	0.11	-0.01	-11.0	$0.66 \times \text{DAOD}_C + 0.01$
Europe	156	4359	0.27	0.03	0.55	0.05	0.01	18.2	$0.70 \times \text{DAOD}_M$
			0.35	0.00	0.53	0.04	-0.02	-48.6	$0.47 \times \text{DAOD}_C + 0.02$
Americas	319	6656	0.29	0.02	0.54	0.04	0.01	25.5	$0.77 \times \text{DAOD}_M$
			0.33	0.00	0.31	0.03	-0.02	-55.8	$0.26 \times \text{DAOD}_C + 0.02$
Australia	12	507	0.51	0.0	0.57	0.03	-0.02	-43.9	$0.37 \times \text{DAOD}_M + 0.02$
			0.28	0.0	0.32	0.04	-0.02	-59.3	$0.34 \times \text{DAOD}_C + 0.03$

895

Table 4. Global (60° S-60° N) seasonal mean DAOD and TAOD based on MODIS and CALIOP (2007~2019) dust retrievals. Since Earth is a sphere, grid-cell surface area decreases toward the poles. We weight each grid-cell surface area into ocean, land and global DAOD average.

		MAM		JJA		SON		DJF		Annual	
		DAOD	TAOD	DAOD	TAOD	DAOD	TAOD	DAOD	TAOD	DAOD	TAOD
MODIS	Ocean	0.057	0.151	0.062	0.153	0.047	0.143	0.052	0.144	0.055	0.148
	Land	0.131	0.283	0.119	0.270	0.079	0.206	0.085	0.217	0.103	0.244
	Global	0.075	0.183	0.077	0.183	0.055	0.159	0.059	0.160	0.067	0.171
	Land /Ocean	2.27	1.87	1.90	1.77	1.67	1.44	1.64	1.51	1.89	1.65
CALIOP	Ocean	0.022	0.098	0.025	0.104	0.015	0.092	0.016	0.090	0.020	0.096
	Land	0.086	0.196	0.086	0.228	0.051	0.186	0.047	0.157	0.068	0.192
	Global	0.039	0.124	0.041	0.137	0.025	0.117	0.024	0.107	0.032	0.121
	Land /Ocean	3.90	1.99	3.45	2.20	3.40	2.03	2.87	1.74	3.45	2.00

900

Table 5. DAOD inter-seasonal trend over major dust-laden regions based on MODIS and CALIOP observations. The changing rate of DAOD trend is shown in a sequence of annual/spring/summer/fall/winter in each cell of the table. Those statistically meaningful trends with $p < 0.05$ are shown in bold.

	MODIS [% yr ⁻¹]					CALIOP [% yr ⁻¹]				
	Annual	MAM	JJA	SON	DJF	Annual	MAM	JJA	SON	DJF
Sahara Desert (a)	-0.04	-0.84	0.21	0.29	0.51	-0.09	-0.93	0.34	-0.52	0.55
Middle East (b)	0.32	-0.61	-0.02	1.80	1.37	-1.84	-2.36	-1.86	-2.46	-0.09
Eastern Asia (c)	-1.74	-3.48	-0.28	-0.33	-0.56	-1.70	-2.46	-1.99	-0.45	-1.42
TAT (d)	0.34	-0.68	-0.03	1.68	1.32	-0.25	-1.41	-0.07	0.91	-0.09
CRB (e)	1.10	0.78	0.94	1.59	1.97	-0.40	-1.39	-0.34	0.79	-1.09
MED (f)	0.10	0.32	0.49	-0.71	0.03	-1.09	-1.07	-1.63	-1.20	-0.52
NWP (g)	-1.67	-2.33	-1.93	0.63	-1.35	-1.58	-3.01	-2.89	-0.40	-0.19
ARB (h)	-1.42	-0.72	-1.81	-1.85	-0.31	-1.17	-1.70	-0.46	-3.60	-0.06
IND (i)	-0.09	-0.51	0.40	0.38	-0.89	-1.96	-2.92	-2.43	-0.21	-0.54

905




Figure 1. Scatter density histogram comparing monthly (from 2007 to 2019) MODIS DAOD (on the left panel) and CALIOP DAOD (on the right panel) with monthly coarse mode COD_{SDA} retrieved at 550nm for MODIS comparison and at 532nm for CALIOP comparison from the Level 2 (cloud screened and quality assured) Spectral Deconvolution Algorithm (SDA) version 4.1 (O'Neill et al., 2003). The 1 to 1 line and linear regression line are shown by dotted and solid lines, respectively. The number of sites (Sites), of matchups (N), correlation (R), slope (S), constant (C), and root mean square error (RMSE) of the linear regression as well as absolute bias (Ba) and relative Bias (Br) are indicated in the lower right of the panel. Ba, Br and RMSE are defined as:

$$B_a = \overline{DAOD}_{C \text{ or } M} - \overline{COD}_{SDA}, B_r = \overline{DAOD}_{C \text{ or } M} / \overline{COD}_{SDA} - 1, RMSE = \sqrt{\frac{\sum_i (\overline{DAOD}_{C \text{ or } M, i} - \overline{COD}_{SDA, i})^2}{N}}$$




Figure 2 Left panel: the x-axis is the MODIS derived DAOD, y-axis is the absolute MODIS DAOD–AERONET COD difference (without scaling by AMF). Data are sorted by bins of 100 values (we have 16653 matchups in total; therefore, the last bin has 53 values). The means and standard deviations of the MODIS DAOD_M are the centers and half widths of the boxes in the horizontal. The mean, medians, and lower to upper quartile interval of the MODIS –AERONET SDA differences are the red dots, the center, and top-bottom intervals of the boxes. The dotted line is the error estimated from the least squares linear fit of the 68th percentiles for each box. The right panel is the same except for CALIOP DAOD.





Figure 3 The same as Figure 1 except for over ocean.



940 Figure 4 The same as Figure 2 except for over ocean.




Figure 5. Spatial distribution of the seasonal mean CALIOP-based DAOD, MODIS-based DAOD and the fraction of DAOD with respect to the TAOD based on CALIOP and MODIS respectively for the globe at a $5^\circ \text{ longitude} \times 2^\circ \text{ latitude}$ resolution based on 13-year (2007-2019) CALIOP measurements. **DJF**: December from previous year-January-February; **MAM**: March-April-May; **JJA**: June-July-August; **SON**: September-October-November.




Figure 6. Conditional DAOD (the first column), climatological DAOD (the second column) based on CALIOP dust retrieval from 2007 to 2019. The third column shows the relative difference between conditionally sampled DAOD and climatological DAOD with respect to the climatological DAOD expressed in fraction.

955




Figure 7. The difference between MODIS and CALIOP for seasonal mean TAOD (the first column), DAOD (the second column), and the fraction of DAOD in TAOD (the third column) on a basis of 13-year (2007-2019) average. The fourth column is the seasonal mean cloud fraction from MODIS L3 product.




Figure 8. Major dust-laden regions including three dust source regions on land (a ~ c) and six outflow regions over ocean (e ~ i). (a) Sahara Desert (14°N-30°N, 15°W-30°E), (b) Middle East (10°N-35°N, 40°E-85°E) and (c) Eastern Asia (30°N-50°N, 75°E-130°E) (d) the tropical Atlantic Ocean–TAT (0°-30°N, 15°W-60°W), (e) the Caribbean Sea–CRB (6°N-22°N, 60°W-90°W), (f) the Mediterranean Sea–MED (30°N-46°N, 5°W-35°E), (g) the northwest Pacific Ocean–NWP (30°N-55°N, 120°E-160°E), (h) the Arabian Sea–ARB (0°-26°N, 45°E-80°E) and (i) the tropical Indian Ocean and the Bay of Bengal–IND (10°S-22°N, 75°E-100°E). Note we only consider grids over land for the three dust source regions and grids over ocean for the six dust outflow regions.




Figure 9. Comparison of CALIOP DAOD against MODIS DAOD over dust-laden regions indicated in Figure 8. Color represents the probability density using gaussian kernel density estimation. Grey points represent data points within the lowest 5% of data density. Those grey points are excluded in the linear regression analysis. The blackline and blue shadow are the linear regression for $LR=44 \pm 9sr$, the red line and red shadow in (a) and (d) represent the linear regression for $LR=58 \pm 8sr$. Red text in panel (a) and (d) is the linear fit based on $LR=58 sr$. Black text in each panel is the linear fit based on $LR=44 sr$. R is Pearson's linear correlation coefficient between MODIS and CALIOP DAOD.




Figure 10. Monthly variation of DAOD from CALIOP (green) and MODIS (red) for major dust-laden regions indicated in Figure 8. Vertical line represents ± 1 sigma (standard deviation) over the 13-year period.




Figure 11. Vertical profiles of seasonal mean dust extinction coefficient (Mm^{-1}) in 9 dust-laden regions indicated in Figure 8. Different colors represent different seasons. The numbers on each plot are the seasonal mean DAOD for the region.




Figure 12. Six dust potential source subregions in Northern Africa (NAF) and Eastern Asia (EAS) based on Fig.1. and Fig. 2. in Formenti, et al., 2011. PSA NAF-1(30N-36N, 0-9E), PSA NAF-2 (16N-28N, 10W-15W), PSA NAF-3 (18N-26N, 5W-5E), PSA NAF-4 (24N-30N, 15E-20E), PSA NAF-5 (14N-20N, 15E-20E), PSA NAF-6 (14N-24N, 25E-35E); EAS-1: (34N-40N, 75E-90E) ; EAS-2: (44N-46N, 85E-90E); EAS-3: (40N-42N,90E-95E and 42N-44N, 85E-90E); EAS-4: (42N-46N, 100E-115E); EAS-5: (38N-42N, 100E-110E); EAS-6: (42N-46N, 115E-125E and 48N-50N, 115E-120E)




Figure 13. Annual cycle of 13-year (2007-2019) monthly mean DAOD over the six PSAs of North African dust. The CALIOP DAOD annual cycle shown in the figure is derived from backscatter coefficients using LR of 44sr. The mean bias (B) is computed as the average of CALIOP DAOD / MODIS DAOD ratios of all data pairs. B =1, >1, <1 indicates no bias, high bias and low bias. The mean bias (B) associated with CALIOP DAOD based on LR=44sr and 58sr are shown in the upper left of each panel.




Figure 14. Global map of DAOD interannual trend based on CALIOP (left) and MODIS (right) dust climatology data over 2007-2019 period. Red and blue represents increasing and decreasing trend, respectively. Symbol '+' denotes trends with p-value < 0.05, which are considered as statistically meaningful trend.




Figure 15. DAOD interannual variability over main dust source regions (a-c) and dust outflow regions (d-i) revealed by CALIOP (green curve) and MODIS (red curve) observations.




Figure 16. Interannual variability of CALIOP (green) and MODIS (red) DAOD in the six potential dust source areas in Eastern Asia (refer to Figure 12).




Figure 17. Correlation between DAOD in EAS-5 (Southern Gobi Desert) and DAOD in NWP for annual mean (left) and springtime average (right).

1040




Figure 18. The inter-annual trend of Enhance Vegetation Index (EVI), surface wind speed and precipitation and their correlation with DAOD in spring, EAS5 region. The 1st row shows inter-annual trend of EVI, surface wind speed and precipitation. The 2nd shows the correlation of EVI, surface wind speed and precipitation with MODIS-based DAOD and CALIOP-based DAOD respectively. In addition, the time series of EVI versus DAOD, wind speed versus DAOD, precipitation versus DAOD is shown in Figure S10 in the supplement.

1045

References

- Abish, B. and K. Mohanakumar. 2013. "Absorbing Aerosol Variability over the Indian Subcontinent and Its Increasing Dependence on ENSO." *Global and Planetary Change* 106:13–19.
- Ackerman, A. S., O. B. Toon, D. E. Stevens, A. J. Heymsfield, V. Ramanathan, and E. J. Welton. 2000. "Reduction of Tropical Cloudiness by Soot." *Science* 288(5468):1042–47.
- Albrecht, B. A. 1989. "Aerosols, Cloud Microphysics, and Fractional Cloudiness." *Science* 245(4923):1227–30.
- Amiridis, V., U. Wandinger, E. Marinou, E. Giannakaki, A. Tsekeri, S. Basart, S. Kazadzis, A. Gkikas, M. Taylor, J. Baldasano, and A. Ansmann. 2013. "Optimizing CALIPSO Saharan Dust Retrievals." *Atmospheric Chemistry and Physics* 13(23):12089–106.
- An, Linchang, Huizheng Che, Min Xue, Tianhang Zhang, Hong Wang, Yaqiang Wang, Chunhong Zhou, Hujia Zhao, Ke Gui, and Yu Zheng. 2018. "Temporal and Spatial Variations in Sand and Dust Storm Events in East Asia from 2007 to 2016: Relationships with Surface Conditions and Climate Change." *Science of The Total Environment* 633:452–62.
- Anderson, Theodore L., Yonghua Wu, D. Allen Chu, Beat Schmid, Jens Redemann, and Oleg Dubovik. 2005. "Testing the MODIS Satellite Retrieval of Aerosol Fine-Mode Fraction." *Journal of Geophysical Research D: Atmospheres* 110(18):1–16.
- Ansmann, A., P. Seifert, Matthias Tesche, and U. Wandinger. 2012. "Profiling of Fine and Coarse Particle Mass: Case Studies of Saharan Dust and Eyjafjallajökull/Grimsvötn Volcanic Plumes." *Atmospheric Chemistry and Physics*.
- Ansmann, A., M. Tesche, P. Seifert, S. Groß, V. Freudenthaler, A. Apituley, K. M. Wilson, I. Serikov, H. Linné, B. Heinold, A. Hiebsch, F. Schnell, J. Schmidt, I. Mattis, U. Wandinger, and M. Wiegner. 2011. "Ash and Fine-Mode Particle Mass Profiles from EARLINET-AERONET Observations over Central Europe after the Eruptions of the Eyjafjallajökull Volcano in 2010." *Journal of Geophysical Research Atmospheres* 116(12).
- Baars, Holger, Thomas Kanitz, Ronny Engelmann, Dietrich Althausen, Birgit Heese, Mika Komppula, Jana Preißler, Matthias Tesche, Albert Ansmann, and Ulla Wandinger. 2016. "An Overview of the First Decade of PollyNET: An Emerging Network of Automated Raman-Polarization Lidars for Continuous Aerosol Profiling."
- Di Biagio, C., Y. Balkanski, S. Albani, O. Boucher, and P. Formenti. 2020. "Direct Radiative Effect by Mineral Dust Aerosols Constrained by New Microphysical and Spectral Optical Data." *Geophysical Research Letters* 47(2):e2019GL086186.
- Boucher, O., D. Randall, P. Artaxo, C. Bretherton, G. Feingold, P. Forster, V.-M. Kerminen, Y. Kondo, H. Liao, U. Lohmann, P. Rasch, S.K. Satheesh, S. Sherwood, B. Stevens and X. Y. Zhang. 2013. "Clouds and Aerosols." *Climate Change 2013 The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.
- Burton, S. P., R. A. Ferrare, C. A. Hostetler, J. W. Hair, R. R. Rogers, M. D. Obland, C. F. Butler, A. L. Cook, D. B. Harper, and K. D. Froyd. 2012. "Aerosol Classification Using Airborne High Spectral Resolution Lidar Measurements-Methodology and Examples." *Atmospheric Measurement Techniques* 5(1):73.

- Chen, Cheng, Oleg Dubovik, Daven K. Henze, Tatyana Lapyonak, Mian Chin, Fabrice Ducos, Pavel Litvinov, Xin Huang, and Lei Li. 2018. "Retrieval of Desert Dust and Carbonaceous Aerosol Emissions over Africa from POLDER/PARASOL Products Generated by the GRASP Algorithm." *Atmospheric Chemistry and Physics* 18(16):12551–80.
- 1100 Chimot, Julien, J. Pepijn Veefkind, Tim Vlemmix, Johan F. de Haan, Vassilis Amiridis, Emmanouil Proestakis, Eleni Marinou, and Pieternel F. Levelt. 2017. "An Exploratory Study on the Aerosol Height Retrieval from OMI Measurements of the 477 Nm O.Sub.2 - O.Sub.2 Spectral Band Using a Neural Network Approach." *Atmospheric Measurement Techniques* 10:783.
- 1105 Clarisse, Lieven, Cathy Clerbaux, Bruno Franco, Juliette Hadji-Lazaro, Simon Whitburn, A. K. Kopp, Daniel Hurtmans, and P-F Coheur. 2019. "A Decadal Data Set of Global Atmospheric Dust Retrieved from IASI Satellite Measurements." *Journal of Geophysical Research: Atmospheres* 124(3):1618–47.
- 1110 Dubovik, O., A. Sinyuk, T. Lapyonok, B. N. Holben, M. Mishchenko, P. Yang, T. F. Eck, H. Volten, O. Munoz, B. Veihelmann, W. J. van der Zande, J. F. Leon, M. Sorokin, and I. Slutsker. 2006. "Application of Spheroid Models to Account for Aerosol Particle Nonsphericity in Remote Sensing of Desert Dust." *Journal of Geophysical Research-Atmospheres* 111(D11).
- 1115 Eck, T. F., B. N. Holben, J. S. Reid, O. Dubovik, A. Smirnov, N. T. O'Neill, I. Slutsker, and S. Kinne. 1999. "Wavelength Dependence of the Optical Depth of Biomass Burning, Urban, and Desert Dust Aerosols." *Journal of Geophysical Research Atmospheres* 104(D24):31333–49.
- 1120 Esselborn, Michael, Martin Wirth, Andreas Fix, Bernadett Weinzierl, Katharina Rasp, Matthias Tesche, and Andreas Petzold. 2009. "Spatial Distribution and Optical Properties of Saharan Dust Observed by Airborne High Spectral Resolution Lidar during SAMUM 2006." *Tellus B: Chemical and Physical Meteorology* 61(1):131–43.
- Evan, A. T., J. Dunion, J. A. Foley, A. K. Heidinger, and C. S. Velden. 2006. "New Evidence for a Relationship between Atlantic Tropical Cyclone Activity and African Dust Outbreaks." *Geophysical Research Letters* 33(19).
- 1125 Fan, Bihang, Li Guo, Ning Li, Jin Chen, Henry Lin, Xiaoyang Zhang, Miaogen Shen, Yuhao Rao, Cong Wang, and Lei Ma. 2014. "Earlier Vegetation Green-up Has Reduced Spring Dust Storms." *Scientific Reports* 4:1–6.
- 1130 Fiebig, Markus, Andreas Petzold, Ulla Wandinger, Manfred Wendisch, Christoph Kiemle, Armin Stifter, Martin Ebert, Tom Rother, and Ulrich Leiterer. 2002. "Optical Closure for an Aerosol Column: Method, Accuracy, and Inferable Properties Applied to a Biomass-burning Aerosol and Its Radiative Forcing." *Journal of Geophysical Research: Atmospheres* 107(D21):LAC-12.
- 1135 Formenti, P., L. Schutz, Y. Balkanski, K. Desboeufs, M. Ebert, K. Kandler, A. Petzold, D. Scheuven, S. Weinbruch, and D. Zhang. 2011. "Recent Progress in Understanding Physical and Chemical Properties of African and Asian Mineral Dust." *Atmospheric Chemistry and Physics* 11(16):8231–56.
- 1140 Gasteiger, Josef, Matthias Wiegner, Silke Groß, Volker Freudenthaler, Carlos Toledano, Matthias Tesche, and Konrad Kandler. 2011. "Modelling Lidar-Relevant Optical Properties of Complex Mineral Dust Aerosols." *Tellus, Series B: Chemical and Physical Meteorology* 63(4):725–41.
- Ge, J. M., J. P. Huang, C. P. Xu, Y. L. Qi, and H. Y. Liu. 2014. "Characteristics of Taklimakan

- Dust Emission and Distribution: A Satellite and Reanalysis Field Perspective.” *Journal of Geophysical Research: Atmospheres* 119(20):11,711-772,783.
- 1145 Gelaro, Ronald, Will McCarty, Max J. Suárez, Ricardo Todling, Andrea Molod, Lawrence Takacs, Cynthia A. Randles, Anton Darmenov, Michael G. Bosilovich, Rolf Reichle, Krzysztof Wargan, Lawrence Coy, Richard Cullather, Clara Draper, Santha Akella, Virginie Buchard, Austin Conaty, Arlindo M. da Silva, Wei Gu, Gi Kong Kim, Randal Koster, Robert Lucchesi, Dagmar Merkova, Jon Eric Nielsen, Gary Partyka, Steven Pawson, William Putman, Michele Rienecker, Siegfried D. Schubert, Meta Sienkiewicz, and Bin
- 1150 Zhao. 2017. “The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2).” *Journal of Climate* 30(14):5419–54.
- Getzewich, Brian J., Mark A. Vaughan, William H. Hunt, Melody A. Avery, Kathleen A. Powell, Jason L. Tackett, David M. Winker, Jayanta Kar, Kam Pui Lee, and Travis D. Toth. 2018. “CALIPSO Lidar Calibration at 532 Nm: Version 4 Daytime Algorithm.”
- 1155 *Atmospheric Measurement Techniques* 11(11):6309–26.
- Ginoux, Paul, Dmitri Garbuzov, and N. Christina Hsu. 2010. “Identification of Anthropogenic and Natural Dust Sources Using Moderate Resolution Imaging Spectroradiometer (MODIS) Deep Blue Level 2 Data.” *Journal of Geophysical Research Atmospheres* 115(5):1–10.
- 1160 Ginoux, Paul, Joseph M. Prospero, Thomas E. Gill, N. Christina Hsu, and Ming Zhao. 2012. “Global-Scale Attribution of Anthropogenic and Natural Dust Sources and Their Emission Rates Based on MODIS Deep Blue Aerosol Products.” *Reviews of Geophysics* 50(3).
- Gkikas, Antonis, Emmanouil Proestakis, Vassilis Amiridis, Stelios Kazadzis, Enza Di Tomaso, Alexandra Tsekeri, Eleni Marinou, Nikos Hatzianastassiou, and Carlos Pérez García-Pando. 2021. “ModIs Dust AeroSol (MIDAS): A Global Fine-Resolution Dust Optical Depth Data Set.” *Atmospheric Measurement Techniques* 14(1):309–34.
- 1165 Gong, S. L., X. Y. Zhang, T. L. Zhao, # X B Zhang, L. A. Barrie, I. G. Mckendry, and C. S. Zhao. 2005. *A Simulated Climatology of Asian Dust Aerosol and Its Trans-Pacific Transport. Part II: Interannual Variability and Climate Connections*.
- Griffin, Dale W. 2007. “Atmospheric Movement of Microorganisms in Clouds of Desert Dust and Implications for Human Health.” *Clinical Microbiology Reviews* 20(3):459 LP – 477.
- 1170 Grousset, Francis E., Paul Ginoux, Aloys Bory, and Pierre E. Biscaye. 2003. “Case Study of a Chinese Dust Plume Reaching the French Alps.” *Geophysical Research Letters* 30(6).
- Hansen, J., M. Sato, and R. Ruedy. 1997. “Radiative Forcing and Climate Response.” *Journal of Geophysical Research-Atmospheres* 102(D6):6831–64.
- 1175 Hayasaka, Tadahiro, Shinsuke Satake, Atsushi Shimizu, Nobuo Sugimoto, Ichiro Matsui, Kazuma Aoki, and Yoshitaka Muraji. 2007. “Vertical Distribution and Optical Properties of Aerosols Observed over Japan during the Atmospheric Brown Clouds-East Asia Regional Experiment 2005.” *Journal of Geophysical Research Atmospheres* 112(22).
- 1180 Hsu, N. C., M. J. Jeong, C. Bettenhausen, A. M. Sayer, R. Hansell, C. S. Seftor, J. Huang, and S. C. Tsay. 2013. “Enhanced Deep Blue Aerosol Retrieval Algorithm: The Second Generation.” *Journal of Geophysical Research: Atmospheres* 118(16):9296–9315.
- Hsu, N. Christina, Si Chee Tsay, Michael D. King, and Jay R. Herman. 2004. “Aerosol Properties over Bright-Reflecting Source Regions.” *IEEE Transactions on Geoscience and Remote Sensing* 42(3):557–69.
- 1185 Huang, Jianping, Patrick Minnis, Bin Chen, Zhongwei Huang, Zhaoyan Liu, Qingyun Zhao, Yuhong Yi, and J. Kirk Ayers. 2008. “Long-Range Transport and Vertical Structure of Asian Dust from CALIPSO and Surface Measurements during PACDEX.” *Journal of*

Geophysical Research: Atmospheres 113(D23).

- 1190 Huang, Jianping, Patrick Minnis, Yuhong Yi, Qiang Tang, Xin Wang, Yongxiang Hu, Zhaoyan Liu, Kirk Ayers, Charles Trepte, and David Winker. 2007. "Summer Dust Aerosols Detected from CALIPSO over the Tibetan Plateau." *Geophysical Research Letters* 34(18).
- 1195 Huang, Yue, Jasper F. Kok, Konrad Kandler, Hannakaisa Lindqvist, Timo Nousiainen, Tetsu Sakai, Adeyemi Adebisi, and Olli Jokinen. 2020. "Climate Models and Remote Sensing Retrievals Neglect Substantial Desert Dust Asphericity." *Geophysical Research Letters* 47(6).
- Järvinen, E., O. Kemppinen, T. Nousiainen, T. Kociok, O. Möhler, T. Leisner, and M. Schnaiter. 2016. "Laboratory Investigations of Mineral Dust Near-Backscattering Depolarization Ratios." *Journal of Quantitative Spectroscopy and Radiative Transfer* 178:192–208.
- 1200 Jickells, T. D., Z. S. An, K. K. Andersen, A. R. Baker, G. Bergametti, N. Brooks, J. J. Cao, P. W. Boyd, R. A. Duce, K. A. Hunter, H. Kawahata, N. Kubilay, J. Laroche, P. S. Liss, N. Mahowald, J. M. Prospero, A. J. Ridgwell, I. Tegen, and R. Torres. 2005. "Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate." *Science* (308):67–71.
- 1205 Kalashnikova, O. V, R. Kahn, I. N. Sokolik, and Wen-Hao Li. 2005. "Ability of Multiangle Remote Sensing Observations to Identify and Distinguish Mineral Dust Types: Optical Models and Retrievals of Optically Thick Plumes." *Journal of Geophysical Research: Atmospheres* 110(D18).
- 1210 Kaufman, Y. J., I. Koren, L. A. Remer, D. Tanré, P. Ginoux, and S. Fan. 2005. "Dust Transport and Deposition Observed from the Terra-Moderate Resolution Imaging Spectroradiometer (MODIS) Spacecraft over the Atlantic Ocean." *Journal of Geophysical Research D: Atmospheres* 110(10):1–16.
- 1215 Kim, Dongchul, Mian Chin, Hongbin Yu, Xiaohua Pan, Huisheng Bian, Qian Tan, Ralph A. Kahn, Kostas Tsigaridis, Susanne E. Bauer, Toshihiko Takemura, Luca Pozzoli, Nicolas Bellouin, and Michael Schulz. 2019. "Asian and Trans-Pacific Dust: A Multimodel and Multiremote Sensing Observation Analysis." *Journal of Geophysical Research: Atmospheres* 124(23):13534–59.
- 1220 Kim, Man Hae, Ali H. Omar, Jason L. Tackett, Mark A. Vaughan, David M. Winker, Charles R. Trepte, Yongxiang Hu, Zhaoyan Liu, Lamont R. Poole, Michael C. Pitts, Jayanta Kar, and Brian E. Magill. 2018. "The CALIPSO Version 4 Automated Aerosol Classification and Lidar Ratio Selection Algorithm." *Atmospheric Measurement Techniques* 11(11):6107–35.
- Klüser, L., D. Martynenko, and T. Holzer-Popp. 2011. "Thermal Infrared Remote Sensing of Mineral Dust over Land and Ocean: A Spectral SVD Based Retrieval Approach for IASI." *Atmospheric Measurement Techniques* 4(5):757–73.
- 1225 Kok, Jasper F., David A. Ridley, Qing Zhou, Ron L. Miller, Chun Zhao, Colette L. Heald, Daniel S. Ward, Samuel Albani, and Karsten Haustein. 2017. "Smaller Desert Dust Cooling Effect Estimated from Analysis of Dust Size and Abundance." *Nature Geoscience* 10(4):274–78.
- Koren, I., Y. J. Kaufman, L. A. Remer, and J. V. Martins. 2004. "Measurement of the Effect of Amazon Smoke on Inhibition of Cloud Formation." *Science* 303(5662):1342–45.
- 1230 Kurosaki, Yasunori and Masao Mikami. 2003. "Recent Frequent Dust Events and Their Relation to Surface Wind in East Asia." *Geophysical Research Letters* 30(14).
- Lau, K. M. and K. M. Kim. 2007. "Cooling of the Atlantic by Saharan Dust." *Geophysical Research Letters* 34(23).

- 1235 Lee, Eun Hee and Byung Ju Sohn. 2011. "Recent Increasing Trend in Dust Frequency over Mongolia and Inner Mongolia Regions and Its Association with Climate and Surface Condition Change." *Atmospheric Environment* 45(27):4611–16.
- Levy, R. C., S. Mattoo, L. A. Munchak, L. A. Remer, A. M. Sayer, F. Patadia, and N. C. Hsu. 2013. "The Collection 6 MODIS Aerosol Products over Land and Ocean." *Atmospheric Measurement Techniques* 6(11):2989–3034.
- 1240 Levy, Robert C., Shana Mattoo, Virginia Sawyer, Yingxi Shi, Peter R. Colarco, Alexei I. Lyapustin, Yujie Wang, and Lorraine A. Remer. 2018. "Exploring Systematic Offsets between Aerosol Products from the Two MODIS Sensors." *Atmospheric Measurement Techniques* 11(7):4073–92.
- 1245 Li, W. J. and L. Y. Shao. 2009. "Observation of Nitrate Coatings on Atmospheric Mineral Dust Particles." *Atmospheric Chemistry and Physics* 9(6):1863–71.
- Liu, Zhaoyan, Nobuo Sugimoto, and Toshiyuki Murayama. 2002. "Extinction-to-Backscatter Ratio of Asian Dust Observed with High-Spectral-Resolution Lidar and Raman Lidar." *Applied Optics* 41(15):2760–67.
- 1250 Marinou, Eleni, Vassilis Amiridis, Ioannis Biniotoglou, Athanasios Tsikerdekis, Stavros Solomos, Emannouil Proestakis, Dimitra Konsta, Nikolaos Papagiannopoulos, Alexandra Tsekeri, Georgia Vlastou, Prodromos Zanis, Dimitrios Balis, Ulla Wandinger, and Albert Ansmann. 2017. "Three-Dimensional Evolution of Saharan Dust Transport towards Europe Based on a 9-Year EARLINET-Optimized CALIPSO Dataset." *Atmospheric Chemistry and Physics* 17(9):5893–5919.
- 1255 Martins, José Vanderlei, Didier Tanré, Lorraine Remer, Yoram Kaufman, Shana Mattoo, and Robert Levy. 2002. "MODIS Cloud Screening for Remote Sensing of Aerosols over Oceans Using Spatial Variability." *Geophysical Research Letters* 29(12):MOD4-1-MOD4-4.
- Mbourou, G. N'Tchayi, J. J. Bertrand, and S. E. Nicholson. 1997. "The Diurnal and Seasonal Cycles of Wind-Borne Dust over Africa North of the Equator." *Journal of Applied Meteorology* 36(7):868–82.
- 1260 Mielonen, T., A. Arola, M. Komppula, J. Kukkonen, J. Koskinen, G. De Leeuw, and K. E. J. Lehtinen. 2009. "Comparison of CALIOP Level 2 Aerosol Subtypes to Aerosol Types Derived from AERONET Inversion Data." *Geophysical Research Letters* 36(18).
- 1265 Miller, R. L. and I. Tegen. 1998. "Climate Response to Soil Dust Aerosols." *Journal of Climate* 11(12):3247–67.
- Müller, Detlef, Albert Ansmann, Ina Mattis, Matthias Tesche, Ulla Wandinger, Dietrich Althausen, and G. Pisani. 2007. "Aerosol-type-dependent Lidar Ratios Observed with Raman Lidar." *Journal of Geophysical Research: Atmospheres* 112(D16).
- 1270 N'Tchayi Mbourou, G., J. J. Bertrand, and S. E. Nicholson. 1997. "The Diurnal and Seasonal Cycles of Wind-Borne Dust over Africa North of the Equator." *Journal of Applied Meteorology* 36(7):868–82.
- O'Neill, N. T., T. F. Eck, A. Smirnov, B. N. Holben, and S. Thulasiraman. 2003. "Spectral Discrimination of Coarse and Fine Mode Optical Depth." *Journal of Geophysical Research: Atmospheres* 108(17).
- 1275 Omar, A. H., D. M. Winker, J. L. Tackett, D. M. Giles, J. Kar, Z. Liu, M. A. Vaughan, K. A. Powell, and C. R. Trepte. 2013. "CALIOP and AERONET Aerosol Optical Depth Comparisons: One Size Fits None." *Journal of Geophysical Research Atmospheres* 118(10):4748–66.
- Omar, Ali H., David M. Winker, Chieko Kittaka, Mark A. Vaughan, Zhaoyan Liu, Yongxiang

- 1280 Hu, Charles R. Trepte, Raymond R. Rogers, Richard A. Ferrare, Kam Pui Lee, Ralph E. Kuehn, and Chris A. Hostetler. 2009. "The CALIPSO Automated Aerosol Classification and Lidar Ratio Selection Algorithm." *Journal of Atmospheric and Oceanic Technology* 26(10):1994–2014.
- 1285 Parrington, Josef R., WILLIAM H. ZOLLER, and NAMIK K. ARAS. 1983. "Asian Dust: Seasonal Transport to the Hawaiian Islands." *Science* 220(4593):195 LP – 197.
- Proestakis, Emmanouil, Vassilis Amiridis, Eleni Marinou, Aristeidis K. Georgoulas, Stavros Solomos, Stelios Kazadzis, Julien Chimot, Huizheng Che, Georgia Alexandri, Ioannis Biniotoglou, Vasiliki Daskalopoulou, Konstantinos A. Kourtidis, Gerrit De Leeuw, and Ronald J. Van Der A. 2018. "Nine-Year Spatial and Temporal Evolution of Desert Dust Aerosols over South and East Asia as Revealed by CALIOP." *Atmospheric Chemistry and Physics* 18(2):1337–62.
- 1290 Prospero, Joseph M., Paul Ginoux, Omar Torres, Sharon E. Nicholson, and Thomas E. Gill. 2002. "Environmental Characterization of Global Sources of Atmospheric Soil Dust Identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) Absorbing Aerosol Product." *Reviews of Geophysics* 40(1):2-1-2–31.
- 1295 Pu, Bing and Paul Ginoux. 2018. "How Reliable Are CMIP5 Models in Simulating Dust Optical Depth?" *Atmospheric Chemistry and Physics* 18(16):12491–510.
- Qian, Weihong, Lingshen Quan, and Shaoyin Shi. 2002. "Variations of the Dust Storm in China and Its Climatic Control." *Journal of Climate* 15(10):1216–29.
- 1300 Querol, Xavier, Aurelio Tobías, Noemí Pérez, A. Karanasiou, Fulvio Amato, Massimo Stafoggia, C. Pérez García-Pando, P. Ginoux, Francesco Forastiere, and Sophie Gumy. 2019. "Monitoring the Impact of Desert Dust Outbreaks for Air Quality for Health Studies." *Environment International* 130:104867.
- 1305 Rajapakshe, Chamara, Zhibo Zhang, John E. Yorks, Hongbin Yu, Qian Tan, Kerry Meyer, Steven Platnick, and David M. Winker. 2017. "Seasonally Transported Aerosol Layers over Southeast Atlantic Are Closer to Underlying Clouds than Previously Reported." *Geophysical Research Letters* 44(11):5818–25.
- Remer, L A, Y. J. Kaufman, D. Tanre, S. Mattoo, D. A. Chu, J. V Martins, R. R. Li, C. Ichoku, R. C. Levy, R. G. Kleidman, T. F. Eck, E. Vermote, and B. N. Holben. 2005. "The MODIS Aerosol Algorithm, Products, and Validation." *Journal of the Atmospheric Sciences* 62(4):947–73.
- 1310 Remer, Lorraine A, Y. J. Kaufman, D. Tanré, S. Mattoo, D. A. Chu, J. V Martins, R. R. Li, C. Ichoku, R. C. Levy, R. G. Kleidman, T. F. Eck, E. Vermote, and B. N. Holben. 2005. "The MODIS Aerosol Algorithm, Products, and Validation." *Journal of the Atmospheric Sciences* 62(4):947–73.
- 1315 Ridley, A. David, L. Colette Heald, F. Jasper Kok, and Chun Zhao. 2016. "An Observationally Constrained Estimate of Global Dust Aerosol Optical Depth." *Atmospheric Chemistry and Physics* 16(23):15097–117.
- 1320 Rosenfeld, D. and I. M. Lensky. 1998. "Satellite-Based Insights into Precipitation Formation Processes in Continental and Maritime Convective Clouds." *Bulletin of the American Meteorological Society* 79(11):2457–76.
- Sakai, Tetsu, Tomohiro Nagai, Yuji Zaizen, and Yuzo Mano. 2010. "Backscattering Linear Depolarization Ratio Measurements of Mineral, Sea-Salt, and Ammonium Sulfate Particles Simulated in a Laboratory Chamber." *Applied Optics* 49(23):4441–49.
- 1325 Sayer, A. M., N. C. Hsu, C. Bettenhausen, and M. J. Jeong. 2013. "Validation and Uncertainty

- Estimates for MODIS Collection 6 ‘Deep Blue’ Aerosol Data.” *Journal of Geophysical Research Atmospheres* 118(14):7864–72.
- Schuster, G. L., M. Vaughan, D. MacDonnell, W. Su, D. Winker, O. Dubovik, T. Lapyonok, and C. Trepte. 2012. “Comparison of CALIPSO Aerosol Optical Depth Retrievals to
1330 AERONET Measurements, and a Climatology for the Lidar Ratio of Dust.” *Atmospheric Chemistry and Physics* 12(16):7431–52.
- Shao, Y. P., K. H. Wyrwoll, A. Chappell, J. P. Huang, Z. H. Lin, G. H. McTainsh, M. Mikami, T. Y. Tanaka, X. L. Wang, and S. Yoon. 2011. “Dust Cycle: An Emerging Core Theme in Earth System Science.” *Aeolian Research* 2(4):181–204.
- 1335 Shimizu, Atsushi, Nobuo Sugimoto, Ichiro Matsui, Kimio Arao, Itsushi Uno, Toshiyuki Murayama, Naoki Kagawa, Kazuma Aoki, Akihiro Uchiyama, and A. Akihiro Yamazaki. 2004. “Continuous Observations of Asian Dust and Other Aerosols by Polarization Lidars in China and Japan during ACE-Asia.” *Journal of Geophysical Research D: Atmospheres* 109(19).
- 1340 Shimizu, Atsushi, Nobuo Sugimoto, Tomoaki Nishizawa, Yoshitaka Jin, and Dashdondog Batdorj. 2017. “Variations of Dust Extinction Coefficient Estimated by Lidar Observations over Japan, 2007–2016.” *Scientific Online Letters on the Atmosphere* 13:205–8.
- Song, Hongquan, Kesheng Zhang, Shilong Piao, and Shiqiang Wan. 2016. “Spatial and Temporal Variations of Spring Dust Emissions in Northern China over the Last 30 Years.”
1345 *Atmospheric Environment* 126:117–27.
- Song, Q., Z. Zhang, H. Yu, S. Kato, P. Yang, P. Colarco, L. A. Remer, and C. L. Ryder. 2018. “Net Radiative Effects of Dust in the Tropical North Atlantic Based on Integrated Satellite Observations and in Situ Measurements.” *Atmospheric Chemistry and Physics* 18(15).
- Sternberg, Troy, Henri Rueff, and Nick Middleton. 2015. “Contraction of the Gobi Desert, 2000–
1350 2012.” *Remote Sensing* 7(2):1346–58.
- Su, L. and O. B. Toon. 2011. “Saharan and Asian Dust: Similarities and Differences Determined by CALIPSO, AERONET, and a Coupled Climate-Aerosol Microphysical Model.” *Atmospheric Chemistry and Physics* 11(7):3263–80.
- Tang, Mingjin, Huanhuan Zhang, Wenjun Gu, Jie Gao, Xing Jian, Guoliang Shi, Bingqi Zhu, Luhua Xie, Liya Guo, Xiaoyan Gao, Zhe Wang, Guohua Zhang, and Xinming Wang. 2019.
1355 “Hygroscopic Properties of Saline Mineral Dust From Different Regions in China: Geographical Variations, Compositional Dependence, and Atmospheric Implications.” *Journal of Geophysical Research: Atmospheres* 124(20):10844–57.
- Tesche, M., A. Ansmann, D. Müller, D. Althausen, R. Engelmann, V. Freudenthaler, and S. Groß. 2009. “Vertically Resolved Separation of Dust and Smoke over Cape Verde Using Multiwavelength Raman and Polarization Lidars during Saharan Mineral Dust Experiment 2008.” *Journal of Geophysical Research Atmospheres* 114(13).
- 1360 Textor, C., M. Schulz, S. Guibert, S. Kinne, Y. Balkanski, S. Bauer, T. Berntsen, T. Berglen, O. Boucher, M. Chin, F. Dentener, T. Diehl, R. Easter, H. Feichter, D. Fillmore, S. Ghan, P. Ginoux, S. Gong, A. Grini, J. Hendricks, L. Horowitz, P. Huang, I. Isaksen, T. Iversen, S. Kloster, D. Koch, A. Kirkevåg, J. E. Kristjansson, M. Krol, A. Lauer, J. F. Lamarque, X. Liu, V. Montanaro, G. Myhre, J. Penner, G. Pitari, S. Reddy, Seland, P. Stier, T. Takemura, and X. Tie. 2006. “Analysis and Quantification of the Diversities of Aerosol Life Cycles within AeroCom.” *Atmospheric Chemistry and Physics* 6(7):1777–1813.
- 1370 Thorsen, Tyler J. and Qiang Fu. 2015. “CALIPSO-Inferred Aerosol Direct Radiative Effects: Bias Estimates Using Ground-Based Raman Lidars.” *Journal of Geophysical Research*

- 120(23):12,209-12,220.
- Twomey, Sean. 1977. "The Influence of Pollution on the Shortwave Albedo of Clouds." *Journal of the Atmospheric Sciences* 34(7):1149–52.
- 1375 Uno, Itsushi, Kenta Eguchi, Keiya Yumimoto, Toshihiko Takemura, Atsushi Shimizu, Mitsuo Uematsu, Zhaoyan Liu, Zifa Wang, Yukari Hara, and Nobuo Sugimoto. 2009. "Asian Dust Transported One Full Circuit around Theglobe." *Nature Geoscience* 2(8):557–60.
- Voss, Kara K. and Amato T. Evan. 2020. "A New Satellite-Based Global Climatology of Dust Aerosol Optical Depth." *Journal of Applied Meteorology and Climatology* 59(1):83–102.
- 1380 Voss, Kenneth J., Ellsworth J. Welton, Patricia K. Quinn, James Johnson, Anne M. Thompson, and Howard R. Gordon. 2001. "Lidar Measurements during Aerosols99." *Journal of Geophysical Research: Atmospheres* 106(D18):20821–31.
- Winker, D. M., J. L. Tackett, B. J. Getzewich, Z. Liu, M. A. Vaughan, and R. R. Rogers. 2013. "The Global 3-D Distribution of Tropospheric Aerosols as Characterized by CALIOP." *Atmospheric Chemistry and Physics* 13:3345.
- 1385 Winker, David M., Mark A. Vaughan, Ali Omar, Yongxiang Hu, Kathleen A. Powell, Zhaoyan Liu, William H. Hunt, and Stuart A. Young. 2009. "Overview of the CALIPSO Mission and CALIOP Data Processing Algorithms." *Journal of Atmospheric and Oceanic Technology* 26(11):2310–23.
- 1390 Wu, Tong, Zhanqing Li, Jun Chen, Yuying Wang, Hao Wu, Xiao'ai Jin, Chen Liang, Shangze Li, Wei Wang, and Maureen Cribb. 2020. "Hygroscopicity of Different Types of Aerosol Particles: Case Studies Using Multi-Instrument Data in Megacity Beijing, China." *Remote Sensing* 12(5).
- Xu, Hui, Fengjie Zheng, and Wenhao Zhang. 2016. "Variability in Dust Observed over China Using A-Train Caliop Instrument." *Advances in Meteorology* 2016(2015).
- 1395 Yang, W., A. Marshak, O. V Kalashnikova, and A. B. Kostinski. 2012. "CALIPSO Observations of Transatlantic Dust: Vertical Stratification and Effect of Clouds." *Atmospheric Chemistry and Physics* 12:11339.
- Yorks, J. E., S. P. Palm, D. L. Hlavka, M. J. McGill, E. Nowottnick, P. Selmer, and W. D. Hart. 2015. "The Cloud-Aerosol Transport System (CATS) Algorithm Theoretical Basis Document."
- 1400 Young, Stuart A., Mark A. Vaughan, Anne Garnier, Jason L. Tackett, James D. Lambeth, and Kathleen A. Powell. 2018. "Extinction and Optical Depth Retrievals for CALIPSO's Version 4 Data Release." *Atmospheric Measurement Techniques* 11:5701.
- 1405 Yu, H. B., M. Chin, H. S. Bian, T. L. Yuan, J. M. Prospero, A. H. Omar, L. A. Remer, D. M. Winker, Y. K. Yang, Y. Zhang, and Z. B. Zhang. 2015a. "Quantification of Trans-Atlantic Dust Transport from Seven-Year (2007-2013) Record of CALIPSO Lidar Measurements." *Remote Sensing of Environment* 159:232–49.
- Yu, H. B., M. Chin, T. L. Yuan, H. S. Bian, L. A. Remer, J. M. Prospero, A. Omar, D. Winker, Y. K. Yang, Y. Zhang, Z. B. Zhang, and C. Zhao. 2015b. "The Fertilizing Role of African Dust in the Amazon Rainforest: A First Multiyear Assessment Based on Data from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations." *Geophysical Research Letters* 42(6):1984–91.
- 1410 Yu, Hongbin, Mian Chin, Lorraine A. Remer, Richard G. Kleidman, Nicolas Bellouin, Huisheng Bian, and Thomas Diehl. 2009. "Variability of Marine Aerosol Fine-Mode Fraction and Estimates of Anthropogenic Aerosol Component over Cloud-Free Oceans from the Moderate Resolution Imaging Spectroradiometer (MODIS)." *Journal of Geophysical*
- 1415

Research Atmospheres 114(10):1–11.

- 1420 Yu, Hongbin, Mian Chin, David M. Winker, Ali H. Omar, Zhaoyan Liu, Chieko Kittaka, and Thomas Diehl. 2010. “Global View of Aerosol Vertical Distributions from CALIPSO Lidar Measurements and GOCART Simulations: Regional and Seasonal Variations.” *Journal of Geophysical Research Atmospheres* 115(4):1–19.
- 1425 Yu, Hongbin, Lorraine A. Remer, Mian Chin, Huisheng Bian, Qian Tan, Tianle Yuan, and Yan Zhang. 2012. “Aerosols from Overseas Rival Domestic Emissions over North America.” *Science* 337(6094):566–69.
- Yu, Hongbin, Lorraine A. Remer, Ralph A. Kahn, Mian Chin, and Yan Zhang. 2013. “Satellite Perspective of Aerosol Intercontinental Transport: From Qualitative Tracking to Quantitative Characterization.” *Atmospheric Research* 124:73–100.
- 1430 Yu, Hongbin, Qian Tan, Mian Chin, Lorraine A. Remer, Ralph A. Kahn, Huisheng Bian, Dongchul Kim, Zhibo Zhang, Tianle Yuan, Ali H. Omar, David M. Winker, Robert C. Levy, Olga Kalashnikova, Laurent Crepeau, Virginie Capelle, and Alain Chédin. 2019. “Estimates of African Dust Deposition Along the Trans-Atlantic Transit Using the Decadelong Record of Aerosol Measurements from CALIOP, MODIS, MISR, and IASI.” *Journal of Geophysical Research: Atmospheres* 124(14):7975–96.
- 1435 Yu, Hongbin, Yang Yang, Hailong Wang, Qian Tan, Mian Chin, Robert Levy, Lorraine Remer, Steven Smith, Tianle Yuan, and Yingxi Shi. 2020. “Interannual Variability and Trends of Combustion Aerosol and Dust in Major Continental Outflows Revealed by MODIS Retrievals and CAM5 Simulations During 2003–2017.” *Atmospheric Chemistry and Physics Discussions* 1–38.
- 1440 Yu, Yan, Olga V. Kalashnikova, Michael J. Garay, Huikyo Lee, Myungje Choi, Gregory S. Okin, John E. Yorks, James R. Campbell, and Jared Marquis. 2021. “A Global Analysis of Diurnal Variability in Dust and Dust Mixture Using CATS Observations.” *Atmospheric Chemistry and Physics* 21(3):1427–47.
- 1445 Yu, Yan, Olga V Kalashnikova, Michael J. Garay, and Michael Notaro. 2019. “Climatology of Asian Dust Activation and Transport Potential Based on MISR Satellite Observations and Trajectory Analysis.” *Atmospheric Chemistry & Physics* 19(1).
- Yue, Xu, Huijun Wang, Zifa Wang, and Ke Fan. 2009. “Simulation of Dust Aerosol Radiative Feedback Using the Global Transport Model of Dust: 1. Dust Cycle and Validation.” *Journal of Geophysical Research Atmospheres* 114(10).
- 1450