

Tropospheric ozone production and chemical regime analysis during the COVID-19 lockdown over Europe

Clara M. Nussbaumer¹, Andrea Pozzer¹, Ivan Tadic¹, Lenard Röder¹, Florian Obersteiner²,
Hartwig Harder¹, Jos Lelieveld^{1,3}, and Horst Fischer¹

¹Max Planck Institute for Chemistry, Department of Atmospheric Chemistry, 55128 Mainz, Germany

²Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany

³Climate and Atmosphere Research Center, The Cyprus Institute, Nicosia, Cyprus

Correspondence: Clara Nussbaumer (clara.nussbaumer@mpic.de)

Abstract.

The COVID-19 (Coronavirus disease 2019) European lockdowns have led to a significant reduction in the emissions of primary pollutants such as NO (nitric oxide) and NO₂ (nitrogen dioxide). As most photochemical processes are related to nitrogen oxide (NO_x ≡ NO + NO₂) chemistry, this event has presented an exceptional opportunity to investigate its effects on
5 air quality and secondary pollutants, such as tropospheric ozone (O₃). In this study, we present the effects of the COVID-19 lockdown on atmospheric trace gas concentrations, net ozone production rates (NOPR) and the dominant chemical regime throughout the troposphere based on three different research aircraft campaigns across Europe. These are the UTOPIHAN campaigns in 2003 and 2004, the HOOVER campaigns in 2006 and 2007 and the BLUESKY campaign in 2020, the latter performed during the COVID-19 lockdown. We present in situ observations and simulation results from the ECHAM5/MESSy
10 Atmospheric Chemistry model which allows for scenario calculations with business as usual emissions during the BLUESKY campaign, referred to as "no-lockdown scenario". We show that the COVID-19 lockdown reduced NO and NO₂ mixing ratios in the upper troposphere by around 55 % compared to the no-lockdown scenario due to reduced air traffic. O₃ production and loss terms reflected this reduction with a deceleration in O₃ cycling due to reduced mixing ratios of NO_x while NOPRs were largely unaffected. We also study the role of methyl peroxyradicals forming HCHO ($\alpha_{CH_3O_2}$) to show that the COVID-19
15 lockdown shifted the chemistry in the upper troposphere/tropopause region to a NO_x limited regime during BLUESKY. In comparison, we find a VOC limited regime to be dominant during UTOPIHAN.

1 Introduction

COVID-19 (Coronavirus disease 2019) describes the disease accompanying an infection with the SARS-CoV-2 (severe acute respiratory syndrome coronavirus-2) virus. The disease is highly infectious and can have severe health consequences, including
20 premature death, particularly for elderly and people with pre-existing conditions (WHO, 2021). On 11 March 2020, COVID-19 was declared a pandemic by the World Health Organization (WHO, 2020a,b). As a response, in many countries worldwide - including the European continent - governments initiated a shutdown of the daily life for minimizing the spread of the virus, which is referred to as COVID-19 lockdown. Among others, this included a reduction in vehicular and industrial activities as

well as sharp restrictions on air travel accompanied by a reduction in atmospheric pollutants such as nitrogen oxides ($\text{NO}_x \equiv$
25 $\text{NO} + \text{NO}_2$) (Venter et al., 2020; Kroll et al., 2020; Chossière et al., 2021; Onyeaka et al., 2021; Salma et al., 2020; Matthias
et al., 2021; Forster et al., 2020).

NO and NO_2 are important atmospheric trace gases as they are involved in almost all photochemical processes taking place
in the earth's atmosphere. NO_x directly impacts the production of tropospheric ozone (O_3) which is a hazard to human and plant
health (Nuvolone et al., 2018; Mills et al., 2018). Together with volatile organic compound (VOC) oxidation, NO forms NO_2
30 within the HO_x cycle, catalyzed by an OH radical. Under the influence of sunlight, NO_2 can subsequently form O_3 through
the reaction with molecular oxygen as shown in Reaction (R1) (Leighton, 1961; Crutzen, 1988; Lelieveld and Dentener, 2000;
Pusede and Cohen, 2012; Pusede et al., 2015; Nussbaumer and Cohen, 2020).



Various termination reactions such as the formation of HNO_3 from OH and NO_2 or other radical recombinations cause ozone
35 chemistry to be non-linear, which means that a reduction in ambient NO_x can either increase or decrease O_3 production
(Calvert and Stockwell, 1983; Pusede et al., 2015). For low ambient NO_x levels, a NO_x reduction usually causes a decrease in
 O_3 production which is referred to as a NO_x limited chemical regime. In contrast, a NO_x reduction increases O_3 production
when a VOC limited chemical regime is dominant - usually at high ambient NO_x levels (Sillman et al., 1990; National Re-
search Council, 1991; Pusede and Cohen, 2012). In the transition region between the two regimes, changes in NO_x do not (or
40 only slightly) impact O_3 production rates (Wang et al., 2018). Earlier studies on evaluating correlations of NO_x and O_3 in the
troposphere include Liu et al. (1987); Logan (1985) and Lin et al. (1988), reporting a non-linear dependence that varies with
ambient levels of hydrocarbons and NO_x .

Many different methods enable the determination of the dominant chemical regime, such as the use of the weekend ozone
effect which considers the response of O_3 to NO_x reductions on weekends or the ratio of HCHO to NO_2 with various ap-
45 proaches from in situ observations, remote sensing and model simulations (e.g. Jin et al. (2020); Pusede and Cohen (2012);
Nussbaumer and Cohen (2020); Duncan et al. (2010)). We have recently shown that the fraction α of methyl peroxyradicals
(CH_3O_2) forming formaldehyde (HCHO) in correlation with ambient NO concentrations is capable of indicating the dominant
chemical regime based on three different field campaigns across Europe in Finland (HUMPPA 2012), Germany (HOPE 2012)
and Cyprus (CYPHEX 2014) (Nussbaumer et al., 2021). CH_3O_2 formed from e.g. the oxidation of acetaldehyde (CH_3CHO) or
50 methane (CH_4) can either react with NO or OH radicals to form HCHO or undergo the competing reaction with HO_2 to form
methyl hydroperoxide (CH_3OOH). For more details, please see Figure 1 in Nussbaumer et al. (2021). $\alpha_{\text{CH}_3\text{O}_2}$ consequently
depends on the ambient concentrations of NO , OH and HO_2 and the respective rate constants for the reaction with CH_3O_2 ,
the latter of which were taken from the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (2021). Self-

reaction of CH_3O_2 as a contributor to CH_3O_2 loss forming HCHO is negligible. The calculation of $\alpha_{\text{CH}_3\text{O}_2}$ is presented in Equation (1).

$$\alpha_{\text{CH}_3\text{O}_2} = \frac{k_{\text{CH}_3\text{O}_2+\text{NO}} \times [\text{NO}] + k_{\text{CH}_3\text{O}_2+\text{OH}} \times [\text{OH}]}{k_{\text{CH}_3\text{O}_2+\text{NO}} \times [\text{NO}] + k_{\text{CH}_3\text{O}_2+\text{OH}} \times [\text{OH}] + k_{\text{CH}_3\text{O}_2+\text{HO}_2} \times [\text{HO}_2]} \quad (1)$$

Low values for $\alpha_{\text{CH}_3\text{O}_2}$ with a high response to NO are an indicator for a NO_x limited regime whereas high values for $\alpha_{\text{CH}_3\text{O}_2}$ with little response to changing NO represent a VOC limitation (Figure 11 in Nussbaumer et al. (2021)). Investigating the dominant chemical regime is an important method for analyzing photochemical processes and air quality.

Previous studies have explored changes in air quality, trace gas emissions and the dominant chemical regime during the COVID-19 lockdown in Europe. Menut et al. (2020) reported NO_2 reductions between 30 and 50 % for various Western European countries in the course of March 2020 with both decreasing and increasing O_3 concentrations in response, depending on the location, based on surface in situ observations and model simulations. Ordóñez et al. (2020) observed decreased NO_2 and increased O_3 concentrations in Central Europe in March and April 2020 based on in situ observations compared to 2015 - 2019. While they found NO_2 reductions to be mainly attributed to the COVID-19 lockdown, O_3 enhancements were predominantly affected by meteorological changes. Chossière et al. (2021) presented evidence on NO_2 reductions during the COVID-19 lockdown in Europe and O_3 changes dependent on the dominant chemical regime through investigation of HCHO/ NO_2 ratios based on in situ and satellite observations. Similar studies were performed by Matthias et al. (2021); Mertens et al. (2021); Balamurugan et al. (2021); Grange et al. (2021) and many more. Besides the changes within the dominant chemical regime through NO_x reductions, i.e. increasing ozone within a VOC limited regime and decreasing ozone within a NO_x limited regime, the COVID-19 lockdown could have potentially changed the dominant chemical regime from VOC to NO_x limited as pointed out by Kroll et al. (2020) and Gaubert et al. (2021). Cazorla et al. (2021) found a lockdown induced change from a VOC to a NO_x limited regime in Quito (Ecuador) based on the share of precursor loss to HNO_3 and H_2O_2 . The latter is dominant for NO_x limited chemistry (Kleinman et al., 2001). A change from a VOC to a NO_x limited regime was also reported by Zhu et al. (2021) in China based on HCHO to NO_2 ratios (NO_x limitation for ratios above 2 according to Duncan et al. (2010)). Most of the literature on pollutant reductions during the COVID-19 lockdown focuses on near-surface air quality and only few studies consider the free troposphere. Steinbrecht et al. (2021), Chang et al. (2022) and Bouarar et al. (2021) reported decreases in O_3 concentrations in the free troposphere based on in situ observations and modeling studies in the northern hemisphere. Bouarar et al. (2021) found that reduced air traffic - a unique incidence after strongly increasing aircraft activities over the past decades as shown by Lee et al. (2021) - can explain around a third of the observed O_3 decrease in 2020, the remaining contributions coming from ground-level reductions and meteorological differences. Reduced O_3 in the free troposphere was also reported by Clark et al. (2021) around Frankfurt airport. Cristofanelli et al. (2021) reported lower O_3 concentrations above the PBL in 2020 compared to the 1996 - 2019 average at Mount Cimone in Italy which is in line with findings by the World Meteorological Organization (2021), extended to include two mountain sites in Germany.

In this study, we present atmospheric trace gas concentrations, net ozone production rates and an analysis on the dominant chemical regime based on in situ observations during the research aircraft campaign BLUESKY which took place in May

and June 2020 over Europe, and model simulations. During this time period, aircraft activity was still strongly limited due to the COVID-19 lockdown. We compare the results to model simulations assuming business as usual emissions not impacted by government restrictions which we refer to as "no-lockdown scenario". Additionally, we present results on two previous aircraft campaigns which are UTOPIHAN (Upper Tropospheric Ozone: Processes Involving HO_x and NO_x) in 2003/2004 and HOOVER (HO_x over Europe) in 2006/2007. While many studies have been published on emissions reductions and the effect on secondary pollutants during the COVID-19 lockdown, only a few studies have investigated changes in the dominant chemical regime and to our knowledge we are the first to report a shift to NO_x limited chemistry in the upper troposphere. This can demonstrate the consequences of emission changes of VOCs (including methane) and NO_x on tropospheric ozone.

95 2 Observations and methods

2.1 Calculations of net ozone production rates (NOPR)

Besides the chemical regime, production and loss processes of O₃ are effective tools in exploring relevant photochemistry. As already demonstrated in Reaction (R1), O₃ is formed via NO₂ photolysis. Under the assumption of photostationary state, this term can be equated with the reactions of NO with O₃, HO₂ and RO₂ (Hosaynali Beygi et al., 2011). The resulting term for O₃ production P(O₃) is shown in Equation (2) (Tadic et al., 2020; Leighton, 1961). $j(NO_2)$ is the photolysis frequency of NO₂ and k describes the respective rate constant (for this work taken from the IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (2021)).

$$P(O_3) = [NO_2] \times j(NO_2) = [NO] \times (k_{O_3+NO} \times [O_3] + k_{NO+HO_2} \times [HO_2] + \sum_z k_{NO+R_zO_2} \times [R_zO_2]) \quad (2)$$

We assume R_zO₂ (the sum of all peroxy radicals) to be represented by CH₃O₂ which we find to be a reasonable approximation when comparing modeled CH₃O₂ to the overall modeled RO₂ as shown in Figure S1 of the Supplement, exemplarily for the BLUESKY campaign. Above 800 hPa, CH₃O₂ represents more than 90 % of RO₂. Below 800 hPa, it still accounts for more than 70 % on average. CH₃O₂ can be calculated via Equation (3) as derived by Bozem et al. (2017a). While the model can simulate CH₃O₂ mixing ratios, Equation (3) is required when working with experimental data as CH₃O₂ was not directly measured.

$$110 \quad [CH_3O_2] = \frac{k_{CH_4+OH} \times [CH_4]}{k_{CO+OH} \times [CO]} \times [HO_2] \quad (3)$$

O₃ loss occurs via the reaction with NO, OH and HO₂ and via photolysis and can be calculated as presented in Equation (4). The photolysis of O₃ first yields O¹D which reacts back to O₃ through collision with O₂ or N₂, and causes an O₃ loss through reaction with H₂O. The share of O₃ that is effectively lost through O₃ photolysis is described by α_{O^1D} in Equation (5) (Bozem

et al., 2017a). Additional loss due to reactions of O_3 with alkenes and the loss of NO_2 due to formation of HNO_3 or peroxy
 115 nitrates are negligibly small, particularly in the upper troposphere.

$$L(O_3) = [O_3] \times (k_{O_3+NO} \times [NO] + k_{O_3+HO_2} \times [HO_2] + k_{O_3+OH} \times [OH] + \alpha_{O^1D} \times j(O^1D)) \quad (4)$$

$$\alpha_{O^1D} = \frac{k_{O^1D+H_2O} \times [H_2O]}{k_{O^1D+N_2} \times [N_2] + k_{O^1D+O_2} \times [O_2] + k_{O^1D+H_2O} \times [H_2O]} \quad (5)$$

Net ozone production rates (NOPR) are then calculated from the difference in $P(O_3)$ and $L(O_3)$ whereas $P(O_3)$ can either be
 expressed via NO_2 or NO reaction terms. The term $k_{O_3+NO} \times [O_3] \times [NO]$ can be neglected for the latter as it is equally
 120 present in $P(O_3)$ and $L(O_3)$.

$$\begin{aligned} NOPR &= P(O_3) - L(O_3) \\ &= [NO_2] \times j(NO_2) - [O_3] \times (k_{O_3+NO} \times [NO] + k_{O_3+HO_2} \times [HO_2] + k_{O_3+OH} \times [OH] + \alpha_{O^1D} \times j(O^1D)) \\ &= [NO] \times (k_{NO+HO_2} \times [HO_2] + k_{NO+CH_3O_2} \times [CH_3O_2]) \\ &\quad - [O_3] \times (k_{O_3+HO_2} \times [HO_2] + k_{O_3+OH} \times [OH] + \alpha_{O^1D} \times j(O^1D)) \end{aligned} \quad (6)$$

2.2 Field experiments

We have investigated in situ trace gas observations from three different research aircraft campaigns which are the UTOPIHAN
 campaigns in 2003/2004, the HOOVER campaigns in 2006/2007 and the BLUESKY campaign in 2020. Figure 1 shows
 125 an overview of the flight tracks over Europe. We have filtered the data for the tropospheric region by help of the modeled
 tropopause pressure (see Section 2.3) and south of $60^\circ N$ as there were no data points for the BLUESKY campaign further
 north. Dashed lines show the complete flight tracks during each campaign and solid lines show the data which we have
 considered in this study. The experimental data were obtained with a time resolution of 1 minute and subsequently adjusted to
 fit the model resolution of 6 minutes. For this, each sixth experimental data point (which fit the model time scale) and the data
 130 points from ± 2 minutes were averaged. The remaining data points were discarded.

2.2.1 UTOPIHAN 2003/04

The UTOPIHAN (Upper Tropospheric Ozone: Processes Involving HO_x and NO_x) campaigns took place in June/July 2003
 and March 2004 starting from Oberpfaffenhofen airport in Germany ($48.08^\circ N$, $11.28^\circ E$) with the GFD (Gesellschaft für
 Flugzieldarstellung, Hohn, Germany) research aircraft Learjet 35A (Colomb et al., 2006; Klippel et al., 2011; Stickler et al.,
 135 2006). NO and O_3 were measured via chemiluminescent detection (CLD 790 SR, ECO Physics, Dürnten, Switzerland). NO
 data have a precision of 6.5 %, an accuracy of ≤ 25 % and a detection limit of < 0.01 ppbv. O_3 data have a precision of 1 % and
 an accuracy of 5 %. $j(NO_2)$ was determined via filter radiometers (Meterologie Consult GmbH, Königstein, Germany) with a




Figure 1. Overview of the flight tracks of the considered aircraft campaigns UTOPIHAN (2003 & 2004) in green, HOOVER (2006 & 2007) in blue and BLUESKY (2020) in red. Solid lines present the data considered in this study (filtered for troposphere and south of 60 °N) and dashed lines show the complete flight tracks.

precision of 1 % and an accuracy of 15 %. CO measurements were obtained from a tunable diode laser absorption spectrometer with a detection limit of 0.26 ppbv (30 s time resolution) and an accuracy of 3.6 % (6 s time resolution) (Kormann et al., 2005).

140 2.2.2 HOOVER 2006/07

The HOOVER (HO_x over Europe) campaigns took place in October 2006 and July 2007 using the GFD research aircraft Learjet 35A with the campaign base in Hohn, Germany (54.31 °N, 9.53 °E) (Klippel et al., 2011; Bozem et al., 2017b,a; Regelin et al., 2013). NO and O₃ measurements were performed via chemiluminescence (CLD 790 SR, ECO Physics, Dürnten, Switzerland) with a precision of 7 and 4 %, an accuracy of 12 and 2 % and a detection limit of 0.2 and 2 ppbv, respectively
 145 (30 s time resolution) (Hosaynali Beygi et al., 2011). CO and CH₄ were measured via quantum cascade laser absorption spectroscopy with an accuracy of 1.1 and 0.6 % and detection limits of 0.2 and 6 ppbv, respectively (2 s time resolution) (Schiller et al., 2008). OH and HO₂ measurements were performed via laser-induced fluorescence with the HORUS (HydrOxyl Radical measurement Unit based on fluorescence Spectroscopy) instrument with an accuracy of 18 % and detection limits of 0.016 and 0.33 pptv, respectively (1 min time resolution) (Regelin et al., 2013). Photolysis frequencies were measured using
 150 filter radiometers (Meterologie Consult GmbH, Königstein, Germany) with a precision of 1 % and an accuracy of 15 % (1 s time resolution). H₂O was measured via IR-absorption with a typical accuracy of 1 % (modified LI-6262, LI-COR Inc., Lincoln, USA) (Gurk et al., 2008; LI-COR, Inc., 1996).

2.2.3 BLUESKY 2020

The BLUESKY campaign took place in May and June 2020 over Europe. Eight flights were carried out using the HALO (High
155 Altitude Long Range) research aircraft starting from the campaign base in Oberpfaffenhofen, Germany. The goal of the campaign was to examine the effects of the COVID-19 lockdown on the troposphere and lower stratosphere over European cities, rural areas and the transatlantic flight corridor. More details can be found in Reifenberg et al. (2021) and Voigt et al. (2021). While most restrictions across Europe were in place in March and April 2020, May and June emissions, particularly from air travel but also from ground-based sources such as onroad traffic, were still affected by the COVID-19 lockdowns (Schlosser
160 et al., 2020; Brockmann Lab, 2022; Hasegawa, 2022; EUROCONTROL, 2022). NO was measured via chemiluminescence (CLD 790 SR, ECO Physics, Dürnten, Switzerland) with a total uncertainty of 15 % and a detection limit of 5 pptv (1 min time resolution) (Tadic et al., 2020).

O₃ measurements were performed with the FAIRO (Fast AIRborne Ozone) instrument, which allows fast detection via chemiluminescence that is calibrated in situ by UV photometry (2.5 % combined uncertainty, 5 Hz time resolution) (Zahn
165 et al., 2012). CO was measured via the quantum cascade laser spectrometer TRISTAR (Tracer In Situ Tdlas for Atmospheric Research) with an uncertainty of 3 % (1 min time resolution) (Schiller et al., 2008).

2.3 Modeling study

The modeled data were obtained from the ECHAM5 (5th generation European Centre Hamburg general circulation model, version 5.3.02)/MESSy2 (2nd generation Modular Earth Submodel System, version 2.54.0) Atmospheric Chemistry (EMAC)
170 model which is described in Jöckel et al. (2016) and Reifenberg et al. (2021).

We use data of NO, NO₂, O₃, OH, HO₂, CO, CH₄, CH₃O₂, H₂O, j(NO₂), j(O¹D) temperature and pressure, modeled along the flight tracks of the described research aircraft campaigns UTOPIHAN, HOOVER and BLUESKY. The data were filtered for the troposphere using the modeled tropopause pressure. Stratospheric data were discarded. In order to evaluate the impact of reduced emissions during the COVID-19 lockdown, the model was used to simulate a scenario with usual emissions for the
175 BLUESKY campaign which we refer to as "no-lockdown scenario". For details of the model set-up please see the paper by Reifenberg et al. (2021).

3 Results and Discussion

This analysis is structured as follows: As a full set of in situ observations necessary for a regime analysis and calculating net ozone production rates, which includes NO, O₃, OH, HO₂, CO, CH₄, H₂O, j(NO₂) and j(O¹D), is only available for the
180 HOOVER campaign, we first show that the model and experimental data are in close agreement for this campaign. We conclude from this finding that the model is generally capable of reproducing the experimental data and therefore use the model data in our following analysis. In the second step, we provide a comparison between the three campaigns as well as the no-lockdown

scenario regarding the individual trace gases and net ozone production rates. We finally present our results of the analysis of the dominant chemical regime, based on $\alpha_{CH_3O_2}$.

185 3.1 Comparison of Model and Experiment

Figure 2 shows a comparison of in situ observations (orange) and model simulations (blue) for the HOOVER campaign as vertical profiles. The shaded areas present the 1σ standard deviations and the numbers of data points available for each altitude bin are shown in Table S1 and S2 of the Supplement.

Figure 2a presents the vertical profile of NO which shows the typical tropospheric C-shape distribution with the highest 190 values at the surface (e.g. vehicle and industrial emissions) and the upper troposphere (e.g. aircraft and lightning emissions). Ground-level mixing ratios (0 - 1000 m) were around 0.4 ppbv and decreased with altitude to values of 37 ± 27 (1σ) pptv and 47 ± 32 pptv for the model and the experiment, respectively, between 3 and 9 km altitude. The only relevant deviation of model and experiment was between 10 and 11 km altitude with mixing ratios of 0.20 ± 0.03 ppbv and 0.39 ± 0.32 ppbv, respectively.

Figure 2b shows the measured and modeled O_3 mixing ratios which were lowest at ground levels with 43.7 ± 14.5 ppbv 195 and 36.4 ± 12.8 ppbv for model and experiment and increased with altitude up to 128.1 ± 22.7 ppbv and 97.5 ± 15.6 ppbv, respectively. Model values were approximately 20 % higher compared to the measured data, but showed the same vertical shape. The observed positive O_3 bias of the modeled data is an issue almost all global models suffer from in the northern hemisphere and which has not been entirely understood yet (Revell et al., 2018; Young et al., 2018; Jöckel et al., 2016; Parrish et al., 2014).

CO vertical profiles are shown in Figure 2c which were highest at the surface with 146.4 ± 63.2 ppbv and 128.0 ± 42.3 ppbv 200 for model and experiment, respectively, and decreased with altitude to around 70 ppbv in the upper troposphere. HO_x ($\equiv OH + HO_2$) are presented in Figure 2d and e. HO_2 mixing ratios showed a maximum value of around 20 pptv between 2 and 3 km altitude and decreased aloft to values of around 2 pptv in the upper troposphere. Model and experiment showed close agreement. OH mixing ratios were mostly below 1 pptv. Similar to NO, the main deviation between model and experiment was 205 between 10 and 11 km altitude where measured values were higher by around 0.5 pptv. Nevertheless, the error bars representing the 1σ standard deviation of the averages overlapped at all altitudes.

Figure 2f shows the vertical profiles of CH_4 which did not show any particular gradient with altitude. Mixing ratios were 1809 ± 19 ppbv for the model simulation and 1815 ± 40 ppbv for the experiment throughout the campaign. CH_4 is needed for calculating CH_3O_2 via Equation (3), which we show in Figure 2g in orange compared to the model simulation of CH_3O_2 . 210 Figure 2h and i present the photolysis frequencies $j(NO_2)$ and $j(O^1D)$ which show close agreement for model and experiment. We show the vertical profiles for H_2O , temperature and pressure in Figure S2 of the Supplement. Again, model simulation can represent the experimental data well.

For the UTOPIHAN and the BLUESKY campaigns only a limited number of observations is available. Similar to the HOOVER campaigns, NO, O_3 and CO can be well approximated by the model simulations which we present in Figure S3 215 and S4 of the Supplement. Tropospheric ozone is slightly overestimated, which we attribute to the simplified representation of multiphase chemistry (clouds) in the present model version, which underpredicts chemical ozone loss (Rosanka et al., 2021).




Figure 2. Vertical profiles of in situ observations and model data of the atmospheric trace gases (a) NO, (b) O₃, (c) CO, (d) HO₂, (e) OH, (f) CH₄ and (g) CH₃O₂ and the photolysis rates (h) $j(\text{NO}_2)$ and (i) $j(\text{O}^1\text{D})$ during the HOOVER campaign for estimating the model performance. Blue colors show modeled data by EMAC along the HOOVER campaign flight track (Model) and orange colors show experimental data (Experiment). The orange trace in panel (g) shows the calculation of CH₃O₂ from experimental CH₄, CO and HO₂ via Equation (3). The shaded areas represent the 1σ standard deviation from averaging the data points at each altitude bin. The numbers of data points averaged per altitude bin are displayed in Table S1 and S2 of the Supplement.

Based on these results, we conclude that the model is generally capable of well representing the in situ observations and use the model data for all following analyses.

3.2 Campaign Comparison

220 3.2.1 Trace gases




Figure 3. Vertical profiles of the atmospheric trace gases (a) NO, (b) O_3 , (c) CO, (d) NO_2 , (e) HO_2 and (f) OH for the campaigns UTOPIHAN (green), HOOVER (blue), BLUESKY (red) and the no-lockdown (NL) scenario (yellow). All data shown here are from EMAC model simulations along the flight track of each research campaign. Two separate simulations were run on the flight path of BLUESKY, one with lockdown and one with business-as-usual emissions. The shaded areas represent the 1σ standard deviation from averaging the data points at each altitude bin. The numbers of data points averaged per altitude bin are displayed in Table S2 of the Supplement.

Figure 3 presents the vertical profiles of some selected trace gases during the research aircraft campaigns UTOPIHAN (green), HOOVER (blue) and BLUESKY (red) which were obtained from model simulations. Yellow lines show the no-lockdown (NL) scenario for the BLUESKY campaign in 2020.

The vertical profiles of NO are presented in Figure 3a. For all campaigns, we observe the typical C-shape as described
225 for the HOOVER campaigns in Section 3.1. Surface (0- 1000 m) mixing ratios were similar for UTOPIHAN and HOOVER
with 0.45 ± 0.37 (1σ) ppbv and 0.43 ± 0.74 ppbv, respectively. In comparison, the ground-level concentration of NO during
BLUESKY was 0.12 ± 0.11 ppbv. The differences in NO mixing ratios between the campaigns are the outcome of the general
emission reduction due to legislative limitation of nitrogen oxides and other hazardous pollutants over the past decades, as the
campaigns took place 15- 20 years apart. We show the decrease in NO_x emissions in the model over the past two decades
230 in Figure S5 of the Supplement. Assuming the no-lockdown scenario during BLUESKY, NO ground level mixing ratios
were 0.15 ± 0.14 ppbv and therefore 25 % higher compared to actual mixing ratios (20 % emission reduction). This difference
between lockdown and no-lockdown mixing ratios is slightly lower compared to the findings by other studies, for example by
Donzelli et al. (2021) who found a NO decrease by 35 - 65 % in Valencia, Spain or by Higham et al. (2021) who reported a
NO decrease by 55 % in the UK compared to 2019. A possible reason can be that the BLUESKY aircraft campaign took place
235 in May and June 2020 whereas the main lockdown period across Europe occurred rather in March and April. Emission were
still reduced in the following months, but likely to a smaller extent. NO was low and similar for all campaigns between 3 and
8 km altitude, a region without any particular NO sources, with most values below 50 pptv. Above 10 km, NO mixing ratios
were 0.29 ± 0.19 ppbv for UTOPIHAN, 0.21 ± 0.03 ppbv for HOOVER and 0.08 ± 0.04 ppbv for BLUESKY. In comparison,
NO mixing ratios for the no-lockdown scenario were 0.17 ± 0.08 ppbv above 10 km altitude. This corresponds to an emission
240 reduction of 55 % and results in both absolute and relative NO reductions in the upper troposphere being much higher compared
to ground-level reductions. The observed NO reduction in the upper troposphere can be attributed to reduced air traffic which
we show in Figure 4. In addition to the vertical profiles of NO for BLUESKY (red) and BLUESKY-NL (yellow), we present the
modeled BLUESKY-NL scenario without aircraft emissions in blue. In the lower troposphere, where aircraft emissions do not
play a significant role, this profile is identical to the BLUESKY-NL scenario. In the upper troposphere, it is very similar to the
245 BLUESKY scenario (including the air travel restrictions), showing that reduced air traffic causes the observed NO decrease.

Figure 3b presents the O_3 vertical profiles. For all campaigns, O_3 mixing ratios were lowest at ground levels with values of
around 50 ppbv and increased with increasing altitude up to around 140 ppbv above 10 km altitude. No significant differences
between the campaigns can be observed. While ozone concentrations are dependent on various effects such as precursor levels
(including NO_x and VOCs) or meteorology, seasonal variations with a maximum around summer time and a minimum during
250 winter months are also of importance (Logan, 1985). The here shown campaigns include different seasons: the HOOVER
campaigns took place in October and July and the UTOPIHAN campaigns include data from July and March. Figure S6 of
the Supplement shows the vertical profiles of ozone separated into different seasons, for both modeled and measured data.
Comparing late spring / early summer data of the three field campaigns reveals that O_3 levels during BLUESKY were lower
compared to HOOVER and UTOPIHAN which is in line with findings from Clark et al. (2021), Chang et al. (2022), Bouarar
255 et al. (2021) and Miyazaki et al. (2021).

CO vertical profiles can be seen in 3c. Ground level mixing ratios were 181.4 ± 39.4 ppbv for UTOPIHAN, 146.4 ± 63.2 ppbv
for HOOVER and slightly lower with 103.2 ± 9.2 ppbv for BLUESKY. Mixing ratios slightly decreased with altitude. Above
3 km altitude, CO for HOOVER was lower compared to the other campaigns (mostly between 70 and 80 ppbv). Mixing ratios




Figure 4. NO vertical profiles for BLUESKY (red), for the BLUESKY no-lockdown scenario (yellow) and for the BLUESKY no-lockdown scenario without aircraft emissions (blue) (model data). Upper tropospheric NO reductions observed for BLUESKY can be attributed to reduced air traffic during the COVID-19 lockdowns.

for UTOPIHAN were slightly higher up to 11 km altitude (between 90 and 110 ppbv) compared to BLUESKY (between 80 and 100 ppbv), but generally, significant differences are not evident.

Figure 3d shows the vertical profiles of NO_2 mixing ratios. Similar to NO, ground level NO_2 mixing ratios were highest for UTOPIHAN and HOOVER with 1.57 ± 0.77 ppbv and 2.58 ± 2.72 ppbv, respectively. In contrast, mixing ratios for BLUESKY were 0.39 ± 0.30 ppbv and 0.49 ± 0.38 ppbv considering the no-lockdown scenario which yields a 20 % NO_2 lockdown reduction, as observed for NO. We show the NO_2 range 0 - 1 ppbv for enabling the campaign distinction at low mixing ratios and present the full range in Figure S7 of the Supplement. As expected for NO_2 , mixing ratios decreased with increasing altitude. No differences between the campaigns can be observed for mid-range altitudes. In the upper troposphere, NO_2 mixing ratios for the individual campaigns showed the same behavior as for NO. Above 10 km altitude, NO_2 was on average 100.6 ± 93.2 pptv for UTOPIHAN, 70.5 ± 13.5 pptv for HOOVER and 43.1 ± 23.1 pptv for the no-lockdown scenario for BLUESKY. In comparison, BLUESKY NO_2 mixing ratios were 19.9 ± 9.8 pptv which corresponds to a 55 % reduction. In contrast to NO, NO_2 reductions were relatively higher in the upper troposphere, but absolutely higher at the surface.

Figure 3e and f show the vertical profiles of HO_x . HO_2 mixing ratios were highest at mid-range altitudes (2 - 6 km) with values up to 20 pptv and decreased aloft. OH mixing ratios were lowest at the surface (0.1 - 0.2 pptv) and increased with altitude. Above 10 km altitude, OH mixing ratios were 0.62 ± 0.38 pptv for UTOPIHAN, 0.40 ± 0.24 pptv for HOOVER, 0.30 ± 0.06 pptv for BLUESKY and 0.39 ± 0.08 pptv for the no-lockdown scenario.




Figure 5. Vertical profiles of (a) net ozone production rates, (b) O_3 production via NO_2 photolysis, (c) O_3 loss via reaction with NO , (d) O_3 loss via photolysis, (e) O_3 loss via reaction with HO_2 and (f) O_3 loss via reaction with OH for the campaigns UTOPIHAN (green), HOOVER (blue), BLUESKY (red) and the no-lockdown (NL) scenario (yellow). The numbers of data points averaged per altitude bin are displayed in Table S2 of the Supplement.

Figure 5 shows the vertical profiles of O_3 production and loss terms. All calculations were performed using model data (justified by the findings from Section 3.1) as a full set of in situ observations is only available for HOOVER, but not for UTOPIHAN and BLUESKY. Figure 5a presents net ozone production rates, which were highest at the surface with values between 1 and 2 ppbv h^{-1} , but had large atmospheric variabilities, represented by the 1σ variability shades from the vertical 280 bin averaging. NOPRs then decreased with increasing altitude. For the HOOVER campaigns, O_3 loss dominated between 3 and 6 km altitude with NOPRs of $-58.9 \pm 73.4 \text{ pptv h}^{-1}$. Negative NOPRs were also found for BLUESKY between 4 and 7 km with $-18.7 \pm 12.9 \text{ pptv h}^{-1}$ and for UTOPIHAN as well as the no-lockdown BLUESKY scenario between 5 and 6 km. NOPRs were mostly positive and constant aloft. Above 10 km altitude, NOPRs were $91.7 \pm 260.9 \text{ pptv h}^{-1}$ for UTOPIHAN (51 data points), $71.2 \pm 151.5 \text{ pptv h}^{-1}$ for HOOVER (25 data points), $60.7 \pm 39.7 \text{ pptv h}^{-1}$ for BLUESKY (130 data points)




Figure 6. Modeled and experimental vertical profiles of $P(O_3)$ for HOOVER. Modeled $P(O_3)$ was calculated via NO_2 photolysis and experimental $P(O_3)$ was calculated via the extended Leighton ratio as shown in Equation (2).

285 and 61.4 ± 99.8 pptv h^{-1} for the no-lockdown scenario. The error ranges are large and overlapping and therefore, significant differences between the campaigns cannot be observed.

Figure 5b shows O_3 production. We calculated the $P(O_3)$ via the photolysis of NO_2 . In contrast, NO_2 is not available experimentally for the HOOVER campaign in which case the approximation via the extended Leighton ratio as shown in Equation (2) is necessary. Modeled $P(O_3)$ via NO_2 photolysis and measured $P(O_3)$ via reaction of NO with O_3 , OH and
 290 HO_2 are in good agreement which we show in Figure 6. The only relevant deviation is observed at ground levels, where the experimental value is significantly higher compared to the modeled value. However, only three data points were available for the calculation with a 1σ standard deviation of the averaging of $>100\%$. Similar to NOPRs in Figure 5a, ground-level $P(O_3)$ shows large variability with absolute values of around 10 ppbv h^{-1} for BLUESKY and values of around 20 ppbv h^{-1} for UTOPIHAN and HOOVER. The production term then decreased with altitude for each campaign. Significant differences
 295 between the campaigns can only be observed at high altitudes. Above 10 km, $P(O_3)$ was 4.55 ± 3.82 ppbv h^{-1} for UTOPIHAN (51 data points) and 2.68 ± 0.90 ppbv h^{-1} for HOOVER (25 data points). For BLUESKY with the no-lockdown scenario, $P(O_3)$ was 2.17 ± 0.95 ppbv h^{-1} (130 data points) and in comparison, lockdown values were on average 0.97 ± 0.41 ppbv h^{-1} which corresponds to a 55% reduction in ozone production. We observed the same relative reduction as for NO and NO_2 mixing ratios.

300 Figure 5c presents the vertical profiles of O_3 loss via the reaction with NO which show a similar course compared to the $P(O_3)$ profiles. Above 10 km, O_3 loss via reaction with NO was largest for UTOPIAN with 4.37 ± 3.82 ppbv h^{-1} , followed by HOOVER with 2.56 ± 0.87 ppbv h^{-1} . For BLUESKY, a loss of 0.86 ± 0.42 ppbv h^{-1} was observed during the lockdown and




Figure 7. $\alpha_{CH_3O_2}$ for the campaigns UTOPIHAN (green), HOOVER (blue), BLUESKY (red) and the no-lockdown (NL) scenario (yellow) (a) as vertical profile, (b) in correlation with NO below 2 km and (c) in correlation with NO above 10 km.

a loss of 2.05 ± 1.02 ppbv h^{-1} for the no-lockdown scenario. Figures 5d-f present additional considered loss pathways for O_3 via photolysis and via the reactions with HO_2 and OH. It can be seen that these O_3 losses are negligibly small in comparison to the loss via NO and no significant differences between the campaigns were present.

Consequently, net production of ozone was dominated by NO_x chemistry for all campaigns and variations in production and loss terms corresponded to the mixing ratios of NO and NO_2 as presented in Figure 3. In the campaign comparison, higher NO_x concentrations (as for example for UTOPIHAN) lead to higher production and loss terms of O_3 and vice versa. For the BLUESKY campaign, this analysis shows that the lockdown did not affect net ozone production rates, but instead impacted the cycling of O_3 such that both production and loss rates were decreased through the reduced availability of NO and NO_2 in the upper troposphere.

3.3 Chemical regime

As described above, the share of methyl peroxyradicals forming formaldehyde $\alpha_{CH_3O_2}$ can be a measure for the dominant chemical regime when correlated with NO mixing ratios. We have previously validated this method in a comparison to the established method of analyzing the HCHO/ NO_2 ratio (Nussbaumer et al., 2021). HCHO can be formed by almost any hydrocarbon and is therefore a proxy for VOCs which are often not measured in their entirety. Likewise, $\alpha_{CH_3O_2}$ - representing the HCHO yield from methyl peroxy radicals - is capable of revealing the dominant chemical regime without the knowledge of ambient VOC levels. Figure 7a shows the vertical profiles of $\alpha_{CH_3O_2}$ for all available data point for all campaigns based on the model simulation. $\alpha_{CH_3O_2}$ values were close to 1 at the surface and decreased with altitude up to around 5 km where values of around 0.6 were observed, with no significant differences between the campaigns. $\alpha_{CH_3O_2}$ increased again aloft whereas it was lowest for the BLUESKY campaign. Above 10 km, $\alpha_{CH_3O_2}$ was 0.97 ± 0.03 for UTOPIHAN, 0.98 ± 0.01 for HOOVER and 0.96 ± 0.04 for the no-lockdown scenario for BLUESKY. In comparison, $\alpha_{CH_3O_2}$ was lower for BLUESKY with 0.90 ± 0.06 .

Figures 7b and c present $\alpha_{CH_3O_2}$ in correlation with NO mixing ratios below 2 km altitude and above 10 km altitude, respectively, based on model results. Below 2 km altitude, $\alpha_{CH_3O_2}$ ranged between 0.5 and 1.0 over the NO range of 0 - 1 ppbv. No significant trends or differences can be observed. We show $\alpha_{CH_3O_2}$ between 2 and 10 km altitude in Figure S8 of the Supplement which does not present any differences between the campaigns either. In contrast above 10 km altitude, tropospheric $\alpha_{CH_3O_2}$ showed a different behavior for each campaign. For an easier distinction, we show each campaign in an individual panel in Figure S9 of the Supplement. For UTOPIHAN, $\alpha_{CH_3O_2}$ was high and almost non-responsive to changing NO mixing ratios with a slope of $\Delta\alpha/\Delta NO = 0.09 \pm 0.02 \text{ ppbv}^{-1}$. In contrast, $\alpha_{CH_3O_2}$ for BLUESKY was between 0.75 and 1. Small changes in NO mixing ratios caused large changes in $\alpha_{CH_3O_2}$ with a slope of $1.12 \pm 0.08 \text{ ppbv}^{-1}$. For the no-lockdown scenario the response of $\alpha_{CH_3O_2}$ to NO was intermediate between UTOPIHAN and BLUESKY with a slope of $0.37 \pm 0.03 \text{ ppbv}^{-1}$. These observations suggest that a VOC limited chemical regime was present during the UTOPIHAN campaign in the upper troposphere and a transition regime during the BLUESKY no-lockdown scenario, likely due to emission control over time. For BLUESKY, we observe a distinct NO_x limitation in the upper troposphere which is related to the lockdown conditions. Aircraft NO_x emissions are much larger than aircraft VOC emissions (Schumann, 2002). We can therefore expect reduced air traffic to effect lower NO_x/VOC ratios shifting chemistry towards a NO_x limited regime. Lamprecht et al. (2021) reported ground-level reductions of several aromatic VOCs during the COVID-19 lockdown to be comparable to NO_x reductions in Europe, implicating a steady NO_x/VOC level and therefore no changes in the dominating chemical regime, which is in line with our findings for the lower troposphere. Only few data points were available for HOOVER which were observed at similar NO levels and the response of $\alpha_{CH_3O_2}$ to NO can therefore not be investigated. While the NOPR did not change under lockdown conditions due to compensating effects in the NO_x chemistry, we can expect impacts on tropospheric ozone from changes in VOCs (including CH_4) relevant for future emission scenarios. The effects of NO_x aircraft emissions on O_3 and CH_4 have been previously discussed for pre-lockdown conditions in Khodayari et al. (2014) and Khodayari et al. (2015) who present increased methane loss rates and a shorter lifetime as a response to increased OH concentrations from aviation as well as higher ozone production rates. Having investigated NOPR in this study, lockdown effects on CH_4 loss in the UT induced by reduced air traffic could be subject to future studies.

4 Conclusions

In this study, we have presented in situ observations of atmospheric trace gases and model simulations from the EMAC model for three different aircraft campaigns across Europe, the UTOPIHAN campaigns in 2003/04, the HOOVER campaigns in 2006/07 and the BLUESKY campaign in 2020, including a modeled "no-lockdown scenario" with business as usual emissions for the latter. We found that model results can reproduce in situ observations well and thus could be used for further analysis which benefits from a more complete set of parameters and a higher data coverage. While observations for O_3 , CO and HO_x were very similar for all campaigns, NO_x showed significant differences, particularly in the upper troposphere, where mixing ratios were highest for UTOPIHAN and HOOVER, followed by the no-lockdown scenario for BLUESKY. Observed NO and NO_2 emissions during the BLUESKY campaign were approximately 55 % lower compared to the modeled no-lockdown

scenario which are attributed to reduced aircraft activity at these altitudes due to the COVID-19 travel restrictions. We found a similar trend in production and loss terms of O_3 which were dominated by NO_x chemistry. The COVID-19 lockdown caused a significant deceleration in O_3 cycling whereas net ozone production rates were not affected by the emission reductions. Finally, we showed that chemistry in the upper troposphere was VOC limited during the UTOPIHAN campaign, NO_x limited during
360 the BLUESKY campaign and in a transition regime for the BLUESKY no-lockdown scenario. While ground-level chemistry regimes were not found to be affected, the COVID-19 lockdown caused the predominant chemistry to shift from a transition regime to a clear NO_x limited regime at high altitudes.

We found that the three aircraft campaigns, performed over a period of 17 years, represent the range from VOC to NO_x limited tropospheric ozone chemistry, which can help analyze the impacts of anthropogenic emission scenarios. We encourage
365 future studies to investigate the dominating chemical regime in the upper troposphere, a topic which has not received much attention in literature so far, in order to get a deeper understanding of photochemical processes and the dominant ozone chemistry in a range of the atmosphere which receives its main NO_x emissions from air traffic and lightning. The COVID-19 lockdown has been a unique opportunity to examine the effect of sharp reductions in primary pollutants on our atmosphere and could be a guidepost for future air policy in an effort to decrease anthropogenic emissions and to decelerate global warming.

370 *Data availability.* Data measured during the flight campaigns BLUESKY, UTOPIHAN and HOOVER are available upon request at <https://keeper.mpdl.mpg.de/> to all scientists agreeing to the respective data protocols. The model results used in this study are available upon request to the author.

Author contributions. CMN and HF had the idea and designed the study. CMN analyzed the data and wrote the manuscript. AP provided the modeling data. IT provided NO data for BLUESKY. CO data for BLUESKY were received from LR. O_3 data for BLUESKY were obtained
375 from FO. HH provided HO_x data for HOOVER. JL and HF were significantly involved in planning and operating the research campaign.

Competing interests. Andrea Pozzer is a member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements. We acknowledge Simon Reifenberg for preparing the input data for EMAC. This work was supported by the Max Planck Graduate Center (MPGC) with the Johannes Gutenberg-Universität Mainz.

References

- 380 Balamurugan, V., Chen, J., Qu, Z., Bi, X., Gensheimer, J., Shekhar, A., Bhattacharjee, S., and Keutsch, F. N.: Tropospheric NO₂ and O₃ response to COVID-19 lockdown restrictions at the national and urban scales in Germany, *Journal of Geophysical Research: Atmospheres*, 126, 1–15, <https://doi.org/10.1029/2021JD035440>, 2021.
- Bouarar, I., Gaubert, B., Brasseur, G. P., Steinbrecht, W., Doumbia, T., Tilmes, S., Liu, Y., Stavrakou, T., Deroubaix, A., Darras, S., et al.: Ozone Anomalies in the Free Troposphere During the COVID-19 Pandemic, *Geophysical Research Letters*, 48, e2021GL094204, <https://doi.org/10.1029/2021GL094204>, 2021.
- 385 Bozem, H., Butler, T. M., Lawrence, M. G., Harder, H., Martinez, M., Kubistin, D., Lelieveld, J., and Fischer, H.: Chemical processes related to net ozone tendencies in the free troposphere, *Atmospheric Chemistry and Physics*, 17, 10 565–10 582, <https://doi.org/10.5194/acp-17-10565-2017>, 2017a.
- Bozem, H., Pozzer, A., Harder, H., Martinez, M., Williams, J., Lelieveld, J., and Fischer, H.: The influence of deep convection on HCHO and H₂O₂ in the upper troposphere over Europe, *Atmospheric Chemistry and Physics*, 17, 11 835–11 848, <https://doi.org/10.5194/acp-17-11835-2017>, 2017b.
- 390 Brockmann Lab: Covid-19 Mobility Project, <http://covid-19-mobility.org/>, "accessed on 2022-03-09", 2022.
- Calvert, J. G. and Stockwell, W. R.: Deviations from the O₃–NO–NO₂ photostationary state in tropospheric chemistry, *Canadian Journal of Chemistry*, 61, 983–992, <https://doi.org/10.1139/v83-174>, 1983.
- 395 Cazorla, M., Herrera, E., Palomeque, E., and Saud, N.: What the COVID-19 lockdown revealed about photochemistry and ozone production in Quito, Ecuador, *Atmospheric pollution research*, 12, 124–133, <https://doi.org/10.1016/j.apr.2020.08.028>, 2021.
- Chang, K.-I., Cooper, O. R., Gaudel, A., Allaart, M., Ancellet, G., Clark, H., Godin-Beekmann, S., Leblanc, T., Van Malderen, R., Nédélec, P., Petropavlovskikh, I., Steinbrecht, W., Stübi, R., Tarasick, D. W., and Torres, C.: Impact of the COVID-19 Economic Downturn on Tropospheric Ozone Trends: An Uncertainty Weighted Data Synthesis for Quantifying Regional Anomalies Above Western North America and Europe, *AGU Advances*, 3, e2021AV000 542, <https://doi.org/10.1029/2021AV000542>, 2022.
- 400 Chossière, G. P., Xu, H., Dixit, Y., Isaacs, S., Eastham, S. D., Allroggen, F., Speth, R. L., and Barrett, S. R.: Air pollution impacts of COVID-19–related containment measures, *Science Advances*, 7, 1–10, <https://doi.org/10.1126/sciadv.abe1178>, 2021.
- Clark, H., Bennouna, Y., Tsvilidou, M., Wolff, P., Sauvage, B., Barret, B., Le Flochmoën, E., Blot, R., Boulanger, D., Cousin, J.-M., Nédélec, P., Petzold, A., and Thouret, V.: The Effects of the COVID-19 Lockdowns on the Composition of the Troposphere as Seen by IAGOS, *Atmospheric Chemistry and Physics Discussions*, pp. 1–33, <https://doi.org/10.5194/acp-21-16237-2021>, 2021.
- 405 Colomb, A., Williams, J., Crowley, J., Gros, V., Hofmann, R., Salisbury, G., Klüpfel, T., Kormann, R., Stickler, A., Forster, C., and Lelieveld, J.: Airborne measurements of trace organic species in the upper troposphere over Europe: the impact of deep convection, *Environmental Chemistry*, 3, 244–259, <https://doi.org/10.1071/EN06020>, 2006.
- Cristofanelli, P., Arduni, J., Serva, F., Calzolari, F., Bonasoni, P., Busetto, M., Maione, M., Sprenger, M., Trisolino, P., and Putero, D.: Negative ozone anomalies at a high mountain site in northern Italy during 2020: a possible role of COVID-19 lockdowns?, *Environmental Research Letters*, 16, 074 029, <https://doi.org/10.1088/1748-9326/ac0b6a>, 2021.
- 410 Crutzen, P. J.: Tropospheric ozone: An overview, Springer, https://doi.org/10.1007/978-94-009-2913-5_1, 1988.
- Donzelli, G., Cioni, L., Cancellieri, M., Llopis-Morales, A., and Morales-Suárez-Varela, M.: Relations between air quality and COVID-19 lockdown measures in Valencia, Spain, *International Journal of Environmental Research and Public Health*, 18, 2296, <https://doi.org/10.3390/ijerph18052296>, 2021.
- 415

- Duncan, B. N., Yoshida, Y., Olson, J. R., Sillman, S., Martin, R. V., Lamsal, L., Hu, Y., Pickering, K. E., Retscher, C., Allen, D. J., and Crawford, J. H.: Application of OMI observations to a space-based indicator of NO_x and VOC controls on surface ozone formation, *Atmospheric Environment*, 44, 2213–2223, <https://doi.org/10.1016/j.atmosenv.2010.03.010>, 2010.
- EUROCONTROL: COVID-19 impact on the European air traffic network, <https://www.eurocontrol.int/covid19>, "accessed on 2022-03-09", 420 2022.
- Forster, P. M., Forster, H. I., Evans, M. J., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Le Quéré, C., Rogelj, J., Rosen, D., Schleussner, C.-F., Richardson, T. B., Smith, C. J., and Turnock, S. T.: Current and future global climate impacts resulting from COVID-19, *Nature Climate Change*, 10, 913–919, <https://doi.org/10.1038/s41558-020-0883-0>, 2020.
- Gaubert, B., Bouarar, I., Doumbia, T., Liu, Y., Stavrakou, T., Deroubaix, A., Darras, S., Elguindi, N., Granier, C., Lacey, F., Müller, J.-F., Shi, 425 X., Tilmes, S., Wang, T., and Brasseur, G. P.: Global changes in secondary atmospheric pollutants during the 2020 COVID-19 pandemic, *Journal of Geophysical Research: Atmospheres*, 126, 1–22, <https://doi.org/10.1029/2020JD034213>, 2021.
- Grange, S. K., Lee, J. D., Drysdale, W. S., Lewis, A. C., Hueglin, C., Emmenegger, L., and Carslaw, D. C.: COVID-19 lockdowns highlight a risk of increasing ozone pollution in European urban areas, *Atmospheric Chemistry and Physics*, 21, 4169–4185, <https://doi.org/10.5194/acp-21-4169-2021>, 2021.
- 430 Gurk, C., Fischer, H., Hoor, P., Lawrence, M., Lelieveld, J., and Wernli, H.: Airborne in-situ measurements of vertical, seasonal and latitudinal distributions of carbon dioxide over Europe, *Atmospheric Chemistry and Physics*, 8, 6395–6403, <https://doi.org/10.5194/acp-8-6395-2008>, 2008.
- Hasegawa, T.: Effects of Novel Coronavirus (COVID-19) on Civil Aviation: Economic Impact Analysis, https://www.icao.int/sustainability/Documents/Covid-19/ICAO_coronavirus_Econ_Impact.pdf, "accessed on 2022-03-09", 2022.
- 435 Higham, J., Ramírez, C. A., Green, M., and Morse, A.: UK COVID-19 lockdown: 100 days of air pollution reduction?, *Air Quality, Atmosphere & Health*, 14, 325–332, <https://doi.org/10.1007/s11869-020-00937-0>, 2021.
- Hosaynali Beygi, Z., Fischer, H., Harder, H., Martinez, M., Sander, R., Williams, J., Brookes, D., Monks, P., and Lelieveld, J.: Oxidation photochemistry in the Southern Atlantic boundary layer: unexpected deviations of photochemical steady state, *Atmospheric Chemistry and Physics*, 11, 8497–8513, <https://doi.org/10.5194/acp-11-8497-2011>, 2011.
- 440 IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation: Evaluated Kinetic Data, <http://iupac.pole-ether.fr>, "accessed on 2021-11-03", 2021.
- Jin, X., Fiore, A., Boersma, K. F., Smedt, I. D., and Valin, L.: Inferring Changes in Summertime Surface Ozone–NO_x–VOC Chemistry over US Urban Areas from Two Decades of Satellite and Ground-Based Observations, *Environmental science & technology*, 54, 6518–6529, <https://doi.org/10.1021/acs.est.9b07785>, 2020.
- 445 Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C. A., Brinkop, S., Cai, D. S., Dyroff, C., Eckstein, J., Frank, F., Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, M., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth system chemistry integrated modelling (ESCiMo) with the modular earth submodel system (MESSy) version 2.51, *Geoscientific Model Development*, 9, 1153–1200, <https://doi.org/10.5194/gmd-9-1153-2016>, 2016.
- 450 Khodayari, A., Tilmes, S., Olsen, S., Phoenix, D., Wuebbles, D., Lamarque, J.-F., and Chen, C.-C.: Aviation 2006 NO_x-induced effects on atmospheric ozone and HO_x in Community Earth System Model (CESM), *Atmospheric Chemistry and Physics*, 14, 9925–9939, <https://doi.org/10.5194/acp-14-9925-2014>, 2014.

- Khodayari, A., Olsen, S. C., Wuebbles, D. J., and Phoenix, D. B.: Aviation NO_x-induced CH₄ effect: Fixed mixing ratio boundary conditions versus flux boundary conditions, *Atmospheric Environment*, 113, 135–139, <https://doi.org/10.1016/j.atmosenv.2015.04.070>, 2015.
- 455 Kleinman, L. I., Daum, P. H., Lee, Y.-N., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J., and Rudolph, J.: Sensitivity of ozone production rate to ozone precursors, *Geophysical Research Letters*, 28, 2903–2906, <https://doi.org/10.1029/2000GL012597>, 2001.
- Klippel, T., Fischer, H., Bozem, H., Lawrence, M. G., Butler, T., Jöckel, P., Tost, H., Martinez, M., Harder, H., Regelin, E., Sander, R., Schiller, C. L., Stickler, A., and Lelieveld, J.: Distribution of hydrogen peroxide and formaldehyde over Central Europe during the HOOVER project, *Atmospheric chemistry and physics*, 11, 4391–4410, <https://doi.org/10.5194/acp-11-4391-2011>, 2011.
- 460 Kormann, R., Königstedt, R., Parchatka, U., Lelieveld, J., and Fischer, H.: QUALITAS: A mid-infrared spectrometer for sensitive trace gas measurements based on quantum cascade lasers in CW operation, *Review of scientific instruments*, 76, 075 102, <https://doi.org/10.1063/1.1931233>, 2005.
- Kroll, J. H., Heald, C. L., Cappa, C. D., Farmer, D. K., Fry, J. L., Murphy, J. G., and Steiner, A. L.: The complex chemical effects of COVID-19 shutdowns on air quality, *Nature Chemistry*, 12, 777–779, <https://doi.org/10.1038/s41557-020-0535-z>, 2020.
- 465 Lamprecht, C., Graus, M., Striednig, M., Sticherer, M., and Karl, T.: Decoupling of urban CO₂ and air pollutant emission reductions during the European SARS-CoV-2 lockdown, *Atmospheric Chemistry and Physics*, 21, 3091–3102, <https://doi.org/10.5194/acp-21-3091-2021>, 2021.
- Lee, D. S., Fahey, D., Skowron, A., Allen, M., Burkhardt, U., Chen, Q., Doherty, S., Freeman, S., Forster, P., Fuglestedt, J., Gettelman, A., Leon, R. D., Lim, L., Lund, M., Millar, R., Owen, B., Penner, J., Pitari, G., Prather, M., Sausen, R., and Wilcox, L.:
470 The contribution of global aviation to anthropogenic climate forcing for 2000 to 2018, *Atmospheric Environment*, 244, 117 834, <https://doi.org/10.1016/j.atmosenv.2020.117834>, 2021.
- Leighton, P.: Photochemistry of air pollution, Academic Press, Inc., New York, 1961.
- Lelieveld, J. and Dentener, F. J.: What controls tropospheric ozone?, *Journal of Geophysical Research: Atmospheres*, 105, 3531–3551, <https://doi.org/10.1029/1999JD901011>, 2000.
- 475 LI-COR, Inc.: LI-6262 CO₂/H₂O Analyzer Operating and Service Manual, <https://www.licor.com/documents/umazybbyhzalz840pf63a1137qz7ofib>, 1996.
- Lin, X., Trainer, M., and Liu, S.: On the nonlinearity of the tropospheric ozone production, *Journal of Geophysical Research: Atmospheres*, 93, 15 879–15 888, <https://doi.org/10.1029/JD093iD12p15879>, 1988.
- Liu, S., Trainer, M., Fehsenfeld, F., Parrish, D., Williams, E., Fahey, D. W., Hübler, G., and Murphy, P. C.: Ozone production in the rural
480 troposphere and the implications for regional and global ozone distributions, *Journal of Geophysical Research: Atmospheres*, 92, 4191–4207, <https://doi.org/10.1029/JD092iD04p04191>, 1987.
- Logan, J. A.: Tropospheric ozone: Seasonal behavior, trends, and anthropogenic influence, *Journal of Geophysical Research: Atmospheres*, 90, 10 463–10 482, <https://doi.org/10.1029/JD090iD06p10463>, 1985.
- Matthias, V., Quante, M., Arndt, J. A., Badeke, R., Fink, L., Petrik, R., Feldner, J., Schwarzkopf, D., Link, E.-M., Ramacher, M. O., and
485 Wedemann, R.: The role of emission reductions and the meteorological situation for air quality improvements during the COVID-19 lockdown period in central Europe, *Atmospheric Chemistry and Physics*, 21, 13 931–13 971, <https://doi.org/10.5194/acp-21-13931-2021>, 2021.
- Menut, L., Bessagnet, B., Siour, G., Mailler, S., Pennel, R., and Cholakian, A.: Impact of lockdown measures to combat Covid-19 on air quality over western Europe, *Science of the Total Environment*, 741, 140 426, <https://doi.org/10.1016/j.scitotenv.2020.140426>, 2020.

- 490 Mertens, M., Jöckel, P., Matthes, S., Nützel, M., Grewe, V., and Sausen, R.: COVID-19 induced lower-tropospheric ozone changes, *Environmental Research Letters*, 16, 1–9, <https://doi.org/10.1088/1748-9326/abf191>, 2021.
- Mills, G., Pleijel, H., Malley, C. S., Sinha, B., Cooper, O. R., Schultz, M. G., Neufeld, H. S., Simpson, D., Sharps, K., Feng, Z., Gerosa, G., Harmens, H., Kobayashi, K., Saxena, P., Paoletti, E., Sinha, V., and Xu, X.: Tropospheric Ozone Assessment Report: Present-day tropospheric ozone distribution and trends relevant to vegetation, *Elementa: Science of the Anthropocene*, 6, <https://doi.org/10.1525/elementa.302>, 2018.
- 495 Miyazaki, K., Bowman, K., Sekiya, T., Takigawa, M., Neu, J. L., Sudo, K., Osterman, G., and Eskes, H.: Global tropospheric ozone responses to reduced NO_x emissions linked to the COVID-19 worldwide lockdowns, *Science Advances*, 7, 1 – 14, <https://doi.org/10.1126/sciadv.abf7460>, 2021.
- National Research Council, C. o. T. O.: Rethinking the ozone problem in urban and regional air pollution, National Academy Press, Washington, D.C., 1991.
- 500 Nussbaumer, C. M. and Cohen, R. C.: The Role of Temperature and NO_x in Ozone Trends in the Los Angeles Basin, *Environmental Science & Technology*, 54, 15 652–15 659, <https://doi.org/10.1021/acs.est.0c04910>, 2020.
- Nussbaumer, C. M., Crowley, J. N., Schuladen, J., Williams, J., Hafermann, S., Reiffs, A., Axinte, R., Harder, H., Ernest, C., Novelli, A., Sala, K., Martinez, M., Mallik, C., Tomsche, L., Plass-Dülmer, C., Bohn, B., Lelieveld, J., and Fischer, H.: Measurement report: Photochemical production and loss rates of formaldehyde and ozone across Europe, *Atmospheric Chemistry and Physics Discussions*, pp. 1–29, <https://doi.org/10.5194/acp-2021-694>, 2021.
- 505 Nuvolone, D., Petri, D., and Voller, F.: The effects of ozone on human health, *Environmental Science and Pollution Research*, 25, 8074–8088, <https://doi.org/10.1007/s11356-017-9239-3>, 2018.
- Onyeaka, H., Anumudu, C. K., Al-Sharify, Z. T., Egele-Godswill, E., and Mbaegbu, P.: COVID-19 pandemic: A review of the global lockdown and its far-reaching effects, *Science Progress*, 104, 1–18, <https://doi.org/10.1177/00368504211019854>, 2021.
- 510 Ordóñez, C., Garrido-Perez, J. M., and García-Herrera, R.: Early spring near-surface ozone in Europe during the COVID-19 shutdown: Meteorological effects outweigh emission changes, *Science of the total environment*, 747, 141 322, <https://doi.org/10.1016/j.scitotenv.2020.141322>, 2020.
- Parrish, D., Lamarque, J.-F., Naik, V., Horowitz, L., Shindell, D., Staehelin, J., Derwent, R., Cooper, O., Tanimoto, H., Volz-Thomas, A., Gilge, S., Scheel, H.-E., Steinbacher, M., and Fröhlich, M.: Long-term changes in lower tropospheric baseline ozone concentrations: Comparing chemistry-climate models and observations at northern midlatitudes, *Journal of Geophysical Research: Atmospheres*, 119, 5719–5736, <https://doi.org/10.1002/2013JD021435>, 2014.
- 515 Pusede, S. and Cohen, R.: On the observed response of ozone to NO_x and VOC reactivity reductions in San Joaquin Valley California 1995–present, *Atmospheric Chemistry and Physics*, 12, 8323–8339, <https://doi.org/10.5194/acp-12-8323-2012>, 2012.
- 520 Pusede, S. E., Steiner, A. L., and Cohen, R. C.: Temperature and recent trends in the chemistry of continental surface ozone, *Chemical reviews*, 115, 3898–3918, <https://doi.org/10.1021/cr5006815>, 2015.
- Regelin, E., Harder, H., Martinez, M., Kubistin, D., Tatum Ernest, C., Bozem, H., Klippel, T., Hosaynali-Beygi, Z., Fischer, H., Sander, R., et al.: HO_x measurements in the summertime upper troposphere over Europe: a comparison of observations to a box model and a 3-D model, *Atmospheric Chemistry and Physics*, 13, 10 703–10 720, <https://doi.org/10.5194/acp-13-10703-2013>, 2013.
- 525 Reifenberg, S. F., Martin, A., Kohl, M., Hamryszczak, Z., Tadic, I., Röder, L., Crowley, D., Fischer, H., Kaiser, K., Schneider, J., Dörich, R., Crowley, J. N., Tomsche, L., Marsing, A., Voigt, C., Zahn, A., Pöhlker, C., Hollanda, B., Krüger, O., Pöschl, U., Pöhlker, M., Lelieveld,

- J., and Pozzer, A.: Impact of reduced emissions on direct and indirect aerosol radiative forcing during COVID-19 lockdown in Europe, manuscript in preparation, 2021.
- 530 Revell, L. E., Stenke, A., Tummon, F., Feinberg, A., Rozanov, E., Peter, T., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Butchart, N., Deushi, M., Jöckel, P., Kinnison, D., Michou, M., Morgenstern, O., O'Connor, F. M., Oman, L. D., Pitari, G., Plummer, D. A., Schofield, R., Stone, K., Tilmes, S., Visionsi, D., Yamashita, Y., and Zeng, G.: Tropospheric ozone in CCMI models and Gaussian process emulation to understand biases in the SOCOLv3 chemistry–climate model, *Atmospheric chemistry and physics*, 18, 16 155–16 172, <https://doi.org/10.5194/acp-18-16155-2018>, 2018.
- 535 Rosanka, S., Sander, R., Franco, B., Wespes, C., Wahner, A., and Taraborrelli, D.: Oxidation of low-molecular-weight organic compounds in cloud droplets: global impact on tropospheric oxidants, *Atmospheric Chemistry and Physics*, 21, 9909–9930, <https://doi.org/10.5194/acp-21-9909-2021>, 2021.
- Salma, I., Vörösmarty, M., Gyöngyösi, A. Z., Thén, W., and Weidinger, T.: What can we learn about urban air quality with regard to the first outbreak of the COVID-19 pandemic? A case study from central Europe, *Atmospheric Chemistry and Physics*, 20, 15 725–15 742, <https://doi.org/10.5194/acp-20-15725-2020>, 2020.
- 540 Schiller, C., Bozem, H., Gurk, C., Parchatka, U., Königstedt, R., Harris, G., Lelieveld, J., and Fischer, H.: Applications of quantum cascade lasers for sensitive trace gas measurements of CO, CH₄, N₂O and HCHO, *Applied Physics B*, 92, 419–430, <https://doi.org/10.1007/s00340-008-3125-0>, 2008.
- Schlosser, F., Maier, B. F., Jack, O., Hinrichs, D., Zachariae, A., and Brockmann, D.: COVID-19 lockdown induces disease-mitigating structural changes in mobility networks, *Proceedings of the National Academy of Sciences*, 117, 32 883–32 890, <https://doi.org/10.1073/pnas.2012326117>, 2020.
- 545 Schumann, U.: Aircraft emissions, *Encyclopedia of global environmental change*, 3, 178–186, 2002.
- Sillman, S., Logan, J. A., and Wofsy, S. C.: The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes, *Journal of Geophysical Research: Atmospheres*, 95, 1837–1851, <https://doi.org/10.1029/JD095iD02p01837>, 1990.
- 550 Steinbrecht, W., Kubistin, D., Plass-Dülmer, C., Davies, J., Tarasick, D. W., Gathen, P. v. d., Deckelmann, H., Jepsen, N., Kivi, R., Lyall, N., Palm, M., Notholt, J., Kois, B., Oelsner, P., Allaart, M., PETERS, A., Gill, M., Malderen, R. V., Delcloo, A. W., Sussmann, R., Mahieu, E., Servais, C., Romanens, G., Stübi, R., Ancellet, G., Godin-Beekmann, S., Yamanouchi, S., Strong, K., Johnson, B., Cullis, P., Petropavlovskikh, I., Hannigan, J. W., Hernandez, J.-L., Rodriguez, A. D., Nakano, T., Chouza, F., Leblanc, T., Torres, C., Garcia, O., Röhling, A. N., Schneider, M., Blumenstock, T., Tully, M., Paton-Walsh, C., Jones, N., Querel, R., Strahan, S., Stauffer, R. M., Thompson, A. M., Inness, A., Engelen, R., Chang, K.-L., and Cooper, O. R.: COVID-19 crisis reduces free tropospheric ozone across the Northern
- 555 Hemisphere, *Geophysical Research Letters*, 48, 1–11, <https://doi.org/10.1029/2020GL091987>, 2021.
- Stickler, A., Fischer, H., Williams, J., De Reus, M., Sander, R., Lawrence, M., Crowley, J., and Lelieveld, J.: Influence of summertime deep convection on formaldehyde in the middle and upper troposphere over Europe, *Journal of Geophysical Research: Atmospheres*, 111, 1–17, <https://doi.org/10.1029/2005JD007001>, 2006.
- Tadic, I., Crowley, J. N., Dienhart, D., Eger, P., Harder, H., Hottmann, B., Martinez, M., Parchatka, U., Paris, J.-D., Pozzer, A., Rohloff, R., Schuladen, J., Shenolikar, J., Tauer, S., Lelieveld, J., and Fischer, H.: Net ozone production and its relationship to nitrogen oxides and volatile organic compounds in the marine boundary layer around the Arabian Peninsula, *Atmospheric Chemistry and Physics*, 20, 6769–6787, <https://doi.org/10.5194/acp-20-6769-2020>, 2020.
- 560 Venter, Z. S., Aunan, K., Chowdhury, S., and Lelieveld, J.: COVID-19 lockdowns cause global air pollution declines, *Proceedings of the National Academy of Sciences*, 117, 18 984–18 990, <https://doi.org/10.1073/pnas.2006853117>, 2020.

- 565 Voigt, C., Lelieveld, J., Schlager, H., Schneider, J., Curtius, J., Meerkötter, R., Sauer, D., Bugliaro, L., Bohn, B., Crowley, J. N., Erbertseder, T., Groß, S., Li, Q., Mertens, M., Pöhlker, M., Pozzer, A., Schumann, U., Tomsche, L., Williams, J., Zahn, A., Andreae, M., Borrmann, S., Bräuer, T., Dörich, R., Dörnbrack, A., Edtbauer, A., Ernle, L., Fischer, H., Giez, A., Granzin, M., Grewe, V., Hahn, V., Harder, H., Heinritzi, M., Holanda, B., Jöckel, P., Kaiser, K., Krüger, O., Lucke, J., Marsing, A., Martin, A., Matthes, S., Pöhlker, C., Pöschl, U., Reifenberg, S., Ringsdorf, A., Scheibe, M., Tadic, I., Zauner-Wieczorek, M., Henke, R., and Rapp, M.: BLUESKY aircraft mission reveals reduction
570 in atmospheric pollution during the 2020 Corona lockdown, *Bulletin of the American Meteorological Society*, manuscript under review, 2021.
- Wang, P., Chen, Y., Hu, J., Zhang, H., and Ying, Q.: Attribution of tropospheric ozone to NO_x and VOC emissions: considering ozone formation in the transition regime, *Environmental science & technology*, 53, 1404–1412, <https://doi.org/10.1021/acs.est.8b05981>, 2018.
- WHO: WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020, <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>, "accessed on 2021-11-02", 2020a.
- WHO: Coronavirus disease 2019 (COVID-19) Situation Report – 51, https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200311-sitrep-51-covid-19.pdf?sfvrsn=1ba62e57_10, "accessed on 2021-11-02", 2020b.
- WHO: Coronavirus disease 2019 (COVID-19), https://www.who.int/health-topics/coronavirus#tab=tab_1, "accessed on 2021-11-02", 2021.
- 580 World Meteorological Organization: WMO Air Quality and Climate Bulletin, https://library.wmo.int/index.php?lvl=notice_display&id=21942#.YTIzN9, editors: O. R. Cooper, R. S. Sokhi, J. M. Nicely, G. Carmichael, A. Darmenov, P. Laj and J. Liggio, accessed on 2022-03-18, 2021.
- Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M., Neu, J., Parrish, D., Rieder, H., Schnell, J., et al.: Tropospheric Ozone Assessment Report: Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends, *Elementa: Science of the Anthropocene*, 6, <https://doi.org/10.1525/elementa.265>, 2018.
- 585 Zahn, A., Weppner, J., Widmann, H., Schlote-Holubek, K., Burger, B., Kühner, T., and Franke, H.: A fast and precise chemiluminescence ozone detector for eddy flux and airborne application, *Atmospheric Measurement Techniques*, 5, 363–375, <https://doi.org/10.5194/amt-5-363-2012>, 2012.
- Zhu, S., Poetscher, J., Shen, J., Wang, S., Wang, P., and Zhang, H.: Comprehensive insights into O₃ changes during the COVID-19 from
590 O₃ formation regime and atmospheric oxidation capacity, *Geophysical research letters*, 48, 1–11, <https://doi.org/10.1029/2021GL093668>, 2021.