

Spatial and temporal variations of CO₂ mole fractions observed at Beijing, Xianghe and Xinglong in North China

Yang Yang^{1,2,7}, Minqiang Zhou^{2,3,7}, Ting Wang^{2,7}, Bo Yao⁶, Pengfei Han⁵, Denghui Ji^{2,7}, Wei Zhou^{4,7}, Yele Sun^{4,7}, Gengchen Wang^{2,7}, and Pucai Wang^{2,7}

¹Shanghai Ecological Forecasting and Remote Sensing Center, Shanghai, China

²Key Laboratory of Middle Atmosphere and Global Environment Observation (LAGEO), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

³Royal Belgian Institute for Space Aeronomy, Brussels, Belgium

⁴State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry (LAPC), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

⁵State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics (LASG), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

⁶The China Meteorological Administration, Meteorological Observation Centre

⁷University of Chinese Academy of Sciences, Beijing, China

Correspondence: Minqiang Zhou (minqiang.zhou@aeronomie.be); Ting Wang (wangting@mail.iap.ac.cn)

Abstract. Atmospheric CO₂ mole fractions are observed at Beijing (BJ), Xianghe (XH), and Xinglong (XL) in North China using the Picarro G2301 Cavity Ring-Down Spectroscopy instruments. The measurement system is described comprehensively for the first time. The geo-distances among these three sites are within 200 km, but they have very different surrounding environments: BJ is inside the megacity; XH is in the suburban area; XL is in the countryside on a mountain. The mean and standard deviation of CO₂ mole fractions at BJ, XH, and XL between October 2018 and September 2019 are 448.4±12.8 ppm, 436.0±9.2 ppm and 420.6±8.2 ppm, respectively. The seasonal variations of CO₂ at these three sites are similar, with a maximum in winter and a minimum in summer, which is dominated by the terrestrial ecosystem. However, the seasonal variations of CO₂ at BJ and XH are more affected by human activities as compared to XL. By using CO₂ at XL as the background, CO₂ enhancements are observed simultaneously at BJ and XH. The diurnal variations of CO₂ are driven by the boundary layer height, photosynthesis and human activities at BJ, XH and XL. [We also compare the CO₂ measurements at BJ, XH and XL with 5 urban sites in US, and it is found that the CO₂ mean concentration at BJ is the largest.](#) Moreover, we address the impact of the wind on the CO₂ mole fractions at BJ and XL. This study provides an insight into the spatial and temporal variations of CO₂ mole fractions in North China.

1 Introduction

Carbon dioxide (CO₂) is the largest contributor to the total positive radiative forcing of the earth among anthropogenic gases. CO₂ has reached up to 140% relative to the pre-industrial level mainly due to fossil fuel combustion and land-use change (IPCC, 2013). The increase in CO₂ has led to an imbalance of $0.58 \pm 0.15 \text{ Wm}^{-2}$ in energy budget between 2005 and 2010 at the top of atmosphere (Hansen et al., 2011), resulting into changes in the atmospheric temperature, the sea level, and the hydrology.

Urban areas only take up around 2% of global land cover, while they emit more than 70% of CO₂ emissions from burning fossil fuels (Churkina, 2016). According to Gao et al. (2018), CO₂ emissions in metropolitan regions increased continuously from 1985 to 2006. Dhakal (2009) showed that China's urbanization rate has already reached 40% in 2005 and it is predicted to reach up to the level of 60% in 2030. This kind of increase certainly demands large quantities of energy consuming, leading to a large amount of CO₂ emissions.

It is important to understand atmospheric CO₂ variations in urban, suburban and rural areas. Previous studies carried out in urban areas, such as Phoenix, USA (Idso et al., 2013) and Copenhagen, Denmark (Soegaard and Møller-Jensen, 2003) show that CO₂ mole fractions are larger in the city center as compared to the outskirts, which is called "urban CO₂ dome". Various underlying surfaces, such as buildings, roads, trees, croplands, and grasslands cause complicate CO₂ characteristics (Cheng et al., 2018). George et al. (2007) pointed out that the horizontal gradients of CO₂ mole fractions among urban, suburban and rural areas are caused by different population densities and traffic volumes.

The Beijing-Tianjin-Hebei (BTH) area is an economically dynamic region, located in North China, with highly urbanized cities, suburban cities and rural areas (Figure 1). During the last two decades, the population in Beijing has increased from 13.64 million in 2000 to 21.54 million in 2018, the car amount increases from 1.04 million in 2000 to 5.74 million in 2018 (<http://data.stats.gov.cn/>). In the BTH area, the major CO₂ emissions are coming from industry, residential emissions, power plant and transportation (Song et al., 2013; Feng et al., 2019). In order to reduce the carbon emissions, Beijing has adopted a number of vehicle emission control strategies since the mid-1990s, for example, emission control on new and in-use vehicles, fuel quality improvements, alternative-fuel and advanced vehicles and public transport (Wu et al., 2011). During the China's 12th (2011-2015) and 13th (2016-2020) Five-Year Plan periods, comprehensive work programs have been implemented for energy conservation and emission reduction in Beijing. More recently, Beijing has also launched the short-term 'the three-year blue-sky defense battle of Beijing' between 2018 and 2020. Regional networks incorporated with high-accuracy CO₂ measurements can be used to retrieve carbon emissions and sinks in the horizontal gradient. The vertical gradient of CO₂ mole fractions can also be observed at several different heights at the same location (Bakwin et al., 1998).

To better understand the characteristics of CO₂ variations in the BTH area, 3 Cavity Ring-down Spectroscopy (CRDS) analyzers (Picarro G2301) within 200 km were installed at Beijing (BJ), Xianghe (XH), and Xinglong (XL). The three sites have very different surrounding environments: BJ is inside the megacity, XH is in the suburban area, and XL is in the countryside on a mountain. The measurements between June 2018 and April 2020 at the three sites allow us to better understand the differences among the urban, suburban and rural sites about the seasonal, synoptic and diurnal variations of CO₂ mole fractions. Section 2 describes the site locations as well as the measurement system. The results and discussions are presented in Section 3. Finally, the conclusions are drawn in Section 4.




Figure 1. (a): the Location of three sites at Beijing (BJ, 39.96 °N, 116.36 °E, 49 m a.s.l.), Xianghe (XH, 39.75 °N, 116.96 °E, 30 m a.s.l.) and Xinglong (XL, 40.40°N, 117.50 °E, 940 m a.s.l.), together with the land cover in this area. The red bars are the carbon dioxide emissions at the 3 sites based on the EDGAR data. The map within ~2 km of BJ (b), XH (c) and XL (c) are coming from © Google Maps (<https://www.google.com/maps>).

2 Measurements

50 2.1 Sites

The locations of the three sites at BJ (39.96 °N, 116.36 °E, 49 m above sea level (a.s.l.)), XH (39.75 °N, 116.96 °E, 30 m a.s.l.) and XL (40.40°N, 117.50 °E, 940 m a.s.l.) are shown in Figure 1. The red bars above the sites are the anthropogenic carbon dioxide emissions in 2015 from the Emission Database for Global Atmospheric Research (EDGAR) v5.0 (Crippa et al., 2019). The CO₂ fluxes are 2.02×10^{-6} , 1.12×10^{-7} , and 2.45×10^{-8} kg m⁻² s⁻¹ at BJ, XH and XL, respectively.

55 The BJ site is located in a highly urbanized area, with dense buildings, shopping centers, roads and residential districts. To the east of the site, there is the Beijing-Tibet expressway (G6) carrying a heavy volume of traffic. Within 1 km of the site,




Figure 2. The schematic diagram of measurement system, including a meteorological tower at BJ/XH or observation building at XL, an intake system, a calibration unit and a CRDS analyzer.

the heights of trees are about 15-20 m, and the heights of buildings are about 70-200 m (Cheng et al., 2018). The vegetation fractions around the BJ site are between 10% and 18% (Liu et al., 2012).

The XH site is in a suburban area about 50 km to the southeast of Beijing. XH is surrounded by croplands and irrigated croplands. Within 1 km of the XH site, the residential houses are mainly home-built, with an average height of ~20 m. The center of Xianghe county is about 2 km to the east of the site.

The XL site is located on a mountain, inside the Xinglong Observatory of the National Astronomical Observatories, Chinese Academy of Sciences (NAOC) (<https://www.xinglong-naoc.org/html/en/>), which is about 120 km to the northeast of Beijing. XL is located in a highly vegetated area.

65 2.2 Instrumentation Measurement system

The Picarro Cavity Ring-Down Spectroscopy (CRDS) G2301 analyzers were installed at BJ, XH, and XL to measure CO₂, CH₄, and H₂O mole fractions. The same measurement system is operated at these three sites, which is composed of an intake system, a calibration unit, and a Picarro analyzer (Figure 2). Note that there are two sampling heights at BJ (80 and 280 m above ground level (a.g.l.)) and XH (60 and 80 m a.g.l.), but only one sampling height (10 m a.g.l.) at XL. The measurements start in June 2018 at BJ and XH, and in May 2016 at XL. To compare the CO₂ measurements among these sites, we focus on the data after June 2018 in this study.

2.2.1 Intake system

The surrounding air is sampled by a vacuum pump (DA7002D) with a maximum flux of 20 L.min⁻¹ through an inlet (Figure 2). The sample air is then introduced into a 10 mm-diameter tube (SYNFLEX 1300), mounted with a capsule filter (Whatman, USA) to filter out the solid particle with a diameter larger than 2 μm and liquid particles with a diameter larger than 0.03 mm. In addition, a 7 mm sintered filter (Membrane) is installed to filter out the solid particle with a diameter larger than 7 μm. Moreover, an air compressor and a dry machine together with a single Nafion tubing selectively permeable membrane dryer

(MD-110-72P-4; Perma Pure, Halma, UK) in self-purge are installed to remove water vapor. The sample dew-point temperature can reach down to -25 °C, corresponding to a relative humidity of 1-20 %. The flux of the Nafion outflow is 200-400 ml min⁻¹.

80 The outflow is then vented to the unloading valve (Figure 2), which guarantees that the air fed to the Picarro G2301 analyzer is controlled at near-ambient pressure. Before the ambient CO₂ measurements, the sampled air is introduced to the calibration unit to check the precision and stability of the system, which will be introduced in detail in Section 2.3.

2.2.1 Calibration unit

~~The~~ The last part of the measurement system is the Picarro analyzer, which is composed of a laser, a high-finesse optical cavity, and a detector. The sample air is first introduced into the cavity. After that, the laser passes through the sample air and the intensity of the laser arriving at the detector is monitored as I . Then, the 'ring-down' measurements start as the laser rapidly shuts down. Meanwhile, the sample gas is measured by recording the decay of the laser intensity with time. This decay depends on the optical path inside of the cavity, which is in correlation with the absorption and scattering of the sample air. The analyzer continuously scans the laser over CO₂ spectral features and records the absorption loss at a wavelength of 1603 nm to form the spectrum. As a result, CO₂ mole fractions are derived from these spectra and collected by the Data Acquisition part.

85
90

2.3 Calibration method

As is shown in Figure 2, the intake system is connected to an 8-position valve, which is used to choose the air coming from the sample air, the target gas, or the calibration gas. The target and calibration gases are pressurized in 29.5 L treated aluminum alloy cylinders, which are scaled to the WMO X2007 standard by the China Meteorological Administration, Meteorological
95 Observation Centre. The same calibration procedure is operated at these three sites: 1) 3-hours sample air; 2) 5-minutes calibration gas; 3) 3-hours sample air; 4) 5-minutes target gas. This process repeats every 6 hours and 10 minutes. Note that, the airs coming from two levels at XH and BJ are switched every 5 minutes during the 3-hours sample air period. As the remaining volume in the tubes needs time for flushing, the response of the analyzer turns to be stable about 1 minute after each switching. In order to reduce the uncertainty, we do not consider the first 3-minutes measurements after each switching.

100 The calibration gas is to calculate the calibration factor (cf),

$$cf = \overline{CO_{2,mcal}} / CO_{2,cal}, \quad (1)$$

where $CO_{2,mcal}$ is the CO₂ mole fraction measured by the Picarro analyzer from the calibration gas and $CO_{2,cal}$ is the standard CO₂ mole fraction of the calibration cylinder. cf is applied to correct the sample air during the next 6 hours,

$$CO_{2,c} = cf \times CO_{2,m}, \quad (2)$$

105 where $CO_{2,m}$ is the CO₂ mole fraction measured by the Picarro analyzer and $CO_{2,c}$ is the calibrated CO₂ mole fraction.

The target gas is used to check the precision and stability of the system. The T value are calculated as follows,

$$T = cf \times CO_{2,mtar} - CO_{2,tar}, \quad (3)$$

where $CO_{2,tar}$ is the standard CO₂ mole fraction of the target gas cylinder, $CO_{2,mtar}$ is the CO₂ mole fraction measured by the Picarro analyzer from the target gas. ~~In this study, we only select the period-~~

110 To keep the CRDS stable over time, only the periods with T value within ± 0.1 ppm are selected (Fang et al., 2014). The measurement uncertainties of the Picarro instrument at the three sites are calculated as the standard deviation (std) of T, which are 0.01, 0.06, and 0.02 ppm at BJ, XH, and XL respectively. ~~What's more, to reduce the uncertainties of the inlet filter, pump, and extreme weather, we manually flag and filter out the abnormal-~~

2.4 Data quality control

115 Besides the calibration procedure mentioned in Section 2.3, we also do auto and manual flagging of the raw data. In each 1-hour CO₂ measurements at these three sites.

2.4.1 ~~Picarro analyzer~~

~~The Picarro analyzer is composed of a laser, a high-finesse optical cavity, and a detector. The sample air is first introduced into the cavity. After that, the laser passes through the sample air and the intensity of the laser arriving at the detector is monitored as I . Then, the 'ring-down' measurements start as the laser rapidly shuts down. Meanwhile, the sample gas is measured by recording the decay of the laser intensity with time. This decay depends on the optical path inside of the cavity, which is in correlation with the absorption and scattering of the sample air. The analyzer continuously scans the laser over measurement window, auto-flags are assigned when deviations from CO₂ mean are found larger than 2-times hourly CO₂ std. Furthermore, manual flags are assigned by technicians at each site according to the logbook to exclude no-valid data resulted from the inlet filter, pump, and extreme weather issues. In addition, as the CRDS measurement system records CO₂ spectral features and records the absorption loss at a wavelength of 1603 nm to form the spectrum. As a result, and CH₄ simultaneously, the variations of these two gases are checked together to manually flag CO₂ mole fractions are derived from these spectra and collected by the Data Acquisition part. The temporal resolution of the CO₂ measurements is 2 seconds. CH₄ outliers.~~

120
125

2.5 Meteorological fields

130 The CO₂ variations are additionally characterized by specific meteorological parameters, such as local wind and temperature fields. The ~~local~~ meteorological sensors at BJ are installed at the same tower as the Picarro on 120 m a.g.l., and the meteorological sensors at XL are ~5 m northwest to the Picarro sample tube. The meteorological fields at XH are not discussed here as there is a technical issue with the wind sensor.

Figure 3 shows the wind frequencies at BJ and XL in each season, which are binned with a resolution of 2 m.s⁻¹ for the wind speed and 10° for the wind direction. At BJ, two dominant wind regimes are observed throughout the whole year: north (northwest to northeast clockwise) and southwest. The percentage of wind frequency in the north region is 34%, 36%, 50% and 60% respectively from spring to winter. The wind speed varies from 0.63 m.s⁻¹ on 10 May 2019 to 14.98 m.s⁻¹ on 20 December 2018, with a mean of 3.92 m.s⁻¹. From spring to autumn, more winds are with a low wind speed. However,

135




Figure 3. Wind frequency as a function of wind speed (m.s^{-1}) and wind direction ($^{\circ}$) in each season spring (MAM), summer (JJA), autumn (SON) and winter (DJF) at BJ (a) and XL (b) from October 2018 to September 2019.

in winter, the prevailing northwest wind contributes to high wind frequencies with the increase of wind speed. At XL, the dominant winds are mainly from the west (southwest to northwest clockwise), together with some winds from the southeast. The percentage of wind frequency in the west region is 52%, 33%, 56% and 57% respectively from spring to winter. The wind speed varies from near-zero on 18 August 2019 to 10.75 m.s^{-1} on 17 April 2019, with a mean of 2.52 m.s^{-1} .

The atmospheric boundary layer height (BLH) is an another important parameter to characterize the diurnal variation of CO_2 (Li et al., 2014; Culf et al., 1997). In this study, we use the BLH hourly data of the ERA5 reanalysis data from the European Centre for Medium-Range Weather Forecasts (ECMWF) with a spatial resolution of $0.25^{\circ} \times 0.25^{\circ}$ (Hersbach et al., 2020).

3 Results and discussions

3.1 CO_2 time series and comparison with other urban sites

Figure 3-4 shows the time series of hourly CO_2 mole fractions at the three sites between June 2018 and March 2020. The two-levels (80 m and 280 m) measurements at BJ are marked as BJ L1 and BJ L2, and the two-levels (60 m and 80 m) measurements at XH are marked as XH L1 and XH L2. The gaps in the CO_2 time series are due to the malfunctions of the instruments. To better understand the influence of the wind on CO_2 , we classify the CO_2 mole fractions at XL and BJ L1 based on the wind information into five classes respectively (Fig. 4a and b). The BJ L1 is used here as it is closer to the wind sensor as compared to BJ L2. The local class is defined as wind speed less than 2 m.s^{-1} , while the wind speed larger than 2 m.s^{-1} are classified into four sections according to the wind direction: northwest (NW), northeast (NE), southwest (SW) and southeast (SE).




Figure 4. The time series of the CO₂ measurements at XL (a), BJ L1 (b), BJ L2 (c), XH L1 (d) and XH L2 (e) between June 2018 and March 2020. The CO₂ measurements at XL (a) and BJ LI (b) are colored by wind classes discussed in the text.

155 As expected, the urban BJ site observes a much higher CO₂ level than the suburban XH and rural XL sites. The CO₂ measurements at the urban site BJ L1 (Fig. 4b) are influenced by the wind speeds and wind directions. High CO₂ mole fractions generally appear in local class throughout the whole year, indicating the strong local anthropogenic emissions. The north sectors (NS and NE) usually contribute low CO₂ mole fractions during the autumn-winter period. However, in spring and summer, the SW sector contributes lower CO₂, indicating the low CO₂ varies with the wind direction season by season at
 160 BJ. Different from BJ, the CO₂ mole fraction in the local class at XL covers all the data range throughout the whole year. In spring and summer, the wind from the south (SE and SW) makes CO₂ increase at XL.




Figure 5. (a) Monthly means of CO₂ at BJ (L1), XH (L1), XL between October 2018 and September 2019, at BU, CRA, COM, IMC and SF during the latest 1 year and (b) the diurnal cycles of CO₂.

As XL is located in a rural area that is less influenced by human activities. Comparisons with other five urban sites in USA with a similar latitude of BJ are also discussed in this section. All these five sites belong to the CO₂ Urban Synthesis and Analysis (CO₂-USA) Data Synthesis Network (Feng et al., 2016). The site locations, elevations, inlet heights, and references are listed in Table 1. As the CO₂ measurements at these five sites do not cover the period between October 2018 and September 2019, we use the latest 1-year available CO₂ measurements. The monthly means and diurnal cycles of CO₂ at BJ (L1), XH (L1), and 5 American urban sites are shown in Figure 5. It is found that the phases of the seasonal CO₂ cycles at BU, CRA, COM, IMC and SF are consistent with the observations at BJ (L1), XH (L1) and XL, with a high value in autumn-winter and a low value in summer. Among the five American sites, the highest CO₂ concentration is observed at IMC. The IMC site is inside a commercial zone and the CO₂ measurements over there are more strongly influenced by local emissions over there (Bares et al., 2019). The CO₂ concentration is also high at COM, because the Los Angeles megacity is one of the largest fossil fuel CO₂ emitters in the world (Matthäus et al., 2021). Figure 5 (a) shows that the CO₂ concentrations at COM and IMC are in the same level with the one at XH, but are less than the CO₂ concentration at BJ. The CO₂ concentrations at SF, BU and CRA are much lower as compared to BJ and XH, because of lower anthropogenic emissions at these sites (McKain et al., 2015; Lauvaux et al., 2016; Shusterman et al., 2016).

Figure 5 (b) shows the diurnal variations of CO₂, with the amplitudes of 22.4, 19.4, 6.6, 16.3, 14.8, 41.5, 41.1 and 37.2 ppm at BJ (L1), XH (L1), XL, BU, CRA, COM, IMC and SF, respectively. The amplitudes of the diurnal variation at COM, IMC and SF are higher than that at BJ, although the yearly mean CO₂ at-levels at these sites are smaller than that at BJ. As the sampling heights at these sites and BJ are similar, the large amplitudes of the diurnal variation indicate that stronger variation in the local emissions and/or sinks exists at these three American sites as compared to BJ.

Table 1. Site characteristics of BJ, XH and XL in North China, BU, CRA, COM, IMC and SF in USA from the CO₂ Urban Synthesis and Analysis (CO₂-USA) Data Synthesis Network.

Site Code	Site Name	Lat (°N)	Lon (°E)	Elevation (m a.s.l.)	Inlet Height (m a.g.l.)	City	Reference
BJ	Beijing	39.96	116.36	49	80/280	Beijing	Cheng et al. (2018)
XH	Xianghe	39.75	116.96	30	60/80	Xianghe	Yang et al. (2020)
XL	Xinglong	40.40	117.50	940	10	Xinglong	Yang et al. (2019)
BU	Boston University	42.35	-71.10	4	29	Boston	Sargent et al. (2018) McKain et al. (2015)
CRA	Crawfordsville	39.99	-86.74	264	76	Indianapolis	Lauvaux et al. (2016) Richardson et al. (2017)
COM	Compton	33.87	-118.28	9	45	Los Angeles	Verhulst et al. (2017)
IMC	Intermountain Medical Center	40.67	-111.89	1316	66	Salt Lake City	Mitchell et al. (2018) Bares et al. (2019)
SF	SF Hospital Bldg 5	37.76	-122.41	23.9	52	San Francisco	Shusterman et al. (2016)

3.2 Contribution of main CO₂ sources

We use the CarbonTracker model, version CT-NRT.v2021-3 (Peters et al., 2005) to evaluate the influence of anthropogenic, biogenic, oceanic and fire sources at these three sites respectively. The CarbonTracker is a data assimilation system developed by the National Oceanic and Atmospheric Administration (NOAA) to keep track of sources and sinks of atmospheric CO₂ around the world. Four tracers (biosphere, ocean, fire and fossil fuel) are treated separately to simulate atmospheric CO₂ mole fractions. Mustafa et al. (2020) evaluated the CarbonTracker model in Asia by comparing with satellite measurements, and they found that the CarbonTracker model captures the variation of CO₂ well. The model provides 3-hourly CO₂ data at 25 levels from surface to ~123 km, and the spatial resolution of the global CarbonTracker CO₂ simulation is 3° × 2° (longitude × latitude). As BJ and XH are in the same model grid, we note the CO₂ simulations in the BJ/XH grid as BJ.

Figure 6 shows the time series of CO₂ simulations from fossil fuel (CO_{2,ff}), biosphere (CO_{2,bio}), fire (CO_{2,fire}) and ocean (CO_{2,oce}) modules at BJ/XH and XL between October 2018 and September 2019. It is found that the fire and ocean CO₂ at BJ/XH and XL are close to each other throughout the whole year. According to the Global Fire Assimilation System (GFAS) (<https://www.ecmwf.int/en/forecasts/dataset/global-fire-assimilation-system/>) wildfire emissions, there is almost no biomass burning CO₂ emissions at BJ, XH and XL sites. The CarbonTracker model simulations confirm that fire CO₂ concentrations in this region are almost the same, and the simulated fire CO₂ at these sites are transported by the wildfire emissions at other places. What's more, the CarbonTracker model suggests that the fire CO₂ at these sites only take up a small proportion of the observed CO₂ (less than 5%). The biogenic CO₂ at BJ/XH and XL have a similar level between October 2018 and June 2019, and become slightly different in summer 2019. However the difference in biogenic CO₂ is much less than that of

the anthropogenic CO₂ differences. The high CO₂ concentrations at BJ and XH in winter are apparently dominated by the enhancement of fossil fuel. The variation of the fossil fuel CO₂ at XL is much less than that at BJ/XH. Therefore, by using the CO₂ measurements at XL as the background, we can significantly reduce the influence from fire, biosphere and ocean, and extract the signal of the anthropogenic CO₂ differences.

The CO₂ enhancement at BJ or XH relative to XL is then calculated as

$$\Delta CO_{2,BJ/XH} = CO_{2,BJ/XH} - CO_{2,XL} \quad (4)$$

The time series of hourly $\Delta CO_{2,BJ/XH}$ are presented in Figure 7a. The ΔCO_2 has a maximum in winter and a minimum in summer at both BJ and XH. The high value is probably related to more combustion of fossil fuel from traffic and heating systems in winter (Liu et al., 2012). The daily ΔCO_2 can reach up to 106.8 ppm in December 2018 at BJ and 78.5 ppm in January 2019 at XH. The mean ΔCO_2 at BJ and XH are 26.2 ± 20.6 ppm and 15.2 ± 13.6 ppm, respectively.

There are 271 days when ΔCO_2 are observed at both BJ and XH (Figure 7b). The correlation efficiency (R) of 0.81 is found between the ΔCO_2 at BJ and XH, indicating that the ΔCO_2 s change simultaneously at BJ and XH. The slope of the linear fitting suggests that the ΔCO_2 at BJ is 1.23 times larger than that of XH.

3.3 Seasonal variations

The seasonal cycles of CO₂ are derived from the measurements at the lower levels at BJ and XH, and the measurements at XL. The lower levels at BJ and XH are used here as they reflect more information about surface fluxes. Figure 8a shows the CO₂ monthly means between October 2018 and September 2019, together with the temperature at BJ and leaf area index (LAI). The LAI monthly data are from the Copernicus Global Land Service (<https://land.copernicus.eu/global/products/lai>) with a spatial resolution of 1 km. Figure 8a shows the LAI monthly means in the region of Fig. 1.

Between October 2018 and September 2019, the mean of CO₂ mole fractions at BJ is 448.4 \pm 12.8 ppm, which is larger than those at XH (436.0 \pm 9.2 ppm) and XL (420.6 \pm 8.2 ppm). The phases of the seasonal cycle of CO₂ at BJ, XH and XL are similar, with a high value in autumn-winter and a low value in summer, which is consistent with other observations in North Hemisphere (Nevison et al., 2008). It is expected mainly due to the seasonal cycle of the biosphere fluxes (LAI). The increased temperature in summer is favorable for plant growth, leading to larger photosynthesis. In winter, the respiration of plants and the anthropogenic heating emissions contribute to a high CO₂ level. The amplitudes of the seasonal variation of CO₂ at BJ, XH and XL are 60.3-41.2 ppm, 36.1 ppm and 29.3 ppm, respectively. According to the CarbonTracker simulation (Figure 6), the CO₂ seasonal cycle in this region is mainly driven by the biogenic and anthropogenic CO₂. At XL, the anthropogenic CO₂ is almost constant through the whole year, while the biogenic CO₂ is low in summer and high in winter. For BJ/XH, apart from the similar biogenic CO₂ seasonal variation, the anthropogenic CO₂ is also high in winter and lower in summer. Therefore, combining the effect from the biosphere and human activities, the amplitude of CO₂ seasonal variation at BJ/XH is larger than that at XL. What's more, as the anthropogenic emission at BJ is much larger than that at XH, indicated by the EDGAR emission dataset, we thus observe the largest amplitude of the seasonal variation at BJ.




Figure 6. The time series of CO₂ simulations from fossil fuel (CO_{2,ff}), biosphere (CO_{2,bio}), fire (CO_{2,fire}) and ocean (CO_{2,oce}) modules at BJ/XH and XL.

Figure 8b, c and d show the CO₂ monthly means together with the monthly 1σ standard deviation at each site. We take the days when measurements are available at all three sites or the days when measurements are available at XH and XL. The CO₂ variability (1σ) is highest at BJ and lowest at XL. The seasonal CO₂ variation and 1σ standard deviation at each site are further assessed in the following.

235 **Autumn.** At each site, monthly mean CO₂ mole fractions are increasing with the decrease of LAI. The increase rates of CO₂ at BJ, XH and XL are 30, 19 and 9 ppm/month, respectively. The 1σ standard deviation of each month at BJ is generally larger than that of XH, then followed by XL.

240 **Winter.** The CO₂ removed by the photosynthesis is weak in this region as the LAI is low. The CO₂ change simultaneously at BJ and XH, increasing from December 2018 to January 2019 and decreasing afterwards. Similar to autumn, the month-to-month variation of CO₂ at BJ is larger than those at BJ and XL, together with the largest 1σ at BJ. The 1σ at BJ and XH is larger in winter as compared to other seasons.




Figure 7. (a): the time series of daily CO₂ enhancements at BJ and XH relative to XL between October 2018 and September 2019. The blue and black lines are the monthly means of CO₂ enhancements at BJ and XH, respectively. (b): the correlation between daily mean CO₂ enhancements at BJ and XH.

Spring. The decrease of CO₂ in March 2019 is highly related to the temperature increase. As the heating is officially stopped in the middle of March, the anthropogenic emissions are much reduced (Shi et al., 2020). In April and May, the LAI increases significantly, leading to the decrease of CO₂, especially at XL. The regional biosphere activity affects more on CO₂ mole fractions at XL, while the large anthropogenic emissions at BJ and XH may reduce the influence from the photosynthesis.

Summer. At all the sites, the minimum CO₂ is observed in August with the maximum LAI corresponding to the largest photosynthesis CO₂ absorption activity. The month-to-month variation of 1σ is small at BJ and XH.

3.4 Diurnal variations

The diurnal variations of CO₂ at BJ, XH and XL between October 2018 and September 2019 are shown in Figure 9. The amplitudes of the diurnal variations are between 16.4 ppm and 44.1 ppm at BJ. The relatively large amplitudes are observed in summer and winter compared to spring and autumn. The phase of the diurnal variation at BJ varies with season. There are one peak in the early morning (4:00-7:00) and one trough in the afternoon (14:00-16:00) in spring and summer. However, there are two peaks (8:00-9:00, 22:00-1:00), and two troughs (4:00-7:00, 14:00-16:00) in late autumn and winter. At XH, there are one peak (4:00-7:00) and one trough (14:00-16:00) throughout the whole year. The amplitude of the diurnal variation at XH is about 6-20 ppm smaller than that at BJ between November 2018 and May 2019. At XL, the peak of CO₂ occurs around 4:00-7:00, and the trough occurs in the afternoon around 12:00-14:00. The amplitudes of diurnal variations at XL are larger in summer as compared to other seasons. Moreover, the amplitudes of diurnal variations at XL are much smaller as compared to those at BJ and XH, especially in winter.

The diurnal variations of CO₂ are mainly affected by the BLH, photosynthesis, and local human activities (Chan et al., 2008; Denning et al., 1999). Generally, the increase of sunlight enhances the plant photosynthetic rate, vice versa. There is no




Figure 8. (a): the monthly means of CO₂ at BJ L1, XH L1 and XL between October 2018 and September 2019. The monthly mean air temperature at BJ and regional mean Leaf Area Index (LAI) of the area in Figure 1a during the same period are also displayed. (b-d): monthly means of CO₂ together with the 1 σ standard deviation at BJ L1, XH L1 and XL between June 2018 and February 2020. The gap at BJ L1 is due to the instrument failure. The shadow is the measurement period displayed in Figure 8a.




Figure 9. The diurnal cycles of CO₂ variations at BJ L1, XH L1 and XL in each month between October 2018 and September 2019. The collocated days are displayed (N). The error bars are the hourly standard deviations of CO₂.




Figure 10. (a-c): mean diurnal cycles of BLH from ERA5 and mean diurnal CO₂ variations at BJ L1 (a), XH L1 (b) and XL (c) in each season between October 2018 and September 2019.

photosynthetic CO₂ sink before sunrise or after sunset (Lv et al., 2020; Bagley et al., 2015). To better understand the influence of the BLH on the diurnal CO₂ variations, we show the CO₂ diurnal cycles for each season at BJ L1, XH L1 and XL, together with the BLH hourly means.

BJ L1. The increase of the BLH after sunrise (5:00 - 8:00) and the photosynthetic uptake during the day make the CO₂ mole fraction decrease. The CO₂ mole fraction reaches a minimum in the afternoon around 16:00-17:00, corresponding to the maximum BLH. After that, the BLH decreases resulting into the accumulation of CO₂. In spring and summer, the CO₂ mole fraction keeps increasing until the next day (5:00-8:00) before sunrise, and in autumn and winter, the CO₂ mole fraction starts decreasing at midnight. Note that the enhancement of CO₂ around 9:00 in winter is not related to the BLH, which is probably due to the rush traffic emission.

XH L1. Similar to BJ, the variation of the CO₂ mole fraction is dominated by the BLH during the day. The CO₂ mole fraction decreases with the increase of BLH. The CO₂ mole fraction reaches a minimum in the afternoon around 16:00-17:00, corresponding to the highest BLH. However, at night, the variation of CO₂ at XH is not the same as that at BJ, especially in autumn and winter. In autumn, the CO₂ mole fraction keeps increasing until next day before sunrise (5:00-8:00), and in winter, the CO₂ mole fraction stays stable after midnight. Similar to BJ, the peak CO₂ around 9:00-10:00 in winter may be due to the traffic emission in the rush hour.

XL. Different from BJ and XH, the minimum of the CO₂ mole fraction occurs earlier than the maximum of BLH in spring and summer. For example, the minimum of the CO₂ mole fraction is around 12:00 and the maximum of BLH occurs around 16:00. The solar radiation is strongest at noon, which leads to the largest photosynthesis removing CO₂. The diurnal variation of CO₂ at daytime is then strongly affected by the plants in these two seasons. However, in autumn and winter, the minimum of the CO₂ mole fraction occurs close to the maximum of the BLH, which is probably also dominated by the change of PBL due to the low LAI in these two seasons (Mohotti and Lawlor, 2002; Newman et al., 2013).

3.5 CO₂ variations with the wind

Wind speed and wind direction are the two key factors in modulating the dispersion of CO₂ emissions (Turnbull et al., 2015; Lac et al., 2013; ángeles García et al., 2012). The influence of wind on CO₂ mole fraction at BJ and XL is discussed specifically in this section. To minimize the influence from the diurnal variation, we focus on the measurements between 14:00 and 16:00 during daytime for the highest BLH, and between 22:00 and 02:00 during nighttime for the lowest BLH. Besides, we reduce the impact from the seasonal variation of CO₂ by applying the following method. First, we calculate the mean of CO₂ over 10 days ($CO_{2,10d}$). Second, the ratio between the $CO_{2,10d}$ and the annual mean of original CO₂ is derived ($Index_{10d} = CO_{2,10d}/CO_{2,mean}$), and the $Index_h$ is interpolated from the $Index_{10d}$ at an hourly scale. Finally, The deseasonalized CO₂ is calculated as $CO_{2,de} = CO_2/Index_h$. In summary, we use the deseasonalized CO₂ during the daytime (14:00-16:00) and the nighttime (22:00-02:00) separately to understand the influence of the wind.

Figure 11 shows the daytime and nighttime wind roses of CO₂ mole fractions at BJ and XL, with a resolution of 1 m.s⁻¹ wind speed and of 10 °wind direction. Note that only the bins with the measurement number larger than 3 at BJ or 5 at XL are shown here.

BJ. At BJ, the wind mainly comes from the southwest and the northwest, with more winds come from the southwest during the day and more winds come from the northwest at night. The high CO₂ mole fractions are observed with a low wind speed (<2 m.s⁻¹). For the wind with a relatively large speed (>2 m.s⁻¹), it is found that the CO₂ with the wind coming from the southwest is about ~21 ppm larger than those with the wind coming from the northwest during the day.

XL. The wind speed at XL is generally smaller as compared to BJ. The wind at XL is mainly coming from the southeast-northwest sector in a clockwise direction. During the day, the high CO₂ mole fractions are observed along with a relatively large wind speed (>2 m.s⁻¹). This can be attributed to the impact of remote emissions advocated from the south, where large cities, such as Beijing and Tianjin, are located. At night, although the dominant wind shifts to the west, the high CO₂ mole fractions can be observed in almost all the directions with wind speeds ranging from 0 to 3 m.s⁻¹.

3.6 Two-levels measurements at BJ and XH

Figure 12 shows the CO₂ hourly means observed at two levels at BJ and XH between October 2018 and September 2019. Note that, we select measurements when the hourly means are available at both levels.

At BJ, CO₂ mole fractions at L1 are generally higher than L2, ~~but the~~ as L1 is closer to near-ground human emissions. At BJ L1 (80 m a.g.l.), we can observe a peak in the early morning, which is corresponding to the transportation rush hour. The valley of CO₂ mole fractions at at BJ L1 are lower than that at occurs at 16:00-17:00 because of the maximum PBL resulting from the unstable atmosphere. After that, the atmosphere changes from unstable to stable during the night, leading to the CO₂ peak again. At BJ L2 occasionally in the afternoon in August and September 2019. The diurnal variations (280 m a.g.l.), the diurnal variation of CO₂ generally follows that at L1 and-. Note that the peak of the CO₂ at L2 are similar in spring and summer, but not in autumn and winter. occurs in the early morning later than that at L1 as the CO₂ at the ground level moved upward with the increase in convective PBL, with a large difference in winter and a small difference in summer. The CO₂ diurnal variations




Figure 11. (1): binned CO₂ mole fraction as a function of wind speed (m.s⁻¹) and wind direction (°) at BJ L1 (a, b) and XL (c, d) based on daytime (14:00-16:00 LTC) and nighttime (22:00 -1:00 LTC) data between October 2018 and September 2019. (2): mean 1σ standard deviation of the CO₂ mole fractions in each bin. (3): the CO₂ measurement counts in each bin.

315 from two-layers Picarro measurements in 2018 and 2019 in our study are consistent with the seven open-path infrared gas analyzers (Model LI-7500A; at 8, 16, 47, 80, 140, 200 and 280 m a.g.l.) measurements between 2013 and 2016 at the same site (Cheng et al., 2018). In summer, the temperature is high due to a larger solar irradiance, the atmosphere becomes unstable quickly accelerating the uplifting of the PBL. In winter, the uplifting of the PBL is slow because of the stable atmosphere.

At XH, the CO₂ mole ~~fraction-fractions~~ at L1 ~~is slightly larger than that at and~~ L2 ~~, especially at night. The diurnal variations~~
320 ~~at two levels at XH are close are~~ closer to each other ~~, which is due to the fact that as compared to the two-layers measurements~~
~~at BJ, because~~ the difference in the ~~altitudes between the two levels~~ vertical distance of two layers at XH is only 20 m. Nevertheless, we can still observe that the peak of the CO₂ at L2 occurs in the early morning later than that at L1 as the CO₂ at the ground level moved upward with the increase in convective PBL, with a large difference in winter and a small difference
~~in summer.~~

325 To compare the vertical distribution of CO₂ at BJ and XH, we calculate the CO₂ gradient ($\delta CO_2 = \frac{CO_{2,L1} - CO_{2,L2}}{Alt_{L2} - Alt_{L1}}$) (Figure 12c), The diurnal variations of δCO_2 at BJ and XH have a similar pattern: close-zero during the day and positive at night. The maximum δCO_2 can reach to 0.6 ppm/m at XH in 2018 August and 0.2 ppm/m at BJ in 2018 November. The larger height difference at BJ (120 m) as compared to XH (20 m) may contribute to the smaller δCO_2 .

3.7 Weekday-weekend variation

330 Figure 13 shows the ~~hourly means and stds of the deseasonalized average hourly means of~~ CO₂ ~~mole fractions at BJ, XH and~~
~~XL on weekdays and weekends. There is no clear difference in the~~ on weekday, weekend and all days at BJ (L1), XH (L1), XL
~~between October 2018 and September 2019, and BU (Boston) between April 2018 and April 2019. At BJ (L1), the nighttime~~
~~CO₂ measurements on weekend from 20 pm to 6 am next morning are generally ~5 ppm larger than those on weekday. XH~~
~~(L1) and XL CO₂ measurements on weekend are ~2 ppm than those on weekday throughout the whole day respectively. On~~
335 ~~the contrary, BU CO₂ mole fraction between weekdays and weekends. However, we find that the measurements on weekday~~
~~are ~8 ppm larger than those on weekend between 4 and 6 am. The CO₂ mole fractions have a minimum on Tuesday at~~
~~BJ and XH. It is indicated that the anthropogenic emission is lowest on Tuesday~~ differences on weekday and weekend at
BU turn smaller after sunrise. The mean CO₂ at BJ (L1), XH (L1), XL and BU is 447.6, 436.2, 420.3 and 429.8 ppm
~~respectively on weekday, and 449.2, 437.6, 421.4 and 427.5 ppm respectively on weekend. The weekday-weekend varia-~~
340 ~~tions at BJ and XH are different from similar to~~ that at Nanjing China (Gao et al., 2018), where CO₂ mole fractions are
higher on ~~weekends, or weekend, but different~~ from Boston USA, London UK and Tamil Nadu India, where the CO₂ mole
fractions are higher on ~~weekdays (Hernández-Paniagua et al., 2015; Kumar and Nagendra, 2015; Briber et al., 2013)~~ weekday
~~(Hernández-Paniagua et al., 2015; Kishore Kumar and Shiva Nagendra, 2015; Briber et al., 2013).~~

4 Conclusion

345 In this study, we show the CO₂ measurements from the in situ Picarro instruments at BJ, XH, and XL between June 2018 and March 2020. It is the first time to investigate CO₂ variations at these sites. BJ is inside the megacity, XH is in the suburban




Figure 12. (a): the CO₂ measurements of BJ L1 and BJ L2 between October 2018 and September 2019. The error bars are the hourly standard deviations of CO₂. (b): the CO₂ measurements of XH L1 and XH L2 between August 2018 and July 2019. (c): the hourly δ CO₂ [ppm/m] in each month at BJ and XH.

area, and XL is in the countryside on a mountain. The uncertainties of the CO₂ are 0.01, 0.06 and 0.02 ppm at BJ, XH and XL, respectively. The means and stds of CO₂ mole fractions are 448.4 ± 12.8 ppm, 436.0 ± 9.2 ppm and 420.6 ± 8.2 ppm at BJ (L1), XH (L1) and XL, respectively. The [CarbonTracker simulations at these three sites show fire, ocean and biogenic CO₂ are close to each other throughout the whole year, and the variation of the fossil fuel CO₂ at XL is much less than that at BJ/XH. The CO₂ measurements at XL are used to represent the background and we find that there is a good relationship between the CO₂ enhancements at BJ and XH. BJ and XH are affected by CO₂ emissions and transports simultaneously. Comparison with other urban sites in US shows that the CO₂ concentration is the largest at BJ.](#)

The variations of CO₂ at BJ, XH, and XL are discussed on diurnal and seasonal scales. It is found that the seasonal cycles of CO₂ at these three sites are similar, with a high value in winter and a low value in summer, which is closely related to air temperature and LAI. However, the amplitudes of seasonal variations are different, with the values of ~~60.3~~ [41.2](#) ppm, 36.1 ppm and 29.3 ppm at BJ, XH and XL, respectively. For the diurnal variation, the CO₂ is relatively low during the day and high at night. The diurnal variation of CO₂ at BJ, XH and XL is affected by the BLH, photosynthesis and human activities, and the impact of photosynthesis is more significant at XL.




Figure 13. The average hourly means (a) and standard deviations (b) of CO₂ on weekdays-weekday, weekend and weekends-all days at (a) BJ (L1), BJ-L2, (b) XH (L1), XH-L2 and (c) XL and (d) BU (Boston) between October 2018 and September 2019. The light gray shaded area represents one standard deviation from the mean for all days.

360 The CO₂ measurements are compared against the local wind data at BJ and XL. At BJ, high CO₂ mole fractions are observed with low wind speeds (< 2 m.s⁻¹). At XL, the high CO₂ mole fractions during daytime are observed with the wind coming from the south, where the urban area is located.

The two-levels measurements at BJ and XH show that the CO₂ mole fractions at lower and upper levels are close to each other during the day. The CO₂ mole fraction at the lower level is larger than that at the upper level at night with a vertical
 365 gradient up to 0.6 ppm/m at XH and 0.2 ppb/m at BJ. There is no clear difference in the-The CO₂ mole fraction-between weekdays-and-weekends-fractions on weekend at BJ, XH and XL are found slightly higher than the ones on weekday.

Author contributions. MZ, TW, PW and GW designed the experiment. YY performed the data curation. YY and MZ wrote the manuscript, and all authors read and provided comments on the paper.

Competing interests. The authors declare that they have no conflict of interest.

370 References

- Bagley, J., Rosenthal, D. M., Ruiz-Vera, U. M., Siebers, M. H., Kumar, P., Ort, D. R., and Bernacchi, C. J.: The influence of photosynthetic acclimation to rising CO₂ and warmer temperatures on leaf and canopy photosynthesis models, *Global Biogeochem. Cycles*, 29, 194–206, <https://doi.org/10.1002/2014GB004848>, 2015.
- Bakwin, P. S., Tans, P. P., Hurst, D. F., and Zhao, C.: Measurements of carbon dioxide on very tall towers: results of the NOAA/CMDL
375 program, *Tellus B*, 50, 401–415, <https://doi.org/10.1034/j.1600-0889.1998.t01-4-00001.x>, 1998.
- Bares, R., Mitchell, L., Fasoli, B., Bowling, D. R., Catharine, D., Garcia, M., Eng, B., Ehleringer, J., and Lin, J. C.: The Utah urban carbon dioxide (UUCON) and Uintah Basin greenhouse gas networks: instrumentation, data, and measurement uncertainty, *Earth Syst. Sci. Data.*, 11, 1291–1308, <https://doi.org/10.5194/essd-11-1291-2019>, 2019.
- Briber, B. M., Hutyra, L. R., Dunn, A. L., Raciti, S. M., and Munger, J. W.: Variations in Atmospheric CO₂ Mixing Ratios across a Boston,
380 MA Urban to Rural Gradient, *Land*, 2, 304–327, <https://doi.org/10.3390/land2030304>, 2013.
- Chan, D., Ishizawa, M., Higuchi, K., Maksyutov, S., and Chen, J.: Seasonal CO₂ rectifier effect and large-scale extratropical atmospheric transport, *J. Geophys. Res. Atmos.*, 113, <https://doi.org/10.1029/2007JD009443>, 2008.
- Cheng, X. L., Liu, X. M., Liu, Y. J., and Hu, F.: Characteristics of CO₂ Concentration and Flux in the Beijing Urban Area, *J. Geophys. Res. Atmos.*, 123, 1785–1801, <https://doi.org/10.1002/2017JD027409>, 2018.
- 385 Churkina, G.: The role of urbanization in the global carbon cycle., *Front. Ecol. Evol.*, 3, <https://doi.org/10.3389/fevo.2015.00144>., 2016.
- Crippa, M., Oreggioni, G., Guizzardi, D., Muntean, M., Schaaf, E., Vullo, E. L., Solazzo, E., Monforti-Ferrario, F., Olivier, J., and Vignati, E.: Fossil CO₂ and GHG emissions of all world countries - 2019 Report, Publications Office of the European Union, Luxembourg, pp. ISBN 978–92–76–11 100–9, <https://doi.org/10.2760/687800>, 2019.
- Culf, A., Fisch, G., Malhi, Y., and Nobre, C.: The influence of the atmospheric boundary layer on carbon dioxide concentrations over a
390 tropical forest, *Agric. For. Meteorol.*, 85, 149 – 158, [https://doi.org/10.1016/S0168-1923\(96\)02412-4](https://doi.org/10.1016/S0168-1923(96)02412-4), 1997.
- Denning, A. S., Takahashi, T., and Friedlingstein, P.: Can a strong atmospheric CO₂ rectifier effect be reconciled with a “reasonable” carbon budget?, *Tellus B*, 51, 249–253, <https://doi.org/10.1034/j.1600-0889.1999.t01-1-00010.x>, 1999.
- Dhokal, S.: Urban energy use and carbon emissions from cities in China and policy implications, *Energ. Policy.*, 37, 4208–4219, <https://doi.org/10.1016/j.enpol.2009.05.020>, 2009.
- 395 Fang, S. X., Zhou, L. X., Tans, P. P., Ciais, P., Steinbacher, M., Xu, L., and Luan, T.: In situ measurement of atmospheric CO₂ at the four WMO/GAW stations in China, *Atmos. Chem. Phys.*, 14, 2541–2554, <https://doi.org/10.5194/acp-14-2541-2014>, 2014.
- Feng, S., Lauvaux, T., Newman, S., Rao, P., Ahmadov, R., Deng, A., Díaz-Isaac, L. I., Duren, R. M., Fischer, M. L., Gerbig, C., Gurney, K. R., Huang, J., Jeong, S., Li, Z., Miller, C. E., O’Keeffe, D., Patarasuk, R., Sander, S. P., Song, Y., Wong, K. W., and Yung, Y. L.: Los Angeles megacity: a high-resolution land–atmosphere modelling system for urban CO₂ emissions, *Atmos. Chem. Phys.*, 16, 9019–9045,
400 <https://doi.org/10.5194/acp-16-9019-2016>, <https://acp.copernicus.org/articles/16/9019/2016/>, 2016.
- Feng, T., Zhou, W., Wu, S., Niu, Z., Cheng, P., Xiong, X., and Li, G.: High-resolution simulation of wintertime fossil fuel CO₂ in Beijing, China: Characteristics, sources, and regional transport, *Atmos. Environ.*, 198, 226–235, <https://doi.org/10.1016/j.atmosenv.2018.10.054>, 2019.
- Gao, Y. Q., Lee, X. H., Liu, S. D., Hu, N., Wei, X., Hu, C., Liu, C., Zhang, Z., and Yang, Y. C.: Spatio-temporal variability of the
405 near-surface CO₂ concentration across an industrial-urban-rural transect, Nanjing, China, *Sci. Total Environ.*, 631-632, 1192–1200, <https://doi.org/10.1016/j.scitotenv.2018.03.126>, 2018.

- George, K., Ziska, L., Bunce, J., and Quebedeaux, B.: Elevated atmospheric CO₂ concentration and temperature across an urban-rural transect, *Atmo. Environ.*, 41, 7654–7665, <https://doi.org/10.1016/j.atmosenv.2007.08.018>, 2007.
- Hansen, J., Sato, M., Kharecha, P., and von Schuckmann, K.: Earth’s energy imbalance and implications, *Atmos. Chem. Phys.*, 11, 13421–13449, <https://doi.org/10.5194/acp-11-13421-2011>, 2011.
- Hernández-Paniagua, I. Y., Lowry, D., Clemitshaw, K. C., Fisher, R. E., France, J. L., Lanoisellé, M., Ramonet, M., and Nisbet, E. G.: Diurnal, seasonal, and annual trends in atmospheric CO₂ at southwest London during 2000–2012: Wind sector analysis and comparison with Mace Head, Ireland, *Atmos. Environ.*, 105, 138 – 147, <https://doi.org/10.1016/j.atmosenv.2015.01.021>, 2015.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Idso, C., Idso, S., and Balling, R.: The urban CO₂ dome of Phoenix, Arizona, *Phys. Geogr.*, 19, 95–108, <https://doi.org/10.1080/02723646.1998.10642642>, 2013.
- IPCC: Climate change 2013: The physical science basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, <https://www.ipcc.ch/report/ar5/wg1/>, 2013.
- Kishore Kumar, M. and Shiva Nagendra, S.: Characteristics of ground level CO₂ concentrations over contrasting land uses in a tropical urban environment, *Atmos. Environ.*, 115, 286–294, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2015.05.044>, <https://www.sciencedirect.com/science/article/pii/S1352231015301151>, 2015.
- Kumar, M. K. and Nagendra, S. M. S.: Characteristics of ground level CO₂ concentrations over contrasting land uses in a tropical urban environment, *Atmos. Environ.*, 115, 286–294, <https://doi.org/10.1016/j.atmosenv.2015.05.044>, 2015.
- Lac, C., Donnelly, R. P., Masson, V., Pal, S., Riette, S., Donier, S., Queguiner, S., Tanguy, G., Ammoura, L., and Xueref-Remy, I.: CO₂ dispersion modelling over Paris region within the CO₂-MEGAPARIS project, *Atmos. Chem. Phys.*, 13, 4941–4961, <https://doi.org/10.5194/acp-13-4941-2013>, 2013.
- Lauvaux, T., Miles, N. L., Deng, A., Richardson, S. J., Cambaliza, M. O., Davis, K. J., Gaudet, B., Gurney, K. R., Huang, J., O’Keefe, D., Song, Y., Karion, A., Oda, T., Patarasuk, R., Razlivanov, I., Sarmiento, D., Shepson, P., Sweeney, C., Turnbull, J., and Wu, K.: High-resolution atmospheric inversion of urban CO₂ emissions during the dormant season of the Indianapolis Flux Experiment (INFLUX), *J. Geophys. Res. Atmos.*, 121, 5213–5236, <https://doi.org/10.1002/2015JD024473>, 2016.
- Li, Y., Deng, J., Mu, C., Xing, Z., and Du, K.: Vertical distribution of CO₂ in the atmospheric boundary layer: Characteristics and impact of meteorological variables, *Atmos. Environ.*, 91, 110 – 117, <https://doi.org/10.1016/j.atmosenv.2014.03.067>, 2014.
- Liu, H. Z., Feng, J. W., Järvi, L., and Vesala, T.: Four-year (2006–2009) eddy covariance measurements of CO₂ flux over an urban area in Beijing, *Atmos. Chem. Phys.*, 12, 7881–7892, <https://doi.org/10.5194/acp-12-7881-2012>, 2012.
- Ly, Z., Shi, Y., Zang, S., and Sun, L.: Spatial and Temporal Variations of Atmospheric CO₂ Concentration in China and Its Influencing Factors, *Atmos.*, 11, 231, <https://doi.org/10.3390/atmos11030231>, 2020.
- Matthäus, K., Annmarie, E., Dustin, R., Lin, J., Feng, S., Lei, R. X., Lauvaux, T., Oda, T., Roehl, C., Blavier, J. F., and Iraci, L.: Urban-focused satellite CO₂ observations from the Orbiting Carbon Observatory-3: A first look at the Los Angeles megacity, *Remote Sens. Environ.*, 258, 112314, <https://doi.org/10.1016/j.rse.2021.112314>, 2021.

- 445 McKain, K., Down, A., Raciti, S. M., Budney, J., Hutyra, L. R., Floerchinger, C., Herndon, S. C., Nehrkorn, T., Zahniser, M. S., Jackson, R. B., Phillips, N., and Wofsy, S. C.: Methane emissions from natural gas infrastructure and use in the urban region of Boston, Massachusetts, *Proc. Natl. Acad. Sci. U. S. A.*, 112, 1941–1946, <https://doi.org/10.1073/pnas.1416261112>, 2015.
- Mitchell, L. E., Lin, J. C., Bowling, D. R., Pataki, D. E., Strong, C., Schauer, A. J., Bares, R., Bush, S. E., Stephens, B. B., Mendoza, D., Mallia, D., Holland, L., Gurney, K. R., and Ehleringer, J. R.: Long-term urban carbon dioxide observations reveal spatial and temporal dynamics related to urban characteristics and growth, *Proc. Natl. Acad. Sci. U. S. A.*, 115, 2912–2917, 450 <https://doi.org/10.1073/pnas.1702393115>, 2018.
- Mohotti, A. J. and Lawlor, D. W.: Diurnal variation of photosynthesis and photoinhibition in tea: effects of irradiance and nitrogen supply during growth in the field, *J. Exp. Bot. Copiar*, pp. 313–322, 2002.
- Mustafa, F., Bu, L. B., Wang, Q., Ali, M. A., Bilal, M., Shahzaman, M., and Qiu, Z. F.: Multi-Year Comparison of CO₂ Concentration from NOAA Carbon Tracker Reanalysis Model with Data from GOSAT and OCO-2 over Asia, *Rem. Sens.*, 12, 2498, 455 <https://doi.org/10.3390/rs12152498>, 2020.
- Nevison, C. D., Mahowald, N. M., Doney, S. C., Lima, I. D., Werf, G. R. V. D., Randerson, J. T., Baker, D. F., And, P. K., and Mckinley, G. A.: Contribution of ocean, fossil fuel, land biosphere, and biomass burning carbon fluxes to seasonal and interannual variability in atmospheric CO₂, *J. Geophys. Res. Biogeo.*, 113, <https://doi.org/10.1029/2007JG000408>, 2008.
- Newman, S., Jeong, S., Fischer, M. L., Xu, X., Haman, C. L., Lefer, B., Alvarez, S., Rappenglueck, B., Kort, E. A., and Andrews, A. E.: 460 Diurnal tracking of anthropogenic CO₂ emissions in the Los Angeles basin megacity during spring 2010, *Atmos. Chem. Phys.*, 13, 4359–4372, 2013.
- ángeles García, M., Sánchez, M. L., and Pérez, I. A.: Differences between carbon dioxide levels over suburban and rural sites in Northern Spain, *Environ. Sci. Pollut. Res.*, 19, 432–439, <https://doi.org/10.1007/s11356-011-0575-4>, 2012.
- Peters, W., Miller, J. B., Whitaker, J., Denning, A. S., Hirsch, A., Krol, M. C., Zupanski, D., Bruhwiler, L., and Tans, P. P.: An ensemble 465 data assimilation system to estimate CO₂ surface fluxes from atmospheric trace gas observations, *J. Geophys. Res. Atmos.*, 110, <https://doi.org/10.1029/2005JD006157>, 2005.
- Richardson, S., Miles, N., Davis, K. J., Lauvaux, T., Martins, D. K., Turnbull, J. C., McKain, K., Sweeney, C., and L., M. O.: Tower measurement network of in-situ CO₂, CH₄, and CO in support of the Indianapolis FLUX (INFLUX) Experiment., *Elementa: Science of the Anthropocene*, 5, 59, <https://doi.org/10.1525/elementa.140>, 2017.
- 470 Sargent, M., Barrera, Y., Nehrkorn, T., Hutyra, L. R., Gately, C. K., Jones, T., McKain, K., Sweeney, C., Hegarty, J., Hardiman, B., Wang, J. A., and Wofsy, S. C.: Anthropogenic and biogenic CO₂ fluxes in the Boston urban region, *Proc. Natl. Acad. Sci. U. S. A.*, 115, 7491–7496, <https://doi.org/10.1073/pnas.1803715115>, 2018.
- Shi, Y., Xi, Z., Simayi, M., Li, J., and Xie, S.: Scattered coal is the largest source of ambient volatile organic compounds during the heating season in Beijing, *Atmos. Chem. Phys.*, 20, 9351–9369, <https://doi.org/10.5194/acp-20-9351-2020>, 2020.
- 475 Shusterman, A. A., Teige, V. E., Turner, A. J., Newman, C., Kim, J., and Cohen, R. C.: The BERkeley Atmospheric CO₂ Observation Network: initial evaluation, *Atmos. Chem. Phys.*, 16, 13 449–13 463, <https://doi.org/10.5194/acp-16-13449-2016>, 2016.
- Soegaard, H. and Møller-Jensen, L.: Towards a spatial CO₂ budget of a metropolitan region based on textural image classification and flux measurements, *Remote Sens. Environ.*, 87, 283–294, [https://doi.org/10.1016/S0034-4257\(03\)00185-8](https://doi.org/10.1016/S0034-4257(03)00185-8), 2003.
- Song, T., Wang, Y., and Sun, Y.: Estimation of carbon dioxide flux and source partitioning over Beijing, China, *J. Environ. Sci.*, 25, 2429– 480 2434, [https://doi.org/10.1016/S1001-0742\(12\)60336-2](https://doi.org/10.1016/S1001-0742(12)60336-2), 2013.

- Turnbull, J. C., Sweeney, C., Karion, A., Newberger, T., Lehman, S. J., Tans, P. P., Davis, K. J., Lauvaux, T., Miles, N. L., Richardson, S. J., Cambaliza, M. O., Shepson, P. B., Gurney, K., Patarasuk, R., and Razlivanov, I.: Toward quantification and source sector identification of fossil fuel CO₂ emissions from an urban area: Results from the INFLUX experiment, *J. Geophys. Res. Atmos.*, 120, 292–312, <https://doi.org/10.1002/2014JD022555>, 2015.
- 485 Verhulst, K. R., Karion, A., Kim, J., Salameh, P. K., Keeling, R. F., Newman, S., Miller, J., Sloop, C., Pongetti, T., Rao, P., Wong, C., Hopkins, F. M., Yadav, V., Weiss, R. F., Duren, R. M., and Miller, C. E.: Carbon dioxide and methane measurements from the Los Angeles Megacity Carbon Project – Part 1: calibration, urban enhancements, and uncertainty estimates, *Atmos. Chem. Phys.*, 17, 8313–8341, <https://doi.org/10.5194/acp-17-8313-2017>, 2017.
- Wu, Y., Wang, R., Zhou, Y., Lin, B., Fu, L., He, K., and Hao, J.: On-Road Vehicle Emission Control in Beijing: Past, Present, and Future, 490 *Environ. Sci. Technol.*, 45, 147–153, <https://doi.org/10.1021/es1014289>, 2011.
- Yang, Y., Wang, T., Wang, P. C., Zhou, M. Q., and Yao, B.: In-situ measurement of CO₂ at the Xinglong regional background station over North China, *Atmospheric and Oceanic Science Letters*, 12, 385–391, <https://doi.org/10.1080/16742834.2019.1644949>, 2019.
- Yang, Y., Zhou, M., Langerock, B., Sha, M. K., Hermans, C., Wang, T., Ji, D., Vigouroux, C., Kumps, N., Wang, G., De Maz-
495 ière, M., and Wang, P.: New ground-based Fourier-transform near-infrared solar absorption measurements of XCO₂, XCH₄ and XCO at Xianghe, China, *Earth Syst. Sci. Data.*, 12, 1679–1696, <https://doi.org/10.5194/essd-12-1679-2020>, 2020.