

Secondary organic aerosol phase behaviour in chamber photo-oxidation of mixed precursors

Yu Wang¹, Aristeidis Voliotis¹, Yunqi Shao¹, Taomou Zong², Xiangxinyue Meng², Mao Du¹, Dawei Hu¹, Ying Chen^{3,#}, Zhijun Wu^{2,4,5}, M. Rami Alfarra^{1,6}, Gordon McFiggans^{1,*}

5 ¹Centre for Atmospheric Science, Department of Earth and Environmental Sciences, The University of Manchester, Manchester M13 9PL, UK

²State Key Joint Laboratory of Environmental Simulation and Pollution Control, International Joint Laboratory for Regional Pollution Control, Ministry of Education (IJRC), College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

10 ³Lancaster Environment Centre, Lancaster University, LA1 4YQ, UK

⁴International Joint Laboratory for Regional Pollution Control, 52425 Jülich, Germany, and Beijing 100871, China

⁵Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Nanjing University of Information Science and Technology, Nanjing 210044, China

15 ⁶National Centre for Atmospheric Science, School of Earth and Environmental Sciences, The University of Manchester, Manchester, M13 9PL, UK

Currently at Exeter Climate Systems, University of Exeter, Exeter, EX4 4QE, UK

*Correspondence to: Gordon McFiggans (g.mcfiggans@manchester.ac.uk)

Abstract.

The phase behaviour of aerosol particles plays a profound role in atmospheric physicochemical processes, influencing their physical and optical properties and further impacting climate and air quality. However, understanding of aerosol phase behaviour is still incomplete, especially that of multicomponent particles
25 which contain inorganic compounds and secondary organic aerosol (SOA) from mixed volatile organic compound (VOC) precursors. We report measurements conducted in the Manchester Aerosol Chamber (MAC) to investigate the aerosol rebounding tendency, measured as “bounce fraction”, as a surrogate of particle phase behaviour during SOA formation from photo-oxidation of biogenic (α -pinene, isoprene) and ~~anthropogenic-anthropogenic~~ (*o*-cresol) VOCs and their binary mixtures on deliquescent ammonium
30 sulphate seed.

Aerosol phase behaviour is RH and chemical composition dependent (key factors determining aerosol water uptake). Liquid (bounce fraction, $BF < 0.2$) at $RH > 80\%$ and non-liquid behaviour ($BF > 0.8$) at $RH < 30\%$ were observed, with a liquid-to-nonliquid transition with decreasing RH between $30\% \sim 80\%$. This RH-dependent phase behaviour ($RH_{BF=0.2, 0.5, 0.8}$) increased towards a maximum with
35 increasing organic-inorganic-mass ratio ($MR_{org/inorg}$) during SOA formation evolution in all investigated VOC systems. With the use of comparable initial ammonium sulphate seed concentration, the SOA production rate of the VOC systems determines the $MR_{org/inorg}$, and consequently the change of the phase behaviour. Although less important than RH and $MR_{org/inorg}$, the SOA composition plays a second-order role, with differences in liquid-to-nonliquid transition at moderate $MR_{org/inorg}$ of ~ 1 observed between
40 biogenic-only (~~anthropogenic-free~~) and ~~anthropogenic-anthropogenic~~-containing VOC systems. Considering the combining role of the RH and chemical composition in aerosol phase behaviour, the BF decreased monotonically with increasing hygroscopic growth factor (GF) and the BF was ~ 0 when GF was larger than 1.15. The real atmospheric consequences of our results are that any processes changing ambient RH or $MR_{org/inorg}$ (aerosol liquid water) will influence their particle phase behaviour. Where
45 abundant anthropogenic VOCs contribute to SOA, compositional changes of SOA may influence phase behaviour at moderate organic mass fraction ($\sim 50\%$) compared with purely biogenic SOA. Further studies are needed on more complex and realistic atmospheric mixtures.

50 **1 Introduction**

Aerosol particles are ubiquitous in the atmosphere and can act as reaction vessels where physicochemical processes occur. As one of the key physical properties of aerosol particles, the aerosol phase behaviour can significantly impact those physicochemical processes (Martin, 2000). Following the pioneering work
55 of Virtanen et al. (2010), there has been considerable recent efforts to resolve the influences of particle phase behaviour from a number of perspectives mainly relating to retardation of diffusion or mobile components, including water. The viscous solid particles have potential impacts on physicochemical processes, such as constraining gas-particle partitioning of semi-volatile organic species (Vaden et al., 2011; Shiraiwa et al., 2011; Shiraiwa and Seinfeld, 2012; Zaveri et al., 2014; Renbaum-Wolff et al., 2013),
60 heterogeneous reactions or liquid phase reactions (Shiraiwa et al., 2011; Koop et al., 2011; Kuwata and Martin, 2012; Zhang et al., 2018; Martin, 2000). These processes affect secondary organic and inorganic particulate matters formation in the atmosphere, further impacting their optical properties and air quality. Moreover, the ice nucleation abilities in the upper tropospheric conditions and cloud condensation nuclei activation of aerosol particles are affected by the phase behaviour (Pöschl, 2011; Murray et al.,
65 2010; Murray, 2008; Reid et al., 2018; Shiraiwa et al., 2017; Ignatius et al., 2016; Slade et al., 2017), further impacting cloud formation and regional climate. ~~A~~Better understanding the phase behaviour of atmospheric particles is important for understanding physicochemical processes in the atmosphere, and aerosol-cloud interactions.

The particle rebounding property has been widely used to study phase behaviour of aerosol particles
70 (Dahneke, 1971; Stein et al., 1994). In a real atmosphere, the phase behaviour of aerosol particles varied significantly under various environments, depending on the ambient relative humidity (RH) and aerosol chemical composition. For example, background atmospheric particles in the tropical rainforest over central Amazonia, mainly composed of isoprene-derived secondary organic aerosol (SOA), were initially

liquid for ambient RH > 80% and temperature of 23--27 °C (Bateman et al., 2015b). In contrast, when
75 the measurement site was influenced by anthropogenic air mass from urban pollution and biomass
burning, non-liquid PM fraction increased to 60% at 95% RH (Bateman et al., 2017). However, in the
boreal forest of northern Finland, atmospheric particles (mainly monoterpene-derived SOA) showed
amorphous, solid-like phase state (Virtanen et al., 2010). An enhanced fraction of particulate sulphate can
lead to loss of particle bounce and atmospheric particles with high fraction of inorganic compounds
80 showed a liquid-like phase state under moderate and high ambient RH (e.g. urban area (Liu et al., 2017),
subtropical coastal megacity (Liu et al., 2019), south-eastern US rural site (Pajunoja et al., 2016), North-
eastern near forest area (Slade et al., 2019)). When the terpene-dominant SOA increased during night
time (Slade et al., 2019) or ambient RH dropped under 60% (Liu et al., 2017), the non-liquid PM
increased.

85 Consistent with the main findings in the field studies, the particle bounce of pure SOA from the oxidation
of representative biogenic or anthropogenic VOC (e.g. isoprene, α -pinene, toluene) decreased with
elevated RH and varied with SOA composition (Bateman et al., 2015a; Saukko et al., 2012). The
discrepancy between the Bateman et al. (2015a) and Saukko et al. (2012) was the RH at which aerosol
particles fully adhered to the substrate (with < 20% aerosol particles rebounding; referred as the adhesion
90 RH). In a well-calibrated rebounding impactor system, the rebound or adhering behaviour is highly related
to the aerosol phase state, which is determined by material softening. Bateman et al. (2015a) found the
complete adhesion RH for the isoprene SOA, α -pinene SOA, 2:1 isoprene/ α -pinene mixture and toluene
SOA were > 65%, > 95%, > 80% and > 80%. However, Saukko et al. (2012) observed that SOA from
isoprene and α -pinene oxidation, do not fully adhere to the substrate even RH of up to 90%. This
95 discrepancy might due to the slightly different instrumentation design (pressure drop in Saukko et al.
(2012) vs atmospheric pressure in Bateman et al. (2015a)) or SOA composition difference from different
oxidation conditions. As observed in the real atmosphere, aerosol particles are usually a mixture with
organic and inorganic species (e.g. sulphate and nitrate) (Jimenez et al., 2009). Since the inorganic species
have lower glass transition temperature than atmospheric-relevant organics (Pedernera, 2008; Koop et al.,
100 2011), the presence of inorganic species could theoretically lower the glass transition temperature of the
estimated SOA compounds mixtures. Furthermore, inorganic species are hydrophilic and the absorbed

water molecules can act as a plasticiser, which effectively lower the glass transition temperature and soften the aerosol (Koop et al., 2011; Martin, 2000). Therefore, ignoring the mixing with inorganic species, or assuming external mixing as in Shiraiwa et al. (2017), could bias our understanding on the phase behaviour of aerosols containing abundant inorganic compounds.

There are few studies on phase behaviour of multicomponent aerosol particles. Saukko et al. (2012) found increasing sulphate fraction mixing with SOA produced by longifolene oxidation can reduce the particle rebounding significantly. Saukko et al. (2015) extended this to study deliquescence hysteresis of ammonium sulphate with condensed SOA from α -pinene and isoprene oxidation as manifested by particle rebound. They found the α -pinene SOA condensing on ammonium sulphate seed particles showed no influences on their deliquescence but significantly attenuated the efflorescence behaviour. In contrast, the isoprene SOA system resulted in a loss of sharp deliquescence and efflorescence behaviour in comparison to pure ammonium sulphate (Saukko et al., 2015). These results are partly consistent with similar studies using a different instrument (Hygroscopicity Tandem Differential Mobility Analyzer) (Smith et al., 2011; Smith et al., 2012). They found that the α -pinene SOA on ammonium sulphate seed can slightly shift the deliquescence and efflorescence RH by a few percent (Smith et al., 2011) while the isoprene SOA can significantly decrease the deliquescence and efflorescence RH depending on the organic fraction (Smith et al., 2012). These contrasting findings above indicate that aerosol phase behaviour could differ between different SOA precursor systems as well as in the presence of inorganic compounds. However, it is still unclear how the phase behaviour (and any associated diffusion limiting behaviour) will be influenced during the formation and evolution of SOA from an increased complexity of mixed precursors in the presence of inorganic seed.

We designed a series of aerosol simulation chamber experiments to study SOA formed from representative biogenic (α -pinene, isoprene) and ~~anthropogenic~~ anthropogenic (*o*-cresol) VOC photochemistry. The experiments studied single VOC precursors and their binary mixtures under modest-NO_x condition on deliquescent ammonium sulphate seed particles. We frame our results around the following hypotheses:

1) Aerosol phase behaviour for SOA mixture is driven by RH and organic-inorganic-mass ratio

2) The difference in SOA composition is less important in determining the phase behaviour than the SOA

130 production rate, which changes the organic-inorganic ratio.

The main objective of this paper is to test the above two hypotheses and discuss their potential atmospheric implications.

2 Measurements and Methods

2.1 Reaction chamber and experimental setup

135 The aerosol particles for the experiments were produced in the Manchester Aerosol Chamber (MAC).

Detailed description of MAC can be found in Shao et al. (2021) and a brief introduction is given as below.

The facility is run as a batch reactor with an 18 m³ (3_m (H) * 3_m (L) * 2_m (W)) FEP Teflon bag supported by three aluminium frames, in which the upper and the lower frame can move freely with the expansion or collapsing as sampling air is introduced to or extracted from the chamber. The Teflon bag

140 is enclosed inside a housing with temperature and relative humidity controlled by an air conditioning

system. Two 6_kW Xenon arc lamps (XBO 6000 W/HSLA OFR, Osram) and a series of halogen bulbs arranged in 7 rows containing 16 bulb each (Solux 50W/4700K, Solux MR16, USA) are mounted inside of the enclosure housing. The combination of 5 rows of halogen bulbs (2 rows spare) and two Xenon arc

lamps is chosen to be a good representative to mimic the solar spectrum in the wavelength of 290_--_800

145 nm (Alfarra et al., 2012). The calculated photolysis rate of NO₂ (J_{NO_2}) from O₃-NO-NO₂ photostationary

state was $(1.8_--_3) \times 10^{-3} \text{ s}^{-1}$ during the experiment period. The simulated irradiation spectrum intensity

in our chamber is around a third of comparable to the measured solar spectrum at the mid-day on a clear sky day in June of Manchester (<https://www.eurochamp.org/Facilities/SimulationChambers/MAC-MICC.aspx>).

150 To ensure chamber cleanliness and data reproducibility, an automatic fill/flush cycle is conducted pre-

and post-experiment using 3 m³ min⁻¹ purified clean air. 5_~_6 cycles are routinely conducted between experiments, with total number concentration of aerosol particles usually lower than 10 p_/cce⁻¹ after

cleaning. Air is scrubbed using a series of filters, including Purafil (Purafil Inc., USA) and charcoal removing reactive gaseous compounds and HEPA (Donaldson Filtration, USA) removing particles and a dryer. To remove reactive compounds, the chamber is soaked in high concentrations of O₃ (~1 ppm) overnight between experiments which is removed during pre-experiment fill/flush cycles on the subsequent day. An additional harsh cleaning procedure is conducted weekly with high O₃ (~1 ppm) and UV for 4~5 hours to consume the remaining reactive compounds.

Seed particles, VOC, NO_x and water vapour are injected into chamber before illumination. Deliquescent ammonium sulphate (AS) seed (Puratronic, 99.999% purity) are nebulized and introduced into a drum to mix before flushing into chamber. Liquid VOC precursors (α -pinene, isoprene, *o*-cresol; Sigma Aldrich, GC grade \geq 99.99% purity) are injected using a syringe into a heated glass bulb in which the VOCs are instantaneously vaporized. The vaporized VOCs are flushed into the chamber with a flow of ~0.5 bar high purity nitrogen (ECD grade, 99.997%). Here, the experiments will investigate single and ~~iso-reactive~~ binary mixtures under modest-NO_x condition (VOC/NO_x of ~~5-3~~ ~10). NO_x (mostly as NO₂ in this study) is introduced through a cylinder with a flow of ~0.3 bar high purity nitrogen (ECD grade, 99.997%) and the O₃-NO-NO₂ photostationary state will establish immediately after illumination. The detailed initial conditions for chamber setup are shown in Table 1 ~~and the experimental design will be explained~~ in details in Sec. 2.3.

Table 1. Summary of the initial conditions of chamber experiments.

Exp. Date	VOC type	[VOC] ₀ (ppbV)	VOC/NO _x	T (°C)	RH (%)	AS Seed conc. ($\mu\text{g m}^{-3}$) ^a	O:C ratio
28-Mar-2019	α -pinene	309	7.7	26.7	50.5	72.6	0.36 \pm 0.03
17-Apr-2019	α -pinene	155	4.4	25.9	55.0	47.8	0.45 \pm 0.04
02-Apr-2019	Isoprene	164	6.8	27.2	47.3	64.1	n.a.
12-Apr-2019	<i>o</i> -cresol	400	n.a.	27.3	53.3	47.8	0.64 \pm 0.03
19-Apr-2019	<i>o</i> -cresol	200	5.0	26.9	51.3	51.3	0.66 \pm 0.05
08-Apr-2019	α -pinene/isoprene	237 (155/82)	9.9	27.0	48.4	62.0	0.43 \pm 0.05

23-Apr-2019	α -pinene/ <i>o</i> -cresol	355 (155/200)	n.a.	25.6	55.8	42.5	0.48 ± 0.05
18-Apr-2019	<i>o</i> -cresol/isoprene	282 (200/82)	8.3	27.1	52.7	49.6	0.69 ± 0.05

^a calculated mass concentration from volume concentration from DMPS with a density of 1.77 g cm^{-3} .

2.2 Instrumentation

175 A series of instruments are equipped for gas-phase and particle-phase measurements in MAC. NO, NO₂ and NO_x are recorded by NO-NO₂-NO_x analyser (Model 42i, Thermo, USA) and O₃ is measured by O₃ analyser (Model 49C, Thermo, USA). RH and T inside of MAC are recorded by an Edgetech dewpoint hygrometer (DM-C1-DS2-MH-13, USA) and two Sensirion SHT75 sensors (Farnell 413-0698, USA). The aerosol number size distribution (20 ~ 550 nm) is measured by a Differential Mobility Particle Sizer (DMPS). Hygroscopic growth factor (GF) at 90% RH of submicron aerosol particles (75 ~ 250 nm) was recorded by a custom-made Hygroscopicity Tandem Differential Mobility Analyser (HTDMA) (Good et al., 2010), which can be used to calculate the GF at given RH using κ -Köhler approximation (Petters and Kreidenweis, 2007).

185 The particle bounce behaviour (bounce fraction, BF) was measured by a three-arm particle rebound apparatus with RH adjustment system. A brief instrumental description is provided below and more details can be found in Bateman et al. (2014) and Liu et al. (2017) along with a schematic diagram in Figure S1 of the latter. Three single-stage impactors operated in parallel in the system combined with a condensation particle counter (CPC, Model: 3772, TSI, USA). The first impactor is not equipped with a plate (step 1 in Figure S1 in Liu et al. (2017)), so particles can pass through the first impactor directly, measuring the total particle population (N₁). The second impactor is equipped with a smooth plate (step 190 2 in Figure S1 in Liu et al. (2017)), which provides a solid surface and allows particles rebounding from the impactor. The particle population measured after the second impactor represents the sum of particles that do not strike the impactor and that strike but rebound from the impactor (N₂). The third impactor is equipped with a grease-coated plate (step 3 in Figure S1 in Liu et al. (2017)). The coated grease is quite sticky and all particles striking on the plate will be stuck. Therefore, the particle number population after 195 grease-coated plate provides a measure of the particles that don't strike the impactor (N₃). The rebound fraction BF is defined in equation [1]. During the experiments, a Differential Mobility Analyzer (DMA) was used to select a monodisperse aerosol particles from chamber, with mobility diameter of 100 ~ 200 nm following the growth of the aerosol particles. The selected particle sizes are larger than the 50 %

transmission diameter of the used impactor (84.9 ± 5.4 nm) (Bateman et al., 2014) to ensure a reliable bounce fraction measurement.

$$BF = \frac{N_2 - N_3}{N_1 - N_3} \quad [1]$$

BF can be used as a proxy of aerosol phase behaviour, with limited capability representing semi-solid or solid particles over 100 Pa s in viscosity (Bateman et al., 2015a; Reid et al., 2018). Nevertheless, it provides insights into the transition process between liquid and solid/semi-solid phase, referred to as liquid-to-nonliquid transition below.

The chemical composition of the non-refractory PM₁ components (NH₄⁺, NO₃⁻, SO₄²⁻, and SOA) was recorded by a High-Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS, Aerodyne Research Inc., USA). A detailed introduction of the instrument, calibration procedures and quantification of the aerosol concentrations were described previously (DeCarlo et al., 2006; Jayne et al., 2000; Allan et al., 2003; Allan et al., 2004). The instrument was operated in ‘V mode’, and recorded with a time resolution of 1 min (30 s mass spectrum (MS) + 30 s particle-time-of-flight (PToF)). The elemental ratio of O:C used for the proxy of organic oxidation state and the ions of NO⁺ and NO₂⁺ used for the organic nitrate fraction calculation were derived from the high-resolution fitting on V-mode data.

Calibrations were performed before and after the campaign using monodisperse (350 nm) ammonium nitrate and ammonium sulphate particles following the standard procedure in Jayne et al. (2000) and Jimenez et al. (2003). An averaged ionization efficiency of ammonium nitrate was $9.38 \times 10^{8-8}$ ions molecule⁻¹ from the two calibrations. According to the ion balance of ammonium nitrate and ammonium sulphate in the calibrations, the specific relative ionization efficiencies (RIE) for NH₄⁺ and SO₄²⁻ are determined as 3.57 ± 0.02 and 1.28 ± 0.01 , respectively. The RIE of all organic compounds used the default value of 1.4 (Alfarra et al., 2004). In this study, real-time collection efficiency was applied to the data analysis by comparing NR-PM₁ mass with DMPS mass. In this study, the AMS mass concentrations were corrected by comparing to the real-time DMPS unit mass multiplied by concurrent calculated density from AMS species. The mass ratio of AMS/(DMPS*density) is 0.4 ~1.0 in all investigated VOC systems.

2.3 Rationale behind the choice of precursor

225 The real atmosphere comprises a complex mixture of VOCs with various reactivities, many of which may act as SOA precursors with varying degrees of efficiency. Recent chamber studies have started using relatively simple VOC mixtures to investigate complex interactions in their ability to form SOA in the presence of inorganic seed particles (McFiggans et al., 2019; Shilling et al., 2019). Extending these previous studies, a project was designed aiming at characterising chemical mechanisms, yield, and

230 physicochemical properties (volatility, hygroscopicity, CCN activity, and phase behaviour) of SOA formed from initially iso-reactive biogenic/anthropogenic VOC photochemistry on ammonium sulphate seed and exploring potential implications to the real atmosphere. Building on the McFiggans et al. (2019), we added an representative anthropogenic VOC (*o*-cresol) into initially designed binary mixtures of biogenic VOCs (isoprene & α -pinene). *o*-cresol can be emitted into atmosphere both directly

235 from biomass burning (Coggon et al., 2019; Koss et al., 2018) and indirectly from the oxidation of toluene (e.g. motor vehicles, solvent use (Fishbein, 1985)). Therefore, anthropogenic source is one of the main contributors to the *o*-cresol but worth noting that the natural biomass burning can also be an important contributor. *o*-cresol is chosen by virtue of its comparable $\cdot\text{OH}$ reactivity ~~comparable~~ with the chosen biogenic VOCs (Coeur-Tourneur et al., 2006), which enables comparable initial concentration to have an

240 equal reactivity with $\cdot\text{OH}$ at the beginning of experiment (referred as iso-reactive). The modest SOA yield of *o*-cresol (Henry et al., 2008) gives a good contrast with the low-yield isoprene and high-yield α -pinene. The overall experiment design enables us to explore SOA formation in initially iso-reactive single, binary and ternary VOC mixtures oxidation, details can be found in Voliotis et al. (to be submitted to this issue). This manuscript focuses on aerosol phase behaviour of seeded SOA from iso-reactive single and

245 binary VOCs photochemistry. Unfortunately, the bounce impactor was unavailable for the ternary mixture experiment.

3 Results and Discussion

3.1 BF dependence on RH and organic-inorganic-mass ratio

Figure 1 shows the rebound curves of the 100 ~ 200 nm multicomponent aerosol particles formed in various VOC systems for the period when organic mass fraction in NR-PM₁ is larger than 0.05. For all investigated VOC systems, the aerosol particles exhibited BF > 0.8 at RH < 30% and BF < 0.2 at RH > 80% at room temperature (18 °C). Between 30% and 80% RH, the BF monotonically decreased with the increasing RH, indicating a gradual transition in BF (usually within 15 ~ 25% RH width for BF declining from > 0.8 to < 0.2). Assuming the aerosol particles to be non-liquid if their BF > 0.8 and liquid if BF < 0.2 (Bateman et al., 2015a; Liu et al., 2017), this implies a gradual transition with RH in all investigated VOC systems in contrast to the rapid dissolution corresponding to deliquescence of inorganic salt particles (Tang and Munkelwitz, 1993; Kreidenweis and Asa-Awuku, 2014). Owing to limitations of the technique in differentiating particle viscosity at high values (Bateman et al., 2014), the non-liquid phase could represent semi-solid or solid phase. In addition, the rebound curves varied along with SOA formation and subsequent evolution as the photochemistry continues in all the investigated VOC systems (as shown in Figure 1). To illustrate the influences of chemical composition, the overview of rebound curves as a function of organic-inorganic-mass ratio ($MR_{org/inorg}$) in all VOC systems is shown in Figure S1, indicating the potential important role of $MR_{org/inorg}$ (and maybe OA composition) in determining the phase behaviour as RH.

To clearer describe the phase behaviour during SOA formation evolution, the $RH_{BF=0.2}$, $RH_{BF=0.5}$, and $RH_{BF=0.8}$ are determined representing the RH at which the BF is close to 0.2 (± 0.15), 0.5 (± 0.15), and 0.8 (± 0.15). It is worth noting that the determination of $RH_{BF=0.2, 0.5, 0.8}$ carry a maximum error of $\pm 5\%$ owing to the measurement resolution and resultant number of data points. By tracking the variation of $RH_{BF=0.2, 0.5, 0.8}$ during SOA formation evolution, we can understand the tendency of phase behaviour change and insight into the liquid-to-nonliquid transition at the variation of RH at which BF changes from 0.8 to 0.2 among VOC systems. As different VOCs have different reactivity with oxidant and yield, the formed SOA mass ~~from different VOCs~~ after six-hour photo-oxidation, and consequently the $MR_{org/inorg}$, are-were different among VOC systems. For *o*-cresol/isoprene, 50% reactivity *o*-cresol, *o*-cresol, α -

pinene/isoprene, α -pinene/*o*-cresol, 50% reactivity α -pinene and α -pinene systems, the $MR_{org/inorg}$ reached
275 up to 0.53 ± 0.02 , 1.27 ± 0.05 , 2.47 ± 0.06 , 3.77 ± 0.11 , 3.77 ± 0.01 , 4.45 ± 0.04 , and 7.37 ± 0.06 for the
last rebound curve at the end of experiments, respectively, ~~for the last rebound curve at the end of~~
~~experiments~~. As shown in Figure 2, the $RH_{BF=0.2, 0.5, 0.8}$ increased towards a maximum as an increase of
the $MR_{org/inorg}$ in all investigated VOC systems, indicating an increase of rebound tendency during SOA
evolution on deliquescent sulphate seed. Figure 2_a-c ~~shows~~ showed a co-increasing trend between
280 $RH_{BF=0.2, 0.5, 0.8}$ and $MR_{org/inorg}$ as $MR_{org/inorg} < 2$, thereafter $RH_{BF=0.2, 0.5, 0.8}$ ~~remains~~ remained constant at 70
~ 75%, 65 ~ 70% and 50 ~ 65%, respectively, in all ~~investigated~~ VOC systems. Herein, the reason for
the decrease of $RH_{BF=0.5}$ from 60% to 50% at $MR_{org/inorg} \sim 1$ in the *o*-cresol system ~~is~~ was not known and
the same behaviour was not observed for $RH_{BF=0.2, 0.8}$.

As expected, the change in $MR_{org/inorg}$ during SOA formation and evolution differed in various VOC
285 systems (as shown in Figure 3), depending on their SOA production rate (as shown in Figure S2), noting
the comparable initial sulphate seed concentrations. The order of the SOA production rate (from low to
high) in all the investigated VOC systems ~~is~~ was ~~are~~, *o*-cresol/isoprene \leq , 50% reactivity *o*-cresol, \leq *o*-
cresol \leq , α -pinene/isoprene \leq , α -pinene/*o*-cresol, \leq 50% reactivity α -pinene \leq and α -pinene (note that
insufficient mass was generated from isoprene ~~in the presence of a neutral ammonium sulphate~~
290 ~~seed system~~). It is worth noting that the SOA production rate in α -pinene/isoprene, α -pinene/*o*-cresol and
50% reactivity α -pinene systems are very similar, and their order is determined by the minor difference
of the slopes in Figure S2. Clearly, a higher SOA production rate ~~increases~~ increased the $MR_{org/inorg}$ faster
and consequently the phase behaviour change. As shown in Figure 4, $RH_{BF=0.2, 0.5, 0.8}$ showed an increasing
trend towards a maximum across the various VOC systems, highly related to how fast the SOA ~~are~~ were
295 formed. With an increasing SOA formation rate from the lowest *o*-cresol/isoprene to highest α -pinene
system, a shorter time was taken for $RH_{BF=0.2, 0.5, 0.8}$ to reach the maximum. Take $RH_{BF=0.5}$ as an example,
it can be seen from Figure 1 that the $RH_{BF=0.5}$ at the beginning of the photochemistry ~~is~~ was lower than
40% for all investigated VOC systems. ~~Panel Figure 4 ba)~~ ~~shows~~ showed that, for the *o*-cresol/isoprene
system with the lowest SOA formation rate, it took ~ 3.5 h for $RH_{BF=0.5}$ to increase to 50% and ~ 5 h to
300 approach 60%. For comparison, for α -pinene/isoprene (the moderate) and α -pinene (the highest) systems,

it takes only ~ 1.5 h and 0.5 h for the $RH_{BF=0.5}$ increasing to 50% and 70% , respectively. Similar results can also be found for the cases in $RH_{BF=0.2, 0.8}$.

It can be seen that in all of the investigated VOC systems, when the SOA mass fraction was > 0.05 , the phase dependence on $MR_{org/inorg}$ was qualitatively similar irrespective of the single/binary biogenic/~~anthropogenic-anthropogenic~~ SOA precursors. That is, the $RH_{BF=0.2, 0.5, 0.8}$ increased towards a maximum value with an increase of the $MR_{org/inorg}$ during SOA formation evolution. In regard to the ~~pure~~ SOA from biogenic or anthropogenic VOCs oxidation showed amorphous solid property rather than the expected liquid phase as deliquescent inorganic particles (Bateman et al., 2015a; Virtanen et al., 2010; Saukko et al., 2015; Saukko et al., 2012), such as the $RH_{BF=0.2, 0.5, 0.8}$ of ~~pure~~-SOA formed from α -pinene and toluene photo-oxidation were $80 \sim 90\%$, $80 \sim 85\%$, and $\sim 65\%$, respectively (Bateman et al., 2015a). It is expected that the increasing aerosol rebounding tendency can happen with increasing SOA mass condensing on the deliquescent sulphate seed. This speculation is supported by our results that an increase of $RH_{BF=0.2, 0.5, 0.8}$ toward the maximum with more SOA condensation. Additionally, the maximum $RH_{BF=0.2, 0.5, 0.8}$ at high $MR_{org/inorg}$ ~~up to of~~ ~ 8 in our study were $10 \sim 15\%$ lower than ~~pure~~-SOA mentioned above, indicating the presence of small mass fraction of inorganic compounds ($\sim 10\%$) makes multicomponent aerosol particles bounce less than ~~pure-the~~ SOA. Moreover, the time taken for $RH_{BF=0.2, 0.5, 0.8}$ to reach the maximum ~~dependeds~~ on how rapidly the SOA was formed, shorter time for the faster SOA production rate of the investigated VOC systems. This general behaviour was independent of the yield of the VOC and whether the precursor was biogenic or anthropogenic. In addition, the individual VOC systems behaved in the same way as the VOC mixtures.

As shown in Figure 3, it should be noted that, to avoid high noise-signal-ratio for low organic mass loading measured by HR-ToF-AMS, the data points with organic mass fraction larger than 0.05 (assuming uniform chemical composition) were selected for consideration. It was observed that particulate nitrate (Max. $5\% \sim 16\%$ of $NR-PM_1$) ~~is-was~~ formed during photochemistry in all investigated VOC systems. Nitrate can be formed either from oxidation of NO_2 followed by neutralisation by NH_3 or organic oxidation product (organic nitrate). The organic nitrate fraction in total nitrate signal can be estimated following the NO^+/NO_2^+ ratio method proposed by Farmer et al. (2010), given the differentiation of

NO⁺/NO₂⁺ ratio for pure NH₄NO₃ (2.56, from calibrations) and organic nitrate of 10~15 (Bruns et al., 2010; Fry et al., 2009; Kiendler-Scharr et al., 2016; Reyes-Villegas et al., 2018). As shown in Figure S3, organic nitrate fraction in total nitrate signal was lower than 1520% in almost all investigated VOC systems except for the last two hour of *o*-cresol system (up to ~2026%). This NO⁺/NO₂⁺ ratio method can indicate organic nitrate statistically only if the organic nitrate fraction is larger than 15% (Bruns et al., 2010). Thus the observed particulate nitrate is mainly inorganic nitrate with small contribution of organic nitrate (< 20%) for all VOC systems in this study. The small contribution of organic nitrate mass have little influences on the onset of organic mass fraction > 0.05. Moreover, Li et al. (2017) found that the pure NH₄NO₃ particles adhered on the impactor even the RH has been reduced to ~ 5%. It is worth noting that the variable particulate nitrate across all VOC systems in this study might have some influence on the phase behaviour of the multicomponent aerosol particles during SOA formation evolution and the influence of changing inorganic component ratios on the BF in multicomponent mixtures should be the focus of this work.

3.2 BF dependence on OA composition

In addition to the control of the BF by MR_{org/inorg}, there is an indication that the SOA composition across the various VOC systems may also impact aerosol phase behaviour. As indicated in Figure 2 b-c, the increase of the RH_{BF=0.5,0.8} as a function of MR_{org/inorg} is less rapid in the α -pinene/isoprene, 50% reactivity α -pinene and α -pinene system (referred as BVOC₋ systems, anthropogenic-free) than the anthropogenic-containing VOC systems (AVOC-containing systems, including *o*-cresol, 50% reactivity *o*-cresol, *o*-cresol/isoprene, α -pinene/*o*-cresol). It can be seen that, for the AVOC-containing systems, the RH_{BF=0.5} and RH_{BF=0.8} were 65~70% and 40~55% when MR_{org/inorg} approached ~1, whereas only 40~45% and ~25% in the 50% reactivity α -pinene and α -pinene/isoprene systems (BVOC₋ systems). Interestingly, with more SOA condensing, the discrepancy disappeared and the RH_{BF=0.5} and RH_{BF=0.8} converged for BVOC₋ and AVOC-containing systems as the MR_{org/inorg} > 2. In contrast, the RH_{BF=0.2} as a function of MR_{org/inorg} was the same for BVOC₋ and AVOC-containing mixtures. This indicates the decreased RH to achieve liquid-to-nonliquid transition for BVOC₋ and AVOC-containing systems was different at moderate MR_{org/inorg} of ~1 but quantitatively similar at low and high MR_{org/inorg}.

355 The above evidence indicates VOC type (BVOC- or AVOC-containing), thereafter SOA composition, can play a second-order role in the phase behaviour of multicomponent aerosol particles. Here, we used degree of oxidation of the SOA (O:C atomic ratio derived from HR-ToF-AMS) as a proxy of SOA composition to explore its ~~relation~~relationship with phase behaviour. As shown in Table 1, for BVOC- systems, the averaged atomic O:C ratio of SOA in α -pinene/isoprene, 50% reactivity α -pinene, and α -pinene systems, ~~are~~were 0.43 ± 0.05 , 0.45 ± 0.04 , 0.36 ± 0.03 , respectively. For AVOC-containing systems, the O:C ratio of SOA was 10~50% higher than BVOC systems, with 0.69 ± 0.05 , 0.66 ± 0.05 , 0.64 ± 0.03 , 0.48 ± 0.05 in *o*-cresol/isoprene, 50% reactivity *o*-cresol, *o*-cresol and α -pinene/*o*-cresol systems, respectively. In addition, the $RH_{BF=0.2, 0.5, 0.8}$ response to the $MR_{org/inorg}$ ~~was~~is coloured according to O:C ratio in Figure 2. ~~It can be seen that the~~The O:C ratio ~~is~~was characteristic for the SOA precursors and ~~shows~~showed little variation through individual experiments. No direct relationship between O:C ratio and the phase behaviour change was observed during SOA formation evolution among all investigated VOC systems. Previous studies have shown a tentative dependence on the O:C ratio of the organic particles with phase behaviour (e.g. glass transition temperature) (Koop et al., 2011; Shiraiwa et al., 2017). In contrast, Saukko et al. (2012) showed the O:C ratio only influenced rebounding behaviour of SOA formed from *n*-heptadecane in a Potential Aerosol Mass (PAM) reactor, with no influence in the α -pinene, longifolene, and naphthalene systems. There ~~is~~was some evidence showing that aerosol morphology (single well-mixed phase or phase separation) ~~is~~was closely related to O:C ratio of SOA and organic-inorganic-mass ratio (Bertram et al., 2011; Krieger et al., 2012; You et al., 2014; Freedman, 2017; Song et al., 2012; Smith et al., 2013). It is unknown whether the morphology plays a role in the phase behaviour discrepancy between BVOC- and AVOC-containing systems at moderate $MR_{org/inorg}$ considering their O:C difference, which is of interest and needs further investigations.

3.3 Mixing role of chemical composition and RH on BF

As we know that the chemical composition and RH are key factors influencing aerosol water uptake at given size (Kreidenweis and Asa-Awuku, 2014). To discuss the mixing role of chemical composition and RH on BF, the GF at the RH of BF measurement was calculated from HTDMA measurement using κ -Köhler equation, and the relation between the BF and the GF was plotted in Figure 5. It can be seen that

385 the BF showed a monotonic decrease from ~ 1 to ~ 0 with the increasing GF from ~ 1 to ~ 1.15 in all VOC systems. As the GF is larger than 1.15, the aerosol particles kept in liquid phase (BF ~ 0). This evidence indicated the key role of increasing aerosol water uptake on the phase transition from the non-liquid to the liquid. The BF-GF relation of the varying multicomponent aerosol particles including SOA and inorganic compounds is comparable with the previous study (Bateman et al., 2015a). They measured the BF of the SOA from α -pinene, toluene and isoprene and found that the BF is nearly 0 when the GF is larger than > 1.15 (Bateman et al., 2015a)(Bateman et al., 2015a).

390 **4 Conclusions and implications**

Our experiments support the validity of our two proposed hypotheses: 1) Aerosol phase behaviour for SOA mixture is determined by RH and organic-inorganic-mass ratio ($MR_{org/inorg}$). 2) The difference in SOA composition is less important in determining the phase behaviour than the rate of SOA production, which changes the $MR_{org/inorg}$.

395 First, aerosol phase behaviour is clearly RH-dependent as already widely known. Multicomponent aerosol particles were always found to exhibit liquid-like behaviour ($BF < 0.2$) above 80% RH, and nonliquid-like behaviour ($BF > 0.8$) at below 30% RH. The bounce measurements always indicated continuously increasing nonliquid-like behaviour as RH decreased from 80% to 30%. These RH-dependent rebound curves strongly depended on the increase in $MR_{org/inorg}$ during SOA formation in all investigated VOC systems. The identified $RH_{BF=0.2, 0.5, 0.8}$ increased towards a maximum with the increase of the $MR_{org/inorg}$, and the increase rate of $RH_{BF=0.2, 0.5, 0.8}$ is determined by the SOA production rate on sulphate seed across the VOC systems. This general behaviour was independent of the yield of the SOA precursors and whether the precursor was biogenic or anthropogenic. In some ways, this is an obvious result that follows from the rate of SOA mass increase in the system under investigation. However, this production rate will
405 be dependent on the reactivity and yield of VOCs in the mixture, their concentrations and interactions

influencing SOA mass formation. This set of complex dependencies will control the changes in particle $MR_{org/inorg}$ in mixtures and hence in phase behaviour.

410 Although less important than the RH and $MR_{org/inorg}$, the SOA composition plays a secondary role affecting the phase behaviour. It was observed that the $RH_{BF=0.2}$ as a function of $MR_{org/inorg}$ was the same for BVOC (AVOC-free) and AVOC-containing systems, however the decreased RH to achieve liquid-to-
415 nonliquid transition (BF from ~ 0.2 to ~ 0.8) was different at moderate $MR_{org/inorg}$ of ~ 1 but quantitatively similar at low and high $MR_{org/inorg}$. For example, at moderate $MR_{org/inorg}$ of ~ 1 , the $RH_{BF=0.2}$ for BVOC and AVOC-containing systems was $65 \sim 70\%$. To achieve liquid-to-nonliquid transition with BF of ~ 0.8 , the RH needed to decrease to $40 \sim 55\%$ in AVOC-containing systems but to $\sim 25\%$ in BVOC systems.
420 This discrepancy cannot be explained by the atomic O:C ratio of SOA during SOA formation evolution. It should be the focus of future work to investigate whether SOA composition has a more pronounced effect outside the dynamic O:C range of the mixtures in our experiments. Additionally, by combining the chemical composition and RH, we calculated the hygroscopic growth factor (GF) and found its key role in impacting phase behaviour. The multicomponent aerosol particles were liquid in all VOC systems when the GF is higher than 1.15 at room temperature and transmitted from liquid to non-liquid when the GF decreased to ~ 1 .

MM Multicomponent aerosol phase behaviour depends on RH and organic-inorganic-mass ratio in a particle and in the atmosphere as well as our chamber. Any interactions influencing these two factors will therefore influence the phase behaviour. For example, a fast formation of SOA or significant ambient RH
425 change in real atmosphere could change phase behaviour of particles, and consequently influencing atmospheric physicochemical processes. There is an additional indication that an increased anthropogenic VOC contribution to the mixture could give different phase behaviour at certain moderate organic mass fraction ($\sim 50\%$), hence there will be some second-order differences depending on the relative contributions of anthropogenic and biogenic VOC. Further studies should be carried out on more complex
430 and realistic atmospheric mixtures.

Data availability

The observational dataset of this study ~~will be is~~-available on the open dataset of EUROCHAMP-2020 programme ~~shortly~~ (<https://data.eurochamp.org/data-access/chamber-experiments/>).

440 Author contributions

~~Y.W. conceived idea of this study and~~ G.M., M.R.A., Y.W., A.V. and Y.S. ~~conceived-co-designed the experiments~~~~the study~~. G.M. and Z.W. co-applied Trans-National Access (TNA) funding from EUROCHAMP for the phase behaviour measurement. Y.W., A.V., Y.S. and D.M. conducted the chamber experiments and collected the dataset. T.Z. and X.M. operated the rebound impactor apparatus. Y.C. and
445 D.H. provided helpful discussions. Y.W. performed data integration, data analysis and wrote the manuscript with the inputs from all co-authors under the guidance of G.M. and M.R.A.

Acknowledgement

Z.W. acknowledges National Natural Science Foundation of China (41875149). Y.W. acknowledges the ~~joint scholarship~~~~joint scholarship~~ of The University of Manchester and Chinese Scholarship Council.

450 M.R.A. acknowledges support by UK National Centre for Atmospheric Sciences (NACS) funding. A.V.
acknowledges the Natural Environment Research Council (NERC) EAO Doctoral Training Partnership
funding. This work is supported by the following projects: Trans-National Access (TNA) of
EUROCHAMP-2020. We ~~acknowledges~~acknowledge AMF/AMOF for providing SMPS instrument
(AMF_25072016114543 and AMF_04012017142558).

455 **Competing interests**

All authors declare no conflict of interest.

References

- 460 Alfarra, M. R., Coe, H., Allan, J. D., Bower, K. N., Boudries, H., Canagaratna, M. R., Jimenez, J. L., Jayne, J. T., Garforth, A.
A., Li, S.-M., and Worsnop, D. R.: Characterization of urban and rural organic particulate in the Lower Fraser Valley using
two Aerodyne Aerosol Mass Spectrometers, *Atmospheric Environment*, 38, 5745-5758,
<https://doi.org/10.1016/j.atmosenv.2004.01.054>, 2004.
- 465 Alfarra, M. R., Hamilton, J. F., Wyche, K. P., Good, N., Ward, M. W., Carr, T., Barley, M. H., Monks, P. S., Jenkin, M. E.,
Lewis, A. C., and McFiggans, G. B.: The effect of photochemical ageing and initial precursor concentration on the composition
and hygroscopic properties of β -caryophyllene secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 6417-6436, 10.5194/acp-
12-6417-2012, 2012.
- Allan, J. D., Jimenez, J. L., Williams, P. I., Alfarra, M. R., Bower, K. N., Jayne, J. T., Coe, H., and Worsnop, D. R.: Quantitative
sampling using an Aerodyne aerosol mass spectrometer 1. Techniques of data interpretation and error analysis, *Journal of
Geophysical Research: Atmospheres*, 108, 10.1029/2002jd002358, 2003.
- 470 Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A. M., Drewnick, F., Onasch,
T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A generalised method for the extraction of chemically resolved
mass spectra from Aerodyne aerosol mass spectrometer data, *Journal of Aerosol Science*, 35, 909-922,
<https://doi.org/10.1016/j.jaerosci.2004.02.007>, 2004.
- Bateman, A. P., Belassein, H., and Martin, S. T.: Impactor Apparatus for the Study of Particle Rebound: Relative Humidity
and Capillary Forces, *Aerosol Science and Technology*, 48, 42-52, 10.1080/02786826.2013.853866, 2014.
- 475 Bateman, A. P., Bertram, A. K., and Martin, S. T.: Hygroscopic Influence on the Semisolid-to-Liquid Transition of Secondary
Organic Materials, *The Journal of Physical Chemistry A*, 119, 4386-4395, 10.1021/jp508521c, 2015a.









- Bateman, A. P., Gong, Z., Liu, P., Sato, B., Cirino, G., Zhang, Y., Artaxo, P., Bertram, A. K., Manzi, A. O., Rizzo, L. V., Souza, R. A. F., Zaveri, R. A., and Martin, S. T.: Sub-micrometre particulate matter is primarily in liquid form over Amazon rainforest, *Nature Geoscience*, 9, 34, 10.1038/ngeo2599
- 480 <https://www.nature.com/articles/ngeo2599#supplementary-information>, 2015b.
- Bateman, A. P., Gong, Z., Harder, T. H., de Sá, S. S., Wang, B., Castillo, P., China, S., Liu, Y., O'Brien, R. E., Palm, B. B., Shiu, H. W., Cirino, G. G., Thalman, R., Adachi, K., Alexander, M. L., Artaxo, P., Bertram, A. K., Buseck, P. R., Gilles, M. K., Jimenez, J. L., Laskin, A., Manzi, A. O., Sedlacek, A., Souza, R. A. F., Wang, J., Zaveri, R., and Martin, S. T.: Anthropogenic influences on the physical state of submicron particulate matter over a tropical forest, *Atmos. Chem. Phys.*, 17, 1759-1773, 10.5194/acp-17-1759-2017, 2017.
- 485
- Bertram, A. K., Martin, S. T., Hanna, S. J., Smith, M. L., Bodsworth, A., Chen, Q., Kuwata, M., Liu, A., You, Y., and Zorn, S. R.: Predicting the relative humidities of liquid-liquid phase separation, efflorescence, and deliquescence of mixed particles of ammonium sulfate, organic material, and water using the organic-to-sulfate mass ratio of the particle and the oxygen-to-carbon elemental ratio of the organic component, *Atmos. Chem. Phys.*, 11, 10995-11006, 10.5194/acp-11-10995-2011, 2011.
- 490
- Bruns, E. A., Perraud, V., Zelenyuk, A., Ezell, M. J., Johnson, S. N., Yu, Y., Imre, D., Finlayson-Pitts, B. J., and Alexander, M. L.: Comparison of FTIR and Particle Mass Spectrometry for the Measurement of Particulate Organic Nitrates, *Environmental Science & Technology*, 44, 1056-1061, 10.1021/es9029864, 2010.
- Coeur-Tourneur, C., Henry, F., Janquin, M.-A., and Brutier, L.: Gas-phase reaction of hydroxyl radicals with m-, o- and p-cresol, *International Journal of Chemical Kinetics*, 38, 553-562, 10.1002/kin.20186, 2006.
- 495
- Coggon, M. M., Lim, C. Y., Koss, A. R., Sekimoto, K., Yuan, B., Gilman, J. B., Hagan, D. H., Selimovic, V., Zarzana, K. J., Brown, S. S., Roberts, J. M., Müller, M., Yokelson, R., Wisthaler, A., Krechmer, J. E., Jimenez, J. L., Cappa, C., Kroll, J. H., de Gouw, J., and Warneke, C.: OH chemistry of non-methane organic gases (NMOGs) emitted from laboratory and ambient biomass burning smoke: evaluating the influence of furans and oxygenated aromatics on ozone and secondary NMOG formation, *Atmos. Chem. Phys.*, 19, 14875-14899, 10.5194/acp-19-14875-2019, 2019.
- 500
- Dahneke, B.: The capture of aerosol particles by surfaces, *Journal of Colloid and Interface Science*, 37, 342-353, [https://doi.org/10.1016/0021-9797\(71\)90302-X](https://doi.org/10.1016/0021-9797(71)90302-X), 1971.
- DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, *Analytical Chemistry*, 78, 8281-8289, 10.1021/ac061249n, 2006.
- 505
- Farmer, D. K., Matsunaga, A., Docherty, K. S., Surratt, J. D., Seinfeld, J. H., Ziemann, P. J., and Jimenez, J. L.: Response of an aerosol mass spectrometer to organonitrates and organosulfates and implications for atmospheric chemistry, *Proceedings of the National Academy of Sciences*, 107, 6670-6675, 10.1073/pnas.0912340107, 2010.
- Fishbein, L.: An overview of environmental and toxicological aspects of aromatic hydrocarbons II. Toluene, *Science of The Total Environment*, 42, 267-288, [https://doi.org/10.1016/0048-9697\(85\)90062-2](https://doi.org/10.1016/0048-9697(85)90062-2), 1985.
- 510
- Freedman, M. A.: Phase separation in organic aerosol, *Chemical Society Reviews*, 46, 7694-7705, 10.1039/C6CS00783J, 2017.
- 515
- Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H., Dubé, W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H. P., Brauers, T., and Cohen, R. C.: Organic nitrate and secondary organic aerosol yield from NO₃ oxidation of β -pinene evaluated using a gas-phase kinetics/aerosol partitioning model, *Atmos. Chem. Phys.*, 9, 1431-1449, 10.5194/acp-9-1431-2009, 2009.

- Good, N., Coe, H., and McFiggans, G.: Instrumentational operation and analytical methodology for the reconciliation of aerosol water uptake under sub- and supersaturated conditions, *Atmos. Meas. Tech.*, 3, 1241-1254, 10.5194/amt-3-1241-2010, 2010.
- 520 Henry, F., Coeur-Tourneur, C., Ledoux, F., Tomas, A., and Menu, D.: Secondary organic aerosol formation from the gas phase reaction of hydroxyl radicals with m-, o- and p-cresol, *Atmospheric Environment*, 42, 3035-3045, <https://doi.org/10.1016/j.atmosenv.2007.12.043>, 2008.
- 525 Ignatius, K., Kristensen, T. B., Järvinen, E., Niehman, L., Fuchs, C., Gordon, H., Herenz, P., Hoyle, C. R., Duplissy, J., Garimella, S., Dias, A., Frege, C., Höppel, N., Tröstl, J., Wagner, R., Yan, C., Amorim, A., Baltensperger, U., Curtius, J., Donahue, N. M., Gallagher, M. W., Kirkby, J., Kulmala, M., Möhler, O., Saathoff, H., Schnaiter, M., Tomé, A., Virtanen, A., Worsnop, D., and Stratmann, F.: Heterogeneous ice nucleation of viscous secondary organic aerosol produced from ozonolysis of α -pinene, *Atmos. Chem. Phys.*, 16, 6495-6509, 10.5194/acp-16-6495-2016, 2016.
- Jayne, J. T., Leard, D. C., Zhang, X., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development of an Aerosol Mass Spectrometer for Size and Composition Analysis of Submicron Particles, *Aerosol Science and Technology*, 33, 49-70, 10.1080/027868200410840, 2000.
- 530 Jimenez, J. L., Jayne, J. T., Shi, Q., Kolb, C. E., Worsnop, D. R., Yourshaw, I., Seinfeld, J. H., Flagan, R. C., Zhang, X., Smith, K. A., Morris, J. W., and Davidovits, P.: Ambient aerosol sampling using the Aerodyne Aerosol Mass Spectrometer, *Journal of Geophysical Research: Atmospheres*, 108, doi:10.1029/2001JD001213, 2003.
- 535 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525-1529, 10.1126/science.1180353, 2009.
- 540 Kiendler-Scharr, A., Mensah, A. A., Friese, E., Topping, D., Nemitz, E., Prevot, A. S. H., Äijälä, M., Allan, J., Canonaco, F., Canagaratna, M., Carbone, S., Crippa, M., Dall'Osto, M., Day, D. A., De Carlo, P., Di Marco, C. F., Elbern, H., Eriksson, A., Freney, E., Hao, L., Herrmann, H., Hildebrandt, L., Hillamo, R., Jimenez, J. L., Laaksonen, A., McFiggans, G., Mohr, C., O'Dowd, C., Otjes, R., Ovadnevaite, J., Pandis, S. N., Poulain, L., Schlag, P., Sellegri, K., Swietlicki, E., Tiitta, P., Vermeulen, A., Wahner, A., Worsnop, D., and Wu, H.-C.: Ubiquity of organic nitrates from nighttime chemistry in the European submicron aerosol, *Geophysical Research Letters*, 43, 7735-7744, 10.1002/2016gl069239, 2016.
- 545 Koop, T., Bookhold, J., Shiraiwa, M., and Poschl, U.: Glass transition and phase state of organic compounds: dependency on molecular properties and implications for secondary organic aerosols in the atmosphere, *Physical Chemistry Chemical Physics*, 13, 19238-19255, 10.1039/C1CP22617G, 2011.
- 550 Koss, A. R., Sekimoto, K., Gilman, J. B., Selimovic, V., Coggon, M. M., Zarzana, K. J., Yuan, B., Lerner, B. M., Brown, S. S., Jimenez, J. L., Krechmer, J., Roberts, J. M., Warneke, C., Yokelson, R. J., and de Gouw, J.: Non-methane organic gas emissions from biomass burning: identification, quantification, and emission factors from PTR-ToF during the FIREX 2016 laboratory experiment, *Atmos. Chem. Phys.*, 18, 3299-3319, 10.5194/acp-18-3299-2018, 2018.
- 555 Kreidenweis, S., and Asa-Awuku, A.: Aerosol Hygroscopicity: Particle Water Content and Its Role in Atmospheric Processes, 331-361 pp., 2014.

- Krieger, U. K., Marcolli, C., and Reid, J. P.: Exploring the complexity of aerosol particle properties and processes using single particle techniques, *Chemical Society Reviews*, 41, 6631-6662, 10.1039/C2CS35082C, 2012.
- 560 Kuwata, M., and Martin, S. T.: Phase of atmospheric secondary organic material affects its reactivity, *Proceedings of the National Academy of Sciences*, 109, 17354-17359, 10.1073/pnas.1209071109, 2012.
- Li, Y. J., Liu, P. F., Bergoend, C., Bateman, A. P., and Martin, S. T.: Rebounding hygroscopic inorganic aerosol particles: Liquids, gels, and hydrates, *Aerosol Science and Technology*, 51, 388-396, 10.1080/02786826.2016.1263384, 2017.
- 565 Liu, Y., Wu, Z., Wang, Y., Xiao, Y., Gu, F., Zheng, J., Tan, T., Shang, D., Wu, Y., Zeng, L., Hu, M., Bateman, A. P., and Martin, S. T.: Submicrometer Particles Are in the Liquid State during Heavy Haze Episodes in the Urban Atmosphere of Beijing, China, *Environmental Science & Technology Letters*, 4, 427-432, 10.1021/acs.estlett.7b00352, 2017.
- Liu, Y., Wu, Z., Huang, X., Shen, H., Bai, Y., Qiao, K., Meng, X., Hu, W., Tang, M., and He, L.: Aerosol Phase State and Its Link to Chemical Composition and Liquid Water Content in a Subtropical Coastal Megacity, *Environmental Science & Technology*, 53, 5027-5033, 10.1021/acs.est.9b01196, 2019.
- 570 Martin, S. T.: Phase Transitions of Aqueous Atmospheric Particles, *Chemical Reviews*, 100, 3403-3454, 10.1021/cr990034t, 2000.
- McFiggans, G., Mentel, T. F., Wildt, J., Pullinen, I., Kang, S., Kleist, E., Schmitt, S., Springer, M., Tillmann, R., Wu, C., Zhao, D., Hallquist, M., Faxon, C., Le Breton, M., Hallquist, Å. M., Simpson, D., Bergström, R., Jenkin, M. E., Ehn, M., Thornton, J. A., Alfarra, M. R., Bannan, T. J., Percival, C. J., Priestley, M., Topping, D., and Kiendler-Scharr, A.: Secondary organic aerosol reduced by mixture of atmospheric vapours, *Nature*, 565, 587-593, 10.1038/s41586-018-0871-y, 2019.
- 575 Murray, B. J.: Inhibition of ice crystallisation in highly viscous aqueous organic acid droplets, *Atmos. Chem. Phys.*, 8, 5423-5433, 10.5194/acp-8-5423-2008, 2008.
- Murray, B. J., Wilson, T. W., Dobbie, S., Cui, Z., Al-Jumur, S. M. R. K., Möhler, O., Schnaiter, M., Wagner, R., Benz, S., Niemand, M., Saathoff, H., Ebert, V., Wagner, S., and Kärcher, B.: Heterogeneous nucleation of ice particles on glassy aerosols under cirrus conditions, *Nature Geoscience*, 3, 233, 10.1038/ngeo817
580 <https://www.nature.com/articles/ngeo817#supplementary-information>, 2010.
- Pajunoja, A., Hu, W., Leong, Y. J., Taylor, N. F., Miettinen, P., Palm, B. B., Mikkonen, S., Collins, D. R., Jimenez, J. L., and Virtanen, A.: Phase state of ambient aerosol linked with water uptake and chemical aging in the southeastern US, *Atmos. Chem. Phys.*, 16, 11163-11176, 10.5194/acp-16-11163-2016, 2016.
- 585 Pedernera, D. A.: Glass formation in upper tropospheric aerosol particles, Ph.D. thesis (in German), Bielefeld University, <http://pub.uni-bielefeld.de/publication/2303351>, 2008.
- Petters, M. D., and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961-1971, 10.5194/acp-7-1961-2007, 2007.
- 590 Pöschl, U.: Gas-particle interactions of tropospheric aerosols: Kinetic and thermodynamic perspectives of multiphase chemical reactions, amorphous organic substances, and the activation of cloud condensation nuclei, *Atmospheric Research*, 101, 562-573, <https://doi.org/10.1016/j.atmosres.2010.12.018>, 2011.
- Reid, J. P., Bertram, A. K., Topping, D. O., Laskin, A., Martin, S. T., Petters, M. D., Pope, F. D., and Rovelli, G.: The viscosity of atmospherically relevant organic particles, *Nature Communications*, 9, 956, 10.1038/s41467-018-03027-z, 2018.

- Renbaum-Wolff, L., Grayson, J. W., Bateman, A. P., Kuwata, M., Sellier, M., Murray, B. J., Shilling, J. E., Martin, S. T., and Bertram, A. K.: Viscosity of α -pinene secondary organic material and implications for particle growth and reactivity, *Proceedings of the National Academy of Sciences*, 110, 8014-8019, 10.1073/pnas.1219548110, 2013.
- Reyes-Villegas, E., Priestley, M., Ting, Y. C., Haslett, S., Bannan, T., Le Breton, M., Williams, P. I., Bacak, A., Flynn, M. J., Coe, H., Percival, C., and Allan, J. D.: Simultaneous aerosol mass spectrometry and chemical ionisation mass spectrometry measurements during a biomass burning event in the UK: insights into nitrate chemistry, *Atmos. Chem. Phys.*, 18, 4093-4111, 10.5194/acp-18-4093-2018, 2018.
- Saukko, E., Lambe, A. T., Massoli, P., Koop, T., Wright, J. P., Croasdale, D. R., Pedernera, D. A., Onasch, T. B., Laaksonen, A., Davidovits, P., Worsnop, D. R., and Virtanen, A.: Humidity-dependent phase state of SOA particles from biogenic and anthropogenic precursors, *Atmos. Chem. Phys.*, 12, 7517-7529, 10.5194/acp-12-7517-2012, 2012.
- Saukko, E., Zorn, S., Kuwata, M., Keskinen, J., and Virtanen, A.: Phase State and Deliquescence Hysteresis of Ammonium-Sulfate-Seeded Secondary Organic Aerosol, *Aerosol Science and Technology*, 49, 531-537, 10.1080/02786826.2015.1050085, 2015.
- Shao, Y., Wang, Y., Du, M., Voliotis, A., Alfarra, M. R., Turner, S. F., and McFiggans, G.: Characterisation of the Manchester Aerosol Chamber facility, *Atmos. Meas. Tech. Discuss.*, 2021, 1-50, 10.5194/amt-2021-147, 2021.
- Shilling, J. E., Zawadowicz, M. A., Liu, J., Zaveri, R. A., and Zelenyuk, A.: Photochemical Aging Alters Secondary Organic Aerosol Partitioning Behavior, *ACS Earth and Space Chemistry*, 3, 2704-2716, 10.1021/acsearthspacechem.9b00248, 2019.
- Shiraiwa, M., Ammann, M., Koop, T., and Pöschl, U.: Gas uptake and chemical aging of semisolid organic aerosol particles, *Proceedings of the National Academy of Sciences*, 108, 11003-11008, 10.1073/pnas.1103045108, 2011.
- Shiraiwa, M., and Seinfeld, J. H.: Equilibration timescale of atmospheric secondary organic aerosol partitioning, *Geophysical Research Letters*, 39, 10.1029/2012gl054008, 2012.
- Shiraiwa, M., Li, Y., Tsimpidi, A. P., Karydis, V. A., Berkemeier, T., Pandis, S. N., Lelieveld, J., Koop, T., and Pöschl, U.: Global distribution of particle phase state in atmospheric secondary organic aerosols, *Nature Communications*, 8, 15002, 10.1038/ncomms15002, 2017.
- Slade, J. H., Shiraiwa, M., Arangio, A., Su, H., Pöschl, U., Wang, J., and Knopf, D. A.: Cloud droplet activation through oxidation of organic aerosol influenced by temperature and particle phase state, *Geophysical Research Letters*, 44, 1583-1591, 10.1002/2016gl072424, 2017.
- Slade, J. H., Ault, A. P., Bui, A. T., Ditto, J. C., Lei, Z., Bondy, A. L., Olson, N. E., Cook, R. D., Desrochers, S. J., Harvey, R. M., Erickson, M. H., Wallace, H. W., Alvarez, S. L., Flynn, J. H., Boor, B. E., Petrucci, G. A., Gentner, D. R., Griffin, R. J., and Shepson, P. B.: Bouncer Particles at Night: Biogenic Secondary Organic Aerosol Chemistry and Sulfate Drive Diel Variations in the Aerosol Phase in a Mixed Forest, *Environmental Science & Technology*, 53, 4977-4987, 10.1021/acs.est.8b07319, 2019.
- Smith, M. L., Kuwata, M., and Martin, S. T.: Secondary Organic Material Produced by the Dark Ozonolysis of α -Pinene Minimally Affects the Deliquescence and Efflorescence of Ammonium Sulfate, *Aerosol Science and Technology*, 45, 244-261, 10.1080/02786826.2010.532178, 2011.
- Smith, M. L., Bertram, A. K., and Martin, S. T.: Deliquescence, efflorescence, and phase miscibility of mixed particles of ammonium sulfate and isoprene-derived secondary organic material, *Atmos. Chem. Phys.*, 12, 9613-9628, 10.5194/acp-12-9613-2012, 2012.

- Smith, M. L., You, Y., Kuwata, M., Bertram, A. K., and Martin, S. T.: Phase Transitions and Phase Miscibility of Mixed Particles of Ammonium Sulfate, Toluene-Derived Secondary Organic Material, and Water, *The Journal of Physical Chemistry A*, 117, 8895-8906, 10.1021/jp405095e, 2013.
- 635 Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid-liquid phase separation in aerosol particles: Dependence on O:C, organic functionalities, and compositional complexity, *Geophysical Research Letters*, 39, 10.1029/2012GL052807, 2012.
- Stein, S. W., Turpin, B. J., Cai, X., Huang, P.-F., and McMurry, P. H.: Measurements of relative humidity-dependent bounce and density for atmospheric particles using the DMA-impactor technique, *Atmospheric Environment*, 28, 1739-1746, [https://doi.org/10.1016/1352-2310\(94\)90136-8](https://doi.org/10.1016/1352-2310(94)90136-8), 1994.
- 640 Tang, I. N., and Munkelwitz, H. R.: Composition and temperature dependence of the deliquescence properties of hygroscopic aerosols, *Atmospheric Environment. Part A. General Topics*, 27, 467-473, [https://doi.org/10.1016/0960-1686\(93\)90204-C](https://doi.org/10.1016/0960-1686(93)90204-C), 1993.
- Vaden, T. D., Imre, D., Beránek, J., Shrivastava, M., and Zelenyuk, A.: Evaporation kinetics and phase of laboratory and ambient secondary organic aerosol, *Proceedings of the National Academy of Sciences*, 108, 2190-2195, 645 10.1073/pnas.1013391108, 2011.
- Virtanen, A., Joutsensaari, J., Koop, T., Kannosto, J., Yli-Pirilä, P., Leskinen, J., Mäkelä, J. M., Holopainen, J. K., Pöschl, U., Kulmala, M., Worsnop, D. R., and Laaksonen, A.: An amorphous solid state of biogenic secondary organic aerosol particles, *Nature*, 467, 824, 10.1038/nature09455 <https://www.nature.com/articles/nature09455#supplementary-information>, 2010.
- 650 You, Y., Smith, M. L., Song, M., Martin, S. T., and Bertram, A. K.: Liquid-liquid phase separation in atmospherically relevant particles consisting of organic species and inorganic salts, *International Reviews in Physical Chemistry*, 33, 43-77, 10.1080/0144235X.2014.890786, 2014.
- Zaveri, R. A., Easter, R. C., Shilling, J. E., and Seinfeld, J. H.: Modeling kinetic partitioning of secondary organic aerosol and size distribution dynamics: representing effects of volatility, phase state, and particle-phase reaction, *Atmos. Chem. Phys.*, 14, 655 5153-5181, 10.5194/acp-14-5153-2014, 2014.
- Zhang, Y., Chen, Y., Lambe, A. T., Olson, N. E., Lei, Z., Craig, R. L., Zhang, Z., Gold, A., Onasch, T. B., Jayne, J. T., Worsnop, D. R., Gaston, C. J., Thornton, J. A., Vizuete, W., Ault, A. P., and Surratt, J. D.: Effect of the Aerosol-Phase State on Secondary Organic Aerosol Formation from the Reactive Uptake of Isoprene-Derived Epoxydiols (IEPOX), *Environmental Science & Technology Letters*, 5, 167-174, 10.1021/acs.estlett.8b00044, 2018.
- 660







Figure 1. Time series of the Bounce fraction (BF) as a function of RH measured by three-arm particle rebound apparatus with RH adjustment system of the 100-200 nm secondary organic aerosol (SOA) formed from various iso-reactive single/binary biogenic/anthropogenic VOCs (volatile organic compounds) photochemistry under relative low- NO_x condition on deliquescent ammonium sulphate seed (as the SOA mass fraction in NR- PM_{10} is larger than 0.05).







Figure 2. The measured (a) $RH_{BF=0.2}$, (b) $RH_{BF=0.5}$ and (c) $RH_{BF=0.8}$ as a function of organic-inorganic ratio ($MR_{org/inorg}$) and coloured by atomic O:C ratio in various VOC systems photochemistry on deliquescent ammonium sulphate seed. Black box points out the $RH_{BF=0.2}$, 0.5, 0.8 at moderate $MR_{org/inorg}$ of ~ 1 for biogenic VOC systems (α -pinene/isoprene, 50% reactivity α -pinene).






Figure 3. Summary of mass fraction of chemical species (SO₄, NO₃, NH₄, SOA) in bulk NR-PM₁ measured by HR-ToF-AMS in various VOC systems photochemistry on deliquescent ammonium sulphate seed. The black dashed line represents the defined time where SOA mass fraction is 0.05.




Figure 4. The time series of measured (a) $RH_{BF=0.2}$, (b) $RH_{BF=0.5}$ and (c) $RH_{BF=0.8}$ of the multi-component aerosol particles in various VOC systems photochemistry on deliquescent ammonium sulphate seed.




Figure 5. The relation of BF and hygroscopic GF of aerosol particles during six-hour photochemistry experiments in various VOC systems.