

# First insights into Northern Africa high-altitude background aerosol chemical composition and source influences

a mis en forme : Police :17 pt

Nabil Deabji<sup>1,2</sup>, Khanneh Wadinga Fomba<sup>1</sup>, Laurent Poulain<sup>1</sup>, Sebastian Zeppenfeld<sup>1</sup>, Souad El Hajjaji<sup>2</sup>, Abdelwahid Mellouki<sup>3</sup>

a mis en forme : Police :12 pt

5 and Hartmut Herrmann<sup>1</sup>

<sup>1</sup>Leibniz Institute for Tropospheric Research (TROPOS), Atmospheric Chemistry Department (ACD), Permoserstraße 15, 04318, Leipzig, Germany

<sup>2</sup>LS3MN3E-CERNE2D, Faculty of Science, Mohammed V University in Rabat, 4 Avenue Ibn Battouta, B.P. 1040, 10100 Rabat, Morocco

10 <sup>3</sup>Institut de Combustion Aérothermique Réactivité et Environnement/OSUC-CNRS, 1C Avenue de la Recherche Scientifique, 45071 Orléans Cedex 2, France

Correspondence to: Hartmut Herrmann (herrmann@tropos.de)

**Abstract.** Field measurements were conducted to determine aerosol chemical composition in a newly established remote high-altitude site in North Africa, at the Atlas Mohammed V atmospheric observatory (AM5) located in the Middle-Atlas Mountains. The main objectives of AM5 are and to investigate the variations in the aerosol composition, and better assess global and regional changes in atmospheric composition in north Africa. A total of 200 Particulate matter (PM<sub>10</sub>) filter samples were collected in Morocco using a high-volume (HV) collector in a 12h sampling interval from August to December 2017. The chemical composition of the samples was analyzed for trace metals, water soluble ions, and organic carbon (OC/EC), aliphatic hydrocarbons, and polycyclic aromatic hydrocarbon (PAHs) contents.

15 The results indicate that high-altitudes aerosol composition is influenced by both regional as well as trans-regional transport of emissions. However, local sources play an important role, especially during low wind speed periods, as observed for November and December. During background conditions characterized by low wind speeds (Av. 20 3 m s<sup>-1</sup>) and mass concentrations in the ranged from 9.8 to 1220  $\mu\text{g m}^{-3}$ . The chemical composition is found to be dominated by inorganic elements, mainly suspended dust (61.47%) and ionic species (7%), followed by organic matter (7.15%), water content (12.12%), and unidentified mass (11.9%). Despite the proximity of the site to the Sahara Desert, its influence on the atmospheric composition at this high-altitude site was mainly seasonal and accounted for only 22% of the sampling duration. Biogenic organics contributed up to 7% of the organic matter 25 with high contributions from compounds such as heneicosane, hentriacontane, and nonacosane. The AM5 site is dominated by four main air mass inflow, which often leads to different aerosol chemical compositions. Mineral dust influenced was seasonal and ranged between 210 and 740% of the PM mass with peaks observed during the summer and was accompanied by high concentrations of SO<sub>4</sub><sup>2-</sup> of up to 4.33.0  $\mu\text{g m}^{-3}$ . During winter, PM<sub>10</sub> concentrations are low (< 30  $\mu\text{g m}^{-3}$ ), the influence of the desert is weaker, and the marine air masses (64.53%) 30 are more dominant with a mixture of sea salt and polluted aerosol from the coastal regions (Rabat and Casablanca). During the daytime, mineral dust contribution to PM increased by about 42% because of road dust resuspension. In contrast, during night-time, an increase in the concentrations of alkanes, PAHs, alkane-2-oneketones, and anthropogenic metals such as Pb, Ni, and Cu was found due to variations in the boundary layer height. The results 35

a mis en forme : Non Surlignage

a mis en forme : Non Surlignage

a mis en forme : Exposant

a mis en forme : Non Surlignage

a mis en forme : Non Surlignage

a mis en forme : Non Surlignage

provide the first insights detailed seasonal and diurnal variation of the aerosol chemical composition,  
40 valuableuseful for long-term assessment of climate and regional influence of air pollution in North Africa.

## 1 Introduction

Aerosols are important constituents of the atmosphere due to their role in controlling climate processes and their impact on air quality, the environment, and ecosystems. They can have adverse effects on human health and have been associated with respiratory disorders, strokes, pulmonary and cardiovascular diseases (Du et al., 2016; Pope  
45 et al., 2018; Song et al., 2014). Aerosol particles can serve as cloud condensation nuclei and as substrates for heterogeneous reactions (Leng et al., 2014). Their chemical composition affects aerosol-cloud interaction and may exert a warming or a cooling influence on the atmosphere due to direct and radiative forcing (King et al., 2003; Satheesh and Krishna Moorthy, 2005). Therefore, the study of aerosol chemical properties is essential for a better understanding of atmospheric processes.

50 Atmospheric aerosol particle composition depends on local and regional emission sources as well as transboundary pollution. The particles are emitted directly into the atmosphere from natural sources such as sea salt or mineral dust and anthropogenic activities such as industrial or traffic emissions, constituting primary emissions. They can also be formed in the atmosphere through gas-to-particle conversion or particle-phase reactions, constituting secondary aerosols (Carter et al., 2005). After emission, these particles are exposed to  
55 changing humidity, temperature, pressure, and solar radiation in the atmosphere that alters their properties through different aging and oxidative processes during atmospheric transport. Consequently, high-altitude sites provide the required infrastructure for investigating and characterizing the possible atmospheric aerosol interactions associated with the particles.

High-altitude sites in remote regions are less affected by direct local anthropogenic emissions. Their high altitude  
60 allows the study of aerosol particles in the free atmosphere and provides a good impression of aerosol background concentrations. The topography, meteorological conditions, and changing boundary layer heights provide various pathways for aerosol interactions, which could influence inversion processes and enhance biogenic particle formation. Such sites are hence unique for monitoring the temporal variation in the aerosol chemical compositions over longer periods and provide a better understanding of various factors, such as meteorology, climate, and  
65 environmental changes that may in the long-term effectaffect the local and regional air composition (Okamoto and Tanimoto, 2016).

There is increasing interest in atmospheric aerosol studies at high-altitude sites. Studies have investigated the aerosol chemical composition in mountainous regions, highlighting the influence of mountain valleys, night-time  
70 mountain breeze, and topography in dispersing polluted air masses to the free troposphere (Zhang et al., 2009; Alastuey et al., 2005; Buchunde et al., 2019; Leena et al., 2017; Glasius et al., 2018; Lugauer et al., 1998; Mukherjee et al., 2020). Furthermore, other studies at the Northeastern Himalayas, India (Chatterjee et al., 2010),  
75 the Bachelor Observatory Mountain in Oregon, USA, (Ambrose et al., 2011), a Mountain site at Lulang on the southeast Tibetan Plateau, China (Zhao et al., 2013) have reported the importance of the aerosol chemical composition at mountain sites in the identification of potential source regions of anthropogenic pollutants and their mechanism of transport. Measurement and simulation at sSome observations and somemodels have elaborated the emissions forof some trace gases such as CO and O<sub>3</sub> from the boundary layer into the free

Code de champ modifié

a mis en forme : Français

a mis en forme : Français

troposphere by convective, frontal, and orographic lifting at mountain sites (Bey et al., 2001; Ding et al., 2015; Liang et al., 2004; Weiss-Penzias et al., 2006). Nevertheless, the effects of these mechanisms have rarely been studied at the aerosol chemical composition scale. Despite the increasing interest in high-altitude aerosol research, most studies [have reported measurements of](#) tracer gases over [the long-term periods](#), but limited studies have addressed the interaction between natural emissions such as mineral dust, biogenic compounds, and anthropogenic emissions in the free troposphere ([Fiore et al. 2009; Jonson et al. 2010; Logan et al. 2012; Gilge et al. 2010; Kumar et al. 2013](#)).

Moreover, such studies have been reported mostly in central Europe, Asia, and North America ([Okamoto and Tanimoto, 2016](#)). A few attempts in Africa have been made to investigate the [microphysical microphysical and optical properties](#) of mineral dust transport in North-Africa ([Kandler et al. 2009; Müller et al. 2012; Ryder et al. 2011; Schladitz et al. 2009; Veselovskii et al. 2016](#)). Other studies have focused on polluted regions (Benchrif et al., 2018; Inchaouh, 2017; Tahri et al., 2013), such that information about the chemical composition of particulate matter at high-altitudes is limited. Likewise, background aerosol information, essential in assessing long-term regional changes in atmospheric composition in this region remains scarce and difficult to assess due to lack of the necessary infrastructure. This poor state of knowledge limits transregional investigation on the effect of different sources and source regions on the chemical composition of aerosol particles over sensitive regions such as the Atlas Mountains in North Africa.

The Middle-Atlas region located in the North of Morocco is typically considered as an area with high rainfall. Still, according to the report from the Moroccan ministry of environment, the annual average rainfall decreased by about 100 mm, with an increase in temperature by about 1.5 °C (Royaume du Maroc, 2009) within the past 50 years. These are indicators of the sensitive nature of the Middle-Atlas region to a changing climate. The Ifrane national park, which is located in the Middle-Atlas, suffers from intense pressure due to forest degradation, and overgrazing resulting in significant climatic consequences (Campbell et al., 2017). At the same time, it is classified as a site of biological and ecological interest. Recently, soil erosion was particularly intense in some clay-dominated valleys (Mounir et al., 2019). This change can lead to several consequences such as an increase in aridity, reduction of precipitation, change in primary emissions, and atmospheric composition. Thus, the observations from this region could provide new knowledge into atmospheric composition changes over time, related to different climatic and anthropogenic dynamics. The evaluation of local regional, trans-regional, and climate change effects can help assess air quality and climate-relevant mitigation strategies.

The aim of this study was, therefore, to i) quantify and characterize the variability of PM<sub>10</sub> mass concentration in the high-altitudes of the Middle-Atlas region, ii) determine their chemical composition, iii) identify the possible sources of the aerosol particles, and iv) evaluate the relative contributions of the source regions to the observed concentrations. Within the present study, chemical composition at the high altitude AM5 observatory is presented. Chemical components such as trace metals, OC, EC, ionic, and organics species were investigated, and meteorological and back trajectory analysis was performed. Moreover, the influence of dust on the chemical composition and the day and night variation of the PM<sub>10</sub> concentration were investigated.

## 2 Experimental

### 115 2.1 Site description and particles sampling

The Atlas Mohammed V (AM5) atmospheric research station situated in a strategic location in the Middle Atlas was founded in 2017. It is operated by the Centre National de la Recherche Scientifique (CNRS-ICARE, Orléans-France), the Mohammed V University (Rabat-Morocco), and the Leibniz Institute for Tropospheric Research (TROPOS, Leipzig-Germany). The observatory is located at Michlifen in the Middle Atlas region at an altitude of 2100 m a.s.l in a remote hilly site (33°24'22.2"N 5°06'12.0"W) ~~characterized by its strategic location~~. On one side, ~~are~~ the plains and plateaus of central Atlantic Morocco, ~~and~~ on the other side, the arid areas. It is about 300 km north of the ~~Sahara desert~~ Sahara desert, about 230 km east of the Atlantic Ocean and the populated and industrial regions of Casablanca and Rabat, and about 340 km south of the Mediterranean Sea. The orientation of the Middle Atlas Mountains diagonally extends from southwest to northeast over a distance of 450 km. The AM5 station is surrounded by cedar forests and pastureland that come to life in spring and summer, which is a sharp contrast to the hot, dry climate surrounding it. The nearest urban towns are Ifrane, and Azrou, which ~~is-are~~ about 22 km away, and Fes city located 82 km north of the AM5 station. Due to its remote location, the influence of anthropogenic emissions from the Ifrane area is low.

130 Aerosol particles were sampled using a PM<sub>10</sub> high volume Digitel (DHA-80, Switzerland) with a flow rate of 500 l/min on quartz fiber filters (Munktell, MK 360). The collection period was from August to December 2017, during which 200 filters have been collected in a day and night-time (12 h) sampling routine. At the end of the sampling, the collected filters were placed in a refrigerator at a temperature of 5°C and subsequently frozen at -20°C. After storage, the filters were transported at -20°C to TROPOS (Leipzig-Germany) in aluminum cans for 135 chemical analysis. In September due to power failure and instrument outage fewer samples were collected (n=20) Due to the lower number of samples, ~~the missing days of sampling were replaced by the average concentration of the collected samples in September.~~

### 2.2 Local meteorology and station characteristics

Meteorological data were collected from summer 2017 to spring 2018 for major parameters such as temperature, 140 relative humidity, wind speed and direction, atmospheric pressure, visibility, and precipitation at a sampling rate of 1 min using an automated weather station (Bresser AWS, Germany). As shown in Table 1, in summer, temperatures are moderate or warm during the day, cool at night. Winter is much colder, and the daily ~~temperature~~ amplitudes are lower ~~because t-~~ The valleys only receive the sun's rays in the middle of the day. In winter, the 145 station remains entirely in the shade for several weeks. Thermal contrasts between slopes are important when the topographic terrain is oriented east-west. ~~In the valleys in-during~~ high-pressure weather, thermal breezes are common. The temperature varies seasonally, especially during the transition from summer to winter, with maximum and minimum values of 26°C and -1°C, respectively. The annual average temperature is approximately 14°C, with a sharp decrease during the night.

150 In contrast, the visibility varies slightly, with intense UV radiation during summer when the sky is often clear. Fog occurrence is high during autumn and winter. The wind comes from all directions, but it is dominated by air

mass from the west, as shown in Fig. 2. The average wind speed at AM5 was about ~~21.5.8 km/h~~<sup>1</sup> but reached a maximum of ~~19.74 km/h~~<sup>1</sup> due to turbulence in the mountain region, especially during winter. Over the summer, the minimum wind speed was about ~~1.66 m s<sup>-1</sup>~~<sup>1</sup>, and the relative humidity (RH) was low. In autumn, RH was on average about 36% and reached up to 97% under the influence of marine air mass in winter. The Middle-Atlas region is considered a very humid and temperate climate. Indeed, the water balance required for plants is mainly positive in winter, about 141 mm, while annual precipitation is 300 mm. Rainfall occurs mainly during winter, with heavy thunderstorms and a lot of snow coverage. The northern part of the Middle Atlas Mountains is the wettest region in Morocco after the Rif mountain regions, according to Nourelbait et al. (2016). Precipitation increases in frequency and intensity during the winter. Indeed, the mountain imposes an ascent of air masses, which results in cooling, the formation of clouds, and the condensation of water vapor. The proportion of snowfall also increases rapidly because of the altitude, especially in winter. On the other hand, no difference was observed for the visibility during the seasons which shows an average value of 10 km.

a mis en forme : Exposant

155 RH was on average about 36% and reached up to 97% under the influence of marine air mass in winter. The Middle-Atlas region is considered a very humid and temperate climate. Indeed, the water balance required for plants is mainly positive in winter, about 141 mm, while annual precipitation is 300 mm. Rainfall occurs mainly during winter, with heavy thunderstorms and a lot of snow coverage. The northern part of the Middle Atlas Mountains is the wettest region in Morocco after the Rif mountain regions, according to Nourelbait et al. (2016).  
160 Precipitation increases in frequency and intensity during the winter. Indeed, the mountain imposes an ascent of air masses, which results in cooling, the formation of clouds, and the condensation of water vapor. The proportion of snowfall also increases rapidly because of the altitude, especially in winter. On the other hand, no difference was observed for the visibility during the seasons which shows an average value of 10 km.

165 There is a wide variation in local wind distribution during the summer over seasons season where, and the wind comes from all directions, except from the north. A predominance of southwestern winds distinguishes it. On the one hand, During summer, southeast winds have a higher frequency but a lower average speed of about 6.5 m s<sup>-1</sup>. Similarly, southeast winds are dominant with a slight decrease in wind frequency during the fall season. While western winds are dominant by up to 30% during winter and spring. During this period, there is a strong occurrence of westerly winds which are often characterized by high wind speeds (stiff breeze) of up to 20 m s<sup>-1</sup>. To conclude, the wind frequently comes from the west and southwest during the winter months, in contrast to the dominated south winds in summer.

a mis en forme : Non Surlignage

a mis en forme : Couleur de police : Automatique

### 2.3 Aerosol particle chemical analysis

#### Particle mass

175 The collected filters were weighed on a microbalance with an accuracy of 10 µg (Mod. AT261 Delta Range, Mettler) after being stabilized for 72 hours at constant temperature ( $20\pm1^{\circ}\text{C}$ ) and humidity ( $50\pm5\%$ ), before and after sampling. The difference between the weights was determined and divided by the total sampling volume to obtain the mass concentrations. After the determination of the particulate matter concentration, both organic and inorganic analyses were carried out at the TROPOS laboratories.

180

#### Carbon compounds

Organic and elemental carbon were analyzed by a thermo-optical method (Sunset Laboratory Inc. U.S.A) at a maximum Temperature of  $850^{\circ}\text{C}$  with the normalized temperature program EUSAAR2 (EUropean Supersites for Atmospheric Aerosol Research), as described in the literature (Cavalli et al., 2010; Yttri et al., 2019). The method is in line with the standard proposed by the European networks (ACTRIS, EMEP). Samples were thermally desorbed from the filter medium under an inert He-atmosphere followed by an oxidizing O<sub>2</sub>/He-atmosphere using carefully controlled heating ramps. A flame ionization detector is used to quantify methane after catalytic methanation of CO<sub>2</sub>. First, the sample is heated up to about  $870^{\circ}\text{C}$  in an inert atmosphere of pure helium. These conditions allow the organic carbon to volatilize and to be fed into the second furnace filled with Mn<sub>2</sub>O (oxidation catalyst), where it is quantitatively oxidized into CO<sub>2</sub>. As a second step, the sample is placed in an oxidizing

atmosphere (helium/oxygen), leading to the oxidation and volatilization of the refractory elemental carbon remaining on the filter (van Pinxteren et al., 2015). Charring processes lead to the overestimation of EC and an underestimation of OC, resulting in lower OC/EC ratios. Therefore, an optical correction was applied for the charring processes.

195 The correction value optical correction of charring for pyrolytic carbon is obtained by measuring the transmission of the sample with a laser (wavelength 678 nm). The limit of detection for OC/EC measurement was 0.2 µg/cm<sup>2</sup>. Organic matter (OM) was estimated based on the  $f_{OM/OC}$  conversion factor, according to Turpin and Lim (2001). The total carbon TC was considered as the sum of organic carbon and elemental carbon (TC=OC+EC). Organic matter was considered to be was about twice the organic carbon (OM = 2.1 \*OC). Since the conversion factor depends on the specific proportion of each site, the factor  $f_{OM/OC} = 2.1$  is suggested because it takes into account aged aerosols (Turpin and Lim, 2001). The primary organic carbon (POAC?) fraction in the PM<sub>10</sub> was estimated using EC as a tracer, by taking the minimum OC/EC ratio for the entire study period and multiplying it to the EC content, as in the following equation POC = (OC/EC)<sub>min</sub> × EC. Consequently, the secondary organic carbon (SOC) contribution to the total OC can be estimated as the difference between the total organic carbon and total primary organic carbon concentrations (SOC = OC - POC).

200  
205  
210  
Organic compounds such as n-alkanes oxygenated polycyclic aromatic hydrocarbons (oxy-PAHs), and n-alkane-2-onesketones were detected using a Curie-point pyrolyzer (JPS-350, JAI Inc., Japan) coupled with a GC-MS system (6890 N GC, 5973 inert MSD, Agilent Technologies, CA, USA) as described by Neusüss et al., (2000). Saccharidic compounds such as mannitol, glucose, levoglucosan, and arabitol were determined using high-performance anion-exchange chromatography with pulsed amperometric detection (HPAEC-PAD) as described by Inuma et al., 2009.

#### Trace metals

215 Trace metals were determined using the Total Reflection X-Ray Fluorescence technique (TXRF), whereby 3 spots of 8 mm in diameter each were digested in 1.125 mL HNO<sub>3</sub> and 0.375 HCl using the Mars 6 (CEM, Germany) microwave. 50 µL of the digested solution was deposited on previously siliconized quartz carriers, and 10 ng of Gallium was added onto the sample as an internal standard. The samples were subsequently measured using an S2-PICOFOX (Bruker AXS Microanalysis GmbH, Germany) instrument. Further details of the technique and measurement procedure have been reported elsewhere (Fomba et al., 2020).

220

#### Water-soluble ions

225 The major ionic constituents were analyzed using a standard ion chromatography technique (ICS3000, Dionex, USA) equipped with automatic eluent generation (KOH for anions and methanesulfonic acid (MSA) for cations) and a micro membrane removal unit. Ion analysis was performed for Na<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>, Ca<sup>2+</sup> cations, and Cl<sup>-</sup> and Br<sup>-</sup>, NO<sup>-</sup>, SO<sub>4</sub><sup>2-</sup> and C<sub>2</sub>O<sub>4</sub><sup>2-</sup> anions. For these analyses, 3 spots of 2 cm each in diameter were extracted from the filter in deionized water via shaking for 2 h. The extract was filtered through a 0.45 µm unidirectional syringe filter to remove insoluble matter, and the filtrate was analyzed. The blank field filters were analyzed using similar procedures and were subtracted from the sample concentrations following the methodology described by Inuma et al. (2009).

230

### *Estimation of Sea salt components*

Sea salt concentrations were calculated by adding chloride to sodium, and the sea salt (ss) contributions of potassium, calcium, magnesium, and sulfate, where ss-K<sup>+</sup>, ss-Ca<sup>2+</sup>, ss-Mg<sup>2+</sup>, and ss-SO<sub>4</sub><sup>2-</sup>, respectively, have been estimated as 0.03, 0.5, 0.12, and 0.25 fractions of the measured Na<sup>+</sup>, respectively (Marenco et al., 2006). However, the estimation of sea salt assumes that all Na and Cl are coming from marine contribution without taking into consideration other potential sources. The estimation of non-sea salt sulfate (nss-SO<sub>4</sub><sup>2-</sup>) was estimated by subtracting the contribution of ss-SO<sub>4</sub><sup>2-</sup> from the total SO<sub>4</sub><sup>2-</sup> mass concentration (Amadio et al., 2014). The water content of the samples was estimated according to the E-AIM (Extended Aerosol Inorganics Model) III of Clegg et al. (1998).

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Normal, Gauche

### **2.4 Determination of mineral dust**

Mineral dust (MD) is a significant contributor to atmospheric particulate matter especially in North-Africa, where the average percentage can vary from 7% to 62% (Gherboudj et al., 2017). The high spatiotemporal variability of the dust emission can sometimes be difficult to quantify correctly and may lead to some uncertainty. Therefore, the estimation of MD can be subjective because several estimation methods are available in the literature. The most common methods were applied to the samples collected at AM5 station. Within the present study, the aim was first to evaluate and then select an appropriate method for the interpretation of the results. Four methods were highlighted that are representative of those used in the literature. The first method implemented by Fomba et al., (2014) consists of subtracting the total PM<sub>10</sub> mass concentration from the analyzed mass of the other elements, representing the upper limit of the possible MD concentration in the samples. This method is an interesting approach especially when the elemental analysis is not available. However, to improve on the MD quantification using available MD-related elements, applying a stoichiometric equation reduces the uncertainty of the values. Using this method, an average MD concentration of about 24.5  $\mu\text{g m}^{-3}$  was obtained. In contrast, methods 2 and 3 use different approaches. They estimate MD base on given stoichiometry and apply different elemental concentrations such as Al or Ca, Fe, Ti to estimate the MD load. Method 2 uses the factor (1.16) to compensate for the exclusion of MgO, Na<sub>2</sub>O, K<sub>2</sub>O, and H<sub>2</sub>O from the crustal mass calculation, as shown in Table 2 (Maenhaut et al., 2005). Whereas method 3 considers carbonate such as calcite, dolomite, and other oxides such as TiO<sub>2</sub>, Fe<sub>2</sub>O<sub>3</sub>, and MnO<sub>2</sub> (Minguillón et al., 2007; Nerriere et al., 2007). As a result, the average MD concentration using methods 2 and 3 were similar, about 19.9 and 18.9  $\mu\text{g m}^{-3}$ , respectively. Method 4 takes into account that sea salt significantly affects these concentrations, and the non-sea salt (nss) content of the elements, such as nss-Ca<sup>2+</sup> or nss-Mg<sup>2+</sup> are used to replace the total Ca and Mg concentrations. This is more accurate in a sea-salt-dominated environment but could underestimate the calcium contribution as sodium also has a crustal origin (Cesari et al., 2012; Perrino et al., 2014). The average MD concentration from this method was about 15.5  $\mu\text{g m}^{-3}$ .

In conclusion, differences of up to 37% were obtained between these methods. The obtained MD concentrations were high for method 1 and lowest for method 4 and similar between methods 3 and 4, indicating their robustness. Thus, this approach was applied in this study to estimate the MD content. Method 3 does not consider quantifying silicates SiO<sub>2</sub> and aluminosilicate Al<sub>2</sub>SiO<sub>5</sub>, a major component in natural mineral dust. While However, method 2 allows to take into account the overall mineral composition and therefore was applied in this study. Due to

270 limitations in quantifying Si from quartz fiber, MD was finally estimated by replacing "Si" with "Ca" base on the established soil stoichiometric ratio of the average upper continental crust (Si=10.3 Ca), according to Wedepohl (1995) in the equation used in Method 2. The final equation used for mineral dust estimation is given as follows:

275 
$$\text{Mineral dust} = 1.16(1.90 \text{ Al} + 23.3 \text{ Ca} + 2.09 \text{ Fe} + 1.67 \text{ Ti}), \quad (1)$$

## 2.5 Back trajectory analysis

The origin of the air masses reaching the station, 96h back trajectories were estimated using the NOAA HYSPLIT (Hybrid Single-Particle Lagrangian Trajectory, HYSPLIT 4) model (<http://www.ready.noaa.gov/ready/hysplit4.html>; Draxler and Hess, 2004), using the 1° resolution Global Data Assimilation System (GDAS) input data. Due to the resolution of the input data, the exact altitude of the mountain is not properly represented and according to the HYSPLIT model at the AM5 site, the terrain height is at 1000 m only. Therefore, trajectories were calculated every hour for the altitude of 1000 m above model ground level. The profile of the air mass altitude during transport was applied for the interpretation of the data for each air mass.

285 In order To group the back trajectories into distinct transport patterns, a manual classification approach has been used. This method consists of grouping 12-hourly ensemble trajectories with the time interval between adjacent nodes of 14 h and calculated for 96 h, and attributing them to a specific air mass category. The assignment of each pathway trajectories was based on their crossing over given latitude-longitude grids attributed to given geographical sectors. To assign a sample to a specific air mass category, 60% of the trajectories must have a similar profile. Some samples (n=15 samples) have with trajectories from mixed origins (e.g., marine air mass over Europe and the desert) have been excluded from the classification of the air masses in this study. In total, the air masses of 175 samples could be grouped into related to the four distinctive air masses will categories which were Atlantic Ocean Background Air mass (BAMAOB), Atlantic Coast Europe (ACE), Mediterranean Coast Europe (MCE), Saharan Dust (SD).

## 3 Results and discussion

### 3.1 Variation of PM<sub>10</sub> mass

The PM<sub>10</sub> mass concentration time series at the AM5 station varied from 9.5 µg m<sup>-3</sup> to 145.6 µg m<sup>-3</sup> with averaged of 29.2 ± 17.3 µg m<sup>-3</sup>, as shown in Fig. 3a. The PM<sub>10</sub> mass shows a strong seasonal variation during the five months of measurement from August to December 2017, as illustrated in Fig. 3a. The highest mean monthly concentration was observed in August (49.9 ± 25.9 µg m<sup>-3</sup>) and continuously decreased until December (15.9 ± 5.6 µg m<sup>-3</sup>), as shown in figure Fig. 3b. The observed temporal variation in the concentration is related most likely to factors such as meteorological conditions and the air mass arriving at the station. To understand the seasonal variation of the particulate matter load, PM<sub>10</sub> mass wereas, combined with wind speed and wind direction to create the polar plots as presented in Fig. 3c, which illustrates the variation of PM<sub>10</sub> concentrations as a function of wind speed and direction.

During August, the high PM<sub>10</sub> concentrations were mostly related to high wind speeds from the southeast-wind direction. For example, PM<sub>10</sub> mass concentration often exceeded 50 µg m<sup>-3</sup> and sometimes even reached up to 145

a mis en forme

[1]

310 µg m<sup>-3</sup> during August, when the wind speed was stronger than 9 m s<sup>-1</sup>. The high PM<sub>10</sub> concentration recorded was due to the influence of Saharan dust events during periods of air mass influence from the southern sector located in the southeast of the AM5 station, as indicated in Figure 3e. The Middle Atlas region is marked by particular meteorological conditions during the summer with the low humidity and often low precipitation (avg. 37 mm), as shown in Table 1. These hot and arid conditions are known to favor the transport of dust particles from the Saharan

315 desert to the Atlas Mountains (Rodríguez et al., 2011). Furthermore, high concentrations of up to 40-50 µg m<sup>-3</sup> were observed as well with westerly winds, especially during northwest and southwest winds. High PM<sub>10</sub>

a mis en forme

[2]

320 concentrations were observed during strong westerly winds of up to > 7 m s<sup>-1</sup>. The back trajectory analysis suggests that the high concentrations during this period were most likely associated with the long-range transport of aerosol particles from the western coast of the Iberian Peninsula. In contrast to the summer

325 period, PM<sub>10</sub> mass concentrations were lower during the fall period, despite some temporal peaks. The PM<sub>10</sub> concentrations were generally lower in September ( $24.2 \pm 5.1 \mu\text{g m}^{-3}$ ) and October ( $30.5 \pm 10.8 \mu\text{g m}^{-3}$ ). During this period, winds originated from the northeast clockwise suggesting the influence of air mass transport from the Mediterranean Sea coast. A sharp fall in PM<sub>10</sub> concentrations was noticed in November ( $22.8 \pm 7.9 \mu\text{g m}^{-3}$ ) and December ( $15.9 \pm 5.6 \mu\text{g m}^{-3}$ ). Overall, PM<sub>10</sub> concentration decreased from the summer to winter period by 32%.

330 This trend is most likely due to the increased amount of precipitation (peaks of 852 mm) during fall and winter, which can lead to the wash-out effect of aerosol and its components (Holst et al., 2008).

To establish a reference baseline and evaluate the background conditions at the site, the lower 5<sup>th</sup> percentile from all of our data points (PM<sub>10</sub> < 12 µg m<sup>-3</sup>) was found to be representative of remote

a mis en forme

[3]

335 background aerosol conditions. The PM<sub>10</sub> frequency and probability density function as shown in Fig. S3 XX confirmed this observation. The samples within this PM concentration range had similar air mass trajectories and typical meteorological conditions with low wind speeds < 3 m s<sup>-1</sup>. The air masses typically traveled in the free troposphere at about 1000 m above sea level, crossing the North-Atlantic Ocean before arriving at the site within the past 96 h. These conditions were, however, not free from local and regional pollution from point sources such as dust resuspension from the cars assessing the site.

a mis en forme

[4]

340 Consequently, the average background PM<sub>10</sub> mass concentration at the AM5 was  $10.9 \mu\text{g m}^{-3}$ , which was found to be stable and representative of periods of little external influence. In comparison, Benchrif et al. (2018) reported background PM<sub>10</sub> values for Northern Morocco with an average of  $12.2 \mu\text{g m}^{-3}$ , which is very similar to the concentrations determined in this study,  $10.9 \mu\text{g m}^{-3}$ .

345 The mean concentration recorded at AM5 from this study ( $29.2 \pm 17.3 \mu\text{g m}^{-3}$ ) agreed well with the PM<sub>10</sub> concentration of other remote high-altitude sites, such as Darjeeling in Northeastern Himalayas ( $29 \mu\text{g m}^{-3}$ ; Chatterjee et al., 2010), Lhasa in Tibet ( $37 \mu\text{g m}^{-3}$ ; Wang et al., 2015), and Mahabaleshwar in India ( $37 \mu\text{g m}^{-3}$ ; Leena et al., 2017) as presented in Table 3. Other high-altitude stations, such as Izaña, in Canary Islands ( $46 \mu\text{g m}^{-3}$ ) showed much higher PM<sub>10</sub>, most likely due to the exposure to strong Saharan dust events (García et al., 2017). In contrast, the PM<sub>10</sub> concentrations at AM5 were considerably higher than the PM<sub>10</sub> levels recorded in

European and Asian high-altitude sites. For example, the average~~average or maximum~~ PM<sub>10</sub> mean value recorded in this study was about factor 2 greater twice that of in Mount Cimone, Italy (16  $\mu\text{g m}^{-3}$ ; (Marenco et al., 2006) and factor 6 greater than the PM<sub>10</sub> in Everest Mountain (Decesari et al., 2010) as well as and in Puy de Dôme, France (6  $\mu\text{g m}^{-3}$ ; Bourcier et al. 2012), and, ~~approxima~~ was also approximately 10 times greater than the average level in Jungfraujoch (3  $\mu\text{g m}^{-3}$ ; Cozic et al., 2008). Other Moroccan sites, such as Marrakech, Meknes, and Agadir, which are exposed to strong urban emissions, usually show PM<sub>10</sub> concentrations between 50 and 110  $\mu\text{g m}^{-3}$ , which is much higher than the concentration found at AM5 in this study (Inchaouh, 2017; Tahri et al., 2013, 2017). These results highlight the cleanlinesse better air quality at AM5 in comparison to many sites ~~of the air and indicate~~<sup>on</sup>, that the AM5 station apparently can serves as a good remote reference station for defining background concentrations in Morocco and possibly the whole of North Africa.

### 3.13.2 Air mass origins

The calculation of back trajectories using the Hysplit model allowed the identification of several remote sources of PM<sub>10</sub> at the station. Four main<sup>s</sup> air masses were classified into four categories were identified<sup>s</sup>, as shown in Fig. 54. i) Air masses that The background air mass which spent the last 96h over the Atlantic Ocean -at high-altitude (1000 m.asl), representative of typical background air mass (BAM) conditions, which influenced about during transport represents 5.3% of all samples; ii) The Atlantic a Air masses originating<sup>ed</sup> from the Atlantic and crossing over the Coast of Europe<sup>an</sup> (ACE), especially Spain and over Moroccan industrial cities located at the North Atlantic coast, located about 426 km north influencing about accounts for 26.8% of all samples; iii) The ACE-a Air masses from Europe crossing over the influence of the Mediterranean Coast and Europe (MCE) as well as over crossing Coastal area in North Moroccan cities, as shown in Fig. 4c~~x~~ and influencing o-was higher (about 37.4% of all samples and iv) in comparison to the ACE air mass. Air masses ,which originating<sup>ed</sup> from Southern and/or Eastern Sahara crossing the desert (SD), at different altitudes before arriving at the AM5 and influencing about accounted for 22.6% of all samples. The back trajectories represent mixing scenarios (7.9% of all samples) were not assigned to any of the four major classes as mentioned above.

### 3.23.3 Characterization of aerosol chemical composition

The statistics results of the measured PM<sub>10</sub> chemical components during the four air mass categories are shown in Table 4, including the average concentrations and their<sup>s</sup> and variations (-standard deviation (Std). Likewise, Fig ure 6 shows the time series of the investigated chemical species investigated within the sampling period colored with periods of the different air mass influence.

#### 3.3.1 Mineral dust

During the 5 months of PM collection at the AM5 site, the average of mineral dust concentration was about 17.7  $\pm 7.4 \mu\text{g m}^{-3}$  and varied strongly between 0.05  $\mu\text{g m}^{-3}$  and 107  $\mu\text{g m}^{-3}$ . The highest mean concentrations were observed in August (39  $\mu\text{g m}^{-3}$ ) and the lowest in December (3.7  $\mu\text{g m}^{-3}$ ). Low concentrations were observed during days with low wind speeds (< 2 m s<sup>-1</sup>), low<sup>e</sup> Saharan dust air mass inflow, and after precipitation events, which typically occurred in the fall and winter. The influence of the Saharan dust on the Middle Atlas region remains relatively dependent on meteorological conditions. Firstly, the direction and speed of the wind, as theThe transport of Saharan dust was dependent on wind direction. Typically Saharan dust events winds were observed

a mis en forme : Couleur de police : Texte 1, Non Surlignage

a mis en forme : Police :Non Italique

a mis en forme : Indice

a mis en forme : Anglais (E.U.)

a mis en forme : Normal, Sans numérotation ni puces

a mis en forme : Police :Couleur de police : Noir, Bordure :: (Pas de bordure)

a mis en forme : Couleur de police : Accent 1

a mis en forme : Normal, Gauche

a mis en forme : Exposant

385 during low wind speed periods from south and southeast. Secondly, their progression depends mainly on  
favorable weather conditions for transport, the difference in temperature between day and night, humidity, and  
especially the scarcity of rainfall. Thirdly, The low wind speed from the south can be explained by the high altitude  
(4000 m) of the High-Atlas mountains situated at 4000 m of altitude acts as a barrier to Saharan dust transport  
which forces the winds to deviate from their path which could lead to low wind speed and acts as a barrier to  
Saharan dust transport. All these factors influence the transport of large particles from the Sahara to the Middle-  
Atlas during the different seasons.

390  
395 Nevertheless, even during days of low wind speed, mineral dust still dominated the background aerosol  
composition and contributed up to 51% of the total mass, as shown in Fig. S4, for typical background conditions  
chemical composition. The highest aerosol mass was observed during days of Saharan dust events from the 10<sup>th</sup>  
to the 13<sup>th</sup> of August when air mass crossed the Sahara (SD) prior to their arrival before arriving at the AM5 site.  
The duration of Saharan dust events varied from 1 to 3 days, with the longest event (also supported by back  
400 trajectory analysis) also observed during this event August. Most dust events occurred during the summer, as  
indicated by the strong increase of the typical crustal elements (Guimot et al. 2007; Arimoto et al., 2006) such as  
nss-Ca<sup>2+</sup>, Fe, and -(Fig. 622) a of mineral dust, which is present on average at a concentration of  $5.5 \pm 3.5 \mu\text{g}$   
 $\text{m}^{-3}$  during remote background conditions, was found to be more than 7 times higher ( $37.9 \pm 25.3 \mu\text{g m}^{-3}$ ) during  
dust events (SD), in comparison to remote background conditions (BAM) with an average concentration of  $5.5 \pm$   
405  $3.5 \mu\text{g m}^{-3}$ , as observed in Table 4. Other less intense Saharan dust storms occurred during the summer season  
between the 21<sup>st</sup> and 24<sup>th</sup> of August with a similar impact where high dust concentrations that was 5 times higher  
as than dust background dust concentrations. The duration of Saharan dust events varied from 1 to 3 days,  
with the longest event also supported by back trajectory analysis observed from 10<sup>th</sup> to the 13<sup>th</sup> of August. The  
presence of mineral dust is relatively low but still significant for air masses other than SD, such as during the  
410 ACE ( $13.3 \pm 5.2 \mu\text{g m}^{-3}$ ) and MCE ( $19.9 \pm 11.9 \mu\text{g m}^{-3}$ ) air mass influences, as shown in Table 4. This suggests  
that the air masses were very often loaded with mineral dust originating from regional sources in the Middle Atlas.  
The Fe/Ca ratio was used to distinguish between different mineral dust sources. While Fe/Ca ratios close to 0.4  
415 indicate dust from the south, while Fe/Ca ratios greater than 1 indicate dust from east. The Fe/Ca ratio is about  
0.5 during BAM conditions, indicating local sources emitted by road dust or resuspension of agricultural activities.  
While Fe/Ca ratios close to 0.4 indicate dust from the south, Fe/Ca ratios greater than 1 indicate dust from east.  
Mineral dust can be transported over long distances particularly from the North African source region to the  
420 Mediterranean basin and Europe (Schepanski et al., 2016). Table 5. For instance, the high-altitude site in Mt.  
Cimone, Italy recorded several days with African dust transport which influenced the chemical composition  
(Marenco et al., 2006). However, the Saharan dust concentration at AM5 ( $17.7 \mu\text{g m}^{-3}$ ) is approximately 4 times  
higher than at Mt. Cimone ( $4 \mu\text{g m}^{-3}$ ). Nevertheless, the average concentrations of elements, such as Al, Fe, Ti,  
and Mn, are comparable with the values reported in Mt. Cimone, Italy (Marenco et al., 2006). Table 5. However,  
the calcium concentration at station the AM5 ( $0.65 \pm 0.58 \mu\text{g m}^{-3}$ ), was 2 times higher than the concentration  
recorded in Mt. Cimone. Furthermore, the calcium concentration was 5 times higher than the concentration  
recorded at other high-altitude station such as Mt. Himalaya and Mt. Everest (Chatterjee et al. 2010; d; De Cesari

a mis en forme : Couleur de police : Texte 1  
a mis en forme : Couleur de police : Texte 1  
a mis en forme : Couleur de police : Texte 1  
a mis en forme : Couleur de police : Texte 1

425 et al., 2010). Although, and than This suggests that the AM5 experiences higher amounts of calcium-rich dust in comparison to other sites. Some studies have reported the high content in calcite in the soils of Northern Morocco  $1.07 \text{ ng m}^{-3}$  which confirms the predominance of calcium-rich in the Atlas regions (Desboeufs and Cautenet, 2005; Kandler et al., 2009; Benchrif et al., 2018).

a mis en forme : Anglais (E.U.)

### 3.3.2 OM-OC and EC

430 Organic carbon (OC) and elemental carbon (EC) showed strong variation and a distinct seasonal variation differences with an average of  $1.1 \pm 0.8 \mu\text{g m}^{-3}$  and  $0.25 \pm 0.13 \mu\text{g m}^{-3}$ , respectively, and strong seasonal variation. The OC has both primary and secondary origin, and can be formed from primarily emitted substances through condensation or chemical reactions among them (Sarkar et al., 2019). The OC concentration reached a maximum  $4.5 \mu\text{g m}^{-3}$  during summer, whereas the lowest concentration was recorded observed during winter at about  $0.03 \mu\text{g m}^{-3}$ . The average concentration of OC progressively decreased from summer ( $2.1 \pm 0.89 \mu\text{g m}^{-3}$ ) until to the 435 winter ( $0.34 \pm 0.26 \mu\text{g m}^{-3}$ ). The abundant contribution of organic matter in summer was more abundant in summer can be due to high biogenic emissions in the Middle-Atlas. However, a slight increase in OC was also observed during dust events, as shown in Fig. 5, which suggests that the dust deposited at AM5 also contained has often been loaded with biogenic material from the surrounding-s of the Middle-Atlas region. The average 440 concentration of POC was found to be  $0.27 \pm 0.35 \mu\text{g m}^{-3}$  (26% of OC concentration), whereas SOC was estimated as  $0.83 \pm 0.75 \mu\text{g m}^{-3}$  (74% of OC).

a mis en forme : Exposant

445 The EC concentration showed little temporal variation except for a few pollution episodes during which could be observed during the sampling period, as shown by the peaks in the EC concentrations could be observed (in Fig. 5). The sudden increase in EC observed during summer ( $0.6 \mu\text{g m}^{-3}$ ) and autumn ( $0.7 \mu\text{g m}^{-3}$ ) were characterized by two different sources. Firstly, from Europe through the Atlantic Coast during ACE long-range transport air mass influence during in summer and. Secondly, from nearby urban regions especially in the evenings when the temperature is low (with an average of  $5^\circ\text{C}$ ) and high values of anthropogenic metals, such as  $\text{Pb}$ ,  $\text{Cu}$ , and  $\text{Ni}$ , observed. The wind direction and back trajectory analysis indicates that the most likely sources 450 of pollution are the urban cities of Fes and Meknes, located about 85 and 50 km in the North from the station. However, the winter period was marked by low EC concentrations, with an average of  $0.1 \pm 0.06 \mu\text{g m}^{-3}$ . The elemental carbon concentration EC concentrations ( $0.24 \mu\text{g m}^{-3}$ ) during Saharan dust air masses were not significantly low indicating that the influenced of by urban pollution during dust events was low, as shown by the unchanging elemental carbon concentration EC concentrations ( $0.21 \mu\text{g m}^{-3}$ ) associated with their period of influence at the AM5 site. Few studies observed that the transport of North African dust was often loaded with pollutants (Gangoiti et al., 2006; Kalderon-Asael et al., 2009; Astitha et al., 2010). Nevertheless, according to the 455 back trajectories, the transport of mineral dust takes place directly from the Sahara Desert situated in the south of the AM5 site, without passing through cities with intense anthropogenic activities. The study of Decesari et al. 460 2010 reported similar concentrations of OC ( $0.8 \pm 0.6 \mu\text{g m}^{-3}$ ) in  $\text{PM}_{10}$  at the high altitude Himalayan station in Nepal, whereas and lower EC ( $0.11 \pm 0.13 \mu\text{g m}^{-3}$ ) concentrations in  $\text{PM}_{10}$  at the Himalayan high-altitude Himalayan station in Nepal have been found. Furthermore, Sharma et al., 2020 reported higher OC ( $5.4 \pm 2.0 \mu\text{g}$

a mis en forme : Non Surlignage

a mis en forme : Couleur de police : Texte 1

Code de champ modifié

a mis en forme : Couleur de police : Texte 1, Français

a mis en forme : Couleur de police : Texte 1

$\text{m}^{-3}$ ) and EC ( $2.2 \pm 2.0 \mu\text{g m}^{-3}$ ) at the ~~high altitude~~ high-altitude site of Darjeeling, India most likely due to the higher influence of anthropogenic activities at the site.

The OC/EC ratio contributes to assessing the ageing of aerosols during long-range transport as well as the impact of the combustion source producing EC. The average OC/EC ratio was computed as  $4.8 \pm 5.4 \mu\text{g}$  with a range of  $0.22\text{--}56.1$ . Conversely, the OC/EC ratio tends to ~~decrease~~ increase from summer ( $11.2 \pm 9.7$ ) to winter to ( $2.2 \pm 1.4$ ). The decreasing trend of OC/EC ratio can be due to the formation of secondary organic aerosols in summer by photochemical processes, as shown by the high secondary organic carbon (SOC) content observed during August ( $1.9 \pm 0.89 \mu\text{g m}^{-3}$ ) in contrast to December ( $0.12 \pm 0.24 \mu\text{g m}^{-3}$ ). At the same time, the increase of wood combustion from nearby urban regions during colder periods (autumn-winter) combined with the changes in the meteorological conditions, prevents the transport of pollutants (Chu et al., 2005). The highest OC/EC ratio ( $6.3 \pm 7.51$ ) was observed for MCE air masses, while the lowest ratio was recorded for BAM at about  $2.2 \pm 1.10$ , as shown in Table 4. The OC/EC ratio observed at AM5 for BAM were as similar to those found in local samples at Northern Morocco with an average of 1.9 (Benchrif et al. 2018). But a Moreover, the OC/EC ratio shows a slight difference with those observed in Mt. Everest (Decesari et al., 2010) whose ratios varied from 5 to 9. To conclude, carbonaceous aerosols show strong season and temporal variation with high OC values observed during summer due to biogenic emissions, mineral dust, and long-range transport, in contrast, to EC which shows low variability except during ~~some regional~~ few regional anthropogenic pollution events.

### 3.3.3 Sea salt

The Middle Atlas region is influenced by two maritime sources of sea salt, more often from the Atlantic Ocean, and sometimes from the Mediterranean Sea. During the study period, the average concentration of sea salt remains low ( $0.45 \pm 0.55 \mu\text{g m}^{-3}$ ) and contributed only 1.6% of the total PM<sub>10</sub> concentration. The highest concentrations were recorded during August when sea salt concentrations reached a maximum of  $3.4 \mu\text{g m}^{-3}$ . The sea salt then decreased gradually, reaching a minimum concentration of  $0.06 \mu\text{g m}^{-3}$  during December. The sea salt concentration was high when wind speed exceeded  $6 \text{ m s}^{-1}$ , indicating that sea salt is strongly dependent on the meteorological conditions and air mass sources.

a mis en forme : Exposant

The higher concentrations of Na<sup>+</sup> ( $0.66 \pm 0.37 \mu\text{g m}^{-3}$ ) and Cl<sup>-</sup> ( $0.48 \pm 0.32 \mu\text{g m}^{-3}$ ) in the hot season (Aug-Sep) could be due to a larger contribution of marine aerosol. During this period, sea salt made up 11% ~~from~~ of the total PM<sub>10</sub> mass, especially when the air mass came from the Atlantic Ocean (ACE) in comparison to ~~compared to less than~~  $< 1\%$  of the PM<sub>10</sub> mass when the air masses were from the Sahara Desert (SD). However, no significant difference was noticed ~~of~~ in sea salt concentrations found in the Atlantic Ocean ACE ( $0.46 \pm 0.72 \mu\text{g m}^{-3}$ ) and Mediterranean Sea MCE samples ( $0.45 \pm 0.55 \mu\text{g m}^{-3}$ ), as shown in Table 4. Na<sup>+</sup> concentration was high during a pollution episode on the 16<sup>th</sup> of August, which ~~went along~~ coincided with, high concentrations of EC, SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup> ~~as well~~. This was due to the influence of ACE air masses with high EC content that made up about 2% of the PM<sub>10</sub> mass at the AM5 site. The average ratio of Cl<sup>-</sup>/Na<sup>+</sup> in PM<sub>10</sub> was found to be lower (0.43) than the value typically observed in seawater (1.81) (McInnes et al., 1994; Prodi et al., 2009). This points out to chlorine depletion due to chemical reactions that involve NaCl and HNO<sub>3</sub> or H<sub>2</sub>SO<sub>4</sub> leading to the formation of NaNO<sub>3</sub> or Na<sub>2</sub>SO<sub>4</sub> and gaseous HCl (McInnes et al., 1994). The estimated chlorine depletion was 42% for ACE

and 49% for MCE air masses comparable with reported values from 28% to 63% (Avg. 48%) observed in PM<sub>10</sub> in the Atlantic Ocean and Mediterranean Sea (Contini et al., 2010, 2014). ~~be on~~ Consequently, sea salt was mainly present as aged sea salt at AM5 during the sampling period. A detailed discussion on the correlation between Na<sup>+</sup> and Cl<sup>-</sup> will be elaborated on in section 3.4.

a mis en forme : Anglais (E.U.)

505

The comparison of sodium and chloride concentrations with other high-altitude studies is shown in table 54. On one hand, the observed that the concentrations of Na<sup>+</sup> ( $1.8 \pm 2.31 \mu\text{g m}^{-3}$ ) and Cl<sup>-</sup> ( $0.80 \pm 1.33 \mu\text{g m}^{-3}$ ) are several times lower than those at hill station at Darjeeling in India which has a concentration of Na<sup>+</sup> and Cl<sup>-</sup>, of  $2.2 \pm 2.0 \mu\text{g m}^{-3}$  and  $2.3 \pm 1.5 \mu\text{g m}^{-3}$ , respectively. On the other hand, the concentration of Na<sup>+</sup> and Cl<sup>-</sup> were 4 to 8 times higher than the values reported at Mt. Everest station located at an altitude of 5079 m asl. In addition, the concentration of chloride was in a good agreement with those observed value about  $82 \pm 98 \mu\text{g m}^{-3}$  in Mt. Cimone, Italy,  $0.82 \pm 0.98 \mu\text{g m}^{-3}$ . Sea salt concentration observed at the AM5 ( $0.45 \mu\text{g m}^{-3}$ ) was 5 times lower than at Tetouan ( $2.46 \mu\text{g m}^{-3}$ ), a coastal Mediterranean city in northern Morocco, and approximately 20 times lower than Cap Verde Atmospheric Observatory (CVAO) located in the tropical Atlantic Ocean (Benchrif et al. 2018; Fomba et al. 2014). The contribution of marine aerosols originating from the Atlantic Ocean remains relatively low compared to other sites, but nevertheless existent.

### 3.3.4 Ammonium, nitrate, and sulfate

520

A significant part of aerosol PM composition was associated with the formation of secondary inorganic aerosols (SIA), about 7.2% of PM<sub>10</sub> mass which are mainly composed by of sulfate, nitrate, and ammonium. They made up about 7.2% of the PM<sub>10</sub> mass. The temporal variation during the sampling period of SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, and NH<sub>4</sub><sup>+</sup> is presented in Fig. 6, with average concentrations of  $0.9 \pm 0.8 \mu\text{g m}^{-3}$ ,  $0.8 \pm 0.6 \mu\text{g m}^{-3}$ , and  $0.3 \pm 0.2 \mu\text{g m}^{-3}$ , observed, respectively. In summer, the concentrations were relatively high during the few days in August, as shown in Fig. 6. The with the observation of the highest sulfate, nitrate, and ammonium concentrations with maximum concentrations of up to  $6.1 \mu\text{g m}^{-3}$ ,  $4.4 \mu\text{g m}^{-3}$ , and  $1.2 \mu\text{g m}^{-3}$ , respectively (Fig. 6). This was due to the transport of polluted MCE air masses through the Mediterranean Sea and across cities in the North of Morocco leading to high PM loaded aerosols. High peaks concentration Peaks due to air mass influence On average, the influence of long-range transport during the coming from the Atlantic Ocean ACE remains relatively similar to the Mediterranean Sea and MCE air masses for sulfate ( $2.8 \mu\text{g m}^{-3}$ ) and nitrate ( $2.3 \mu\text{g m}^{-3}$ ) were similar. However, the contribution of ammonium ( $1.7 \mu\text{g m}^{-3}$ ) to particulate matter was particularly higher for MCE air mass. Additionally, other peaks were also observed in aerosol concentrations both for SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> during August. This could be attributed to the long-range transport of dust aerosol from the Saharan desert in Southern Morocco. The intervening subsequent months demonstrate a clear decreasing trend of SIA from high concentrations in summer ( $3.8 \mu\text{g m}^{-3}$ ), to relatively low concentrations during winter ( $1.01 \mu\text{g m}^{-3}$ ). In average, the sulfate concentration for MCE ( $1.2 \pm 0.9 \mu\text{g m}^{-3}$ ) was about 5 times higher than background sulfate concentrations ( $0.2 \pm 0.2 \mu\text{g m}^{-3}$ ). In addition, non-sea salt sulfate (nss-SO<sub>4</sub><sup>2-</sup>) represented about 95% of the total SO<sub>4</sub><sup>2-</sup> at AM5 and it had a strong correlation with NH<sub>4</sub><sup>+</sup>, suggesting that secondary sulfate was mainly present as ammonium sulfate (detailed discussion later in section 2.3.4??).

a mis en forme : Couleur de police : Texte 1

Secondary inorganic aerosol over the Atlas Mountains have been compared with the data reported in other high-altitude stations (Table 54). The average concentration of nitrate ( $0.8 \pm 0.6 \mu\text{g m}^{-3}$ ) at AM5 was comparable with those reported in Mt. Himalaya and Mt. Cimone,  $0.9 \pm 0.2 \mu\text{g m}^{-3}$  and  $0.8 \pm 0.7 \mu\text{g m}^{-3}$ , respectively (Chatterjee et al., 2010; (Marenco et al., 2006). However, the concentration of  $\text{NO}_3^-$  were found to be approximately 2 times higher than the value reported in Mt. Puy de Dôme (Bourcier et al., 2012), as shown in Table 5. Two factors could be responsible for the high nitrate concentration recorded at AM5. Firstly, the mineral dust particles present in the Middle Atlas region contain calcium carbonates (calcite and dolomite) which can react with nitric acid gas in the atmosphere to form nitrate salts (Krueger et al., 2004; Khrissi et al., 2018). Second, the difference could be explained by long-range transport of polluted air mass from MCE and ACE which enhanced the nitrate concentration. Similar concentrations of ammonium at AM5 ( $0.3 \pm 0.2 \mu\text{g m}^{-3}$ ) were found at Puy de Dôme, France ( $0.3 \pm 0.2 \mu\text{g m}^{-3}$ ), whereas the concentration was 5 times lower than those reported in other high-altitude sites such as the Mt. Himalaya (Chatterjee et al., 2010). This indicates that the influence of ammonium remains relatively low despite the proximity of the approximate site of the agricultural activities located in the surroundings of Meknes. The concentrations of sulfate ( $0.9 \pm 0.8 \mu\text{g m}^{-3}$ ) over AM5 were comparable with those at Puy de Dôme ( $1.3 \pm 1.1 \mu\text{g m}^{-3}$ ), but were almost 4-5 times lower than all the other hilly stations except Mt. Everest (Decesari et al., 2010).

a mis en forme : Normal, Interligne : 1,5 ligne

540  
545  
550  
555

### 3.3.5 Organic compounds

The identification of the organic chemical compounds enables a better understanding of the organic fraction in the composition of aerosols and the quantification of the contribution of biogenic and as well as anthropogenic emissions (Jaenicke, 2005). Therefore, a large number of individual organic chemical compounds were analyzed. Figure 68 shows the temporal variation of organic compounds, including n-alkanes, PAHs, and ketones n-alkan-2-ones and sugars.

#### n-alkanes

Individual n-alkanes with C atom numbers in the range 20-34 were analyzed. The distinguishing aspect of alkanes is their specific source and their ability to provide information about their origins (Pietrogrande et al., 2010). Individual n-alkanes with C atom numbers in the range 19-34 were analyzed. Figure 68 shows the temporal variation of the n-alkanes—the background composition, which is not necessarily constant revealing strong variations over the seasons at with an average concentration of about  $86.417 \pm 7.1 \text{ ng m}^{-3}$ . In contrast, the variation of the average concentration under the influence of air masses decreases from summer ( $16.148 \pm 8.9 \text{ ng m}^{-3}$ ) to winter ( $24.687 \pm 2.0 \text{ ng m}^{-3}$ ). On average, the alkanes during BAM conditions accounted was about for  $4.9 \pm 3.2 \text{ ng m}^{-3}$  and was dominated by biogenic species such as heptacosane, hentriacontane, and nonacosane which made up about accounts for 60% of the total alkanes, as shown in Fig. S10. During SD air mass influence, such as in August, high concentrations of alkanes were observed (about  $50.9 \text{ ng m}^{-3}$ ). The predominant compounds during dust events were pentacosane, hexacosane, heptacosane, and nonacosane suggesting that OC was loaded with biogenic matter (Pio et al., 2001). MCE air mass influence revealed considerably high ( $10.5 \pm 7.7 \text{ ng m}^{-3}$ ) n-alkane concentration with elevated concentrations of typical anthropogenic tracers found on the samples such as nonadecane, and tricosane. This indicates that the long-range transport of MCE air masses was often loaded with anthropogenic material. While the contribution of MCE and SD air masses to alkanes is two

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

a mis en forme : Police : Italique

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Non Surlignage

a mis en forme : Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

times higher ~~as than~~ during the background conditions, the concentration recorded for ACE  $5.6 \pm 3.7 \text{ ng m}^{-3}$  remains relatively similar to BAM conditions ~~by~~. Additionally, organic compounds such as pristane and phytane, considered as typical molecular markers of traffic emissions were almost not found or rarely found in most of the samples. This indicates that the influence of traffic was considerably low.

a mis en forme

[... [5]]

To distinguish between natural biogenic emissions from plants and incomplete combustion, the carbon preference index (CPI) was also calculated and used as a marker (Alves et al., 2012; Pietrogrande et al., 2011). The part of n-alkanes with an ~~odd-even~~ number of C-atoms exceeding the distribution of the average concentration of n-alkanes can be considered as coming from ~~plant~~ vegetable waxes. However, odd C-atom-numbers can ~~also~~ originate from incomplete biomass combustion (Iinuma et al., 2007). Table 4 presents the CPI values calculated according to each air mass. The average CPI value was  $3.8 \pm 2.4$  and ranged from 0.77 to 18.63. However, high CPI ( $>1$ ) was observed for all air masses, which indicates that the alkanes originated from plants waxes, as presented in Table 4 (Kavouras, 2002). In contrast, ~~not~~ values of CPI were recorded close to 1, which shows the minor influence of anthropogenic activities and traffic emission at AMS. The average concentration of alkanes was ~~is~~ dominated by odd C-atoms with a concentration of  $6.4 \text{ ng m}^{-3}$ , compared to  $1.9 \text{ ng m}^{-3}$  for even C-atoms. During summer, higher concentrations were observed, approximately  $39.2$  and  $10.4 \text{ ng m}^{-3}$  for n-alkanes with odd and even C-atoms, respectively. Therefore, the average CPI increases during dust events during SD air mass influence during dust event ( $4.0 \pm 3.1$ ) was due to ~~and~~ the higher contribution of odd C-atoms alkanes. Similar to SD, the average CPI for MCE was about  $3.9 \pm 1.9$  during MCE air mass influence was due to higher contribution of even C-atoms alkanes. However, a slight decrease of the mean CPIs ( $3.3 \pm 0.80$ ) was observed during the BAM conditions occurring mainly during autumn and winter, as ~~conditions~~ the biogenic activity was relatively low compared to the summer. Overall, the most dominant n-alkanes such as nonacosane and hentriacontane ~~ever all~~ were observed during all the air masses indicating high local influence and they ~~and~~ were typically associated with the biogenic activity.

a mis en forme

[... [6]]

a mis en forme : Anglais (E.U.)

a mis en forme

[... [7]]

a mis en forme : Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

### PAHs

In the present ~~his~~ study, 32 polycyclic aromatic hydrocarbons (PAHs) with 3 to 7 rings were quantified. In Fig. 8 the temporal variation in the form of the sum of the 20 identified PAH compounds of PAH identified in the particles is presented in Figure 6. The contribution of PAHs was much lower than alkanes with an average concentration of  $0.68 \pm 0.83 \text{ ng m}^{-3}$  over the whole study period. The mean concentration of PAHs was about  $6.9 \text{ ng m}^{-3}$ . Contrary to what has been observed for alkanes, the PAH concentrations determined during the autumn months were more important ~~higher~~ than those during the summer and winter ~~ones~~. The highest amount of PAH was detected during October ~~autumn~~, approximately  $53.784 \text{ ng m}^{-3}$  due to long-range transport of MCE air masses, as shown in Figure 6. The minimum concentration was observed during winter, of about  $0.605 \text{ ng m}^{-3}$ . The average background concentration of PAHs was  $0.4 \pm 0.5 \text{ ng m}^{-3}$ , which was ~~Their influence remained~~ low in comparison to other organic compounds ~~likely~~ because of high evaporation on warm days (Cincinelli et al., 2007). During MCE air mass, the PAH concentrations increased by 52% compared to the BAM concentration, as shown in Table 5. The most abundant PAHs found in the BAM samples were fluorene and retene, which represent 75% of total background PAH concentrations. The abundance of fluorene (Fa) and retene (Rete) ~~which~~ were found in samples from different air masses suggesting that they potentially ~~come~~ originate from similar sources ~~potentially~~ local or

a mis en forme : Police : Italique

a mis en forme

[... [8]]

a mis en forme : Anglais (E.U.)

a mis en forme

[... [9]]

620 regional emissions sources. Moreover, fluorene and retene are marker compounds for wood combustion of heating or combustion of organic substances, but it is also found in trace concentrations in the combustion of gasoline or diesel (Spindler et al., 2012). Other compounds such as 17b(H),21b(H)-hepane coronene (6.4%), 2,6-dimethylphenanthrene dibenzo(a,h)anthracene, (6.5%), or 3,5-dimethylphenanthrene phenanthrene (6.2%) were observed during long-range transport of polluted air masses during ACE or NAO-MCE air mass influence. The contribution of PAHs remains relatively lower for ACE ( $0.4 \pm 0.4 \mu\text{g m}^{-3}$ ) air masses than for MCE air mass influence, which indicates that not all long-range transport was loaded with combustion tracers. The average PAH concentration was higher during SD than ACE air masses of about  $0.7 \pm 0.7 \mu\text{g m}^{-3}$ , however The most abundant compounds were fluorene and 9H-fluorenone, such organic compounds which were found within a similar concentration ranged during conditions. Therefore, mineral dust transport was not affected by the combustion processes. As a result, the observation of PAH concentration shows a strong variation with high biogenic activities in the surroundings during summer and high anthropogenic PAHs during pollution episodes from combustion processes in autumn.

a mis en forme

[... [10]]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme

[... [11]]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme

[... [12]]

### *n-alkan-2-ones*

a mis en forme : Couleur de police : Texte 1

a mis en forme : Police : Italique

a mis en forme : Police : Gras, Soulignement , Couleur de police : Texte 1

a mis en forme : Normal

[... [13]]

In total, 5 n-alkan-2-ones ketones were detected in this study, as shown in Fig S8. The n-alkan-2-one ketones concentrations increased significantly from summer ( $1.8 \text{ ng m}^{-3}$ ) to autumn ( $9.7 \text{ ng m}^{-3}$ ), then decreased continuously to winter ( $6.3 \text{ ng m}^{-3}$ ), by a factor of 2, with an average of  $6.69 \text{ ng m}^{-3}$  for the whole sampling period  $\text{ng/m}^3$ . The minimum concentration was recorded during the summer of about  $0.60 \text{ ng m}^{-3}$ . In contrast, the maximum concentration was reached during autumn of about  $52 \text{ ng m}^{-3}$  due to ACE air mass transport influence, as shown in Fig. 6. The sum of n-alkane-2-one was between  $0.67$  to  $13.2 \text{ ng m}^{-3}$ . The same relative composition of n-alkan-2-one concentrations was observed in both seasons, suggesting that they came from similar sources. However, the levels of n-alkan-2-one were much lower in concentration<sup>22</sup> than those of n-alkanes. The average background concentration of the total n-alkan-2-one was  $5.9 \pm 5.5 \text{ ng m}^{-3}$ . During this period, major n-alkan-2-one constituents recorded at AM5 were 2-Nonadecanone, 2-Heptadecanone, 2-Octadecanone, which represented 29%, 25%, and 18% of the total detected n-alkan-2-one, respectively. These organic compounds that appear in fossil fuel burning events were found in low concentrations at the AM5. The remaining part was made up of 2-hexadecanone and 2-octadecanone, which counts made up for 15.6%, and 6.2%, of the total n-alkan-2-ones, respectively. The discrepancy of n-alkan-2-ones concentrations in comparison to BAM chemical composition was low from August to October. The trend is completely reversed, as the average concentration was significantly elevated during November and December. During these two months, the concentration increased by 35%. Some spikes detected in November and December are characterized by a change in wind direction, and high wind speeds. These samples, with higher concentrations for 2-heptadecanone and 2-nonadecanone show a strong correlation with temperature. Indeed, this period marks the beginning of winter, with average temperatures dropping as low as  $4.5^\circ\text{C}$ , as shown in Table 1. No correlation between alkane-2-ones was found with elemental carbon, but the analysis of the data shows a strong correlation only for the night samples with As and K<sup>+</sup>. This indicates that a possible source of the n-alkan-2-ones is combustion due to residential heating. These data present the first measurement made that shows the influence of combustion in North Africa on n-alkan-2-ones concentrations. Similar conclusions were reported by Müller, (1997) who highlighted the anthropogenic source of alkane-2-ones during winter.

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme

[... [14]]

Two main sources are known to be responsible for the presence of n-alkane-2-one in the air: Incomplete combustion and in-situ microbial  $\alpha$ -oxidation of the carbon chain (Khedidji et al., 2020). The first source often creates n-alkane-2-one with a predominance of odd-numbered carbon atoms; on the other hand, the last source gives rise to the opposite. In the present study, a strong predominance of odd n-alkane-2-one rather than pairs was clearly observed for these compounds, suggesting that the incomplete combustion of organic material was the principal ambient source in this region. In particular, heptadecane-2-one (K17) was the most abundant in all the samples and reached a maximum 5.23 ng m<sup>-3</sup>.

a mis en forme : Anglais (E.U.)

660  
665 **Sugar alcohols**  
Sugar alcohols contributed to the organic fraction where Three main sugar alcohols that are compounds of sugar levoglucosan, arabitol and glucose were identified between August 2011 and December 2017 in the AM5 particle samples from Atlas Mountains. The sugar concentration levels in the aerosol samples ranged from 0.02 to 39.6 ng m<sup>-3</sup>. The average concentrations of sugar compounds were higher during summer (7.9 ng m<sup>-3</sup>) and decreased continually until November (1.2 ng m<sup>-3</sup>). During December, the concentrations were relatively higher than November (2.1 ng m<sup>-3</sup>) where some high peaks were observed. Glucose sugars were was about five times higher in summer than in winter. Notably, there were three extreme days observed for sugars were where high sugar concentrations were observed, as shown in Fig. 6.7: 13<sup>th</sup> August 2017 (39.6 ng m<sup>-3</sup>), 18<sup>th</sup> September 2017 (ng m<sup>-3</sup>), and 20<sup>th</sup> October (28.6 ng m<sup>-3</sup>). The high values observed during These peaks are due to the long-range transport from the coast of Europe (ACE) and the coast of the Mediterranean Sea (MCE). Their average sugar concentrations during these air mass influences were the higher throughout the sampling period: ACE (3.70  $\pm$  6.05 ng m<sup>-3</sup>) and MCE (5.26  $\pm$  7.76 ng m<sup>-3</sup>), as listed in Table 4. In contrast, sugar compounds were relatively low in SD (1.85  $\pm$  2.94 ng m<sup>-3</sup>) air masses and were not found in the background PM<sub>10</sub> conditions. Levoglucosan which is considered as a good tracer of biomass burning emissions in aerosol particulate matter, was particularly higher for ACE (2.07 ng m<sup>-3</sup>) and MCE (1.65 ng m<sup>-3</sup>) air mass influence, as displayed in Fig. S810. (Bauer et al., 2008). While arabitol shows a similar concentration for MCE and ACE with a mean of 1.05 ng m<sup>-3</sup> which suggesting that particles were loaded with primary biological aerosols such as pollen, fungal spores, vegetative debris, viruses, and bacteria were transport from the marine coast (Fu et al., 2012). Whereas glucose remained relatively high during MCE air mass influence in comparison with to other air masses influence. During SD air mass influence. Moreover, the the concentration of arabitol was extremely low in SD air mass with a concentration less than 0.08 ng m<sup>-3</sup>. However, glucose showed a higher concentration of about 0.74 ng m<sup>-3</sup> but remains 3 times lower than MCE concentrations. This proves indicates that the mineral dust sugars were most likely originated from marine air masses. was not loaded with biological material as previously reported.

a mis en forme : Couleur de police : Texte 1

a mis en forme : Police : Italique

a mis en forme : Interligne : 1,5 ligne

a mis en forme : Police : Italique, Non souligné, Couleur de police : Texte 1

a mis en forme

[15]

670  
675  
680  
685  
690 To conclude, Sugars have potentially two major sources at AM5: a natural biological source from marine air masses including MCE and ACE and an anthropogenic source from biomass burning potentially from urban cities close to the site. The contribution of arabitol and glucose was significantly higher during the summer, linked to more developed vegetation and a higher biogenic activity, and in contrast to winter, where levoglucosane appears to be was higher. Sugars have potentially two major sources at AM5: a natural biological source from marine air masses including MCE and ACE and an anthropogenic source from biomass burning potentially from urban cities close to the site.

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme

[16]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme

[17]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1

[18]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

### 3.3.6 Crustal enrichment factor

Analysis of the crustal enrichment factor (EF) has been used to estimate the contributions of crustal matter to the ambient PM<sub>10</sub> particles at AM5. For this study, titanium (Ti) was used as a reference element due to the low recovery of Al and high recovery of Ti, and as it is also considered a suitable tracer for mineral dust (Fomba et al., 2013). Furthermore, Al or Fe has more anthropogenic sources than Ti, and Fe has more anthropogenic sources, which makes Ti suitable as reference element. (Fomba et al., 2013). However, the comparison of EFs for Ti and Al as reference elements shows a similar trend, with only slight differences observed in the absolute values. The average upper continental crust composition, according to Wedepohl (1995) was used for the calculation of the enrichment factors. The EF relative to Ti was calculated using equation 2 as follow:

$$EF = \frac{(Z_{Ti})_{sample}}{(Z_{Ti})_{crust}} \quad (2)$$

a mis en forme : Indice

The enrichment factor provides the ability to classify metals based on their enrichment to the soil. Elements with an EF under 2 are considered to have a similar composition to the reference soil values. An enrichment factor above 2 but below 10 is assumed to have low enrichment with a possible mixture of both crustal and non-crustal sources. Elements with an EF above 10 are considered enriched, while enrichment factors above 100 are considered highly enriched, suggesting that the elements are from non-crustal and more likely anthropogenic sources. The enrichment factor does not take into account each pollution episode but is a general approach to the classification of metals according to their crustal origin. Within the present study, the elemental enrichment factors showed similar trends for the different air mass inflow to the station. Three groups of elements could be identified from the elemental enrichment factors. Figure 7 shows the average PM<sub>10</sub> elemental crustal enrichment factors (EF) of aerosol PM<sub>10</sub> evaluated for the different trace metal elements at AM5 according to their respective air mass origins.

a mis en forme : Anglais (E.U.)

Group I includes elements such as Al, Ba, Rb, K, and Fe with enrichment factors between 0.8 and 2. Their enrichment factors suggest that these elements are associated with particulate matter from the resuspension of soil or other crustal sources. Elements such as Al, Fe, and Mn show little dispersion, and their variation seems to be constant across different air masses, clearly indicating that the source was soil. As suggested by other studies, these metals could also have an anthropogenic source, but in this study, they clearly showed crustal matter origin (Viana et al., 2008; Birmili et al., 2006; Contini et al., 2012). In On one hand, no correlation was found between Al, Fe, and anthropogenic tracers such as EC or other heavy metals, which indicates their natural origin. On the other hand, K showed slightly higher EFs for air masses from the Atlantic Ocean, suggesting that sea salt and sources other than mineral dust, such as biomass combustion, might have contributed to its presence.

Group II elements include heavy metals such as Sr, Ca, Cu, Mn, Ce, V, La, Co, and As. These elements had enrichment factors ranging from 2 to 10, indicating the possibility of having mixed origin from both crustal and anthropogenic sources. The lowest enrichment factors were observed during In the case of background samples,

**conditions** suggesting that the elements may have been of crustal origin, **the lowest enrichment factors were observed**. In contrast, the highest enrichment factors were mainly observed in air masses that originated from the coast of Europe (ACE) and crossed major urban cities such as Rabat/Salé/Kenitra and Casablanca before arriving at Atlas station M5. In this case, it is assumed that these elements were probably influenced during transport by anthropogenic emissions. In contrast, the Mediterranean Sea air mass appears to remain relatively unaffected by anthropogenic emissions. In addition to its atmospheric crustal origin, V had a high enrichment factor mainly due to residual oil combustion, especially at night. Particles from oil combustion processes were often observed in high concentrations during winter due to their size and long lifetime in the atmosphere and the combustion activities in the nearby urban cities.

740

745

Group III contains the elements with EF from 10 to 1000, including heavy metals such as Cr, Zn, Ni, Pb, as well as Br, Se, and Sb. These elements showed high enrichment factors in all air mass directions. They are mainly present in the marine air masses of the Atlantic, but also the Mediterranean air masses. An increase in heavy metal concentration has been observed during winter, and at night when the temperature drops **and**, the air mass inflow from the cities towards the mountain prevails. Atmospheric Ni and Cr are released during combustion processes, while Pb is mainly released from smelters or the combustion of unleaded petrol, waste, and coal (Pacyna et al., 2007). Combustion processes are generally the main contributors to these anthropogenic metals. Zn had a weak correlation with Pb and Ni, suggesting that its origin is also anthropogenic. The nearest urban cities are Meknes and Fes, where anthropogenic activities such as waste incineration, and road traffic pollution are common. Furthermore, the V/Ni ratio was observed higher for MCE air mass, about 2.8 which is considered typical for heavy fuel oil combustion (Mazzei et al., 2008; Pandolfi et al., 2009; Bove et al., 2014).

750

755

a mis en forme : Anglais (E.U.)

### **3.4 Inter-relationship between aerosol components**

**The inter-relationship between the different species and the scatter plots are analysis of the correlation matrices between the different species presented in Fig. 8.** The analysis of the single correlation coefficients allows obtaining information about the possible common sources of aerosol.

760

765

770

#### **3.4.1 Nitrate and nss-sulfate**

The correlation between  $\text{NO}_3^-$  and  $\text{nss-SO}_4^{2-}$  ( $r^2=0.76$ ) indicates their possible common origin. The correlation was more pronounced **particularly observed** for MCE air masses ( $r^2=0.80$ ) in contrast to ACE ( $r^2=0.43$ ) air masses, **suggesting an enhanced the transport of the secondary anthropogenic aerosol from the Mediterranean-coast** (Liu et al., 2017) to the AM5 site. The nss-sulfate concentrations were slightly correlating with Vanadium, which is associated with the emissions of oil combustion, ships emissions as well as iron and steel industrial settlements emissions (Pandolfi et al., 2011). A strong correlation of  $\text{NO}_3^-$  and  $\text{nss-SO}_4^{2-}$  was observed with oxalate ( $\text{C}_2\text{H}_4\text{O}_4^{2-}$ ), which could indicate that they have a common source and that they can eomeoriginate from biomass burning and secondary transformations. However,  $\text{nss-SO}_4^{2-}$  also originated attributed to from crustal sources especially as soil-elevated origin concentrations were observed during dust events. This assertion was supported by a good correlation of  $\text{nss-SO}_4^{2-}$  with  $\text{nss-Ca}^{2+}$  (Fig. S7), indicating the likely presence of calcite particles most often from the of crustal origin. A similar observation was reported by Okada and Kai, (2004), who observed that Desert dust was associated with sulfur compounds and organic matter from surrounding agricultural areas. Indeed,

a mis en forme

[19]

a mis en forme : Anglais (E.U.)

[20]

a mis en forme : Anglais (E.U.)

[21]

a mis en forme : Couleur de police : Rouge foncé

[22]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[23]

a mis en forme : Normal, Interligne : 1,5 ligne

[24]

a mis en forme

[25]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[26]

a mis en forme : Police :Gras, Anglais (E.U.)

[27]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[28]

a mis en forme : Normal, Interligne : 1,5 ligne

[29]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[30]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[31]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[32]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[33]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[34]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[35]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[36]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[37]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[38]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[39]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[40]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[41]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[42]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[43]

a mis en forme : Couleur de police : Texte 1, Anglais (E.U.)

[44]

775 the particles with high sulfate content were accompanied by Ca and were assigned as gypsum particles, also which  
suggesting that the sulfur in these particles originated from a sedimentary source (Falkovich et al.,  
2001). Ammonium was less enhanced during dust storms. The correlation with ammonium will be discussed in  
detail in the next section.

### **3.4.2 Ammonium nitrate and ammonium sulfate**

The analysis of the correlation matrices between nss- $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  with ammonium ( $\text{NH}_4^+$ ) was applied to better understand the inter-relationship between the secondary inorganic species. A correlation between nss- $\text{SO}_4^{2-}$  and  $\text{NH}_4^+$  ( $r^2 = 0.90$ ) supported the hypothesis of dominant ammonium sulfate particles ( $(\text{NH}_4)_2\text{SO}_4$ ) in the summer especially when air masses were coming from ACE, as shown in Fig. 8. During this period, a strong correlation was found between sulfate and solar radiation which suggests that nss- $\text{SO}_4^{2-}$  was produced via photochemical reaction (Baker and Scheff, 2007). Nevertheless, the transport of nss- $\text{SO}_4^{2-}$  from the Atlantic coast also contributes to the formation of ammonium sulfate. However, the trend is more towards ammonium nitrate ( $\text{NH}_4\text{NO}_3$ ) in winter, given that the main correlation of  $\text{NH}_4^+$  with  $\text{NO}_3^-$  ( $r^2 = 0.95$ ) mainly present in MCE air masses. Nitrate shows a strong dependency on the temperature at AM5, most likely due to the stability of ammonium nitrate in the atmosphere at low temperatures (Squizzato et al., 2013). The predominance of nitrates over sulfates during winter, where nitrates and ammonium remain high, is probably due to the influence of temperature that prevents the dissociation of ammonium nitrate particles (Ricciardelli et al., 2017). Moreover, a similar pattern of  $\text{NO}_3^-$  and  $\text{NH}_4^+$  as observed by Querol et al., 2004, in the Mediterranean coast with a summer minimum, and suggested that it could be due to the low thermal stability of the nitrate in the hot season.

#### 3.4.4 Sodium and chlorine

The evolution of the sea salt constituents and their relationship with the most important aerosol acidic species such as  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  was investigated according to their air mass origins (Fig. 8). A correlation between sodium and chloride was observed ( $r^2=0.76$ ), as shown in Fig. 8. The scatter plot of molar equivalent concentrations of  $\text{Na}^+$  and  $\text{Cl}^-$  shows a strong correlation specifically for ACE and SD air masses. However, the data points are below the seawater reference line and only approach this line when the  $\text{Cl}^-$  concentration is combined with  $\text{NO}_3^-$  and nss- $\text{SO}_4^{2-}$ . This indicates that chloride was depleted in the sea salt particles due to the displacement of chloride by sulfate from sulfuric acid when air masses were coming from MCE and ACE, especially as since photochemical processes favor sulfate formation during summer. The same scenario has been observed for  $\text{NO}_3^-$  with a considerable difference during the winter. Indeed, the correlation between sodium and the sum of chloride and nitrate shows the chloride depletion and indicates that the Mediterranean Sea air mass was loaded with aged sea salt. Similar results were observed in the North of Morocco where the mass fraction of nitrate was higher in the coarse fraction which is indeed corresponding to aged sea salt (Benchrif et al. 2018). No correlation between  $\text{Na}^+$  and  $\text{Cl}^-$  was observed in the BAM conditions. The influence of sea salt remains a significant consequence despite the distance that separates the station from the ocean coast.

### 3.33.5 Day and night-time variation

810 Inorganic ions, mineral dust, and organic matter carbon (OCM)

Diurnal variations of various PM<sub>10</sub> chemical species were analyzed to understand the influence of day and night variations on their concentrations. Figure 9.11 shows the variation between day and for given chemical species.

**a mis en forme** : Police :Gras, Anglais (E.U.), Barré

**a mis en forme** : Indice

**a mis en forme** : Exposant

**a mis en forme** : Indice

**a mis en forme** : Exposant

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Exposant

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1, Anglais (E.U.)

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Anglais (E.U.)

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1, Exposant

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Anglais (E.U.)

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1, Exposant

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1, Indice

**a mis en forme** : Couleur de police : Texte 1, Exposant

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Indice

**a mis en forme** : Exposant

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Couleur de police : Texte 1

**a mis en forme** : Non Surlignage

**a mis en forme** : Couleur de police : Rouge foncé

The OC increase from night-time ( $0.9 \mu\text{g m}^{-3}$ ) to the day ( $1.2 \mu\text{g m}^{-3}$ ) and was accompanied by a slight increase in alkanes such as pristane, docosane, and nonacosane, as observed in Fig. 9a. These alkanes indicate that the organic fraction was dominated by biogenic sources during the day. The highest concentrations of biogenic compounds were reached during the summer. During summer, higher concentrations of Al ( $0.6 \mu\text{g m}^{-3}$ ) and Fe ( $0.7 \mu\text{g m}^{-3}$ ) were observed during the day, compared to the night. This increase seems to be related to an additional source from the resuspension of road dust, due to car traffic during the day. Nevertheless, the transport of mineral dust tracers from the Saharan dust could be controlled by other factors. For example, the study done by Khan et al., (2015) indicates that the penetration of dust into the free troposphere in the Atlas Mountains can also be due to orographic lifting, convection on the mountain slopes, and updrafts in the breeze front. While the  $\text{Ca}^{2+}$  concentration which is approximately  $0.6 \mu\text{g m}^{-3}$  seems stable between day and night. The continuous presence of calcium indicates that it comes from a different source most likely local or regional. This difference in dust trace metal concentrations shows the important role that mineral dust calcium plays as a local source.

a mis en forme : Exposant

#### Elemental carbon (EC), anthropogenic metals, and PAHs

The composition of  $\text{PM}_{10}$  in the evening is characterized by a high concentration of anthropogenic trace elements. The EC concentrations slightly increased in the evening from  $0.22$  to  $0.26$   $\mu\text{g m}^{-3}$ . Besides, an increase of PAHs and ketones~~alkane2-one~~ concentrations such as fluorene, retene, and 2-nonadecanone was observed during the night-time. The PAH concentrations during the day and night-time were  $0.68$  and  $0.84 \mu\text{g m}^{-3}$ , respectively. In particular, the Fluorene was the most abundant PAH in all the night samples and reached a maximum of  $2.6 \text{ ng m}^{-3}$ . A correlation ( $r^2=0.67$ ) was found between PAHs and EC during night-time indicating their anthropogenic origins. In addition, anthropogenic metals such as Pb, Cr, V, Cr, Ni, and Cu associated with combustion and traffic emissions, increase by a factor of 1.8 during the evening (Fig 9b).

a mis en forme : Non Surlignage

Indeed, the anthropogenic influence at the AM5 site occurs during the evening due to two important factors. First, the site is located in a mountainous region influenced by the temperature fluctuation between day and night. The rapid cooling between day and night was accompanied by a change of aerosol sources. This phenomenon is widespread, especially in summer. Second, the variation of the air mass, combined with a change in the height of the boundary layer, contributes to the transport of pollutants from urban sites Fes and Meknes to the AM5 site.

a mis en forme : Non Surlignage

#### Influence of meteorology

The variation of the meteorological parameters between day and night is a critical factor that can indeed influence the chemical composition of the particles. First, a significant difference in  $\text{PM}_{10}$  was observed on days when the day and night temperature difference was substantial, for example, on the 12th of August. The concentration decreased during the day ( $113.01 \mu\text{g m}^{-3}$ ) to  $80.4 \mu\text{g m}^{-3}$  at night. second, the influence of meteorology on s Secondary inorganic aerosols, such as sulfates and nitrate, were characterized by different variations between day and night. On the one hand, the sulfate is highly slightly enhanced from night ( $0.94 \mu\text{g m}^{-3}$ ) to during the day ( $1.21 \mu\text{g m}^{-3}$ ). This increase during the day could be explained partly by Sulfate is also source sulfate originating from dust resuspension, which explains its increase during the day. However, a correlation between solar radiation and sulfate suggests that photo-oxidation during the day could also be a source of the sulfate increase, while increase. Whereas, nitrate recorded during the day ( $0.80 \mu\text{g m}^{-3}$ ), shows higher concentrations during the

night especially ( $1.06 \mu\text{g m}^{-3}$ ). The drop in temperature between the day and the night, especially during the winter, allows for a rapid formation of ammonium nitrate.

855 *Mechanism of day-night variation*

In principle, two mechanisms control the variation between day and night: The wind direction and the boundary layer height. The wind direction plays an important role because it introduces air mass transported from different sources. Indeed, the winds come from all directions, but it is dominated from the ~~west east section~~ during the day, and by the ~~South East~~west during the night, as shown in Fig .S10. High speeds were recorded during the night, up to  $17.5 \text{ m s}^{-1}$ , mostly associated with marine air masses. This suggests that long-distant transport often occurred during the night while the wind speed during the day was relatively lower. The lower wind speed during the day indicates that the influence of local sources is important. Furthermore, the topography, as well as the embedded valleys, also play a role in the pollution transport during the daytime as shown by Lang et al., (2015). Mountains can give rise to daytime upslope winds and night-time downslope winds. The valley bottom warms during the day, warm air rises the slopes of the surrounding mountains and hills to create a valley breeze. During night-time, radiation from the earth's surface cools the slopes, causing cooler, denser air to drain into the valley. In addition, local boundary layer processes and long-range transport contribute to chemical composition changes (Nair et al., 2007). A similar impact of mountain-valley circulations on air pollution was observed by Bei et al. (2018). Studies at high-altitude sites in southwest India, also found that diurnal variations of aerosol particle concentrations were related to mountain valley winds and the variation in a planetary boundary layer height (Buchunde et al., 2019).

a mis en forme : Couleur de police : Rouge foncé

**3.4.3.6 Differences in chemical composition between dust and non-dust events**

To investigate the impact of the dust event on the  $\text{PM}_{10}$  chemical composition, the data has been segregated into two categories dust and non-dust episode. Only selected days with high influence of Saharan air mass with Al or  $\text{Ti} > 1 \mu\text{g m}^{-3}$  xxx, were considered to be representative for dust event days backward trajectories. This approach was previously used by Kocak et al., 2012 to investigate the influence of mineral dust in the eastern Mediterranean. Whereas the non-dust samples were categorized on the basis of based on aerosol Al concentrations ( $\text{Al} < 1 \mu\text{g m}^{-3}$ ). All the samples that constitute a mixing scenario were excluded. Figure 10 shows the average concentrations of (A)  $\text{PM}_{10}$  mass, (B) Fe/Al and Fe/Ca ratios, (C) OC and major ionic species, (D) EC and minor ionic species, (E) organic compounds during non-dust and dust events.

a mis en forme : Non Surlignage

880

*Mass*

The Long-range transport of mineral dust showed a significant impact on  $\text{PM}_{10}$  composition. During dust events,  $\text{PM}_{10}$  concentrations were on average 3 times higher, and up to a maximum of 10 times higher, in comparison to non-dust days. Mineral dust was about 5 times higher in comparison to background conditionsnon-dust samples (Fig. 10a). In addition, RH was lower during dust event days ranging from 20% to 45%, whereas it was 50% to 70% on the non-dust days. Similar results were observed by (Mukherjee et al., 2020) showing the impact of dust on the local meteorological conditions.

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1

a mis en forme : Exposant

a mis en forme : Couleur de police : Texte 1

a mis en forme : Indice

*Minerals, metals, and ions*

890 [Aerosol inorganic species demonstrate distinct differences in chemical composition between dust and non-dust events](#). Although the North African mineral dust is mainly made up of clay minerals and quartz, the content of calcium carbonates varies depending on the North Africa source (Chiapello et al., 1997; Glaccum and Prospero, 1980). A comparison between Fe/Al and Fe/Ca ratios were used to provide the potential geographical origin of mineral dust according to their chemical composition (Formenti et al., 2014). The variability of the Fe/Al ratio was relatively low (0.92) during non-dust events and decreased to about 0.69 during the dust events, as shown in Fig 10b. The Fe/Ca ratio was also used to make distinctions amongst sources. It was found that [on average](#), the Fe/Ca ratio was 1.4, [higher than 1](#) during dust events, [while Fe/Ca was and averagely 0.4 during background conditions](#) [non-dust events samples](#). These ratios, which are robust indicators of large-scale mineral dust source variation, were supported by air mass backward trajectories.

900 The [back trajectory analysis indicate](#) [indicates that](#) particles during dust events came from the Saharan region of Mauritania and southern Morocco as also [indicated highlighted](#) by their high Fe/Ca ratio while mineral dust during [background conditions](#) [BAM](#) as [described indicated](#) above was of local sources emitted from road-dust or resuspension from agricultural activities. [Fe/Ca ratios close to 0.4 indicate that the dust comes from the south, while Fe/Ca ratios higher than 1 indicate that the dust comes from the East. The average Fe/Ca ratio obtained at the station was 0.54, which suggests that the dust often originated from south Morocco. During the dust event, the Fe/Ca ratio reached 1.9, which suggests that long-distance transport of the dust from the eastern Saharan regions was observed, which is in agreement with the pollution rose, which indicated high concentrations from southeast winds.](#)

910 An increase in the concentration of many ions was also observed during dust events. sulfate, nitrate, calcium, ammonium, showed an increase in the average concentration of about a factor of 4.5 while chloride, magnesium, and potassium experienced an increase in their concentrations by a factor of 5 (Fig 10c, 10d). During dust events, sodium concentration also experienced an increase by a factor of 3. Its correlation with iron and aluminum suggested its possible soil origin. Other metals of anthropogenic origin (not plotted), such as Cu, Ni, and Pb, showed no significant difference between dust and non-dust events.

#### Organics

920 Figure (10c) reveals that organic carbon increased averagely during the dust event from 0.5 to 3.5  $\mu\text{g m}^{-3}$ . Samples collected during the dust period showed a strong correlation between organic matter and other elements of crustal origin such as nss-Ca<sup>2+</sup> and nss-Mg<sup>2+</sup>. The OC/EC ratio was in the range of 4-6 during the dust period, while the ratio was lower than 3 during [background conditions](#) [BAM](#) which suggests that the organic fraction was affected by desert dust particles. [Correlation of mineral dust and OC](#) Specifically, there is a clear correlation between Fe, Mn, Al, K<sup>+</sup>, Ca<sup>2+</sup> and OC<sup>22</sup> to a less extent, Mg<sup>2+</sup>. During this period, a correlation was observed between OC as well as some organic species Nonacosane and Heptacosane with mineral elements such as calcium ( $r^2=0.81$ ) and magnesium ( $r^2=0.73$ ). During the winter, this relationship became practically insignificant ( $r^2=0.15$ ). This suggests a possible common origin of this species that is the crustal mineral aerosol as it is also confirmed by the low enrichment factor of all the species mentioned. Some organic compounds increase during dust events,

930 especially odd alkanes such as nonacosane, hentriacontane, heptacosane, and tricosane by a factor of 4. In contrast, elemental carbon remains globally constant with marginal changes (Fig 10e). The PAH fluorene, showed similar concentrations during dust events and background conditions, indicating that Saharan dust was not a significant source. An increase of about 65% was observed for pentacosane, octacosane, hexacosane and decosane. Due to the biodiversity of several plant species and remarkable microbiological activity surrounding  
935 the Atlas regions, an increase of some organic compounds and organic matter fraction suggests that mineral dust was loaded with biogenic compounds during dust transport.

#### 4 Conclusion

In the present study, PM<sub>10</sub> particulate matter was chemically characterized at the newly established AM5 research station located in the Middle Atlas region (Morocco) at an altitude of 2100 m from August to December 2017.  
940 The aerosol chemical composition was evaluated during remote background conditions and the main air mass origins were identified. The data shows an overview of the background chemical composition and the different sources affecting aerosol composition at such a remote high-altitude site. The influence of desert dust was investigated as the site location is close to the Sahara Desert.

Despite the proximity of the site to the Saharan Desert, the influence of the desert on the atmospheric composition  
945 at this altitude was only seasonal. PM<sub>10</sub> mass concentration showed a decreasing trend with high concentrations during summer due to dust events and significantly reduced during autumn due to the washout effect from enhanced rainfall. Four main air mass inflows at the site were identified using back trajectory analysis, with each air mass distinguished by different chemical compositions. The influence of marine air mass from the [Atlantic Ocean \(NAO\)](#) [Mediterranean Sea](#) is prevalent at AM5 and [made up made-up](#) [3753%](#) of all air masses. [Background](#)  
950 [The](#) chemical composition in the Middle-Atlas during [remote conditions sampling period](#) ([60 samples](#)) is mainly dominated by locally emitted dust ([4761%](#)) with high contribution from road dust, [Organic Matter \(15%\)](#), ionic species ([716%](#)), organic matter ([745%](#)), water content (12%), and indeterminate mass ([119%](#)). Biogenic organics contributed up to 7% of the organic matter. Organic matter increased during dust events due to biogenic crustal material emissions. Diurnal variation of PM was related to the variation in a planetary boundary layer, mountain-valley winds as well as changes in different local sources. Mineral dust influenced was seasonal and ranged between 20 and [740%](#) of the mass concentration on PM<sub>10</sub> with peaks observed during the summer, accompanied  
955 by high concentrations of SO<sub>4</sub><sup>2-</sup> of up to [3.04-3  \$\mu\text{g m}^{-3}\$   \$\mu\text{g/m}^3\$](#) .

a mis en forme : Indice

a mis en forme : Non Surlignage

During winter, PM<sub>10</sub> concentrations are low, the influence of the desert is weaker, and the marine air masses are  
960 more dominant with a mixture of polluted aerosol from the coastal regions of Rabat and Casablanca and sea salt's observed. High concentrations of mineral dust were observed during the daytime due to the resuspension of road dust, while an increase of PAHs and anthropogenic metals such as Pb, Ni, and Cu were found during night-time because of the boundary layer variation. Data show that proximity to [the](#) desert does not necessarily imply constant exposure to mineral dust. Furthermore, topography and temperature variation at mountain sites control PM concentrations.

This is the first high altitude aerosol characterization study in North Africa which fills an important gap in the African region presently not available. The data from the AM5 sites thus present a reference for aerosol particle

composition under regional background conditions as well as during the influence of continental air masses.

970 Several other studies are needed to better understand the influence of the desert on the chemical composition but also the microphysical properties in the Middle Atlas region. In this study, only the chemical composition of bulk PM<sub>10</sub> particles was investigated, however, the size distribution remains an important factor and should be ~~taken into account~~ considered in further studies. An additional study on the chemical composition of the urban cities near the AM5, especially in Fes, would allow a better understanding of the anthropogenic influence in the North 975 of Morocco.

**Data availability.** All data will be made available upon request by the authors.

980 **Author contributions.** WM, HH, and SE designed the experiment at the AM5 station, KWF and ND performed the fieldwork, collected the samples, and performed the data analysis. ND performed the laboratory investigations, compiled the final Figures, and wrote the article. LP has contributed to the back trajectories calculation and the classification of air masses. ND and KWF undertook the results interpretation to which HH contributed. SZ contributed to text improvement for certain sections of the manuscript. All authors reviewed, edited, and contributed to the article. WM, HH, and SE designed the experiment at the AM5 station; KWF and ND collected the samples and performed the data analysis. ND performed the laboratory investigations, compiled the final figures, and prepared the article. All authors reviewed and edited the article and contributed to the discussion.

985

**Competing interests.** The authors declare that they have no conflict of interest.

990 **Acknowledgements.** The authors would like to thank Ibrahim Ouchen, Sayf El Islam Barcha, and Mehdi El Baramoussi for assistance in sample collection at the AM5. The authors would also acknowledge the support of Kangwei Li, Julia Wilk, Susanne Fuchs, Sylvia Haferkorn, and Cornelia Pielok for their support with the trace metal analysis.

**Financial support.** This research has been supported by the European Union's Horizon 2020 research and innovation programme (MARSU, grant no. 690958)

995 The publication of this article was funded by the Open Access Fund of the Leibniz Association.

## 5 References

- Aas, W., Tsyro, S., Bieber, E., Bergström, R., Ceburnis, D., Ellermann, T., Fagerli, H., Frölich, M., Gehrig, R., Makkonen, U., Nemitz, E., Otjes, R., Perez, N., Perrino, C., Prévôt, A. S. H., Putaud, J.-P., Simpson, D., Spindler, G., Vana, M., and Yttri, K. E.: Lessons learnt from the first EMEP intensive measurement periods, 12, 8073–8094, <https://doi.org/10.5194/acp-12-8073-2012>, 2012.
- Ajdour, A., Leghrif, R., Chaoufi, J., Chirmata, A., Menut, L., and Mailler, S.: Towards air quality modeling in Agadir City (Morocco), Materials Today: Proceedings, S2214785319326227, <https://doi.org/10.1016/j.matpr.2019.07.438>, 2019.
- Alves, C., Vicente, A., Pio, C., Kiss, G., Hoffer, A., Decesari, S., Prevôt, A. S. H., Minguillón, M. C., Querol, X., Hillamo, R., Spindler, G., and Swietlicki, E.: Organic compounds in aerosols from selected European sites – Biogenic versus anthropogenic sources, Atmospheric Environment, 59, 243–255, <https://doi.org/10.1016/j.atmosenv.2012.06.013>, 2012.
- Ambrose, J. L., Reidmiller, D. R., and Jaffe, D. A.: Causes of high O<sub>3</sub> in the lower free troposphere over the Pacific Northwest as observed at the Mt. Bachelor Observatory, Atmospheric Environment, 45, 5302–5315, <https://doi.org/10.1016/j.atmosenv.2011.06.056>, 2011.
- Amodio, M., Catino, S., Dambruoso, P. R., de Gennaro, G., Di Gilio, A., Giungato, P., Laiola, E., Marzocca, A., Mazzone, A., Sardaro, A., and Tutino, M.: Atmospheric Deposition: Sampling Procedures, Analytical Methods, and Main Recent Findings from the Scientific Literature, Advances in Meteorology, 2014, 1–27, <https://doi.org/10.1155/2014/161730>, 2014.
- Ansmann, A., Petzold, A., Kandler, K., Tegen, I., Wendisch, M., Müller, D., Weinzierl, B., Müller, T., and Heintzenberg, J.: Saharan Mineral Dust Experiments SAMUM-1 and SAMUM-2: what have we learned?, 63, 403–429, <https://doi.org/10.1111/j.1600-0889.2011.00555.x>, 2011.
- Ansmann, A., Tesche, M., Knippertz, P., Bierwirth, E., Althausen, D., MüLLER, D., and Schulz, O.: Vertical profiling of convective dust plumes in southern Morocco during SAMUM, 61, 340–353, <https://doi.org/10.1111/j.1600-0889.2008.00384.x>, 2009.
- Arimoto, R., Kim, Y. J., Kim, Y. P., Quinn, P. K., Bates, T. S., Anderson, T. L., Gong, S., Uno, I., Chin, M., Huebert, B. J., Clarke, A. D., Shinohara, Y., Weber, R. J., Anderson, J. R., Guazzotti, S. A., Sullivan, R. C., Sodeman, D. A., Prather, K. A., and Sokolik, I. N.: Characterization of Asian Dust during ACE-Asia, Global and Planetary Change, 52, 23–56, <https://doi.org/10.1016/j.gloplacha.2006.02.013>, 2006.
- Astitha, M., Kallos, G., Spyrou, C., O'Hirok, W., Lelieveld, J., and Denier van der Gon, H. A. C.: Modelling the chemically aged and mixed aerosols over the eastern central Atlantic Ocean – potential impacts, Atmos. Chem. Phys., 10, 5797–5822, <https://doi.org/10.5194/acp-10-5797-2010>, 2010.
- Baker, K. and Scheff, P.: Photochemical model performance for PM2.5 sulfate, nitrate, ammonium, and precursor species SO<sub>2</sub>, HNO<sub>3</sub>, and NH<sub>3</sub> at background monitor locations in the central and eastern United States, Atmospheric Environment, 41, 6185–6195, <https://doi.org/10.1016/j.atmosenv.2007.04.006>, 2007.
- Bauer, H., Schueller, E., Weinke, G., Berger, A., Hitzenberger, R., Marr, I. L., and Puxbaum, H.: Significant contributions of fungal spores to the organic carbon and to the aerosol mass balance of the urban atmospheric aerosol, Atmospheric Environment, 42, 5542–5549, <https://doi.org/10.1016/j.atmosenv.2008.03.019>, 2008.
- Bei, N., Zhao, L., Wu, J., Li, X., Feng, T., and Li, G.: Impacts of sea-land and mountain-valley circulations on the air pollution in Beijing-Tianjin-Hebei (BTH): A case study, Environmental Pollution, 234, 429–438, <https://doi.org/10.1016/j.envpol.2017.11.066>, 2018.
- Benchrif, A., Guinot, B., Bounakhla, M., Cachier, H., Dammati, B., and Baghdad, B.: Aerosols in Northern Morocco: Input pathways and their chemical fingerprint, Atmospheric Environment, 174, 140–147, <https://doi.org/10.1016/j.atmosenv.2017.11.047>, 2018.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, J. Geophys. Res., 106, 23073–23095, <https://doi.org/10.1029/2001JD000807>, 2001.
- Bonasoni, P., Laj, P., Marinoni, A., Sprenger, M., Angelini, F., Arduini, J., Bonafè, U., Calzolari, F., Colombo, T., De Cesari, S., Di Biagio, C., di Sarra, A. G., Evangelisti, F., Duchi, R., Facchini, Mc.,

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

- Fuzzi, S., Gobbi, G. P., Maione, M., Panday, A., Roccato, F., Sellegri, K., Venzac, H., Verza, Gp., Villani, P., Vuillermoz, E., and Cristofanelli, P.: Atmospheric Brown Clouds in the Himalayas: first two years of continuous observations at the Nepal Climate Observatory-Pyramid (5079 m), *Atmos. Chem. Phys.*, 10, 7515–7531, <https://doi.org/10.5194/acp-10-7515-2010>, 2010.
- 1055 Tahri, M., Benchraf, A., Bounakhla, M., Benyaich, F., and Noack, Y.: Seasonal variation and risk assessment of PM2.5 and PM2.5–10 in the ambient air of Kenitra, Morocco, *Environ. Sci.: Processes Impacts*, 19, 1427–1436, <https://doi.org/10.1039/C7EM00286F>, 2017.
- 1060 Bourcier, L., Sellegri, K., Chausse, P., Pichon, J. M., and Laj, P.: Seasonal variation of water-soluble inorganic components in aerosols size-segregated at the puy de Dôme station (1,465 m a.s.l.), France, *J Atmos Chem*, 69, 47–66, <https://doi.org/10.1007/s10874-012-9229-2>, 2012.
- Bove, M. C., Brotto, P., Cassola, F., Cuccia, E., Massabò, D., Mazzino, A., Piazzalunga, A., and Prati, P.: An integrated PM2.5 source apportionment study: Positive Matrix Factorisation vs. the chemical transport model CAMx, *Atmospheric Environment*, 94, 274–286, <https://doi.org/10.1016/j.atmosenv.2014.05.039>, 2014.
- 1065 Calvert, J. G.: Glossary of atmospheric chemistry terms (Recommendations 1990), 62, 2167–2219, <https://doi.org/10.1351/pac199062112167>, 1990.
- Campbell, J. F. E., Fletcher, W. J., Joannin, S., Hughes, P. D., Rhanem, M., and Zielhofer, C.: Environmental Drivers of Holocene Forest Development in the Middle Atlas, Morocco, *Front. Ecol. Evol.*, 5, 113, <https://doi.org/10.3389/fevo.2017.00113>, 2017.
- 1070 Cavalli, F., Viana, M., Yttri, K. E., and Genberg, J.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, 11, 2010.
- Cesari, D., Contini, D., Genga, A., Siciliano, M., Elefante, C., Baglivi, F., and Daniele, L.: Analysis of raw soils and their re-suspended PM10 fractions: Characterisation of source profiles and enrichment factors, *Applied Geochemistry*, 27, 1238–1246, <https://doi.org/10.1016/j.apgeochem.2012.02.029>, 2012.
- 1075 Chatterjee, A., Adak, A., Singh, A. K., Srivastava, M. K., Ghosh, S. K., Tiwari, S., Devara, P. C. S., and Raha, S.: Aerosol Chemistry over a High Altitude Station at Northeastern Himalayas, India, *PLoS ONE*, 5, e11122, <https://doi.org/10.1371/journal.pone.0011122>, 2010.
- Chiapello, I., Bergametti, G., Chatenet, B., Bousquet, P., Dulac, F., and Soares, E. S.: Origins of African dust transported over the northeastern tropical Atlantic, *J. Geophys. Res.*, 102, 13701–13709, <https://doi.org/10.1029/97JD00259>, 1997.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, 59, 23, 2002.
- 1085 Chu, A. K. M., Kwok, R. C. W., and Yu, K. N.: Study of pollution dispersion in urban areas using Computational Fluid Dynamics (CFD) and Geographic Information System (GIS), *Environmental Modelling & Software*, 20, 273–277, <https://doi.org/10.1016/j.envsoft.2004.05.007>, 2005.
- Cincinelli, A., Bubba, M. D., Martellini, T., Gambaro, A., and Lepri, L.: Gas-particle concentration and distribution of n-alkanes and polycyclic aromatic hydrocarbons in the atmosphere of Prato (Italy), *Chemosphere*, 68, 472–478, <https://doi.org/10.1016/j.chemosphere.2006.12.089>, 2007.
- Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic Model of the System  $H+ - NH_4+ - SO_4^{2-} - NO_3^- - H_2O$  at Tropospheric Temperatures, *J. Phys. Chem. A*, 102, 2137–2154, <https://doi.org/10.1021/jp973042r>, 1998.
- 1090 Contini, D., Cesari, D., Donateo, A., Chirizzi, D., and Belosi, F.: Characterization of PM10 and PM2.5 and Their Metals Content in Different Typologies of Sites in South-Eastern Italy, *Atmosphere*, 5, 435–453, <https://doi.org/10.3390/atmos5020435>, 2014.
- Contini, D., Genga, A., Cesari, D., Siciliano, M., Donateo, A., Bove, M. C., and Guascito, M. R.: Characterisation and source apportionment of PM10 in an urban background site in Lecce, *Atmospheric Research*, 95, 40–54, <https://doi.org/10.1016/j.atmosres.2009.07.010>, 2010.
- 1095 Cozic, J., Verheggen, B., Weingartner, E., Crosier, J., Bower, K. N., Flynn, M., Coe, H., Henning, S., Steinbacher, M., Henne, S., Coen, M. C., Petzold, A., and Baltensperger, U.: Chemical composition of free tropospheric aerosol for PM1 and coarse mode at the high alpine site Jungfraujoch, 18, 2008.
- Decesari, S., Facchini, M. C., Carbone, C., Julianelli, L., Rinaldi, M., Finessi, E., Fuzzi, S., Marinoni, A., Cristofanelli, P., Duchi, R., Bonasoni, P., Vuillermoz, E., Cozic, J., Jaffrezo, J. L., and Laj, P.:

a mis en forme : Anglais (E.U.)

- 105 Chemical composition of PM<sub>2.5</sub>; and PM<sub>10</sub>; at the high-altitude Himalayan station Nepal Climate Observatory-Pyramid (NCO-P) (5079 m a.s.l.), Atmos. Chem. Phys., 10, 4583–4596, <https://doi.org/10.5194/acp-10-4583-2010>, 2010.
- Desboeufs, K. V. and Cautenet, G.: Transport and mixing zone of desert dust and sulphate over Tropical Africa and the Atlantic Ocean region, <https://doi.org/10.5194/acpd-5-5615-2005>, 2005.
- 110 Ding, K., Liu, J., Ding, A., Liu, Q., Zhao, T. L., Shi, J., Han, Y., Wang, H., and Jiang, F.: Uplifting of carbon monoxide from biomass burning and anthropogenic sources to the free troposphere in East Asia, Atmos. Chem. Phys., 15, 2843–2866, <https://doi.org/10.5194/acp-15-2843-2015>, 2015.
- Draxler, R., and Hess, G.: Description of the HYSPLIT4 modeling system, NOAA Technical Memorandum, ERL, ARL-224, 2004.
- 115 Du, Y., Xu, X., Chu, M., Guo, Y., and Wang, J.: Air particulate matter and cardiovascular disease: the epidemiological, biomedical and clinical evidence, 8, 12, doi: 10.3978/j.issn.2072-1439.2015.11.37 2016.
- Falkovich, A. H., Ganor, E., Levin, Z., Formenti, P., and Rudich, Y.: Chemical and mineralogical analysis of individual mineral dust particles, J. Geophys. Res., 106, 18029–18036, <https://doi.org/10.1029/2000JD900430>, 2001.
- 120 Fast, J. D., Gustafson, W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, J. Geophys. Res., 111, D21305, <https://doi.org/10.1029/2005JD006721>, 2006.
- Fomba, K. W., Deabji, N., Barcha, S. E. I., Ouchen, I., Elbaramoussi, E. M., El Moursli, R. C., Harnafi, M., El Hajjaji, S., Mellouki, A., and Herrmann, H.: Application of TXRF in monitoring trace metals in particulate matter and cloud water, Atmos. Meas. Tech., 13, 4773–4790, <https://doi.org/10.5194/amt-13-4773-2020>, 2020.
- 125 Fomba, K. W., Müller, K., van Pinxteren, D., and Herrmann, H.: Aerosol size-resolved trace metal composition in remote northern tropical Atlantic marine environment: case study Cape Verde islands, 13, 4801–4814, <https://doi.org/10.5194/acp-13-4801-2013>, 2013.
- 130 Fomba, K. W., Müller, K., van Pinxteren, D., Poulain, L., van Pinxteren, M., and Herrmann, H.: Long-term chemical characterization of tropical and marine aerosols at the Cape Verde Atmospheric Observatory (CVAO) from 2007 to 2011, 14, 8883–8904, <https://doi.org/10.5194/acp-14-8883-2014>, 2014.
- 135 Formenti, P., Caquineau, S., Desboeufs, K., Klaver, A., Chevaillier, S., Journet, E., and Rajot, J. L.: Mapping the physico-chemical properties of mineral dust in western Africa: mineralogical composition, Atmos. Chem. Phys., 14, 10663–10686, <https://doi.org/10.5194/acp-14-10663-2014>, 2014.
- 140 Fu, P., Kawamura, K., Kobayashi, M., and Simoneit, B. R. T.: Seasonal variations of sugars in atmospheric particulate matter from Gosan, Jeju Island: Significant contributions of airborne pollen and Asian dust in spring, Atmospheric Environment, 55, 234–239, <https://doi.org/10.1016/j.atmosenv.2012.02.061>, 2012.
- Gangoiti, G., Alonso, L., Navazo, M., Garcia, J. A., and Millán, M. M.: North African soil dust and European pollution transport to America during the warm season: Hidden links shown by a passive tracer simulation: EUROPEAN POLLUTION TRANSPORT TO AMERICA, J. Geophys. Res., 111, n/a-n/a, <https://doi.org/10.1029/2005JD005941>, 2006.
- 145 García, M. I., Rodríguez, S., and Alastuey, A.: Impact of North America on the aerosol composition in the North Atlantic free troposphere, Atmos. Chem. Phys., 17, 7387–7404, <https://doi.org/10.5194/acp-17-7387-2017>, 2017.
- 150 Gelencsér, A., May, B., Simpson, D., Sánchez-Ochoa, A., Kasper-Giebl, A., Puxbaum, H., Caseiro, A., Pio, C., and Legrand, M.: Source apportionment of PM<sub>2.5</sub> organic aerosol over Europe: Primary/secondary, natural/anthropogenic, and fossil/biogenic origin, J. Geophys. Res., 112, D23S04, <https://doi.org/10.1029/2006JD008094>, 2007.
- Glaccum, R. A. and Prospero, J. M.: Saharan aerosols over the tropical North Atlantic — Mineralogy, Marine Geology, 37, 295–321, [https://doi.org/10.1016/0025-3227\(80\)90107-3](https://doi.org/10.1016/0025-3227(80)90107-3), 1980.
- 155 Glasius, M., Hansen, A. M. K., Claeys, M., Henzing, J. S., Jedynska, A. D., Kasper-Giebl, A., Kistler, M., Kristensen, K., Martinsson, J., Maenhaut, W., Nejgaard, J. K., Spindler, G., Stenström, K. E., Swietlicki, E., Szidat, S., Simpson, D., and Yttri, K. E.: Composition and sources of carbonaceous aerosols in Northern Europe during winter, Atmospheric Environment, 173, 127–141, <https://doi.org/10.1016/j.atmosenv.2017.11.005>, 2018.

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

a mis en forme : Anglais (E.U.)

- Golly, B., Waked, A., Weber, S., Samake, A., Jacob, V., Conil, S., Rangognio, J., Chrétien, E., Vagnot, M.-P., Robic, P.-Y., Besombes, J.-L., and Jaffrezo, J.-L.: Organic markers and OC source apportionment for seasonal variations of PM<sub>2.5</sub> at 5 rural sites in France, *Atmospheric Environment*, 198, 142–157, <https://doi.org/10.1016/j.atmosenv.2018.10.027>, 2019.
- 1165 Hien, P. D., Bac, V. T., Tham, H. C., Nhan, D. D., and Vinh, L. D.: Influence of meteorological conditions on PM<sub>2.5</sub> and PM<sub>2.5</sub>–<sub>10</sub> concentrations during the monsoon season in Hanoi, Vietnam, 12, [https://doi.org/10.1016/S1352-2310\(02\)00295-9](https://doi.org/10.1016/S1352-2310(02)00295-9), 2002.
- Holst, J., Mayer, H., and Holst, T.: Effect of meteorological exchange conditions on PM<sub>10</sub> concentration, *metz*, 17, 273–282, <https://doi.org/10.1127/0941-2948/2008/0283>, 2008.
- 1170 Iinuma, Y., Brüggemann, E., Gnauk, T., Müller, K., Andreae, M. O., Helas, G., Parmar, R., and Herrmann, H.: Source characterization of biomass burning particles: The combustion of selected European conifers, African hardwood, savanna grass, and German and Indonesian peat, *J. Geophys. Res.*, 112, D08209, <https://doi.org/10.1029/2006JD007120>, 2007.
- 1175 Iinuma, Y., Engling, G., Puxbaum, H., and Herrmann, H.: A highly resolved anion-exchange chromatographic method for determination of saccharide tracers for biomass combustion and primary bio-particles in atmospheric aerosol, *Atmospheric Environment*, 43, 1367–1371, <https://doi.org/10.1016/j.atmosenv.2008.11.020>, 2009.
- Inchaouh, M.: state of ambient air quality in marrakech city (morocco) over the period 2009 – 2012, *geomate*, <https://doi.org/10.21660/2017.29.1254>, 2017.
- 1180 Jaenicke, R.: Abundance of Cellular Material and Proteins in the Atmosphere, 308, 73–73, <https://doi.org/10.1126/science.1106335>, 2005.
- Kalderon-Asael, B., Erel, Y., Sandler, A., and Dayan, U.: Mineralogical and chemical characterization of suspended atmospheric particles over the east Mediterranean based on synoptic-scale circulation patterns, *Atmospheric Environment*, 43, 3963–3970, <https://doi.org/10.1016/j.atmosenv.2009.03.057>, 2009.
- 1185 Kandler, K., Benker, N., Bundke, U., Cuevas, E., Ebert, M., Knippertz, P., Rodríguez, S., Schütz, L., and Weinbruch, S.: Chemical composition and complex refractive index of Saharan Mineral Dust at Izaña, Tenerife (Spain) derived by electron microscopy, *Atmospheric Environment*, 41, 8058–8074, <https://doi.org/10.1016/j.atmosenv.2007.06.047>, 2007.
- 1190 Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., JäCKEL, S., Jaenicke, R., Knippertz, P., Lieke, K., Massling, A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at Tinfou, Morocco, during SAMUM 2006, *Tellus B: Chemical and Physical Meteorology*, 61, 32–50, <https://doi.org/10.1111/j.1600-0889.2008.00385.x>, 2009.
- 1195 Kavouras, I. G.: Particle size distribution of organic primary and secondary aerosol constituents in urban, background marine, and forest atmosphere, *J. Geophys. Res.*, 107, 4069, <https://doi.org/10.1029/2000JD000278>, 2002.
- Kchih, H., Perrino, C., and Cherif, S.: Investigation of Desert Dust Contribution to Source Apportionment of PM<sub>10</sub> and PM<sub>2.5</sub> from a Southern Mediterranean Coast, *Aerosol Air Qual. Res.*, 15, 454–464, <https://doi.org/10.4209/aaqr.2014.10.0255>, 2015.
- 1200 Khan, B., Stenckov, G., Weinzierl, B., Kalenderski, S., and Osipov, S.: Dust plume formation in the free troposphere and aerosol size distribution during the Saharan Mineral Dust Experiment in North Africa, *Tellus B: Chemical and Physical Meteorology*, 67, 27170, <https://doi.org/10.3402/tellusb.v67.27170>, 2015.
- 1205 Khedidji, S., Müller, K., Rabhi, L., Spindler, G., Fomba, K. W., Pinxteren, D. van, Yassaa, N., and Herrmann, H.: Chemical Characterization of Marine Aerosols in a South Mediterranean Coastal Area Located in Bou Ismaïl, Algeria, *Aerosol Air Qual. Res.*, 20, 2448–2473, <https://doi.org/10.4209/aaqr.2019.09.0458>, 2020.
- 1210 Khrissi, S., Bejjit, L., Haddad, M., Falguères, C., Ait Lyazidi, S., and El Amraoui, M.: Study of marbles from Middle Atlas (Morocco): elemental, mineralogical and structural analysis, *IOP Conf. Ser.: Mater. Sci. Eng.*, 353, 012013, <https://doi.org/10.1088/1757-899X/353/1/012013>, 2018.
- King, M. D., Menzel, W. P., Kaufman, Y. J., Tanre, D., Bo-Cai Gao, Platnick, S., Ackerman, S. A., Remer, L. A., Pincus, R., and Hubanks, P. A.: Cloud and aerosol properties, precipitable water, and

a mis en forme : Anglais (E.U.)

- 1215 profiles of temperature and water vapor from MODIS, IEEE Trans. Geosci. Remote Sensing, 41, 442–458, <https://doi.org/10.1109/TGRS.2002.808226>, 2003.
- 1220 Knippertz, P., Deutscher, C., Kandler, K., Müller, T., Schulz, O., and Schütz, L.: Dust mobilization due to density currents in the Atlas region: Observations from the Saharan Mineral Dust Experiment 2006 field campaign, J. Geophys. Res., 112, D21109, <https://doi.org/10.1029/2007JD008774>, 2007.
- Kocak, M., Theodosi, C., Zarmpas, P., Séguret, M. J. M., Herut, B., Kallos, G., Mihalopoulos, N., Kubilay, N., and Nimmo, M.: Influence of mineral dust transport on the chemical composition and physical properties of the Eastern Mediterranean aerosol, Atmospheric Environment, 57, 266–277, <https://doi.org/10.1016/j.atmosenv.2012.04.006>, 2012.
- 1225 Kreidenweis, S. M., Petters, M. D., and DeMott, P. J.: Single-parameter estimates of aerosol water content, Environ. Res. Lett., 3, 035002, <https://doi.org/10.1088/1748-9326/3/3/035002>, 2008.
- Krueger, B. J., Grassian, V. H., Cowin, J. P., and Laskin, A.: Heterogeneous chemistry of individual mineral dust particles from different dust source regions: the importance of particle mineralogy, Atmospheric Environment, 38, 6253–6261, <https://doi.org/10.1016/j.atmosenv.2004.07.010>, 2004.
- 1230 Lang, M. N., Gohm, A., and Wagner, J. S.: The impact of embedded valleys on daytime pollution transport over a mountain range, 17, <https://doi.org/10.5194/acp-15-11981-2015>, 2015.
- Leena, P. P., Vijayakumar, K., Anilkumar, V., and Pandithurai, G.: Analysing temporal variability of particulate matter and possible contributing factors over Mahabaleshwar, a high-altitude station in Western Ghats, India, Journal of Atmospheric and Solar-Terrestrial Physics, 164, 105–115, <https://doi.org/10.1016/j.jastp.2017.08.013>, 2017.
- 1235 Leena, P. P., Vijayakumar, K., Anilkumar, V., and Pandithurai, G.: Analysing temporal variability of particulate matter and possible contributing factors over Mahabaleshwar, a high-altitude station in Western Ghats, India, Journal of Atmospheric and Solar-Terrestrial Physics, 164, 105–115, <https://doi.org/10.1016/j.jastp.2017.08.013>, 2017.
- 1240 Leng, C., Zhang, Q., Tao, J., Zhang, H., Zhang, D., Xu, C., Li, X., Kong, L., Cheng, T., Zhang, R., Yang, X., Chen, J., Qiao, L., Lou, S., Wang, H., and Chen, C.: Impacts of new particle formation on aerosol cloud condensation nuclei (CCN) activity in Shanghai: case study, Atmos. Chem. Phys., 14, 11353–11365, <https://doi.org/10.5194/acp-14-11353-2014>, 2014.
- Liang, Q., Jaeglé, L., Jaffe, D. A., Weiss-Penzias, P., Heckman, A., and Snow, J. A.: Long-range transport of Asian pollution to the northeast Pacific: Seasonal variations and transport pathways of carbon monoxide: transport pathways to the northeast pacific, J. Geophys. Res., 109, <https://doi.org/10.1029/2003JD004402>, 2004.
- 1245 Liu, J., Russell, L. M., Lee, A. K. Y., McKinney, K. A., Surratt, J. D., and Ziemann, P. J.: Observational evidence for pollution-influenced selective uptake contributing to biogenic secondary organic aerosols in the southeastern U.S.: Evidence for Selective Uptake of bSOA, Geophys. Res. Lett., 44, 8056–8064, <https://doi.org/10.1002/2017GL074665>, 2017.
- Maenhaut, W., Raes, N., Chi, X., Cafmeyer, J., Wang, W., and Salma, I.: Chemical composition and mass closure for fine and coarse aerosols at a kerbside in Budapest, Hungary, in spring 2002, 34, 290–296, <https://doi.org/10.1002/xrs.820>, 2005.
- 1250 Marenci, F., Bonasoni, P., Calzolari, F., Ceriani, M., Chiari, M., Cristofanelli, P., D'Alessandro, A., Fermo, P., Lucarelli, F., Mazzei, F., Nava, S., Piazzalunga, A., Prati, P., Valli, G., and Vecchi, R.: Characterization of atmospheric aerosols at Monte Cimone, Italy, during summer 2004: Source apportionment and transport mechanisms, J. Geophys. Res., 111, D24202, <https://doi.org/10.1029/2006JD007145>, 2006.
- 1255 Marzi, R., Torkelson, B. E., and Olson, R. K.: A revised carbon preference index, Organic Geochemistry, 20, 1303–1306, [https://doi.org/10.1016/0146-6380\(93\)90016-5](https://doi.org/10.1016/0146-6380(93)90016-5), 1993.
- Mazzei, F., D'Alessandro, A., Lucarelli, F., Nava, S., Prati, P., Valli, G., and Vecchi, R.: Characterization of particulate matter sources in an urban environment, Sci Total Environ, 401, 81–89, <https://doi.org/10.1016/j.scitotenv.2008.03.008>, 2008.
- McInnes, L. M., Covert, D. S., Quinn, P. K., and Germani, M. S.: Measurements of chloride depletion and sulfur enrichment in individual sea salt particles collected from the remote marine boundary layer, 12, <https://doi.org/93JD03453>, 1994.
- Miche, H., Saracco, G., Mayer, A., Qarqori, K., Rouai, M., Dekayir, A., Chalikakis, K., and Emblanch, C.: Hydrochemical constraints between the karst Tabular Middle Atlas Causses and the Saïs basin

a mis en forme : Anglais (E.U.)

- 1270 (Morocco): implications of groundwater circulation, *Hydrogeol J*, 26, 71–87, <https://doi.org/10.1007/s10040-017-1675-0>, 2018.
- Minguillón, M. C., Querol, X., Alastuey, A., Monfort, E., and Miró, J. V.: PM sources in a highly industrialised area in the process of implementing PM abatement technology. Quantification and evolution, 9, 1071–1081, <https://doi.org/10.1039/B705474B>, 2007.
- 1275 Mounir, S., Saoud, N., Charroud, M., Mounir, K., and Choukrad, J.: The Middle Atlas Geological karsts forms: Towards Geosites characterization, *Oil Gas Sci. Technol. – Rev. IFP Energies nouvelles*, 74, 17, <https://doi.org/10.2516/ogst/2018089>, 2019.
- Müller, K.: Determination of aldehydes and ketones in the atmosphere — A comparative long time study at an urban and a rural site in Eastern Germany, *Chemosphere*, 35, 2093–2106, [https://doi.org/10.1016/S0045-6535\(97\)00267-1](https://doi.org/10.1016/S0045-6535(97)00267-1), 1997.
- 1280 Nair, V. S., Moorthy, K. K., Alappattu, D. P., Kunhikrishnan, P. K., George, S., Nair, P. R., Babu, S. S., Abish, B., Satheesh, S. K., Tripathi, S. N., Nirjanan, K., Madhavan, B. L., Srikant, V., Dutt, C. B. S., Badarinath, K. V. S., and Reddy, R. R.: Wintertime aerosol characteristics over the Indo-Gangetic Plain (IGP): Impacts of local boundary layer processes and long-range transport: winter aerosols over indo-gangetic plain, *J. Geophys. Res.*, 112, n/a-n/a, <https://doi.org/10.1029/2006JD008099>, 2007.
- Nerriere, É., Guegan, H., Bordigoni, B., Hautemaniere, A., Momas, I., Ladner, J., Target, A., Lameloise, P., Delmas, V., Personnaz, M.-B., Koutrakis, P., and Zmirou-Navier, D.: Spatial heterogeneity of personal exposure to airborne metals in French urban areas, *Science of The Total Environment*, 373, 49–56, <https://doi.org/10.1016/j.scitotenv.2006.10.042>, 2007.
- 1290 Neusüss, C., Pelzing, M., Plewka, A., and Herrmann, H.: A new analytical approach for size-resolved speciation of organic compounds in atmospheric aerosol particles: Methods and first results, *J. Geophys. Res.*, 105, 4513–4527, <https://doi.org/10.1029/1999JD901038>, 2000.
- Noulas, C., Tziouvalekas, M., and Karyotis, T.: Zinc in soils, water and food crops, *Journal of Trace Elements in Medicine and Biology*, 49, 252–260, <https://doi.org/10.1016/j.jtemb.2018.02.009>, 2018.
- Nourelbait, M., Rhoujati, A., Benkaddour, A., Carré, M., Eynaud, F., Martinez, P., and Cheddadi, R.: Climate change and ecosystems dynamics over the last 6000 years in the Middle Atlas, Morocco, *Clim. Past*, 12, 1029–1042, <https://doi.org/10.5194/cp-12-1029-2016>, 2016.
- 1300 Okada, K. and Kai, K.: Atmospheric mineral particles collected at Qira in the Taklamakan Desert, China, *Atmospheric Environment*, 38, 6927–6935, <https://doi.org/10.1016/j.atmosenv.2004.03.078>, 2004.
- Pacyna, E. G., Pacyna, J. M., Fudala, J., Strzelecka-Jastrzab, E., Hlawiczka, S., Panasiuk, D., Nitter, S., Pregger, T., Pfeiffer, H., and Friedrich, R.: Current and future emissions of selected heavy metals to the atmosphere from anthropogenic sources in Europe, *Atmospheric Environment*, 41, 8557–8566, <https://doi.org/10.1016/j.atmosenv.2007.07.040>, 2007.
- 1305 Pandolfi, M., Gonzalez-Castañedo, Y., Alastuey, A., de la Rosa, J. D., Mantilla, E., de la Campa, A. S., Querol, X., Pey, J., Amato, F., and Moreno, T.: Source apportionment of PM10 and PM2.5 at multiple sites in the strait of Gibraltar by PMF: impact of shipping emissions, *Environ Sci Pollut Res*, 18, 260–269, <https://doi.org/10.1007/s11356-010-0373-4>, 2011.
- Parrish, D. D., Law, K. S., Staehelin, J., Derwent, R., Cooper, O. R., Tanimoto, H., Volz-Thomas, A., Gilge, S., Scheel, H.-E., Steinbacher, M., and Chan, E.: Long-term changes in lower tropospheric baseline ozone concentrations at northern mid-latitudes, *Atmos. Chem. Phys.*, 12, 11485–11504, <https://doi.org/10.5194/acp-12-11485-2012>, 2012.
- 1315 Perrino, C., Catrambone, M., Dalla Torre, S., Rantica, E., Sargolini, T., and Canepari, S.: Seasonal variations in the chemical composition of particulate matter: a case study in the Po Valley. Part I: macro-components and mass closure, *Environ Sci Pollut Res*, 21, 3999–4009, <https://doi.org/10.1007/s11356-013-2067-1>, 2014.
- Pietrogrande, M. C., Abbaszade, G., Schnelle-Kreis, J., Bacco, D., Mercuriali, M., and Zimmermann, R.: Seasonal variation and source estimation of organic compounds in urban aerosol of Augsburg, Germany, *Environmental Pollution*, 159, 1861–1868, <https://doi.org/10.1016/j.envpol.2011.03.023>, 2011.
- Pietrogrande, M. C., Mercuriali, M., Perrone, M. G., Ferrero, L., Sangiorgi, G., and Bolzacchini, E.: Distribution of n-Alkanes in the Northern Italy Aerosols: Data Handling of GC-MS Signals for

a mis en forme : Anglais (E.U.)

- 1325 Homologous Series Characterization, Environ. Sci. Technol., 44, 4232–4240, https://doi.org/10.1021/es1001242, 2010.
- Pio, C. A., Alves, C. A., and Duarte, A. C.: Identification, abundance and origin of atmospheric organic particulate matter in a Portuguese rural area, Atmospheric Environment, 35, 1365–1375, https://doi.org/10.1016/S1352-2310(00)00391-5, 2001. a mis en forme : Anglais (E.U.)
- 1330 Pope, C. A., Cohen, A. J., and Burnett, R. T.: Cardiovascular Disease and Fine Particulate Matter: Lessons and Limitations of an Integrated Exposure–Response Approach, Circ Res, 122, 1645–1647, https://doi.org/10.1161/circresaha.118.312956, 2018.
- Prodi, F., Belosi, F., Contini, D., Santachiara, G., Matteo, L. D., Gambaro, A., Donateo, A., and Cesari, D.: Aerosol fine fraction in the Venice Lagoon: Particle composition and sources, 10, https://doi.org/10.1016/j.atmosres.2008.09.020, 2009. a mis en forme : Anglais (E.U.)
- 1335 Putero, D., Cristofanelli, P., Marinoni, A., Adhikary, B., Duchi, R., Shrestha, S. D., Verza, G. P., Landi, T. C., Calzolari, F., Busetto, M., Agnello, G., Biancofiore, F., Di Carlo, P., Panday, A. K., Rupakheti, M., and Bonasoni, P.: Seasonal variation of ozone and black carbon observed at Paknajol, an urban site in the Kathmandu Valley, Nepal, Atmos. Chem. Phys., 15, 13957–13971, https://doi.org/10.5194/acp-15-13957-2015, 2015. a mis en forme : Anglais (E.U.)
- 340 Querol, X., Alastuey, A., Ruiz, C. R., Artiñano, B., Hansson, H. C., Harrison, R. M., Buringh, E., ten Brink, H. M., Lutz, M., Bruckmann, P., Straehl, P., and Schneider, J.: Speciation and origin of PM10 and PM2.5 in selected European cities, Atmospheric Environment, 38, 6547–6555, https://doi.org/10.1016/j.atmosenv.2004.08.037, 2004.
- 345 Ricciardelli, I., Bacco, D., Rinaldi, M., Bonafè, G., Scotto, F., Trentini, A., Bertacci, G., Ugolini, P., Zigola, C., Rovere, F., Maccone, C., Pironi, C., and Poluzzi, V.: A three-year investigation of daily PM2.5 main chemical components in four sites: the routine measurement program of the Supersito Project (Po Valley, Italy), Atmospheric Environment, 152, 418–430, https://doi.org/10.1016/j.atmosenv.2016.12.052, 2017.
- 350 Rodriguez, S., Alastuey, A., Alonso-Pérez, S., Querol, X., Cuevas, E., Abreu-Afonso, J., Viana, M., Pérez, N., Pandolfi, M., and de la Rosa, J.: Transport of desert dust mixed with North African industrial pollutants in the subtropical Saharan Air Layer, Atmos. Chem. Phys., 11, 6663–6685, https://doi.org/10.5194/acp-11-6663-2011, 2011.
- 355 Rodriguez, S., Gonzalez, Y., Cuevas, E., Ramos, R., Romero, P. M., Abreu-Afonso, J., and Redondas, A.: Atmospheric nanoparticle observations in the low free troposphere during upward orographic flows at Izan˜a Mountain Observatory, 17, https://doi.org/10.5194/acp-9-6319-2009, 2009.
- Royaume du Maroc: Plan national de lutte contre le réchauffement climatique, Ministère de l’Énergie, des Mines, de l’Eau et de l’Environnement, Rabat, 2009.
- 360 Sarkar, S., Chauhan, A., Kumar, R., and Singh, R. P.: Impact of Deadly Dust Storms (May 2018) on Air Quality, Meteorological, and Atmospheric Parameters Over the Northern Parts of India, GeoHealth, 3, 67–80, https://doi.org/10.1029/2018GH000170, 2019. a mis en forme : Anglais (E.U.)
- Satheesh, S. K. and Krishna Moorthy, K.: Radiative effects of natural aerosols: A review, Atmospheric Environment, 39, 2089–2110, https://doi.org/10.1016/j.atmosenv.2004.12.029, 2005.
- 365 Satsangi, P. G., Chavan, S. P., Rao, P. S. P., and Safai, P. D.: Chemical characterization of particulate matter at Sinhagad, a high altitude station in Pune, India, 9, 2014.
- Scerri, M. M., Kandler, K., and Weinbruch, S.: Disentangling the contribution of Saharan dust and marine aerosol to PM10 levels in the Central Mediterranean, Atmospheric Environment, 147, 395–408, https://doi.org/10.1016/j.atmosenv.2016.10.028, 2016.
- 370 Schepanski, K., Mallet, M., Heinold, B., and Ulrich, M.: North African dust transport toward the western Mediterranean basin: atmospheric controls on dust source activation and transport pathways during June–July 2013, Atmos. Chem. Phys., 16, 14147–14168, https://doi.org/10.5194/acp-16-14147-2016, 2016. a mis en forme : Anglais (E.U.)
- 375 Sharma, S. K., Choudhary, N., Kotnala, G., Das, D., Mukherjee, S., Ghosh, A., Vijayan, N., Rai, A., Chatterjee, A., and Mandal, T. K.: Wintertime carbonaceous species and trace metals in PM10 in Darjeeling: A high altitude town in the eastern Himalayas, Urban Climate, 34, 100668, https://doi.org/10.1016/j.uclim.2020.100668, 2020.
- Sokolik, I. N. and Toon, O. B.: Direct radiative forcing by anthropogenic airborne mineral aerosols, Nature, 381, 681–683, https://doi.org/10.1038/381681a0, 1996.

- 1380 Song, Q., Christiani, D., Xiaorong Wang, and Ren, J.: The Global Contribution of Outdoor Air Pollution to the Incidence, Prevalence, Mortality and Hospital Admission for Chronic Obstructive Pulmonary Disease: A Systematic Review and Meta-Analysis, *IJERPH*, 11, 11822–11832, <https://doi.org/10.3390/ijerph111111822>, 2014.
- 1385 Spindler, G., Brüggemann, E., Gnauk, T., Grüner, A., Müller, K., and Herrmann, H.: A four-year size-segregated characterization study of particles PM10, PM2.5 and PM1 depending on air mass origin at Melpitz, *Atmospheric Environment*, 44, 164–173, <https://doi.org/10.1016/j.atmosenv.2009.10.015>, 2010.
- 1390 Spindler, G., Gnauk, T., Grüner, A., Iinuma, Y., Müller, K., Scheinhardt, S., and Herrmann, H.: Size-segregated characterization of PM10 at the EMEP site Melpitz (Germany) using a five-stage impactor: a six year study, *J Atmos Chem*, 69, 127–157, <https://doi.org/10.1007/s10874-012-9233-6>, 2012.
- Squizzato, S., Masiol, M., Brunelli, A., Pistollato, S., Tarabotti, E., Rampazzo, G., and Pavoni, B.: Factors determining the formation of secondary inorganic aerosol: a case study in the Po Valley (Italy), *Atmos. Chem. Phys.*, 13, 1927–1939, <https://doi.org/10.5194/acp-13-1927-2013>, 2013.
- 1395 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bull. Amer. Meteor. Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- Tahri, M., Bounakhlia, M., Zghaïd, M., Benchraf, A., Zahry, F., Noack, Y., and Benyâich, F.: TXRF characterization and source identification by positive matrix factorization of airborne particulate matter sampled in Kenitra City (Morocco): TXRF characterization and source identification in Kenitra City, Morocco, 42, 284–289, <https://doi.org/10.1002/xrs.2484>, 2013.
- Tiwari, S., Hopke, P. K., Pipal, A. S., Srivastava, A. K., Bisht, D. S., Tiwari, S., Singh, A. K., Soni, V. K., and Attri, S. D.: Intra-urban variability of particulate matter (PM2.5 and PM10) and its relationship with optical properties of aerosols over Delhi, India, *Atmospheric Research*, 166, 223–232, <https://doi.org/10.1016/j.atmosres.2015.07.007>, 2015.
- 1400 1405 Turpin, B. J. and Lim, H.-J.: Species Contributions to PM2.5 Mass Concentrations: Revisiting Common Assumptions for Estimating Organic Mass, *Aerosol Science and Technology*, 35, 602–610, <https://doi.org/10.1080/02786820119445>, 2001.
- Van Pinxteren, D., Brüggemann, E., Gnauk, T., Müller, K., Thiel, C., and Herrmann, H.: A GIS based approach to back trajectory analysis for the source apportionment of aerosol constituents and its first application, *J Atmos Chem*, 67, 1–28, <https://doi.org/10.1007/s10874-011-9199-9>, 2010.
- 1410 Wang, Y. Q., Zhang, X. Y., Sun, J. Y., Zhang, X. C., Che, H. Z., and Li, Y.: Spatial and temporal variations of the concentrations of PM 10 , PM 2.5 and PM 1 in China, *Atmos. Chem. Phys.*, 15, 13585–13598, <https://doi.org/10.5194/acp-15-13585-2015>, 2015.
- Waples, D. W.: Correlations, in: *Geochemistry in Petroleum Exploration*, edited by: Waples, D. W., Springer Netherlands, Dordrecht, 155–180, [https://doi.org/10.1007/978-94-009-5436-6\\_10](https://doi.org/10.1007/978-94-009-5436-6_10), 1985.
- 1415 Wedepohl, K. H.: The composition of the continental crust, 16, 1995.
- Weiss-Penzias, P., Jaffe, D. A., Swartzendruber, P., Dennison, J. B., Chand, D., Hafner, W., and Prestbo, E.: Observations of Asian air pollution in the free troposphere at Mount Bachelor Observatory during the spring of 2004: observations at mt. bachelor observatory, *j. Geophys. Res.*, 111, n/a-n/a, <https://doi.org/10.1029/2005JD006522>, 2006.
- 1420 Yttri, K. E., Simpson, D., Bergström, R., Kiss, G., Szidat, S., Cebrunis, D., Eckhardt, S., Hueglin, C., Nøjgaard, J. K., Perrino, C., Pisso, I., Prevot, A. S. H., Putaud, J.-P., Spindler, G., Vana, M., Zhang, Y.-L., and Aas, W.: The EMEP Intensive Measurement Period campaign, 2008–2009: characterizing carbonaceous aerosol at nine rural sites in Europe, *Atmos. Chem. Phys.*, 19, 4211–4233, <https://doi.org/10.5194/acp-19-4211-2019>, 2019.
- Zhang, J. M., Wang, T., Ding, A. J., Zhou, X. H., Xue, L. K., Poon, C. N., Wu, W. S., Gao, J., Zuo, H. C., Chen, J. M., Zhang, X. C., and Fan, S. J.: Continuous measurement of peroxyacetyl nitrate (PAN) in suburban and remote areas of western China, *Atmospheric Environment*, 43, 228–237, <https://doi.org/10.1016/j.atmosenv.2008.09.070>, 2009.
- 1425 1430 Zhao, Z., Cao, J., Shen, Z., Xu, B., Zhu, C., Chen, L.-W. A., Su, X., Liu, S., Han, Y., Wang, G., and Ho, K.: Aerosol particles at a high-altitude site on the Southeast Tibetan Plateau, China: Implications for pollution transport from South Asia: AEROSOL PARTICLES IN SOUTHEAST TP, *J. Geophys. Res. Atmos.*, 118, 11,360-11,375, <https://doi.org/10.1002/jgrd.50599>, 2013.

a mis en forme : Anglais (E.U.)



Figure 1. (A) Location map of AM5 site in the Middle-Atlas; (B) Photo of the AM5 site





Figure 2. Seasonal wind rose plots at AM5

1440




**Figure 3.** (A) Time series of daily  $\text{PM}_{10}$  mass; (B) Box plot of monthly averages of  $\text{PM}_{10}$  mass; (C) Pollution rose of  $\text{PM}_{10}$  mass. The presented data were separated according to each season into summer (Aug), autumn (Sep-Nov), and winter (Dec).


a mis en forme : Retrait : Gauche : 0,75 cm

a mis en forme : Couleur de police : Texte 1

a mis en forme : Couleur de police : Texte 1



**Figure 3.** (A) Time series of daily  $\text{PM}_{10}$  mass and Box plot of monthly averages of  $\text{PM}_{10}$  mass; (B) Pollution rose of  $\text{PM}_{10}$  mass



**Figure 45.** Typical 96h air mass back trajectory ensembles performed for AM5 during routine samples periods; aerosol type and PM<sub>10</sub> mass concentration are given in parentheses: (a) 18 December 2017: air mass from the North Atlantic Ocean considered to be representative of background conditions (BAM,  $m=10.9 \mu\text{g m}^{-3}$ ); (b) 10 October 2017: air mass from Europe crossing coastline of North Morocco (Atlantic Coast Europe,  $m=44.1 \mu\text{g m}^{-3}$ ); (c) 2 November 2017: slightly polluted air mass from North East crossing Mediterranean Sea Morocco (Mediterranean Coast Europe,  $m=26.4 \mu\text{g m}^{-3}$ ) ; (d) 13 August 2017: dust loaded air mass coming from Sahara desert (Saharan Dust,  $m=94.1 \mu\text{g m}^{-3}$ ).

a mis en forme : Retrait : Gauche : 0,75 cm

a mis en forme : Police : Italique


a mis en forme : Exposant

a mis en forme : Police : Non Gras

a mis en forme : Police : Non Gras

a mis en forme : Police : Non Gras


a mis en forme : Interligne : 1,5 ligne




**Figure 56.** Time series of major aerosol chemical constituents in PM<sub>10</sub> filter samples collected from August to December 2017; The color of the symbols displayed for each sample represents a specific air mass origin: Background (blue); ACE (red); MCE (green); SD (yellow).

a mis en forme : Retrait : Gauche : 1 cm

a mis en forme : Centré




**Figure 6.** Time series of organic compounds in  $\text{PM}_{10}$  filter samples collected from August to December 2017 at AM5; The color of the symbols displayed for each sample represents a specific air mass origin: Background (blue); ACE (red); MCE (green); SD (yellow).



**Figure 77.** Crustal enrichment factors (EF) of aerosol PM<sub>10</sub> analysis evaluated for the different trace metal elements at AM5; The averaged values are plotted according to their respective air mass origins.

a mis en forme : Centré



**Figure 89.** Correlations-Scatter plot of (A)  $\text{NH}_4^+$  with  $\text{NO}_3^-$  and  $\text{nss-SO}_4^{2-}$  during summer; (B)  $\text{NH}_4^+$  with  $\text{NO}_3^-$  and  $\text{nss-SO}_4^{2-}$  during autumn-winter; (C)  $\text{Na}^+$  and  $\text{Cl}^- + \text{SO}_4^{2-}$  during ACE air mass; (D)  $\text{Na}^+$  and  $\text{Cl}^- + \text{NO}_3^-$  during MCE air mass at the AM5 site.

between various compound and elements according to seasons

a mis en forme : Non Surlinage

a mis en forme : Non Surlinage

a mis en forme : Indice

a mis en forme : Retrait : Gauche : 1 cm

a mis en forme : Non Surlinage

a mis en forme : Non Surlinage

a mis en forme : Non Surlinage

a mis en forme : Normal, Retrait : Gauche : 0 cm, Interligne : simple

a mis en forme : Interligne : 1,5 ligne

a mis en forme : Normal, Retrait : Gauche : 0 cm

a mis en forme : Justifié, Retrait : Gauche : 0,75 cm, Interligne : 1,5 ligne

a mis en forme : Indice

a mis en forme : Exposant


a mis en forme : Indice

a mis en forme : Exposant

a mis en forme : Indice

a mis en forme : Exposant


a mis en forme : Exposant



**Figure 911.** Day and night-time variation of (A) OC, EC, ionic chemical species, (B) anthropogenic metals, and (C) Organic compounds such as alkanes, PAHs and alkane-2-ones ketones.

a mis en forme : Retrait : Gauche : 0,75 cm

a mis en forme : Normal, Retrait : Gauche : 0 cm



**Figure 10.** Boxplot in the period of dust and non-dust events for concentrations of (A) PM<sub>10</sub> mass and mineral dust; (B) Fe/Al and Fe/Ca ratio (C) OC and major water-soluble ions, (D) EC and minor water-soluble ions; and (E) organic compounds.

a mis en forme : Retrait : Gauche : 0,75 cm

**Table 1.** Meteorological parameters over Middle-Atlas from summer 2017 to spring 2018

Meteorological Parameter	Period			
	Summer 2017	Autumn 2017	Winter 2018	Spring 2018
Temperature (°C)				
Mean	24	17	5	10
Min	19	10	-1	3
Max	26	21	7	13
Wind speed ( $m\ s^{-1}\ km\ h$ )				
Mean	185	185	206	267
Min	61	82	72	123
Max	298	5920	5920	7418
Relative humidity (%)				
Mean	37	39	74	62
Max	80	92	97	94
Rainfall (mm)	37	63	141	56
Wind direction (degrees)	182	185	205	221
Visibility (km)	10	10	9	9
Pressure (mbar)	1 022	1 025	1 034	1 027

1510

**Table 2.** Comparison of different methods for dust estimation

Method	Mean	Max	Min	Std	Equation	Reference
Method 1	24.5	132.1	5.19	18.2	$MD_1 = PM10\ mass - (\sum All\ detected\ element)$	(Maenhaut et al., 2005))
Method 2	19.9	112.3	0.46	17.8	$MD_2 = 1.16(1.90 Al + 2.15 Si + 1.41 Ca + 2.09 Fe + 1.67 Ti)$	<a href="#">Maenhaut et al., 2005</a>
Method 3	18.9	104.9	0.20	18.9	$MD_3 = 4\left(\frac{Al}{27}\right)51 + 100\left(\frac{Ca}{40}\right) + 84\left(\frac{Mg}{24}\right) + 80\left(\frac{Ti}{48}\right) + 87\left(\frac{Mn}{55}\right) + 80\left(\frac{Fe}{56}\right)$	<a href="#">Minguillón et al., 2007; Nerriere et al., 2007</a>
Method 4	15.5	91.4	0.13	13.9	$MD_4 = 2.1 Al + 2.9 Si + 1.4 nss - Ca^{2+} + 1.4 nss - Mg^{2+} + 1.43 Fe + 1.55 K + 1.58 Mn$	<a href="#">Cesari et al., 2012; Marenco et al., 2006; Perrino et al., 2014</a>

1515  
520  
525  
1530

**Table 3.** Average mass concentration of PM<sub>10</sub> from other [high-altitude sites and urban sites in Morocco](#) sites reported in the literature according to altitude.

Nº	Site	Site type	Sampling Period	Altitude (m)	PM <sub>10</sub> ( $\mu\text{g m}^{-3}$ )	References
1	Mt. Everest, Nepal	High altitude	Feb 2006-May 2008	5079	6	Decesari et al., 2010
2	Lhasa, Tibet	High altitude	Jan-Feb 2006	3663	37.7	Wang et al., 2015
3	Jungfraujoch, Switzerland	High altitude	Feb-Mar 2005	3580	240	Cozic et al., 2008
4	Izaña, Canary Islands	High altitude	Feb 2008-Aug 2013	2400	46	Garcia et al., 2017
5	Mount Cimone, Italy	High altitude	Jun-Aug 2004	2165	220	Marenco et al., 2006
6	Atlas (AM5), Morocco	High altitude	Aug-Dec 2017	2100	29	Present study
7	Puy de Dome, France	High altitude	Apr 2006-Apr 2007	1465	6	Bourcier et al. 2012
8	Sinhagad, India	High altitude	Nov 2008-Apr 2009	1455	35.8	Satsangi et al., 2014
9	Mahabaleshw ar, India	High altitude	Jun 2012-May 2013	1348	140	Leena et al., 2017
10	Dharjeeling, India	High altitude	Feb 2013-Jan 2014	9	Putero et al., 2017	
11	Marrakech, Morocco	Urban	2009-2012	465	1348	Inchaouh et al., 2017
12	Morocco Maha	Urban	May 2013	55	37.7	Leena et al., 2017
13	Meknes, Morocco	Urban	Mar 2007-Apr 2008	546	47	Tahri et al., 2017
14	Tetouan, Morocco	Urban	May 2011-Apr 2012	105	216	Benchrif et al., 2018
15	Morocco Debb	Urban	2012-Dec 2014-Jun 2015	105	31.6	Benchrif et al., 2018
16	Gezo, Malta	Marine	Mar 2012-May 2013	114	18.2	Secri et al., 2016
17	Kenitra, Morocco	Urban	Feb 2007-Feb 2008	26	110	Tahri et al., 2013
18	Tunis-Tunisia	Urban	Jun-Jul 2008	40	58.9	
19	CVAO—Cap Verde	Cap	Jan 2007-Dec 2011	10	47.2	F

- a mis en forme ... [27]
- a mis en forme le tableau ... [28]
- a mis en forme ... [29]
- a mis en forme ... [30]
- a mis en forme ... [31]
- a mis en forme ... [33]
- a mis en forme ... [32]
- a mis en forme ... [34]
- a mis en forme ... [35]
- a mis en forme ... [36]
- a mis en forme ... [38]
- a mis en forme ... [39]
- a mis en forme ... [40]
- a mis en forme ... [41]
- a mis en forme ... [37]
- a mis en forme ... [42]
- a mis en forme ... [43]
- a mis en forme ... [45]
- a mis en forme ... [44]
- a mis en forme ... [46]
- a mis en forme ... [47]
- a mis en forme ... [48]
- a mis en forme ... [49]
- a mis en forme ... [50]
- a mis en forme ... [51]
- a mis en forme ... [52]
- a mis en forme ... [53]
- a mis en forme ... [54]
- a mis en forme ... [55]
- a mis en forme ... [56]
- a mis en forme ... [57]
- a mis en forme ... [58]
- a mis en forme ... [59]
- a mis en forme ... [60]
- a mis en forme ... [61]
- a mis en forme ... [62]
- a mis en forme ... [63]
- a mis en forme ... [64]
- a mis en forme ... [65]
- a mis en forme ... [66]
- a mis en forme ... [67]
- a mis en forme ... [68]
- a mis en forme ... [69]
- a mis en forme ... [70]
- a mis en forme ... [71]
- a mis en forme ... [72]
- a mis en forme ... [73]
- a mis en forme ... [74]
- a mis en forme ... [75]
- a mis en forme ... [76]
- a mis en forme ... [77]
- a mis en forme ... [78]
- a mis en forme ... [79]
- a mis en forme ... [80]
- a mis en forme ... [81]
- a mis en forme ... [82]
- a mis en forme ... [83]
- a mis en forme ... [84]
- a mis en forme ... [86]
- a mis en forme ... [87]
- a mis en forme ... [85]
- a mis en forme ... [88]
- a mis en forme ... [91]
- a mis en forme ... [89]

**Table 4** Concentrations of main aerosol chemical species in PM<sub>10</sub> according to each air mass ( $\mu\text{g m}^{-3}$ ) at AMS;  
The organic composition is given in  $\text{ng m}^{-3}$ ; The number of samples is written in parentheses.

Aerosol components	Air mass			
	BAM (n=10)	ACE (n=51)	MCE (n=71)	SD (n=43)
Mass load	<u>10.9 ± 0.9</u>	<u>20.4 ± 6.3</u>	<u>33.8 ± 14.5</u>	<u>37.9 ± 25.3</u>
Dust	<u>5.5 ± 3.5</u>	<u>13.3 ± 5.2</u>	<u>19.9 ± 11.9</u>	<u>29.1 ± 22.6</u>
Sea salt	<u>0.05 ± 0.06</u>	<u>0.3 ± 0.5</u>	<u>0.3 ± 0.4</u>	<u>0.2 ± 0.2</u>
OM	<u>1.0 ± 0.6</u>	<u>1.4 ± 1.1</u>	<u>2.7 ± 1.7</u>	<u>2.3 ± 1.8</u>
EC	<u>0.2 ± 0.1</u>	<u>0.2 ± 0.1</u>	<u>0.2 ± 0.1</u>	<u>0.2 ± 0.1</u>
POC	<u>0.1 ± 0.06</u>	<u>0.2 ± 0.3</u>	<u>0.3 ± 0.3</u>	<u>0.2 ± 0.2</u>
SOC	<u>0.3 ± 0.2</u>	<u>0.4 ± 0.4</u>	<u>1.0 ± 0.8</u>	<u>0.9 ± 0.8</u>
NO <sub>3</sub> <sup>-</sup>	<u>0.5 ± 0.6</u>	<u>0.6 ± 0.7</u>	<u>1.0 ± 0.7</u>	<u>0.9 ± 0.4</u>
nss-SO <sub>4</sub> <sup>2-</sup>	<u>0.2 ± 0.2</u>	<u>0.5 ± 0.5</u>	<u>1.2 ± 0.9</u>	<u>1.0 ± 0.6</u>
NH <sub>4</sub> <sup>+</sup>	<u>0.2 ± 0.2</u>	<u>0.2 ± 0.2</u>	<u>0.3 ± 0.2</u>	<u>0.2 ± 0.1</u>
Ca <sub>2</sub> <sup>+</sup>	<u>0.2 ± 0.2</u>	<u>0.2 ± 0.2</u>	<u>0.8 ± 0.5</u>	<u>0.9 ± 0.6</u>
Alkanes	<u>4.9 ± 3.2</u>	<u>5.6 ± 3.7</u>	<u>10.5 ± 7.7</u>	<u>9.2 ± 8.6</u>
PAHs	<u>0.4 ± 0.5</u>	<u>0.4 ± 0.4</u>	<u>0.9 ± 2.1</u>	<u>0.7 ± 0.7</u>
Alkan-2-ones	<u>7.8 ± 6.9</u>	<u>5.9 ± 5.5</u>	<u>5.5 ± 5.0</u>	<u>6.6 ± 4.1</u>
Sugars	-	<u>3.7 ± 6.0</u>	<u>5.2 ± 7.7</u>	<u>1.8 ± 2.9</u>
Oxalate	<u>44 ± 26</u>	<u>73 ± 58</u>	<u>129 ± 58</u>	<u>107 ± 63</u>
pH	<u>5.6 ± 0.2</u>	<u>6.0 ± 0.4</u>	<u>6.5 ± 0.4</u>	<u>6.5 ± 0.4</u>
OC/EC	<u>2.2 ± 1.1</u>	<u>3.3 ± 1.9</u>	<u>6.3 ± 7.5</u>	<u>4.6 ± 4.6</u>
CPI	<u>3.3 ± 0.8</u>	<u>3.5 ± 2.4</u>	<u>3.9 ± 1.9</u>	<u>4.0 ± 3.1</u>

**Table 5.** Concentrations of main aerosol chemical species in PM<sub>10</sub> (ng m<sup>-3</sup>) at AM5 compared to other high altitude mountain stations. Data are reported in the format average (mean ± standard deviation) and NA: not available. Notice that the concentrations of PM<sub>10</sub> mass are given in µg m<sup>-3</sup>. <sup>a</sup> Present study; <sup>b</sup> Bourcier et al. 2012; <sup>c</sup> Chatterjee et al. 2010; <sup>d</sup> Marenco et al. 2006; <sup>e</sup> Decesari et al., 2010.

Elements	Mt. Atlas Morocco <sup>a</sup>	Mt. Puy de Dome, France <sup>b</sup>	Mt. Himalaya, India <sup>c</sup>	Mt. Cimone, Italy <sup>d</sup>	Mt. Everest, Nepal <sup>e</sup>
Altitude (m a.s.l)	2100 m	1465 m	2194 m	2165 m	5079 m
Samples	190	NA	111	57	99
Period	Aug-Dec 2017	Apr 2006-Apr 2007	Jan-Dec 2005	Jun-Aug 2004	Apr 2006 - May 2008
Mass load	29.1 ± 17.3	5.6 ± 4.6	29.5 ± 20.8	16.1 ± 9.8	5.6 ± 4.6
OC	1069 ± 818	NA	NA	NA	800 ± 637
EC	247 ± 134	NA	NA	NA	115 ± 132
Na <sup>+</sup>	186 ± 231	NA	2200 ± 2000	NA	24.2 ± 22.5
K <sup>+</sup>	42 ± 35	NA	310 ± 210	NA	34 ± 32
Ca <sup>2+</sup>	649 ± 579	15.5 ± 10.2	130 ± 10	360 ± 550	138 ± 90
Mg <sup>2+</sup>	60 ± 50	NA	120 ± 60	NA	19.3 ± 7.2
Cl <sup>-</sup>	80 ± 133	NA	2350 ± 1500	82 ± 98	22 ± 46
NH <sub>4</sub> <sup>+</sup>	298 ± 220	297 ± 276	50 ± 40	1400 ± 800	175 ± 183
NO <sub>3</sub> <sup>-</sup>	859 ± 687	510 ± 980	950 ± 200	840 ± 770	170 ± 223
SO <sub>4</sub> <sup>2-</sup>	941 ± 848	1380 ± 1160	3500 ± 2100	3500 ± 2000	394 ± 329
Al	443 ± 830	NA	NA	300 ± 460	740
Fe	486 ± 728	NA	NA	260 ± 440	NA
Ti	37 ± 45	NA	NA	30 ± 50	NA
V	3.5 ± 12.2	NA	NA	3.1 ± 1.5	NA
K	174 ± 156	NA	NA	160 ± 210	NA
Cr	4.3 ± 5.2	NA	NA	NA	NA
Ni	2.4 ± 3.1	NA	NA	1.4 ± 0.5	NA
Cu	1.2 ± 3.1	NA	NA	2.9 ± 3.1	NA
Zn	8.6 ± 6.2	NA	NA	9.9 ± 6.6	NA
Pb	4.8 ± 4.5	NA	NA	3.9 ± 2.4	NA
Mn	12.4 ± 39.3	NA	NA	6.2 ± 7.0	NA

a mis en forme

... [110]

a mis en forme : Police :12 pt, Couleur de police : Automatique

... [111]

a mis en forme

... [112]

a mis en forme

... [113]

a mis en forme

... [114]

a mis en forme

... [115]

a mis en forme

... [116]

a mis en forme

... [117]

a mis en forme

... [118]

a mis en forme : Couleur de police : Texte 1

... [119]

a mis en forme

... [120]

a mis en forme

... [121]

a mis en forme

... [122]

a mis en forme

... [123]

a mis en forme : Couleur de police : Texte 1

... [124]

a mis en forme : Couleur de police : Texte 1

... [125]

a mis en forme : Couleur de police : Texte 1

... [126]

a mis en forme : Couleur de police : Texte 1

... [127]

a mis en forme : Couleur de police : Texte 1

... [128]

a mis en forme : Couleur de police : Texte 1

... [129]

a mis en forme : Couleur de police : Texte 1

... [130]

a mis en forme : Couleur de police : Texte 1

... [131]

a mis en forme : Couleur de police : Texte 1

... [132]

a mis en forme : Couleur de police : Texte 1

... [133]

a mis en forme : Couleur de police : Texte 1

... [134]

a mis en forme : Couleur de police : Texte 1

... [135]

a mis en forme : Couleur de police : Texte 1

... [136]

a mis en forme : Couleur de police : Texte 1

... [137]

a mis en forme : Couleur de police : Texte 1

... [138]

a mis en forme : Couleur de police : Texte 1

... [139]

a mis en forme : Couleur de police : Texte 1

... [140]

a mis en forme : Couleur de police : Texte 1

... [141]

a mis en forme : Couleur de police : Texte 1

... [142]

a mis en forme : Couleur de police : Texte 1

... [143]

a mis en forme : Couleur de police : Texte 1

... [144]

a mis en forme : Couleur de police : Texte 1

... [145]

a mis en forme : Couleur de police : Texte 1

... [146]

a mis en forme : Couleur de police : Texte 1

... [147]

a mis en forme : Couleur de police : Texte 1

... [148]

a mis en forme : Couleur de police : Texte 1

... [149]

a mis en forme : Couleur de police : Texte 1

... [150]

a mis en forme : Couleur de police : Texte 1

... [151]

a mis en forme : Couleur de police : Texte 1

... [152]

a mis en forme : Couleur de police : Texte 1

... [153]

a mis en forme : Couleur de police : Texte 1

... [154]

a mis en forme : Couleur de police : Texte 1

... [155]

a mis en forme : Couleur de police : Texte 1

... [156]

a mis en forme : Couleur de police : Texte 1

... [157]

a mis en forme : Couleur de police : Texte 1

... [158]

a mis en forme : Couleur de police : Texte 1

... [159]

a mis en forme : Couleur de police : Texte 1

... [160]

a mis en forme : Couleur de police : Texte 1

... [161]

a mis en forme : Couleur de police : Texte 1

... [162]

a mis en forme : Couleur de police : Texte 1

... [163]

a mis en forme : Couleur de police : Texte 1

... [164]

a mis en forme : Couleur de police : Texte 1

... [165]

a mis en forme : Couleur de police : Texte 1

... [166]

a mis en forme : Couleur de police : Texte 1

... [167]

a mis en forme : Couleur de police : Texte 1

... [168]

a mis en forme : Couleur de police : Texte 1

... [169]

a mis en forme : Couleur de police : Texte 1

... [170]

a mis en forme : Couleur de police : Texte 1

... [171]

a mis en forme : Couleur de police : Texte 1

... [172]

a mis en forme : Couleur de police : Texte 1

... [173]

a mis en forme : Couleur de police : Texte 1

... [174]

a mis en forme : Couleur de police : Texte 1

... [175]

a mis en forme : Couleur de police : Texte 1

... [176]

a mis en forme : Couleur de police : Texte 1

... [177]

a mis en forme : Couleur de police : Texte 1

... [178]

a mis en forme : Couleur de police : Texte 1

... [179]

a mis en forme : Couleur de police : Texte 1

... [180]

a mis en forme : Couleur de police : Texte 1

... [181]

a mis en forme : Couleur de police : Texte 1

... [182]

a mis en forme : Couleur de police : Texte 1

... [183]

a mis en forme : Couleur de police : Texte 1

... [184]

a mis en forme : Couleur de police : Texte 1

... [185]

a mis en forme : Couleur de police : Texte 1

... [186]

a mis en forme : Couleur de police : Texte 1

... [187]

a mis en forme : Couleur de police : Texte 1

... [188]

a mis en forme : Couleur de police : Texte 1

... [189]

a mis en forme : Couleur de police : Texte 1

... [190]

a mis en forme : Couleur de police : Texte 1

... [191]

a mis en forme : Couleur de police : Texte 1

... [192]

a mis en forme : Couleur de police : Texte 1

... [193]

a mis en forme : Couleur de police : Texte 1

... [194]

a mis en forme : Couleur de police : Texte 1

... [195]

a mis en forme : Couleur de police : Texte 1

... [196]

a mis en forme : Couleur de police : Texte 1

... [197]

a mis en forme : Couleur de police : Texte 1

... [198]

a mis en forme : Couleur de police : Texte 1

... [199]

a mis en forme : Couleur de police : Texte 1

... [200]

a mis en forme : Couleur de police : Texte 1

... [201]

a mis en forme : Couleur de police : Texte 1

... [202]

a mis en forme : Couleur de police : Texte 1

... [203]

a mis en forme : Couleur de police : Texte 1

... [204]

a mis en forme : Couleur de police : Texte 1

... [205]

a mis en forme : Couleur de police : Texte 1

... [206]

a mis en forme : Couleur de police : Texte 1

... [207]

a mis en forme : Couleur de police : Texte 1

... [208]

a mis en forme : Couleur de police : Texte 1

... [209]

a mis en forme : Couleur de police : Texte 1

... [210]

a mis en forme : Couleur de police : Texte 1

... [211]

a mis en forme : Couleur de police : Texte 1

... [212]

a mis en forme : Couleur de police : Texte 1

... [213]

a mis en forme : Couleur de police : Texte 1

... [214]

a mis en forme : Couleur de police : Texte 1

... [215]

a mis en forme : Couleur de police : Texte 1

... [216]

a mis en forme : Couleur de police : Texte 1

... [217]

a mis en forme : Couleur de police : Texte 1

... [218]

a mis en forme : Couleur de police : Texte 1

... [219]

a mis en forme : Couleur de police : Texte 1

... [220]

a mis en forme : Couleur de police : Texte 1

... [221]

a mis en forme : Couleur de police : Texte 1









<b>Page 16 : [5] a mis en forme</b>	Nabil Deabji	18/08/2021 13:06:00
Couleur de police : Texte 1, Anglais (E.U.)		
<b>Page 16 : [5] a mis en forme</b>	Nabil Deabji	18/08/2021 13:06:00
Couleur de police : Texte 1, Anglais (E.U.)		
<b>Page 16 : [5] a mis en forme</b>	Nabil Deabji	18/08/2021 13:06:00
Couleur de police : Texte 1, Anglais (E.U.)		
<b>Page 16 : [6] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [6] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [6] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [7] a mis en forme</b>	Nabil Deabji	18/08/2021 14:16:00
Couleur de police : Texte 1		
<b>Page 16 : [8] a mis en forme</b>	Nabil Deabji	18/08/2021 22:38:00
Couleur de police : Texte 1		
<b>Page 16 : [8] a mis en forme</b>	Nabil Deabji	18/08/2021 22:38:00
Couleur de police : Texte 1		
<b>Page 16 : [8] a mis en forme</b>	Nabil Deabji	18/08/2021 22:38:00
Couleur de police : Texte 1		
<b>Page 16 : [8] a mis en forme</b>	Nabil Deabji	18/08/2021 22:38:00
Couleur de police : Texte 1		
<b>Page 16 : [9] a mis en forme</b>	Nabil Deabji	04/09/2021 14:17:00
Non Surlignage		
<b>Page 16 : [9] a mis en forme</b>	Nabil Deabji	04/09/2021 14:17:00
Non Surlignage		
<b>Page 16 : [9] a mis en forme</b>	Nabil Deabji	04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 16 : [9] a mis en forme** Nabil Deabji 04/09/2021 14:17:00

Non Surlignage

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1

**Page 17 : [10] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 17 : [11] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 17 : [11] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 17 : [11] a mis en forme** Nabil Deabji 18/08/2021 22:38:00

Couleur de police : Texte 1, Anglais (E.U.)











Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [21] a mis en forme** Nabil Deabji 23/08/2021 16:19:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [21] a mis en forme** Nabil Deabji 23/08/2021 16:19:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [21] a mis en forme** Nabil Deabji 23/08/2021 16:19:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [22] a mis en forme** Nabil Deabji 26/08/2021 19:59:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [22] a mis en forme** Nabil Deabji 26/08/2021 19:59:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [23] a mis en forme** Khamneh Wadinga Fomba 31/08/2021 09:13:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [23] a mis en forme** Khamneh Wadinga Fomba 31/08/2021 09:13:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [23] a mis en forme** Khamneh Wadinga Fomba 31/08/2021 09:13:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [24] a mis en forme** Nabil Deabji 26/08/2021 20:18:00

Couleur de police : Texte 1, Anglais (E.U.), Indice

**Page 20 : [24] a mis en forme** Nabil Deabji 26/08/2021 20:18:00

Couleur de police : Texte 1, Anglais (E.U.), Indice

**Page 20 : [24] a mis en forme** Nabil Deabji 26/08/2021 20:18:00

Couleur de police : Texte 1, Anglais (E.U.), Indice

**Page 20 : [24] a mis en forme** Nabil Deabji 26/08/2021 20:18:00

Couleur de police : Texte 1, Anglais (E.U.), Indice

**Page 20 : [25] a mis en forme** Khamneh Wadinga Fomba 31/08/2021 09:13:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [25] a mis en forme** Khamneh Wadinga Fomba 31/08/2021 09:13:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [26] a mis en forme** Nabil Deabji 26/08/2021 19:57:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [26] a mis en forme** Nabil Deabji 26/08/2021 19:57:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [26] a mis en forme** Nabil Deabji 26/08/2021 19:57:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [26] a mis en forme** Nabil Deabji 26/08/2021 19:57:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [26] a mis en forme** Nabil Deabji 26/08/2021 19:57:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 20 : [26] a mis en forme** Nabil Deabji 26/08/2021 19:57:00

Couleur de police : Texte 1, Anglais (E.U.)

**Page 44 : [27] a mis en forme** Nabil Deabji 26/07/2021 18:13:00

Anglais (E.U.)

**Page 44 : [28] a mis en forme le tableau** Nabil Deabji 27/08/2021 12:58:00

a mis en forme le tableau

**Page 44 : [29] a mis en forme** Nabil Deabji 08/08/2021 19:49:00

Anglais (E.U.), Exposant

Page 44 : [30] a mis en forme	Nabil Deabji	27/08/2021 12:55:00
Centré		
Page 44 : [31] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [32] a mis en forme	Nabil Deabji	20/07/2021 14:43:00
Centré		
Page 44 : [33] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [33] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [33] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [34] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [35] a mis en forme	Nabil Deabji	26/08/2021 14:10:00
Gauche		
Page 44 : [36] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Non Surlignage		
Page 44 : [37] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [38] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
Page 44 : [38] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
Page 44 : [39] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [40] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Non Surlignage		
Page 44 : [40] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Non Surlignage		
Page 44 : [41] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [42] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [43] a mis en forme	Nabil Deabji	26/08/2021 14:10:00
Gauche		
Page 44 : [44] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [45] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [45] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
Page 44 : [46] a mis en forme	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		

<b>Page 44 : [47] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [48] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [48] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [49] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [50] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [51] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [52] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [53] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [53] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [54] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [55] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Non Surlignage		
<b>Page 44 : [56] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Anglais (E.U.)		
<b>Page 44 : [57] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Anglais (E.U.)		
<b>Page 44 : [58] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Non Surlignage		
<b>Page 44 : [59] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [60] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [61] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [62] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Police :Non Gras		
<b>Page 44 : [63] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [64] a mis en forme</b>	Nabil Deabji	03/09/2021 11:10:00
Police :Gras		
<b>Page 44 : [65] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00
Anglais (E.U.)		
<b>Page 44 : [66] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [67] a mis en forme</b>	Nabil Deabji	08/08/2021 19:54:00

Anglais (E.U.)

**Page 44 : [68] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.)

**Page 44 : [69] a mis en forme** Nabil Deabji 26/08/2021 14:10:00

Gauche

**Page 44 : [70] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.)

**Page 44 : [71] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.)

**Page 44 : [72] a mis en forme** Nabil Deabji 26/08/2021 14:10:00

Gauche

**Page 44 : [73] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.)

**Page 44 : [73] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.)

**Page 44 : [74] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Non Surlignage

**Page 44 : [75] a mis en forme** Nabil Deabji 26/08/2021 14:10:00

Gauche

**Page 44 : [76] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Non Surlignage

**Page 44 : [77] a mis en forme** Nabil Deabji 26/08/2021 14:10:00

Gauche

**Page 44 : [78] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.), Non Surlignage

**Page 44 : [79] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.), Non Surlignage

**Page 44 : [80] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.), Non Surlignage

**Page 44 : [81] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.), Non Surlignage

**Page 44 : [81] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.), Non Surlignage

**Page 44 : [81] a mis en forme** Nabil Deabji 08/08/2021 19:54:00

Anglais (E.U.), Non Surlignage

**Page 44 : [82] a mis en forme** Nabil Deabji 26/08/2021 14:10:00

Gauche

**Page 44 : [83] a mis en forme** Nabil Deabji 26/07/2021 18:13:00

Anglais (E.U.)

**Page 44 : [84] a mis en forme** Nabil Deabji 26/07/2021 18:13:00

Anglais (E.U.), Non Surlignage

**Page 44 : [85] a mis en forme** Nabil Deabji 26/07/2021 18:13:00

Anglais (E.U.), Non Surlignage

**Page 44 : [85] a mis en forme** Nabil Deabji 26/07/2021 18:13:00

Anglais (E.U.), Non Surlignage

<b>Page 44 : [86] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [87] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [87] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [88] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [89] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [90] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [91] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [91] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [92] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [93] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [94] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [95] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [96] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [97] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [98] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [99] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [100] a mis en forme</b>	Nabil Deabji	26/08/2021 14:10:00
Gauche		
<b>Page 44 : [101] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [102] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [103] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [103] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		
<b>Page 44 : [103] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Non Surlignage		

<b>Page 44 : [104] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [105] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Surlignage		
<b>Page 44 : [106] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [107] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.), Surlignage		
<b>Page 44 : [108] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 44 : [109] a mis en forme</b>	Nabil Deabji	26/07/2021 18:13:00
Anglais (E.U.)		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [110] a mis en forme</b>	Nabil Deabji	08/08/2021 18:00:00
Indice		
<b>Page 46 : [111] a mis en forme</b>	Nabil Deabji	03/09/2021 11:09:00
Police :Gras, Couleur de police : Texte 1		
<b>Page 46 : [111] a mis en forme</b>	Nabil Deabji	03/09/2021 11:09:00
Police :Gras, Couleur de police : Texte 1		
<b>Page 46 : [111] a mis en forme</b>	Nabil Deabji	03/09/2021 11:09:00
Police :Gras, Couleur de police : Texte 1		
<b>Page 46 : [112] a mis en forme</b>	Khanneh Wadinga Fomba	31/08/2021 09:13:00
Couleur de police : Texte 1, Portugais		
<b>Page 46 : [112] a mis en forme</b>	Khanneh Wadinga Fomba	31/08/2021 09:13:00
Couleur de police : Texte 1, Portugais		
<b>Page 46 : [112] a mis en forme</b>	Khanneh Wadinga Fomba	31/08/2021 09:13:00
Couleur de police : Texte 1, Portugais		
<b>Page 46 : [113] a mis en forme</b>	Nabil Deabji	08/08/2021 18:18:00
Couleur de police : Texte 1		
<b>Page 46 : [113] a mis en forme</b>	Nabil Deabji	08/08/2021 18:18:00



**Page 46 : [121] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [121] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [122] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [122] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [122] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [123] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [123] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [123] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1

**Page 46 : [123] a mis en forme** Nabil Deabji 26/08/2021 12:23:00  
Couleur de police : Texte 1