

# Investigation of New Particle Formation mechanisms and aerosol processes

## at the Marambio Station, Antarctic Peninsula.

Lauriane Lucie Josette Quéléver<sup>1</sup>, Lubna Dada<sup>1,2,3</sup>, Eija Asmi<sup>4,5</sup>, Janne Lampilahti<sup>1</sup>, Tommy Chan<sup>1</sup>, Jonathan Ferrara<sup>4</sup>, Gustavo Copes<sup>4</sup>, German Pérez-Fogwill<sup>4</sup>, Luis Barreira<sup>5</sup>, Minna Aurela<sup>5</sup>, Douglas Worsnop<sup>1,6</sup>, Tuija Jokinen<sup>1,7</sup>, and Mikko Sipilä<sup>1</sup>

<sup>1</sup> Institute for Atmospheric and Earth System Research / INAR-Physics, P.O. Box 64, FI-00014 University of Helsinki, Finland.

<sup>2</sup> Extreme Environments Research Laboratory, École Polytechnique Fédérale de Lausanne (EPFL) Valais, Sion, 1951, Switzerland

<sup>3</sup> Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen, Switzerland

<sup>4</sup> Servicio Meteorológico Nacional / SMN, Av. Dorrego, 4019, Buenos Aires, Argentina

<sup>5</sup> Finnish Meteorological Institute / FMI, Erik Palmenin aukio 1, FI-00560 Helsinki, Finland

<sup>6</sup> Aerodyne Research, Inc. Billerica, MA 01821, USA

<sup>7</sup> The Cyprus Institute, The Climate & Atmosphere Research Centre (CARE-C), P.O. Box 27456 Nicosia, Cyprus

*Corresponding Author:*

Lauriane L. J. Quéléver ([Lauriane.quelever@helsinki.fi](mailto:Lauriane.quelever@helsinki.fi)), Mikko Sipilä ([Mikko.sipila@helsinki.fi](mailto:Mikko.sipila@helsinki.fi))

### Abstract

Understanding chemical processes leading to the formation of atmospheric aerosol particles is crucial to improve our capabilities in predicting the future climate. However, those mechanisms are still inadequately characterized, especially in polar regions. In this study, we report observations of neutral and charged aerosol precursor molecules and chemical clusters composition (qualitatively and quantitatively), as well as air ions and aerosol particle number concentrations and size distributions from the Marambio research station (64° 15' S - 56° 38' W), located North of the Antarctic Peninsula. We conducted measurements during the austral summer, between 15 January and 25 February 2018. The scope of this study is to characterize new particle formation (NPF) event parameters and connect our observations of gas phase compounds with the formation of secondary aerosols to resolve the nucleation mechanisms at the molecular scale. NPF occurred on 40 % of

35 measurement days. All NPF events were observed during days with high solar radiation, mostly with  
36 above freezing temperatures and, with low relative humidity. The averaged formation rate for 3 nm  
37 particles ( $J_3$ ) was  $0.686 \text{ cm}^{-3} \text{ s}^{-1}$  and the average particle growth rate ( $GR_{3.8-12 \text{ nm}}$ ) was  $4.2 \text{ nm h}^{-1}$ .  
38 Analysis of neutral aerosol precursor molecules showed measurable concentrations of iodic acid (IA),  
39 sulfuric acid (SA) and methane sulfonic acid (MSA) throughout the entire measurement period with  
40 significant increase of MSA and SA concentrations during NPF events. We highlight SA as a key  
41 contributor to NPF processes, while IA and MSA would likely only contribute to particle growth.  
42 Mechanistically, anion clusters containing ammonia/ dimethylamine (DMA) and SA were identified,  
43 suggesting significant concentration of ammonia and DMA as well. Those species are likely  
44 contributing to NPF events since SA alone is not sufficient to explain observed nucleation rates. Here,  
45 we provide evidence of the marine origin of the measured chemical precursors and discuss their  
46 potential contribution to the aerosol phase.

47

## 48 **1 Introduction**

49

50 Atmospheric aerosol particles impact the planetary energy budget and radiation balance by  
51 influencing cloud optical properties and cloud lifetime (IPCC, 2013). Even though they are crucial  
52 for regulating the climate, aerosol particles remain insufficiently characterized, especially in remote  
53 regions (Schmale et al., 2019). Understanding the atmospheric processes in places where  
54 anthropogenic influence is minimal, such as polar regions, is important for characterizing the pre-  
55 industrial-like atmosphere. At the same time, polar environments are significantly impacted by  
56 human-induced climate change and are warming twice as fast as the global average (Stuecker et al.,  
57 2018). Additionally, polar ecosystems and landscapes are more and more being disturbed by  
58 increasing average temperatures that further affect emissions of trace gases into the atmosphere.

59

60 Aerosol particles have been observed and characterized in many places in the Arctic and Antarctica  
61 (Shaw, 1979; Shaw, 1988; Asmi et al., 2010; Kerminen et al., 2018; Sipilä et al., 2016; Dall'Osto et  
62 al., 2017; Jokinen et al., 2018; Dall'Osto et al., 2018; Herenz et al., 2019; Baccharini et al., 2020;  
63 Dall'Osto et al., 2019; Beck et al., 2021; Brean et al., 2021). Antarctic primary particles, mainly  
64 originating from sea spray or blowing snow, only weakly contribute to total particle number  
65 concentration (Lachlan-Cope et al., 2020). Modelling studies have estimated that primary particles  
66 would only contribute to  $\sim 2 \%$  of the total particle count that the ground level in Antarctica  
67 (Merikanto et al., 2009). Secondary formation of aerosol particles, on the other hand, is believed to  
68 be the principal contributor to cloud condensation nuclei (CCN), especially in the Antarctic peninsula

69 were models showed contributions varying from 75 % up to 100 % (Jokinen et al., 2018; Kerminen  
70 et al., 2018; Merikanto et al., 2009). These secondary aerosols originate from nucleation of gas phase  
71 molecules, typically condensing oxidation products of locally or regionally emitted vapors. Once  
72 formed, neutral or charged molecular clusters can grow by condensation of gases to sizes where they  
73 can act as CCN. The process of aerosol nucleation followed by subsequent growth is called new  
74 particle formation (NPF). Only a few studies have observed nucleation mode particles (sub-10 nm)  
75 from Antarctica - including oceanic, coastal, and continental areas (Asmi et al., 2010; Virkkula et  
76 al., 2009; Järvinen et al., 2013; Kyrö et al., 2013; Weller et al., 2015; Jokinen et al., 2018; Kerminen  
77 et al., 2018; Lachlan-Cope et al., 2020; Brean et al., 2021) - and only two of those report molecular  
78 clusters forming from precursor gases in coastal sites (Jokinen et al., 2018; Brean et al., 2021).

79  
80 Previously identified aerosol precursor vapors include sulfuric acid ( $\text{H}_2\text{SO}_4$ , SA, e.g., Kulmala et al.,  
81 2013; Sipilä et al., 2010), iodic acid ( $\text{HIO}_3$ , IA, Sipilä et al., 2016; Baccarini et al., 2020) and iodous  
82 acid ( $\text{HIO}_2$ , He et al., 2021), or highly oxygenated organic molecules (HOM, e.g., Ehn et al., 2014;  
83 Bianchi et al., 2019; Kirkby et al., 2016). On the other hand, nucleating ion clusters such as bisulfate  
84 ions with SA and ammonia (e.g., Kirkby et al., 2011; Yan et al., 2018; Jokinen et al., 2018; Beck et  
85 al., 2021), bisulfate ions and neutral SA with dimethyl amine ( $(\text{CH}_3)_2\text{NH}$ , DMA, Kürten et al., 2014),  
86 were previously identified to participate in aerosol formation processes.

87  
88 At Marambio, Aboa, and Princess Elisabeth coastal Antarctic stations, air mass trajectories during  
89 NPF events indicated a dominant role of the Southern Ocean as a source of aerosol particles, which  
90 chemical aerosol composition analysis confirmed by the abundance of e.g., marine-originated sulfates  
91 (Asmi et al., 2010; Herenz et al., 2019; Jokinen et al., 2018). These studies suggest that NPF is linked  
92 to sulfur-containing compounds originating from dimethyl sulfide ( $(\text{CH}_3)_2\text{S}$ , DMS). At the Antarctic  
93 coast, oceanic DMS concentrations are the highest during December to January (i.e., austral summer)  
94 with concentration that could exceed 15 nM within the upper 10 m layer of the ocean around the  
95 Peninsula compared to a yearly average of  $\sim 5$  nM (Lana et al., 2011). DMS has two well-known  
96 oxidation products formed from gas-phase reaction with OH radicals: SA (formed via sulfur dioxide,  
97  $\text{SO}_2$ ) and methane sulfonic acid ( $\text{CH}_3\text{SO}_3\text{H}$ , MSA), which can then initiate particle formation and  
98 subsequent particle growth (Barnes et al., 2006; Mardyukov and Schreiner, 2018).

99  
100 IA was also found to significantly contribute to NPF in marine and polar environments (Sipilä et al.,  
101 2016; Baccarini et al., 2020; He et al., 2021). Although the chemical production of IA is not fully  
102 resolved, IA results from the oxidation of reactive iodine (in the form of  $\text{I}_2$ , HIO, or intermediate I,

103 He et al., 2021) sourcing from algae/phytoplankton emissions contained within the sea  
104 water/ice/snow and exchanged into the atmosphere (Saiz-Lopez and Von Glasow, 2012). At the  
105 Antarctic Peninsula, the Weddell Sea – that undergoes consistent and recurrent phytoplankton bloom  
106 episodes every early spring - is a potential reservoir for iodine acid emissions, especially due to slower  
107 ice retreat during the austral summer and colder sea surface temperature than the Southern Ocean or  
108 than the Bellingshausen Sea, on the north and west-side of the peninsula, respectively (Atkinson et  
109 al., 2012; Von Berg et al., 2020).

110

111 This work aims at characterizing the gas-phase aerosol precursors naturally emitted from the marine/ice  
112 ecosystem and from the Antarctic continent with comprehensive gas phase and aerosol measurements  
113 performed at the Marambio research station, Antarctica. We present our results characterizing  
114 atmospheric NPF events observed in the Antarctic Peninsula (in section 3.2). There, we provide an  
115 extensive analysis of number size distributions of atmospheric aerosols and naturally charged ions,  
116 from ~1 nm to 800 nm. We also report the formation rates ( $J$ ) and growth rates ( $GR$ ) of the observed  
117 particle formation events. Our core analysis (in section 3.3) describes gas-phase composition and  
118 NPF precursors by mass spectrometry measurements of the concentration of SA, MSA and IA (cf.  
119 section 3.3.1) and by identifying the composition of ambient ions present during NPF events (cf.  
120 section 3.3.2). Finally, in chapter 4, we combine our results for the chemical composition of both gas-  
121 and aerosol-phases to discuss sources of precursor vapors producing secondary aerosol in the  
122 Antarctic Peninsula.

123

## 124 **2 Methods**


125

### 126 **2.1 Measurement site**

127

128 In-situ atmospheric gas- and particle-phase measurements were performed at the Argentinian  
129 Antarctic Station Marambio (64° 15'S - 56° 38' W) located on the Seymour Island in the  
130 northeastern-most part of the Antarctic Peninsula. Geographically (cf. Fig. 1), the island is on the  
131 north edge of the Weddell Sea. The station is operated year-round, with extensive scientific and  
132 logistic activities during the austral summer. The campaign instrumentation was located inside a  
133 laboratory container, specifically designed for continuous atmospheric composition measurements,  
134 sufficiently away (approx. 800 m) from the station main buildings and ideally placed fore-wind from  
135 the airstrip to avoid interfering contamination signals in our measurements. Comprehensive  
136 measurements of atmospheric composition and meteorology have been carried out in this location

137 since 2013. A description of the Marambio station surroundings, the measurement container, and  
138 some previous results can be found in earlier publications, e.g., Asmi et al. (2018).  
139



140  
141 **Figure 1:** Map of the Antarctic Peninsula, a), with zoom in the north sector, b), and in the Seymour Island with the Marambio station,  
142 in c). The measurement location is indicated by the red pin in every panels.

## 143 144 **2.2 Austral summer campaign 2018**

145  
146 The austral summer campaign 2018 for measuring the precursors contributing to NPF at the  
147 Marambio station was planned for the warm season, when the probability of NPF occurrence is high  
148 (Fiebig et al., 2014; Kerminen et al., 2018). The aim was to characterize NPF events, formation, and  
149 growth parameters, and resolve the aerosol chemical precursors from their source to their chemical  
150 pathways leading to nucleation. We measured concentrations of size segregated aerosol particles (~  
151 1 – 800 nm diameter) from 19 January to 23 February and chemical information of possible gas-phase  
152 precursors from 30 January to 20 February, continuously (whenever possible).

## 153 154 **2.3 Instrumentation**

### 155 156 *Measurements of aerosols and their gas-phase precursors*

157

158 We deployed a large suite of instruments, briefly described below. Instrument principles and  
159 operations are given in detail in the *Supplementary Information*.

160

161 (1) The chemical composition and concentration of neutral molecules was measured by a nitrate-  
162 based chemical ionization atmospheric pressure interface time of flight (CI-APi-TOF, ToFwerk A.G.)  
163 mass spectrometer (Jokinen et al., 2012). Its high-resolution analysis identifies and quantifies  
164 potential ambient gas-phase aerosol precursors, such as SA, MSA, IA, or HOM. In some instances,  
165 we used the APi-TOF to directly measure atmospheric ions (i.e., naturally charged anions or cations),  
166 allowing only atmospheric ions to enter the instruments (i.e., with no added chemical ionization) (cf.  
167 *Supplementary information*).

168

169 (2) A nano condensation nucleus counter (nCNC) combining a particle size magnifier (PSM,  
170 Airmodus Ltd.) with a condensation particle counter (CPC, Airmodus Ltd.) was employed to measure  
171 nano-particle concentration in the diameter range from 1 nm to 3 nm. The instrumentation and data  
172 inversions are extensively described in Vanhanen et al. (2011), Kangasluoma et al. (2016), Chan et  
173 al. (2020) and Lehtipalo et al. (2022) (cf. *Supplementary information*).

174

175 (3) A neutral cluster and air ion spectrometer (NAIS, Airel Ltd., Mirme and Mirme, 2013) provided  
176 number size distributions of neutral particles and naturally charged atmospheric clusters in the size  
177 range from 2 to 42 nm and from 0.8 to 42 nm, respectively. NAIS is well described in the literature  
178 and was operated according to the instructions of Manninen et al., 2016. (cf. *Supplementary*  
179 *information*).

180

181 (4) Aerosol particle size distributions were measured with a differential mobility particle sizer  
182 (DMPS): a Vienna-type differential mobility analyzer for particle size separation (Wiedensohler et  
183 al., 2012), followed by a condensation particle counter (CPC; TSI model 3772) for particle counting.  
184 The particle size distribution was measured in 25 separate size bins at a 6-minute time resolution.  
185 This measurement has been carried out continuously since the year 2013 but only data measured  
186 during the summer 2018 campaign are presented here. (cf. *Supplementary information*).

187

188 The CI-APi-TOF, nCNC and NAIS were operated with individual (horizontal) inlet lines with a  
189 minimum length (1 m, 0.8 m, and 0.9 m, respectively) to minimize losses of nanoparticles. The DMPS  
190 was operated in the station's common inlet (PM<sub>10</sub>, Asmi et al., 2018) and equipped with an additional  
191 PM<sub>1</sub> cyclone to prevent super micron particles from entering the setup.

192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225

### *Ancillary measurements*

Meteorological parameters were measured on the roof of the measurement container and recorded on a data logger (QML201L, Vaisala Ltd.) since 2013. Measured parameters include temperature and relative humidity (HMP155, Vaisala Ltd.), atmospheric pressure (PTB220, Vaisala Ltd.), wind speed and wind direction (ultrasonic anemometer: Thies 2D, Thies Clima), and radiation (pyranometer: CMP11, Kipp&Zonen).

### *Chemical composition of the aerosol phase*

Aerosol samples for chemical analysis were collected using a virtual impactor (VI, Loo and Cork, 1988), in which particles were divided to two size fractions: fine ( $D_p < 2.5 \mu\text{m}$ ) and coarse ( $2.5 \mu\text{m} < D_p < 10 \mu\text{m}$ ). Flow rate of the VI was  $16.7 \text{ L min}^{-1}$ , of which  $15 \text{ L min}^{-1}$  is used to collect the fine particles and the remaining  $1.7 \text{ L min}^{-1}$  is used for collection of the coarse particles. Particles were collected on 47-mm Teflon filters. Sampling time was one week. Collected filters were stored on petri slides and kept in a freezer ( $-18 \text{ }^\circ\text{C}$ ) until analysis.

The mass concentrations of sodium ( $\text{Na}^+$ ), ammonium ( $\text{NH}_4^+$ ), potassium ( $\text{K}^+$ ), magnesium ( $\text{Mg}^{2+}$ ), calcium ( $\text{Ca}^{2+}$ ), chloride ( $\text{Cl}^-$ ), nitrate ( $\text{NO}_3^-$ ), sulphate ( $\text{SO}_4^{2-}$ ), MSA and oxalate ( $\text{C}_2\text{O}_4^{2-}$ ) were determined from the filter substrates. Cation and anion analyses were done simultaneously with two ion chromatography systems (Dionex, ICS-2000). Filters were extracted in 10 mL of Milli-Q water with 10-min gentle rotation just before chemical analysis. The uncertainty of the IC analysis was estimated according to the analysis of standards as 5 % to 10 %, depending on the ion analyzed. In this study chemical composition of the fine particles is only presented.

## **2.4 Characteristics of New Particle Formation events**

NPF events were identified based on the appearance of particles in the nucleation mode (3 - 25 nm) showing signs of growth (e.g., particles reaching  $> 10 \text{ nm}$ ). A classification method was originally proposed by Dal Maso et al. (2005) based on DMPS data. In this study, we combined both DMPS and NAIS data to extend our observations to smaller sizes (Dada et al., 2018).

226 Condensation sink ( $CS$ ,  $s^{-1}$ ), which is a measure of how rapid precursor vapor concentrations are lost  
 227 to pre-existing particles, was calculated using the particle number size distribution measured by the  
 228 DMPS following the method proposed by Kulmala et al. (2012).

229  
 230 Growth rates ( $GR$ ) of particles between 3.8 nm to 12 nm were calculated using the 50 %  
 231 appearance time method (Lehtipalo et al., 2014; Dada et al., 2020). Formation rates ( $J_{1.5}$ ,  $J_3$ ,  $J_5$ , and  
 232  $J_{10}$ ) were calculated using the balance equation where the change in concentration of particles inside  
 233 a size bin is equivalent to the sources ( $J_{Dp}$ ) minus the available sinks ( $F_{Coag} + F_{Growth}$ ), as in Kulmala et  
 234 al. (2012).

$$235$$

$$236 \quad (1) \quad J_{Dp} = \frac{dN_{Dp}}{dt} + CoagS_{Dp} \cdot N_{Dp} + \frac{GR}{\Delta Dp} \cdot N_{Dp}$$

237 where  $Dp$  represents the lower diameter of the bin,  $N_{Dp}$  is the particle number concentration inside the size bin,  
 238 and  $GR$  is the growth rate of particles out of the bin.  $\Delta Dp$  is the difference between the upper and lower ends of  
 239 the size bin of interest.

240  
 241 The coagulation sink ( $CoagS_{Dp}$ ) is a measure of how rapid freshly formed particles of diameter  $Dp$  are lost to  
 242 pre-existing particles by collision or coalescence and is calculated as follows:

$$243$$

$$244 \quad (2) \quad CoagS_{Dp} = \int K(Dp, Dp') n(Dp') dDp' \cong \sum_{Dp'=Dp}^{Dp'=maxK} K(Dp, Dp') N_{Dp'}$$

245  
 246 where  $K(Dp, Dp')$  is the coagulation coefficient of particle sizes  $Dp$  and  $Dp'$ , those inside the bin of  $J_{Dp}$  and  
 247 those of pre-existing particles, respectively.  $N_{Dp'}$  is the number concentration of the pre-existing particles.

248  
 249 In this work, the formation rates of 1.5 nm particles ( $J_{1.5}$ ) were calculated using nCNC data in the size range of  
 250 1.5 nm to 3 nm. Formation rates  $J_3$ ,  $J_5$  and  $J_{10}$  were calculated using NAIS data (total particle mode – using the  
 251 differential mobility analyzer of negative voltage) at the size ranges of 3 - 7 nm, 5 - 9 nm, and 10 - 14 nm,  
 252 respectively. During events when the  $GR$  could not be calculated (i.e., four events in total) – due to the absence  
 253 of continuous growth within the size ranges -, a median growth rate calculated of all the events occurring in  
 254 the same month was used to estimate the formation rate as described by Kulmala et al. (2022).

255  
 256 In addition, the charged particle formation rates ( $J^{\pm 1.5}$  and  $J^{\pm 2}$ ) were calculated using ion number  
 257 concentration measured by NAIS in both polarities, whenever possible, to determine the contribution  
 258 of ion induced nucleation to the overall formation rate, according to the following equation:

259



$$(3) J_{Dp}^{\pm} = \frac{dN_{Dp}^{\pm}}{dt} + CoagS_{Dp} \cdot N_{Dp}^{\pm} + \frac{GR}{\Delta Dp} \cdot N_{Dp}^{\pm} + \alpha \cdot N_{Dp}^{\pm} \cdot N_{<Dp}^{\pm} - \chi \cdot N_{Dp} \cdot N_{<Dp}^{\pm}$$

261

262 where  $N_{Dp}^{\pm}$  is the concentration of ion in a given size range with a lower limit of 1.5 or 2 nm  
 263 (depending on the  $J$  associated size range) and the upper limit  $\Delta Dp$  being larger than that. The  
 264 coefficient  $\alpha$  is the ion-ion recombination coefficient and  $\chi$  is the ion-aerosol attachment coefficient.  
 265  $\alpha$  and  $\chi$  were  $1.6 \times 10^6 \text{ cm}^{-3} \text{ cm}^{-1}$  and  $0.01 \times 10^6 \text{ cm}^{-3} \text{ s}^{-1}$ , respectively (Kulmala et al., 2012).

266


### 267 3 Results

268

#### 269 3.1 Ambient conditions: Meteorological parameters

270

271 Ambient meteorological conditions during the campaign are presented in Figure 2. Many  
 272 sunny days are observed, occurring with above-zero degrees Celsius ambient temperatures, high  
 273 radiation, and relative humidity below 75 %. Even though the measurement site belongs to the  
 274 Antarctic continent, its position - at relatively high latitude ( $64^{\circ} 15' \text{ S}$ ) - gives daily solar cycles with  
 275 zero radiation during the short summer nighttime periods (cf. Fig. 2b). During daytime, higher  
 276 radiation enhances photo-active emissions from the sea/land ecosystems and promotes atmospheric  
 277 photochemistry. This is expected to lead to day-time appearance of gas-phase molecules and/or  
 278 molecular clusters of condensing vapors that can lead to new particle formation and/or aerosol  
 279 growth. Furthermore, since temperatures above freezing also lead to ice melting (especially in the  
 280 Weddell Sea area), they are likely to enhance chemical fluxes between thinning ice, ocean, and air  
 281 (Notz, 2009). These clear sky, warm, sunny days during summertime are known to be optimal  
 282 weather conditions for NPF to occur in the mid-latitude regions (Dada et al., 2017) and in the Arctic  
 283 (e.g., Beck et al. 2021) and Antarctic regions (Weller et al., 2015; Jokinen et al., 2018; Baccarini et  
 284 al., 2020; Brean et al., 2021). Analysis of winds (cf. Fig. 2e) revealed that periods of the highest  
 285 temperature are seen when wind is blowing from north, bringing warmer air across from the Southern  
 286 Ocean. Predominant winds were seen from North-Northwest and South-Southwest sectors. The  
 287 strongest winds were recorded from the South ( $180^{\circ}$ ), also driving primarily cold air from the  
 288 continental plateau, agreeing with Asmi et al. (2018).




289  
 290 **Figure 2:** Time series of meteorological parameters: a) temperature (dark blue), dew point (light blue), b) global radiation (orange), c)  
 291 relative humidity (light green), d) atmospheric pressure (blue green), e) wind direction (dark blue bars) and wind speed (medium blue  
 292 line). The data are presented with a 10-min resolution and reference time set as local standard, UTC-3.

293  
294  
295  
296  
297  
298  
299  
300  
301  
302

### 3.2 Observation & Characterization of NPF events

A total of 13 NPF events (occurring on 12 days) were observed during the 35-day austral summer campaign. Figure 3a shows the continuous size distribution series over the total measurement period from NAIS measurement (size range: 1.8 – 42 nm) and Figure 3b combined the data from nCNC (size shown: 1.2 – 2.7 nm), the NAIS (2.7 nm - 12 nm), and the DMPS (12 nm – 800 nm). Daily number size distribution plots are shown for the 12 event days separately in the *Supplementary Information* (Fig. S3) as well as continuous size distribution series for ion mode (Fig. S4).



303

304 **Figure 3:** a) Number size distribution from NAIS measurement from 1.8 nm to 42 nm over the whole measurement campaign and b)  
305 combined size distribution of aerosol particle from 1.2 nm to 800 nm for 3-7 February 2018. The last surface plot combines data from  
306 nCNC (1.2 - 2.7 nm), NAIS (particle mode, 2.7 - 12 nm) and DMPS (12-800 nm), whenever the data were available and according to  
307 DMPS data flag (only unpolluted data are shown from the DMPS measurements).

308 On each of the NPF events observed during the campaign, high population of sub-3 nm aerosol  
309 particles were seen with concentration larger than  $10^4 \text{ cm}^{-3}$ . The particles were observed to grow to  
310 Aitken mode size range (25 - 100 nm) but rarely reached the accumulation mode sizes (i.e., > 100  
311 nm) (cf. Fig. 3b).

312

313 Strong NPF events were identified before midday - local time - (66 % of events), lasting longer than  
314 events occurring later in the afternoon. Afternoon events (33 % of the total events) never lasted longer  
315 than 2.5 hours. NPF events did vary from day to day: On certain days, growth of the particles was  
316 interrupted by abrupt changes in wind direction - resulting in an inability to retrieve a growth or a  
317 formation rate within reasonable uncertainty (e.g., 28 January) -; on other days multiple NPF events  
318 were observed (e.g., 16 February); or NPF events appeared to have a bumped shape, appearing  
319 sporadically - resembling those observed in the Arctic by Beck et al. (2021) – such as the events  
320 observed on 28 January and 6 February.

321

322 Formation rates were calculated for particle sizes of 1.5, 2, 3, 5 and 10 nm particles, as summarized  
323 in Table 1. These formation rates were calculated for the smallest possible particles but also for bigger  
324 sizes to ease the comparison with the literature.

325

326 Formation rates of 1.5 nm particles (i.e.,  $J_{1.5}$ ) were surprisingly high, reaching the maximum rates of  
327 9.2, 10.0, or even  $19 \text{ cm}^{-3} \text{ s}^{-1}$ , on February 11<sup>th</sup>, 10<sup>th</sup>, and 3<sup>rd</sup>, respectively. These significantly high  
328 values are comparable to those calculated for highly vegetated rural location or even urban areas (e.g.,  
329 Deng et al., 2020; Kontkanen et al., 2016; Yu et al., 2014), which is surprising due to the remote  
330 nature of the environment at our measurement site. For example, formation rates for 1.5 nm particle  
331 on the order of  $0.030 - 0.013 \text{ cm}^{-3} \text{ s}^{-1}$  were estimated for the Aboa research station - on the other side  
332 of the Weddell Sea in East Antarctica (Jokinen et al. 2018). A recent study observing nucleation  
333 phenomena at the peninsula close by our measurement location (Brean et al., 2021) found a  
334 remarkably high  $J_{1.7}$  value reaching up to  $3.07 \text{ cm}^{-3} \text{ s}^{-1}$  but still lower than our observed values.  
335 Although the formation rates at 1.5 nm are higher than those reported previously, the rates for larger  
336 particles (i.e.,  $J_3$ ,  $J_5$ ,  $J_{10}$ ) are comparable with the previously reported studies. This discrepancy  
337 between formations rates of 1.5 nm particles and of bigger particles could be attributed to favorable  
338 nucleation conditions (i.e., sufficient nucleating vapors) but a lack of condensable vapors contributing  
339 to particle growth which, in all, reduces the probability of particle survival.

340

341 The nucleation rates for 3 nm particles in our study ranged from  $0.13 \text{ cm}^{-3} \text{ s}^{-1}$  to  $3 \text{ cm}^{-3} \text{ s}^{-1}$ , which  
342 were, on average, also significantly higher than  $0.03 \text{ cm}^{-3} \text{ s}^{-1} < J_3 < 0.14 \text{ cm}^{-3} \text{ s}^{-1}$  reported from  
343 Jokinen et al. (2018) study or from Kecorius et al. (2019) who showed  $J_3$  values from  $0.080 \text{ cm}^{-3} \text{ s}^{-1}$   
344 to  $0.319 \text{ cm}^{-3} \text{ s}^{-1}$  during a polar cruise in the Arctic.

345 On the other hand, our observation agrees with the  $J_{10}$  reported by Kyrö et al. (2013) ranging from  
346 0.003 to 0.3  $\text{cm}^{-3} \text{s}^{-1}$ , with  $0.006 \text{ cm}^{-3} \text{ s}^{-1} < J_{10} < 0.37 \text{ cm}^{-3} \text{ s}^{-1}$  in our case.

347

348 Additionally, to account for ion induced nucleation, we calculated the formation rates for charged  
349 ions of 1.5 nm ( $J^{+/-1.5}$ ) separately using the ion concentrations measured with the (N)AIS in ion mode.  
350 The formation rates for positive 1.5 ions,  $J^{+1.5}$ , ranged from  $3.6 \times 10^{-4}$  to  $8.3 \times 10^{-2} \text{ cm}^{-3} \text{ s}^{-1}$  and  $J_{1.5}$   
351 was from  $2.2 \times 10^{-3}$  to  $5.4 \times 10^{-2} \text{ cm}^{-3} \text{ s}^{-1}$ . Our results are significantly lower than the maximum  $J_{1.5}$   
352 of  $0.33 \text{ cm}^{-3} \text{ s}^{-1}$  reported in Beck et al. (2021) or from Kyrö et al. (2013) who showed  $0.02 \text{ cm}^{-3} \text{ s}^{-1} <$   
353  $J_{1.6} < 4.2 \text{ cm}^{-3} \text{ s}^{-1}$  - in two coastal Antarctic sites, but in the range reported from the Arctic ocean by  
354 Kecorius et al. (2019, e.g.,  $0.026 \text{ cm}^{-3} \text{ s}^{-1} < J_{1.6} < 0.060 \text{ cm}^{-3} \text{ s}^{-1}$ ). In comparison to 1.5 nm-neutral  
355 particle formation rates, the ion formation rates do not represent a major contribution (ratio  $J_{neutral}$   
356  $/J_{ion} \sim 10^3$ ), implying that the relatively high neutral nucleation rates are not primarily due to ion-  
357 mediated nucleation.

358

359 Particle growth rates were calculated for 3.8 nm to 12 nm diameter size or alternatively up to 11 nm  
360 and 8 nm for event #1 and event #13, respectively, whenever the shape of the number size distribution  
361 was continuously increasing. Averages for each event are also presented in Table 1. Similarly to  
362 formation rates of the smallest particles, the growth rates were remarkably higher ( $1.2 \text{ nm h}^{-1} < GR$   
363  $< 10.9 \text{ nm h}^{-1}$ ) than previously reported for other Antarctic sites. In comparison, Weller et al. (2015)  
364 reported growth rates from 3 to 25 nm particles ranging from 0.06 to 0.9  $\text{nm h}^{-1}$  at Neumayer III  
365 station, Jokinen et al. (2018) showed  $0.26 \text{ nm h}^{-1} < GR < 1.30 \text{ nm h}^{-1}$  at Aboa and Brean et al. (2021)  
366 published growth rates for 4.5 - 10 nm particles of 0.41  $\text{nm h}^{-1}$  to 0.58  $\text{nm h}^{-1}$  measured at the  
367 Peninsula. Additionally, Kerminen et al. (2018) reviewed aerosol characterization from many  
368 different field studies and indicated an upper growth rate estimation of 5.5  $\text{nm h}^{-1}$  (as 95<sup>th</sup> percentile)  
369 in Antarctic sites versus 4.1  $\text{nm h}^{-1}$  in Arctic environments. Recent Arctic studies by Kecorius et al.  
370 (2019) reported  $0.62 \text{ nm h}^{-1} < GR_{(3-7 \text{ nm})} < 4.25 \text{ nm h}^{-1}$  while Collins et al. (2017) reported averaged  
371 growth rates of  $4.3 \pm 4.1 \text{ nm h}^{-1}$ .

372

373 **Table 1:** Summary of NPF event classification, formation and growth parameters calculated during the Marambio austral summer campaign. The pollution flag is determined based on DMPS data  
 374 according to NILU/EBAS format classification at the start time of the event (when data available, 000: clean data, 189: data coming from a possibly polluted sector (e.g., station main buildings),  
 375 599: suspected pollution from unknown source). All formation rates are averaged between the start and end of the characterized events.

376  
 377

#	Date (dd/mm)	Start Time (SLT*, hh.mm)	Duration (hh.mm)	Flag (DMPS)	Formation Rates ( $J_{size} \text{ cm.s}^{-1}$ )					Condensation Sink ( $s^{-1}$ )	Growth Rates ( $Gr_{size} \text{ nm.h}^{-1}$ ) $Gr_{3.8-12}^{**} \pm \text{err.}$
					$J_{1.5}$ ( $J_{1.5}/J_{1.5}^+$ )	$(J_2/J_2^+)$	$J_3$	$J_5$	$J_{10}$		
1	19/01	14.35	0.55	189	-	-	-	-	-	-	$9.7 \pm 3.2$ ( $Gr_{3.8-11}$ )
2	24/01	18.30	2.26	-	$-(1.3e^{-2} / 1.2e^{-2})$	$(1.2e^{-2} / 1.9e^{-2})$	$6.0e^{-1}$	$3.3e^{-2}$	$6.1e^{-3}$	-	$2.4 \pm 0.7$
3	25/01	14.00	2.25	000	$-(1.2e^{-2} / 9.8e^{-3})$	$(4.9e^{-1} / 6.9e^{-1})$	1.2	$4.7e^{-1}$	$3.7e^{-1}$	$7.5e^{-4}$	$10.9 \pm 2.8$
4	27/01	18.50	2.06	189	$4.3$ ( $3.6e^{-4} / 3.6e^{-3}$ )	$(6.5e^{-3} / 2.0e^{-2})$	$6.9e^{-1}$	$7.2e^{-1}$	$1.8e^{-1}$	$3.8e^{-4}$	-
5	28/01	9.13	7.25	000	$3.6$ ( $8.5e^{-3} / 6.7e^{-3}$ )	$(1.9e^{-2} / 3.4e^{-2})$	$1.6e^{-1}$	$4.8e^{-2}$	$1.2e^{-2}$	$2.2e^{-4}$	-
6	03/02	8.13	7.04	~000	$19.0$ ( $3.4e^{-2} / 2.5e^{-2}$ )	$(2.9 / 4.0)$	3.0	$5.5e^{-1}$	$2.9e^{-1}$	$1.1e^{-3}$	$2.4 \pm 0.4$
7	05/02	8.35	6.36	000	$1.5$ ( $7.2e^{-3} / 5.2e^{-3}$ )	$(3.9e^{-3} / 4.3e^{-3})$	$1.8e^{-1}$	$1.3e^{-1}$	$6.8e^{-2}$	$1.3e^{-3}$	$1.6 \pm 0.7$
8	06/02	9.35	4.25	000	$1.1$ ( $8.3e^{-2} / 5.4e^{-2}$ )	$(2.7 / 1.7)$	$1.3e^{-1}$	$9.0e^{-3}$	$1.2e^{-2}$	$4.8e^{-4}$	$2.4 \pm 1.2$
9	10/02	11.16	3.52	~000	$10.0$ ( $4.2e^{-3} / 2.2e^{-3}$ )	$(1.8e^{-1} / 2.1e^{-1})$	$8.3e^{-1}$	$3.6e^{-1}$	$3.6e^{-2}$	$3.7e^{-4}$	-
10	11/02	10.27	6.16	000	9.2 (-/-)	$(1.3 / 8.9e^{-1})$	$9.2e^{-1}$	$3.5e^{-1}$	$2.1e^{-1}$	$4.0e^{-4}$	-
11	12/02	10.40	5.40	189	$-(8.1e^{-3} / 1.9e^{-2})$	$(4.1e^{-1} / 5.9e^{-1})$	2.1	2.7	$1.8e^{-1}$	$2.2e^{-3}$	$1.2 \pm 0.7$
12	16/02	11.18	6.03	599	$3.7$ ( $5.9e^{-2} / 3.2e^{-2}$ )	$(7.2e^{-2} / 4.3e^{-2})$	$5.9e^{-1}$	$4.0e^{-1}$	$1.2e^{-1}$	$2.4e^{-5}$	$3.6 \pm 0.7$
13					-	-	-	-	-	$2.2 \pm 2.9$ ( $Gr_{3.8-8}$ )	

\* Standard local time, UTC -3

\*\* Growth rates were calculated ideally from 3.8 nm to 12 nm. If the growth on that range cannot be determined in that range, the specific range is indicated in subscript as  $Gr_{x-x'}$ .


378

### 3.3 Chemical composition of the gas-phase precursor molecules

#### 3.3.1 Gas-phase contribution to NPF


Measurements utilizing nitrate based chemical ionization mass spectrometry (CI-APi-TOF) detected gas-phase molecules and molecular clusters that have affinity for proton transfer with nitrate reagent ion. These included SA, MSA, and IA - detected as  $\text{HSO}_4^-$  and  $\text{HNO}_3\text{HSO}_4^-$  ( $m/z$  96.9601 Th and 159.9557 Th),  $\text{CH}_3\text{SO}_3^-$  and  $\text{HNO}_3\text{CH}_3\text{SO}_3^-$  ( $m/z$  94.9808 Th and 157.9765 Th),  $\text{IO}_3^-$  and  $\text{HNO}_3\text{IO}_3^-$  ( $m/z$  174.8898 Th and 237.8854 Th), respectively. Concentrations were calculated from high resolution peak fitting and reported below. The time series of those compounds during the whole campaign, according to the data availability of the CI operation, are shown in Figure 4.

The measured gas-phase concentrations of the species of interest showed maxima of  $\sim 2.6 \times 10^7$  molecules  $\text{cm}^{-3}$ ,  $\sim 2.3 \times 10^7$  molecules  $\text{cm}^{-3}$  and  $\sim 3.6 \times 10^6$  molecules  $\text{cm}^{-3}$  for the total SA, MSA, and IA concentrations, respectively. The three gas-phase compounds seemed to evolve with roughly similar behavior, likely due to favorable weather conditions promoting atmospheric chemical reactions and especially by enhancing photochemical oxidation reactions within a stable boundary layer. A significant and simultaneous increase in SA, MSA, IA concentrations was observed during peaking radiation time close to noon-time – coinciding with local zenith time, i.e., solar noon that is slightly later than midday - and prior to many NPF events, though to a smaller extent for IA. This clearly demonstrates the diurnal nature of the emission of those species, and their transport – especially for SA and IA – as shown in Figure 5. Additionally, it is worth mentioning that even though the MSA concentrations were on average higher than those of SA, event days correlated strongly with peaking SA concentration (then becoming higher than MSA concentration) while IA – whose concentration remained unchanged independently of NPF occurrence - was significantly lower than SA or MSA concentrations.



405  
406  
407  
408

**Figure 4:** High resolution time series of  $\text{H}_2\text{SO}_4$  (SA, red),  $\text{CH}_4\text{SO}_3$  (MSA, orange),  $\text{HIO}_3$  (IA, green). Numbers specified on the top of the figure refer to the event number as introduced in the NPF analysis (Cf. Table 1). The data are averaged to 10 min with local time (UTC -3). Note that the missing points are due to switch of the measurement mode and / or data availability.



409  
410  
411

**Figure 5:** Statistical diurnal time series of  $\text{H}_2\text{SO}_4$  (SA, red),  $\text{CH}_4\text{SO}_3$  (MSA, orange),  $\text{HIO}_3$  (IA, green) observed during non-event days a) and event days b). The solid lines represent the median hourly average concentrations, the dotted lines are the concentration means and the colored shaded areas show data points within the 25<sup>th</sup> and the 75<sup>th</sup> percentiles. Nighttime is represented by the gray-shaded area with median theoretical sunsets and sunrises occurring during the campaign. The actual sunset and sunrise values could vary approx.  $\pm 1.25$  hours from the beginning to the end of the campaign.

416  
417  
418  
419  
420

Figure 5 shows the diurnal variations of SA, MSA and IA for event b) and non-event days a), separately. In both figures, we could see a clear rise in concentrations of SA, MSA and IA throughout the day, with maxima observed at midday.



421 While both MSA and SA concentrations are on average increasing after sunrise, IA concentration  
422 started to increase already during nighttime, shortly after 01:00 SLT, suggesting that strong  
423 photochemistry conditions (i.e., direct solar radiation) might not be necessary to produce the observed  
424 IA agreeing with He et al. (2021). Earlier studies have shown that IA is even anticorrelated with (still  
425 omnipresent) solar radiation in more continental Antarctica (Jokinen et al., 2018). It is likely that even  
426 very low levels of solar radiation would be sufficient to saturate the iodic acid production (Beck et  
427 al., 2021). Distinctively, a clear difference in the statistical series can be seen between event and non-  
428 event days, with more than doubled SA concentrations at zenith-time (factor of 2.25). MSA  
429 concentrations are only slightly higher in the morning and afternoon – with an apparent drop  
430 occurring from 14:00 to 16:00 – although the starting concentrations at sunrise and sunset are also  
431 smaller, depicting strong concentration variability as compared to non-event days. On the other hand,  
432 IA is on average lower during event-days – e.g., from  $\sim 4.5 \times 10^5$  molecules  $\text{cm}^{-3}$  vs  $\sim 3 \times 10^5$   
433 molecules  $\text{cm}^{-3}$ , with a factor of 1.5 smaller at zenith-time. This last observation indicates that IA is  
434 likely not a key contributor to particle formation, while the duo SA-MSA could influence the aerosol  
435 phase as already suggested by Beck et al. (2021), Hodshire et al. (2019) and Willis et al. (2016).

436


437

### 438 **3.3.2 Molecular characterization of aerosol / ion precursors:**

#### 439 **Study case on 16 February**

440

441 The aerosol event observed on 16 February was particularly interesting due to the occurrence  
442 of two consecutive NPF events within the same day. An overview of both aerosol concentrations /  
443 parameters and reactive trace gases concentrations for 16 February is shown in Figure 6 with a) the  
444 particle number size distribution series from NAIS measurement, b) time series of total particle  
445 concentration within several size modes (i.e., cluster, nucleation, Aitken and accumulation), c)  
446 estimated  $J$  rates for 1.5, 3, 5 and 10 nm particles and d) time series for SA, MSA and IA.



447

448

449

450

451

452

453

454

455

456


**Figure 6:** Overview of aerosol formation event on 16 February 2018: a) Number size distribution from NAIS measurements (Particle mode). b) Concentrations of particles modes (cluster: sub- 3 nm – medium light blue, nucleation: 3 – 25 nm – light blue, Aitken: 25 – 100 nm – medium dark blue, accumulation: 100 – 1000 nm – dark blue). Note that the missing points for Aitken and Accumulation mode are due to data filtering from suspected pollution from the DMPS data set. c) Formation rate estimation for J<sub>1.5</sub> (neutral - light blue, (-) - blue green, (+) - green), J<sub>2</sub> ((-) - orange, (+) - yellow), J<sub>3</sub> (medium light blue), J<sub>5</sub> (medium dark blue) and J<sub>10</sub> (dark blue). Note that the formation rate calculation depends on the condensation/coagulation sinks calculated from the DMPS data. d) High resolution time series of H<sub>2</sub>SO<sub>4</sub> (red), CH<sub>4</sub>SO<sub>3</sub> (orange), HIO<sub>3</sub> (green). Note that the data gap is due to ion mode of APi-measurement. All data are averaged / estimated with a 10 min time resolution, with standard local time (UTC -3) as clock reference.

457 A clear NPF episode occurred from about 11:15 in the morning until sunset with a net increase of  
458 cluster and nucleation mode particle concentrations. Close to noontime all trace gases of interest were  
459 significantly increasing (Fig. 6d). MSA increased by almost a factor of 10 by 13:00 ( $[MSA]_{t=13.00} =$   
460  $1.3 \times 10^6$  molecules  $\text{cm}^{-3}$ ) and by more than 2 orders of magnitude at 16:00 ( $[MSA]_{t=17.30} = 9.6 \times 10^6$   
461 molecules  $\text{cm}^{-3}$ ) as compared by the minimum baseline of  $\sim 2 \times 10^5$  molecules  $\text{cm}^{-3}$  for both MSA  
462 and SA. SA maximum concentration was  $6.5 \times 10^6$  molecules  $\text{cm}^{-3}$ , also observed at 13:00.

463  
464 IA concentrations were fluctuating throughout the day with maximum concentrations of  $\sim 2.05 \times 10^5$   
465 molecules  $\text{cm}^{-3}$  and  $\sim 2.35 \times 10^5$  molecules  $\text{cm}^{-3}$ , respectively around 05:30 – 06:30 and shortly after  
466 13:00. The net increase of all these gas-phase species occurred quasi-simultaneously as the increase  
467 of formation rate of the smallest particles and ions (Fig. 6c), occurring around noon. Interestingly,  
468 several sudden, short-lasting increases for all SA, MSA, and IA concentrations were seen even before  
469 13:00, matching bumps in nucleation mode particles around 09:00 and 11:00 and one to two orders  
470 of magnitudes increase of cluster mode particles at 11:00. The high concentrations of SA, MSA (and  
471 possibly IA) likely trigger the observed aerosol processes (i.e., nucleation and/or growth of particles).

472  
473 We investigated the mass spectrum further and show a complementary mass spectrum in Figure S2.  
474 Any HOM (Bianchi et al., 2019; Ehn et al., 2014) could be reliably resolved from the mass spectra  
475 analysis. We estimate that the total oxidized organic contribution, possibly condensable HOM, could  
476 not exceed  $5.5 \times 10^6$  molecules  $\text{cm}^{-3}$ , between 16:30 and 17:30; assuming mass spectral peaks found  
477 above 200 Th with positive mass defect could be HOM (*cf. Supplementary*). However, the actual  
478 HOM contribution is likely to be only a fraction of this estimation. Therefore, it is unlikely that HOM  
479 significantly contributed to nucleation or growth of newly formed particles.

480  
481 Specifically for this event, we complemented the CI-APi-TOF measurement with negative APi-TOF  
482 ambient measurement mode, from 14:52 to 15:53, as reported in Table S1. By showing the  
483 divergence of the exact molecular mass of a detected ion in high resolution from its integer mass (i.e.,  
484 defined as mass defect), a mass defect plot illustrates the mass defect of selected ions (in y-axis) over  
485 a studied mass range (in x-axis). Compared to a typical mass spectrum this has the advantage to show  
486 the most significant ion population at once, rather independently of signal intensity which is then  
487 scaled by the marker size, easing the identification of e.g., clustering mechanisms as further  
488 discussed. There, each point of the mass defect plot corresponds to a unique atomic composition. An  
489 example of such representation is shown in Figure 6, for the study case of 16 February.



490

491 **Figure 7:** Mass defect plot for (-) API-TOF measurements, on 16 February from 14:52 to 15:52. The figure type represents the deviation  
 492 of high resolution fitted peaks exact mass normalized to the unit mass detected for each peak y-axis - e.g.  $\text{Br}^{\text{exact mass}} = 78.9189$ ,  $\text{Br}^{\text{unit}}$   
 493  $\text{mass} = 79$ ,  $\text{Br}^{\text{mass defect}} = -0.0811$  – while the x-axis represents the mass range (m/z). Data average is 1 hour for each spectrum to improve  
 494 the signal to noise ratio and the peak shape statistics.

495 The API-TOF ion mode showed TIC maximum values around  $50 \text{ ions s}^{-1}$ , detecting only ions that are  
 496 naturally charged. The highest signals (i.e., intensity shown by marker sizes, cf. Fig. 7) are bisulfate  
 497 and its multimers  $(\text{H}_2\text{SO}_4)_{0-4}\text{HSO}_4^-$ , respectively at  $m/z = 96.9601$ ,  $m/z = 194.9275$ ,  $m/z = 292.8949$   
 498 and  $m/z = 390.8622$ . Along with SA multimers, successive addition of sulfuric acid and ammonia  
 499  $\text{NH}_3$  formed clusters represented with the light-blue dots, within the mass range 400 – 1250 Th. These  
 500 sulfuric acid (bisulfate) - ammonia clusters were found with up to 10 additions of  $\text{H}_2\text{SO}_4$  and 9  $\text{NH}_3$   
 501 on top of the  $\text{HSO}_4^-$  core ion. In line with previous field observations (e.g., Schobesberger et al., 2015;  
 502 Bianchi et al., 2016; Jokinen et al., 2018; Yan et al., 2018; Beck et al., 2021; Sipilä et al., 2021) and  
 503 laboratory studies (e.g., Kirkby et al., 2011; Schobesberger et al., 2015) showed that ammonia is  
 504 detected only in clusters with 3 or more  $\text{H}_2\text{SO}_4$  molecules around the  $\text{HSO}_4^-$  core.

505

506 The numerous clusters containing both SA and ammonia, as well as the high intensity of the  
 507 respective cluster signal (depicted by the marker size in Figure 6) suggest a high concentration of  
 508 such ion group in the gas-phase which also could indicate an ion induced nucleation driven by the  
 509 ternary system SA-ammonia(-water) - similar to observations made by Jokinen et al. (2018) in East

510 Antarctica as well as measured in multiple other locations around the globe (e.g., Bianchi et al., 2016;  
511 Yan et al., 2018; Beck et al., 2021; Sipilä et al., 2021).

512

513 Additional clusters containing SA and DMA (principally as  $(\text{H}_2\text{SO}_4)_2 \cdot (\text{CH}_3)_2\text{NH} \cdot \text{HSO}_4^-$  ( $m/z =$   
514  $337.9527$ ) and as  $(\text{H}_2\text{SO}_4)_3 \cdot (\text{CH}_3)_2\text{NH} \cdot \text{HSO}_4^-$  ( $m/z = 435.9210$ ) but also one larger cluster,  
515  $(\text{H}_2\text{SO}_4)_7 \cdot (\text{CH}_3)_2\text{NH} \cdot \text{HSO}_4^-$  ( $m/z = 827.7893$ ) was identified. Similarly to the finding of Brean et al.  
516 (2021) who detected various SA-amine clusters around the peninsula, SA-DMA- $\text{H}_2\text{O}$  driven  
517 nucleation could also occur leading to formation rates significantly higher than those observed in the  
518 SA- $\text{NH}_3$ - $\text{H}_2\text{O}$  system (Kürten et al., 2014; Almeida et al., 2013). Furthermore, laboratory studies  
519 (e.g., Schobesberger et al., 2013) have shown that (negative) ion composition from APi-TOF analysis  
520 can match the cluster composition of the neutral clusters involved in NPF. Therein, the appearance  
521 of DMA in the SA dimer and trimer suggests sufficient DMA to trigger nucleation. On the other hand,  
522 the large clusters containing  $\text{NH}_3$  is consistent with high  $\text{NH}_3$  gas concentrations that can dominate  
523 nano-particle growth. This highlights that DMA is 1000 times more effective than  $\text{NH}_3$  for nucleation;  
524 so that sub-ppt DMA and 100 – 1000 ppt levels (and above) of  $\text{NH}_3$  could explain the ion cluster  
525 distribution observed in Figure 7.

526


527 MSA-containing ions can be identified (cf. Fig. 7, orange dots) at  $m/z = 94.9808$  and  $m/z = 192.9482$ ,  
528 respectively. Several studies have shown MSA can be involved in atmospheric nucleation (Hodshire  
529 et al., 2019) - especially with clusters containing MSA, SA and DMA (Bork et al., 2014). However,  
530 in our case, MSA-containing clusters were only found as  $\text{CH}_3\text{SO}_3^-$  or as  $\text{CH}_3\text{SO}_3\text{H} \cdot \text{HSO}_4^-$ . MSA was  
531 not observed in large clusters that would contain altogether SA,  $\text{NH}_3$  and/or DMA. Therefore it is un-  
532 likely that MSA would contribute to the nucleation observed in this study case (at least not through  
533 to negative ion pathway).

534

535 Finally, we identified the presence of halogen compounds, (cf. Fig. 7, yellow and green dots), such  
536 as  $\text{Br}^-$  ( $m/z = 78.9189$ ),  $\text{I}^-$  ( $m/z = 126.9050$ ),  $\text{IO}_3^-$  ( $m/z = 174.8898$ ),  $\text{HIO}_3\text{NO}_3^-$  ( $m/z = 237.8854$ ),  
537  $\text{HIO}_3\text{HSO}_4^-$  ( $m/z = 272.8571$ ). The presence of halogenated species as natural ions could also indicate  
538 that these compounds contribute to aerosol processes. However, these ions were only identified as  
539 small halogen clusters of low molecular weight and with only low signal intensity. Considering  
540 previous observations in the Arctic and coastal environment, Sipilä et al. (2016) only shows iodine-  
541 related nucleation as successive addition of multiple iodic acid and water group, it is unlikely that IA  
542 promotes nucleation either through neutral or ion-induced mechanism at this Antarctic site.

543

### 3.3.3 Chemical composition of the aerosol phase



545

546 **Figure 8:** Chemical composition of water-soluble ions from weekly fine aerosol samples (PM<sub>2.5</sub>)

547 Asmi et al. (2018) reported aerosol chemical composition from weekly filter analysis  
 548 containing marine-originated sea salt aerosols and the presence of ammonium, sulfate, nitrate and  
 549 mesylate (CH<sub>3</sub>SO<sub>3</sub><sup>-</sup>). Here, we performed the identical analysis, which agrees with the previously  
 550 reported results.

551

552 Until February 16<sup>th</sup>, the fine aerosol composition remains rather constant with high amounts of non-  
 553 sea salt sulphate (nss-sulphate, i.e., sulphate - 0.246 × sodium - in mass concentration, Brewer (1975))  
 554 which is partly neutralized by ammonia and a significant contribution from secondary marine MSA.  
 555 The presence of Na<sup>+</sup>, Cl<sup>-</sup> and Ca<sup>2+</sup> ions also indicates contribution from primary marine sea salt and  
 556 continental soil minerals. Aerosol composition and mass concentrations are very similar to those  
 557 found in Asmi et al. (2018) for Marambio summer aerosol. The highest concentrations of ions are  
 558 measured during the active nucleation period in the beginning of February. By the end of the month,  
 559 the concentrations of MSA, ammonium and calcium ions decrease. No indication of anthropogenic  
 560 contamination is found in the samples.

561

562 The bulk aerosol chemical composition (Fig. 8) reflects rather well the observed cycle of the  
 563 secondary particle formation and provides further evidence of the important role of sulphate/SA,  
 564 methane sulfonate, and ammonia in the production and especially in the growth of Antarctic  
 565 secondary aerosol. The qualitative agreement of gas-phase and particle-phase compositions indicates  
 566 that it is likely possible that these compounds originate from similar marine/coastal sources.

567

568


### 3.3.4 Source of gas-phase precursors

569

570 Backward trajectories from sampled air masses (Fig. 9) were analyzed and seen to originate  
571 mostly from the West sector passing through the Southern Ocean before surrounding the Peninsula  
572 from its Northwest side - coinciding with the observed wind directions. The observed trajectories  
573 could explain the low concentration of IA, whose variability was independent of NPF occurrence due  
574 to (1) the absence of travel over the Weddell littoral seaside prior to the NPF or (2) because the algae  
575 bloom on the melting sea ice – possibly responsible for an increase reactive iodine of organic origin-  
576 has already occurred earlier in the season. On the other hand, the air is clearly enriched in MSA and  
577 SA when passing over the ocean that has the highest DMS concentration over the Dec-Feb months  
578 (Lana et al., 2011). In some instances, air mass trajectories are seen to turn over the North of the  
579 peninsula, which can indicate a possible enrichment of gas produced by the fauna on the land or at  
580 the shore (e.g., local source of ammonia and amines from penguin colony established at approx. 8 km  
581 South of the sampling site during summertime). With a majority of trajectories originating from the  
582 Bellingshausen Sea,  $\text{NH}_3$  and various type of amines likely originate from the (melting) ocean as  
583 discussed in Dall'Osto et al. (2017).

584

585 Figure 9c also shows the air mass trajectories altitudes versus time. While Kerminen et al. (2018)  
586 reviewed that Antarctic NPF would easily take place within the free troposphere, both the number  
587 size distribution and the indicated altitude flight path point toward nucleation processes occurring at  
588 the surface layer considering the lifetime ( $\lambda$ ) of key gas-phase precursors (e.g.,  $\lambda_{\text{SA}} < 0.5$  h, Fiedler et  
589 al. (2005)). Similar back trajectories were calculated for non-event days, separately for days with  
590 high SA concentrations (i.e.,  $[\text{SA}] > 2 \times 10^6$  molecules  $\text{cm}^{-3}$ ) and for days with lower SA (Fig. S5a-c  
591 and Fig. 5a'-c', respectively). The result of the analysis as well as the model characteristics are shown  
592 in the *Supplementary Information*.



593  
 594 **Figure 9:** Backward trajectories for recorder NPF events a) (numbering according to classification Table 1). b) shows a zoom-in over  
 595 the Antarctic Peninsula and c) shows the trajectory altitudes colored by formation rates  $J_3$  values. In panel a) and b) the black and grey  
 596 lines delimit the sea ice extent as average for January and February 2018, respectively. Sea ice data are from: <ftp://sidads.colorado.edu/DATASETS/NOAA/G02135/south/monthly/> (Fetterer et al., 2017).  
 597

598 Local wind analysis (Fig. S6) was performed to determine possible local emission sources. In essence,  
 599 no prevalent wind direction was found specifically for NPF days. However, in some instances,  
 600 moderate to strong winds were seen to originate from the South/South-West sector where strong  
 601 emission from the fauna is likely to occur by the shore at this season. Those days, 25 January (event  
 602 # 3), 10 February (event # 9), and 11 February (event # 10), were among the highest growth ( $GR_{3.8-12} = 10 \text{ nm h}^{-1}$ ) and the highest formation rates ( $J_{1.5} = 10 \text{ cm}^{-3} \text{ s}^{-1}$  and  $J_{1.5} = 9.2 \text{ cm}^{-3} \text{ s}^{-1}$ ), respectively,  
 603 which agree with our hypothesis attributing high emissions of e.g., ammonia / amines to the bird  
 604 colony. On the other hand, Event #12 has a different trajectory and prevalent wind direction against  
 605 the South sector. Given the fact that the anion mass defect (Fig.7) suggests significant presence of  
 606 ammonia and possibly DMA on this event, specifically, nitrogen-containing base compounds origin  
 607 can also be explained by bio-mechanism from the zoo/phytoplankton in the Antarctic Ocean.  
 608



## 609 4 Discussion

610

611 We observed the presence of many chemical constituents with significant measurable  
612 concentrations especially during aerosol formation events:

613

614 (1) IA daily maximum concentrations showed low variability, independent of NPF occurrence. Its  
615 concentration - only rarely over  $10^6$  molecules  $\text{cm}^{-3}$  - was unlikely sufficient to initiate NPF alone  
616 as compared with previously reported concentrations of  $10^8$  molecules  $\text{cm}^{-3}$  and ( $>$ )  $8 \times 10^6$  mol-  
617 ecules  $\text{cm}^{-3}$  for Arctic and coastal studies (Sipilä et al., 2016; Baccharini et al., 2020; Beck et al.,  
618 2021). Iodine emissions have been connected to sea-ice conditions and photolabile iodine com-  
619 pounds that are rapidly oxidized (Saiz-Lopez et al., 2015; He et al., 2021). The colder Weddell  
620 Sea provides high potential for the emission of iodine and thus iodic acid in the atmosphere. How-  
621 ever, the temporality is a key factor governing such emissions. In fact, many studies have shown  
622 that IA concentration in polar-marine environment are usually peaking in early spring - already  
623 before the sea ice melt onset - (Saiz-Lopez et al., 2007) and during the refreezing transition period  
624 (Baccharini et al., 2020), strongly linking IA emissions with the sea ice state. This alternatively  
625 implies that our measurement period was then not optimum to catch iodine-related vapors maxima  
626 since the neighboring sea ice – key surface emission medium - was already melted around the  
627 Seymour Island. We do not exclude the possibility that IA would contribute to NPF in other sea-  
628 sons, but our data set show that during the austral summer period, from mid-January until the end  
629 of February, IA is likely not substantial.

630

631 The high concentrations of SA suggest that SA plays a key role in nucleation similarly to earlier  
632 observations from Antarctica (Jokinen et al., 2018) and from the Arctic (Beck et al., 2021). The  
633 significantly higher nucleation rates, with similar concentrations of SA, contrast with those studies  
634 that concluded that SA – ammonia ion induced nucleation was the primary pathway to new particle  
635 formation. The reported new particle formation rates were systematically below  $1 \text{ cm}^{-3} \text{ s}^{-1}$ , typi-  
636 cally ca. 2 orders of magnitude lower than observed in our study despite comparable air tempera-  
637 tures.

638

639 Taking the case study of 16 February as an example, we found sulfuric acid – ammonia complexes  
640 with high abundancy from the analysis of natural ion chemical composition (Fig. 7). That would  
641 indicate that at least negative ion-induced nucleation pathway proceeds with sequential addition  
642 of SA and  $\text{NH}_3$ . However, the total nucleation rate was  $3.6 \text{ cm}^{-3} \text{ s}^{-1}$  on this day, much larger than  
643 the negative ion induced nucleation rate of  $0.06 \text{ cm}^{-3} \text{ s}^{-1}$ . This would lead to the conclusion that a  
644 neutral nucleation mechanism could dominate the process in our case.

645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681

Outside the case study, nucleation rates are still remarkably high, often exceeding the ion-induced nucleation rate limit set by ion-production rate of ca.  $2 \text{ ion pairs cm}^{-3} \text{ s}^{-1}$  by galactic cosmic radiation. While the exact nature of neutral mechanism is not directly detected, the parameterization experiments performed in the CERN CLOUD-chamber (Dunne et al., 2016; Kürten, 2019) indicate that very high concentrations of ammonia would be needed to explain the observed nucleation rates – above ppb levels, under assumption that the ternary SA-NH<sub>3</sub>-H<sub>2</sub>O mechanism was solely responsible for initial nucleation.

- (2) Another possibility is that dimethyl amine – as detected in negative ion spectra (cf. Fig. 7) - efficiently nucleates with sulfuric acid and could be primarily responsible for neutral new particle SA-H<sub>2</sub>O-DMA formation pathway, as discussed in several studies (Kürten et al., 2014; Brean et al., 2021). Unfortunately, the deployed instrumentation does not allow the determination of DMA concentrations to assess the exact contribution of DMA on nucleation. Yet, our observation shows significantly higher nucleation rates than the reported values of both studies and, considering the range of measured SA concentrations, this conjecture is not likely to happen alone but rather as a possible synergetic effect of both amines and ammonia in the nucleation with sulfuric acid and water at the encountered temperatures.
- (3) In addition to the high quantity of SA and MSA measured in CI mode, MSA was observed in ion cluster in form of CH<sub>3</sub>SO<sub>3</sub><sup>-</sup> and MSA clusters with bisulfate ion but not in any larger nucleating cluster. Its contribution to a possible ion-induced nucleation mechanism is therefore unlikely in our case. However, we still cannot exclude the possibility that MSA would be weakly bounded to ion clusters and then evaporated inside the mass spectrometer vacuum chambers and lost in semi-energetic collisions within the electric fields used to focus ions inside the system. Additionally, mesylate (i.e., the MSA fragments - CH<sub>3</sub>SO<sub>3</sub><sup>-</sup>) were observed in significant fraction in the chemical speciation of the aerosol phase, which indicates its involvement in particle growth.
- (4) The missing link is the (high) concentration of ammonia and amines required to follow the proposed SA-(NH<sub>3</sub>/DMA)-H<sub>2</sub>O nucleation scheme, likely on the order of ppb for ammonia and ppt levels for DMA. Without direct measurement of neither NH<sub>3</sub> nor DMA, we can only speculate that our observations point toward a strong source, possibly land-based (e.g., from high penguin population during summertime) or marine-based (e.g., emissions from plankton's biological activity in the (melting) ocean). In fact, predominant wind possibly passing over a penguin colony at shore was seen from the wind roses proposed in Figure S6 for event 3, 4, 7, 9, and 10 and confirmed by the air mass trajectory analysis (Fig. 9). Yet, for our case study (event #12) - proving significant contribution of ammonia / DMA to detected in ion clusters -, nor the wind analysis nor

682 the backward trajectory analysis point toward emission sourcing from the bird colony. We do not  
683 exclude the possibility of additional influence from the local fauna, further away from the island,  
684 however this nitrogen-containing species could also be marine-sourced, especially by emission  
685 from the metabolic activity of planktons and accumulation of precursors (e.g., proteins) in the  
686 melting sea ice (Dall'Osto et al., 2017). Ammonia and amine concentrations are key parameters  
687 that need to be assessed to fully understand the nucleation mechanism that our analysis suggests.

688  
689 (5) MSA, together with DMA, NH<sub>3</sub> and SA, could potentially nucleate via a neutral, yet unidentified  
690 mechanism. However, this potential mechanism does not manifest itself in chemical composition  
691 of negative ion clusters, and potentially would only occur as neutral. It is also important to consider  
692 that our measurement relies on clusters that are stable enough to be analyzed in the very low-  
693 pressure chamber of the TOF, and thus might not be detectable with all species or molecular cluster  
694 possibly present in the real atmosphere. Further investigations of nucleation potential of MSA  
695 including the aforementioned compounds as mixture are required before MSA can be attributed  
696 to be responsible for our observed high nucleation rates that we observe at Marambio.

697  
698 (6) To this remains the question of organics; our data set do not show substantial presence of HOM  
699 (as characterized in forest environments), and possible detected compounds entering in our estimation  
700 – yet unidentified chemically, and which can also represent instrumental chemical background –  
701 could be low-volatility organic compounds originating from marine environment that  
702 may contribute to the growth to some extent (Weller et al., 2015; Kecorius et al., 2019).

703  
704 (7) A possible explanation for our observations of higher-than-expected nucleation and growth rates  
705 is that particle formation and growth could occur primarily at colder temperatures at higher altitudes,  
706 which would enhance both SA-NH<sub>3</sub> and SA-DMA nucleation. Nucleated particles would  
707 then be transported to lower altitudes resulting in apparently higher nucleation and growth rates.  
708 This in turn suggest more regional emission sources rather than local emissions of SA, NH<sub>3</sub> or  
709 DMA as our analysis points toward. However, further investigations on vertical mixing would  
710 have been needed to confirm vertical distribution of nucleation and growth rates.

711  
712

## 713 5 Conclusion

714

715 The Austral summer campaign 2018 at the Marambio Antarctic research station revealed very  
716 active aerosol processes and a frequency of new particle formation events of  $\sim 40\%$ . Out of 35 days  
717 of active measurement, a total of 13 NPF events were identified and characterized. In Marambio,  
718 NPF was promoted by key meteorological factors: high radiation (clear sky conditions), low humidity  
719 conditions (incl. the absence of precipitation), and high temperature – close to zero or above. We  
720 retrieved formation and growth rates for neutral and charged particles measured within 1.5 nm up to  
721 800 nm size range - combining nCNC, NAIS and DMPS instrumentation. As a reference, the  
722 formation rate of 3-nm (neutral) particles ( $J_3$ ) during NPF was on average  $0.6863 \text{ cm}^{-3} \text{ s}^{-1}$  and the  
723 growth rate ( $GR_{4-12 \text{ nm}}$ ) was on average  $4.2 \text{ nm}\cdot\text{h}^{-1}$ . Those values are particularly high in comparison  
724 with the literature, the comparison remains even more surprising considering the most active NPF  
725 day where  $J_{1.5}$  reached  $19 \text{ cm}^{-3}\text{s}^{-1}$  and  $GR_{4-12 \text{ nm}}$  was  $10.9 \text{ nm}\cdot\text{h}^{-1}$ . The chemical analysis of gas-phase  
726 aerosol precursors with CI-APi-TOF showed high concentrations of SA and MSA - on average  $5.17$   
727  $\times 10^5 \text{ molecules cm}^{-3}$  and  $1.18 \times 10^6 \text{ molecules cm}^{-3}$ , respectively – but quite low concentration of IA  
728 (on average  $2.06 \times 10^5 \text{ molecules cm}^{-3}$ ). All chemical species reached their maxima during mid-day  
729 (e.g., with maximum concentration of  $2.56 \times 10^7$ ,  $2.32 \times 10^7 \text{ molecules}\cdot\text{cm}^{-3}$ , respectively for SA and  
730 MSA during aerosol formation events). The ubiquitously high SA concentration (peaking at almost  
731  $10^7 \text{ molecules cm}^{-3}$  during many events) and the high abundance of SA-associated ion clusters makes  
732 no doubt on the involvement of SA on NPF. The speciation of negative ion spectra revealed clusters  
733 of dimethylamine – sulfuric acid – bisulfate-ions, as well as various clusters of sulfuric acid -  
734 ammonia- bisulfate-ion - consistent with previously reported studies. However, the systematically  
735 high formation / growth rates cannot be quantitatively explained by SA-(DMA/  $\text{NH}_3$ )- $\text{H}_2\text{O}$  NPF  
736 pathways without direct measurement of DMA and  $\text{NH}_3$  gas phase concentrations. Additionally,  
737 consistently high concentrations of MSA during daytime suggests a possible role of MSA in the  
738 growth, along with SA and ammonia - as confirmed by the bulk aerosol chemical composition.  
739 Herein, the mechanism of forming secondary aerosol in the peninsula involves a combination of SA,  
740 DMA,  $\text{NH}_3$  and MSA (and only relatively low contribution of IA, at least at this time of the year).  
741 Better understanding the synergetic effect of DMA/ $\text{NH}_3$  at these temperatures and environments  
742 require more field measurements and simulated experiments, specifically including direct  
743 measurement of DMA and  $\text{NH}_3$  concentrations.

744

745

746 **Data Availability**

747

748 *The key data set for both aerosol characterization and gas-phase composition are publically*  
749 *available on the Zenodo data repository platform:*

750

751 Quéléver; L. L. J.; Dada, L.; Asmi, E.; Lampilahti, J.; Chan, T.; Ferrara, J.; Copes, G.; Pérez-Fogwill,  
752 G.; Barreira, L.; Aurela, M.; Worsnop, D.; Jokinen, T.; Sipilä, M.; (2022). Investigation of New  
753 Particle Formation mechanisms and aerosol processes at the Marambio Station, Antarctic Peninsula  
754 [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.6560413>

755

756 *For further information please contact Lauriane L. J. Quéléver (lauriane.quelever@helsinki.fi).*

757

758 **Acknowledgments**

759

760 This work was supported by the European Research Council under the following programs: ERC  
761 (GASPARCON – grant n° 714621 & COALA – grant n° 638703) and EMME-CARE (grant  
762 n° 856612). We acknowledge the support from the Academy of Finland (projects n° 296628 and n°  
763 328290), the Center of Excellence program (project n° 307331, 334514, 3282290), and Finnish  
764 Academy ACFA (project n° 335845 and ACCC Flagship funding (337552). We thank the generous  
765 help from the Servicio Meteorológico Nacional - SMN, the Fuerza Aerea Argentina - FAA, and the  
766 FINNish Antarctic Research Program – FINNARP. We are thankful for the technical personnel of  
767 the Marambio base for their support in cooperation, logistics and measurements. We thank Toftools  
768 for providing mass spectrometry analysis tool package for MATLAB.

769

770 **Author's Contribution**

771

772 **EA** and **MS** supervised the project. **EA** initiated the measurement container deployment and is  
773 responsible for the large size aerosol and auxiliary measurements data from the measurement  
774 container. **LLJQ** co-organized the project, prepared, set up and performed the measurements,  
775 analyzed the data, and wrote the manuscript. **LLJQ** operated the instrumentation with the help of  
776 **EA**, **JEF** and **GEC**. **EA**, **MS** and **TJ** supervised the analysis and contributed to the scientific  
777 discussion in collaboration with **DW**. **LLJQ** analyzed the gas-phase measurements and produced  
778 most figures. **LD** processed, analyzed, and discussed the aerosol phase data, prepared by **TC**, **JL**,  
779 **GPF** and **EA**. **LB** and **MA** analyzed the chemical composition of bulk aerosols. **EA** performed the

780 backward trajectory analysis and its relevant figures. All co-authors contributed to the scientific  
781 discussion, commenting and the revision of the manuscript.

782

### 783 **Competing interests**

784

785 *The authors claim no competing interests.*

786

### 787 **Supplementary Information (cf. Separate file)**

788

### 789 **References**

790

791 Almeida, J., Schobesberger, S., Kürten, A., Ortega, I. K., Kupiainen-Määttä, O., Praplan, A. P., Adamov, A.,  
792 Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne,  
793 E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M.,  
794 Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T.,  
795 Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppä, J., Loukonen, V., Makhmutov, V.,  
796 Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F., Riipinen, I.,  
797 Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld,  
798 J. H., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara,  
799 P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer,  
800 D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R.,  
801 Vehkamäki, H., and Kirkby, J.: Molecular understanding of sulphuric acid–amine particle nucleation in the  
802 atmosphere, *Nature*, 502, 359-363, 10.1038/nature12663, 2013.

803 Atkinson, H. M., Huang, R. J., Chance, R., Roscoe, H. K., Hughes, C., Davison, B., Schönhardt, A., Mahajan,  
804 A. S., Saiz-Lopez, A., Hoffmann, T., and Liss, P. S.: Iodine emissions from the sea ice of the Weddell Sea,  
805 *Atmos. Chem. Phys.*, 12, 11229-11244, 10.5194/acp-12-11229-2012, 2012.

806 Asmi, E., Frey, A., Virkkula, A., Ehn, M., Manninen, H. E., Timonen, H., Tolonen-Kivimäki, O., Aurela, M.,  
807 Hillamo, R., and Kulmala, M.: Hygroscopicity and chemical composition of Antarctic sub-micrometre aerosol  
808 particles and observations of new particle formation, *Atmos. Chem. Phys.*, 10, 4253-4271, 10.5194/acp-10-  
809 4253-2010, 2010.

810 Asmi, E., Neitola, K., Teinilä, K., Rodriguez, E., Virkkula, A., Backman, J., Bloss, M., Jokela, J., Lihavainen,  
811 H., de Leeuw, G., Paatero, J., Aaltonen, V., Mei, M., Gambarte, G., Copes, G., Albertini, M., Fogwill, G. P.,  
812 Ferrara, J., Barlasina, M. E., and Sánchez, R.: Primary sources control the variability of aerosol optical  
813 properties in the Antarctic Peninsula, *Tellus B: Chemical and Physical Meteorology*, 70, 1-16,  
814 10.1080/16000889.2017.1414571, 2018.

815 Baccarini, A., Karlsson, L., Dommen, J., Duplessis, P., Vüllers, J., Brooks, I. M., Saiz-Lopez, A., Salter, M.,  
816 Tjernström, M., Baltensperger, U., Zieger, P., and Schmale, J.: Frequent new particle formation over the high  
817 Arctic pack ice by enhanced iodine emissions, *Nature Communications*, 11, 4924, 10.1038/s41467-020-  
818 18551-0, 2020.

819 Barnes, I., Hjorth, J., and Mihalopoulos, N.: Dimethyl Sulfide and Dimethyl Sulfoxide and Their  
820 Oxidation in the Atmosphere, *Chemical Reviews*, 106, 940-975, 10.1021/acs.accounts.cr020529+,  
821 2006.

822

823 Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä,  
824 O., Wimmer, D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, X.-C., Kontkanen, J., Wolf,  
825 K. K. E., Cappelletti, D., Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling,  
826 A., Nøjgaard, J. K., Krejci, R., Karlsson, L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur,  
827 R. C., Leino, K., Kangasluoma, J., Duplissy, E.-M., Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A.,  
828 Petäjä, T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V.-M., and Sipilä, M.: Differing  
829 Mechanisms of New Particle Formation at Two Arctic Sites, *Geophysical Research Letters*, 48,  
830 e2020GL091334, 10.1029/2020GL091334, 2021.

831 Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crouse, J. D., Wennberg,  
832 P. O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A.,  
833 Donahue, N., Kjaergaard, H. G., and Ehn, M.: Highly Oxygenated Organic Molecules (HOM) from Gas-Phase  
834 Autoxidation Involving Peroxy Radicals: A Key Contributor to Atmospheric Aerosol, *Chemical Reviews*, 119,  
835 3472-3509, 10.1021/acs.chemrev.8b00395, 2019.

836 Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov,  
837 A., Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten,  
838 A., Manninen, H. E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius,  
839 J., Worsnop, D. R., Kulmala, M., Dommen, J., and Baltensperger, U.: New particle formation in the free  
840 troposphere: A question of chemistry and timing, *Science*, 352, 1109-1112, 10.1126/science.aad5456, 2016.

841 Bork, N., Elm, J., Olenius, T., and Vehkamäki, H.: Methane sulfonic acid-enhanced formation of molecular  
842 clusters of sulfuric acid and dimethyl amine, *Atmos. Chem. Phys.*, 14, 12023-12030, 10.5194/acp-14-12023-  
843 2014, 2014.

844 Brean, J., Dall'Osto, M., Simó, R., Shi, Z., Beddows, D. C. S., and Harrison, R. M.: Open ocean and coastal  
845 new particle formation from sulfuric acid and amines around the Antarctic Peninsula, *Nature Geoscience*, 14,  
846 383-388, 10.1038/s41561-021-00751-y, 2021.

847 Brewer, G. P.: Minor elements in seawater, *Chemical Oceanography*, 1, 415-496, 1975.

848 Chan, T., Cai, R., Ahonen, L. R., Liu, Y., Zhou, Y., Vanhanen, J., Dada, L., Chao, Y., Liu, Y., Wang, L.,  
849 Kulmala, M., and Kangasluoma, J.: Assessment of particle size magnifier inversion methods to obtain the  
850 particle size distribution from atmospheric measurements, *Atmos. Meas. Tech.*, 13, 4885-4898, 10.5194/amt-  
851 13-4885-2020, 2020.

852 Collins, D. B., Burkart, J., Chang, R. Y. W., Lizotte, M., Boivin-Rioux, A., Blais, M., Mungall, E. L., Boyer,  
853 M., Irish, V. E., Massé, G., Kunkel, D., Tremblay, J. É., Papakyriakou, T., Bertram, A. K., Bozem, H.,  
854 Gosselin, M., Lévasseur, M., and Abbatt, J. P. D.: Frequent ultrafine particle formation and growth in Canadian  
855 Arctic marine and coastal environments, *Atmos. Chem. Phys.*, 17, 13119-13138, 10.5194/acp-17-13119-2017,  
856 2017.

857 Dada, L., Chellapermal, R., Buenrostro Mazon, S., Paasonen, P., Lampilahti, J., Manninen, H. E., Junninen,  
858 H., Petäjä, T., Kerminen, V. M., and Kulmala, M.: Refined classification and characterization of atmospheric  
859 new-particle formation events using air ions, *Atmos. Chem. Phys.*, 18, 17883-17893, 10.5194/acp-18-17883-  
860 2018, 2018.

861 Dada, L., Paasonen, P., Nieminen, T., Buenrostro Mazon, S., Kontkanen, J., Peräkylä, O., Lehtipalo, K.,  
862 Hussein, T., Petäjä, T., Kerminen, V. M., Bäck, J., and Kulmala, M.: Long-term analysis of clear-sky new  
863 particle formation events and nonevents in Hyytiälä, *Atmos. Chem. Phys.*, 17, 6227-6241, 10.5194/acp-17-  
864 6227-2017, 2017.

865

- 866 Dada, L., Lehtipalo, K., Kontkanen, J., Nieminen, T., Baalbaki, R., Ahonen, L., Duplissy, J., Yan, C., Chu, B.,  
867 Petäjä, T., Lehtinen, K., Kerminen, V.-M., Kulmala, M., and Kangasluoma, J.: Formation and growth of sub-  
868 3-nm aerosol particles in experimental chambers, *Nature Protocols*, 15, 1013-1040, 10.1038/s41596-019-  
869 0274-z, 2020.
- 870 Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P., and Lehtinen, K. E.: Formation  
871 and growth of fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II,  
872 Hyytiälä, Finland, *Boreal Environment Research*, 10, 323, 2005.
- 873 Dall'Osto, M., Beddows, D. C. S., Tunved, P., Harrison, R. M., Lupi, A., Vitale, V., Becagli, S., Traversi, R.,  
874 Park, K. T., Yoon, Y. J., Massling, A., Skov, H., Lange, R., Strom, J., and Krejci, R.: Simultaneous  
875 measurements of aerosol size distributions at three sites in the European high Arctic, *Atmos. Chem. Phys.*, 19,  
876 7377-7395, 10.5194/acp-19-7377-2019, 2019.
- 877 Dall'Osto, M., Geels, C., Beddows, D. C. S., Boertmann, D., Lange, R., Nøjgaard, J. K., Harrison, R. M.,  
878 Simo, R., Skov, H., and Massling, A.: Regions of open water and melting sea ice drive new particle formation  
879 in North East Greenland, *Scientific Reports*, 8, 6109, 10.1038/s41598-018-24426-8, 2018.
- 880 Dall'Osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Ström, J., Hansson, H. C., Yoon, Y. J., Park, K.-T.,  
881 Becagli, S., Udusti, R., Onasch, T., O'Dowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to  
882 atmospheric new particle formation, *Scientific Reports*, 7, 3318, 10.1038/s41598-017-03328-1, 2017.
- 883 Deng, C., Fu, Y., Dada, L., Yan, C., Cai, R., Yang, D., Zhou, Y., Yin, R., Lu, Y., Li, X., Qiao, X., Fan, X.,  
884 Nie, W., Kontkanen, J., Kangasluoma, J., Chu, B., Ding, A., Kerminen, V.-M., Paasonen, P., Worsnop, D. R.,  
885 Bianchi, F., Liu, Y., Zheng, J., Wang, L., Kulmala, M., and Jiang, J.: Seasonal Characteristics of New Particle  
886 Formation and Growth in Urban Beijing, *Environmental Science & Technology*, 54, 8547-8557,  
887 10.1021/acs.est.0c00808, 2020.
- 888 Dunne, E. M., Gordon, H., Kürten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle, K. J.,  
889 Adamov, A., Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., Curtius,  
890 J., Dommen, J., Donahue, N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A.,  
891 Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J., Lehtipalo, K.,  
892 Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A.,  
893 Reddington, C. L., Riccobono, F., Richards, N. A., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger,  
894 S., Sengupta, K., Simon, M., Sipilä, M., Smith, J. N., Stozkhov, Y., Tomé, A., Tröstl, J., Wagner, P. E.,  
895 Wimmer, D., Winkler, P. M., Worsnop, D. R., and Carslaw, K. S.: Global atmospheric particle formation from  
896 CERN CLOUD measurements, *Science*, 354, 1119-1124, 10.1126/science.aaf2649, 2016.
- 897 Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann,  
898 R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S.,  
899 Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G.,  
900 Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop,  
901 D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506,  
902 476-479, 10.1038/nature13032, 2014.
- 903 Fetterer, F., K. Knowles, W. N. Meier, M. Savoie, and A. K. Windnagel. 2017, updated daily. *Sea*  
904 *Ice Index, Version 3*. [ <ftp://sidads.colorado.edu/DATASETS/NOAA/G02135/south/monthly/>].  
905 Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center, [10.7265/N5K072F8](https://doi.org/10.7265/N5K072F8), 2021  
906 [last access: August 5, 2021].
- 907  
908 Fiebig, M., Hirdman, D., Lunder, C. R., Ogren, J. A., Solberg, S., Stohl, A., and Thompson, R. L.: Annual  
909 cycle of Antarctic baseline aerosol: controlled by photooxidation-limited aerosol formation, *Atmos. Chem.*  
910 *Phys.*, 14, 3083-3093, 10.5194/acp-14-3083-2014, 2014.



- 911 Fiedler, V., Dal Maso, M., Boy, M., Aufmhoff, H., Hoffmann, J., Schuck, T., Birmili, W., Hanke, M., Uecker,  
912 J., Arnold, F., and Kulmala, M.: The contribution of sulphuric acid to atmospheric particle formation and  
913 growth: a comparison between boundary layers in Northern and Central Europe, *Atmos. Chem. Phys.*, 5, 1773-  
914 1785, 10.5194/acp-5-1773-2005, 2005.
- 915 He, X.-C., Tham, Y. J., Dada, L., Wang, M., Finkenzeller, H., Stolzenburg, D., Iyer, S., Simon, M., Kürten,  
916 A., Shen, J., Rörup, B., Rissanen, M., Schobesberger, S., Baalbaki, R., Wang, D. S., Koenig, T. K., Jokinen,  
917 T., Sarnela, N., Beck, L. J., Almeida, J., Amanatidis, S., Amorim, A., Ataei, F., Baccharini, A., Bertozzi, B.,  
918 Bianchi, F., Brilke, S., Caudillo, L., Chen, D., Chiu, R., Chu, B., Dias, A., Ding, A., Dommen, J., Duplissy, J.,  
919 El Haddad, I., Gonzalez Carracedo, L., Granzin, M., Hansel, A., Heinritzi, M., Hofbauer, V., Junninen, H.,  
920 Kangasluoma, J., Kempainen, D., Kim, C., Kong, W., Krechmer, J. E., Kvashin, A., Laitinen, T.,  
921 Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Li, Z., Makhmutov, V., Manninen, H. E., Marie,  
922 G., Marten, R., Mathot, S., Mauldin, R. L., Mentler, B., Möhler, O., Müller, T., Nie, W., Onnela, A., Petäjä,  
923 T., Pfeifer, J., Philippov, M., Ranjithkumar, A., Saiz-Lopez, A., Salma, I., Scholz, W., Schuchmann, S.,  
924 Schulze, B., Steiner, G., Stozhkov, Y., Tauber, C., Tomé, A., Thakur, R. C., Väisänen, O., Vazquez-Pufleau,  
925 M., Wagner, A. C., Wang, Y., Weber, S. K., Winkler, P. M., Wu, Y., Xiao, M., Yan, C., Ye, Q., Ylisirniö, A.,  
926 Zauner-Wieczorek, M., Zha, Q., Zhou, P., Flagan, R. C., Curtius, J., Baltensperger, U., Kulmala, M.,  
927 Kerminen, V.-M., Kurtén, T., Donahue, N. M., Volkamer, R., Kirkby, J., Worsnop, D. R., and Sipilä, M.: Role  
928 of iodine oxoacids in atmospheric aerosol nucleation, *Science*, 371, 589, 10.1126/science.abe0298, 2021.
- 929 Herenz, P., Wex, H., Mangold, A., Laffineur, Q., Gorodetskaya, I. V., Fleming, Z. L., Panagi, M., and  
930 Stratmann, F.: CCN measurements at the Princess Elisabeth Antarctica research station during three austral  
931 summers, *Atmos. Chem. Phys.*, 19, 275-294, 10.5194/acp-19-275-2019, 2019.
- 932 Hodshire, A. L., Campuzano-Jost, P., Kodros, J. K., Croft, B., Nault, B. A., Schroder, J. C., Jimenez, J. L., and  
933 Pierce, J. R.: The potential role of methanesulfonic acid (MSA) in aerosol formation and growth and the  
934 associated radiative forcings, *Atmos. Chem. Phys.*, 19, 3137-3160, 10.5194/acp-19-3137-2019, 2019.
- 935 IPCC, Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex,  
936 V., and Midgley, P. M. (Eds.): *Climate Change 2013: The Physical Science Basis. Contribution of Working  
937 Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge  
938 University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.,  
939 10.1017/CBO9781107415324, 2013.
- 940 Jokinen, T., Sipilä, M., Junninen, H., Ehn, M., Lönn, G., Hakala, J., Petäjä, T., Mauldin Iii, R. L., Kulmala,  
941 M., and Worsnop, D. R.: Atmospheric sulphuric acid and neutral cluster measurements using CI-APi-TOF,  
942 *Atmos. Chem. Phys.*, 12, 4117-4125, 10.5194/acp-12-4117-2012, 2012.
- 943 Jokinen, T., Sipilä, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E. M., Junninen, H., Kangasluoma, J.,  
944 Manninen, H. E., Petäjä, T., Kulmala, M., Worsnop, D. R., Kirkby, J., Virkkula, A., and Kerminen, V. M.:  
945 Ion-induced sulfuric acid–ammonia nucleation drives particle formation in coastal Antarctica, *Science  
946 Advances*, 4, eaat9744, 10.1126/sciadv.aat9744, 2018.
- 947 Järvinen, E., Virkkula, A., Nieminen, T., Aalto, P. P., Asmi, E., Lanconelli, C., Busetto, M., Lupi, A.,  
948 Schioppo, R., Vitale, V., Mazzola, M., Petäjä, T., Kerminen, V. M., and Kulmala, M.: Seasonal cycle and  
949 modal structure of particle number size distribution at Dome C, Antarctica, *Atmos. Chem. Phys.*, 13, 7473-  
950 7487, 10.5194/acp-13-7473-2013, 2013.
- 951

- 952 Kangasluoma, J., Franchin, A., Duplissy, J., Ahonen, L., Korhonen, F., Attoui, M., Mikkilä, J., Lehtipalo, K.,  
953 Vanhanen, J., Kulmala, M., and Petäjä, T.: Operation of the Airmodus A11 nano Condensation Nucleus  
954 Counter at various inlet pressures and various operation temperatures, and design of a new inlet system, *Atmos.*  
955 *Meas. Tech.*, 9, 2977-2988, 10.5194/amt-9-2977-2016, 2016.
- 956 Kecorius, S., Vogl, T., Paasonen, P., Lampilahti, J., Rothenberg, D., Wex, H., Zeppenfeld, S., van Pinxteren,  
957 M., Hartmann, M., Henning, S., Gong, X., Welti, A., Kulmala, M., Stratmann, F., Herrmann, H., and  
958 Wiedensohler, A.: New particle formation and its effect on cloud condensation nuclei abundance in the  
959 summer Arctic: a case study in the Fram Strait and Barents Sea, *Atmos. Chem. Phys.*, 19, 14339-14364,  
960 10.5194/acp-19-14339-2019, 2019.
- 961 Kerminen, V.-M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi, F.: Atmospheric new particle  
962 formation and growth: review of field observations, *Environmental Research Letters*, 13, 103003,  
963 10.1088/1748-9326/aadf3c, 2018.
- 964 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L.,  
965 Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer,  
966 D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R.  
967 C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A.,  
968 Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S.,  
969 Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y.,  
970 Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E.,  
971 Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of  
972 sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429-433,  
973 10.1038/nature10343, 2011.
- 974 Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan,  
975 C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer,  
976 A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C.,  
977 Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim,  
978 J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A.,  
979 Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen,  
980 M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov,  
981 Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E.,  
982 Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R.,  
983 Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic  
984 particles, *Nature*, 533, 521-526, 10.1038/nature17953, 2016.
- 985 Kontkanen, J., Järvinen, E., Manninen, H. E., Lehtipalo, K., Kangasluoma, J., Decesari, S., Gobbi, G. P.,  
986 Laaksonen, A., Petäjä, T., and Kulmala, M.: High concentrations of sub-3nm clusters and frequent new particle  
987 formation observed in the Po Valley, Italy, during the PEGASOS 2012 campaign, *Atmos. Chem. Phys.*, 16,  
988 1919-1935, 10.5194/acp-16-1919-2016, 2016.
- 989 Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä,  
990 M., Schobesberger, S., and Rantala, P.: Direct observations of atmospheric aerosol nucleation, *Science*, 339,  
991 943-946, 10.1126/science.1227385, 2013.
- 992 Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P.,  
993 Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V.-M.: Measurement  
994 of the nucleation of atmospheric aerosol particles, *Nature Protocols*, 7, 1651-1667, 10.1038/nprot.2012.091,  
995 2012.
- 996

- 997 Kulmala, M., Stolzenburg, D., Dada, L., Cai, R., Kontkanen, J., Yan, C., Kangasluoma, J., Ahonen, L. R.,  
 998 Gonzalez-Carracedo, L., Sulo, J., Tuovinen, S., Deng, C., Li, Y., Lehtipalo, K., Lehtinen, K. E. J., Petäjä, T.,  
 999 Winkler, P. M., Jiang, J., and Kerminen, V.-M.: Towards a concentration closure of sub-6 nm aerosol particles  
 1000 and sub-3 nm atmospheric clusters, *Journal of Aerosol Science*, 159, 105878, 10.1016/j.jaerosci.2021.105878,  
 1001 2022.
- 1002 Kyrö, E. M., Kerminen, V. M., Virkkula, A., Dal Maso, M., Parshintsev, J., Ruiz-Jimenez, J., Forsström, L.,  
 1003 Manninen, H. E., Riekkola, M. L., Heinonen, P., and Kulmala, M.: Antarctic new particle formation from  
 1004 continental biogenic precursors, *Atmos. Chem. Phys.*, 13, 3527-3546, 10.5194/acp-13-3527-2013, 2013.
- 1005 Kürten, A.: New particle formation from sulfuric acid and ammonia: nucleation and growth model based on  
 1006 thermodynamics derived from CLOUD measurements for a wide range of conditions, *Atmos. Chem. Phys.*,  
 1007 19, 5033-5050, 10.5194/acp-19-5033-2019, 2019.
- 1008 Kürten, A., Jokinen, T., Simon, M., Sipilä, M., Sarnela, N., Junninen, H., Adamov, A., Almeida, J., Amorim,  
 1009 A., Bianchi, F., Breitenlechner, M., Dommen, J., Donahue, N. M., Duplissy, J., Ehrhart, S., Flagan, R. C.,  
 1010 Franchin, A., Hakala, J., Hansel, A., Heinritzi, M., Hutterli, M., Kangasluoma, J., Kirkby, J., Laaksonen, A.,  
 1011 Lehtipalo, K., Leiminger, M., Makhmutov, V., Mathot, S., Onnela, A., Petäjä, T., Praplan, A. P., Riccobono,  
 1012 F., Rissanen, M. P., Rondo, L., Schobesberger, S., Seinfeld, J. H., Steiner, G., Tomé, A., Tröstl, J., Winkler,  
 1013 P. M., Williamson, C., Wimmer, D., Ye, P., Baltensperger, U., Carslaw, K. S., Kulmala, M., Worsnop, D. R.,  
 1014 and Curtius, J.: Neutral molecular cluster formation of sulfuric acid–dimethylamine observed in real time under  
 1015 atmospheric conditions, *Proceedings of the National Academy of Sciences*, 111, 15019,  
 1016 10.1073/pnas.1404853111, 2014.
- 1017 Lachlan-Cope, T., Beddows, D. C. S., Brough, N., Jones, A. E., Harrison, R. M., Lupi, A., Yoon, Y. J.,  
 1018 Virkkula, A., and Dall'Osto, M.: On the annual variability of Antarctic aerosol size distributions at Halley  
 1019 Research Station, *Atmos. Chem. Phys.*, 20, 4461-4476, 10.5194/acp-20-4461-2020, 2020.
- 1020 Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman,  
 1021 E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide  
 1022 concentrations and emission fluxes in the global ocean, *Global Biogeochemical Cycles*, 25,  
 1023 10.1029/2010GB003850, 2011.
- 1024 Lehtipalo, K., Leppä, J., Kontkanen, J., Kangasluoma, J., Franchin, A., Wimmer, D., Schobesberger, S.,  
 1025 Junninen, H., Petaja, T., and Sipilä, M.: Methods for determining particle size distribution and growth rates  
 1026 between 1 and 3 nm using the Particle Size Magnifier, *Boreal Environment Research*, 2014.
- 1027 Lehtipalo, K., Ahonen, L. R., Baalbaki, R., Sulo, J., Chan, T., Laurila, T., Dada, L., Duplissy, J., Miettinen,  
 1028 E., Vanhanen, J., Kangasluoma, J., Kulmala, M., Petäjä, T., and Jokinen, T.: The standard operating procedure  
 1029 for Airmodus Particle Size Magnifier and nano-Condensation Nucleus Counter, *Journal of Aerosol Science*,  
 1030 159, 105896, 10.1016/j.jaerosci.2021.105896, 2022.
- 1031 Loo, B. W. and Cork, C. P.: Development of High Efficiency Virtual Impactors, *Aerosol Science and  
 1032 Technology*, 9, 167-176, 10.1080/02786828808959205, 1988.
- 1033 Manninen, H. E., Mirme, S., Mirme, A., Petäjä, T., and Kulmala, M.: How to reliably detect molecular clusters  
 1034 and nucleation mode particles with Neutral cluster and Air Ion Spectrometer (NAIS), *Atmos. Meas. Tech.*, 9,  
 1035 3577-3605, 10.5194/amt-9-3577-2016, 2016.
- 1036 Mardyukov, A. and Schreiner, P. R.: Atmospherically Relevant Radicals Derived from the Oxidation of  
 1037 Dimethyl Sulfide, *Accounts of Chemical Research*, 51, 475-483, 10.1021/acs.accounts.7b00536, 2018.  
 1038
- 1039 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on  
 1040 global CCN, *Atmos. Chem. Phys.*, 9, 8601-8616, 10.5194/acp-9-8601-2009, 2009.

- 1041 Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS – a spectrometer for the  
1042 measurement of cluster ion and nanometer aerosol size distributions, *Atmos. Meas. Tech.*, 6, 1061-1071,  
1043 10.5194/amt-6-1061-2013, 2013.
- 1044 Notz, D.: The future of ice sheets and sea ice: Between reversible retreat and unstoppable loss, *Proceedings of*  
1045 *the National Academy of Sciences*, 106, 20590-20595, 10.1073/pnas.0902356106, 2009.
- 1046 Saiz-Lopez, A. and von Glasow, R.: Reactive halogen chemistry in the troposphere, *Chemical Society*  
1047 *Reviews*, 41, 6448-6472, 10.1039/C2CS35208G, 2012.
- 1048 Saiz-Lopez, A., Blaszczak-Boxe, C. S., and Carpenter, L. J.: A mechanism for biologically induced iodine  
1049 emissions from sea ice, *Atmos. Chem. Phys.*, 15, 9731-9746, 10.5194/acp-15-9731-2015, 2015.
- 1050 Saiz-Lopez, A., Mahajan Anoop, S., Salmon Rhian, A., Bauguitte Stephane, J. B., Jones Anna, E., Roscoe  
1051 Howard, K., and Plane John, M. C.: Boundary Layer Halogens in Coastal Antarctica, *Science*, 317, 348-351,  
1052 10.1126/science.1141408, 2007.
- 1053 Schmale, J., Baccarini, A., Thurnherr, I., Henning, S., Efraim, A., Regayre, L., Bolas, C.,  
1054 Hartmann, M., Welti, A., Lehtipalo, K., Aemisegger, F., Tatzelt, C., Landwehr, S., Modini, R. L.,  
1055 Tummon, F., Johnson, J. S., Harris, N., Schnaiter, M., Toffoli, A., Derkani, M., Bukowiecki, N.,  
1056 Stratmann, F., Dommen, J., Baltensperger, U., Wernli, H., Rosenfeld, D., Gysel-Ber, M., and  
1057 Carslaw, K. S.: Overview of the Antarctic Circumnavigation Expedition: Study of Preindustrial-like  
1058 Aerosols and Their Climate Effects (ACE-SPACE), *Bulletin of the American Meteorological*  
1059 *Society*, 100(11), 2260-2283, 10.1175/BAMS-D-18-0187.1, 2019.
- 1060
- 1061 Schobesberger, S., Junninen, H., Bianchi, F., Lönn, G., Ehn, M., Lehtipalo, K., Dommen, J., Ehrhart, S.,  
1062 Ortega, I., Franchin, A., Nieminen, T., Riccobono, F., Hutterli, M., Duplissy, J., Almeida, J., Amorim, A.,  
1063 Breitenlechner, M., Downard, A., Dunne, E., and Worsnop, D.: Molecular understanding of atmospheric  
1064 particle formation from sulfuric acid and large oxidized organic molecules, *Proceedings of the National*  
1065 *Academy of Sciences of the United States of America*, 10.1073/pnas.1306973110, 2013.
- 1066 Schobesberger, S., Franchin, A., Bianchi, F., Rondo, L., Duplissy, J., Kürten, A., Ortega, I. K., Metzger, A.,  
1067 Schnitzhofer, R., Almeida, J., Amorim, A., Dommen, J., Dunne, E. M., Ehn, M., Gagné, S., Ickes, L., Junninen,  
1068 H., Hansel, A., Kerminen, V. M., Kirkby, J., Kupc, A., Laaksonen, A., Lehtipalo, K., Mathot, S., Onnela, A.,  
1069 Petäjä, T., Riccobono, F., Santos, F. D., Sipilä, M., Tomé, A., Tsagkogeorgas, G., Viisanen, Y., Wagner, P.  
1070 E., Wimmer, D., Curtius, J., Donahue, N. M., Baltensperger, U., Kulmala, M., and Worsnop, D. R.: On the  
1071 composition of ammonia-sulfuric-acid ion clusters during aerosol particle formation, *Atmos. Chem. Phys.*,  
1072 15, 55-78, 10.5194/acp-15-55-2015, 2015.
- 1073 Shaw, G. E.: Considerations on the origin and properties of the Antarctic aerosol, *Reviews of Geophysics*, 17,  
1074 1983-1998, 10.1029/RG017i008p01983, 1979.
- 1075 Shaw, G. E.: Antarctic aerosols: A review, *Reviews of Geophysics*, 26, 89-112, 10.1029/RG026i001p00089,  
1076 1988.
- 1077 Sipilä, M., Berndt, T., Petäjä, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin, R. L.,  
1078 Hyvärinen, A.-P., Lihavainen, H., and Kulmala, M.: The Role of Sulfuric Acid in Atmospheric Nucleation,  
1079 *Science*, 327, 1243, 10.1126/science.1180315, 2010.
- 1080 Sipilä, M., Sarnela, N., Neitola, K., Laitinen, T., Kemppainen, D., Beck, L., Duplissy, E. M., Kuittinen, S.,  
1081 Lehmusjärvi, T., Lampilahti, J., Kerminen, V. M., Lehtipalo, K., Aalto, P. P., Keronen, P., Siivola, E., Rantala,  
1082 P. A., Worsnop, D. R., Kulmala, M., Jokinen, T., and Petäjä, T.: Wintertime sub-arctic new particle formation  
1083 from Kola Peninsula sulphur emissions, *Atmos. Chem. Phys. Discuss.*, 2021, 1-27, 10.5194/acp-2020-1202,  
1084 2021.

- 1085 Sipilä, M., Sarnela, N., Jokinen, T., Henschel, H., Junninen, H., Kontkanen, J., Richters, S., Kangasluoma, J.,  
1086 Franchin, A., Peräkylä, O., Rissanen, M. P., Ehn, M., Vehkamäki, H., Kurten, T., Berndt, T., Petäjä, T.,  
1087 Worsnop, D., Ceburnis, D., Kerminen, V.-M., Kulmala, M., and O'Dowd, C.: Molecular-scale evidence of  
1088 aerosol particle formation via sequential addition of HIO<sub>3</sub>, *Nature*, 537, 532-534, 10.1038/nature19314, 2016.
- 1089 Stuecker, M. F., Bitz, C. M., Armour, K. C., Proistosescu, C., Kang, S. M., Xie, S.-P., Kim, D., McGregor, S.,  
1090 Zhang, W., Zhao, S., Cai, W., Dong, Y., and Jin, F.-F.: Polar amplification dominated by local forcing and  
1091 feedbacks, *Nature Climate Change*, 8, 1076-1081, 10.1038/s41558-018-0339-y, 2018.
- 1092 Vanhanen, J., Mikkilä, J., Lehtipalo, K., Sipilä, M., Manninen, H. E., Siivola, E., Petäjä, T., and Kulmala, M.:  
1093 Particle Size Magnifier for Nano-CN Detection, *Aerosol Science and Technology*, 45, 533-542,  
1094 10.1080/02786826.2010.547889, 2011.
- 1095 Virkkula, A., Asmi, E., Teinilä, K., Frey, A., Aurela, M., Timonen, H., Mäkelä, T., Samuli, A., Hillamo, R.,  
1096 Aalto, P., Kirkwood, S., and Kulmala, M.: Review of Aerosol Research at the Finnish Antarctic Research  
1097 Station Aboa and its Surroundings in Queen Maud Land, Antarctica, *Geophysica*, 45, 2009.
- 1098 Von Berg, L., Prend, C. J., Campbell, E. C., Mazloff, M. R., Talley, L. D., and Gille, S. T.: Weddell Sea  
1099 Phytoplankton Blooms Modulated by Sea Ice Variability and Polynya Formation, *Geophysical Research*  
1100 *Letters*, 47, e2020GL087954, 10.1029/2020GL087954, 2020.
- 1101  
1102 Weller, R., Schmidt, K., Teinilä, K., and Hillamo, R.: Natural new particle formation at the coastal Antarctic  
1103 site Neumayer, *Atmos. Chem. Phys.*, 15, 11399-11410, 10.5194/acp-15-11399-2015, 2015.
- 1104 Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T.,  
1105 Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto,  
1106 P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-Schmidhauser, R.,  
1107 Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Grüning, C., Faloon, K., Beddows, D., Harrison, R.,  
1108 Monahan, C., Jennings, S. G., O'Dowd, C. D., Marinoni, A., Horn, H. G., Keck, L., Jiang, J., Scheckman, J.,  
1109 McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Lösschau, G., and Bastian,  
1110 S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate  
1111 high quality long-term observations of atmospheric particle number size distributions, *Atmos. Meas. Tech.*, 5,  
1112 657-685, 10.5194/amt-5-657-2012, 2012.
- 1113 Willis, M. D., Burkart, J., Thomas, J. L., Köllner, F., Schneider, J., Bozem, H., Hoor, P. M., Aliabadi, A. A.,  
1114 Schulz, H., Herber, A. B., Leaitch, W. R., and Abbatt, J. P. D.: Growth of nucleation mode particles in the  
1115 summertime Arctic: a case study, *Atmos. Chem. Phys.*, 16, 7663-7679, 10.5194/acp-16-7663-2016, 2016.
- 1116 Yan, C., Dada, L., Rose, C., Jokinen, T., Nie, W., Schobesberger, S., Junninen, H., Lehtipalo, K., Sarnela, N.,  
1117 Makkonen, U., Garmash, O., Wang, Y., Zha, Q., Paasonen, P., Bianchi, F., Sipilä, M., Ehn, M., Petäjä, T.,  
1118 Kerminen, V. M., Worsnop, D. R., and Kulmala, M.: The role of H<sub>2</sub>SO<sub>4</sub>-NH<sub>3</sub> anion clusters in ion-induced  
1119 aerosol nucleation mechanisms in the boreal forest, *Atmos. Chem. Phys.*, 18, 13231-13243, 10.5194/acp-18-  
1120 13231-2018, 2018.
- 1121 Yu, H., Ortega, J., Smith, J. N., Guenther, A. B., Kanawade, V. P., You, Y., Liu, Y., Hosman, K., Karl, T.,  
1122 Seco, R., Geron, C., Pallardy, S. G., Gu, L., Mikkilä, J., and Lee, S.-H.: New Particle Formation and Growth  
1123 in an Isoprene-Dominated Ozark Forest: From Sub-5 nm to CCN-Active Sizes, *Aerosol Science and*  
1124 *Technology*, 48, 1285-1298, 10.1080/02786826.2014.984801, 2014.
- 1125