

Supplement of

Different effects of anthropogenic emissions and aging processes on the mixing state of soot particles in the nucleation and accumulation modes

Yuying Wang^{1,2}, Rong Hu^{1,2}, Qiuyan Wang¹, Zhanqing Li³, Maureen Cribb³, Yele Sun⁴, Xiaorui Song¹, Yi Shang¹, Yixuan Wu¹, Xin Huang¹, Yuxiang Wang¹

¹ Key Laboratory for Aerosol-Cloud-Precipitation of China Meteorological Administration, School of Atmospheric Physics, Nanjing University of Information Science & Technology, Nanjing 210044, China

² State Key Laboratory of Remote Sensing Science, College of Global Change and Earth System Science, Beijing Normal University, Beijing 100875, China

³ Earth System Science Interdisciplinary Center, Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD, USA

⁴ State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

Correspondence to: Yuying Wang (yuyingwang@nuist.edu.cn)




Figure S1. Wind rose diagrams for different months during the field campaign. Units for wind speed are m s^{-1} .




Figure S2. Mean particle number size distributions for different months during the field campaign.




Figure S3. Diurnal variation in the particle number size distribution during the field campaign. The dashed lines show where the 40- and 80-nm particles are located on the ordinate.




Figure S4. Diurnal variations in the number fraction of nonvolatile particles (NF_{NV}) in different months for different particle sizes.