

Kinetics, SOA yields and chemical composition of secondary organic aerosol from β -caryophyllene ozonolysis with and without nitrogen oxides between 213 and 313 K

Linyu Gao^{1,2}, Junwei Song^{1,2}, Claudia Mohr³, Wei Huang⁴, Magdalena Vallon¹, Feng Jiang^{1,2},
5 Thomas Leisner^{1,5}, and Harald Saathoff¹

¹ Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

² Institute of Geography and Geoecology, Working Group for Environmental Mineralogy and Environmental System Analysis, Karlsruhe Institute of Technology, Karlsruhe, Germany

³ Department of Environmental Science, Stockholm University, Stockholm, Sweden

10 ⁴ Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Helsinki, Finland

⁵ Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany

Correspondence to: Linyu Gao (linyu.gao@kit.edu) and Dr. Harald Saathoff (harald.saathoff@kit.edu)

Abstract. β -caryophyllene (BCP) is one of the most important sesquiterpenes (SQTs) in the atmosphere, with a
15 large potential contribution to secondary organic aerosol (SOA) formation mainly from reactions with ozone (O_3)
and nitrate radicals (NO_3). In this work, we study the temperature dependence of the kinetics of BCP ozonolysis,
SOA yields, and SOA chemical composition in the dark and in the absence and presence of nitrogen oxides
including nitrate radicals (NO_3). We cover a temperature range of 213K – 313K, representative of tropospheric
conditions. The oxidized components in both gas and particle phases were characterized on a molecular level by
20 a Chemical Ionization Mass Spectrometer equipped with a Filter Inlet for Gases and AEROSols using iodide as
the reagent ion (FIGAERO-iodide-CIMS). The batch mode experiments were conducted in the 84.5 m³ aluminium
simulation chamber AIDA at the Karlsruhe Institute of Technology (KIT). In the absence of nitrogen oxides, the
temperature-dependent rate coefficient of the endocyclic double bond in BCP reacting with ozone between 243 –
313 K are negatively correlated with temperature, corresponding to the following Arrhenius equation: $k = (1.6 \pm 0.4)$
25 $\times 10^{-15} \times \exp((559 \pm 97)/T)$. The SOA yields increase from 16 \pm 5 % to 37 \pm 11% with temperatures decreasing from
313 K to 243 K at a total organic particle mass of 10 $\mu\text{g m}^{-3}$. The variation of the ozonolysis temperature leads to
a substantial impact on the abundance of individual organic molecules. In the absence of nitrogen oxides,
monomers $C_{14-15}H_{22-24}O_{3-7}$ (37.4 %), dimers $C_{28-30}H_{44-48}O_{5-9}$ (53.7 %) and trimers $C_{41-44}H_{62-66}O_{9-11}$ (8.6 %) are
abundant in the particle phase at 213K. At 313K, we observed more oxidized monomers (mainly $C_{14-15}H_{22-24}O_{6-9}$,
30 67.5 %) and dimers (mainly $C_{27-29}H_{42-44}O_{9-11}$, 27.6 %), including highly oxidized molecules (HOMs, $C_{14}H_{22}O_{7,9}$,
 $C_{15}H_{22}O_{7,9}$, $C_{15}H_{24}O_{7,9}$) which can be formed via hydrogen shift mechanisms, but no significant trimers. In presence
of nitrogen oxides, the organonitrate fraction increased from 3 % at 213K to 12 % and 49 % at 243K and 313K,
respectively. Most of the organonitrates were monomers with C_{15} skeletons and only one nitrate group. Higher
oxygenated organonitrates were observed at higher temperatures, with their signal-weighted O:C atomic ratio
35 increasing from 0.41 to 0.51 from 213K to 313K. New dimeric and trimeric organic species without nitrogen
atoms (C_{20} , C_{35}) were formed in presence of nitrogen oxides at 298-313K indicating potential new reaction
pathways. Overall, our results show that increasing temperatures lead to a relatively small decrease of the rate
coefficient of the endocyclic double bond in BCP reacting with ozone, but to a strong decrease in SOA yields. In
contrast, the formation of HOMs and organonitrates increases significantly with temperature.

40 1 Introduction

Biogenic volatile organic compounds (BVOCs), which are emitted mainly from plants, are the largest source of precursors of atmospheric secondary organic aerosol (SOA), which has profound impacts on visibility, air quality, human health, clouds and climate change (Fehsenfeld et al., 1992; Mellouki et al., 2015; Laothawornkitkul et al., 2009; Charnawskas et al., 2017).

45 Sesquiterpenes (SQTs, $C_{15}H_{24}$), are a class of BVOCs mainly emitted from coniferous trees (Kleist et al., 2012; Matsunaga et al., 2013), scots pines (Kivimäenpää et al., 2020; Weigl et al., 2016), and deciduous trees (Li et al., 2019). Despite their lower atmospheric emission concentrations compared to isoprene and monoterpenes (Faiola et al., 2018), SQTs are important in the atmosphere because of their high reactivities with ozone and large aerosol formation potentials (Ciccioli et al., 1999; Lee et al., 2006a; Lee et al., 2006b; Ng et al., 2007). Depending on
50 region and season they can even play a dominating role (Geron and Arnsts, 2010; Tarvainen et al., 2005; Jardine et al., 2011; Shrivastava et al., 2019).

Among all the sesquiterpene species, β -caryophyllene (BCP) is one of the most abundant and is emitted e.g. from various pine trees, but also from plants used for agricultural purposes (Duhl et al., 2008), like foliage of orange trees (Ciccioli et al., 1999; Hansen and Seufert, 2003), potato plants (Agelopoulos et al., 2000), and cotton
55 (Rodriguez-Saona et al., 2001). BCP is a bicyclic compound with two double bonds with different reactivities towards ozone. The rate coefficient for ozone reacting with the endocyclic double bond at 296 K was determined as $(1.2 \pm 0.4) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Shu and Atkinson, 1994; Richters et al., 2015) [{Cox, 2020 #104}](#). It is estimated to be about 100 times larger than the rate coefficient of $(1.1 \pm 0.4) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ for the exocyclic double bond (Shu and Atkinson, 1994) of the first-generation products. Therefore, the atmospheric
60 lifetime of the endocyclic double bond towards ozone is estimated to about 2 minutes, and the first-generation products can exist about 3.5 hours in typical tropospheric ozone levels of 30 ppb (Winterhalter et al., 2009).

The SOA mass yield of BCP ozonolysis is expected to be dependent on organic particle mass concentration (Odum et al., 1996), temperature (Saathoff et al., 2009) and ozone levels (Chen et al., 2012). Previous studies give a relatively large range of SOA yields of 5-70% depending on the experimental conditions (Jaoui et al. (2003), Lee et al. (2006a), Winterhalter et al. (2009), Chen et al. (2012), Tasoglou and Pandis, 2015), for a temperature range
65 of 287 – 298 K. A comparison of these studies is given in the supplementary information (cf. Table S1).

Yields, products, and mechanisms of BCP ozonolysis were investigated extensively for room temperature (Jaoui et al., 2003; Griffin et al., 1999; Lee et al., 2006a; Lee et al., 2006b; Winterhalter et al., 2009; Nguyen et al., 2009; Jenkin et al., 2012; Richters et al., 2016). The oxidation products mainly include aldehydes, small acids, acetone
70 in the gas phase (Grosjean et al., 1993; Calogirou et al., 1999; Zhao et al., 2010; Larsen et al., 1998), and ketones, aldehydes, alcohols and carboxylic acids in the particle phase (Dekermenjian, 1999; Lee et al., 2006a; Jaoui et al., 2003; van Eijck et al., 2013; Jaoui et al., 2007; Li et al., 2011; Alfarra et al., 2012). Major oxidation products identified are e.g. β -caryophyllonic acid ($C_{15}H_{24}O_3$), β -caryophyllinic acid ($C_{14}H_{22}O_4$, BCA), β -hydroxycaryophyllonic acid ($C_{15}H_{24}O_4$) (Table S3). BCA has also been used as a tracer to estimate the emissions
75 of BCP in the real atmosphere (Jaoui et al., 2007; Hu et al., 2008; Parshintsev et al., 2008; Haque et al., 2016; Verma et al., 2021; Cheng et al., 2021). In addition, highly oxidized multifunctional organic molecules (HOMs) were observed from BCP ozonolysis. Richters et al. (2016) studied their formation mechanism at 295 K in a free-jet flow system using isotopic labelling. Major HOMs identified are $C_{14-15}H_{22}O_7$, $C_{14-15}H_{22}O_9$, and $C_{15}H_{22}O_{11,13}$ (Richters et al., 2016).

80 SQT oxidation products are expected to have low or extremely low vapor pressures owing to their long carbon skeletons with multiple functional groups. For example, the saturation vapor pressure of β -caryophyllinic acid was estimated as low as 3.3×10^{-13} Pa (Li et al., 2011), which leads to a high condensation and SOA formation potential. As a consequence, these species are expected to play an important role in new particle formation (NPF) (Kirkby et al., 2016; Kammer et al., 2020; Huang et al., 2021).

85 Especially during night-time, the reaction between nitrate (NO_3) radicals and unsaturated hydrocarbons is of substantial importance and results in SOA formation. BCP has a high reactivity towards NO_3 radicals, with a rate coefficient of $(1.93 \pm 0.35) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Shu and Atkinson, 1995; Winterhalter et al., 2009; Calvert et al., 2008) [{Cox, 2020 #103}](#). SOA formation from BCP ozonolysis in the presence of NO_3 radicals has only been subject of few studies so far. SOA yields from dark reactions between BCP and NO_3 were estimated to be 1.46
90 and 0.91 for SOA mass concentrations of 113.4 and $60.3 \mu\text{g m}^{-3}$, respectively (Jaoui et al., 2013). The formation mechanism of several organic nitrates including $\text{C}_{15}\text{H}_{23,25}\text{O}_4\text{N}$ and $\text{C}_{15}\text{H}_{25}\text{O}_{6,7}\text{N}$ is described in the Master Chemical Mechanism (MCM v3.2) by Jenkin et al. (2012).

The troposphere covers a wide temperature range which depends on the altitude from the planetary boundary layer (PBL) to the upper troposphere, latitude, and seasonal variations. Typically, the tropospheric temperatures
95 range between 300 K to 200 K. In summer, some regions can even reach to 318 K. BVOCs and their oxidation products are proposed to undergo transport from the PBL to the upper troposphere e.g. via convective systems typical for tropical regions (Andreae et al., 2018). In this conceptual picture of a potential aerosol life cycle the temperature dependence of the different processes involved is quite important. Besides, multiple studies have shown that SOA yields typically increase with decreasing temperature, e.g. for α -pinene and limonene ozonolysis
100 (Saathoff et al., 2009), β -pinene ozonolysis (Von Hessberg et al., 2009), and isoprene ozonolysis (Clark et al., 2016). In addition, at lower temperatures also semi volatile ~~components vapours can have a big condensation potential alter their partitioning behavior due to reduced vapour pressures~~ (Stolzenburg et al., 2018; Ye et al., 2019). ~~Considering Despite of~~ the potentially important role of BCP oxidation products in NPF and their high condensation and SOA formation potential ~~as abovementioned~~, ~~studies on~~ the temperature dependence of SOA
105 formation from ozonolysis of BCP is, ~~however, mainly unclear to date still scarce~~.

The objectives of this work were to study the temperature dependence on the kinetics, yields, and chemical composition distribution of the aerosol produced by the reaction of BCP with O_3 in the temperature range between 213 – 313 K. Furthermore, we investigated the impact of the presence of nitrogen oxides including NO_3 radicals on the SOA chemical composition and volatility.

110 2 Methodology

2.1 Experimental conditions

A schematic of the AIDA (Aerosol Interaction and Dynamics in the Atmosphere) simulation chamber at the Karlsruhe Institute of Technology (KIT) and the main instrumentation used in this work are shown in Fig. 1. The chamber is an 84.5 m^3 aluminium vessel equipped with a LED solar radiation simulator and with precisely
115 controlled temperature, humidity, and gas mixtures. It is operated as a continuously stirred reactor with mixing times of 1-2 minutes achieved using a fan about 1 m above the bottom [of the chamber](#) (Saathoff et al., 2009). Wall and gas temperature inside the chamber are controlled at $\pm 0.3 \text{ K}$ over a wide range of temperatures (313-183 K)

(Wagner et al., 2006). The pressure inside the chamber can be varied from 0.01 to 1000 hPa and the relative humidity (RH) can be modified between close to zero to up to 100% and even supersaturated conditions (Möhler et al., 2003). Water vapor is measured in situ by a tuneable diode laser (TDL) hygrometer with an accuracy of $\pm 5\%$ and a calibrated reference dew point mirror hygrometer (MBW373LX, MBW Calibration Ltd.) with an accuracy of $\pm 1\%$ (Fahey et al., 2014).

The results presented in this work are from dark BCP ozonolysis experiments with or without addition of NO_2 from a campaign in November and December 2019 covering five different temperatures between 213 - 313 K.

The experimental conditions are listed in Table 1. Two additional experiments were undertaken in March 2020 to study the rate coefficients of BCP reacting with ozone at 243 K and 258 K. BCP (98%, Carl Roth GmbH) was added to the AIDA chamber with a flow of 0.01 m^3/min of synthetic air ~~saturated~~potentially saturated with its vapour at 298K. Please note that the BCP concentrations for the experiments at 213 K could not be measured due to the low vapour pressure and strong wall losses at lower temperatures (more details are given in section 2.2).

During the experiment at 213K the initially added BCP was lost to the walls, so almost no SOA was formed after the addition of ozone. However, when adding BCP in the presence of ozone, SOA was formed in quantities comparable to the other experiments. Ozone was in all experiments typically in excess and generated by a silent discharge generator (Semozon 030.2, Sorbios) in pure oxygen (99.9999%). The relative humidity ranged from 96% to 13% for experiments at 213 K and 313 K, respectively. This corresponds to water vapor concentrations of 1 Pa ($3.4 \times 10^{14} \text{ cm}^{-3}$) at 213 K and 952 Pa ($2.2 \times 10^{17} \text{ cm}^{-3}$) at 313 K, respectively, and reflects the variability of the water vapour concentrations throughout the troposphere. At the initial phase of each experiment, BCP was depleted completely by ozonolysis and SOA was formed. After depletion of the BCP at a lower ozone level to facilitate the kinetic study we increased the excess of ozone to accelerate the oxidation of remaining double bonds.

Yields and chemical composition are determined and compared for the time period after increasing the ozone level. Then a second addition of BCP generated more SOA mass. Due to the large excess of ozone, BCP could not be measured during the subsequent additions. The corresponding conditions are marked as 1a-5a in Table 1. Subsequently, NO_2 (1000 ppm of 99.5% purity in nitrogen 99.999%, Basi Schöberl GmbH) was added to the reaction mixture still containing an excess of ozone. ~~A second~~Another step of SOA formation was then initialized by adding more BCP in presence of NO_3 radicals. This series of experiments is marked as 1b-5b in Table 1.

Hence, the BCP and ozone concentrations listed in Table 1 for experiments 1b-5b include also the amounts added in experiments 1a-5a. Ozone was at nearly the same concentrations as for the initial experiments without NO_2 , except for the experiments at 273 K. To get a slower decay of BCP to better determine its rate coefficient at 273K, we added 73 ppb ozone into the chamber first, and then added more ozone to the same concentration level (~ 300 ppb) as used for the other temperatures for comparison. All experiments were performed in the dark and no hydroxyl radical scavenger was used. Hence, the OH radicals generated in the ozonolysis reaction with a yield of $(8 \pm 3) \%$

{Cox, 2020 #103}, $(10.4 \pm 2.3) \%$ (Winterhalter et al., 2009) or 6% (Shu and Atkinson, 1994) also contributed to SOA formation as will be detailed in section 3.1. Time zero in the plots refers to the first addition of ozone to the reaction mixture. Please note that we tried to have similar organic particle mass concentrations among the different experiments in order to limit a potential influence of absorptive partitioning on the comparison among different temperatures. The organic aerosol mass levels were in the range of $4.5\text{-}31.5 \mu\text{g m}^{-3}$ for the initial ozonolysis and between 14 and $41 \mu\text{g m}^{-3}$ for the reaction step with NO_3 radicals. Since we had relatively similar and sufficient

[organic aerosol mass levels available for partitioning, the observed differences in chemical composition should be governed by the changing chemistry, mainly in the gas phase.](#)

160 2.2 Instrumentation

The main instruments used for this study are shown in Fig. 1. They were installed outside the AIDA chamber and operated at 296 K. [To avoid potential artefacts, e.g. due to the phase partitioning of semi-volatile species in the sampling lines from the chamber to the instruments, the sampling lines were partially insulated and the residence time was below 1-2 seconds.](#) The concentrations of O₃ and NO₂ were measured using the gas monitors O₃41M and AS32M (both Environment S.A.), respectively.

165 The concentrations of BCP and low-oxygenated gaseous oxidation products were measured by a Proton-Transfer-Reaction-Time-of-Flight-Mass-Spectrometer (PTR-ToF-MS 4000, Ionicon Analytic GmbH). Data was analysed using PTR viewer 3.3.12. The PTR-ToF-MS is also interfaced with a particle inlet (Chemical Analysis of Aerosol Online, CHARON), which allows to measure the semi-volatile particle components. A detailed description of the CHARON-PTR-MS has been provided elsewhere (Müller et al., 2017; Piel et al., 2021). During the period of BCP injection and ozonolysis, the PTR-ToF-MS was operated for measuring gaseous volatile organic compounds (VOCs) only. After BCP was fully depleted, we switched the PTR-ToF-MS to an alternating measurement mode for detecting the low-oxygenated organic molecules in both gas and particle phases. This alternating measurement mode included 3-min HEPA filter measurement for the particle background, 5-min CHARON particle measurement, 1-min transition for instrument equilibration, 5-min of VOC measurement and another 1-min transition. The CHARON-PTR-MS measured the particle phase at a sampling flow of 500 SCCM via a 1/4" silcosteel tube, while the gas phase was measured at a flowrate of 100 SCCM via a 1/16" PEEK tube taken from the particle measurement flow. Furthermore, a flowrate of 3.9 l/min was added to the total flow to minimize the residence time in the sampling tube. For measuring gases, the drift tube of the PTR-MS was kept at 393 K and 2.8 mbar leading to an electric field (E/N) of 127 Td. During alternating measurements, the drift tube was automatically optimized to 100 Td for particle measurement. BCP was calibrated using a liquid calibration unit (LCU-a, Ionicon Analytic GmbH). The PTR-ToF-MS at an E/N of 127 Td showed significant fragmentation of BCP, in agreement with previous studies (Kim et al., 2009; Kari et al., 2018). In this study, we observed that the parent ion (m/z 205.20, C₁₅H₂₅⁺) contributed (29 ± 1) % to total signals of BCP-related ions including m/z 81.07, 170 m/z 95.09, m/z 109.10, m/z 121.10, m/z 137.13, m/z 149.13 and m/z 205.20 (Fig. S1). Therefore, we scaled the concentration of m/z 205.20 (C₁₅H₂₅⁺) by a factor of 3.45 for the quantification of BCP. The total uncertainty of BCP quantification was estimated as $\pm 20\%$ by including all errors mainly related to the uncertainties of the LCU and the fragmentation pattern of BCP. In this study, we calibrated the mixture of toluene and BCP with a solvent of *n*-hexane using LCU. We obtained a similar sensitivity for toluene using the LCU and a gas cylinder with a toluene standard. Then we calculated the sensitivity of BCP relative to the sensitivity of toluene. This resulted in 175 36.2 ncps/ppb for the parent ion (C₁₅H₂₅⁺, m/z 205) of BCP. The total uncertainty of the quantification of BCP was estimated to $\sim 20\%$ by including the uncertainties of toluene in the gas standard ($\sim 10\%$), the LCU calibration procedure ($\sim 15\%$), and the fragmentation pattern of BCP ($\sim 5\%$).

180 Particle size distributions and number concentrations were measured by a scanning mobility particle sizer (SMPS) 195 utilizing a differential mobility analyzer (DMA, 3071 TSI Inc.) connected to a CPC (3772, TSI Inc.). Particle

number concentrations were measured by two condensation particle counters (3022a and 3776, TSI Inc.). The particle number size distributions of the SMPS were corrected for the total number concentration measured by a calibrated CPC and used to calculate the SOA mass concentration by applying an effective particle density. This particle density was determined by comparing the mobility and aerodynamic size distributions measured by SMPS and AMS, calculated increasing from 0.9 ± 0.1 at 243K to 1.1 ± 0.1 at 313K, respectively (Saathoff et al., 2009).

200 A High Resolution-Time-of-Flight-Aerosol Mass Spectrometer (HR-ToF-AMS, Aerodyne Inc.) was used to continuously measure the total organic particle mass with a time resolution of 30 seconds at a total flow rate of $0.0011 \text{ m}^3/\text{min}$ (with only $0.0008 \text{ m}^3/\text{min}$ going into the instrument). The data was analyzed using the PIKA v1.60C software. For calculation of the organic particle concentration from the AMS mass spectra, a collection efficiency of 0.6 (determined by comparison with the SMPS results) and an ionization efficiency of $(1.52 \pm 0.1) \times 10^{-7}$ (calibrated with 300 nm ammonium nitrate particles) were used. [In the absence of inorganic species, we calculated the mass concentration of the organonitrates \(OrgNO₃, i.e., organonitrates\) from AMS data, by assuming an average OrgNO₃ molecular weight of \$331 \text{ g mol}^{-1}\$ and \$363 \text{ g mol}^{-1}\$ based on the most abundant molecules \(C₁₅H₂₅O₇N and C₁₅H₂₅O₉N\) detected by FIGAERO-iodide-CIMS.](#)

205 Furthermore, we used a chemical ionization mass spectrometer (CIMS) equipped with a Filter Inlet for Gases and AEROSols (FIGAERO) (Aerodyne Research Inc.) using iodide as the reagent ion to measure both gas-phase and particle-phase components. The FIGAERO-iodide-CIMS has been described in detail by Lopez-Hilfiker et al. (Lopez-Hilfiker et al., 2014). It allows very sensitive detection of oxygenated compounds avoiding significant fragmentation. In our experiments, the gas-phase compounds were measured online by FIGAERO-iodide-CIMS with a total flow rate of $0.005 \text{ m}^3/\text{min}$, with only $0.002 \text{ m}^3/\text{min}$ going into the instrument. [The bypass flow of \$0.003 \text{ m}^3/\text{min}\$ was used to reduce the residence time in the sampling line.](#) In parallel, particles were deposited offline on prebaked Teflon filters (Polytetrafluoroethylene, PTFE, $1 \mu\text{m}$, SKC Inc.) using a stainless-steel filter holder with a flow rate of $0.006 \text{ m}^3/\text{min}$ for 5 min, the same offline sampling procedure as described by Huang et al. (2019a). After particle deposition, the filter samples were stored at 243 K for analysis after the experiment.

210 These filters were then heated by FIGAERO-iodide-CIMS using a flow of ultra-high purity nitrogen as carrier gas following a thermal desorption procedure from 296 K to a maximum temperature of 473 K with a total desorption time of 35 minutes (Huang et al., 2018). [We collected 4, 3, 3, 2, 2 particle filter samples for the periods when the particle concentrations reached stable levels in experiments at 213K, 243K, 273K, 298K, and 313K, respectively.](#) The data analysis was done with the Tofware software (version 3.1.2). Note that the reagent ion I⁻ ($m/z = 126.9$) was subtracted from the mass-to-charge ratio of all the molecules shown in this work. For the comparison of measured total particle mass concentration between AMS, SMPS and FIGAERO-iodide-CIMS, we use the maximum sensitivity of 22 cps ppt^{-1} to convert the signals to mass concentration (Lee et al., 2014; Lopez-Hilfiker et al., 2016). In addition, we did a mass calibration for β -caryophyllinic acid (95%, Toronto Research Chemicals) resulting in a sensitivity of $(2.4 \pm_{-0.63}^{0.96}) \text{ cps ppt}^{-1}$ (details are given in Fig. S2). [We included I⁻ \(\$m/z = 127 \text{ Th}\$ \), I\(H₂O\)⁻ \(\$m/z = 145 \text{ Th}\$ \), I\(CH₂O₂\)⁻ \(\$m/z = 173 \text{ Th}\$ \), I₃⁻ \(\$m/z = 381 \text{ Th}\$ \), and a dominating product ion C₃₀H₄₈O₅I⁻ \(\$m/z = 615 \text{ Th}\$ \) for our mass calibration. With this procedure we found the mass defect of most compounds in a good linear correlation and without significant deviations at high masses \(cf. Figure S3\). We constrained the peak assignment errors to 20 ppm.](#)

220 Typically, background measurements for both gas and particle phase were done before and after the first addition of BCP to identify any contamination inside the chamber. However, gas background levels were almost negligible

225


230

235

for most experiments and most of the particle background signals were from filter matrix contaminations mainly due to fluorinated constituents. [We subtracted the mass spectra of the background filter samples from those of the particle loaded filter samples for the same experiments.](#)

2.3 Rate coefficient calculation

240 Based on the PTR-MS measurements of the BCP decay as well as the ozone measurements, the rate coefficients of the reaction of BCP with ozone can be determined. As BCP has two double bonds with reactivities with ozone differing by a factor of 100 (Shu and Atkinson, 1994), here we discuss only the rate coefficient for the reaction of the most reactive endocyclic double bond. Since we did not use an OH radical scavenger, the reaction between BCP and OH radicals, which are generated from BCP ozonolysis, was included in our analysis. Employing the
245 following reaction scheme, the observed decays of ozone and BCP were fitted by adjusting only the rate for reaction R1.



For the reaction of OH radicals with BCP, the rate coefficient of $(1.97 \pm 0.25) \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1}$ determined for
250 296 K (Shu and Atkinson, 1995; Winterhalter et al., 2009) [{Mellouki, 2021 #106}](#) was used for all temperatures due to the lack of its temperature dependence.

To integrate this simple model and to fit the rate coefficient for reaction R1 we used the software KinSim (Kinetics Simulator for Chemical-Kinetics and Environmental-Chemistry Teaching, version 4.14) (Peng and Jimenez, 2019). The OH radical yields (γ in R1) varied between 5-15% at different temperatures (see section 3.1).

255 2.4 SOA yields calculation

The SOA yields (Y_{SOA}) were calculated as $Y_{SOA} = \Delta M_{org} / \Delta VOC$, where ΔM_{org} is the SOA mass formed from the reacted mass of BCP (ΔVOC). Similarly, the yields (Y_{BCA}) of BCA, the typical product of BCP ozonolysis, were calculated as $Y_{BCA} = \Delta M_{BCA} / \Delta VOC$, where ΔM_{BCA} is the mass concentration of β -caryophyllinic acid formed from the reacted mass of BCP (ΔVOC). We used a more than 5-fold ozone excess in this study for all temperatures to
260 facilitate oxidation of all double bonds in BCP and its oxidation products. This should lead to more comparable yields for all conditions studied (Li et al., 2011; Chen et al., 2012). Wall losses of particles and semi volatile trace gases were calculated with the aerosol dynamic model COSIMA (Naumann, 2003; Saathoff et al., 2009) and used to correct the yields. The yields were calculated for the initial period of the experiments, which lasted about 90 minutes. During this relatively short time period and due to the large size of the simulation chamber, particle
265 losses contributed typically 6% or less to the total SOA mass. Due to the relatively small particle sizes at the beginning of the experiments, diffusional losses dominated.

3 Results and discussion

In this chapter we discuss the typical course of the experiments, the kinetics of the reaction of BCP with ozone, the SOA mass yields, as well as the SOA chemical composition in absence and in presence of nitrogen oxides.

270 3.1 Overview of the experiments and kinetics of BCP ozonolysis

Figure 2 shows the evolution of trace gases as well as particle mass and size distribution for BCP ozonolysis at 298K first without and then in presence of nitrogen oxides in experiment 4a and 4b. The time point when ozone was firstly added is regarded as the start of each experiment and is set to zero. As shown in Fig. 2, following the nucleation, the particle size increased to 36 nm and grew further to 57 nm and 69 nm after subsequent additions of ozone and BCP, and finally to 122 nm after NO₂ addition. The particle mass increased in presence of the low initial ozone concentration (25 ppb) and stabilized within 20 minutes after increasing the ozone level to above 300 ppb reaching 13.3 μg m⁻³. A second addition of more BCP led to another increase of the particle mass stabilizing at a level of 17.0 μg m⁻³. To this SOA, 42 ppb of NO₂ were added, which reacted with the excess of ozone forming NO₃ radicals and consequently led to a small increase in particle size and mass (1.9 nm and 0.9 μg m⁻³, respectively) due to their reaction with BCP oxidation products. After the third addition of BCP to the reaction mixture, the SOA mass concentration increased to 50.6 μg m⁻³. For the other four temperatures, these values are given in Table 1, and their time evolution is shown in Fig S4. During the final addition of BCP, the existing particles grew quickly and new particle formation ~~were~~ was observed. The discrepancy in mass concentrations obtained from SMPS and HR-TOF-AMS in the initial phase of the experiment are due to the fact that smaller particles are poorly detected by the HR-TOF-AMS due to the lower transmission of sub-100nm particles in the aerodynamic lens (Liu et al., 2007; Williams et al., 2013). After 160 minutes, the particle size reached 122 nm and both mass concentrations agreed well assuming a collection efficiency of 0.6 for the AMS. Please note that the FIGAERO-iodide-CIMS is sensitive to more polar oxidized compounds, thus, the sum of all compounds detected by CIMS can only be a fraction of the total organic aerosol compounds measured by HR-TOF-AMS. The total organic mass concentration detected by FIGAERO-iodide-CIMS, if assuming an average maximum sensitivity for all compounds, corresponds to 36-61 % of the total organic aerosol mass measured by AMS. On average 36-61 % of the total organic mass concentration was detected by FIGAERO iodide CIMS.

After the first addition of an excess of ozone, BCP was depleted within less than 5 minutes. An example of the kinetic model results for the experiment at 313 K is compared to the measured data in Fig. S3-S5 and all kinetic parameters fitted are listed in Table S2. The OH radical yields from the ozonolysis reaction increase from (5±2) % at 243 K to (15±2) % at 313K, and 91-92% of the BCP are calculated to react with ozone under 243-313K.

Figure 3 shows the temperature dependence of the rate coefficients for BCP ozonolysis. The rate coefficient we determined for 298 K of $(1.09 \pm 0.21) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ agrees very well with values from Shu and Atkinson (1994) of $(1.16 \pm 0.43) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and Richters et al. (2015) of $(1.1 \pm 0.3) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ for 296 K. At lower temperatures, the rate coefficient increases. This is in an agreement with the density-functional theory (DFT) quantum chemical calculations by Nguyen et al. (2009) but with a slightly higher slope. The Arrhenius equation fitted to our measured values is $k = (1.6 \pm 0.4) \times 10^{-15} \times \exp((559 \pm 97)/T)$. This corresponds to a reaction enthalpy of $(5.6 \pm 1.0) \text{ KJ mol}^{-1}$ in our analysis. Please note that the experiment at 273 K was not used for this analysis because of an unusual background signal in the PTR-MS measurement (Fig. S5bS7b).

305 3.2 Particle mass yields

Figure 4 presents the yields from BCP ozonolysis in the absence of NO₂ as a function of the temperature for a constant total organic particle mass concentration (M_{org}) of 10 μg m⁻³. The SOA yields decrease with increasing temperatures from 243K to 313K. For this organic particle mass concentration, we determined a SOA yield of

(19.4±6) % at 298K, which was lower by around 55% and 40% than those reported by Chen et al., (2012) and Tasoglou and Pandis (2015), respectively, for a similar particle mass. However, the results are still within the combined uncertainty limits. Please note that these two previous studies used OH radical scavengers, which should lead to lower yields, since OH radicals may oxidize potentially more VOCs causing higher SOA yields [for the ozonolysis without OH scavenger system](#). The temperature dependence we observed is significantly lower than e.g. for α -pinene (Saathoff et al., 2009) and hence reflects the generally lower vapour pressure of the condensable oxidation products compared to the monoterpene. In our study, the SOA formation time of about 90 minutes was longer than the typical lifetime of the first-generation products from BCP ozonolysis (20 minutes) at an ozone level of 300 ppb, thus the difference cannot be explained by potential incomplete reactions. Different initial ozone levels may also contribute to the higher yields reported by Chen et al., (2012) and Tasoglou and Pandis (2015). [For our study the wall losses and corresponding corrections were relatively small and can't explain deviations of 40-55%. Typically, the losses of acidic gases are larger in an aluminium chamber compared to a Teflon chamber and it may be the opposite for particle losses. This depends on the age of the chamber walls, potential electrostatic losses, and the volume to surface ratio of the chamber. It is therefore not easy to determine the impact of the different wall losses and the wall loss corrections. We can only speculate if too high wall loss corrections for the studies in the Teflon chamber contribute to the different yields.](#)

An overview of the SOA yields determined by Lee et al., 2006, Winterhalter et al, 2009, Chen et al., 2012 and Tasoglou and Pandis (2015) is given in Fig. [S4-S6](#) in comparison with our results in the temperature range between 243 and 313 K. The SOA masses formed reflect the different oxidation products of differing volatilities (Chan et al., 2007) formed at low and high temperatures, which will be discussed in the following section.

3.3 Chemical characterization of SOA from BCP ozonolysis without nitrogen oxides

In this section the gas and particle phase molecular composition will be discussed mainly based on the FIGAERO-iodide-CIMS mass spectra, complemented by the CHARON-PTR-MS.

3.3.1 Composition of the gas-phase products

Figure 5 shows the averaged gas-phase mass spectra for all five temperatures in the range 213-313 K, before adding NO₂ and when the SOA concentration was in a relatively stable state. Only compounds with mass-to-charge ratios less than 400 Th contributed significantly to the FIGAERO-iodide-CIMS spectra. The signals were normalized to the total gas-phase C_xH_yO_z compounds at each temperature. In the temperature range of 213 – 273 K, C₁₅H₂₄O₃ (likely β -caryophyllonic acid), C₁₅H₂₄O₄ (likely β -hydroxycaryophyllonic acid), C₁₅H₂₆O₄ (not identified yet), C₁₄H₂₂O₄ (β -nocaryophyllonic acid or β -caryophyllinic acid) were the most dominant monomeric compounds in the gas phase. These compounds have been identified as products from BCP ozonolysis at room temperature in previous studies (Jaoui et al., 2003; Winterhalter et al., 2009; Chan et al., 2011; Li et al., 2011). [The absolute signals of all detected gas phase species are shown in the Fig. S8.](#) In addition, we also observed several abundant ions, such as C₁₄H₂₀O₂, C₁₅H₂₂O₂ and C₁₄H₂₀O₃, in the PTR-MS mass spectra (Fig. [S5S7](#)). As mentioned above, the PTR-MS is more sensitive to less-oxygenated organic compounds and is prone to fragmentation via e.g. losing H₂O. In combination with FIGAERO-iodide-CIMS data, it is reasonable to interpret that C₁₅H₂₂O₂ in PTR-MS mass spectra is most likely the fragmentation ion from C₁₅H₂₄O₃ via losing H₂O. Besides, C₁₅H₂₄O₂ and its potential fragmentation ion C₁₅H₂₂O could be tentatively identified as BCP aldehyde based on

previous studies (Li et al., 2011). This indicates that the low oxidized monomeric compounds ($O < 5$), which are abundant at lower temperatures, can also be formed at higher temperatures (273-313K) but react (cf. Fig. S7S10) to form higher oxygenated and lower volatile compounds. It cannot be excluded that low molecular-weight compounds, like $C_4H_6O_4$ and $C_2H_4O_3$, might be from the similar process.

3.3.2 Composition of the particle-phase products

Figure 6 shows the particle-phase mass spectra from BCP ozonolysis for 213 K and 313 K. For both lowest and highest temperatures, several first- and second-generation oxidation products, some of which have been identified also in previous studies, were observed, e.g. $C_{14}H_{22}O_4$ (β -caryophyllinic acid or β -hydroxynocaryophyllon aldehyde), $C_{15}H_{24}O_3$ (β -hydroxycaryophyllon aldehyde or β -caryophyllonic acid), and $C_{14}H_{22}O_7$ (2,3-dihydroxy-4-[2-(4-hydroxy-3-oxobutyl)-3,3-dimethylcyclobutyl]-4-oxobutanoic acid) (Jaoui et al., 2007; Li et al., 2011; Winterhalter et al., 2009; Chan et al., 2011; Jaoui et al., 2003; Griffin et al., 1999; Lee et al., 2006a; Lee et al., 2006b; Nguyen et al., 2009; Jenkin et al., 2012; Richters et al., 2016). Table S3 lists all major particulate $C_xH_yO_z$ compounds detected by FIGAERO-iodide-CIMS in this work. Furthermore, we also observed e.g., $C_{14}H_{20}O_2 H^+$, $C_{14}H_{20}O_3 H^+$, $C_{14}H_{22}O_3 H^+$, $C_{15}H_{22}O_2 H^+$ and $C_{15}H_{22}O_3 H^+$ with the CHARON-PTR-MS (Fig. S6S9). From previous studies we know that CHARON-PTR-MS measurements are affected by fragmentation, e.g. by losing H_2O , CO and CO_2 , due to the relatively high collisional energy in the drift tube (100-170 Td) (Gkatzelis et al., 2018). In this study, we operated the CHARON-PTR-MS at 100 Td for the particle measurement. Taking fragmentation by losing one H_2O into account, it is reasonable to speculate that $C_{14}H_{20}O_3$ and $C_{15}H_{22}O_2$ are from the fragmentation of $C_{14}H_{22}O_4$ and $C_{15}H_{24}O_3$, respectively. Similarly, $C_{14}H_{20}O_2$ and $C_{15}H_{22}O_3$ in CHARON-PTR-MS mass spectra could be identified as the fragmentation ions from $C_{14}H_{22}O_3$ and $C_{15}H_{24}O_4$, respectively, which were the most abundant ions in the FIGAERO-iodide-CIMS mass spectra as shown in Fig. 6.

At 213 K, monomers, dimers and trimers are clearly visible. The most abundant compounds measurable with our FIGAERO-iodide-CIMS in each group are $C_{15}H_{24}O_3$, $C_{30}H_{48}O_5$ and $C_{44}H_{68}O_9$, respectively. At 313 K, the monomers dominate and only a few dimers are observed, with the most abundant signals by compounds $C_{14}H_{22}O_7$ and $C_{29}H_{44}O_{10}$, respectively.

Compared to 213K, it is evident that the monomeric compounds formed at 313K are more oxygenated and have higher elemental O:C ratios (Fig. 6). It is inferred that at higher temperatures, once the first-generation products are formed in the gas phase, some of them can remain in the gas due to higher saturation vapor pressures and undergo further oxidation reactions of their unsaturated exocyclic double bonds with the excess of ozone. Since the saturation vapor pressures of the compounds at 213K are substantially lower than at 313K, more relatively lower oxidized molecules are found in the particle phase due to rapid condensation. For the compounds remaining in the gas phase, HOMs such as e.g. $C_{14}H_{22}O_7$ could be formed via simple or extended autoxidation (Richters et al. (2016)). In addition to $C_{14}H_{22}O_7$, we also detected several other compounds (e.g. $C_{14}H_{22}O_9$, $C_{15}H_{22}O_{7,9}$, and $C_{15}H_{24}O_{7,9}$) that are likely products of autoxidation reactions (Jokinen et al., 2016). At 313K, the monomeric highly oxidized compounds (MHOC, $C \leq 15$, $O \geq 6$) dominate the monomers with a signal fraction of 42.5% to total organic signals, which exceeds the contribution of other monomeric low oxygenated organic compounds (MLOC, $C \leq 15$, $O < 6$) with a signal fraction of 24.9%. For comparison, at 213K, MHOC and MLOC contribute 6.9% and 30.5%, respectively. Furthermore, the six confirmed HOM have signal fractions of 9.2% at 313K and nearly zero at 213K. Note that higher oxygenated autoxidation products like $C_{15}H_{22}O_{11}$ (detected by Atmospheric Pressure

interface Time-Of-Flight mass spectrometer (CI-API-TOF) using nitrate as the reagent ion (Jokinen et al., 2016)) and $C_{15}H_{22}O_{13}$ could not be detected in this work due to instrument limitations (Riva et al., 2019). To summarize, the relative abundance of HOMs were higher at increasing temperatures (Fig. 8), indicating that the autoxidation showed a strong temperature dependence, slowing down at reduced temperatures.

390 The dimer groups show two completely different patterns at 213K and 313K. At 213K, the dimers mainly consist of low oxygenated organic compounds (DLOC) with 28-30 carbon atoms and 5-8 oxygen atoms, and the total signal fraction of the dimeric compounds is 53.7%. At 313K, dimers are more oxygenated with 9-11 oxygen atoms with the same carbon number of 28-30 (DHOC), and they contribute less to the total normalised organic signal (27.6%). Here, we put forward esterification as a potential pathway for forming the most abundant dimers at both
395 213K and 313K, similar to the dimer formation of other biogenic VOCs suggested by Yasmeeen et al. (2010) and Müller et al. (2008). The dominating dimeric molecule at 213K, $C_{30}H_{48}O_5$, could potentially be formed via esterification of two of the most abundant monomers ($C_{15}H_{26}O_3$ and $C_{15}H_{24}O_3$). In a similar way, the dimer $C_{29}H_{44}O_9$ could be formed at 313K from $C_{14}H_{22}O_7$ and $C_{15}H_{24}O_3$. These reactions are described in Fig. [S8-S11](#) in the supplement.

400 However, we cannot exclude the reaction of a BCP-derived stabilized Criegee intermediate and an abundant acid as a potential pathway for $C_{30}H_{48}O_5$ formation, similar to mechanisms suggested for e.g. α -pinene (Kristensen et al., 2016; Wang et al., 2016; Witkowski and Gierczak, 2014; Lee and Kamens, 2005). The dimers we observed had no more than 11 oxygen atoms, which is due to low sensitivities of FIGAERO-iodide-CIMS to these compounds (Riva et al., 2019).

405 Significant signals of trimers at 213K are assigned to $C_{41-43}H_{62-68}O_{9-11}$, but they are not detected at 313K. The potential formation pathway of $C_{44}H_{68}O_9$ at 213K is also included in Fig. [S8S11](#). Please note, that the assignment for ions at mass to charge ratios of more than 700 Th has significantly higher uncertainties as for smaller mass peaks.

Figure 7 presents the averaged mass spectra of SOA particles formed from BCP ozonolysis before NO_2 addition
410 at all five temperatures. With increasing temperature, the most abundant monomeric compounds shift to higher masses with higher elemental O:C ratios, e.g. from $C_{15}H_{24}O_3$ ($m/z = 252.2$, O:C = 0.20, 213-243K), $C_{15}H_{24}O_4$ ($m/z = 268.2$, O:C = 0.27, 273K) to $C_{14}H_{22}O_6$ ($m/z = 286.1$, O:C = 0.43, 298K) and $C_{14}H_{22}O_7$ ($m/z = 302.1$, O:C = 0.5, 313K). This indicates again that at higher temperatures, after ozonolysis of the endocyclic bond (formation of first-generation oxidation products), the unsaturated compounds can react further with the excess of ozone and
415 form higher oxidized products (e.g. HOM), while those formed at lower temperatures would partition into the condensed phase before further oxidation can occur. As shown in Fig. 8, the monomer groups (MHOC+MLOC) contribute 39.2%, 64.7%, 85.3%, 68.9% and 67.5% to the total signal from 213K to 313K, respectively. Among all the monomers, the signal fraction of the six identified HOM to the total signals of all organic species has a monotonic positive temperature dependence (cf. Fig. 8), increasing from 0.1% to 9.2% for temperatures increasing
420 from 213K to 313K. This is a similar correlation between HOM formation and temperatures as observed by (Bianchi et al., 2019).

Two different dimeric patterns appear in the temperature range of 213-313K ([Figure 6 and 7](#)). One pattern is represented by molecular formulae of $C_{28-30}H_{42-48}O_{5-8}$ at 213-243K (marked as low temperature group, LT-group), and the other pattern is represented by $C_{28-30}H_{36-44}O_{9-11}$ at 273-313K (marked as high temperature group, HT-group). The LT-group contributes 53.7% at 213K and 32.8% at 243K, with a negative temperature dependence,
425

while the dimeric signal fraction of the HT-group is lower than LT-group, contributing 13.8% at 273K, 24.6% at 298K, and 27.6% at 313K, respectively, as shown in Fig. 8. After 273K, dimer formation is enhanced with increasing temperatures. The contribution in gas and particle phase of the major compounds are shown in Fig. [S9S12](#).

430 **3.4 Chemical characterization of SOA from BCP ozonolysis in presence of nitrogen oxides**

In this section we discuss the chemical composition of SOA from BCP ozonolysis in the presence of nitrogen oxides including NO₃ radicals, which refers to the SOA formed after the last addition of BCP, e.g. [180150-190160](#) min in Fig. 2. Please note that the results given here refer to the total SOA which was formed in two steps, first by pure ozonolysis and in a subsequent step including nitrogen oxides. The SOA mass formed in the presence of nitrogen oxides compared to the total SOA mass detected corresponds to 49% at 213K, 34% at 243K, 49% at 435 273K, 65% at 298K, and 63 % at 313K. Due to the excess of ozone besides NO₂, also NO₃ radicals and N₂O₅ were present. Thus, the BCP was now also oxidized by reaction with NO₃ radicals, with a major pathway to produce organonitrates (org-Ns) (Kiendler-Scharr et al., 2016; Wu et al., 2021). The concentrations of NO₃ radicals before the final addition of BCP were estimated for each experiment using a kinetic box model (details in Fig. [S40S13](#)) and the results are given in Table 1.

440 Based on the ozone and NO₃ concentration levels as well as the corresponding reaction rates, we estimated the fraction of NO₃ radicals contributing to the initial BCP oxidation to 84%, 90% and 72% for 273, 298 and 313K. [Please note that these values should be considered as upper limits due to other potential sinks for NO₃ radicals.](#)

3.4.1 Mass spectra from the gas-phase products

445 Figure 9 shows the averaged gas-phase mass spectra including molecules without nitrogen atom (org) and organonitrates (org-Ns) for all temperatures as detected by CIMS after the particle concentration got stable after the last BCP addition (e.g., at 298K, referring to the time period after 160 min in Fig. 2). The gas phase organonitrates showed an abundance increasing with temperature from 20.7% at 213K, 26.0% at 243K, 38.3% at 273K, 46.5% at 298K, to 48.9% at 313K. These compounds consisted of three groups of different carbon numbers (C₅H₇O₆N, C₁₀H₁₅O_{5.7}N and C₁₅H_{23.25}O_{6.8}N). To illustrate more clearly the organonitrate formation, we show the time evolution of the three most abundant org-Ns in these three groups in the right panel of Fig. 9. The signals of C₅H₇O₆N and C₁₀H₁₅O₆N increased after the NO₂ addition immediately, indicating that their formation was related to reactions between BCP oxidation products (BCP+O₃) and NO₃ radicals. After the start of the last BCP addition, C₁₅H₂₅O₇N started to increase significantly, indicating that its formation was linked to the reaction of BCP and 455 NO₃ directly, but not the reaction between the nitrate radicals and oxidation products from BCP ozonolysis.

3.4.2 Mass spectra from the particle phase

Figure 10 shows the particle chemical composition of SOA from BCP ozonolysis in the presence of NO₂ at 213 K (upper panel) and 313 K (lower panel). At 213K (upper panel), org dominated the particle composition, with a signal fraction of 97.1% to total signals (org-Ns + org), similar to the SOA before NO₂ addition at 213K. Org groups of monomers (41.2% in signal fraction), dimers (48.2%) and trimers (7.4%) were observed. The most abundant signals in each group were C₁₅H₂₄O₃, C₃₀H₄₈O₅ and C₄₄H₆₈O₉, respectively. The largest org-Ns signal was from C₁₅H₂₅O₇N, contributing 0.6% to the total signals. It is obvious that only few org-Ns were formed at 460

213K in our study, with a contribution ($[\text{org-Ns}]/[\text{total org.}]$) of 2.8% measured by FIGAERO-iodide-CIMS (normalised signal fraction, Fig. 12) and 0.08 detected by HR-AMS (Fig. S11S14).

465 Compared to the SOA chemical composition at 213K, substantially more org-Ns (48.9% to total signals) were formed at 313K. Monomeric org-Ns ($\text{C}_{15}\text{H}_{23,25}\text{O}_{6-10}\text{N}$) made up 35.5% of total signal (and the AMS measured 61% of org-Ns to total mass concentration of organic components (Fig. S11S14)). The most abundant monomeric org-Ns signals at 313K were $\text{C}_{15}\text{H}_{23,25}\text{O}_{7-9}\text{N}$, which had also been detected by (Wu et al., 2021). In our study, 51.1% of normalised signals were contributed by org species, with 28.3% of monomeric org ($\text{C}_{13-15}\text{H}_{20-24}\text{O}_{4-8}$), 16.6% of dimeric org ($\text{C}_{20}\text{H}_{22-24}\text{O}_{7-9}$) and 6.3% of trimeric org ($\text{C}_{35}\text{H}_{48}\text{O}_{10-13}$).

The particle mass spectra for all temperatures are shown in Fig. 11, with relative signal abundance of individual molecule to total detected species, including pure organic components (org, red) and organonitrates (org-Ns, blue).

~~The most abundant mass spectral peak of all organic compounds without nitrogen at 298-313K was not $\text{C}_{14}\text{H}_{22}\text{O}_7$ as for pure ozonolysis, but $\text{C}_{15}\text{H}_{24}\text{O}_4$ in the presence of NO_2 and NO_3 radicals. This is attributed to additional formation of pure MLOCs. Among all the org, the most abundant signal at 298-313K was not $\text{C}_{14}\text{H}_{22}\text{O}_7$ as for pure ozonolysis, but $\text{C}_{15}\text{H}_{24}\text{O}_4$ (NO_3 -present), due to the additional formation of pure MLOCs. This could be confirmed by comparison of the absolute signals of $\text{C}_{14}\text{H}_{22}\text{O}_7$ and $\text{C}_{15}\text{H}_{24}\text{O}_4$ in the particles before and after the last BCP addition (Fig. S152). Moreover, the dimeric compounds with 20 carbon atom skeletons (i.e. $\text{C}_{20}\text{H}_{24}\text{O}_{7-8}$, insertion in Fig. 10) and the trimeric compounds with 35 carbon atom skeletons (i.e. $\text{C}_{35}\text{H}_{48}\text{O}_{12}$, Fig. 10 and Fig. 11) were newly formed at higher temperatures ($> 273\text{K}$) in the presence of nitrogen oxides. However, it cannot be excluded that they could be formed by the reaction of the oxidation products from pure BCP ozonolysis with those formed in the presence of NO_2 and NO_3 radicals. In contrast, the mass spectra of non-N-containing organic species (org) at 213-243K showed no substantial changes compared to the species from ozonolysis without nitrogen oxides present (cf. Fig.7). One obvious reason for this may be the lower NO_3 radical concentrations at lower temperatures but also changes in the active reaction pathways may play a role. However, it cannot be excluded that they could be formed by reaction of the oxidation products from BCP ozonolysis (nitrogen oxides absent) and the next step (nitrogen oxides present). In contrast, the mass spectra of non-N-containing organic species (org) at 213-243K showed no substantial changes compared to the species from ozonolysis without nitrogen oxides present (cf. Fig. 7). One obvious reason for this is the lower NO_3 radical concentration at lower temperatures but also changes in the active reaction pathways may play a role.~~

In addition, more organonitrates were detected at increased temperatures, ranging from 2.8% (213K) to 51.5% (298K) and 48.9% (313K), dominated by monomeric org-Ns ($\text{C}_{15}\text{H}_{23,25}\text{O}_{6-10}\text{N}$), wherein the most abundant signals were from $\text{C}_{15}\text{H}_{25}\text{O}_7\text{N}$ at 243-273K and $\text{C}_{15}\text{H}_{23}\text{O}_9\text{N}$ at 298-313K. Monomeric organonitrates contributed 1.7% to 40.1% from 213 K to 298K, and 35.5% at 313K to total organic signals, as shown in Fig. 12. Besides, the signal-weighted averaged O:C of organonitrates monotonically increased from 0.41 to 0.51 from 213K to 298K, and 0.50 at 313K. We assume that the positive impact of temperature on the oxygenation and formation of organonitrates could also be relevant for the highest temperature (313K). However, the higher oxidized organonitrates may be out of the detection range of the FIGAERO-iodide-CIMS, resulting in less signal fractions and lower signal-weighted averaged O:C ratios of organonitrates at 313K. This is supported by HR-TOF-AMS measurements (Fig. S11S14), which show increasing organonitrate fractions from 8% to 61% also for 313 K. On the other hand, thermal instability of some N-containing compounds formed at 313K, e.g. peroxy nitrates (Francisco and Krylowski, 2005; Lee et al., 2016), can also be an explanation for the weaker increase of their

[fraction observed for 313K. For example, the potential contributions from thermal decomposition \(\$C_{1-13}H_yO_zN_1I\$ ions\) showed a positive temperature dependence. Also, abundant N-containing ions, such as low molecular-weight molecule \$C_5H_7O_6N_1\$, desorbed substantially between 120-200°C, and had an unexpected high \$T_{max}\$ of about 160°C.”](#)

It should be noted that the nitrate radical levels were estimated to be higher at higher temperatures (Table 1), which partially explains the increasing organonitrates formation at higher temperatures. The organonitrates detected in this study were dominantly with the same carbon atom skeleton (C_{15}) and only one organonitrate functional group ($-ONO_2$). The O:C ratios for organonitrates increase with temperature. Thus, it can be concluded that higher temperatures favour the formation of higher oxygenated organonitrates from the BCP [ozonolysisoxidation](#) in the presence of O_3 , NO_2 , and NO_3 radicals. The contribution in gas and particle phase of the major compounds are shown in Fig. [S13S16](#).

4 Summary and conclusions

In this work, a series of experiments conducted in the dark AIDA chamber with temperatures covering the whole tropospheric range (213-313K) were analysed to investigate the yields, kinetics and chemistry of BCP ozonolysis in the absence and presence of nitrogen oxides. The rate coefficient of the endocyclic double bond in BCP reacting with ozone was determined in the temperature range between 243 and 313K showing decreasing values with increasing temperature. The rate coefficients agree well with literature data at 296 K and with quantum chemical calculations of the temperature dependence. We determined a reaction enthalpy of (5.6 ± 1.0) KJ mol⁻¹ and OH radical yields increasing from (5 ± 2) % at 243K to (15 ± 2) % at 313K.

SOA yields determined for the ozonolysis of BCP show about 50% smaller values than literature data at 298 K, which is still within the combined uncertainty limits. For the first time the temperature dependence of the SOA yield was determined showing values decreasing from (37 ± 11) % to (16 ± 5) % for temperatures increasing from 243 to 313K and a constant organic particle mass of 10 $\mu\text{g m}^{-3}$. This allows to calculate the potential of BCP to contribute to SOA formation e.g. if reaching higher altitudes in the atmosphere.

The chemical characteristics of BCP pure organic species and organonitrates were determined by the FIGAERO-CIMS using I⁻ as the reagent ion. Major products and different chemical composition of gas and particle phase with and without NO_3 present at all temperatures were resolved. The variation of the ozonolysis temperature revealed substantial impact on the abundance of individual pure organic molecules without nitrogen atoms (org) and organonitrate molecules (org-Ns). In the first generated SOA without nitrogen oxides present, monomers (mainly $C_{14-15}H_{22-24}O_{3-7}$) and dimers (mainly $C_{28-30}H_{44-48}O_{5-9}$) were abundant, wherein minor signals of trimers (mainly $C_{41-44}H_{62-66}O_{9-11}$) were mainly detected at 213K. Potential dimer and trimer formation pathways are proposed. With temperature increasing to 313K, monomers and dimers ($C_{14-15}H_{22-24}O_{6-9}$ and $C_{27-29}H_{42-44}O_{9-11}$, respectively) became more oxidized, and no significant trimeric signals were detected.

The positive temperature dependence of the oxygenation of the products was also observed in the BCP organonitrates. In the presence of nitrogen oxides, most of the organonitrates were found as monomers with a C_{15} skeleton with one nitrate group, with their signal-weighted O:C ratio increasing from 0.41 to 0.51 for temperatures in the range between 213K and 313K. Dimeric and trimeric pure organic species without nitrogen atoms (C_{20} , C_{35}) were newly formed in the presence of nitrogen oxides at 298-313K, which substantially changed the chemical composition of pure organic components and indicates more termination ways might exist.

Please note that we cannot exclude completely that the changing humidity in our experiments has no influence on our results. Nonetheless, we think that the SOA composition we have observed is dominated by gas phase reactions with sufficient water concentrations compared to ambient conditions and that the potential impact of relative humidity on condensed phase reactions was minimized by increasing relative humidity with decreasing temperature {Li, 2019 #223}{Maclean, 2021 #224}. maybe except for the lowermost temperature (213 K).

545

This work helps to get a better understanding on the yields, kinetics and chemical composition of SOA from BCP ozonolysis over a relatively wide range of temperatures, 213 K to 313 K, which is representative of the real atmosphere from boundary layer to upper troposphere. In addition, SOA volatility is expected to have a strong impact on the SOA formation, and will be discussed in our upcoming paper focusing on the desorption of BCP SOA from filters but also the warming up of the aerosol in the simulation chamber overnight.

550

Data availability

The data on SOA yields and chemical composition from FIGAERO-iodide-CIMS is accessible at KIT open data (link/DOI, will be completed). Data are available upon request to the corresponding author.

Conflict of interests

555 The authors declare that they have no conflict of interest.

Author contributions

LG and HS designed the study. Chamber experiments were carried out by LG, HS, JS, FJ and MV. Data analysis and interpretation was performed by LG, HS, CM and WH. LG wrote the manuscript, with input from JS and HS. All co-authors commented on the manuscript.

Acknowledgement

560 LG, JS and FJ acknowledge funding by the China Scholarship Council (CSC). This work was supported by the Helmholtz Association within the project MOSES (Modular Observation Solutions for Earth Systems) and H2020 European Research Council (CHAPAs (grant no. 850614). Much Acknowledge also goes to all the technicians in IMK-AAF in KIT for their support on this work.

References

Agelopoulos, N. G., Chamberlain, K., and Pickett, J. A.: Factors Affecting Volatile Emissions of Intact Potato Plants, *Solanum tuberosum*: Variability of Quantities and Stability of Ratios, *Journal of Chemical Ecology*, 26, 497-511, 10.1023/A:1005473825335, 2000.

570 Alfarra, M., Hamilton, J., Wyche, K., Good, N., Ward, M., Carr, T., Barley, M., Monks, P., Jenkin, M., and Lewis, A.: The effect of photochemical ageing and initial precursor concentration on the composition and hygroscopic

- properties of β -caryophyllene secondary organic aerosol, *Atmospheric Chemistry and Physics*, 12, 6417-6436, 2012.
- Andreae, M. O., Afchine, A., Albrecht, R., Holanda, B. A., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Cecchini, M. A., Costa, A., Dollner, M., Fütterer, D., Järvinen, E., Jurkat, T., Klimach, T., Konemann, T., Knote, C., Krämer, M., Krisna, T., Machado, L. A. T., Mertes, S., Minikin, A., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Sauer, D., Schlager, H., Schnaiter, M., Schneider, J., Schulz, C., Spanu, A., Sperling, V. B., Voigt, C., Walser, A., Wang, J., Weinzierl, B., Wendisch, M., and Ziereis, H.: Aerosol characteristics and particle production in the upper troposphere over the Amazon Basin, *Atmos. Chem. Phys.*, 18, 921-961, 10.5194/acp-18-921-2018, 2018.
- 575
- Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crouse, J. D., Wennberg, P. O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A., Donahue, N., Kjaergaard, H. G., and Ehn, M.: Highly Oxygenated Organic Molecules (HOM) from Gas-Phase Autoxidation Involving Peroxy Radicals: A Key Contributor to Atmospheric Aerosol, *Chemical reviews*, 119, 3472-3509, 10.1021/acs.chemrev.8b00395, 2019.
- 580
- Calogirou, A., Larsen, B., and Kotzias, D.: Gas-phase terpene oxidation products: a review, *Atmospheric Environment*, 33, 1423-1439, 1999.
- 585
- Calvert, J. G., Derwent, R. G., Orlando, J. J., Wallington, T. J., and Tyndall, G. S.: Mechanisms of atmospheric oxidation of the alkanes, 2008.
- Chan, A., Kroll, J., Ng, N., and Seinfeld, J.: Kinetic modeling of secondary organic aerosol formation: effects of particle-and gas-phase reactions of semivolatile products, *Atmospheric Chemistry and Physics*, 7, 4135-4147, 2007.
- 590
- Chan, M., Surratt, J., Chan, A., Schilling, K., Offenberg, J., Lewandowski, M., Edney, E., Kleindienst, T., Jaoui, M., and Edgerton, E.: Influence of aerosol acidity on the chemical composition of secondary organic aerosol from β -caryophyllene, *Atmospheric Chemistry and Physics*, 11, 1735-1751, 2011.
- Charnawskas, J. C., Alpert, P. A., Lambe, A. T., Berkemeier, T., O'Brien, R. E., Massoli, P., Onasch, T. B., Shiraiwa, M., Moffet, R. C., and Gilles, M. K.: Condensed-phase biogenic–anthropogenic interactions with implications for cold cloud formation, *Faraday discussions*, 200, 165-194, 2017.
- 595
- Chen, Q., Li, Y., McKinney, K., Kuwata, M., and Martin, S.: Particle mass yield from beta-caryophyllene ozonolysis, *Atmospheric Chemistry and Physics*, 12, 3165-3179, 2012.
- Cheng, Y., Ma, Y., and Hu, D.: Tracer-based source apportioning of atmospheric organic carbon and the influence of anthropogenic emissions on secondary organic aerosol formation in Hong Kong, *Atmos. Chem. Phys.*, 21, 10589-10608, 10.5194/acp-21-10589-2021, 2021.
- 600
- Ciccioli, P., Brancaleoni, E., Frattoni, M., Di Palo, V., Valentini, R., Tirone, G., Seufert, G., Bertin, N., Hansen, U., and Csiky, O.: Emission of reactive terpene compounds from orange orchards and their removal by within-canopy processes, *Journal of Geophysical Research: Atmospheres*, 104, 8077-8094, 1999.

- 605 Clark, C. H., Kacarab, M., Nakao, S., Asa-Awuku, A., Sato, K., and Cocker III, D. R.: Temperature effects on secondary organic aerosol (SOA) from the dark ozonolysis and photo-oxidation of isoprene, *Environmental Science & Technology*, 50, 5564-5571, 2016.
- Dekermenjian, M.: FTIR analysis of aerosol formed in the ozone oxidation of sesquiterpenes, *Aerosol Science & Technology*, 30, 349-363, 1999.
- 610 Duhl, T. R., Helmig, D., and Guenther, A.: Sesquiterpene emissions from vegetation: a review, *Biogeosciences*, 5, 761-777, 10.5194/bg-5-761-2008, 2008.
- Fahey, D. W., Gao, R. S., Möhler, O., Saathoff, H., Schiller, C., Ebert, V., Krämer, M., Peter, T., Amarouche, N., Avallone, L. M., Bauer, R., Bozóki, Z., Christensen, L. E., Davis, S. M., Durrý, G., Dyroff, C., Herman, R. L., Hunsmann, S., Khaykin, S. M., Mackrodt, P., Meyer, J., Smith, J. B., Spelten, N., Troy, R. F., Vömel, H., Wagner, S., and Wienhold, F. G.: The AquaVIT-1 intercomparison of atmospheric water vapor measurement techniques, *Atmos. Meas. Tech.*, 7, 3177-3213, 10.5194/amt-7-3177-2014, 2014.
- 615 Faiola, C. L., Buchholz, A., Kari, E., Yli-Pirilä, P., Holopainen, J. K., Kivimäenpää, M., Miettinen, P., Worsnop, D. R., Lehtinen, K. E. J., and Guenther, A. B.: Terpene composition complexity controls secondary organic aerosol yields from scots pine volatile emissions, *Scientific Reports*, 8, 1-13, 2018.
- 620 Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A. B., Hewitt, C. N., Lamb, B., Liu, S., Trainer, M., and Westberg, H.: Emissions of volatile organic compounds from vegetation and the implications for atmospheric chemistry, *Global Biogeochemical Cycles*, 6, 389-430, 1992.
- Geron, C. D., and Arnts, R. R.: Seasonal monoterpene and sesquiterpene emissions from *Pinus taeda* and *Pinus virginiana*, *Atmospheric Environment*, 44, 4240-4251, 2010.
- 625 Gkatzelis, G. I., Tillmann, R., Hohaus, T., Müller, M., Eichler, P., Xu, K. M., Schlag, P., Schmitt, S. H., Wegener, R., Kaminski, M., Holzinger, R., Wisthaler, A., and Kiendler-Scharr, A.: Comparison of three aerosol chemical characterization techniques utilizing PTR-ToF-MS: a study on freshly formed and aged biogenic SOA, *Atmos. Meas. Tech.*, 11, 1481-1500, 10.5194/amt-11-1481-2018, 2018.
- 630 Griffin, R. J., Cocker III, D. R., Flagan, R. C., and Seinfeld, J. H.: Organic aerosol formation from the oxidation of biogenic hydrocarbons, *Journal of Geophysical Research: Atmospheres*, 104, 3555-3567, 1999.
- Grosjean, D., Williams, E. L., Grosjean, E., Andino, J. M., and Seinfeld, J. H.: Atmospheric oxidation of biogenic hydrocarbons: reaction of ozone with β -pinene, D-limonene and trans-caryophyllene, *Environmental Science & Technology*, 27, 2754-2758, 1993.
- 635 Hansen, U., and Seufert, G.: Temperature and light dependence of β -caryophyllene emission rates, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Haque, M. M., Kawamura, K., and Kim, Y.: Seasonal variations of biogenic secondary organic aerosol tracers in ambient aerosols from Alaska, *Atmospheric Environment*, 130, 95-104, 2016.

- 640 Hu, D., Bian, Q., Li, T. W., Lau, A. K., and Yu, J. Z.: Contributions of isoprene, monoterpenes, β -caryophyllene, and toluene to secondary organic aerosols in Hong Kong during the summer of 2006, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Huang, W., Saathoff, H., Pajunoja, A., Shen, X., Naumann, K. H., Wagner, R., Virtanen, A., Leisner, T., and Mohr, C.: α -Pinene secondary organic aerosol at low temperature: chemical composition and implications for particle viscosity, *Atmos. Chem. Phys.*, 18, 2883-2898, 10.5194/acp-18-2883-2018, 2018.
- 645 Huang, W., Saathoff, H., Shen, X., Ramisetty, R., Leisner, T., and Mohr, C.: Seasonal characteristics of organic aerosol chemical composition and volatility in Stuttgart, Germany, *Atmos. Chem. Phys.*, 19, 11687-11700, 10.5194/acp-19-11687-2019, 2019a.
- Huang, W., Saathoff, H., Shen, X., Ramisetty, R., Leisner, T., and Mohr, C.: Chemical Characterization of Highly Functionalized Organonitrates Contributing to Night-Time Organic Aerosol Mass Loadings and Particle Growth, *Environmental Science & Technology*, 53, 1165-1174, 10.1021/acs.est.8b05826, 2019b.
- 650 Huang, W., Li, H., Sarnela, N., Heikkinen, L., Tham, Y. J., Mikkilä, J., Thomas, S. J., Donahue, N. M., Kulmala, M., and Bianchi, F.: Measurement report: Molecular composition and volatility of gaseous organic compounds in a boreal forest – from volatile organic compounds to highly oxygenated organic molecules, *Atmos. Chem. Phys.*, 21, 8961-8977, 10.5194/acp-21-8961-2021, 2021.
- 655 Jaoui, M., Leungsakul, S., and Kamens, R.: Gas and particle products distribution from the reaction of β -caryophyllene with ozone, *Journal of Atmospheric Chemistry*, 45, 261-287, 2003.
- Jaoui, M., Lewandowski, M., Kleindienst, T. E., Offenberg, J. H., and Edney, E. O.: β -caryophyllinic acid: An atmospheric tracer for β -caryophyllene secondary organic aerosol, *Geophysical Research Letters*, 34, 2007.
- 660 Jaoui, M., Kleindienst, T. E., Docherty, K. S., Lewandowski, M., and Offenberg, J. H.: Secondary organic aerosol formation from the oxidation of a series of sesquiterpenes: α -cedrene, β -caryophyllene, α -humulene and α -farnesene with O₃, OH and NO₃ radicals, *Environmental Chemistry*, 10, 178-193, 2013.
- Jardine, K., Yañez Serrano, A., Arneeth, A., Abrell, L., Jardine, A., Van Haren, J., Artaxo, P., Rizzo, L. V., Ishida, F. Y., and Karl, T.: Within-canopy sesquiterpene ozonolysis in Amazonia, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- 665 Jenkin, M., Wyche, K., Evans, C., Carr, T., Monks, P., Alfarra, M., Barley, M., McFiggans, G., Young, J., and Rickard, A.: Development and chamber evaluation of the MCM v3. 2 degradation scheme for β -caryophyllene, *Atmospheric Chemistry & Physics Discussions*, 12, 2012.
- Jokinen, T., Kausiala, O., Garmash, O., Peräkylä, O., Junninen, H., Schobesberger, S., Yan, C., Sipilä, M., and Rissanen, M. P.: Production of highly oxidized organic compounds from ozonolysis of beta-caryophyllene: laboratory and field measurements, *Boreal Environment Research*, 2016.
- 670 Kammer, J., Flaud, P. M., Chazeaubeny, A., Ciuraru, R., Le Menach, K., Geneste, E., Budzinski, H., Bonnefond, J. M., Lamaud, E., Perraudin, E., and Villenave, E.: Biogenic volatile organic compounds (BVOCs) reactivity

- related to new particle formation (NPF) over the Landes forest, *Atmospheric Research*, 237, 104869, <https://doi.org/10.1016/j.atmosres.2020.104869>, 2020.
- 675 Kari, E., Miettinen, P., Yli-Pirilä, P., Virtanen, A., and Faiola, C. L.: PTR-ToF-MS product ion distributions and humidity-dependence of biogenic volatile organic compounds, *International Journal of Mass Spectrometry*, 430, 87-97, <https://doi.org/10.1016/j.ijms.2018.05.003>, 2018.
- 680 Kiendler-Scharr, A., Mensah, A. A., Friese, E., Topping, D., Nemitz, E., Prevot, A. S. H., Äijälä, M., Allan, J., Canonaco, F., Canagaratna, M., Carbone, S., Crippa, M., Dall'Osto, M., Day, D. A., De Carlo, P., Di Marco, C. F., Elbern, H., Eriksson, A., Freney, E., Hao, L., Herrmann, H., Hildebrandt, L., Hillamo, R., Jimenez, J. L., Laaksonen, A., McFiggans, G., Mohr, C., O'Dowd, C., Otjes, R., Ovadnevaite, J., Pandis, S. N., Poulain, L., Schlag, P., Sellegri, K., Swietlicki, E., Tiitta, P., Vermeulen, A., Wahner, A., Worsnop, D., and Wu, H.-C.: Ubiquity of organic nitrates from nighttime chemistry in the European submicron aerosol, *Geophysical Research Letters*, 43, 7735-7744, <https://doi.org/10.1002/2016GL069239>, 2016.
- 685 Kim, S., Karl, T., Helmig, D., Daly, R., Rasmussen, R., and Guenther, A.: Measurement of atmospheric sesquiterpenes by proton transfer reaction-mass spectrometry (PTR-MS), *Atmos. Meas. Tech.*, 2, 99-112, 10.5194/amt-2-99-2009, 2009.
- 690 Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521-526, 10.1038/nature17953, 2016.
- 700 Kivimäenpää, M., Babalola, A. B., Joutsensaari, J., and Holopainen, J. K.: Methyl Salicylate and Sesquiterpene Emissions Are Indicative for Aphid Infestation on Scots Pine, *Forests*, 11, 573, 2020.
- 705 Kleist, E., Mentel, T., Andres, S., Bohne, A., Folkers, A., Kiendler-Scharr, A., Rudich, Y., Springer, M., Tillmann, R., and Wildt, J.: Irreversible impacts of heat on the emissions of monoterpenes, sesquiterpenes, phenolic BVOC and green leaf volatiles from several tree species, *Biogeosciences*, 9, 5111, 2012.
- Kristensen, K., Watne, Å. K., Hammes, J., Lutz, A., Petäjä, T., Hallquist, M., Bilde, M., and Glasius, M.: High-molecular weight dimer esters are major products in aerosols from α -pinene ozonolysis and the boreal forest, *Environmental Science & Technology Letters*, 3, 280-285, 2016.
- 705 Laothawornkitkul, J., Taylor, J. E., Paul, N. D., and Hewitt, C. N.: Biogenic volatile organic compounds in the Earth system, *New Phytologist*, 183, 27-51, 2009.


- Larsen, B., Lahaniati, M., Calogirou, A., and Kotzias, D.: Atmospheric oxidation products of terpenes: A new nomenclature, *Chemosphere*, 37, 1207-1220, 1998.
- 710 Lee, A., Goldstein, A. H., Keywood, M. D., Gao, S., Varutbangkul, V., Bahreini, R., Ng, N. L., Flagan, R. C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the ozonolysis of ten different terpenes, *Journal of Geophysical Research: Atmospheres*, 111, 2006a.
- Lee, A., Goldstein, A. H., Kroll, J. H., Ng, N. L., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the photooxidation of 16 different terpenes, *Journal of Geophysical*
715 *Research: Atmospheres*, 111, 2006b.
- Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Kurtén, T., Worsnop, D. R., and Thornton, J. A.: An iodide-adduct high-resolution time-of-flight chemical-ionization mass spectrometer: Application to atmospheric inorganic and organic compounds, *Environmental science & technology*, 48, 6309-6317, 2014.
- Lee, S., and Kamens, R. M.: Particle nucleation from the reaction of α -pinene and O₃, *Atmospheric Environment*,
720 39, 6822-6832, 2005.
- Li, S.-J., Yuan, X.-Y., Li, Q., and Feng, Z.-Z.: Inventory and Characteristics of Biogenic Volatile Organic Compounds (BVOCs) for 12 Deciduous Fruit Trees, *Huan jing ke xue= Huanjing kexue*, 40, 2078-2085, 2019.
- Li, Y., Chen, Q., Guzman, M., Chan, C. K., and Martin, S.: Second-generation products contribute substantially to the particle-phase organic material produced by [β]-caryophyllene ozonolysis, *Atmospheric Chemistry and*
725 *Physics*, 11, 121, 2011.
- Liu, P. S. K., Deng, R., Smith, K. A., Williams, L. R., Jayne, J. T., Canagaratna, M. R., Moore, K., Onasch, T. B., Worsnop, D. R., and Deshler, T.: Transmission Efficiency of an Aerodynamic Focusing Lens System: Comparison of Model Calculations and Laboratory Measurements for the Aerodyne Aerosol Mass Spectrometer, *Aerosol Science and Technology*, 41, 721-733, 10.1080/02786820701422278, 2007.
- 730 Lopez-Hilfiker, F. D., Mohr, C., Ehn, M., Rubach, F., Kleist, E., Wildt, J., Mentel, T. F., Lutz, A., Hallquist, M., Worsnop, D., and Thornton, J. A.: A novel method for online analysis of gas and particle composition: description and evaluation of a Filter Inlet for Gases and AEROsols (FIGAERO), *Atmos. Meas. Tech.*, 7, 983-1001, 10.5194/amt-7-983-2014, 2014.
- Lopez-Hilfiker, F. D., Iyer, S., Mohr, C., Lee, B. H., D'Ambro, E. L., Kurtén, T., and Thornton, J. A.: Constraining the sensitivity of iodide adduct chemical ionization mass spectrometry to multifunctional organic molecules using the collision limit and thermodynamic stability of iodide ion adducts, *Atmos. Meas. Tech.*, 9, 1505-1512, 10.5194/amt-9-1505-2016, 2016.
- 735
- Matsunaga, S. N., Niwa, S., Mochizuki, T., Tani, A., Kusumoto, D., Utsumi, Y., Enoki, T., and Hiura, T.: Seasonal variation in basal emission rates and composition of mono- and sesquiterpenes emitted from dominant conifers in
740 Japan, *Atmospheric Environment*, 69, 124-130, 2013.

- Mellouki, A., Wallington, T., and Chen, J.: Atmospheric chemistry of oxygenated volatile organic compounds: impacts on air quality and climate, *Chemical reviews*, 115, 3984-4014, 2015.
- Möhler, O., Stetzer, O., Schaefers, S., Linke, C., Schnaiter, M., Tiede, R., Saathoff, H., Krämer, M., Mangold, A., Budz, P., Zink, P., Schreiner, J., Mauersberger, K., Haag, W., Kärcher, B., and Schurath, U.: Experimental investigation of homogeneous freezing of sulphuric acid particles in the aerosol chamber AIDA, *Atmos. Chem. Phys.*, 3, 211-223, 10.5194/acp-3-211-2003, 2003.
- Müller, L., Reinnig, M. C., Warnke, J., and Hoffmann, T.: Unambiguous identification of esters as oligomers in secondary organic aerosol formed from cyclohexene and cyclohexene/ α -pinene ozonolysis, *Atmos. Chem. Phys.*, 8, 1423-1433, 10.5194/acp-8-1423-2008, 2008.
- 750 Müller, M., Eichler, P., D'Anna, B., Tan, W., and Wisthaler, A.: Direct Sampling and Analysis of Atmospheric Particulate Organic Matter by Proton-Transfer-Reaction Mass Spectrometry, *Analytical Chemistry*, 89, 10889-10897, 10.1021/acs.analchem.7b02582, 2017.
- Naumann, K.-H.: COSIMA—a computer program simulating the dynamics of fractal aerosols, *Journal of Aerosol Science*, 34, 1371-1397, [https://doi.org/10.1016/S0021-8502\(03\)00367-7](https://doi.org/10.1016/S0021-8502(03)00367-7), 2003.
- 755 Ng, N. L., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Kroll, J. H., Kwan, A. J., McCabe, D. C., Wennberg, P. O., Sorooshian, A., and Murphy, S. M.: Effect of NO_x level on secondary organic aerosol (SOA) formation from the photooxidation of terpenes, 2007.
- Nguyen, T. L., Winterhalter, R., Moortgat, G., Kanawati, B., Peeters, J., and Vereecken, L.: The gas-phase ozonolysis of β -caryophyllene (C₁₅H₂₄). Part II: A theoretical study, *Physical Chemistry Chemical Physics*, 11, 4173-4183, 2009.
- 760 Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H.: Gas/Particle Partitioning and Secondary Organic Aerosol Yields, *Environmental Science & Technology*, 30, 2580-2585, 10.1021/es950943+, 1996.
- Parshintsev, J., Nurmi, J., Kilpeläinen, I., Hartonen, K., Kulmala, M., and Riekkola, M.-L.: Preparation of β -caryophyllene oxidation products and their determination in ambient aerosol samples, *Analytical and bioanalytical chemistry*, 390, 913-919, 2008.
- 765 Peng, Z., and Jimenez, J. L.: KinSim: A Research-Grade, User-Friendly, Visual Kinetics Simulator for Chemical-Kinetics and Environmental-Chemistry Teaching, in, ACS Publications, 2019.
- Piel, F., Müller, M., Winkler, K., Skytte af Sättra, J., and Wisthaler, A.: Introducing the extended volatility range proton-transfer-reaction mass spectrometer (EVR PTR-MS), *Atmospheric Measurement Techniques*, 14, 1355-1363, 2021.
- 770 Richters, S., Herrmann, H., and Berndt, T.: Gas-phase rate coefficients of the reaction of ozone with four sesquiterpenes at 295 ± 2 K, *Physical Chemistry Chemical Physics*, 17, 11658-11669, 10.1039/C4CP05542J, 2015.


- 775 Richters, S., Herrmann, H., and Berndt, T.: Different pathways of the formation of highly oxidized multifunctional organic compounds (HOMs) from the gas-phase ozonolysis of β -caryophyllene, *Atmos. Chem. Phys.*, 16, 9831-9845, 2016.
- 780 Riva, M., Rantala, P., Krechmer, J. E., Peräkylä, O., Zhang, Y., Heikkinen, L., Garmash, O., Yan, C., Kulmala, M., Worsnop, D., and Ehn, M.: Evaluating the performance of five different chemical ionization techniques for detecting gaseous oxygenated organic species, *Atmos. Meas. Tech.*, 12, 2403-2421, 10.5194/amt-12-2403-2019, 2019.
- Rodriguez-Saona, C., Crafts-Brandner, S. J., ParÉ, P. W., and Henneberry, T. J.: EXOGENOUS METHYL JASMONATE INDUCES VOLATILE EMISSIONS IN COTTON PLANTS, *Journal of Chemical Ecology*, 27, 679-695, 10.1023/A:1010393700918, 2001.
- 785 Saathoff, H., Naumann, K., Möhler, O., Jonsson, A., Hallquist, M., Kiendler-Scharr, A., Mentel, T. F., Tillmann, R., and Schurath, U.: Temperature dependence of yields of secondary organic aerosols from the ozonolysis of α -pinene and limonene, *Atmos. Chem. Phys.*, 9, 1551-1577, 2009.
- 790 Shrivastava, M., Andreae, M. O., Artaxo, P., Barbosa, H. M. J., Berg, L. K., Brito, J., Ching, J., Easter, R. C., Fan, J., Fast, J. D., Feng, Z., Fuentes, J. D., Glasius, M., Goldstein, A. H., Alves, E. G., Gomes, H., Gu, D., Guenther, A., Jathar, S. H., Kim, S., Liu, Y., Lou, S., Martin, S. T., McNeill, V. F., Medeiros, A., de Sá, S. S., Shilling, J. E., Springston, S. R., Souza, R. A. F., Thornton, J. A., Isaacman-VanWertz, G., Yee, L. D., Ynoue, R., Zaveri, R. A., Zelenyuk, A., and Zhao, C.: Urban pollution greatly enhances formation of natural aerosols over the Amazon rainforest, *Nature Communications*, 10, 1046, 10.1038/s41467-019-08909-4, 2019.
- 795 Shu, Y., and Atkinson, R.: Rate constants for the gas-phase reactions of O₃ with a series of terpenes and OH radical formation from the O₃ reactions with sesquiterpenes at 296±2 K, *International Journal of Chemical Kinetics*, 26, 1193-1205, 1994.
- Shu, Y., and Atkinson, R.: Atmospheric lifetimes and fates of a series of sesquiterpenes, *Journal of Geophysical Research: Atmospheres*, 100, 7275-7281, <https://doi.org/10.1029/95JD00368>, 1995.
- 800 Stolzenburg, D., Fischer, L., Vogel, A. L., Heinritzi, M., Schervish, M., Simon, M., Wagner, A. C., Dada, L., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bianchi, F., Breitenlechner, M., Brilke, S., Buenrostro Mazon, S., Chen, D., Dias, A., Draper, D. C., Duplissy, J., El Haddad, I., Finkenzeller, H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., He, X., Helm, J., Hofbauer, V., Hoyle, C. R., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lampilahti, J., Lawler, M., Lehtipalo, K., Leiminger, M., Mai, H., Mathot, S., Mentler, B., Molteni, U., Nie, W., Nieminen, T., Nowak, J. B., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Quéléver, L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S., Tauber, C., Tomé, A., Wagner, R., Wang, M., 805 Weitz, L., Wimmer, D., Xiao, M., Yan, C., Ye, P., Zha, Q., Baltensperger, U., Curtius, J., Dommen, J., Flagan, R. C., Kulmala, M., Smith, J. N., Worsnop, D. R., Hansel, A., Donahue, N. M., and Winkler, P. M.: Rapid growth of organic aerosol nanoparticles over a wide tropospheric temperature range, *Proceedings of the National Academy of Sciences*, 115, 9122-9127, 10.1073/pnas.1807604115, 2018.

- 810 Tarvainen, V., Hakola, H., Hellén, H., Bäck, J., Hari, P., and Kulmala, M.: Temperature and light dependence of the VOC emissions of Scots pine, *Atmos. Chem. Phys.*, 5, 989-998, 10.5194/acp-5-989-2005, 2005.
- van Eijck, A., Opatz, T., Taraborrelli, D., Sander, R., and Hoffmann, T.: New tracer compounds for secondary organic aerosol formation from β -caryophyllene oxidation, *Atmospheric Environment*, 80, 122-130, 2013.
- 815 Verma, S. K., Kawamura, K., Deshmukh, D. K., Haque, M. M., and Pavuluri, C. M.: Seasonal Characteristics of Biogenic Secondary Organic Aerosols Over Chichijima Island in the Western North Pacific: Impact of Biomass Burning Activity in East Asia, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD032987, <https://doi.org/10.1029/2020JD032987>, 2021.
- Von Hessberg, C., Von Hessberg, P., Pöschl, U., Bilde, M., Nielsen, O., and Moortgat, G.: Temperature and humidity dependence of secondary organic aerosol yield from the ozonolysis of β -pinene, *Atmos. Chem. Phys.*, 9, 3583-3599, 2009.
- 820 Wagner, R., Bunz, H., Linke, C., Möhler, O., Naumann, K.-H., Saathoff, H., Schnaiter, M., and Schurath, U.: Chamber simulations of cloud chemistry: the AIDA Chamber, in: *Environmental simulation chambers: application to atmospheric chemical processes*, Springer, 67-82, 2006.
- Wang, M., Yao, L., Zheng, J., Wang, X., Chen, J., Yang, X., Worsnop, D. R., Donahue, N. M., and Wang, L.: Reactions of atmospheric particulate stabilized Criegee intermediates lead to high-molecular-weight aerosol components, *Environmental science & technology*, 50, 5702-5710, 2016.
- 825 Weikl, F., Ghirardo, A., Schnitzler, J.-P., and Pritsch, K.: Sesquiterpene emissions from *Alternaria alternata* and *Fusarium oxysporum*: effects of age, nutrient availability, and co-cultivation, *Scientific reports*, 6, 22152, 2016.
- Williams, L. R., Gonzalez, L. A., Peck, J., Trimborn, D., McInnis, J., Farrar, M. R., Moore, K. D., Jayne, J. T., Robinson, W. A., Lewis, D. K., Onasch, T. B., Canagaratna, M. R., Trimborn, A., Timko, M. T., Magoon, G., 830 Deng, R., Tang, D., de la Rosa Blanco, E., Prévôt, A. S. H., Smith, K. A., and Worsnop, D. R.: Characterization of an aerodynamic lens for transmitting particles greater than 1 micrometer in diameter into the Aerodyne aerosol mass spectrometer, *Atmos. Meas. Tech.*, 6, 3271-3280, 10.5194/amt-6-3271-2013, 2013.
- Winterhalter, R., Herrmann, F., Kanawati, B., Nguyen, T. L., Peeters, J., Vereecken, L., and Moortgat, G. K.: The gas-phase ozonolysis of β -caryophyllene (C₁₅H₂₄). Part I: an experimental study, *Physical Chemistry Chemical Physics*, 11, 4152-4172, 2009.
- 835 Witkowski, B., and Gierczak, T.: Early stage composition of SOA produced by α -pinene/ozone reaction: α -Acyloxyhydroperoxy aldehydes and acidic dimers, *Atmospheric Environment*, 95, 59-70, 2014.
- Wu, C., Bell, D. M., Graham, E. L., Haslett, S., Riipinen, I., Baltensperger, U., Bertrand, A., Giannoukos, S., Schoonbaert, J., El Haddad, I., Prevot, A. S. H., Huang, W., and Mohr, C.: Photolytically induced changes in composition and volatility of biogenic secondary organic aerosol from nitrate radical oxidation during night-to-day transition, *Atmos. Chem. Phys.*, 21, 14907-14925, 10.5194/acp-21-14907-2021, 2021.
- 840


- 845 Yasmeeen, F., Vermeulen, R., Szmigielski, R., Iinuma, Y., Böge, O., Herrmann, H., Maenhaut, W., and Claeys, M.: Terpenylic acid and related compounds: precursors for dimers in secondary organic aerosol from the ozonolysis of α - and β -pinene, *Atmos. Chem. Phys.*, 10, 9383-9392, 10.5194/acp-10-9383-2010, 2010.
- 850 Ye, Q., Wang, M., Hofbauer, V., Stolzenburg, D., Chen, D., Schervish, M., Vogel, A., Mauldin, R. L., Baalbaki, R., Brilke, S., Dada, L., Dias, A., Duplissy, J., El Haddad, I., Finkenzeller, H., Fischer, L., He, X., Kim, C., Kürten, A., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Manninen, H. E., Marten, R., Mentler, B., Partoll, E., Petäjä, T., Rissanen, M., Schobesberger, S., Schuchmann, S., Simon, M., Tham, Y. J., Vazquez-Pufleau, M., Wagner, A. C., Wang, Y., Wu, Y., Xiao, M., Baltensperger, U., Curtius, J., Flagan, R., Kirkby, J., Kulmala, M., Volkamer, R., Winkler, P. M., Worsnop, D., and Donahue, N. M.: Molecular Composition and Volatility of Nucleated Particles from α -Pinene Oxidation between $-50\text{ }^{\circ}\text{C}$ and $+25\text{ }^{\circ}\text{C}$, *Environmental Science & Technology*, 53, 12357-12365, 10.1021/acs.est.9b03265, 2019.
- 855 Zhao, Y., Zhang, R., Sun, X., He, M., Wang, H., Zhang, Q., and Ru, M.: Theoretical study on mechanism for O₃-initiated atmospheric oxidation reaction of β -caryophyllene, *Journal of Molecular Structure: THEOCHEM*, 947, 68-75, 2010.



860 Fig. 1. Schematic of the AIDA simulation chamber and its instrumentation for this study.



865 Fig. 2. Evolution of trace gases as well as particle mass & size for β -caryophyllene ozonolysis at 298K first without and then in presence of NO_2 . The time axis is relative to the first addition of ozone. Top panel shows the concentration of ozone (blue) and NO_2 (yellow) as well as the experiment steps; middle panel shows the change of β -caryophyllene (pink), particle concentration from SMPS (blue dots) and COSIMA model (black line), AMS (green dots) and CIMS (red triangle); bottom panel shows the particle size distribution over the course of the whole experiment.



870

Fig. 3. Arrhenius plot of the rate coefficients determined for the reaction of ozone with the endocyclic double bond of β -caryophyllene (blue diamonds) compared to the values measured by Shu and Atkinson et al., 1994 (green circle) and Richter et al., 2016, at 296 K (grey triangle) as well as model calculations by Nguyen et al., 2009 (yellow dashed line). The blue dashed line is a fit to the rate coefficients determined in this work.

875




Fig. 4. SOA mass yield from β -caryophyllene ozonolysis at a constant total organic aerosol mass (M_{org}) of $10 \mu\text{g m}^{-3}$ for temperatures between 243 – 313 K from this study in comparison with literature data. The dashed line represents a single exponential fit to the data. Chen et al., 2012 used OH/Cl scavenger and ammonium sulfate as seeds; Tasoglou and Pandis, 2015 used OH/Cl scavenger. Another comparison with literature data is given in Fig. [S4S6](#).

880




Fig. 5. Averaged CIMS gas phase mass spectra (background subtracted) for all temperatures, indicating the gas information when the SOA got stable in the absence of nitrogen oxides. The particle information sampled at the same time is given in Fig. 6 and Fig. 7.

885




Fig. 6. CIMS mass spectra of particle phase compounds obtained from ozonolysis of β -caryophyllene without NO_2 at 213K (upper panel) and 313K (lower panel). Colours refer to monomeric (brown), dimeric (blue) and trimeric (purple) compounds, labels represent the assignment of individual molecules.

890




Fig. 7. CIMS mass spectra for particle phase compounds for all five temperatures (213 - 313K). Colours refer to monomeric (brown), dimeric (blue) and trimeric (purple) compounds. The table lists the signal-weighted bulk O:C ratios for each temperature. The mass to charge ratios in x-axis are subtracted by I⁻ ($m/z = 126.9$).

895






Fig. 8. Product groups contributing to SOA from ozonolysis of β -caryophyllene without NO_2 at temperatures between 213 – 313 K. Autoxidation-products calculation includes $\text{C}_{14}\text{H}_{22}\text{O}_{7,9}$, $\text{C}_{15}\text{H}_{22}\text{O}_{7,9}$ and $\text{C}_{15}\text{H}_{24}\text{O}_{7,9}$ identified in previous studies (Jokinen et al., 2016; Richters et al., 2016).

900




905 | Fig. 9. CIMS gas phase mass spectra for all temperatures in the presence of NO_2 (a). These spectra correspond to the same time as the particle phase spectra shown in Fig. 1210 and Fig. 11. Colours refer to compounds without nitrogen atoms (black) and compounds with one nitrogen atom (blue). The I- ($m/z = 126.9$) was subtracted and here the mass/charge refers to the weight of the molecules. The panel (b) shows the FIGAERO-iodide-CIMS measurement for the signal change of three typical organonitrate molecules for C5 ($\text{C}_5\text{H}_7\text{O}_6\text{N}$, red line), C10 ($\text{C}_{10}\text{H}_{15}\text{O}_6\text{N}$, blue line) and C15 ($\text{C}_{15}\text{H}_{25}\text{O}_7\text{N}$, black line) corresponding to the evolution starting after NO_2 addition at 298K.




910

Fig. 10. CIMS mass spectra of particle phase compounds obtained from ozonolysis of β -caryophyllene in presence of nitrogen oxides at 213K (upper panel) and 313K (lower panel). Colours refer to compounds without nitrogen atoms (red) and compounds with one nitrogen atom (blue).



915

Fig. 11. CIMS mass spectra for particle phase compounds for all five temperatures (213 – 313 K). Colours refer to C_xH_yO_zN₀ (brown) and C_xH_yO_zN₁ (blue) compounds. The table lists the average O:C ratios for each temperature.



920 Fig. 12. Product groups (monomeric org, dimeric org, trimeric org, monomeric org-Ns, dimeric org-Ns, and trimeric org-Ns) contributing to SOA from ozonolysis of β -caryophyllene in presence of nitrogen oxides at temperatures between 213 – 313 K.

Table1. Compilation of experimental conditions, O:C ratios, SOA densities, and yields

Exp No.	T [K]	RH [%]	BCP [$\mu\text{g m}^{-3}$]	Total O ₃ [ppb]	Total NO ₂ [ppb]	NO ₃ radicals [ppb]	SOA mass ⁺ (SMPS) [$\mu\text{g m}^{-3}$]	geo. mean diameter (nm)	O:C (CIM S)	SOA density [g cm^{-3}]	SOA Yield [%]	BCA Yield [%]
1a	213	96	#	320	0	-	14.4±3.6	67.3±1.2	0.26	0.75±0.06	-	-
2a	243	88	15.8±3.1 (15.6±3.1 [†])	317	0	-	4.5±1.1	73.3±1.1	0.32	0.86±0.14	28.5±8	11.1±7.3
3a	273	67	109.5±21.9	73	0	-	31.5±7.9	73.5±1.9	0.38	1.0±0.08	29.7±8	4.1±2.7
4a	298	27	65.0±13 (23.2±4.6 [†])	325	0	-	13.6±3.4	68.5±1.5	0.41	1.09±0.12	20.9±6	0.8 ± 0.5 (4*)
5a	313	13	78.6±15.7	290	0	-	13.7±3.4	52.9±3.6	0.42	1.09±0.16 _{ass}	17.4±5	0.6±0.4
6a	243	59	65.0±13.0	6-505	0	-	-	=	-	-	-	-
7a	258	50	94±19	13-419	0	-	-	=	-	-	-	-
1b	213	96	#	320	27	-	13.9±3.5 (28.3±7.1 [†])	92.7±0.5	0.27	-	-	-
2b	243	88	14.2± 2.8 [†]	317	32	0.8	18.0±4.5 (52.5±13.1 [†])	96.0±1.1	0.31	-	-	-
3b	273	67	109.5±21.9 [†]	361	39	3.5	30.8±7.7 (62.4±15.6 [†])	114.9±1.6	0.40	-	-	-
4b	298	27	27.9± 5.6 [†]	325	42	5.3	32.1±8.0 (49.1±12.3 [†])	106.8±1.2	0.44	-	-	-
5b	313	13	24.6±4.9 [†]	290	46	5.6	21.8±5.5 (34.4±8.6 [†])	87.6±1.7	0.45	-	-	-

925 *measured by (Jaoui et al., 2003); ⁺SOA mass wall loss corrected; [†] following BCP addition calculated assuming a constant BCP addition rate in each experiment; #not detectable due to wall losses. _{ass} assumed to have the same density as at 298K due to its large measurement uncertainty; BCA, β -caryophyllinic acid, regarded as a tracer of β -caryophyllene in the atmosphere.

930 **Supplement for:**

Kinetics, SOA yields and chemical composition of secondary organic aerosol from β -caryophyllene ozonolysis with and without nitrogen oxides between 213 and 313 K

935 Linyu Gao^{1,2*}, Junwei Song^{1,2}, Claudia Mohr³, Wei Huang⁴, Magdalena Vallon¹, Feng Jiang^{1,2}, Thomas Leisner^{1,5}, and Harald Saathoff^{1*}

1. Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

2. Institute of Geography and Geoecology, Working Group for Environmental Mineralogy and Environmental System Analysis, Karlsruhe Institute of Technology, Karlsruhe, Germany


3. Department of Environmental Science, Stockholm University, Stockholm, Sweden

940 4. Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Helsinki, Finland


5. Institute of Environmental Physics, Heidelberg University, Heidelberg, Germany

Correspondence to: Linyu Gao (linyu.gao@kit.edu) and Harald Saathoff (harald.saathoff@kit.edu)

945



950 **Figure S1. (a) Example of the 5-min average mass spectra-spectrum of β -caryophyllene subtracted from the chamber background when the β -caryophyllene ion (m/z 205.20) shows relatively stable signal intensity in the chamber and (b) relative contribution of β -caryophyllene related ions detected by the PTR-MS in this study.**



955 **Figure S2: Calibration of the FIGAERO-CIMS with the β -caryophyllinic acid (BCA).** The β -caryophyllinic acid was dissolved into methanol to 0.2 ng m^{-3} as a standard BCA solution. Different volume of the standard BCA solution was deposited on a PTFE filter and then the deposited filter was heated by FIGAERO -CIMS carried by ultra-high purity nitrogen following a thermal desorption, as described in Section 2.1. The sensitivity of CIMS to β -caryophyllinic acid was calculated as $(2.4^{+0.96}_{-0.63}) \text{ cps ppt}^{-1}$.

960




Figure S3. Plot of mass defect corresponding to the mass of molecules assigned for the BCP SOA in this work.

965

970




Figure S4. Time evolution of experiments at 213K, 243K, 273K, and 313K.

975







Figure S3S5: Example of the fit of the kinetic model to the observation for the experiment at 313K. Three main reactions were considered in the kinetic analysis to fit the observations:



980



where BCP is the beta-caryophyllene, Pro1, Pro2 and Pro3 represent different reaction products, and k_1, k_2, k_3 are the corresponding reaction rate constants.

985

990

995





Figure S4S6: Temperature dependence of SOA yields as a function of the total organic particle mass concentration (M_{org}). Fitted lines are based on volatility distributions.


1000

1005

1010




1015 | Figure S557: (a) Comparison between the average mass spectra detected by the PTR-MS over the 5 min before and after ozonolysis of β -caryophyllene in the experiment at 273 K on 12/11/2019. (b) Time series of β -caryophyllene (m/z 205.20) and several major gas phase product ions (m/z 237.18, m/z 235.17, m/z 219.17, m/z 209.15 and m/z 195.14) as labelled in (a). Two dashed line show the time points of β -caryophyllene addition and first O_3 addition to the chamber, respectively.



020


Figure S8. Averaged CIMS gas phase mass spectra (background subtracted) for all temperatures, with absolute normalised signal vs. mass to charge ratios (m/z).



1025

Figure S6S9: (a) Average mass spectra of particle phase by the CHARON (blue) and HEPA filter (red) measurement over the course of 1 hour (18:00-19:00 as shown in gray area in (b), aerosol mass concentration: $62 \pm 0.3 \mu\text{g m}^{-3}$) in the β -caryophyllene ozonolysis experiment at 273 K in 12/11/2019. Several major product ions (m/z 221, 235, 237, 239 and 251) are labelled. (b) time series of SMPS mass concentration of aerosol particles and two particle phase product ions (m/z 221 and 235) detected by the CHARON-PTR-MS. The time series of geometric mean size of aerosol particles is given to show the reduced transmission efficiency of CHARON inlet for smaller particles.

1030



1035 | Figure S7S10. Evolution of major oxidation products detected by CIMS in the gas phase in the absence of NO₂. The time zero represents the timepoint when ozone was added first.

1040

1045

1050









Figure S8S11. The potential esterification pathways of the formation of C₂₉ and C₄₄ compounds at 213 K. The structure of C₁₅H₂₄O₃, C₁₄H₂₂O₄, C₁₅H₂₆O₄, and C₁₄H₂₂O₇ were identified by (Li et al., 2011; Chan et al., 2011). Panel (a) shows the dimer (C₂₉H₄₄O₆) and trimer (C₄₄H₆₈O₉) formation at 213K; panel (b) represents the dimer (C₂₉H₄₄O₉) at 313K.

1055 Similarly, the formation pathway of C₃₀ and C₃₅ compounds could be speculated as follows:



Note that we observed a short but significant increase of the dimer signal C₃₀H₄₈O₅ in the gas phase at 213K after the second addition of β-caryophyllene into the chamber. This suggests that the dimer formation could happen in the gas phase at 213K. However, no significant signal change of dimeric molecules was observed at 313K in the gas phase over the course of the experiments. One explanation could be the dimeric compounds (e.g. C₂₉H₄₄O₉) are quite highly oxygenated and extremely low volatile, resulting in a fast condensation process before being detected in the gas phase. The possibility cannot be excluded that the dimeric compounds could be formed in the condensed phase as no dimeric signals could be detected visibly even with the time resolution of 1 s.




1070

Figure S9S12. Fractions of six major particle phase compounds to the sum of all $C_xH_yO_z$ compounds as a function of temperature. The particle fraction (solid bars) and the gas fraction (dashed bars) of each compound are shown.




Figure S9-S12 shows the temperature dependent fractions of six typical and most abundant monomers including the β -caryophyllene tracer, β -caryophyllinic acid ($C_{14}H_{22}O_4$), in both gas and particle phase as formed from β -caryophyllene ozonolysis without NO_2 . Among the few compounds in the gas phase, $C_{15}H_{26}O_4$ and $C_{15}H_{24}O_3$ (likely β -caryophyllinic acid) contributed mainly at lower temperatures (213 – 243 K). $C_{14}H_{22}O_4$ was observed at all temperatures even though its contribution was on average only 0.05% to the total mass including gas and particle phase. In the particle phase, $C_{15}H_{24}O_4$ (likely β -hydroxycaryophyllonic acid) was observed at all temperatures and contributed most at 273 K. $C_{15}H_{24}O_3$ and $C_{15}H_{26}O_4$ were mainly present at lower temperatures. The two most abundant auto oxidation products, $C_{14}H_{22}O_6$ (likely hydrated β -oxocaryophyllonic acid) and $C_{14}H_{22}O_7$ (not yet identified) were observed at all temperatures but their signals were significantly increasing with temperature. Their mass contribution in the particle phase were 0.08 to 2.3% and 0.01 to 3.6% from 213 K to 313 K, respectively.

1080



1085 **Figure S10S13.** Modelling the NO₃ radical formation at 298K before the last β-caryophyllene addition. Time evolution of all reactants are modelled (left panel), including NO₂ (black line), O₃ (red line), NO₃ (green line), O₂ (blue line), N₂O₅ (purple line), HNO₃ (grey line) and wall loss (orange line). The right panel shows the NO₃ concentration evolution alone. Time zero is marked as the start of NO₂ addition.

1090 Because of an excess of ozone in the chamber over the course of the whole experiment, the following reactions were considered [to estimate the potential NO₃ radical levels before the addition of another portion of BCP](#):




All the rate constants used at all temperatures are listed below (IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (ipsl.fr), <https://iupac-aeris.ipsl.fr/#>):


1100 [Please note that we did not include the reactions of NO₃ radicals with BCP or its oxidation products in our box model calculations used to estimate the NO₃ radical concentrations. Therefore, these NO₃ radical concentrations must be considered as upper limits.](#)

	k1	k2	k3	k4	k5*
213K	1.3E-18	3.5E-10	8.3E-07	1.00E-22	1.00E-03
243K	5.4E-18	2.0E-10	5.5E-04	1.00E-22	1.00E-03
273K	1.6E-17	1.3E-10	0.09	1.00E-22	1.00E-03
298K	3.5E-17	8.9E-11	2.9	1.00E-22	1.00E-03
313K	5.2E-17	7.3E-11	17.8	1.00E-22	1.00E-03

*Estimated from previous experiments with N₂O₅ measured inside the AIDA chamber by FTIR.



105 Figure S4S14. Comparison of the temperature dependence on the organonitrate fraction formed. The green dots represent the fraction of the organonitrate concentrated detected by HR-AMS, while the blue dots show the signal fraction of organonitrate assigned by FIGAERO-iodide-CIMS.



| 110 **Figure S12S15.** Comparison of the CIMS signals of C₁₄H₂₂O₇ and C₁₅H₂₄O₄ before (dashed) and after (solid) the last β-caryophyllene addition.




Figure S43S16. Mass fractions of major particle phase compounds to the sum of all detected molecules as a function of temperature. The particle fraction (solid bars in the top) and the gas fraction (dashed bars in the bottom) of each compound are shown.

As shown in Fig S43S16, compared to the case in the absence of NO_2 , the mass fraction of $C_{15}H_{24}O_3$ and $C_{15}H_{24}O_4$ decreased in both gas and particle phase. $C_{30}H_{48}O_5$ contributed more at low temperatures (213 – 243 K) in particle and a relatively large portion of its gas signal showed at the coldest experiment (213 K), which indicates the vapor pressure of $C_{30}H_{48}O_5$ was not so low enough to condense to the particle phase completely even though it had a long carbon skeleton and high mass weight. In contrast, $C_{35}H_{48}O_{12}$ had stronger signals at high temperatures (273 – 313 K). However, nearly no gas signal could be detected under all temperatures due to its highly oxidation degree, indicative of those compounds to be extremely low volatile components, such as $C_{30}H_{48}O_5$ and $C_{15}H_{23}O_9N$. In addition, $C_{15}H_{25}O_7N$ acted as another most abundant nitrated compounds showed highest mass fraction at 273 K in both gas and particle phase.

Table S1. Compilation of experimental conditions and particle mass yields reported in the literature for β -caryophyllene ozonolysis

Reference	Temperature [K]	Δ Morg [$\mu\text{g m}^{-3}$]	RH [%]	Scavenger for OH or CI	Seed	Yield [%]
(Tasoglou and Pandis, 2015)	293	0.3-194	<10	2-butanol	None	1.9-104
Winterhalter et al. (2009)	296	50-370	dry	none	None	6-24
Jaoui et al. (2003)	287-290	-	80-85	none	None	39
Lee et al. (2006a)	293	336 ± 3	6.2	Cyclohexane	None	45 ± 2
Lee et al. (2006b)	295	212	56	None	None	68
Chen et al. (2012)	298	0.5-230	40	Cyclohexane	AS	8-70

- 1130 For the ozonolysis of β -caryophyllene without OH scavenger, a SOA mass yield of 27% was observed at 293 K and an organic particle mass loading of $10 \mu\text{g m}^{-3}$ (Tasoglou and Pandis, 2015). Winterhalter et al. (2009) measured yields ranging between 5 - 24% for particle mass loadings of 50 - 440 $\mu\text{g m}^{-3}$ and no Criegee intermediate scavenger, with the yields increasing for higher concentrations of water vapour or formic acid. Jaoui et al. (2003) measured a yield of 39.3 % without OH scavenger at 287-290 K and Lee et al. (2006a) observed a
- 1135 yield of 45% at 293K and a particle mass loading of $336 \mu\text{g m}^{-3}$ using cyclohexane as OH scavenger. Chen et al. (2012b) varied the ozone levels and determined SOA yields between 8 - 46% for organic particle masses of 0.5-170 $\mu\text{g m}^{-3}$, 15.8-60% at 2.1 - 230 $\mu\text{g m}^{-3}$, and 14.6-69.5% at 19 - 77.4 $\mu\text{g m}^{-3}$ for ozone levels of 50 ppb, 100 ppb and 200 ppb, respectively. By using OH scavenger, Tasoglou and Pandis (2015) also quantified SOA mass yields at room temperature to be 64.4% at $17.6 \mu\text{g m}^{-3}$ for 300 ppb of ozone, 59.6% at $16.3 \mu\text{g m}^{-3}$ for 400 ppb of ozone,
- 1140 and 66.9% at $30.5 \mu\text{g m}^{-3}$ for 500 ppb of ozone.

Table S2. Fitted reaction rates k_{O_3} and OH radical yields for temperatures between 243K to 313K

Temperature [K]	k_{O_3} [cm ³ molecule ⁻¹ s ⁻¹]	OH yields
313	(1.01±0.09) E-14	0.15±0.02
298	(1.09±0.21) E-14	0.08±0.01
258	(1.05±0.25) E-14	0.05±0.03
243	(1.99±0.28) E-14	0.05±0.02

1145

1150

1155

1160

1165

|

Table S3. Main β -caryophyllene oxidation products and their phase partitioning

Formula	<i>m/z</i> (Th)	Compound assignment from literature	Ref.	Potential dimer
C2-C4 fragments				
C ₄ H ₆ O ₄	118.0			
C ₂ H ₄ O ₃	76.0			
C14-15 monomers				
C ₁₄ H ₂₂ O ₃	238.2	β -nocaryophyllon aldehyde	a, b, c, d	
C ₁₅ H ₂₄ O ₃	252.2	β -hydroxycaryophyllon aldehyde; β -caryophyllonic acid	a, b, c, d	C ₃₀ H ₄₈ O ₆
C ₁₄ H ₂₂ O ₄	254.2	β -hydroxynocaryophyllon aldehyde; β -nocaryophyllonic acid; β -caryophyllinic acid	a, b, c, d	C ₂₈ H ₄₄ O ₈
C ₁₅ H ₂₄ O ₄	268.2	β -hydroxycaryophyllonic acid	a, d	
C ₁₅ H ₂₆ O ₄	270.2	Hydrated β -caryophyllonic acid	a	
C ₁₄ H ₂₂ O ₆	286.1	Hydrated β -oxocaryophyllonic acid	a	C ₂₈ H ₄₄ O ₁₂
C ₁₄ H ₂₂ O ₇	302.1		b	
C ₁₄ H ₂₄ O ₇	304.2			
C ₁₅ H ₂₆ O ₅	286.2	Hydrated β -hydroxycaryophyllonic acid	a	C ₃₀ H ₅₂ O ₁₀
C ₁₅ H ₂₂ O ₆	298.1			
C ₁₅ H ₂₂ O ₇	314.1		a	
C ₁₄ H ₂₀ O ₈	316.1			
C ₁₄ H ₂₂ O ₈	318.1			
C ₁₅ H ₂₂ O ₈	330.1			
C ₁₄ H ₂₂ O ₉	334.1			
C ₁₅ H ₂₄ O ₉	348.1			
C ₁₅ H ₃₀ O ₉	354.2			
C ₁₅ H ₂₅ NO ₈	347.2		a	C ₃₀ H ₅₀ N ₂ O ₁₆
C ₁₄ H ₂₃ NO ₉	349.1		a	C ₂₈ H ₄₆ N ₂ O ₁₈
C ₁₅ H ₂₇ NO ₈	349.2		a	
C ₁₅ H ₂₅ NO ₉	363.2		a	C ₃₀ H ₅₀ N ₂ O ₁₈
C28-30 dimers				
C ₂₈ H ₄₄ O ₅	460.3			
C ₃₀ H ₄₈ O ₅	488.3			
C ₂₉ H ₄₆ O ₆	490.3			
C ₂₈ H ₄₄ O ₇	492.3			
C ₃₀ H ₄₈ O ₆	504.3			
C ₂₉ H ₄₆ O ₇	506.3			
C ₂₈ H ₄₄ O ₈	504.3			
C ₃₀ H ₄₈ O ₇	520.3			
C ₂₉ H ₄₆ O ₈	552.3			
C ₃₀ H ₄₈ O ₈	536.3			
C ₂₉ H ₄₄ O ₉	536.3			
C ₂₉ H ₄₆ O ₉	538.3			

C ₂₉ H ₄₄ O ₁₀	552.3			
C ₂₈ H ₄₂ O ₁₁	554.3			
C ₂₉ H ₄₄ O ₁₁	568.3			
Others				
C ₁₁ H ₁₈ O ₃	198.1	3,3-dimethyl-2-(3-oxobutyl) cyclobutanecarboxylic acid	b	
C ₁₃ H ₂₀ O ₅	256.1	β-hydroxynornocaryophyllonic acid	a	C ₂₆ H ₄₀ O ₁₀
C ₃₅ H ₄₈ O ₁₂	660.3			
C ₄₄ H ₆₈ O ₉	740.5			

^a(Chan et al., 2011); ^b(Li et al., 2011); ^c(Winterhalter et al., 2009); ^d(Jaoui et al., 2003)

1170

|

Reference

- 1175 Chan, M., Surratt, J., Chan, A., Schilling, K., Offenberg, J., Lewandowski, M., Edney, E., Kleindienst, T., Jaoui, M., and Edgerton, E.: Influence of aerosol acidity on the chemical composition of secondary organic aerosol from β -caryophyllene, *Atmospheric Chemistry and Physics*, 11, 1735-1751, 2011.
- Chen, Q., Li, Y. L., McKinney, K. A., Kuwata, M., and Martin, S. T.: Particle mass yield from β -caryophyllene ozonolysis, *Atmos. Chem. Phys.*, 12, 3165-3179, 10.5194/acp-12-3165-2012, 2012.
- 1180 Jaoui, M., Leungsakul, S., and Kamens, R. M.: Gas and Particle Products Distribution from the Reaction of β -Caryophyllene with Ozone, *Journal of Atmospheric Chemistry*, 45, 261-287, 10.1023/A:1024263430285, 2003.
- Lee, A., Goldstein, A. H., Keywood, M. D., Gao, S., Varutbangkul, V., Bahreini, R., Ng, N. L., Flagan, R. C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the ozonolysis of ten different terpenes, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006437>, 2006a.
- 1185 Lee, A., Goldstein, A. H., Kroll, J. H., Ng, N. L., Varutbangkul, V., Flagan, R. C., and Seinfeld, J. H.: Gas-phase products and secondary aerosol yields from the photooxidation of 16 different terpenes, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2006JD007050>, 2006b.
- Li, Y., Chen, Q., Guzman, M., Chan, C. K., and Martin, S.: Second-generation products contribute substantially to the particle-phase organic material produced by β -caryophyllene ozonolysis, *Atmospheric Chemistry and Physics*, 11, 121, 2011.
- 1190 Tasoglou, A., and Pandis, S.: Formation and chemical aging of secondary organic aerosol during the β -caryophyllene oxidation, *Atmospheric Chemistry & Physics*, 15, 2015.
- Winterhalter, R., Herrmann, F., Kanawati, B., Nguyen, T. L., Peeters, J., Vereecken, L., and Moortgat, G. K.: The gas-phase ozonolysis of β -caryophyllene (C₁₅H₂₄). Part I: an experimental study, *Physical Chemistry Chemical Physics*, 11, 4152-4172, 2009.

1195