

# Satellite soil moisture data assimilation impacts on modeling weather variables and ozone in the southeastern US - Part 2: Sensitivity to dry deposition parameterizations

Min Huang<sup>1,a</sup>, James H. Crawford<sup>2</sup>, Gregory R. Carmichael<sup>3</sup>, Kevin W. Bowman<sup>4</sup>, Sujay V. Kumar<sup>5</sup>, and Colm Sweeney<sup>6</sup>

<sup>1</sup>College of Science, George Mason University, Fairfax, VA, USA

<sup>2</sup>NASA Langley Research Center, Hampton, VA, USA

<sup>3</sup>College of Engineering, The University of Iowa, Iowa City, IA, USA

<sup>4</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

<sup>5</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>6</sup>NOAA Earth System Research Laboratory Global Monitoring Division, Boulder, CO, USA

<sup>a</sup>Now also visiting: National Centers for Environmental Prediction, College Park, MD, USA

Correspondence to: Min Huang (mhuang10@gmu.edu)

**Abstract.** Ozone (O<sub>3</sub>) dry deposition is a major O<sub>3</sub> sink. As a follow-up study of Huang et al. (2021), we quantify the impact of satellite soil moisture (SM) on model representations of this process when different dry deposition parameterizations are implemented, based on which the implications for interpreting O<sub>3</sub> air pollution levels and assessing the O<sub>3</sub> impacts on human and ecosystem health, are provided. The SM<sub>a</sub> data from NASA's Soil Moisture Active Passive mission are assimilated into the Noah-Multiparameterization land surface model within the NASA Land Information System framework, semicoupled with Weather Research and Forecasting model with online Chemistry regional-scale simulations covering the southeastern US. Major changes in the used modeling system include enabling the dynamic vegetation option, adding the irrigation process, and updating the scheme for the surface exchange coefficient. Two dry deposition schemes are implemented, i.e., the Wesely scheme and a "dynamic" scheme, in the latter of which dry deposition parameterization is coupled with photosynthesis and vegetation dynamics. It is demonstrated that, when the "dynamic" scheme is applied, the simulated O<sub>3</sub> dry deposition velocities v<sub>d</sub>, their stomatal and cuticular pathways, as well as the total O<sub>3</sub> fluxes F<sub>t</sub> are overall larger; v<sub>d</sub> and F<sub>t</sub> are 2–3 times more sensitive to the SM changes due to the data assimilation (DA). Further, through case studies at two forested sites with different soil types and hydrological regimes, we highlight that, applying the Community Land Model-type of SM factor controlling stomatal resistance (i.e., β factor) scheme in replacement of the Noah-type β factor scheme reduced the v<sub>d</sub> sensitivity to SM changes by ~75% at one site while doubled this sensitivity at the other site. Referring to multiple evaluation datasets, which may be associated with variable extents of uncertainty, the model performance of vegetation, surface fluxes, weather, and surface O<sub>3</sub> concentrations, shows mixed responses to the DA, some of which display land cover dependency. Finally, using model-derived concentration- and flux-based policy relevant O<sub>3</sub> metrics as well as their matching exposure-response functions, the relative biomass/crop yield losses for several types of vegetation/crops are estimated to be within a wide range of 1–17%. Their sensitivities to the model's dry deposition scheme and the implementation of SM DA are discussed.

**Moved (insertion) [1]**

**Deleted:** Realistically representing

**Deleted:** in models is important

**Deleted:** accurately simulating

**Deleted:** concentrations and exceedances, as well as

**Moved up [1]:** As a follow-up study of Huang et al.

**Deleted:** .

**Deleted:** (2021), soil moisture (SM)

**Deleted:** C<sub>H</sub> (i.e.,

**Deleted:** for heat) scheme.

**Deleted:** different

**Deleted:** modeled

**Deleted:** as well as the total,

**Deleted:** We also

**Deleted:** configuration

**Deleted:** the

**Deleted:** β factor which presents dependencies on soil type and hydrological regime) can strongly affect

**Deleted:** quantitative results.

**Deleted:** observation and observation-derived

**Deleted:** below 20

## 1 Introduction

55 Ground-level ozone ( $O_3$ ) is a regulated secondary air pollutant harmful to human and ecosystem health (Fleming et al., 2018; Mills et al., 2018a,b). It is closely connected with  $O_3$  at higher altitudes where  $O_3$  plays a more important role in the Earth's climate system, by trapping infrared radiation and absorbing ultraviolet radiation (e.g., Lacin et al., 1990). To better protect human health and public welfare, in 2015, the US primary and secondary National Ambient Air Quality Standards were lowered from 75 ppbv to 70 ppbv, in the format of daily maximum 8-h average (MDA8). Several other  $O_3$ -exposure based

60 metrics have also been applied or/and proposed to assess  $O_3$  impacts on vegetation, such as the accumulated  $O_3$  exposure over given thresholds (e.g., SUM40, SUM60, and AOT40), the averaged  $O_3$  exposure during daylight hours (e.g., M7 and M12), and the sigmoidal-weighted W126 cumulative exposure (e.g., Fredericksen et al., 1996; van Dingenen et al., 2009; Hemispheric Transport of Air Pollution, 2010, and references therein; Avnery et al., 2011; Hollaway et al., 2012; Huang et al., 2013; Lapina et al., 2014; Mills et al., 2007, 2018a,b). To help comply with the tighter air quality standards, an improved

65 understanding of the individual processes affecting the (near)-surface  $O_3$  concentrations and exceedances is demanded. Many  $O_3$ -related processes are highly sensitive to environmental and/or biophysical conditions (e.g., Steinkamp and Lawrence, 2011; Strode et al., 2015; Jiang et al., 2018; Huang et al., 2021, and references therein). These  $O_3$ -related processes include dry deposition of  $O_3$  and its precursors, which is a major sink for near-surface  $O_3$  and depends on dry deposition velocities ( $v_d$ ) and the deposited chemicals' concentrations (Baublitz et al. 2020; Huang et al., 2021). As recognized in numerous studies,

70 accurately estimating dry deposition fluxes is critical to understanding  $O_3$  budgets and exceedances in the past, present, and future (e.g., Stevenson et al., 2006; Griffiths et al., 2021); it could also contribute to a more reasonable assessment of the  $O_3$  impacts on vegetation (e.g., Mills et al., 2011; Lombardozzi et al., 2015; Mills et al., 2018b; Ducker et al., 2018; Ronan et al., 2020; Fu et al., 2022), which is relevant to the budgets of other greenhouse gases, weather, and climate as well.

75 Ozone uptake by plants is generally higher in warm/growing seasons and during the daytime when  $O_3$  concentrations and  $v_d$  values peak. As introduced in Huang et al. (2021) as well as references therein, over the land, surface resistance  $r_s$ , which is composed of stomatal-mesophyll ( $r_s-r_m$ ), cuticular ( $r_{cu}$ ), in-canopy, and ground resistance terms, often exerts the strongest effects on the magnitude and variability of  $v_d$ .  $v_d$  also includes the aerodynamic resistance ( $r_a$ ) and quasi-laminar sublayer resistance ( $r_b$ ) terms.

80 Soil moisture (SM) and its variability impact  $v_d$  in the following ways: 1) SM can play a key role in controlling the opening and closing of plants' stomata as well as the mesophyll functioning (Egea et al., 2011; Baillie and Fleming, 2019), and thus it can directly affect the  $r_s$  and  $r_m$  terms of  $v_d$ ; 2) SM is closely linked with vegetation attributes, such as the growing-season aboveground biomass, which is often expressed as leaf area index (LAI) or vegetation optical depth (VOD) and controls the

85 stomatal and cuticular uptake of  $O_3$ -related species; and 3) SM as well as vegetation conditions can affect multiple  $v_d$  terms

Deleted: ¶

Deleted: .

Deleted: ever-tightening

Deleted:  $v_d$

Deleted: .

Deleted:  $v_d$

Deleted:  $R_c$

Deleted:  $R_a-R_m$

Deleted:  $R_{lu}$

Deleted:  $v_d, v_d$

Deleted:  $R_a$

Deleted:  $R_b$

Deleted:  $v_d$

Deleted:  $R_a$

Deleted:  $R_m$

Deleted:  $v_d$

Deleted:  $v_d$

105 through its interactions with other environmental conditions (e.g., temperatures, radiation, precipitation and humidity fields) that modulate these  $v_d$  terms, and such effects are generally stronger over transitional climate zones located between dry and wet climates. The SM impacts on  $v_d$  and atmospheric states through the above-mentioned pathways are likely to continue to grow in future. This is because, according to Intergovernmental Panel on Climate Change (2021), the occurrence and severity of droughts, some of which are characterized by surface and/or column-averaged SM deficits, are projected to increase over many US regions under warmer future environments. Better understanding the potentially enhanced SM dependency of dry deposition and weather conditions under the changing climate is important because O<sub>3</sub> stress, together with heat, water, as well as other stresses, can pose more complex threats to plant health than single stress alone (Otu-Larbi et al., 2020).

- Deleted:  $V_d$
- Deleted: It is expected that the
- Deleted: of SM
- Deleted:  $V_d$
- Deleted: will
- Deleted: increase as
- Deleted: (Intergovernmental Panel on Climate Change, 2021).
- Deleted: Out

115 Single-point models and three-dimensional chemical transport models have long been used to estimate  $v_d$  values and their responses to climate change. In the widely-used, empirical Wesely scheme (Wesely, 1989),  $v_d$  is sensitive to only a few meteorological variables, with SM and plants' physiological effects ignored. In previous studies, Wesely scheme based  $v_d$  fluxes as well as their various terms from different global, regional, and point-scale modeling systems were intercompared or/and evaluated with  $v_d$  and  $v_g$  observations from sparsely-distributed sites (e.g., Val Martin et al., 2014; Hardacre et al., 2015; Clifton et al., 2017; Silva and Heald, 2018; Wu et al., 2018; Lin et al., 2019) in terms of their magnitude and variability. Studies such as Hardacre et al. (2015) show that, even when similar (Wesely and Wesely-like)  $v_d$  schemes were applied, various models behaved differently in calculating  $v_d$ , reflecting the impacts of land use/land cover (LULC) and meteorological fields which depend on the individual models' configurations (e.g., scales, inputs). In almost all above-cited studies, large model-model and model-observation discrepancies (i.e., by a factor of 2 or more) have been found in places, suggesting the strong needs of diagnosing and addressing issues in the models' configurations and dry deposition parameterizations.

- Deleted: Chemical
- Deleted:  $V_d$
- Deleted:  $V_d$
- Deleted:  $V_d$
- Deleted: and
- Deleted: chemistry
- Deleted:  $V_d$
- Deleted:  $R_s$
- Deleted: : Clifton et al., 2020
- Deleted: Even
- Deleted:  $V_d$
- Deleted:  $V_d$
- Deleted: Large
- Deleted:  $V_d$

125 Revised or alternative dry deposition schemes have been applied in an increasing number of global- and regional-scale modeling studies. In some of these works, stomatal conductance is calculated based on the one big-leaf, multiplicative algorithms that are more complicated than the Wesely (1989) approach, in the way that the empirical maximum stomatal conductance is adjusted by more factors, including water availability and vegetation attributes (e.g., Anav et al., 2018; Falk and Søvde Haslerud, 2019). In others,  $v_d$  calculations are coupled with photosynthesis and vegetation phenology (e.g., Val Martin et al., 2014; Wu et al., 2018; Lin et al., 2019; Wong et al., 2019; Clifton et al., 2020), which in this paper are frequently referred as “dynamic” schemes. Such types of modifications have been the recommended directions for improving the estimates of  $v_d$  as well as the  $v_d$  and O<sub>3</sub> responses to climate change, in that they have been demonstrated to be capable of enhancing the dynamics of the modeled  $v_d$  and reducing their systematic biases. However, results based on such updated  $v_d$  schemes are still associated with variable extents of uncertainty due to limitations in model parameterizations (related to structures, empirical parameters and stress functions) or/and configurations. In some existing works that applied the “dynamic” schemes, such uncertainty was quantified and addressed by simply scaling the fluxes resulting from the “dynamic” schemes towards flux measurements available at very limited locations during non-recent time periods (e.g., Val Martin et al., 2014).

- Deleted:  $R_s$
- Deleted: which
- Deleted: multiple
- Deleted:  $V_d$
- Deleted: Lin
- Deleted:  $V_d$
- Deleted:  $V_d$
- Deleted:  $V_d$
- Deleted:  $V_d$

170 These types of modified [dry deposition](#) schemes still require further investigations and optimizations, which can be approached by: 1) quantifying the sensitivities of process-based model variables to SM and other environmental or/and biophysical variables for various LULC and soil types; 2) improving model representations of processes central to SM states and land-atmosphere interactions, such as including irrigation and other human activities, tuning [physics schemes \(e.g., those related to the surface exchange coefficient,  \$C\_{H}\$ \)](#), in land surface models (LSMs), and using available observations to constrain (some of) the [key land variables in models](#); and 3) including a wide range of observations and/or observation-derived carbon, water, and energy fluxes as well as vegetation states in model evaluation for broad geographical regions. Furthermore, it is important to explicitly connect the progress in dry deposition modeling with the impact assessments of O<sub>3</sub> and other air pollutants on ecosystem health, productivity, and diversity.

Deleted:  $V_d$

A regional-scale land modeling and SM data assimilation (DA) framework coupled with weather and atmospheric chemistry modeling by the Weather Research and Forecasting model with online Chemistry (WRF-Chem) is implemented in this work.

180 Using this tool, we quantify and discuss the responses of  $V_d$  and its key components as well as O<sub>3</sub> concentrations and plant uptake to SM changes due to the DA, for different soil texture, LULC and crop types. The central parts of this work rely on the Noah-Multiparameterization (MP, Niu et al., 2011) LSM with dynamic vegetation that enables the implementation of a modified “dynamic” dry deposition scheme. [This implemented “dynamic” scheme couples the  \$r\_s\$  calculation with photosynthesis for sunlit and shaded leaves and the  \$r\_{10}\$  calculation with vegetation phenology](#). With this modified scheme, both the indirect (i.e., via changing weather and vegetation fields) and direct effects of SM on dry deposition are considered in this modeling system. Results based on this modified and the WRF-Chem default Wesely schemes are compared and evaluated with independent datasets. As an extended work of Huang et al. (2021), here we continue to focus on the southeastern US during summer 2016 for which period prior Noah- and Wesely-based model calculations were conducted and aircraft observations are available. This manuscript introduces the applied two dry deposition schemes in Section 2. It then presents [SM and vegetation states \(Section 3.1\), surface fluxes and weather fields \(Section 3.2\)](#) from this Noah-MP based modeling system, in comparison with those from Huang et al. (2021). Discussions on O<sub>3</sub> concentrations and fluxes based on all related WRF-Chem simulations are also connected with the assessment of O<sub>3</sub> impacts on societies, ecosystem health, and crop yield (Section 3.3). Summary and suggestions on future directions are provided in Section 4.

Deleted: (

Deleted: scheme

Deleted: model

Deleted:  $V_d$

Deleted: results

Deleted: )(Sections 3.1–3.2

## 2 Methods

### 195 2.1 Modeling and DA experiments design

The modeling tools and DA experiment design of this study were largely consistent with the Huang et al. (2021) study: we conducted model simulations over the southeastern US in a semi-coupled Land Information System (LIS)/WRF-Chem system without and with the assimilation of the enhanced SM retrievals from NASA’s Soil Moisture Active Passive (SMAP; [Entekhabi et al., 2010](#)) mission. Two dry deposition schemes ([details in Section 2.3](#)), were applied in cases without and with

the SM DA. The 12 km/63 vertical [Lambert conformal grid](#), atmospheric/land initialization and SM DA methods were adapted from our previous study based on the Noah LSM. Major model input datasets, physics and chemistry schemes were kept similar as before except a few aspects relevant to the upgrade of LSM from Noah to Noah-MP (version 3.6) and the implementation of an irrigation scheme to be introduced in Section 2.2.

Deleted: model domain, horizontal and  
Deleted: resolutions

Same as in Huang et al. (2021), the LULC and soil texture type inputs of our coupled modeling system were based on the International Geosphere-Biosphere Programme-modified Moderate Resolution Imaging Spectroradiometer (Table S1) and the State Soil Geographic datasets, respectively. Crop type data from Monfreda et al. (2008) were used in the irrigation scheme and the assessment of the O<sub>3</sub> impacts on vegetation (Figure 1b), which are roughly consistent with the 2016 records from the US Department of Agriculture National Agricultural Statistics Service for several major crops such as maize, soybean and wheat (<https://nassgeodata.gmu.edu/CropScape>, [Jast](#) access: 8 November 2021). In Section 3 of this paper, model results are summarized and/or discussed by groups of grid-dominant LULC and soil type that are shown in Figure 1(a, d). The original 20 LULC types were grouped into urban and non-urban areas, and for vegetation-dominant areas, into forests, croplands, and shrub/grasslands, following the criteria introduced in Table S1. The grid-dominant LULC groups for vegetated regions used in our analysis are vastly similar to independently-developed data products, e.g.: a dataset derived from the European Space Agency–Climate Change Initiative Land Cover project (<https://gwis.jrc.ec.europa.eu/apps/country.profile/overview/USA>, [Jast](#) access: 8 November 2021), and the 2016 National Land Cover Database ([Wickham et al., 2021](#)). Urban-dominant grid cells are well aligned with dense population areas (Figure 1c) based on the Gridded Population of the World version 4.11 (NASA Socioeconomic Data and Applications Center, 2018). Grid-scale discrepancies exist between the used LULC input and independent LULC products, which, however, are not anticipated to considerably impact the results averaged by LULC groups. Three groups of soil are highlighted, namely sand/loamy sand, loam and clay. The original sand and loamy sand categories are combined because of their high sand fractions ([http://www.soilinfo.psu.edu/index.cgi?soil\\_data&conus&data\\_cov&fract&methods](http://www.soilinfo.psu.edu/index.cgi?soil_data&conus&data_cov&fract&methods), [Jast](#) access: 10 December 2021).

Deleted: lass

Deleted: ;

Deleted: lass

Deleted: .

## 2.2 Physics and configurations of the Noah-MP LSM

The Noah-MP LSM includes a number of improvements from Noah, and one of the enhanced features in Noah-MP is that it contains a separate canopy layer that explicitly computes photosynthetically active radiation, canopy temperature, and related energy, water, and carbon fluxes so that it facilitates a dynamic vegetation model. A modified two-stream radiation transfer scheme was used to compute fractions of sunlit and shaded leaves and their absorbed solar radiation. The Ball-Berry type of  $f_s$  scheme (e.g., Ball et al., 1987) was applied as required by the dynamic vegetation option. When this option is used, green vegetation fraction (GVF) does not come from an input dataset as in Huang et al. (2021) but is related to LAI based on (1):

Deleted:  $R_s$

$$GVF = 1 - e^{-0.52LAI} \quad (1)$$

Deleted: GVF

Deleted:  $e^{-0.52LAI}$

Deleted:  $R_s$

Niyogi and Raman (1997) concluded that Ball-Berry along with two other physiological schemes, performed better on  $f_s$  than the multiplicative Jarvis type which has been frequently used with the prescribed vegetation option. Specifically, it helps better

capture the variance in  $\underline{f_s}$  and is more responsive to environmental changes. As described in Appendix B of Niu et al. (2011), this scheme relates stomatal resistance  $r_{s,i}$  of sunlit and shaded leaves  $i$  to the photosynthesis rates ( $A_i$ ) per unit LAI of sunlit and shaded leaves  $i$  separately:

$$\frac{1}{r_{s,i}} = m \frac{A_i}{C_{air}} \frac{e_{air}}{e_{sat}(TV)} P_{air} + g_{min} \quad (2)$$

255 where  $C_{air}$  is CO<sub>2</sub> concentration at the leaf surface. For our study period this was set at 400 ppmv according to the median value of Atmospheric Carbon and Transport (ACT)-America B-200 aircraft near-surface (i.e., >900 hPa) CO<sub>2</sub> observations, which is close to the global monthly-mean CO<sub>2</sub> concentrations in August 2016 (https://gml.noaa.gov/webdata/ccgg/trends/co2/co2\_mm\_gl.txt, [last](#) access: 8 November 2021);  $TV$ ,  $P_{air}$ ,  $e_{air}$ , and  $e_{sat}(TV)$  are canopy temperature, surface air pressure, vapor pressure at the leaf surface, and saturation vapor pressure inside leaf, 260 respectively;  $g_{min}$  and  $m$  are land cover dependent empirical parameters.  $A_i$  is determined by equations (3)–(6):

$$A_i = I_{gs} \min(A_c, A_{L,i}, A_s) \quad (3)$$

$$A_c = \frac{(c_i - c_{cp}) V_{max}}{c_i + K_c (1 + \frac{o_i}{K_o})} \quad (4)$$

$$A_{L,i} = \frac{(c_i - c_{cp})^{1.6} \alpha PAR_i}{c_i + 2c_{cp}} \quad (5)$$

$$A_s = 0.5 V_{max} \quad (6)$$

265 where  $I_{gs}$  is a TV-dependent growing season index,  $A_c$ ,  $A_{L,i}$ , and  $A_s$  are carboxylase-limited, light-limited, and export-limited photosynthesis rates per unit LAI, respectively;  $c_i$  and  $o_i$  are CO<sub>2</sub> concentrations inside leaf cavity which is about 0.7 times of the atmospheric CO<sub>2</sub> concentration and atmospheric O<sub>2</sub> concentration, respectively.  $PAR$  represents the photosynthetically active radiation per unit LAI.  $c_{cp}$  is the CO<sub>2</sub> compensation point and it equals to  $0.5 \frac{K_c}{K_o} 0.21 o_i$ , where  $K_c$  and  $K_o$  are the Michaelis-Menton constants for CO<sub>2</sub> and O<sub>2</sub>, respectively, varying with TV;  $\alpha$  is the quantum efficiency.

270  $V_{max}$  represents the maximum rate of carboxylation, expressed as:

$$V_{max} = V_{max25} \alpha_{vmax}^{\frac{TV-25}{10}} f(N) f(TV) \beta \quad (7)$$

where  $V_{max25}$  is maximum carboxylation rate at 25°C;  $f(TV)$  is a function that mimics thermal breakdown of metabolic processes;  $f(N)$  is a foliage nitrogen factor; and  $\beta$  is the SM factor controlling  $\underline{f_s}$ , which presents strong dependencies on soil type and hydrological regime. In this study model results based on the Noah and the Community Land Model (CLM, [versions](#)

275 [4.5 and earlier](#)) types of  $\beta$  schemes are compared (Table 1), the latter of which is known to often result in sharper and narrower ranges of variation with SM than the former does. [The Noah and CLM types of  \$\beta\$  parameterizations are based on equations \(8\) and \(9\), respectively:](#)

$$\beta = \sum_{i=1}^{N_{root}} \frac{\Delta z_i}{z_{root}} \min(1.0, \frac{\theta_{liq,i} - \theta_{wilt}}{\theta_{ref} - \theta_{wilt}}) \quad (8)$$

$$\beta = \sum_{i=1}^{N_{root}} \frac{\Delta z_i}{z_{root}} \min(1.0, \frac{\psi_{wilt} - \psi_i}{\psi_{wilt} - \psi_{sat}}), \text{ where } \psi_i = \psi_{sat} (\frac{\theta_{liq,i}}{\theta_{sat}})^{-b} \quad (9)$$

Deleted:  $R_s$

Deleted: lass

Deleted:  $V_{max}$

Deleted:  $R_s$

285  $\theta_{liq,i}$ ,  $\theta_{wilt,i}$ ,  $\theta_{ref,i}$ , and  $\theta_{sat}$  are SM at soil layer *i*, wilting point, reference and saturated SM, respectively.  $N_{root}$  and  $z_{root}$  are the  
 290 numbers of soil layers containing roots and total depth of root zone, respectively.  $\psi_{i}$ ,  $\psi_{wilt,i}$ , and  $\psi_{sat}$  are matric potential at  
 soil layer *i*, wilting and saturated matric potential, respectively, and *b* is Clapp-Hornberger parameter. Major parameters for  
 the calculations of  $\beta$  in both schemes are soil type dependent.

Other Noah-MP configurations which can affect the modeled land state and flux variables include: the three-layer snowpack  
 physics and the CLASS snow surface albedo; the Jordan scheme for partitioning precipitation into rainfall and snowfall; the  
 Niu-Yang-2006 frozen soil permeability and supercooled liquid water option; the Simple Groundwater Model runoff scheme;  
 and the Monin-Obukhov  $C_H$  scheme, which is based on more general Monin-Obukhov similarity theory, and unlike Noah's  
 default Chen97 scheme, accounts for the zero-displacement height. Being affected by stability correction and additional effects  
 of planetary boundary layer height on friction velocity, it is likely that the Monin-Obukhov scheme can result in either weaker  
 295 or greater  $C_H$  (i.e., less or more efficient ventilation of the land surface) than the Chen97 scheme during the daytime in summer  
 (Niu et al., 2011; Yang et al., 2011).

Irrigation process was included in all Noah-MP based simulations in this study. The benefit of including irrigation relies on  
 the choice and parameterization of the irrigation scheme, as well as the LSM model's inputs (Lawston et al., 2015). Sprinkler  
 300 scheme was chosen as it was reported as the prevalent irrigation method in 2015 across the US and many of the states within  
 our model domain (Dieter et al., 2018). Irrigation was triggered over irrigated land in growing season within local morning  
 times (6–10 am) when rootzone SM drops below 50% of the soil field capacity. The irrigated land was determined by the  
 model's LULC input and irrigation intensity information in Salmon et al. (2015), and the rootzone area was derived from the  
 maximum root depth, which varies by crop type and GVF.

305 **2.3 Wesely and dynamic  $O_3$  dry deposition schemes**

Dry deposition velocity  $v_d$  is estimated based on the resistance analogy approach:

$$v_d = \frac{1}{r_a + r_b + r_c} \quad (10)$$

$r_a$  and  $r_b$  are aerodynamic resistance and quasi-laminar sublayer resistance, respectively, sensitive to surface properties such  
 as surface roughness, and follow the Monin-Obukhov similarity theory. Over the land, surface resistance  $r_c$ , the major  
 310 component of  $v_d$ , is classified into stomatal-mesophyll resistance ( $r_s$ ,  $r_m$ ), cuticular resistance ( $r_{cu}$ ), in-canopy resistance ( $r_{dc}$   
 and  $r_{cl}$ ), and ground resistance ( $r_{gc}$  and  $r_{gs}$ ):

$$r_c = \frac{1}{\frac{1}{r_s+r_m} + \frac{1}{r_{cu}} + \frac{1}{r_{dc}+r_{cl}} + \frac{1}{r_{gc}+r_{gs}}} \quad (11)$$

Deleted: ,

- Deleted:  $V_d$
- Deleted:  $V_d = \frac{1}{r_a + r_b + r_c} \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow (8) \uparrow$   
 $R_a$
- Deleted:  $R_b$
- Deleted: ,
- Deleted:  $R_c$
- Deleted:  $V_d$
- Deleted:  $R_s$
- Deleted:  $R_m$
- Deleted:  $R_{lu}$
- Deleted:  $R_{dc}$
- Deleted:  $R_{cl}$
- Deleted:  $R_{ac}$
- Deleted:  $R$
- Deleted:  $R_c = \frac{1}{\frac{1}{r_s+r_m} + \frac{1}{r_{cu}} + \frac{1}{r_{dc}+r_{cl}} + \frac{1}{r_{gc}+r_{gs}}} \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow \rightarrow (9) \uparrow$   
where  $R_{dc}$

330 where  $r_{dc}$  is resistance for gas-phase transfer affected by buoyant convection in the canopy when sunlight heats the (near-) surface,  $r_{cl}$  is resistance for leaves, twigs, bark, and others in the lower canopy,  $r_{ac}$  is resistance for transfer that depends mostly on canopy structure, and  $r_{gs}$  is resistance for soil, leaf litter, snow, and others at the ground surface.

Deleted:  $R_{cl}$   
Deleted:  $R_{ac}$   
Deleted:  $R_{gs}$

Two deposition schemes, namely the Wesely and a dynamic scheme, were applied in this study, in which  $r_s$  and  $r_{tu}$  are treated differently. In the Wesely scheme,  $r_s$  and  $r_{tu}$  are calculated based on (12) and (13):

Deleted:  $R_s$   
Deleted:  $R_{tu}$   
Deleted:  $R_s$   
Deleted:  $R_{tu}$   
Deleted: 10  
Deleted: 11

$$r_s = \begin{cases} r_i \left\{ 1 + \left[ \frac{200}{G+0.1} \right]^2 \right\} \left\{ \frac{400}{T_s(40-T_s)} \right\} \frac{D_{H_2O}}{D_x}, & 0^\circ\text{C} \leq T_s \leq 40^\circ\text{C} \\ \sim 9999, & \text{assuming mass transfer through stomata stops, } T_s > 40^\circ\text{C or } < 0^\circ\text{C} \end{cases} \quad (12)$$

$$r_{tu} = \frac{r_{tu,min}}{10^{-5}H+f_0} + 1000e^{-T_s-4}, \text{ for dry surfaces according to humidity and precipitation fields} \quad (13)$$

Deleted:  $R_s = \begin{cases} r_s \left\{ 1 + \left[ \frac{200}{G+0.1} \right]^2 \right\} \left\{ \frac{400}{T_s(40-T_s)} \right\} \frac{D_{H_2O}}{D_x}, & 0^\circ\text{C} \leq T_s \leq 40^\circ\text{C} \\ \sim 9999, & > 40^\circ\text{C or } T_s < 0^\circ\text{C} \end{cases}$   
Deleted:  $R_{tu} = \frac{r_{tu}}{10^{-5}H+f_0}$

340 Where the LULC- and season-dependent constants  $r_i$  and  $r_{tu,min}$  represent the minimum stomatal and cuticular resistances, respectively, which are subject to uncertainty;  $G$  and  $T_s$  are radiation and surface temperature, respectively, whose definitions are different than those of PAR and TV in equations (2)–(7);  $D_{H_2O}$  and  $D_x$  are molecular diffusivities for water vapor and trace gas  $x$  (e.g., O<sub>3</sub>), respectively;  $H$ , which is sensitive to surface temperature, represents the Henry's law constant for the focused trace gas; and  $f_0$  is a reactivity factor for oxidation. The Wesely-scheme related results new from this study and Huang et al. (2021) are compared (Table 1).

Deleted: 11  
Deleted:  $r_s$  and  $r_{tu}$  are  
Formatted: Font color: Black  
Formatted: Font color: Text 1

345 As expressed in equations (14), in the dynamic scheme,  $r_s$  used in dry deposition modeling was taken from what's calculated from Noah-MP's dynamic vegetation model, and thus considers the physiological process of leaf responses to photosynthesis rate, humidity and CO<sub>2</sub> concentrations. The direct effects of SM, as reflected in the  $\beta$  formula, as well as other environmental variables, are included in this method, and this work quantifies the impact of the  $\beta$  factor configurations in Noah-MP (Table 1) on the dynamic-scheme-related results.

Deleted: 12  
Deleted:  $R_s$   
Deleted:  $R_s$

$$350 r_s = \left( \frac{r_{s,sunlit}L_{sunlit} + r_{s,shaded}L_{shaded}}{LAI} \right) \frac{D_{H_2O}}{D_x} \quad (14)$$

Deleted:  $R_s$   
Deleted: 12  
Deleted:  $R_{tu}$   
Deleted:  $\frac{r_{tu}}{LAI \times (10^{-5}H+f_0)}$

where  $r_{s,sunlit}$  and  $r_{s,shaded}$  are computed based on equations (2)–(7),  $L_{sunlit}$  and  $L_{shaded}$  are proportional to the sunlit and shaded fractions of canopy, respectively, calculated based on the modified two-stream radiation transfer scheme.

In the dynamic scheme,  $r_{tu}$  for dry surfaces is modified from the Wesely formula by considering its LAI dependency:

$$r_{tu} = \frac{r_{tu,min}}{LAI \times (10^{-5}H+f_0)} + 1000e^{-T_s-4} \quad (15)$$

Deleted: 13  
Deleted:  $R_{dc}$   
Deleted:  $R_m$   
Deleted:  $R_m$   
Deleted: 14  
Deleted:  $R_{tu}$

355 In both the Wesely and the dynamic schemes,  $r_{dc}$  is sensitive to surface radiation, and  $r_m$  is expressed as:

$$r_m = \frac{1}{\frac{H}{3000} + 100f_0} \quad (16)$$

Deleted: 11  
Deleted: 13  
Deleted:  $R_{gs}$   
Deleted:  $R_{cl}$

Similar to the  $r_{tu}$  calculations in equations (13) and (15), to approximate an effect that coldness sometimes reduces the uptake,  $1000e^{-T_s-4}$  is added to LULC- and season-dependent constants to derive  $r_{gs}$  and  $r_{cl}$ . It is worth mentioning that the direct



effects of water stress on mesophyll resistance have been recognized (e.g., Egea et al., 2011). Yet, in neither scheme we applied, such effects have been incorporated into the  $r_m$  formula as a part of the  $v_d$  calculation.

Deleted:  $R_m$

Deleted:  $V_d$

#### 2.4 Model evaluation, analysis, and O<sub>3</sub> impact assessments

For the cases listed in Table 1, we quantify the impacts of SM DA on the modeled SM, vegetation dynamics, surface fluxes (i.e., gross primary productivity, GPP, which is integrated by LAI from  $A$  in equations (2)–(3), energy fluxes and their partitioning in the format of evaporative fraction, dry deposition flux and individual  $v_d$  terms for O<sub>3</sub> particularly  $v_z$  and  $v_{tu}$  related), meteorological and surface O<sub>3</sub> fields during the 16–28 August 2016 period. The SM DA impacts on most of these model fields are expressed as daily or/and daytime (around 13:00–24:00 UTC) averaged absolute or relative changes referring to the results from the no-DA cases. For O<sub>3</sub> dry deposition fluxes, we also conducted linear regression analyses to determine the relationships between the relative flux changes versus the relative changes in column-averaged initial SM due to the DA. Results of O<sub>3</sub> dry deposition fluxes and the regression analyses (i.e., slopes and their standard errors, correlation coefficient  $r$  values, and  $p$  values) are summarized by grouped LULC types defined in Figure 1a. Case studies were also conducted at two low-elevation forested sites where we investigated in detail the diurnal and daily variability of O<sub>3</sub> dry deposition fluxes from various model cases and an independent dataset.

Deleted:  $V_d$

Deleted:  $R_z$

Deleted:  $R_{tu}$

Deleted: (-

A variety of data products were utilized in this study to assess the model performance in no-DA and DA cases (Table 2). Many of these evaluation datasets have been applied and introduced in detail in Huang et al. (2021), which are: 1) National Centers for Environmental Prediction Global Surface Observational Weather Data as well as weather data collected onboard the NASA B-200 aircraft during the ACT-America campaign; 2) hourly surface O<sub>3</sub> measurements at the US Environmental Protection Agency Clean Air Status and Trends Network (CASTNET) and Air Quality System (AQS) sites; and 3) daily, 0.5°×0.5° FLUXCOM latent and sensible heat fluxes. New evaluation datasets used in this work include: 1) VOD retrievals from the 9 km enhanced SMAP product, which indicates the attenuation of microwave signals by vegetation, proportional to above-ground canopy biomass, and was used together with a 10-day average Copernicus Global Land Service GVF product to derive GVF for the focused 13-day period; 2) daily GPP estimates from the 9 km SMAP level 4 carbon (L4C) product version 6, developed based on the SMAP L4 surface (0–5 cm) and rootzone (0–100 cm) SM together with satellite LULC and vegetation datasets, which was supplemented by two independent GPP proxies (Whelan et al., 2020) of satellite-derived solar-induced chlorophyll fluorescence (SIF) data (Yu et al., 2019) and the Portable Flask Package (Sweeney et al., 2015) carbonyl sulfide (OCS) measurements collected onboard the B-200 and C-130 aircraft during the ACT-America campaign, and other airborne trace gas (e.g., benzene) measurements during this campaign were analyzed together with the OCS data to help distinguish the influences of combustion sources from plant CO<sub>2</sub> uptake on the observed OCS distributions; and 3)  $v_d$  data from two selected CASTNET sites, estimated using a multilayer model (MLM, not supported by CASTNET as of 2017) version 3.0 which has known limitations and biases against eddy covariance flux measurements as well as  $v_d$  estimated using other methods (e.g., Finkelstein et al., 2000; Saylor et al., 2014; Wu et al., 2018). The known limitations of MLM and how they may affect our

Deleted: .

Deleted: the

Deleted: °, daily

Deleted: the 9 km

Deleted: the

Deleted:  $V_d$

Deleted:  $V_d$

model comparisons with the CASTNET  $V_d$  data are discussed. Our  $O_3$  dry deposition results are also compared with eddy covariance measurements reported in independent works for similar climate or/and LULC types during other time periods.

Deleted:  $V_d$

440 This study also evaluates how the SM DA affected the assessments of surface  $O_3$  impacts on human and ecosystem health. Specifically: 1) MDA8  $O_3$  fields over urban and nonurban terrestrial regions were investigated linked to their respective population ranges; and 2) the LULC-specific Phytotoxic Ozone Dose above the critical level of  $\gamma$  nmol  $m^{-2} s^{-1}$  ( $POD_\gamma$ ) as well as the crop-specific AOT40, which are defined in equations (17) and (18), were evaluated.

Deleted: urban

Deleted:  $O_3$  stomatal uptake  $F_s$  in the format of

Deleted: ),

Deleted: were evaluated based on

Deleted: 15

Deleted: 16),

Deleted:  $POD_\gamma$  (mmol  $m^{-2}$ ) =

Deleted: (15

Deleted:  $AOT40$  (ppmh) =  $\sum[(C - 0.04)]$

Deleted: 16

$$POD_\gamma \text{ (mmol } m^{-2}\text{)} = \sum[(F_s - \gamma) \times \frac{3600}{10^6}], \text{ for hourly daytime stomatal uptake } F_s > \gamma \text{ nmol } m^{-2} s^{-1} \quad (17)$$

445  $AOT40$  (ppmh) =  $\sum[(C - 0.04)]$  for hourly daytime  $O_3$  concentration  $C > 0.04$  ppmv (18)

According to Convention on Long-Range Transboundary Air Pollution (CLRTAP, 2017), the stomatal  $O_3$  uptake  $F_s$  needed in  $POD_\gamma$  calculations was derived based on equation (19):

$$F_s = C \text{ (nmol } m^{-3}\text{)} \times g_s \times \frac{r_c}{1.3 \times 150 \times \sqrt{\frac{L}{u}} + r_c} \quad (19)$$

where  $g_s$ ,  $L$ , and  $u$  are stomatal conductance, leaf width (0.04 m in this work) and surface wind speed, respectively.

450 The calculated  $POD_\gamma$  and AOT40 were used to estimate the Relative Biomass Loss (RBL) or Relative Yield Loss (RYL) for several types of vegetation or crops based on dose-response functions reported in literature (Table 3, CLRTAP, 2017; Mills et al., 2007, 2018a). Our 13-day WRF-Chem model results were linearly-extrapolated to approximately three months to derive the  $POD_\gamma$  and AOT40 fields. While we assess the uncertainty due to such linear extrapolations by relating our 13-day/extrapolated surface  $O_3$  and flux results to seasonal (e.g., averaged for three consecutive months) conditions in 2016, we focus on qualitatively interpreting the results and discussing their implications. The outcome from this analysis is also compared with the findings from several independent  $O_3$  impact assessment studies for different time periods.

Deleted: 2, Convention on Long-Range Transboundary Air Pollution...

Deleted: ~

Deleted: We

### 3 Results and discussions

Deleted: ¶

#### 3.1 Modeled SM and vegetation fields

460 Figure 2 compares the horizontal and vertical gradients of the model's initial SM conditions from the Noah\_D and CLM\_D cases defined in Table 1, in which the Noah and CLM types of  $\beta$  factor schemes were applied. At the surface layer (0–10 cm belowground), both cases produced SM horizontal gradients that resemble the Noah-based results presented in Huang et al. (2021). They are moderately correlated with the column-averaged SM fields ( $r=0.875$  and  $0.871$ , respectively), and the mean differences in column-averaged and surface SM from the Noah\_D and CLM\_D cases are  $0.003$  and  $-0.006$   $m^3 m^{-3}$ , respectively.

Deleted: modeled

465 Kumar et al. (2009) have found that, compared to other LSMs such as the Catchment (based on which the SMAP L4 datasets are produced), the 4-soil-layer Noah and 10-soil-layer CLM LSMs display successively weaker surface-subsurface coupling

strengths, and the weakest coupling strength of CLM was primarily attributed to its significantly larger number of soil layers.

485 The slightly weaker surface-subsurface correlations in the CLM\_D case than in the Noah\_D from this work, both are based on  
a 4-soil-layer Noah-MP modeling system, indicate the minor role of the LSM physics, in particular the  $\beta$  factor scheme, in  
controlling the vertical coupling strength of SM conditions.

The modeled SM fields from the Noah\_D and CLM\_D differ on grid scale, particularly in the subsurface zones (Figure 2a, b).

490 For example, in sand-dominant regions that were experiencing drought conditions during this period (e.g., Florida and the  
Texas-Oklahoma border regions, where simulated SM is mostly under  $0.2 \text{ m}^3 \text{ m}^{-3}$ ), column-averaged SM values from the  
CLM\_D case are notably smaller than those from the Noah\_D case. These results contrast with those reported by Niu et al.  
(2011), in which cases Noah-MP with the CLM-type  $\beta$  factor consumed less soil water, resulting in smaller SM variability  
than did the Noah-type  $\beta$  factor during drought periods. In their cases focusing on loam and clay soil that have higher wilting  
495 points when the CLM-type  $\beta$  factor scheme was applied, plant transpiration ceased to save soil water under drought conditions.  
Our results can be explained by the steeper CLM-type  $\beta$ -SM curve than the Noah-type  $\beta$ -SM curve for low SM, sand-dominant  
areas, as illustrated in Figure 3a of Niu et al. (2011). For such conditions, Noah-MP with the CLM-type  $\beta$  factor produces  
stronger evapotranspiration (ET) and consumes more soil water, resulting in drier soil. For wet regions where SM values  
exceed  $0.4 \text{ m}^3 \text{ m}^{-3}$ , such as Louisiana and Arkansas, the CLM- and Noah-type  $\beta$  values are close to 1.0 and insensitive to soil  
500 type and SM variations; therefore, SM and ET produced from the Noah\_D and CLM\_D cases do not diverge. These findings  
corroborate the conclusions by Yang et al. (2011) that the degree of the  $\beta$  impacts on the SM-ET relationship should depend  
on the soil type and hydrological regime, and they are important for understanding the vegetation and surface flux results to  
be presented in the later parts of this paper.

505 Referring to the SMAP SM data, in general, surface SM produced by the no-DA modeling systems show wet biases in non-  
forested regions and dry biases over the forests for the study period. These SMAP-model discrepancies were successfully  
reduced by the DA for all vegetated LULC groups (Figure S1, left), leading to overall slightly drier soil in DA-enabled  
simulations. For both the Noah\_D and CLM\_D cases, the DA adjusted the modeled SM fields across the entire soil columns,  
demonstrating that observational information at the surface was propagated into deep soil layers. The SM responses to the DA  
510 as a function of soil layer from the Noah\_D and CLM\_D cases are roughly similar but different at small spatial scales, which  
reflect the controls of the  $\beta$  factor scheme on the surface-subsurface coupling strengths of the used modeling/DA system. With  
the SMAP DA enabled, the  $r$  values between column-averaged and surface SM from the Noah\_D and CLM\_D cases increased  
to 0.902 and 0.897, respectively.

515 The satellite-derived GVF fields (methods introduced in Figure S2 caption) transition from low-to-moderate ( $<0.6$ ) to high  
( $>0.8$ ) values from the western (mostly shrub/grasslands) to the central and eastern parts (forests/croplands dominant) of the  
study region, and such spatial gradients are highly correlated with the SMAP VOD retrievals (Figure 3a, d). The Noah\_D and

Deleted: -

Deleted: ;

520 CLM\_D cases both moderately well reproduced these spatial patterns. Major differences between these cases are found in dry sandy regions, where, as discussed in previous paragraphs, more soil water was consumed for ET and plant growth in the CLM\_D case and therefore higher GVF values are given. Overall, the DA adjustments to the modeled GVF and SM fields are positively correlated (Figure S1, right), and the relative changes in GVF are smaller. While the SM changes in the Noah\_D and CLM\_D cases are of close magnitudes, GVF responded more strongly in the CLM\_D case except for sandy regions. Referring to the satellite-derived GVF fields which are also subject to large uncertainty (as discussed in Figure S2 caption), the modeled vegetation fields are more effectively improved by the DA over sparsely vegetated regions such as the South-Central Plains. The DA also remarkably reduced the model-satellite mismatches over some of the dense vegetation regions such as the southwestern Ohio. The likely degraded model performance over certain dense vegetation areas can be partially explained by weaknesses related to the SM-vegetation growth feedbacks (more details in Figure S1 caption) in the dynamic vegetation model parameterizations which need to be identified and addressed in future work. It is also suggested that jointly assimilating satellite SM and vegetation phenology products such as the VOD retrievals needs to be experimented which may maximize the positive DA impacts on multiple land variables and their atmospheric feedbacks.

Deleted: The  
Deleted: adjusted  
Deleted: toward similar directions, with  
Deleted: overall  
Deleted: .

### 3.2 Modeled fluxes and weather conditions

#### 3.2.1 Carbon/energy fluxes and weather conditions

535 Figure 4 compares the spatial distributions of the period-mean WRF-Chem carbon and energy fluxes with SMAP L4C and FLUXCOM products which contain observation information, and Table 4 summarizes WRF-Chem and observation-derived flux results by three LULC groups. The observation-derived products indicate the highest GPP and evaporative fraction (EF = daily latent heat/(daily latent heat + daily sensible heat)) over croplands. Without the DA, the Noah-MP related cases outperformed the Noah related P1\_W case on simulating EF, especially over shrub/grassland and cropland regions. This indicates that, from Noah to Noah-MP, the multiple updates in LSM physics related to  $\tau_s$ , irrigation and  $C_{it}$ , are beneficial. Larger GPP and EF values are found in CLM\_D than in Noah\_D, most of these larger values match better with the SMAP L4C and FLUXCOM data. The DA led to increased EF over shrub/grasslands in all model cases as well as over croplands in the Noah\_D case, bringing the model results closer to the FLUXCOM data. The EF values were unfavorably reduced by the DA in the CLM\_D and P1\_W cases over croplands and in all model cases over forests, reflecting the challenges of satellite SM DA over regions with dense vegetation and/or affected by human activities, which have also been reported and discussed in previous studies (e.g., Huang et al., 2021). For the Noah\_D and CLM\_D cases, this may also be due to the possibly degraded vegetation performance discussed in Section 3.1. The modeled GPP in the CLM\_D cases were lowered by the DA overall, which helped reduce the model-SMAP L4C discrepancies over forests and croplands. In the Noah\_D case, GPP was improved by the DA over forests and (slightly) over shrub/grasslands. Based on the evaluation statistics, for this case, the CLM-type  $\beta$  factor scheme is shown slightly superior to the Noah type. Note that the quality of the SMAP L4C and FLUXCOM products may also be strongly LULC dependent, e.g., it has been known that the uncertainty of SMAP L4C data is generally larger for

Deleted: 3  
Deleted: R<sub>s</sub>  
Deleted: which  
Deleted: .  
Deleted:  
Deleted: of

highly productive plant functional types (Kimball et al., 2020). Such evaluation, therefore, has demonstrated the critical role of LULC type in understanding the model performance of carbon and energy fluxes and its responses to satellite SM DA.

565

Additional datasets were also utilized to help understand terrestrial carbon uptake, including satellite SIF and ACT-America aircraft OCS as well as its vertical gradients (Figure S3). Consistent with the SMAP L4C and WRF-Chem based results, the largest SIF values are shown over croplands, especially maize and soybean fields in Illinois and Indiana, 2–3 times as high as those over shrub/grasslands in the South-Central Plains. All these datasets suggest moderate-to-high terrestrial carbon uptake around the Lower Mississippi croplands and the forests/croplands near the Texas-Oklahoma border, which is supported by the large OCS drawdowns (i.e., the free tropospheric-near surface gradients far exceeded 60 pptv) along with other trace gas measurements taken onboard the B-200 and C-130 aircraft.

570

In general, the modeled EF fields as well as their directions of changes due to the DA resemble those of latent heat flux and relative humidity (RH), which are opposite to those of sensible heat and surface temperatures (Figures 5 and S4). The model overall well reproduced the observed spatiotemporal variability of 2 m air temperature (T2) and RH, as well as FLUXCOM latent and sensible heat fluxes. The diagnostic 2 m weather fields and their responses to the DA strongly correlate with the model's surface-level results. The Noah-MP related cases reacted more strongly to the DA than the Noah-related cases, with the responses in the CLM\_D case larger than in the Noah\_D case except for dry, sandy regions, which can be attributed to combined effects of the used  $C_H$  and stomatal resistance schemes. It is important to note that diagnostic temperature and humidity variables are represented differently in Noah and Noah-MP and thus are not directly comparable. Specifically, in Noah, T2 is an explicit function of surface temperature, air density, specific heat of dry air at constant pressure, and 2 m surface exchange coefficient for heat, and 2 m specific humidity is a function of surface specific humidity, upward moisture flux at the surface, air density and 2 m surface exchange coefficient for moisture; whereas in Noah-MP, they are expressed as functions

575

580

of temperatures and water vapor for vegetated land and bare soil being weighed by their respective fractions. We therefore focus on quantitatively evaluating and intercomparing prognostic model weather variables (i.e., the model-level air temperature and humidity) against ACT-America aircraft observations (Figure 6). For air temperature, at all altitudes and near the surface (i.e.,  $\geq 800$  hPa), the CLM\_D case responded most strongly to the DA, and the DA-enabled CLM\_D case outperformed the Noah\_D and P1\_W cases. This performance is qualitatively consistent with the model's sensible heat performance referring to the FLUXCOM data. As for humidity, despite the most significant DA improvements in CLM\_D, the Noah-MP related cases did not perform as well as the Noah related cases, which is also found in the model's latent heat performance in comparison with the FLUXCOM data. However, note that the model's humidity performance is more strongly related to that of  $f_s$  and  $v_d$  in the Noah-MP based cases via the direct impacts of humidity on  $f_s$  calculations (equation 2). The solar radiation fields from all model cases, which play vital roles in controlling the land-atmosphere exchanges of water and trace gases, do not differ remarkably and their responses to the DA are negligible (e.g., Figure 5g-l). This indicates that the DA impacts on the modeled surface fluxes resulted primarily from the changes in the modeled SM, humidity, surface/canopy temperatures,

585

590

595

**Deleted:** Free-tropospheric (>2 km, above ground level, a.g.l.) OCS mixing ratios are mostly higher than those near the surface ( $\leq 2$  km, a.g.l.), except for locations that may be strongly influenced by oceanic and anthropogenic combustion sources according to independent studies (Lennartz et al., 2017; Zumkehr et al., 2018) and other chemical tracers (e.g., benzene) measured onboard. The maximum OCS mixing ratios are higher than 550 pptv, and the OCS drawdowns (i.e., free-tropospheric minus near-surface concentrations) far exceed 60 pptv around the Lower Mississippi cropland regions and the Texas-Oklahoma border where soil was wet and likely an OCS source (Bunk et al., 2017). These values are much larger than those observed in summer 2004 over the eastern US (Campbell et al., 2008), indicating possible higher OCS emissions and stronger terrestrial carbon uptake in summer 2016 than in summer 2004.

**Deleted:** model results at its

**Deleted:**

**Deleted:** skin

**Deleted:**  $R_s$

**Deleted:**  $V_d$

**Deleted:**  $R_s$

**Deleted:** dramatically

**Deleted:** -

620 as well as vegetation fields. In many cases these primary contributing factors to the DA impacts are interdependent, and their relative contributions vary by location and time.

### 3.2.2 Ozone dry deposition velocities and fluxes

Figure 7 presents the period-mean, daily-averaged  $v_d$  and dry deposition flux  $F_t$  (i.e.,  $v_d$  multiplied by concentration at the surface level, Wesely, 1989) for  $O_3$  from all model cases, along with their responses to the SMAP DA. The daytime averages of these fields have similar spatial gradients but of larger magnitudes (not shown in figures). Table 5 summarizes for three LULC groups the daily- and daytime-averaged results. The modeled stomatal–mesophyll and cuticular conductances, as well as their diurnal variability are indicated in Figure 8. All model cases produced lower  $v_d$  and  $F_t$  values over shrub/grasslands than over forests and croplands, qualitatively consistent with results from many existing model- and measurement-based studies (e.g., Val Martin et al., 2014; Hardacre et al., 2015; Silva and Heald, 2018; Lin et al., 2019). The results from Noah\_W and P1\_W, both of which are based on the same scheme (Wesely), are generally similar, with minor differences largely attributed to different surface temperature fields (Figures 5 and S4). The WRF-Chem modeled  $v_d$  and  $F_t$  fluxes were more strongly affected by the upgrade from the Wesely to the dynamic scheme: i.e., with the updated scheme, they show enhanced magnitudes, stronger spatial variability, as well as more intensive responses to the DA, especially over forests and croplands. These results can be mainly explained by the fact that the stomatal–mesophyll and cuticular resistances in the dynamic scheme are sensitive to more environmental and biophysical variables, accounting for both the direct and indirect (i.e., via influencing the weather fields and plants’ physiology) effects of SM on  $v_d$ ,  $V_d$  from the Noah\_D and CLM\_D cases, as well as its major term stomatal–mesophyll conductance, shows strong correlations with the modeled GPP, latent heat, and EF fields which have been discussed in earlier sections. Comparing the cases that implemented the CLM- and Noah-type  $\beta$  schemes,  $O_3$ -related fluxes resulting from the former configuration are of notably larger magnitude, spatial variability and absolute changes due to the DA. The SM impacts on the modeled  $v_d$  and  $F_t$  were further quantified using linear regression analyses between the relative changes in the modeled  $O_3$  fluxes due to the DA versus those in column-averaged SM initial conditions. All regression models yielded low  $p$  values (i.e.,  $\ll 0.01$ ), suggesting good  $\Delta v_d \sim \Delta SM$  and  $\Delta F_t \sim \Delta SM$  relationships. The regression slopes, all with standard errors of  $< 0.01\%$ , are summarized in barplots (Figure 9) by three LULC groups for all model cases in Table 1. For all LULC groups, the slopes based on the two cases that implemented the dynamic scheme are 2–3 times larger than those from the two cases using the Wesely scheme, and the slopes differ most strongly among the cases over forests and croplands. The low  $r$  values ( $< 0.5$ ) associated with several regression models, reflect the stronger nonlinear relationships between the changes in the studied  $O_3$  fluxes and SM. These results emphasize the importance of better understanding and representing in models the SM control on plants’ stomatal behaviors which regulate the land-atmosphere exchanges of water, energy, and trace gases. The earlier evaluation of the period-mean GPP and EF across the domain have demonstrated some advantages of using the CLM-type  $\beta$  scheme, and that the DA more effectively improved the model performance in sparsely vegetated shrub/grassland regions. These conclusions are likely also applicable to the modeled  $O_3$  dry deposition process, particularly its stomatal–mesophyll pathway.

Deleted:  $V_d$

Deleted:  $V_d$

Deleted: 3

Deleted:  $V_d$

Deleted: .

Deleted:  $V_d$

Deleted:  $V_d$ ,  $V_d$

Deleted: show

Deleted: like

Deleted:  $V_d$

Deleted:  $\Delta V_d$

Deleted: (denoted by red “L”s in Figure 9)

Deleted: realistically

Deleted: ET

Deleted: like

670 In all no-DA and DA cases, the diurnal variability of O<sub>3</sub>-related surface fluxes shows clear LULC dependency. Over the shrub/grassland and forests/croplands regions, the daytime averaged  $v_d$  values are 24–31% and 35–50% higher than the 24 h mean, respectively, while the daytime averaged F<sub>t</sub> results are 40–50% and 42–63% higher than the 24 h mean, respectively (Table 5). Such  $v_d$  diurnal cycles are a result of the strongest diurnal variability in stomatal–mesophyll conductance (i.e., its daytime mean values are approximately twice as high as the 24 h mean for all LULC types) being balanced out by weak diurnal variability associated with other  $v_d$  terms. As the most diurnally variable  $v_d$  component, stomatal–mesophyll conductance, on average, contributes less substantially to  $v_d$  for shrub/grassland areas (24 h/daytime: up to ~30%/40%) than for forests/croplands (24 h/daytime: up to ~50%/66%), which helps explain the weaker diurnal variability in the modeled  $v_d$  over shrub/grasslands. The stronger diurnal cycles in F<sub>t</sub> than in  $v_d$  reflect the impacts of higher daytime O<sub>3</sub> surface concentrations used in the F<sub>t</sub> calculations. The DA did not dominantly intensify or dampen the diurnal cycles of these fluxes for any given grouped LULC type. Whether the DA improved the estimated diurnal cycles of fluxes for various LULC types remains to be evaluated, which can benefit from independent observation-constrained flux products of broad spatial coverage and subdaily variability.

A detailed analysis was then conducted at two forest CASTNET sites with different soil types and hydrological regimes. The modeled  $v_d$  and F<sub>t</sub> from various cases are compared with the operational MLM-based calculations produced at a Florida site SUM156 and a Virginia site PED108 (Figures 10a, b, e, f and S5; Table 6) where many/most/all MLM assumptions apply. The dominant soil types at these sites are sand and loam, and the column-averaged SM values from various model cases are approximately 0.15 and 0.20 m<sup>3</sup> m<sup>-3</sup>, respectively. These various datasets show that stomatal–mesophyll conductance,  $v_d$ , and F<sub>t</sub> sharply increase soon after sunrise, reaching their daily maxima in late morning or early afternoon. The slight declines in fluxes around midday based on some simulations can result from the water and heat stresses which cause stomata closures (Figure 10c, d). The water stress starts to get relieved since the mid-afternoon at the SUM156 site under the influences of convective precipitation whereas persists throughout the afternoon at the PED108 site (Figure 10g, h), which helps shape the slightly different afternoon flux dynamics at these two locations. Without the DA, at both sites, the highest daytime fluxes were produced from the CLM\_D case, followed by the Noah\_D and Noah\_W cases, which are 2–3 times as high as the MLM-estimated. The fluxes from all WRF-Chem cases during the nighttime are close, up to >80% lower than their daytime maxima, contributed mostly by  $r_a$ ,  $r_b$  and non-stomatal  $r_c$  pathways as stomatal–mesophyll conductance is shown negligible (Figure 10c, d). Despite the uncertainty possibly introduced by the limitations of the Monin-Obukhov similarity theory, our nighttime  $v_d$  results are close to flux observations at European forest sites during both dry and wet periods in the past decades (Lin et al., 2020). They are, however, dramatically higher than the MLM-based results that are nearly zero. Wu et al. (2018) compared  $v_d$  observations with single-point model calculations based on the operational MLM, Wesely, and the Noah-GEM dynamic scheme, at a Canadian mixed forest site dominated by sand-like soil. Their diverse model results are qualitatively consistent with our findings at the SUM156 and PED108 sites. The remarkably lower  $v_d$  values from the operational MLM calculations

Deleted:  $v_d$

Deleted: 3

Deleted:  $v_d$

Deleted: ~

Deleted:  $v_d$

Deleted:  $v_d$

Deleted: contribute

Deleted:  $v_d$

Deleted:  $v_d$

Deleted:  $v_d$

Deleted:  $v_d$

Deleted: Figure

Deleted: -d

Deleted: 5).

Formatted: Font color: Text 1

Deleted: All

Deleted:  $v_d$

Deleted: .

Deleted: 10e-f

Deleted: but still dramatically higher than the MLM-based results which are nearly zero. Our nighttime  $v_d$

Deleted:  $v_d$

Deleted: a

Deleted: Noah-GEM

Deleted: these two CASTNET

Deleted:  $v_d$

730 can be partially attributed to its simplified approaches of calculating  $r_a$  and  $r_b$  using wind speed and direction, as well as the  
empirical approach of calculating  $r_s$ , which is subject to errors in the season- and LULC-dependent  $r_s$ . The possible uncertainty  
735 in MLM  $v_d$  can also be explained by the lack of continuous, accurate model input data. Specifically, the factual data such as  
plant and canopy attributes used in the MLM calculations are outdated, which, according to the CASTNET database, represent  
the conditions over 15 year ago; and based on the little day-by-day variability found in the MLM  $v_d$  data during the study  
period which contrasts with our WRF-Chem results (Figure S5), it is likely that many but not all of these are filled historical  
average  $v_d$  values due to the lack of meteorological measurements that are needed in the MLM calculation. Additionally, based  
on the surface heterogeneity within the WRF-Chem grids that these sites fall in, representation errors are estimated to be  
pronounced when comparing the point-scale MLM fluxes with our 12 km WRF-Chem results.

Deleted: are

Deleted:  $R_a$

Deleted:  $R_b$

Deleted:  $R_a$ , and

740 Within the respective ranges of the modeled SM at these two sites,  $\beta$  factors based on the CLM-type scheme are both larger  
than those based on the Noah-type  $\beta$  scheme (referring to Niu et al., Figure 3), which helps explain the higher and more variable  
model fluxes from the CLM\_D case than the Noah\_D case without the DA. At SUM156, despite the strongest SM decrease  
745 ( $\sim 0.04 \text{ m}^3 \text{ m}^{-3}$ ) by the DA in case CLM\_D, the modeled fluxes responded least strongly to the DA, in part due to the flattened  
CLM-type SM- $\beta$  curves in contrast to the linear Noah-type SM- $\beta$  function for sand within the  $0.12\text{--}0.16 \text{ m}^3 \text{ m}^{-3}$  SM range.  
At PED108, the modeled SM values from all model cases were lowered by the DA by  $\sim 0.02 \text{ m}^3 \text{ m}^{-3}$ . The stronger reactions of  
750 fluxes (i.e.,  $v_d$ ,  $F_s$ , and their stomatal-mesophyll portions) to the DA from the CLM\_D case than those from the Noah\_D case  
can be partially explained by the steep CLM-type SM- $\beta$  curve versus the linear Noah-type SM- $\beta$  relationship for loam within  
the  $0.18\text{--}0.22 \text{ m}^3 \text{ m}^{-3}$  SM range. Our case studies at these two sites with the same type of LULC emphasize the importance of  
soil type and hydrological regimes for understanding SM controls on dry deposition, which was often omitted or  
underdiscussed in previous dry deposition studies. It is noted that the effectiveness of SM DA in improving the accuracy of  
land surface states and fluxes at point scale is dependent on the representativeness of the assimilated satellite SM data for these  
sites which is expected to increase with the resolutions of the model and the assimilated satellite land product.

Deleted: the

Deleted: site

Deleted: significantly

Deleted: the

Deleted: site

Deleted: flux

Deleted: Case

### 750 3.3 Policy-relevant O<sub>3</sub> metrics and implications for O<sub>3</sub> impact assessments

#### 3.3.1 MDA8 and implications for O<sub>3</sub> health impacts

755 Figure 11 illustrates the impacts of the choice of dry deposition scheme and SM DA on WRF-Chem modeled surface MDA8  
O<sub>3</sub>. During the study period, several warmer- and drier-than-normal Atlantic states experienced high MDA8 at times (i.e.,  $\geq 60$   
ppbv, which can negatively affect lung function, and at  $\geq 70$  ppbv, cause respiratory symptoms and other adverse effects,  
Fleming et al., 2018, and references therein). Numerous populated urban centers reside in these areas. The levels of MDA8  
are shown to be much lower (i.e.,  $< 40$  ppbv) over the southern part of the domain, including several major urban/suburban  
regions such as the Texas Triangle, which was frequently influenced by passing cold fronts and tropical systems from the Gulf  
of Mexico.



770

All model cases reproduced [the observed](#) MDA8 spatial patterns ([Figure 12a](#)) moderately well. Referring to observations at AQS and CASTNET sites, their domain wide mean RMSEs all fall within 6–8.5 ppbv ([Figure 12b](#)). We first intercompare the MDA8 levels from all no-DA cases. Positive and negative differences between the results from Noah\_W and P1\_W, both of which implemented the Wesely scheme, are almost equally distributed across the domain, with the MDA8 from the former case associated with negligibly lower RMSEs (i.e., <0.02 ppbv on average) referring to AQS and CASTNET observations ([Figures 11k and 12b](#)). The differences between these two cases are largely due to the impact of the chosen LSM on the model's meteorological fields, particularly temperatures, which affected the simulations of various O<sub>3</sub>-related processes including dry deposition. As [Figures 11\(i, j\) and 12b](#) show, replacing Wesely with the dynamic dry deposition scheme considerably lowered the calculated MDA8 levels in majority of the model grids, as well as their associated RMSEs (i.e., by >0.5 ppbv on average) relative to surface observations. These reductions in MDA8 are of comparable magnitudes with those due to updating anthropogenic emissions from the National Emission Inventory 2014 to 2016beta (Huang et al., 2021). Comparing the implementations of [the CLM- and Noah-type](#) β schemes, the former led to stronger reductions in the modeled MDA8 fields and their associated uncertainty. These results reflect the impacts of the faster O<sub>3</sub> removal via dry deposition in the dynamic scheme related cases, as well as the different model meteorology. Our findings are qualitatively consistent with the conclusions from several global-scale modeling experiments that compared the Wesely and dynamic schemes (e.g., Val Martin et al., 2014; Lin et al., 2019).

Deleted: these

Deleted: 11m

775

In all model cases, the DA reduced surface and subsurface SM in many of the grids, leading to enhanced MDA8 ([Figure 11e-h](#)). The responses of the period-mean MDA8 to the DA from the Noah\_W and P1\_W cases are mostly within ±4 ppbv. When the dynamic dry deposition scheme was applied, the modeled MDA8 responded several times more strongly to the DA (i.e., by up to 6 ppbv and 8 ppbv in the Noah\_D and CLM\_D cases, respectively), especially over nonurban regions where surface MDA8 on average are several ppbv lower than in urban grids. In urban grids where population densities are ~25 times higher than in nonurban grids ([Figure 1c](#)), the DA impacts on MDA8 reach 3–4 ppbv in places, under the controls of the local-to-regional circulation patterns ([Figure 13a, e](#)). As the no-DA cases are positively biased against surface observations in many places, corresponding to the DA-induced surface O<sub>3</sub> changes, the overall model performance of MDA8 was not improved, or much degraded, by the DA. Over limited areas such as the South-Central Plains, the modeled MDA8 decreased due to the DA by up to >2 ppbv, corresponding to improved performance. The no-DA and DA results based on different LSMs and dry deposition schemes confirm that drier soil conditions exacerbate O<sub>3</sub> air pollution, which, together with heat stress, threatens human health. Such O<sub>3</sub>–SM relationships have also been demonstrated by Falk and Søvde Haslerud (2019) and Anav et al. (2018) using other chemical transport models and multiplicative dry deposition schemes. Our Noah\_W and P1\_W related results indicate the influences of SM on air quality via its feedbacks to weather; and results from the Noah\_D and CLM\_D cases provide valuable information regarding both the indirect (i.e., via adjusting vegetation phenology and weather conditions) and direct SM effects on O<sub>3</sub>. The complex SM impacts on O<sub>3</sub> dry deposition as well as surface O<sub>3</sub> concentrations based on the

Deleted: Figure 11i-m).

Deleted: Figure

Deleted: i;k;m

780

Deleted: types of

785

In all model cases, the DA reduced surface and subsurface SM in many of the grids, leading to enhanced MDA8 ([Figure 11e-h](#)). The responses of the period-mean MDA8 to the DA from the Noah\_W and P1\_W cases are mostly within ±4 ppbv. When the dynamic dry deposition scheme was applied, the modeled MDA8 responded several times more strongly to the DA (i.e., by up to 6 ppbv and 8 ppbv in the Noah\_D and CLM\_D cases, respectively), especially over nonurban regions where surface MDA8 on average are several ppbv lower than in urban grids. In urban grids where population densities are ~25 times higher than in nonurban grids ([Figure 1c](#)), the DA impacts on MDA8 reach 3–4 ppbv in places, under the controls of the local-to-regional circulation patterns ([Figure 13a, e](#)). As the no-DA cases are positively biased against surface observations in many places, corresponding to the DA-induced surface O<sub>3</sub> changes, the overall model performance of MDA8 was not improved, or much degraded, by the DA. Over limited areas such as the South-Central Plains, the modeled MDA8 decreased due to the DA by up to >2 ppbv, corresponding to improved performance. The no-DA and DA results based on different LSMs and dry deposition schemes confirm that drier soil conditions exacerbate O<sub>3</sub> air pollution, which, together with heat stress, threatens human health. Such O<sub>3</sub>–SM relationships have also been demonstrated by Falk and Søvde Haslerud (2019) and Anav et al. (2018) using other chemical transport models and multiplicative dry deposition schemes. Our Noah\_W and P1\_W related results indicate the influences of SM on air quality via its feedbacks to weather; and results from the Noah\_D and CLM\_D cases provide valuable information regarding both the indirect (i.e., via adjusting vegetation phenology and weather conditions) and direct SM effects on O<sub>3</sub>. The complex SM impacts on O<sub>3</sub> dry deposition as well as surface O<sub>3</sub> concentrations based on the

Deleted: 11f-i

Deleted: 12a;

800

coupled photosynthesis- $\tau$  calculations rely heavily on the application of water stress function ( $\beta$  scheme), soil properties and hydrological regime. The WRF-Chem results from this case indicate that, to more accurately simulate MDA8, improving land DA must be combined with **strong** efforts to identify other sources of uncertainty in  $O_3$  modeling (e.g., emissions, chemistry, and extra-regional pollution contributions) and reduce their negative impacts on model performance.

Deleted: -R,

Deleted: aggressive

### 3.3.2 Implications for $O_3$ vegetation impact assessments using concentration- and flux-based metrics

Both  $O_3$  flux- and concentration-based metrics have been applied to assess  $O_3$  impacts on vegetation as well as the associated economic loss. Estimating the plants' stomatal  $O_3$  uptake  $F_s$  is the basis for constructing flux-based  $O_3$  impact assessments. Figure J4 illustrates the period-mean daytime  $F_s$  fields based on all WRF-Chem no-DA cases as well as their responses to the SM DA. Box-and-whisker plots in Figure J3(b, f) summarize these results by three LULC groups. The averaged  $F_s$  values for all three LULC groups exceed their respective critical levels (i.e.,  $1 \text{ nmol m}^{-2} \text{ s}^{-1}$  for forest and grasslands; and  $3 \text{ nmol m}^{-2} \text{ s}^{-1}$  for crops). As a major contributor to  $O_3$  dry deposition flux during the daytime,  $F_s$  fields appear to be closely correlated in space and time with the surface humidity and flux fields (e.g., GPP, latent heat and EF, as well as  $v_d$ ), which differ distinctly from the surface  $O_3$  concentration fields. For example,  $F_s$  hotspots are shown over some low  $O_3$  concentration areas including the humid, Lower Mississippi River regions, and the lowest  $F_s$  values occur in certain high  $O_3$  concentration regions strongly affected by urban pollution (e.g., Georgia) and pollution transport from upwind US states and/or the stratosphere (e.g., western Kansas and Oklahoma, as discussed in Huang et al., 2021). The changes in  $F_s$  and surface  $O_3$  concentrations due to the DA show opposite directions, i.e., drier soil enhances surface  $O_3$  concentrations whereas slows down the plants' stomatal  $O_3$  uptake (Figures J1e-h and J4e-h). This comparison highlights how the choice of  $O_3$  metrics can affect the assessment of  $O_3$  vegetation impacts under the changing climate. As emphasized by Mills et al. (2018a) and Ronan et al. (2020), flux-based metrics have evident advantages over concentration-based metrics. To conduct reliable impact assessments using these flux-based metrics, accurate information on stomatal and non-stomatal fluxes as well as the various environmental and biophysical variables that they are sensitive to become increasingly important.

Deleted: 13

Deleted: 12

Deleted: :

Deleted:  $V_d$

Deleted: 11f-i

Deleted: 13e-

An assessment of  $O_3$  vegetation impacts was conducted based on the results from various model cases and different metrics, namely  $POD_y$  (where  $y$  is LULC-dependent critical level) and AOT40. For this demonstration, the 13-day model results were linearly extrapolated to **approximately** three months. This also assumed similar DA adjustments to SM dynamics (driven by factors such as clouds/radiation, rainfall, and irrigation for cropland-dominant regions) at **seasonal** time scale. Based on the **seasonal** variability of surface  $O_3$  and **surface fluxes** in the study region, in 2016 (Figure S6), the linearly scaled  $POD_y$  and AOT40 values **are** overall underestimated referring to the 2016 peak AOT40 and surface fluxes occurring during April-May-June and June-July-August, respectively. These overall underestimations may be invalid if the (sub)seasonal variability of surface  $O_3$  and surface fluxes of other years was referred to. We therefore focus on discussing the results qualitatively and highlighting their implications for  $O_3$  impact assessments using long-term records. Statistics of the derived  $POD_y$  and AOT40 fields are summarized by  $O_3$  sensitive LULC and crop types in Figure J3(c, d, g, h). Figures 15 and J2(c,d) present the

Deleted: ~

Deleted: the

Deleted: of ~three months.

Deleted: known

Deleted:  $V_d$

Deleted: ,

Deleted: may have been

Deleted: 12

Deleted: -

Deleted: ;

Deleted: -

Deleted: ),

Deleted: Figure 14 presents

870 estimated AOT40 fields, [the evaluation of them](#), as well as their responses to the SM DA for cropland-dominant grids. The highs and lows in AOT40-related results are found over maize and wheat dominant fields, respectively. Among the three focused LULC types, the highest and lowest POD<sub>y</sub> values are estimated for forests and grasslands, respectively. Largely driven by daytime peak O<sub>3</sub> concentrations, the spatial variability and biases (referring to AQS and CASTNET observations) of the model-derived AOT40 fields, as well as their responses to the DA, match the MDA8-based (Figure 11). In contrast, the spatial variability of POD<sub>y</sub> and F<sub>s</sub> aligns well, so are their responses to the DA. Both POD<sub>y</sub> and AOT40 reacted several times more intensively in the cases that implemented the dynamic dry deposition scheme, especially the CLM\_D case.

875 For selected LULC and crop types, the WRF-Chem derived POD<sub>y</sub> and AOT40 fields were used together with dose-response functions in literature to evaluate the RBL/RYL due to O<sub>3</sub> exposure and uptake. As reported in Figure [13\(c, g\)](#), with the SM DA enabled, the mean RBLs based on Noah\_D and CLM\_D derived POD<sub>y</sub> are [0.05–0.08, 0.01–0.02](#), and 0.04 for deciduous forest, grasslands and wheat, respectively, which are >33% lower than the Noah\_W and P1\_W based RBL estimates. It is shown that, in response to the DA which lowered SM in many places, the Noah\_W and P1\_W based RBL estimates did not drop as strongly as the Noah\_D and CLM\_D based, and even increased [by 0.01](#) for grasslands and wheat. For wheat, one of 880 the most O<sub>3</sub>-sensitive crops, the estimated RYL values based on [the](#) POD<sub>y</sub> and AOT40 approaches differ by up to a factor of 2–3, and the DA had contrasting effects on these estimates (Figure [13c, d, g, h](#)). The POD<sub>y</sub>- and AOT40-based RYL values differ more significantly when the model-derived POD<sub>y</sub> and AOT40 fields came from the Noah\_D and CLM\_D cases. Using the model-derived AOT40 and different AOT40 dose-response functions (Mills et al., 2007, 2018a, Table [3](#)), the estimated RYLs and their changes due to the DA are nonnegligible (Figure [13d, h](#)). Our estimated RBL/RYL results for various LULC 885 and crop types mostly fall within the ranges reported in previous studies which applied model-derived O<sub>3</sub> metrics and dose-response functions (e.g., Avnery et al., 2011; Mills et al., 2007, 2018a). Our results emphasize that the selected O<sub>3</sub> impact assessment metrics for various LULC/crop types and their matching dose-response functions, as well as the model results used to derive the chosen O<sub>3</sub> metrics which are sensitive to dry deposition schemes and SM, all introduce uncertainty to the estimated O<sub>3</sub> impacts on vegetation. The widely-used dose-response functions are considered appropriate for studying North America 890 and Europe, but they may not be applicable to other regions (Emberson et al., 2009). Therefore, updating and developing dose-response relationships for a larger number of vegetation types in different regions of the world are needed, which may require new experiments to be conducted. Yue and Unger (2014) and Lombardozzi et al. (2015) as well as follow-on investigations parameterized the O<sub>3</sub> impacts on several types of vegetation using the relationships between cumulative O<sub>3</sub> uptake and O<sub>3</sub> damage factors for photosynthesis and conductance from empirical and experimental studies. Based on [multidecadal](#) model 895 simulations, they reported <20% changes of biomass, GPP, and energy fluxes due to O<sub>3</sub>, which are roughly consistent with our RBL/RYL results in Figure [13](#). Such approaches that dynamically assess the impacts of O<sub>3</sub> along with other factors (e.g., non-O<sub>3</sub> pollutants and environmental stresses), as highlighted in Emberson et al. (2018), will be considered in future work.

- Deleted: 12
- Deleted: :
- Deleted: 06
- Deleted: 10
- Deleted: 02
- Deleted: 03
- Deleted: -0.05
- Deleted: 12c-
- Deleted: :
- Deleted: -
- Deleted: 2
- Deleted: 12d;

Deleted: multi-decadal

Deleted: 12

We note that, revising the dry deposition scheme and constraining the modeled SM fields with observations would not only better be combined with adding O<sub>3</sub> injury to vegetation but also multistress impacts on biogenic emissions. Considering O<sub>3</sub> injury to vegetation would affect more evidently longer-term climate simulations via feedbacks to biomass, surface fluxes, weather and weather-driven emissions. As for biogenic emissions, Figure S7 shows SM anomalies during the study period determined by our Noah-MP modeling system as well as drought stress activity factor  $\gamma_d$  estimated from  $\beta$  of a multiyear, independent CLM (version 4.5) simulation by Jiang et al. (2018). Based on these, we estimate that, depending on soil type, hydrological regime, as well as  $\beta$  configurations, omitting the direct impacts of water stress on biogenic emissions, may have introduced larger uncertainty (i.e., >30%) to biogenic emission and O<sub>3</sub> modeling over several states experiencing drier-than-normal conditions, particularly South Carolina, Georgia, and Alabama. Quantitatively understanding the interplay between these processes and O<sub>3</sub> pollution levels is recommended for more accurate air quality modeling and O<sub>3</sub> impact assessments.

Deleted: S5

Deleted: such as Tennessee,

Deleted: , and West Virginia

#### 4 Summary and suggestions on future directions

This paper described a follow-up study of Huang et al. (2021). It presented how the choice of O<sub>3</sub> dry deposition scheme affected our evaluation of SMAP SM DA impacts on coupled WRF-Chem modeling over the southeastern US in August 2016. In new Noah-MP LSM related simulations, two dry deposition schemes were implemented, namely the WRF-Chem default Wesely scheme and a dynamic scheme, in the latter of which the calculation of  $\gamma_d$  (particularly its stomatal and cuticular terms) was modified to be coupled with photosynthesis and vegetation phenology. We showed that dry deposition parameterizations significantly affected the modeled O<sub>3</sub> dry deposition process, as well as its response to the DA. Comparing the no-DA cases, it was found that, when the dynamic scheme was applied, overall, the modeled O<sub>3</sub> dry deposition velocities and fluxes were larger and surface O<sub>3</sub> concentrations were lower. The modeled O<sub>3</sub> fluxes responded 2–3 times more strongly to the SM changes due to the DA, which can be mainly explained by the fact that both the direct and indirect (i.e., via influencing weather and vegetation fields) effects of SM on O<sub>3</sub> dry deposition modeling are considered in the dynamic scheme. Depending on soil type and hydrological regime, the selection of SM factor controlling  $\gamma_s$  (i.e.,  $\beta$  factor, a key variable representing the direct effects of SM on the modeled surface fluxes) scheme can strongly affect the quantitative results. The Wesely-scheme derived dry deposition results driven by meteorological fields from Noah-MP and Noah (from Huang et al., 2021) LSM based WRF-Chem simulations displayed much smaller differences than those due to updating the dry deposition parameterizations. While we note that accounting for physiological effects in dry deposition modeling can be beneficial, the Ball-Berry  $\gamma_s$  scheme applied in land surface and dry deposition modeling in this work needs to be compared with other semi-empirical  $\gamma_s$  schemes, for a better understanding of their respective strengths and weaknesses. Alternative schemes include the Medlyn scheme which has been integrated into the CLM version 5. Model intercomparison efforts such as the ongoing AQMEII4 activity (Galmarini et al., 2021) can also help determine areas for improvement in commonly-used dry deposition modeling approaches for studying 2016 and other years, over North America and other regions of the world.

Deleted:  $\gamma_d$

Deleted: were

Deleted:  $R_s$

Deleted:  $R_s$

Deleted:  $R_s$

By analyzing the model responses to the SM DA from these various cases, we conclude that, in coupled modeling systems that consider the direct and indirect influences of SM on O<sub>3</sub> dry deposition, the accuracy of SM is particularly critical to dry deposition and O<sub>3</sub> modeling, as well as the scientific analyses and impact assessments based on model simulations. The usefulness of SM DA for improving the modeled state and flux variables was evaluated by multiple observation(-derived) data products. Referring to in situ measurements, key meteorological variables relevant to  $\psi_d$  calculations such as surface temperature and humidity are shown to be improved by the DA by up to ~9%. Referring to satellite(-derived) datasets which may be associated with high uncertainty, the model performance of vegetation phenology, GPP, as well as energy fluxes and their partitioning, showed mixed, LULC-dependent reactions to the DA. According to the evaluation statistics, for this case, the CLM<sub>4</sub> type  $\beta$  factor scheme was slightly superior to the Noah<sub>1.2</sub> type *one*. The modeled carbon and energy fields as well as their DA-related changes, correlated strongly with the modeled  $\psi_d$  fields, implying that the DA impacts on the accuracy of  $\psi_d$  were also possibly complicated which is difficult to verify due to the lack of high-accuracy, independent  $\psi_d$  evaluation datasets. [a point that has also been brought up in previous dry deposition modeling works \(e.g., Baublitz et al., 2020; Clifton et al., 2020\)](#). Observation(-derived)  $\psi_d$  datasets covering diverse LULC types nested in broad geographical regions and through more recent periods are in strong need. In places, the likely ineffectiveness of SM DA on vegetation and surface fluxes can not only be attributed to the quality of satellite SM retrievals and the used DA approach as discussed in previous Noah LSM based DA experiments, but also shortcomings in the Noah-MP LSM and its dynamic vegetation scheme regarding its surface-subsurface coupling and representation of SM-vegetation growth feedbacks. Continued efforts on advancing land measurement/retrieval skills, identifying and addressing deficits in LSMs as well as practicing multivariate land DA are recommended in future work.

This study also demonstrated that, model-driven assessments of O<sub>3</sub> impacts on human health and various types of vegetation can be significantly affected by the applied O<sub>3</sub> dry deposition scheme, the implementation of land DA, the chosen O<sub>3</sub> metrics and their matching exposure-response functions. Various model cases showed that, the DA impacts on MDA8 were more evident in nonurban areas where the mean MDA8 was ~5 ppbv lower and the averaged population density is <1/25 of those in urban areas. Using concentration- and flux-based metrics AOT40 and POD<sub>y</sub>, the mean RYLs of maize, soybean, and wheat fell within ranges of 0.01–0.04, 0.10–0.17, and 0.04–0.14, respectively. [The multiple no-DA and DA cases helped us better understand the indirect or/and direct effects of SM on O<sub>3</sub> dry deposition process, which have important implications for O<sub>3</sub> impact assessments. It is also recognized that, the DA often exacerbated the positive surface O<sub>3</sub> biases in free-running systems which has been a common issue shared by numerous regional and global models for this study region/season. It is necessary to combine land DA with efforts to identify, quantify, and reduce other sources of uncertainty in O<sub>3</sub> modeling. These should include reasonably representing the impacts of O<sub>3</sub> along with other factors on vegetation, the direct impacts of water stress on biogenic emissions of volatile organic compounds and nitrogen species, as well as the reduction of photolysis reaction rates and the modification of vertical transport due to the presence of foliage \(Li et al., 2016; Jiang et al., 2018; Makar et al., 2017\).](#)

Deleted:  $\psi_d$

Deleted:

Deleted: of

Deleted:

Deleted:  $\psi_d$

Deleted:  $\psi_d$

Deleted:  $\psi_d$

Deleted: .

Deleted:  $\psi_d$

Deleted: O<sub>3</sub>

Deleted: O<sub>7</sub>

Deleted: While the

Deleted: has

Deleted: , it

Deleted: as well as

Deleted: .

### Code and data availability

Dry deposition related updates to LIS/WRF-Chem since Huang et al. (2021) are undergoing reporting processes via NASA's New Technology Reporting System (<https://invention.nasa.gov>). Observations and observation-derived [evaluation](#) data sets [emphasized](#) in this work but not in Huang et al. (2021) can be found at the following locations:  
1005 <https://land.copernicus.eu/global/products/fcover> (Copernicus Global Land Service, 2020, last access: 10 April 2022)  
<https://doi.org/10.5067/L6C9EY1O8VIC> (Kimball et al., 2021), <https://doi.org/10.7927/H49C6VHW> (NASA Socioeconomic Data and Applications Center, 2018), <https://www-air.larc.nasa.gov/cgi-bin/ArcView/actamerica.2016> (NASA, 2020, last access: 8 November 2021), <https://java.epa.gov/castnet/clearsession.do> (US Environmental Protection Agency, 2021, last access: 8 November 2021), and <https://doi.org/10.3334/ORNLDAAC/1696> (Yu et al., 2019).

Deleted: used

Deleted: 2019

### 1010 Author contributions

MH led the design and execution of the study as well as the paper writing, benefitting from discussions with JHC, GRC, KWB, and SVK, with the feedback from the Atmos. Chem. Phys. Editorial Board and reviewers for Huang et al. (2021) also accounted for. CS contributed to data collection during the ACT-America campaign. All authors helped finalize the paper.

### Competing interests

1015 The authors declare that they have no conflict of interest.

### Acknowledgements

We acknowledge NASA SUSMAP sponsorship, as well as ACT-America Science Team and NASA's high-end computing systems and services at Ames and Goddard. We thank Jiang et al. (2018) for developing the  $\gamma_d$  dataset shown in Figure S7. We also greatly appreciate active and relevant discussions with multiple colleagues from the Air Quality Model Evaluation  
1020 International Initiative 4 and the Tropospheric Ozone Assessment Report II communities during and after recent conferences and workshops, particularly: C. Hogrefe, J. Pleim, P. Makar, L. Emberson, B. Sinha, D. Lombardozzi, O. Clifton, L. Emmons, T. Emmerichs, and D. Taraborrelli.

Deleted: S5

## References

Anav, A., Proietti, C., Menut, L., Camicelli, S., De Marco, A., and Paoletti, E.: Sensitivity of stomatal conductance to soil moisture: implications for tropospheric ozone, *Atmos. Chem. Phys.*, 18, 5747–5763, <https://doi.org/10.5194/acp-18-5747-2018>, 2018.

1030

Avnery, S., Mauzerall, D. L., Liu, J., and Horowitz, L. W.: Global Crop Yield Reductions due to Surface Ozone Exposure: 1. Year 2000 Crop Production Losses and Economic Damage, *Atmos. Environ.*, 45, 2284–2296, <https://doi.org/10.1016/j.atmosenv.2010.11.045>, 2011.

1035

Baillie, A. L. and Fleming, A. J.: The developmental relationship between stomata and mesophyll airspace, *New Phytol.*, 225, 1120–1126, <https://doi.org/10.1111/nph.16341>, 2019.

Ball, J. T., Woodrow, I. E., and Berry, J. A. (1987), A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, *Process in Photosynthesis Research*, 1, edited by J. Biggins, 221–234, Martinus Nijhoff, Dordrecht, the Netherlands.

1040

~~Baublitz, C. B., Fiore, A. M., Clifton, O. E., Mao, J., Li, J., Correa, G., Westervelt, D. M., Horowitz, L. W., Paulot, F., and Williams, A. P.: Sensitivity of Tropospheric Ozone Over the Southeast USA to Dry Deposition, *Geophys. Res. Lett.*, 47, e2020GL087158, <https://doi.org/10.1029/2020GL087158>, 2020.~~

1045



~~Clifton, O. E., Fiore, A. M., Munger, J. W., Malyshev, S., Horowitz, L. W., Shevliakova, E., Paulot, F., Murray, L. T., and Griffin, K. L.: Interannual variability in ozone removal by a temperate deciduous forest, *Geophys. Res. Lett.*, 44, 542–552, <https://doi.org/10.1002/2016GL070923>, 2017.~~

1050

Clifton, O. E., Paulot, F., Fiore, A. M., Horowitz, L. W., Correa, G., Baublitz, C. B., Fares, S., Goded, I., Goldstein, A. H., Gruening, C., Hogg, A. J., Loubet, B., Mammarella, I., Munger, J. W., Neil, L., Stella, P., Uddling, J., Vesala, T., and Weng, E.: Influence of dynamic ozone dry deposition on ozone pollution, *J. Geophys. Res.-Atmos.*, 125, e2020JD032398, <https://doi.org/10.1029/2020JD032398>, 2020.

1055

Convention on Long-Range Transboundary Air Pollution (CLRTAP): Mapping Critical Levels for Vegetation, Chapter 3 of Manual for modelling and mapping critical loads and levels, available at: <https://www.umweltbundesamt.de/en/manual-for-modelling-mapping-critical-loads-levels> (last access: 10 December 2021), 2017.

- Moved down [2]: R.,
- Moved down [3]: T.,
- Moved (insertion) [4]
- Moved (insertion) [5]
- Moved (insertion) [6]
- Deleted: Bunk,
- Deleted: Behrendt,
- Deleted: Yi, Z., Andrea,
- Deleted: .
- Moved (insertion) [7]
- Deleted: Kesselmeier, J. (2017), Exchange
- Deleted: carbonyl sulfide (OCS) between soils and atmosphere under various CO<sub>2</sub> concentrations, J.
- Moved (insertion) [8]
- Moved (insertion) [9]
- Moved (insertion) [3]
- Moved up [7]: Res.
- Deleted: Biogeosci., 122, 1343–1358
- Deleted: 2016JG003678. 
- Campbell, J.
- Moved up [4]: E.,
- Moved down [10]: T.,
- Moved up [5]: M.,
- Moved down [11]: J.,
- Deleted: Carmichael, G. R., Chai,
- Deleted: Mena-Carrasco,
- Deleted: Tang, Y., Blake, D. R., Blake, N. J., Vay, S. A., Collatz, G.,
- Deleted: Baker, I., Berry, J. A., Montzka, S. A., Sweeney, C., Schnoor, J. L., and Stanier, C. O.: Photosynthetic control of atmospheric carbonyl sulfide during the growing season, *Science*, 322, 1085–1088, <https://doi.org/10.1126/science.1164015>, 2008. 

1085 [Copernicus Global Land Service: Fraction of green vegetation cover 1 km version 2 data, Flemish Institute for Technological Research on behalf of the European Commission Joint Research Centre \[data set\], Boeretang, Belgium, available at: https://land.copernicus.eu/global/products/fcover \(last access: 10 April 2022\), 2020.](https://land.copernicus.eu/global/products/fcover)

Dieter, C. A., Maupin, M. A., Caldwell, R. R., Harris, M. A., Ivahnenko, T. I., Lovelace, J. K., Barber, N. L., and Linsey, K. S., Estimated use of water in the United States in 2015: U. S. Geological Survey Circular 1441, 65, 1090 <https://doi.org/10.3133/cir1441>, 2018.

Ducker, J. A., Holmes, C. D., Keenan, T. F., Fares, S., Goldstein, A. H., Mammarella, I., Munger, J. W., and Schnell, J.: Synthetic ozone deposition and stomatal uptake at flux tower sites, *Biogeosciences*, 15, 5395–5413, <https://doi.org/10.5194/bg-15-5395-2018>, 2018. 1095

Egea, G., Verhoef, A., and Vidale, P. L.: Towards an improved and more flexible representation of water stress in coupled photosynthesis–stomatal conductance models, *Agric. For. Meteorol.*, 151, 1370–1384. <https://doi.org/10.1016/j.agrformet.2011.05.019>, 2011. 1100

Emberson, L. D., Büker, P., Ashmore, M. R., Mills, G., Jackson, L., Agrawal, M., Atikuzzaman, M. D., Cinderby, S., Engardt, M., Jamir, C., Kobayashi, K., Oanh, N. T. K., Quadir, Q. F., and Wahid, A.: A comparison of North American and Asian exposure-response data for ozone effects on yield, *Environ. Pollut.*, 43, 1945–1953, <https://doi.org/10.1016/j.atmosenv.2009.01.005>, 2009. 1105

Emberson, L. D., Pleijel, H., Ainsworth, E. A., van den Berg, M., Ren, W., Osborne, S., Mills, G., Pandey, D., Dentener, F., Büker, P., Ewert, F., Koeble, R. and Van Dingenen, R.: Ozone effects on crops and consideration in crop models, *Eur. J. Agron.*, 100, 19–34, <https://doi.org/10.1016/j.eja.2018.06.002>, 2018.

Emmerichs, T., Kerkweg, A., Ouwersloot, H., Fares, S., Mammarella, I., and Taraborrelli, D.: A revised dry deposition scheme for land–atmosphere exchange of trace gases in ECHAM/MESSy v2.54, *Geosci. Model Dev.*, 14, 495–519, <https://doi.org/10.5194/gmd-14-495-2021>, 2021. 1110

[Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., Entin, J. K., Goodman, S. D., Jackson, T. J., Johnson, J., Kimball, J., Piepmeier, J. R., Koster, R. D., Martin, N., McDonald, K. C., Moghaddam, M., Moran, S., Reichle, R., Shi, J. C., Spencer, M. W., Thurman, S. W., Tsang, L., and van Zyl, J.: The Soil Moisture Active Passive \(SMAP\) Mission, \*Proc. IEEE\*, 98, 704–716, <https://doi.org/10.1109/JPROC.2010.2043918>, 2010.](#) 1115

Moved (insertion) [10]

Moved (insertion) [11]

Moved (insertion) [2]

Moved (insertion) [12]



1120 Falk, S. and Søvdé Haslerud, A.: Update and evaluation of the ozone dry deposition in Oslo CTM3 v1.0, *Geosci. Model Dev.*,  
12, 4705–4728, <https://doi.org/10.5194/gmd-12-4705-2019>, 2019.

Finkelstein, P. L., Ellestad, T. G., Clarke, J. F., Meyers, T. P., Schwede, D. B., Hebert, E. O., and Neal, J. A.: Ozone and sulfur  
dioxide dry deposition to forests: Observations and model evaluation, *J. Geophys. Res.*, 105, 15365–15377,  
doi:10.1029/2000JD900185, 2000.

1125 Fleming, Z. L., Doherty, R. M., von Schneidmesser, E., Malley, C. S., Cooper, O. R., Pinto, J. P., Colette, A., Xu, X., Simpson,  
D., Schultz, M. G., Lefohn, A. S., Hamad, S., Moolla, R., Solberg, S., and Feng, Z.: Tropospheric Ozone Assessment Report:  
Present-day ozone distribution and trends relevant to human health, *Elem. Sci. Anth.*, 6, 12,  
<https://doi.org/10.1525/elementa.273>, 2018.

1130 Fredericksen, T. S., Skelly, J. M., Steiner, K. C., Kolb, T. E., Kouterick, K. B.: Size-mediated foliar response to ozone in black  
cherry trees, *Environ. Pollut.*, 91, 53–63, [https://doi.org/10.1016/0269-7491\(95\)00032-m](https://doi.org/10.1016/0269-7491(95)00032-m), 1996.

1135 [Fu, J. S., Carmichael, G. R., Dentener, F., Aas, W., Andersson, C., Barrie, I., A., Cole, A., Galy-Lacaux, C., Geddes, J.,  
Itahashi, S., Kanakidou, M., Labrador, L., Paulot, F., Schwede, D., Tan, J., and Vet, R., \*Environ. Sci. Technol.\*, 56, 2134–  
2142, <https://doi.org/10.1021/acs.est.1c05929>, 2022.](https://doi.org/10.1021/acs.est.1c05929)

Galmarini, S., Makar, P., Clifton, O. E., Hogrefe, C., Bash, J. O., Bellasio, R., Bianconi, R., Bieser, J., Butler, T., Ducker, J.,  
Flemming, J., Hodzic, A., Holmes, C. D., Kioutsioukis, I., Kranenburg, R., Lupascu, A., Perez-Camanyo, J. L., Pleim, J., Ryu,  
1140 Y.-H., San Jose, R., Schwede, D., Silva, S., and Wolke, R.: Technical note: AQMEII4 Activity 1: evaluation of wet and dry  
deposition schemes as an integral part of regional-scale air quality models, *Atmos. Chem. Phys.*, 21, 15663–15697,  
<https://doi.org/10.5194/acp-21-15663-2021>, 2021.

Griffiths, P. T., Murray, L. T., Zeng, G., Shin, Y. M., Abraham, N. L., Archibald, A. T., Deushi, M., Emmons, L. K., Galbally,  
1145 I. E., Hassler, B., Horowitz, L. W., Keeble, J., Liu, J., Moeini, O., Naik, V., O'Connor, F. M., Oshima, N., Tarasick, D., Tilmes,  
S., Turnock, S. T., Wild, O., Young, P. J., and Zanis, P.: Tropospheric ozone in CMIP6 simulations, *Atmos. Chem. Phys.*, 21,  
4187–4218, <https://doi.org/10.5194/acp-21-4187-2021>, 2021.

Hardacre, C., Wild, O., and Emberson, L.: An evaluation of ozone dry deposition in global scale chemistry climate models,  
1150 *Atmos. Chem. Phys.*, 15, 6419–6436, <https://doi.org/10.5194/acp-15-6419-2015>, 2015.

Moved (insertion) [13]

Moved (insertion) [14]

- Hemispheric Transport of Air Pollution: Impacts on Health, Ecosystems, and Climate, Chapter 5 of 2010 Final Assessment report, Part A: Ozone and particulate matter, available at: [http://htapold.kaskada.tk/publications/2010\\_report/2010\\_Final\\_Report/HTAP\\_2010\\_Part\\_A\\_110407.pdf](http://htapold.kaskada.tk/publications/2010_report/2010_Final_Report/HTAP_2010_Part_A_110407.pdf) (last access: 16 November 2021), 2010.
- Hollaway, M. J., Arnold, S. R., Challinor, A. J., and Emberson, L. D.: Intercontinental trans-boundary contributions to ozone-induced crop yield losses in the Northern Hemisphere, *Biogeosciences*, 9, 271–292, <https://doi.org/10.5194/bg-9-271-2012>, 2012.
- Huang, M., Carmichael, G. R., Chai, T., Pierce, R. B., Oltmans, S. J., Jaffe, D. A., Bowman, K. W., Kaduwela, A., Cai, C., Spak, S. N., Weinheimer, A. J., Huey, L. G., and Diskin, G. S.: Impacts of transported background pollutants on summertime western US air quality: model evaluation, sensitivity analysis and data assimilation, *Atmos. Chem. Phys.*, 13, 359–391, <https://doi.org/10.5194/acp-13-359-2013>, 2013.
- Huang, M., Crawford, J. H., DiGangi, J. P., Carmichael, G. R., Bowman, K. W., Kumar, S. V., and Zhan, X.: Satellite soil moisture data assimilation impacts on modeling weather variables and ozone in the southeastern US – Part 1: An overview, *Atmos. Chem. Phys.*, 21, 11013–11040, <https://doi.org/10.5194/acp-21-11013-2021>, 2021.
- Intergovernmental Panel on Climate Change: the Sixth Assessment Report, Summary for Policymakers, available at: <https://www.ipcc.ch/report/ar6/wg1> (last access: 24 November 2021), 2021.
- Jiang, X., Guenther, A., Potosnak, M., Geron, C., Seco, R., Karl, T., Kim, S., Gu, L., and Pallardy, S.: Isoprene emission response to drought and the impact on global atmospheric chemistry, *Atmos. Environ.*, 183, 69–83, <https://doi.org/10.1016/j.atmosenv.2018.01.026>, 2018.
- Kimball, J. S., Jones, L. A., Endsley, A., Kundig, T., and Reichle, R.: SMAP L4 Global Daily 9 km EASE-Grid Carbon Net Ecosystem Exchange, Version 6, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], Boulder, Colorado, USA, <https://doi.org/10.5067/L6C9EY1O8VIC>, 2021.
- Kumar, S. V., Reichle, R. H., Koster, R. D., Crow, W. T., and Peters-Lidard, C. D.: Role of subsurface physics in the assimilation of surface soil moisture observations, *J. Hydrometeorol.*, 10, 1534–1547, <https://doi.org/10.1175/2009JHM1134.1>, 2009.

185 [Lacis, A. A., Wuebbles, D. J., and Logan, J. A.: Radiative forcing of climate by changes in the vertical distribution of ozone, J. Geophys. Res., 95, 9971–9981, <https://doi.org/10.1029/JD095iD07p09971>, 1990.](#)

Moved (insertion) [15]

Lapina, K., Henze, D. K., Milford, J. B., Huang, M., Lin, M., Fiore, A. M., Carmichael, G., Pfister, G. G., and Bowman, K.: Assessment of source contributions to seasonal vegetative exposure to ozone in the U.S., J. Geophys. Res.-Atmos., 119, 324–340, <https://doi.org/10.1002/2013JD020905>, 2014.

Lawston, P. M., Santanello, J. A., Zaitchik, B. F., and Rodell, M.: Impact of irrigation methods on land surface model spinup and initialization of WRF forecasts, J. Hydrometeor., 16, 1135–1154, <https://doi.org/10.1175/JHM-D-14-0203.1>, 2015.

195 [Li, J., Mahalov, A., and Hyde, P.: Simulating the impacts of chronic ozone exposure on plant conductance and photosynthesis, and on the regional hydroclimate using WRF/Chem, Environ. Res. Lett., 11, 114017, <http://dx.doi.org/10.1088/1748-9326/11/11/114017>, 2016.](#)

Deleted: Lennartz, S. T., Marandino, C

1200 [Lin, M., Malyshev, S., Shevliakova, E., Paulot, F., Horowitz, L. W., Fares, S., Mikkelsen, T. N., and Zhang, L.: Sensitivity of ozone dry deposition to ecosystem-atmosphere interactions: A critical appraisal of observations and simulations, Global Biogeochem. Cy., 33, 1264–1288, <https://doi.org/10.1029/2018GB006157>, 2019.](#)

Moved up [14]: . A.,

Moved up [9]: M.,

Moved up [13]: R.,

Moved up [12]: C.,

Moved up [8]: E.,

Deleted: von Hobe,

Deleted: Cortes, P., Quack, B., Simo,

Deleted: Booge, D., Pozzer, A., Steinhoff, T., Arevalo-Martinez, D. L., Kloss,

Deleted: Bracher, A., Röttgers, R., Atlas,

Deleted: and Krüger, K.: Direct oceanic emissions unlikely to account for the missing source of atmospheric carbonyl sulfide, Atmos. Chem. Phys., 17, 385–402, <https://doi.org/10.5194/acp-17-385-2017>, 2017.

Lin, M., Horowitz, L.W., Xie, Y., Paulot, F., Malyshev, S., Shevlickova, E., Finco, A., Gerosa, G., Kubinstin, D., and Pilegaard, K.: Vegetation feedbacks during drought exacerbate ozone air pollution extremes in Europe, Nat. Clim. Chang., 10, 444–451, <https://doi.org/10.1038/s41558-020-0743-y>, 2020.

Lombardozi, D., Levis, S., Bonan, G., Hess, P. G., and Sparks, J. P.: The Influence of Chronic Ozone Exposure on Global Carbon and Water Cycles, J. Climate, 28, 292–305, <https://doi.org/10.1175/JCLI-D14-00223.1>, 2015.

210 [Makar, P. A., Staebler, R. M., Akingunola, A., Zhang, J., McLinden, C., Kharol, S. K., Pabla, B., Cheung, P., and Zheng, Q.: The effects of forest canopy shading and turbulence on boundary layer ozone, Nat. Commun., 8, 1–14, <https://doi.org/10.1038/ncomms15243>, 2017.](#)

1215 Mills, G., Buse, A., Gimeno, B., Bemejo, V., Holland, M., Emberson, L., and Pleijel, H.: A synthesis of AOT40-based response functions and critical levels of ozone for agricultural and horticultural crops, Atmos. Environ., 41, 2630–2643, <https://doi.org/10.1016/j.atmosenv.2006.11.016>, 2007.

- 1235 Mills, G., Hayes, F., Simpson, D., Emberson, L., Norris, D., Harmens, H., and Büker, P.: Evidence of widespread effects of ozone on crops and (semi-) natural vegetation in Europe (1990–2006) in relation to AOT40-and flux-based risk maps, *Global Change Biol.*, 17, 592–613, <https://doi.org/10.1111/j.1365-2486.2010.02217.x>, 2011.
- 1240 Mills, G., Sharps, K., Simpson, D., Pleijel, H., Broberg, M., Uddling, J., Jaramillo, F., Davies, W. J., Dentener, F., van den Berg, M., Agrawal, M., Agrawal, S. B., Ainsworth, E. A., Buker, P., Emberson, L., Feng, Z., Harmens, H., Hayes, F., Kopbayashi, K., Paoletti, E., and Van Dingenen, R.: Ozone pollution will compromise efforts to increase global wheat production, *Global Change Biol.*, 24, 3560–3574, <https://doi.org/10.1111/gcb.14157>, 2018a.
- 1245 Mills, G., Pleijel, H., Malley, C. S., Sinha, B., Cooper, O. R., Schultz, M. G., Neufeld, H. S., Simpson, D., Sharps, K., Feng, Z., Gerosa, G., Harmens, H., Kobayashi, K., Saxena, P., Paoletti, E., Sinha, V., Xu, X.: Tropospheric Ozone Assessment Report: Present-day tropospheric ozone distribution and trends relevant to vegetation, *Elem. Sci. Anth.*, 6, 47, <https://doi.org/10.1525/elementa.302>, 2018b.
- 1250 Monfreda, C., Ramankutty, N., and Foley, J. A.: Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Global Biogeochem. Cycles*, 22, GB1022, <https://doi.org/10.1029/2007GB002947>, 2008.
- NASA: ACT-America 2016 1-minute Merged B-200/C-130 Data, Version R1/R4, NASA Langley Research Center Airborne Science Data for Atmospheric Composition [data set], Hampton, Virginia, USA, available at: <https://www-air.larc.nasa.gov/cgi-bin/ArcView/actamerica.2016>, (last access: 8 November 2021), 2020.
- 1255 NASA Socioeconomic Data and Applications Center: Gridded Population of the World, Version 4.11, Columbia University Center for International Earth Science Information Network [data set], Palisades, New York, USA, <https://doi.org/10.7927/H49C6VHW>, 2018.
- 1260 Niu, G. Y., Yang, Z. L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D., Rosero, E., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements, *J. Geophys. Res.-Atmos.*, 116, 1–19, <https://doi.org/10.1029/2010JD015139>, 2011.
- 1265 Niyogi, D. S. and Raman, S.: Comparison of Four Different Stomatal Resistance Schemes Using FIFE Observations, *J. Appl. Meteorol. Climatol.*, 36, 903–917, [https://doi.org/10.1175/1520-0450\(1997\)036%3C0903:COFDSR%3E2.0.CO;2](https://doi.org/10.1175/1520-0450(1997)036%3C0903:COFDSR%3E2.0.CO;2), 1997.

Deleted: .

- 1270 [Otu-Larbi, F., Conte, A., Fares, S., Wild, O., and Ashworth, K.: Current and future impacts of drought and ozone stress on Northern Hemisphere forests, \*Global Change Biol.\*, 26, 6218–6234, <https://doi.org/10.1111/gcb.15339>, 2020.](#)
- Ronan, A. C., Ducker, J. A., Schnell, J. L., and Holmes, C. D.: Have improvements in ozone air quality reduced ozone uptake into plants? *Elem. Sci. Anth.*, 8, 2, <https://doi.org/10.1525/elementa.399>, 2020.
- 1275 Salmon, J. M., Friedl, M. A., Frohling, S., Wisser, D., and Douglas, E. M.: Global rain-fed, irrigated, and paddy croplands: A new high resolution map derived from remote sensing, crop inventories and climate data, *Int. J. Appl. Earth Obs.*, 38, 321–334, <https://doi.org/10.1016/j.jag.2015.01.014>, 2015.
- 1280 Saylor, R. D., Wolfe, G. M., Meyers, T. P., Hicks, B. B.: A corrected formulation of the Multilayer Model (MLM) for inferring gaseous dry deposition to vegetated surfaces, *Atmos. Environ.*, 92, 141–145. <https://doi.org/10.1016/j.atmosenv.2014.03.056>, 2014.
- Silva, S. J. and Heald, C. L.: Investigating dry deposition of ozone to vegetation, *J. Geophys. Res.-Atmos.*, 123, 559–573, <https://doi.org/10.1002/2017JD027278>, 2018.
- 1285 [Steinkamp, J. and Lawrence, M. G.: Improvement and evaluation of simulated global biogenic soil NO emissions in an AC-GCM, \*Atmos. Chem. Phys.\*, 11, 6063–6082, <https://doi.org/10.5194/acp-11-6063-2011>, 2011.](#)
- 1290 Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J. F., Lawrence, M. G., Montanaro, V., Müller, J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K., Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res. Atmos.*, 111, <https://doi.org/10.1029/2005JD006338>, 2006.
- 1295 [Strode, S. A., Rodriguez, J. M., Logan, J. A., Cooper, O. R., Witte, J. C., Lamsal, L. N., Damon, M., Van Aartsen, B., Steenrod, S. D., and Strahan, S. E.: Trends and variability in surface ozone over the United States, \*J. Geophys. Res. Atmos.\*, 120, 9020–9042, <https://doi.org/10.1002/2014JD022784>, 2015.](#)
- 1300 Sweeney, C., Karion, A., Wolter, S., Newberger, T., Guenther, D., Higgs, J. A., Andrews, A. E., Lang, P. M., Neff, D., Dlugokencky, E., Miller, J. B., Montzka, S. A., Miller, B. R., Masarie, K. A., Biraud, S. C., Novelli, P. C., Crotnell, M., Crotnell, A. M., Thoning, K., and Tans, P. P.: Seasonal climatology of CO<sub>2</sub> across North America from aircraft measurements

- in the NOAA/ESRL Global Greenhouse Gas Reference Network, *J. Geophys. Res.-Atmos.*, 120, 5155–5190, <https://doi.org/10.1002/2014jd022591>, 2015.
- 1305 US Environmental Protection Agency: CASTNET Historical Deposition Data, US EPA Office of Atmospheric Programs [data set], Research Triangle Park, North Carolina, USA, available at: <https://java.epa.gov/castnet/clearsession.do>, last access: 8 November 2021.
- 1310 Val Martin, M., Heald, C. L., and Arnold, S. R.: Coupling dry deposition to vegetation phenology in the Community Earth System Model: Implications for the simulation of surface O<sub>3</sub>, *Geophys. Res. Lett.*, 41, 2988–2996, <https://doi.org/10.1002/2014GL059651>, 2014.
- Van Dingenen, R., Dentener, F., Raes, F., Krol, M. C., Emberson, L., and Cofala, J.: The global impact of ozone on agricultural crop yields under current and future air quality legislation, *Atmos. Environ.*, 43, 604–618, <https://doi.org/10.1016/j.atmosenv.2008.10.033>, 2009.
- 1315 Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.*, 41, 52–63, <https://doi.org/10.1016/j.atmosenv.2007.10.058>, 1989.
- 1320 Whelan, M. E., Anderegg, L. D., Badgley, G., Campbell, J. E., Commene, R., Frankenberg, C., Hilton, T. W., Kuai, L., Parazoo, N., Shiga, Y., Wang, Y., and Worden, J.: Scientific Communities Striving for a Common Cause: Innovations in Carbon Cycle Science, *Bull. Amer. Meteor.*, 101, E1537–E1543, <https://doi.org/10.1175/BAMS-D-19-0306.1>, 2020.
- 1325 [Wickham, J., Stehman, S. V., Sorenson, D. G., Gass, L., and Dewitz, J. A.: Thematic accuracy assessment of the NLCD 2016 land cover for the conterminous United States, \*Remote Sens. Environ.\*, 257, 2021, 112357, <https://doi.org/10.1016/j.rse.2021.112357>, 2021.](https://doi.org/10.1016/j.rse.2021.112357)
- 1330 [Wong, A. Y. H., Geddes, J. A., Tai, A. P. K., and Silva, S. J.: Importance of dry deposition parameterization choice in global simulations of surface ozone, \*Atmos. Chem. Phys.\*, 19, 14365–14385, <https://doi.org/10.5194/acp-19-14365-2019>, 2019.](https://doi.org/10.5194/acp-19-14365-2019)
- Wu, Z., Schwede, D. B., Vet, R., Walker, J. T., Shaw, M., Staebler, R., and Zhang, L.: Evaluation and Intercomparison of Five North American Dry Deposition Algorithms at a Mixed Forest Site, *J. Adv. Model. Earth Sy.*, 10, 1571–1586, <https://doi.org/10.1029/2017MS001231>, 2018.
- 1335

Yang, Z.-L., Niu, G.-Y., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Longuevergne, L., Manning, K., Niyogi, D., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 2. Evaluation over global river basins, *J. Geophys. Res.*, 116, D12110, <https://doi.org/10.1029/2010JD015140>, 2011.

1340 Yu, L., Wen, J., Chang, C. Y., Frankenberg, C., and Y. Sun: High Resolution Global Contiguous SIF Estimates Derived from OCO-2 SIF and MODIS, Oak Ridge National Laboratory Distributed Active Archive Center [data set], Oak Ridge, Tennessee, USA, <https://doi.org/10.3334/ORNLDAAAC/1696>, 2019.

1345 Yue, X. and Unger, N.: Ozone vegetation damage effects on gross primary productivity in the United States, *Atmos. Chem. Phys.*, 14, 9137–9153, <https://doi.org/10.5194/acp-14-9137-2014>, 2014.

## Tables

**Table 1: Model cases and their configurations relevant to the discussions of this study.**

Case name	Land surface model	Stomatal resistance scheme	SM factor controlling $f_s$ ( $\beta$ )	Surface exchange coefficient for heat ( $C_H$ ) scheme	Irrigation scheme	Dry deposition scheme	Note
Noah_D	Noah-MP	Ball-Berry	Noah-type	Monin-Obukhov	Sprinkler	Dynamic	new in this study
CLM_D	Noah-MP	Ball-Berry	CLM <sub>v</sub> (version 4.5)-type	Monin-Obukhov	Sprinkler	Dynamic	
Noah_W	Noah-MP	Ball-Berry	Noah-type	Monin-Obukhov	Sprinkler	Wesely	
P1_W	Noah	Jarvis	Noah	Chen97	not included	Wesely	from Part 1

Moved up [15]: A.,

Moved up [6]: . W.,

Deleted: ¶  
Zumkehr,

Deleted: Hilton, T

Deleted: Whelan, M. E., Smith, S., Kuai, L., Worden, J., and Campbell, J. E.: Global Gridded Anthropogenic Emissions Inventory of Carbonyl Sulfide, *Atmos. Environ.*, 183, 11–19, <https://doi.org/10.1016/j.atmosenv.2018.03.063>, 2018.¶

Deleted:  $R_s$

Deleted: -

**Table 2: Evaluation datasets relevant to this study, along with their key attributes. References of these products can be found in the data availability sections of this work and Huang et al. (2021).**

Measurement platform, network or name of dataset	Measured or derived variable	Type of dataset	Spatial resolution	Temporal resolution; coverage of the used dataset	Note
SMAP	VOD	satellite retrieval	9 km	twice-daily; morning time data during August 2015–2019	new in this study, but available in the SMAP enhanced product introduced in part 1
SMAP L4C	GPP	observation derived	9 km	daily; April–September 2016	new in this study

Deleted: Table 2

<a href="#">OCO-2</a>	<a href="#">SIF</a>	<a href="#">observation derived</a>	<a href="#">0.05°×0.05°</a>	<a href="#">approximately biweekly; April–September 2016</a>	
<a href="#">NASA B-200 and C-130 aircraft</a>	<a href="#">OCS</a>	<a href="#">flask observation</a>	<a href="#">variable</a>	<a href="#">variable; 16–28 August 2016</a>	
<a href="#">CASTNET</a>	<a href="#">O<sub>3</sub> dry deposition velocity <math>V_{d(ozone)}</math></a>	<a href="#">modeled</a>	<a href="#">at the SUM156 and PED108 sites</a>	<a href="#">hourly; 16–28 August 2016</a>	
	<a href="#">O<sub>3</sub> flux <math>F_{(ozone)}</math></a>	<a href="#">modeled multiplied by observed</a>			
<a href="#">European Space Agency PROBA-V, via the Copernicus Global Land Service</a>	<a href="#">GVF</a>	<a href="#">satellite retrieval</a>	<a href="#">1 km</a>	<a href="#">ten-day average; August 2015–2019</a>	<a href="#">used as a model input in part 1</a>
<a href="#">Land and water surface reports operationally collected by the National Centers for Environmental Prediction; and NASA B-200 aircraft</a>	<a href="#">air temperature and humidity</a>	<a href="#">in situ observation</a>	<a href="#">variable</a>	<a href="#">variable; 16–28 August 2016</a>	<a href="#">also used as evaluation datasets in part 1</a>
<a href="#">AQS and CASTNET</a>	<a href="#">surface O<sub>3</sub> concentration</a>	<a href="#">in situ observation</a>	<a href="#">variable</a>	<a href="#">hourly; April–September 2016</a>	
<a href="#">FLUXCOM</a>	<a href="#">latent and sensible heat</a>	<a href="#">observation derived</a>	<a href="#">0.5°×0.5°</a>	<a href="#">daily; April–September 2016</a>	

Acronyms: AQS: Air Quality System; CASTNET: Clean Air Status and Trends Network; GPP: gross primary productivity; GVF: green vegetation fraction; L4C: level 4 carbon; OCO-2: Orbiting Carbon Observatory-2; OCS: carbonyl sulfide; PROBA-V: Project for On-Board Autonomy-Vegetation; SIF: solar-induced chlorophyll fluorescence; SMAP: Soil Moisture Active Passive; VOD: vegetation optical depth.

**Table 3:** Dose-response functions used to estimate the LULC- and crop-specific Relative Yield Losses (i.e., 1 - Relative Yield, RY) due to O<sub>3</sub> exposure and uptake, along with their references.

LULC type	Crop type	Dose-response function (references)	
		Based on Phytotoxic Ozone Dose above the critical level $y$ $\text{nmol m}^{-2} \text{s}^{-1}$ ( $\text{POD}_y$ , in $\text{mmol m}^{-2}$ )	Based on AOT40 in $\text{ppmh}$
Deciduous Forest	/	$\text{RY} = -0.0154 \text{POD}_1 + 1.012$ (CLRTAP, 2017)	/
Grasslands	/	$\text{RY} = -0.0074 \text{POD}_1 + 0.982$ (CLRTAP, 2017)	/
Croplands	Maize	/	$\text{RY} = -0.0036 \text{AOT40} + 1.02$ (Mills et al., 2007)
	Soybean	/	$\text{RY} = -0.0116 \text{AOT40} + 1.02$ (Mills et al., 2007)
	Wheat	$\text{RY} = -0.0064 \text{POD}_3 + 0.9756$ (Mills et al., 2018a; CLRTAP, 2017)	$\text{RY} = -0.0161 \text{AOT40} + 0.99$ (Mills et al., 2007) $\text{RY} = -0.009 \text{AOT40} + 0.969$ (Mills et al., 2018a)

Deleted: CLTRAP

Deleted: CLTRAP

Deleted: CLTRAP



**Table 4:** Evaluation of daily-averaged WRF-Chem gross primary productivity and evaporative fraction, referring to the SMAP L4C and FLUXCOM datasets.

Flux variable (unit)	LULC type	Reference datasets (observation- derived)	Model case					
			Noah_D		CLM_D		P1_W	
			No DA	DA	No DA	DA	No DA	DA
Gross primary productivity (g m <sup>-2</sup> d <sup>-1</sup> )	forests	7.39	7.88	7.08	9.06	6.94	/	
	shrub/grass	5.11	3.28	3.29	4.74	3.89		
	croplands	8.94	7.64	7.40	9.77	8.13		
Evaporative fraction (unitless)	forests	0.75	0.65	0.60	0.67	0.60	0.66	0.63
	shrub/grass	0.67	0.53	0.58	0.57	0.61	0.48	0.48 <sup>a</sup>
	croplands	0.79	0.67	0.67 <sup>a</sup>	0.71	0.68	0.63	0.62

<sup>a</sup>The increases from no-DA cases, which led to improved model performance, are <0.005.

Deleted: 1

Deleted: 3

Deleted: /weather variables

Deleted: Observation(-

**Table 5:** The 24 h and daytime mean O<sub>3</sub> deposition velocity ( $v_{d[O_3]}$ ) and flux ( $F_{[O_3]}$ ) for three LULC groups.

LULC type/Model case	Noah D		CLM D		Noah W		P1 W	
	No DA	DA	No DA	DA	No DA	DA	No DA	DA
24 h mean $v_{d[O_3]}$ (cm s <sup>-1</sup> )								
Forests	0.64	0.56	0.68	0.51	0.54	0.53	0.49	0.48
Shrub/Grass	0.48	0.45	0.53	0.45	0.47	0.48	0.46	0.46
Croplands	0.62	0.54	0.67	0.54	0.58	0.58	0.56	0.56
24 h mean $F_{[O_3]}$ (nmol m <sup>-2</sup> s <sup>-1</sup> )								
Forests	7.11	6.38	7.47	6.35	6.31	6.24	5.75	5.68
Shrub/Grass	4.79	4.48	5.21	4.54	4.76	4.79	4.62	4.63
Croplands	6.90	6.11	7.39	6.06	6.69	6.64	6.44	6.42
Daytime mean $v_{d[O_3]}$ (cm s <sup>-1</sup> )								
Forests	0.94	0.80	1.02	0.71	0.79	0.77	0.70	0.69

Deleted: 4

Deleted: V<sub>d</sub>)

Deleted: )

Deleted: V<sub>d</sub>

Deleted: V<sub>d</sub>

Shrub/Grass	0.63	0.56	0.72	0.58	0.61	0.63	0.58	0.58
Croplands	0.88	0.74	0.99	0.73	0.83	0.83	0.80	0.79
Daytime mean $F_{[ozone]}$ (nmol m <sup>-2</sup> s <sup>-1</sup> )								
Forests	11.51	10.04	12.25	8.99	10.05	9.93	9.04	8.88
Shrub/Grass	6.91	6.32	7.77	6.43	6.83	6.99	6.52	6.49
Croplands	10.99	9.42	12.04	9.31	10.61	10.57	10.17	10.07

**Table 6: Period-mean (16–28 August 2016) soil moisture and surface fluxes at two CASTNET sites shown in Figure 1d. Standard deviations calculated based on the hourly O<sub>3</sub> dry deposition velocity  $V_{d[O_3]}$  and flux  $F_{[O_3]}$  results are also included. Daytime is defined as approximately 8:00–19:00 local standard time.**

CASTNET sites (soil type; LULC type; elevation/terrain)	SUM156, Florida (sand; forest; 16 m/flat)		PEDI08, Virginia (loam; forest; 149 m/rolling)	
	No DA	DA	No DA	DA
Modeled soil moisture initial condition, column-averaged (m <sup>3</sup> m <sup>-3</sup> )				
Noah D	0.15	0.12	0.22	0.20
CLM D	0.16	0.12	0.20	0.18
SMAP L4C daily gross primary productivity (g m <sup>-2</sup> d <sup>-1</sup> )	7.30		8.10	
Modeled daily gross primary productivity (g m <sup>-2</sup> d <sup>-1</sup> )	No DA	DA	No DA	DA
Noah D	4.70	3.83	7.42	5.45
CLM D	5.84	5.88	10.10	4.51
CASTNET (MLM-calculated) daytime $V_{d[O_3]}$ (cm s <sup>-1</sup> )	0.39 ± 0.15		0.39 ± 0.18	
Modeled daytime $V_{d[O_3]}$ (cm s <sup>-1</sup> )	No DA	DA	No DA	DA
Noah D	0.68 ± 0.13	0.64 ± 0.11	0.84 ± 0.23	0.65 ± 0.14
CLM D	0.73 ± 0.13	0.74 ± 0.14	1.01 ± 0.29	0.50 ± 0.09
Noah W	0.63 ± 0.11	0.61 ± 0.10	0.78 ± 0.22	0.75 ± 0.22
CASTNET daytime $F_{[O_3]}$ (nmol m <sup>-2</sup> s <sup>-1</sup> )	3.81 ± 2.02		5.02 ± 2.83	
Modeled daytime $F_{[O_3]}$ (nmol m <sup>-2</sup> s <sup>-1</sup> )	No DA	DA	No DA	DA
Noah D	7.23 ± 1.71	6.91 ± 1.56	12.21 ± 3.88	9.67 ± 2.45
CLM D	7.60 ± 1.74	7.63 ± 1.84	14.27 ± 5.01	7.67 ± 1.88
Noah W	6.81 ± 1.56	6.64 ± 1.45	11.74 ± 3.74	11.18 ± 3.56

Deleted: 1

Deleted: 5: Soil

Deleted: V<sub>d</sub>

Deleted: V<sub>d</sub>

Deleted: 70

Deleted: 66

Deleted: 89

Deleted: 68

Deleted: 74

Deleted: 75

Deleted: 07

Deleted: 52

Deleted: 62

Deleted: 82

Deleted: 79

Deleted: 8.67

Deleted: 11.66

Deleted: 17.95

Deleted: 17.15

Deleted: 30.59

Deleted: 24.27

Deleted: 18.85

Deleted: 18.93

Deleted: 35.63

Deleted: 19.24

Deleted: 16.90

Deleted: 16.48

Deleted: 29.44

Deleted: 28.09

415 Figures

Deleted:




Figure 1: (a) Grid-dominant land use/land cover types grouped from the original 20-category model input (Figure 1c in Huang et al., 2021) based on the method in Table S1; (b) grid-dominant crop type over cropland-dominant regions; (c) gridded population density in 2015; (d) highlighted grid-dominant soil types of sand/loamy sand, loam, and clay which are most relevant to discussions in this paper. The original soil type input from the State Soil Geographic database is shown in Figure S1 in Huang et al. (2021). Locations of the two CASTNET sites for the case studies are denoted in green.

420




Figure 2: Period-mean (16–28 August 2016) WRF-Chem (a, b) column-averaged and (c, d) surface-layer soil moisture fields at initial times and (e–h) their relative changes in % due to the SMAP DA. Results based on the Noah\_D and CLM\_D cases are shown in (a, c, e, g) and (b, d, f, h), respectively.

425

Deleted:

Deleted: -

Deleted: -

Deleted: -

Deleted: ;


Deleted: ;

Deleted: ;


Deleted: ;

Deleted: ;

Deleted: ;




445 Figure 3: Period-mean (16–28 August 2016) green vegetation fraction (a) derived from the Copernicus Global Land Service product and the SMAP morning-time (AM) vegetation optical depth (VOD) using the method described in Figure S2; (b, c, e, f) based on WRF-Chem calculations as well as their responses to the SMAP DA. Results from the Noah\_D and CLM\_D cases are shown in (b, c) and (e, f), respectively. Period-mean SMAP AM VOD is shown in (d).




450 Figure 4: Period-mean (16–28 August 2016) WRF-Chem calculated (b–e) gross primary productivity (GPP); and (g–j) evaporative fraction as well as their responses to the SMAP DA. Results based on the Noah\_D and CLM\_D cases are shown in (b, d, g, i) and (c, e, h, j), respectively. Period-mean SMAP L4C GPP and FLUXCOM evaporative fraction are shown in (a) and (f), respectively, which are also used to evaluate the model results (Table 4).

Deleted: ;  
 Deleted: ;  
 Deleted: ;  
 Deleted: -  
 Deleted: -




Deleted: -  
 Deleted: -  
 Deleted: ;  
 Deleted: ;d;  
 Deleted: ;  
 Deleted: ;e;  
 Deleted: ;  
 Deleted: 3



470 Figure 5: Period-mean (16–28 August 2016) WRF-Chem calculated daytime (a–c) surface air temperature and (g–i) surface radiation as well as (d–f, j–l) their responses to the SMAP DA. Results based on the Noah\_D, CLM\_D, and P1\_W cases are shown in (a, d, g, j), (b, e, h, k), and (c, f, i, l), respectively. Overall, the weather fields from Noah\_D and Noah\_W (not shown in figures) cases are nearly the same. Grey lines in (a) indicate the B-200 flight paths over the southeastern US during the 2016 ACT-America campaign.

1475


- Deleted: -
- Deleted: -
- Deleted: -
- Deleted: ;
- Deleted: -
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;
- Deleted: ;



1490

1495



Figure 6: Evaluation of (a) air temperature and (b) water vapor mixing ratios from several WRF-Chem simulations with the B-200 aircraft observations during the 2016 ACT-America campaign. The RMSEs are summarized in barplots based on model comparisons against observations at all altitudes and near the surface (*i.e.*,  $\geq 800$  hPa). Colored texts above the barplots indicate the SMAP DA impacts on RMSEs. The B-200 flight paths are indicated in Figure 5a.











Deleted:

Deleted: -

Deleted: c-d

Deleted: ;

Deleted: ;

Deleted: (e-f

Deleted: and

Deleted: Results

570 Figure 10: Period-mean (16–28 August 2016) diurnal cycles of (a, b)  $O_3$  dry deposition velocity  $v_d$  and (c, d)  $O_3$  dry  
 575 deposition flux  $F_d$  based on the CASTNET dataset (in black solid lines) and their WRF-Chem counterparts (in purple,  
 blue and brown lines) at the (a, c, e, g) SUM156 and (b, d, f, h) PED108 sites whose locations are shown in Figure 1d.  
 (c, d) and (g, h) indicate the diurnal variability of WRF-Chem stomatal-mesophyll conductance  $g_m$  and column-  
 averaged soil moisture (normalized) at these two sites, respectively. The grey vertical lines in (g, h) denote the initial  
 times of WRF-Chem. WRF-Chem results from the no-DA and DA cases are indicated in solid and dashed lines,  
 respectively. Additional time series plots indicating the daily variability of these fluxes are shown in Figure S5.

585





Figure 11: Period-mean (16–28 August 2016) **WRF-Chem** (a–d) surface MDA8 O<sub>3</sub> fields and (e–h) their responses to the SMAP DA. Results based on the Noah\_D, CLM\_D, Noah\_W, and P1\_W cases are shown in (a, e), (b, f), (c, g) and (d, h), respectively, and the differences between the Noah-MP related cases and the P1\_W case are shown in (i–k).



- Deleted: (e) observed and modeled
- Deleted: -
- Deleted: f-i
- Deleted: ;
- Deleted: b;
- Deleted: ), (c,h
- Deleted: ;i
- Deleted: j-l). Mean MDA8 RMSEs from various cases referring to surface observations are summarized in (m).

590






Figure 12: (a) Period-mean (16–28 August 2016) observed surface MDA8 O<sub>3</sub> and (c) AOT40 in cropland-dominant model grids derived from surface observations during 16–28 August 2016. The RMSEs of modeled MDA8 and model-derived AOT40 from various WRF-Chem cases referring to (a) and (b) are summarized in (c) and (d), respectively.




Deleted: -  
 Deleted: -  
 Deleted: ;  
 Deleted: -  
 Deleted: ;



Deleted:  
 Deleted: 13  
 Deleted: -  
 Deleted: non-urban  
 Deleted: -  
 Deleted: ;  
 Deleted: ;  
 Deleted: ;  
 Deleted: ;

610 **Figure 13:** Box-and-whisker plots of WRF-Chem (a) MDA8  $O_3$ ; (b) daytime stomatal  $O_3$  uptake  $F_{s[Ozone]}$ ; (c) derived  $POD_y$ ; and (d) derived AOT40, summarized by LULC and crop types from all DA-enabled cases. The impacts of the SMAP DA on these model fields are shown in (e-h). Red filled circles indicate the mean values. The mean relative biomass/crop yield losses estimated based on all DA-enabled cases, as well as the SMAP DA impacts on these values, are included in (c, d, g, h) in blue text. The crop yield losses for wheat, estimated based on the derived AOT40 and two dose-response functions (M07: Mills et al., 2007; M18: Mills et al., 2018a) are included in (d, h).



615 **Figure 14:** Period-mean (16–28 August 2016) WRF-Chem (a-d) daytime stomatal  $O_3$  uptake  $F_{s[Ozone]}$  fields over terrestrial regions that do not belong to the urban category in Figure 1a and (e-h) their responses in Figure 1a. Results based on the (a, e) Noah\_D, (b, f) CLM\_D, (c, g) Noah\_W, and (d, h) P1\_W cases are shown.





Figure 15: WRF-Chem based AOT40, derived from the modeled surface O<sub>3</sub> fields during 16–28 August 2016, as well as their responses to the SMAP DA. (a, e), (b, f), (c, g), and (d, h) show results derived from the Noah\_D, CLM\_D, Noah\_W, and P1\_W cases, respectively, in cropland-dominant model grids.



Deleted: 14:

Deleted: in cropland-dominant model grids

Deleted: (e) indicates the observation-derived, and those based on WRF-Chem results

Deleted: are shown in (a:f), (b:g), (c:h), and (d:i).