

# Air-sea exchange of acetone, acetaldehyde, DMS and isoprene at a UK coastal site.

Daniel P. Phillips<sup>1,2</sup>, Frances E. Hopkins<sup>1</sup>, Thomas G. Bell<sup>1</sup>, Peter S. Liss<sup>2</sup>, Philip D. Nightingale<sup>1,2,3</sup>, Claire E. Reeves<sup>2</sup>, Charel Wohl<sup>1,2</sup>, Mingxi Yang<sup>1</sup>

5 <sup>1</sup>Plymouth Marine Laboratory, Plymouth PL1 3DH, UK

<sup>2</sup>Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK

<sup>3</sup>Sustainable Agriculture Systems, Rothamsted Research, Devon EX20 2SB, UK

*Correspondence to: Daniel Phillips (dph@pml.ac.uk) and Mingxi Yang (miya@pml.ac.uk)*

## 10 Abstract

Volatile organic compounds (VOCs) are ubiquitous in the atmosphere and are important for atmospheric chemistry. Large uncertainties remain in the role of the ocean in the atmospheric VOC budget because of poorly constrained marine sources and sinks. There are very few direct measurements of air-sea VOC fluxes near the coast, where natural marine emissions could influence coastal air quality (i.e. ozone, aerosols) and terrestrial gaseous emissions could be taken up by the coastal seas.

15 To address this, we present air-sea flux measurements of acetone, acetaldehyde and dimethylsulfide (DMS) at the coastal Penlee Point Atmospheric Observatory (PPAO) in the ~~S~~~~s~~outh-~~W~~~~w~~est UK during the spring (Apr-May 2018). Fluxes of these gases ~~are quantified~~~~were measured~~ simultaneously by eddy covariance (EC) using a proton transfer reaction quadrupole mass spectrometer. Comparisons are made between two wind sectors representative of different air-water exchange regimes: the ~~open\_~~~~water~~~~open~~~~water~~ sector facing the North Atlantic Ocean and the ~~fetch~~~~limited~~~~terrestrially~~~~influenced~~ Plymouth Sound  
20 fed by two estuaries.

Mean EC ( $\pm 1$  standard error) fluxes of acetone, acetaldehyde and DMS from the ~~open\_~~~~water~~~~open~~~~water~~ wind sector were  $-8.0 \pm 0.8$ ,  $-1.6 \pm 1.4$  and  $4.7 \pm 0.6 \mu\text{mol m}^{-2} \text{d}^{-1}$  respectively (- sign indicates net air-to-sea deposition). These measurements are generally comparable (same order of magnitude) to previous measurements in the Eastern North Atlantic Ocean at the same latitude. In comparison, the ~~terrestrially~~~~influenced~~ Plymouth Sound wind sector showed respective fluxes of  $-12.9 \pm 1.4$ ,  
25  $-4.5 \pm 1.7$  and  $1.8 \pm 0.8 \mu\text{mol m}^{-2} \text{d}^{-1}$ . The greater deposition fluxes of acetone and acetaldehyde within the Plymouth Sound were likely to a large degree driven by higher atmospheric concentrations from the terrestrial wind sector. The reduced DMS emission from the Plymouth Sound was caused by a combination of lower wind speed and likely lower dissolved concentrations as a result of the ~~freshwater~~ estuarine influence (i.e. dilution).

In addition, we measured the near surface seawater concentrations of acetone, acetaldehyde, DMS and isoprene from a  
30 marine station 6 km offshore. Comparisons are made between EC fluxes from the ~~open\_~~~~water~~~~open~~~~water~~ and bulk air-sea VOC fluxes calculated using air/water concentrations with a two-layer (TL) model of gas transfer. The calculated TL fluxes ~~are largely consistent~~~~agree~~ with the EC measurements with respect to the directions and magnitudes of fluxes, ~~implying-~~ ~~that~~ ~~any~~ ~~recently~~ ~~proposed~~ ~~surface~~ ~~emissions~~ ~~of~~ ~~acetone~~ and ~~acetaldehyde~~ ~~would~~ ~~be~~ ~~within~~ ~~the~~ ~~propagated~~ ~~uncertainty~~ ~~of~~ ~~2.6~~  ~~$\mu\text{mol m}^{-2} \text{d}^{-1}$~~ . ~~Accordingly,~~ ~~†~~ The computed transfer velocities of DMS, ~~-and~~ acetone, and acetaldehyde from the EC fluxes and  
35 air/water concentrations are largely consistent with previous transfer velocity estimates from the open ocean. ~~This suggests~~ ~~that~~ ~~wind,~~ ~~rather~~ ~~than~~ ~~bottom-driven~~ ~~turbulence~~ ~~and~~ ~~current~~ ~~velocity,~~ ~~is~~ ~~the~~ ~~main~~ ~~driver~~ ~~for~~ ~~gas~~ ~~exchange~~ ~~within~~ ~~the~~ ~~open~~ ~~water~~ ~~sector~~ ~~at~~ ~~PPAO~~ ~~(depth~~ ~~of~~  ~~$\sim$~~  ~~20~~ ~~m)~~.

## 1 Introduction

40 Volatile organic compounds (VOCs) are ubiquitous in the atmosphere and play an important role in atmospheric chemistry and carbon cycling in the biosphere (Heald et al., 2008). Many VOCs can influence the oxidative capacity of the atmosphere by acting as a source or sink of atmospheric ozone (O<sub>3</sub>) and hydroxyl radicals (Atkinson, 2000; Lewis et al., 2005), ~~and subsequently could indirectly thereby~~ influencing local air quality. The lower-volatility oxidation products, produced from reactions of some VOCs with atmospheric oxidants, can condense into particulates and form cloud condensation nuclei (CCN) (Charlson et al., 1987; Blando and Turpin, 2000; Henze and Seinfeld, 2006), affecting the Earth's radiative forcing and climate.

45 The terrestrial environment is the largest source of VOCs to the atmosphere with the ~~flux~~-emission dominated by isoprene from plant foliage (Guenther et al., 2006). Terrestrial biological processes also produce carbonyls including ketones and aldehydes in varying amounts (Kesselmeier and Staudt, 1999). Large uncertainties exist however with respect to the role that the ocean plays as a net source or sink of these gases (Broadgate et al., 1997; Arnold et al., 2009; Millet et al., 2010; Fischer et al., 2012; Wang et al., 2019). This is due to poorly quantified air-sea fluxes as well as uncertainties in the biogeochemical and physical processes that control them.

50 Acetone and acetaldehyde are carbonyl-based VOCs that have been shown to make up to ~57 % of the carbon mass of all non-methane organic carbon compounds in remote marine air over the North Atlantic Ocean (Lewis et al., 2005). These VOCs have been detected in the surface ocean at concentrations of up to tens of nM (Zhou and Mopper, 1997; Williams et al., 2004), with ~~known~~ oceanic sources including the photochemical degradation of dissolved organic matter (DOM) in bulk seawater (Kieber et al., 1990; Zhou and Mopper, 1997; Zhu and Kieber, 2018) and possibly autotrophic/heterotrophic biological processes (Halsey et al., 2017; Schlundt et al., 2017). The heterogeneous oxidation of DOM at the sea surface has ~~also recently~~ been identified as a source of carbonyl-containing VOCs (Zhou et al., 2014; Ciuraru et al., 2015), however the significance of this is currently unknown.

60 Dimethylsulfide (DMS) is a biogenic sulfur-containing VOC that constitutes the majority of the organic sulfur in the atmosphere (Andreae et al., 1985). It is produced in the surface ocean from the degradation of the algal osmolyte dimethylsulfoniopropionate (Kiene et al., 2000) and subsequently emitted in to the atmosphere. DMS ~~remains is often~~ the dominant source of sulfur in the marine atmosphere (Yang et al., 2011b), and the oxidation products of DMS can act as CCN (Charlson et al., 1987; Veres et al., 2020).

65 Isoprene is an unsaturated terpene-based VOC that is produced in the ocean by a broad range of phytoplankton as a secondary metabolic product (Moore et al., 1994; Shaw et al., 2003; Exton et al., 2013; Booge et al., 2016; Dani and Loreto, 2017). Oceanic sources of isoprene are important for the remote marine atmosphere (Lewis et al., 2001; Arnold et al., 2009; Booge et al., 2016) where transport from terrestrial sources is negligible due to ~~the isoprene's~~ very short atmospheric lifetime (~30 min; Carslaw et al., 2000).

70 The main oceanic sink of most VOCs is biological metabolism at varying rates (Dixon et al., 2014; Royer et al., 2016; Halsey et al., 2017). Surface seawater concentrations of acetone and acetaldehyde tend not to be very sensitive towards their air-sea fluxes (Beale et al., 2015). Emission to the atmosphere is generally considered to be a small loss for seawater DMS (Yang et al., 2013b) but a much larger loss ~~relatively~~ for seawater isoprene (Booge et al., 2018).

75 There have been very few direct measurements of air-sea VOC fluxes with the eddy covariance (EC) technique (e.g. Marandino et al., 2005; Yang et al., 2013c, 2014b; Kim et al., 2017). More often, the fluxes are estimated with the bulk method using air/sea concentrations in a two-layer (TL) model (e.g. Baker et al., 2000; Beale et al., 2013; Wohl et al., 2020). Most studies focus on ~~the~~ open ocean air-sea exchange, while air-sea VOC flux measurements from the coast are essentially non-existent (EC or bulk). Compared to the open ocean, the coastal waters tend to be very ~~biogeochemically~~ dynamic ~~biogeochemically~~ (Borges et al., 2005; Bauer et al., 2013) ~~and physically~~. For example, riverine runoffs carry nutrients and organic carbon into the coastal seas, which could stimulate intense biological cycling (Cloern et al., 2014). ~~Furthermore,~~ compared to the remote marine atmosphere, coastal air is affected by terrestrial emissions.

The air–sea gas transfer velocity can be derived by ~~assuming equating that~~ the EC ~~method flux and with the~~ bulk TL ~~method yield identical fluxes~~ two-layer flux (e.g. Blomquist et al., 2010; Bell et al., 2013; Yang et al., 2013a, 2014a). Limited studies have been undertaken to identify whether open-ocean derived parameterisations of the transfer velocity are applicable to coastal systems (Borges et al., 2004; Yang et al., 2019a). Turbulent processes (i.e. wave-breaking, tidal currents, bottom driven  
85 turbulence) are expected to behave differently in these environments (Upstill-Goddard, 2006), ~~with potential impacts on gas exchange.~~

Here we present air–sea fluxes of acetone, acetaldehyde and DMS determined using the EC technique at a coastal observatory in the south–west UK. Using seawater concentrations measured from a marine station 6 km offshore, we further compute the TL fluxes of these VOCs as well as that of isoprene. We compare our coastal flux measurements with previous  
90 observations of open ocean fluxes as well as global model estimates. ~~We further compare the concurrent~~ From the EC and TL fluxes ~~and air/sea concentrations, – and we~~ derive the gas transfer velocities of DMS, ~~and~~ acetone, ~~and acetaldehyde and compare them against previous observations from the open ocean.~~

## 2 Method

### 2.1 Location

The Penlee Point Atmospheric Observatory (PPAO) is a long-term monitoring station and a part of the Western Channel Observatory (WCO; <https://www.westernchannelobservatory.org.uk/penlee/>) on the ~~S~~outh–~~W~~est (SW) coast of the UK (50.318° N, -4.189° E). The suitability of the observatory for direct air–sea exchange measurements (momentum, heat, greenhouse gases, sea spray aerosols,  $O_3$ ) has been discussed in detail before (Yang et al., 2016a, 2016c, 2019a, 2019b; Loades et al., 2020). The PPAO is located on an exposed headland ~~which benefits from~~ that is exposed to two distinct wind sectors  
100 representative of different air–sea exchange regimes. The SW ~~open water~~ open water sector (depth of ~20 m within the flux footprint) faces the western English Channel and North Atlantic Ocean. The ~~N~~orth–~~E~~ast (NE) Plymouth Sound sector (fetch-limited with a depth of ~10 m) is influenced by estuarine output from the rivers Tamar and Plym (Uncles et al., 2015) as well as natural terrestrial and anthropogenic atmospheric emissions. The theoretical extent of the flux footprint at ~~the~~ PPAO has been discussed previously (Yang et al., 2019a; Loades et al., 2020).

### 2.2 Set-up and measurements

The EC system here principally consists of a high sensitivity proton transfer reaction – mass spectrometer (PTR-quadrupole-MS, Ionicon Analytik) and a sonic anemometer (Gill Windmaster Pro). The measurement setup closely followed previous PPAO flux campaigns (Yang et al., 2016a, 2016d), with a few adaptations made to accommodate the PTR-MS requirements. A simple gas flow diagram is shown in ~~Fig. 1~~ Figure 1.

The sonic anemometer was mounted on a mast ~19 m above mean sea level and run at 10 Hz. The air inlet consisted of a downwards-facing 90 ° union, mounted 30 cm below the anemometer, connected via a 10 m tube to a union tee upstream of the air pump (UT1). All PPAO air sampling instruments (including the PTR-MS) sub-sampled ambient air from this union tee. A dry pump (Gast 1023 series) was used to draw ambient air into the observatory. The total flow rate was ~13.5 L min<sup>-1</sup>, as calculated from the continuously monitored flow of the pump (Bronkhurst EL-FLOW series) and the sum of independent flows  
115 of connected instruments. All unions and tubing before UT1 were 9.5 mm internal diameter (ID).

Downstream of UT1 (i.e. closer to the instruments) was a 0.3 m, 3.2 mm ID tube and union tee (UT2), which split the sample air between the PTR-MS and other equipment (see Loades et al. (2020) for simultaneous PPAO ~~ozone~~  $O_3$  fluxes). Downstream of UT2 (i.e. closer to the PTR-MS) was a 0.3 m, 1.6 mm ID tube, which connected through a solenoid valve (Takasago Electric, Inc.) to the 1.2 m, 0.8 mm ID PTR-MS inlet tubing. All tubing, unions, fittings between the mast inlet and

120 PTR-MS inlet were made from perfluoroalkoxy [Swagelok](#), while the solenoid valve was polytetrafluoroethylene and the PTR-MS inlet was polyetheretherketone (heated to 80 °C to limit surface adsorption).

A solenoid valve, controlled from the PTR-MS, was used to automate routine hourly blanking for the VOC measurements. At the beginning of the campaign, sample gas (ambient air) was diverted through a platinum catalyst (heated to 450 °C) to produce VOC-free air (full oxidation to CO<sub>2</sub>; Yang and Fleming, 2019). However, the use of the catalyst caused overheating  
125 within the PPAO building and was replaced with an activated charcoal filter, which unfortunately proved to be inefficient at removing VOCs. The charcoal filter was replaced with compressed clean air (BOC synthetic air) towards the end of the campaign (27/04/18–03/05/18), which yielded the most consistent blanks. ~~Post-campaign experiments show that the acetaldehyde background when measuring dry, CO<sub>2</sub>-free compressed air is lower than measuring moist atmospheric air scrubbed by the catalyst by about 0.33 ppbv (Wohl et al., 2020). This difference is due to the different levels of CO<sub>2</sub> (accounting for ~0.3 ppbv) and to a lesser extent water vapor (accounting for ~0.03 ppbv) between the compressed air and ambient air, qualitatively consistent with previous works (Warneke et al., 2003; Schwarz et al., 2009). Our calculation of atmospheric acetaldehyde mixing ratio accounts for these sensitivities. The sensitivity of the background as a function of sample humidity was determined from post-campaign laboratory experiments (Wohl et al., 2019) and applied to the field data.~~ Overall, the determination of the VOC backgrounds were more uncertain during this campaign than in previous measurements (Yang et al., 2013c). ~~How~~ however, this is not expected to significantly influence the EC fluxes since the air concentrations were detrended during the flux calculation (Sect. 2.4).

The PTR-MS was set to multiple ion detection mode with four VOCs of interest: acetaldehyde, acetone, DMS and isoprene (initially at  $m/z$  45, 59, 63 and 69 respectively). Previous experiments (Schwarz et al., 2009) and our laboratory [results-tests](#) (Wohl et al., 2019) show that substantial isoprene fragmentation occurs at the voltage used in these measurements (690 V, 166  
140 Td). Importantly, the  $m/z$  41 fragment ion was shown to provide a larger and more stable signal than that of the isoprene parent ion ( $m/z$  69). As a result, a week into the campaign the monitoring of isoprene was changed to  $m/z$  41. The calibration (Sect. 2.5) and thus calculation of the isoprene mixing ratio accounted for the fragmentation and the change in ion monitored. The quadrupole mass dwell time was set to 50 ms for H<sub>3</sub>O<sup>+</sup> and 100 ms for each VOC. In total, the full ion cycle was just over 450 ms, which resulted in a sampling frequency of 2.2 Hz. Dwell time for each VOC was a compromise between measurement  
145 frequency and instrument noise. The PTR-MS parameters are listed in Table A31. Data from the sonic anemometer, flow meter and PTR-MS were all recorded on the same computer to avoid desynchronisation.

## 2.4 Eddy covariance flux calculation

EC gas fluxes ( $F$ ) are determined from the correlation between rapid changes (at least a few Hz) in vertical wind velocity ( $w$ ) and the gas mixing ratio ( $x$ );  $F = \rho_a \overline{w'x'}$ , where  $\rho_a$  represents dry air density, the primes represent fluctuations from the mean and the overbar represents temporal averaging. Air density was computed using the PPAO air temperature, pressure and  
150 humidity data (Gill Instruments Metpak Pro). Wind measurements were linearly interpolated from 10 Hz to match VOC measurement frequency.

A double wind rotation (Tanner and Thurtell, 1969; Hyson et al., 1977) was applied to each 10 min segment of wind data ( $u$  and  $v$  = horizontal,  $w$  = vertical), such that  $u$  became the wind speed ( $U$ ) in the mean wind direction and  $v$  and  $w$  each  
155 averaged zero. This was done to minimise the effect of flow distortion caused by the headland.

A lag time of approximately 3.5 s between wind and PTR-MS measurements was calculated from the dimensions of the tubing and flow rate. The [exact latency lag time](#) was [further](#) determined hourly using a lag correlation analysis between  $w$  and  $x$  over a window of +10 s. Acetone had the strongest flux signal of the four VOCs measured and was used to determine the [latency lag time](#).

160 Lag-adjusted PTR-MS data were used to calculate fluxes in 10 min segments, the sampling interval chosen as the best compromise between maintaining sufficient flux signal-to-noise ratio and satisfying the stationarity criteria in this dynamic region (Yang et al., 2016a, 2016c, 2019a, 2019b).

The sampling frequency of the PTR-MS is relatively low (2.2 Hz), and the instrument's response time determined from laboratory tests is just under 0.5 s. The computed fluxes were corrected for high frequency signal loss due to: 1) sampling at 165 2.2 Hz, and thus missing the flux above the Nyquist frequency of 1.1 Hz and 2) attenuation due to the tubing, using a combined wind speed dependent attenuation factor (mean 1.09, max 1.18). This signal attenuation was estimated from the instrument's response time following the approach of Yang et al. (2013a).

The 10 min flux segments that met the quality control criteria (Table A2 for criteria details) were averaged into 1 h or 3 h fluxes, which reduces random noise by a factor of  $\sim\sqrt{6}$  and  $\sim\sqrt{18}$ , respectively. Over the entire duration of the campaign, 61  
170 % of the data were discarded mainly due to in-appropriate wind directions (i.e. from land) or large variability in wind direction (more common at low wind speeds). Diurnal variability was apparent in the [open-water/open water](#) wind sector data (SW winds) with respect to the number of acceptable flux segments. About twice as many flux segments passed quality control in the daylight hours compared to at night, which is likely due to a diurnal sea-breeze effect. ~~Therefore, the mean open water fluxes are weighted towards the daytime by a ratio of 2/3. When EC fluxes were averaged across 5 weeks of measurements,~~  
175 ~~no difference was observed per hour of the day.~~ No diurnal cycle could be seen in the Plymouth Sound wind sector data (NE winds), likely in part due to the limited data size.

Cospectra of acetone and DMS averaged over all quality-controlled periods are shown in [Fig. 2](#) ~~Figure 2~~ for both the [open-water/open water](#) and Plymouth Sound wind sectors. Here the areas between the cospectral curves and zero represent the magnitudes of the fluxes. Theoretical cospectral fits constrained by the actual wind speed and measurement height (Kaimal et al., 1972) suggest that the measured cospectral shapes were reasonable. The Kaimal fits also showed the high frequency flux loss ( $>\sim 1$  Hz) was small, consistent with the estimated high frequency signal loss correction. Acetaldehyde and isoprene are not shown here because the bidirectionality in acetaldehyde flux and the low isoprene flux magnitude (Sect. 3.1) cause very noisy cospectra.

## 2.5 Seawater measurements and two-layer flux calculation

185 Discrete seawater samples were collected from the L4 marine time-series station of the WCO, which is  $\sim 6$  km S-SW of the PPAO headland. The near surface VOC concentrations are presumed to be similar between L4 and the PPAO [open-water/open water](#) flux footprint despite the outflow from the Tamar estuary that hugs the PPAO headland (Uncles et al., 2015); we revisit this assumption in Sect. 3.3. Seawater acetaldehyde, acetone, DMS and isoprene concentrations were determined using a segmented flow coil equilibrator coupled to the PTR-MS before and after the PPAO campaign (following Wohl et al. (2019)).  
190 Additional seawater DMS concentration measurements were made weekly at L4 using gas chromatography (following Hopkins and Archer (2014)) for the Mar–May 2018 period.

Atmospheric VOC mixing ratios were blank-corrected using the synthetic gas cylinder measurements (27/04/18 onwards, see Sect. 2.2), accounting for ~~the~~ humidity dependence in the background ~~for of~~ DMS (~~mean correction 0.22 and 0.12 ppbv, for SW and NE air respectively, estimated from Wohl et al. (2019)) and the CO<sub>2</sub> plus humidity dependences in the background~~  
195 ~~of acetaldehyde (constant correction 0.33 ppbv estimated from data from Wohl et al. (2020)).~~ We have greater confidence in the measured atmospheric VOC mixing ratios after switching to synthetic air for the blank measurement. Thus, the EC fluxes and the two layer (TL) bulk fluxes were compared over this period (27/04/18–03/05/18) only. Gas phase calibrations of the PTR-MS (using a certified gas standard, Apel-Riemer Environmental Inc) before and after the deployment were averaged and applied to both the atmospheric and seawater VOC data.

200 Surface saturation values of DMS and acetaldehyde were calculated using the recommended solubility from the literature (Burkholder et al., 2015; Sander, 2015) as a function of temperature and adjusted for salinity (Johnson, 2010). Recent

calibrations at environmentally relevant concentrations in seawater suggest that acetone is less soluble than previously thought (Wohl et al., 2020); thus the acetone air-over-water solubility recommended by Burkholder et al. (2015) was by divided by 1.4 as recommended by Wohl et al. (2020).

205 The calculation of the bulk TL fluxes requires the use of wind speed dependent gas transfer velocity parametrisations. The measured wind speed at the PPAO was corrected for flow acceleration due to the headland using results from a comparison with a wind sensor on the L4 buoy (Yang et al., 2019a). This scaled the wind speeds from 19 m (height of the PPAO mast) to 4.9 m (height of L4 wind sensor), and also removed the effect of wind acceleration from the sloped PPAO headland. The corrected winds were then scaled to 10 m;  $U_{10} = U_{4.9}(\ln(10/z_0)/\ln(4.9/z_0))$ , where  $z_0$  represents the aerodynamic  
 210 roughness length calculated from  $U$  using the COAREG 3.5 model (Edson et al., 2013). Overall, the mean correction factor was 0.93 (min 0.79 and max 1.39) for the two wind sectors. The relationship between the corrected  $U_{10}$  and EC friction velocity ( $u_* = ((\overline{u'w'})^2 + (\overline{v'w'})^2)^{0.25}$ ) shows reasonable agreement with the COAREG 3.5 model as well as with Mackay and Yeun (1983) (Figure S1), which validate these wind corrections.

Bulk fluxes ( $F$ ) were calculated following Liss and Slater (1974) and Johnson (2010);

$$215 \quad F = -K_a(C_a - HC_w) \quad (1)$$

$$K_a = \left(\frac{1}{k_a} + \frac{H}{k_w}\right)^{-1} \quad (2)$$

where  $K_a$  represents total airside transfer velocity,  $C_a$  and  $C_w$  represent air and waterside concentrations respectively,  $H$  represents air-over-water dimensionless Henry solubility and  $k_a$  and  $k_w$  represent the individual air and waterside transfer velocities, respectively. Note that total waterside transfer velocity ( $K_w$ ), more commonly used for waterside controlled gases  
 220 such as DMS and isoprene, is equal to  $HK_a$ .  $k_a$  was calculated from the COAREG 3.5 model (Edson et al., 2013), which can be approximated as  $k_a \approx -0.32884U_{10}^3 + 27.428U_{10}^2 + 34.936U_{10} + 553.71$ . The VOCs targeted here span only a small range in diffusivity in air, and are thus assumed to have the same  $k_a$  for simplicity (Yang et al., 2014a, 2016b).  $k_w$  for acetone, acetaldehyde and DMS was calculated using the fit from Yang et al. (2011a); this was an average of DMS observations from five cruises:  $k_w \approx (-0.00797U_{10}^3 + 0.208U_{10}^2 + 0.484U_{10})(S_{Cw}/660)^{-0.5}$ , where  $S_{Cw}$  represents waterside Schmidt  
 225 number.  $k_w$  for isoprene was calculated using the fit from Nightingale et al. (2000);  $k_w \approx (0.222U_{10}^2 + 0.333U_{10})(S_{Cw}/600)^{-0.5}$ . At a  $U_{10}$  of 10 m s<sup>-1</sup>, the Nightingale et al. (2000) parametrization would overestimate DMS  $k_w$  by ~37 % because bubble-mediated gas exchange is less important for the moderately soluble DMS (see Bell et al., 2017; Blomquist et al., 2017). The waterside Schmidt number for all gases were computed following Johnson (2010) at the ambient temperature and salinity.

Appendix B provides further information on the chosen gas solubilities, waterside data and calculations.

## 230 3 Results and Discussion

### 3.1 EC fluxes for the ~~open-water~~ open water and Plymouth Sound wind sectors

Flux measurements were made between 05/04/18 and 03/05/18. During this period, winds arrived from the ~~open-water~~ open water wind sector (180–240° N) 30 % of the time and from the Plymouth Sound sector (30–70° N) 8 % of the time. The wind directions defining these two marine flux sectors were re-established (see Appendix C) compared to previous PPAO studies  
 235 after raising the mast height by ~1 m at the beginning of this campaign.

A time series of VOC fluxes (3 h average) is shown in ~~Fig. 3~~ Figure 3. While acetone and DMS had clear unidirectional fluxes (deposition and emission respectively), acetaldehyde showed bidirectional fluxes throughout the campaign. EC fluxes of acetone, acetaldehyde, DMS and isoprene from the two marine wind sectors averaged over the entire campaign are summarised in Table 1, with positive fluxes representing net sea-to-air emission.

240 For the ~~open-water~~ open water sector, acetone and DMS fluxes had relative standard errors <12 % (from 1 h bins), which are confidently represented by the measurements in the mean thanks to the robust signal-to-noise ratio. Acetaldehyde had

a much greater relative standard error (74 %) because the fluxes often changed in direction and were smaller in magnitude (closer to the limits of detection, LOD); acetaldehyde standard error is lowered to 51 % when the 573 10 min flux segments are instead averaged over the campaign. The isoprene flux was not resolvable with the EC method using our PTR-MS due to the low signal-to-noise ratio. The only direct measurement of isoprene flux from the sea (Kim et al. (2017) in the North Atlantic Ocean) used a chemical ionisation mass spectrometer that is significantly more sensitive than the PTR-MS here.

Overall, 68, 33, 62 and 35 % of 3 h averages of acetone, acetaldehyde, DMS and isoprene, respectively, were above the LOD (DMS and isoprene had the additional condition of a positive flux above LOD) for the open-water sector. The estimation of LOD is shown in Appendix A.

The greater air-to-sea deposition fluxes of acetone and acetaldehyde in the Plymouth Sound sector, compared to the open-water sector, were likely driven by the higher gas mixing ratios in terrestrially-influenced air. Distributions of atmospheric acetone mixing ratio from the two wind sectors are shown in Figure 4. Seawater concentrations of acetone and acetaldehyde within the Plymouth Sound flux footprint were not measured, and thus we are unable to comment on the quantitative effect of riverine outflow on the dissolved VOC concentrations. The weaker sea-to-air flux of DMS in the Plymouth Sound sector compared to the open-water sector was likely due to a combination of low DMS water concentrations associated with freshwater outflow (Uncles et al., 2015) and lower wind speed (Figure 4) that results in reduced waterside-gas transfer velocity ( $k_w$ ; Nightingale et al., 2000; Yang et al., 2011a).

### 3.2 Seawater concentration and atmospheric mixing ratios of VOCs

The seawater concentrations of acetone, acetaldehyde and DMS measured around the campaign (Table A4) are broadly similar ( $\leq 48$  % difference) to previous measurements in the Western English Channel (Archer et al., 2009; Beale et al., 2015), a highly variable and seasonal environment. No seawater measurements of isoprene have been published in this region, so comparisons are made to previous observations from the Eastern North Atlantic ( $\sim 0.02$  nM; Broadgate et al., 2004; Hackenberg et al., 2017), which are  $\sim 79$  % lower than measurements here. The Eastern North Atlantic measurements were conducted in Sep–Oct, thus likely represent a period of lower isoprene productivity. An annual study in the North Sea found the highest isoprene seawater concentrations in May ( $\sim 0.06$  nM; Broadgate et al., 1997), which were closer to our L4 measurements but still lower ( $\sim 41$  %).

The mean atmospheric mixing ratios of acetone and acetaldehyde from the open-water sector ( $0.82 \pm 0.16$  and  $0.51 \pm 0.07$  ppbv respectively, Table 2), mainly observed over one day, are greater than previous measurements around the sample site, possibly in part due to our measurements occurring mainly over one day (01/05/18 1<sup>st</sup> May 2018; see Fig. 6 later for timeseries) thus a multiday average was not possible. Remote marine Atlantic Ocean measurements from the ATom-4 campaign (Apr–May 2018),  $0.31 \pm 0.01$  and  $0.15 \pm 0.04$  ppbv respectively ( $\pm$ SD), from the ATom-4 campaign (Apr–May 2018) within 0.5 km of the sea surface and at latitudes of 40–60° N, observed  $0.31 \pm 0.01$  ppbv acetone ( $\pm$ SD) and  $0.15 \pm 0.04$  ppbv acetaldehyde (Wofsy et al., 2018). Better agreement was seen with shipboard measurements between 40 and 50° N in the Eastern North Atlantic Ocean (Yang et al., 2014b) and coastal measurements at the Mace Head Observatory (Lewis et al., 2005) with mean acetone mixing ratios of  $\sim 0.65$  and  $0.50$  ppbv, respectively, and acetaldehyde mixing ratios of  $\sim 0.16$  and  $0.44$  ppbv, respectively. Finally, rooftop measurements at PML ( $\sim 350$  m north of the Plymouth Sound,  $\sim 6$  km NE of the PPAO, and affected by local ship/port activities) showed daytime-night time atmospheric mixing ratios of  $\sim 0.3$ – $1.14$  and  $\sim 0.1$ – $0.4$  ppbv, respectively for acetone and acetaldehyde (Yang et al., 2013c). A summary of these concentrations, in coordinate form, can be found in Fig. S3. The atmospheric carbonyl measurements at the PPAO from this day might not represent purely oceanic conditions even though wind was from the open-water sector. Hysplit (Stein et al., 2015; Rolph et al., 2017) backward trajectories suggest that the air mass sampled on 1<sup>st</sup> May 2018 this day made contact with land (mainland UK)  $\sim 24$ – $48$  h prior to arriving at the PPAO (Figure S3S4). Lewis et al. (2005) show that similar air-mass trajectories over the UK and continental Europe contained high acetone levels (max of  $\sim 1.67$  ppbv).

Our mean DMS mixing ratio ( $0.20 \pm 0.09$ , Table 2) is in agreement with shipboard measurements in the ~~Eastern North~~ Atlantic Ocean in Jun/Jul ( $0.16 \pm 0.14$  and  $0.12 \pm 0.08$  ppbv (Huebert et al., 2010), with and without a phytoplankton bloom, respectively) and the ~~Western North Atlantic Ocean~~ in March ( $\sim 0.13 \pm 0.04$ ; Quinn et al., 2019). Our mean isoprene mixing ratio ( $0.18 \pm 0.05$ ) is higher than observations in autumn from the ~~western North-West~~ ( $0.035 \pm 0.025$  ppbv; Kim et al., 2017) and ~~eastern North-East~~ ( $0.0044 \pm 0.0083$  ppbv; Hackenberg et al., 2017) ~~North-Atlantic ocean~~. The PPAO isoprene mixing ratio is more similar to measurements at the Mace Head Observatory (Carslaw et al., 2000), which further implies some influence from terrestrial land sources at the PPAO site. We note that the TL air-sea isoprene flux is almost purely governed by its seawater concentration where it is grossly supersaturated; the flux is insensitive to the elevated atmospheric mixing ratio.

### 3.3 Comparison between EC and bulk TL fluxes and derivation of the gas transfer velocities

Here we compare EC fluxes of acetone, acetaldehyde and DMS from the ~~open-water~~ sector with fluxes computed with the TL model (using concurrent atmospheric mixing ratios from the PPAO and linearly interpolated seawater concentrations from L4). This is to assess two assumptions in the TL flux calculation: 1) that gas transfer velocity parameterisations from deeper water (i.e. the open ocean or shelf seas) are applicable to this shallower coastal environment, and 2) seawater VOC concentrations in the PPAO flux footprint and at the L4 station are similar.

Acetone and acetaldehyde were both undersaturated in seawater in the ~~open-water~~ sector (Table 2), while DMS and isoprene were both supersaturated. The mean TL fluxes of acetone, acetaldehyde, DMS and isoprene ( $n=20$ ) are provided in Table 2 with corresponding EC measurements from the ~~open-water~~ wind sector for the last week of the campaign (27/04/18–03/05/18). For this short period with satisfactory VOC blanking, the ~~open-water~~ EC fluxes were fairly large in part because of a 17 % higher mean wind speed compared to the rest of the campaign.

Figure 5 compares the VOC EC and TL fluxes for the ~~open-water~~ sector (not inc. isoprene), and ~~Figure 6~~ shows the same data as a time series. Acetone, acetaldehyde and DMS fluxes exhibit reasonable agreement between the EC and TL methods, with both covarying with wind speed. The data points furthest from the 1:1 line in ~~Figure 5~~ are hourly averages with  $\leq 2$  10 min flux segments, suggesting that the discrepancy is mainly due to random noise in the flux measurement. We do not attempt to calculate TL fluxes for the Plymouth Sound sector because we do not have concurrent seawater measurements and have limited EC measurements for comparison.

To assess the wind speed dependence in gas transfer velocities at the PPAO,  $K_a$  is calculated by equating the EC flux with TL flux:  $-\rho_a \overline{w'x'} = -K_a(C_a - HC_w)$ . Figure 7 shows the calculated  $K_a$  and  $K_w$  ( $K_w = HK_a$ ) values for ~~acetone, acetaldehyde and DMS~~ using 1 h flux data, compared to calculated  $K$  from parameterisations of  $k_a$  and  $k_w$ . In the mean, the expected solubility-dependence in these VOCs is observed. ~~Overall,~~ the measurements show mean  $K_a$  of 2121, 1894 and 183  $\text{cm h}^{-1}$  for acetone, acetaldehyde and DMS (in order of decreasing solubility), respectively. These translate to  $K_w$  values of 2.0, 2.6 and 9.9  $\text{cm h}^{-1}$ .

For DMS, to facilitate comparison with previous measurements, the ~~measured-derived~~  $K_w$  is adjusted to a standard Schmidt number of 660 ( $K_{660} = K_w(660/S_{C_w})^{-0.5}$ ). The mean  $K_{660}$  for DMS ( $16.7 \pm 2.7$   $\text{cm h}^{-1}$ ) is in good agreement when compared to published measurements ( $\sim 17$   $\text{cm h}^{-1}$ ; Blomquist et al., 2006; Yang et al., 2011a; Bell et al., 2013) at similar wind speeds in the North Atlantic Ocean. Data from this short campaign suggest that the ~~mean~~ wind speed dependence in DMS transfer velocity at the PPAO (~~open-water~~ sector, depth of  $\sim 20$  m) is largely comparable to deeper water, in agreement with previous  $\text{CO}_2$  gas exchange measurements at this site (Yang et al., 2019a). ~~Our measured acetone and acetaldehyde  $K_a$  are also of the expected magnitudes.~~ Processes important for shallow estuaries, such as bottom-driven turbulence and tidal current, do not appear to be the main controlling factors in gas exchange in this environment.

~~There is a large amount of scatter in Figure 7, especially for acetone and acetaldehyde. This is likely in part due to the poor temporal resolution (weekly) in seawater concentration measurements that do not capture any rapid changes in this dynamic region (e.g. changes in riverine outflow, biological bloom). In addition, Our DMS exchange-transfer velocity~~



measurements imply that the outflow of the estuary Tamar does not substantially ~~change-dilute~~ the seawater DMS concentration within the ~~open-water~~ flux footprint compared to L4, ~~while~~. There is a large amount of scatter in Figure 7, likely in part due to the poor resolution (weekly) in seawater concentration measurements. ~~Further higher resolution measurements are needed to better constrain the influence of riverine outflow.~~

330 Our measured acetone and acetaldehyde  $K_a$  appear to be very scattered but are of the magnitude as the expected values. Current and previous PPAO observations demonstrate that the air-sea exchange of momentum (Figure S1) and the transfer velocity of DMS (Figure 7) and  $\text{CO}_2$  (Yang et al., 2019a) are generally comparable with open-ocean values. It is possible that the measured seawater carbonyl concentrations at L4 are not representative of the concentrations in the flux footprint of the PPAO. This could be because there are spatial gradients in the carbonyl concentrations ~~could be more significant relative to their air/sea concentration differences,~~ or because there are large temporal changes in seawater concentration due to environmental effects (e.g. changes in riverine outflow, biological bloom) that are missed by the simple linear interpolation of weekly seawater measurements. ~~Further h~~Higher resolution measurements within the flux footprints in the future are needed to better constrain the influence of riverine outflow.

340 As mentioned in Sect. 2.5, this paper uses a revised acetone solubility from Wohl et al. (2020). Focusing on the last week of the campaign (Table 2), we compute a mean TL acetone flux of  $-19.0 \pm 2.2$  and  $-21.5 \pm 2.5 \mu\text{mol m}^{-2} \text{d}^{-1}$  ( $\pm$ SD, propagated error in concentration only) with and without the solubility revision. The flux with revised solubility is in better agreement with the EC measurement ( $-14.9 \pm 2.1 \mu\text{mol m}^{-2} \text{d}^{-1}$ ).

### 3.4 Comparison to other flux estimates

345 The acetone fluxes presented here are in good agreement with previous EC measurements ( $-7.0 \pm 2.2 \mu\text{mol m}^{-2} \text{d}^{-1}$ ) from  $50^\circ$  N on an Atlantic Meridional Transect (AMT) research cruise in Autumn 2012 (Yang et al., 2014b). However, the acetaldehyde flux here shows an opposite mean direction to their  $50^\circ$  N average ( $1.8 \pm 1.3 \mu\text{mol m}^{-2} \text{d}^{-1}$ ). Acetone and acetaldehyde bulk fluxes from  $50^\circ$  N during an AMT cruise in 2009, estimated using CAM-Chem modelled atmospheric concentrations, were  $-5.7 \pm 1.1$  and  $-2.4 \pm 0.7 \mu\text{mol m}^{-2} \text{d}^{-1}$  respectively (Beale et al., 2013). The PPAO fluxes are in reasonable agreement with these earlier indirect estimates. Further acetone and acetaldehyde bulk fluxes estimated from a time series of L4 water measurements and a static atmospheric mixing ratios (0.66 and 0.40 ppbv respectively) were  $-7.4 \pm 1.4$  and  $1.4 \pm 1.0 \mu\text{mol m}^{-2} \text{d}^{-1}$ , respectively, for spring 2011 and  $-6.4 \pm 0.1$  and  $-1.5 \pm 0.1 \mu\text{mol m}^{-2} \text{d}^{-1}$ , respectively, as annual averages. The spring acetone fluxes of Beale et al. (2015) are in good agreement with the PPAO direct measurements, however their acetaldehyde flux was in the opposite direction. Importantly, Beale et al. (2015) identified that the L4 water saturation of acetaldehyde was highly sensitive to the atmospheric mixing ratio (changing it between 0.1–0.4 ppbv was enough to switch the direction of flux). Thus variability and uncertainty in the atmospheric acetaldehyde mixing ratio are probably among the main causes for the discrepancies in these air-sea flux estimates (Yang et al. (2014b) ~~(discussed earlier)~~, Beale et al. (2013, 2015), and PPAO measurements presented here).

360 The PPAO DMS flux is in agreement with previous EC DMS flux observations from the Eastern North Atlantic Ocean during summer 2007 ( $2.6 \pm 0.2$  and  $4.4 \pm 0.4 \mu\text{mol m}^{-2} \text{d}^{-1}$  with and without a short, intense phytoplankton bloom; Huebert et al., 2010). Our campaign ended before the spring phytoplankton bloom started at L4 (Archer et al., 2009), which typically starts in May/June. Yang et al. (2016c) estimated bulk DMS fluxes of 3 and  $10 \mu\text{mol m}^{-2} \text{d}^{-1}$  in the winter and summer, respectively, using the L4 annual seawater DMS concentration time series (Archer et al., 2009). The DMS flux measurement at the PPAO is also in reasonable agreement with previous observations in the North Atlantic Ocean without the phytoplankton bloom ( $5.58 \pm 0.15 \mu\text{mol m}^{-2} \text{d}^{-1}$ ; Bell et al., 2013).

365 Although isoprene EC fluxes were unresolvable, the estimated open water TL flux was similar to NECS April estimates of  $\sim 0.8 \mu\text{mol m}^{-2} \text{d}^{-1}$  (Palmer and Shaw, 2005) and  $\sim 1.0 \mu\text{mol m}^{-2} \text{d}^{-1}$  (Booge et al., 2016, supp.) using MODIS satellite observations of chlorophyll-a for production models. A mean direct flux ( $0.7 \mu\text{mol m}^{-2} \text{d}^{-1}$ ) was observed in the eastern North

Atlantic Ocean (Kim et al., 2017); they estimated seawater concentrations (~0.02 nM) that were ~78 % lower than measured here but also experienced much higher wind speed. Importantly, discrepancies between top-down and bottom-up isoprene budget analyses show the ocean fluxes necessary to explain remote marine boundary layer (MBL) concentrations of isoprene are on the order of a magnitude higher than predicted from seawater concentrations alone (Arnold et al., 2009; Booge et al., 2016). Future simultaneous comparisons between EC and TL isoprene fluxes might observe this difference.

We also compare our direct VOC fluxes to global model predictions at this location. Typical global models and climatologies operate with a 1° latitudinal and longitudinal resolution (e.g. Wang et al., 2019, 2020a, 2020b). At the PPAO this equates to a grid size of ~110 km N-S and ~85 km E-W, which is much coarser compared to the flux footprint areas (on the order of a few km<sup>2</sup>). Nevertheless, this comparison allows us to qualitatively assess the spatial representativeness of the PPAO fluxes and identify potential shortcomings in the models.

The GEOS-Chem and CAM-Chem models predict a mean annual acetone flux in the Eastern North Atlantic Ocean coastal shelves (NECS; biogeochemical province that includes the PPAO coastal site) of ~2.5 μmol m<sup>-2</sup> d<sup>-1</sup> (Fischer et al., 2012) and Mar-May flux of ~2.9 μmol m<sup>-2</sup> d<sup>-1</sup> (Wang et al., 2020a), respectively. These two predictions are much lower than measured in the PPAO open-water sector (-8.01±0.77 μmol m<sup>-2</sup> d<sup>-1</sup>, Table 1). ~~Both of these models estimate lower fluxes because they predict a smaller air-sea concentration difference. Fisher et al. (2012) and Wang et al. (2020a) used water concentrations of 15 (static) and ~8 nM respectively (12482 % and ~19-50 % higher than L4 concentrations) and a modelled atmospheric mixing ratio of ~0.45 ppbv (~45 % lower than the PPAO concentration). Seawater acetone has a highly variable lifetime (Beale et al., 2013; de Bruyn et al., 2013; Dixon et al., 2013) so the higher estimated seawater concentration in the Wang et al. (2020a) model may be because of a slower removal rate. We measured an atmospheric acetone mixing ratio that is higher than in the model likely because our measurement, even from the open-water sector, has some terrestrial influence (e.g. via the diurnal sea-breeze effect or advection of air masses previously in contact with land).~~

The small negative acetaldehyde fluxes measured at the PPAO imply a net-weak ocean sink (mean of -1.55±1.14 μmol m<sup>-2</sup> d<sup>-1</sup>, Table 1), which does not agree with global model results. The NECS acetaldehyde flux is ~0.22 μmol m<sup>-2</sup> d<sup>-1</sup> in GEOS-Chem (Millet et al., 2010) and ~1.4 μmol m<sup>-2</sup> d<sup>-1</sup> in CAM-Chem (Wang et al., 2019) in the Mar-May period. The seawater concentrations used by Wang et al. (2019) (~11 nM) are higher than L4 measurements for the same period (79-54 % higher than L4 conc.). In the model, the short marine lifetime of acetaldehyde (de Bruyn et al., 2013, 2017; Dixon et al., 2013) is coupled with an estimate of its production rate to predict the seawater concentration. Wang et al. (2019) use of a constant lifetime of 0.3 d for the global ocean, whereas previous measurements at L4 suggest a much shorter acetaldehyde lifetime (0.01–0.13 d; Beale et al., 2015). de Bruyn et al. (2017) showed shorter seawater acetaldehyde lifetimes at another coastal location directly after rainfall or during the wet season. It is possible that the shorter lifetime at L4/PPAO, and hence lower seawater acetaldehyde concentration, may be related to riverine input. The atmospheric acetaldehyde mixing ratios from the models are also lower than the coastal observations presented here, similar to acetone as discussed above.

~~The PPAO DMS flux is in agreement with previous EC DMS flux observations from the Eastern North Atlantic Ocean during summer 2007 (2.6±0.2 and 4.4±0.4 μmol m<sup>-2</sup> d<sup>-1</sup> excluding with and without a short, intense phytoplankton bloom; Huebert et al., 2010). Our campaign ended before the spring phytoplankton bloom started at L4 (Archer et al., 2009), which typically starts in May/June. With a short bloom included, Huebert et al. (2010) observed a 4.4±0.4 μmol m<sup>-2</sup> d<sup>-1</sup> flux which is indicative of the lower net DMS productivity of the Eastern North Atlantic Ocean (Lana et al., 2011) compared to the PPAO footprint. Yang et al. (2016c) estimated bulk DMS fluxes of 3 and 10 μmol m<sup>-2</sup> d<sup>-1</sup> in the winter and summer, respectively, using the L4 annual seawater DMS concentration time series (Archer et al., 2009). The DMS flux measurement at Penlee is also in reasonable agreement with previous observations in the North Atlantic (5.58±0.15 μmol m<sup>-2</sup> d<sup>-1</sup> without the phytoplankton bloom (Bell et al., 2013); 1.5 μmol m<sup>-2</sup> d<sup>-1</sup> (Kim et al., 2017)).~~

From the perspective of climatology and global modelling, the NECS was predicted to have an April DMS flux of ~14 μmol m<sup>-2</sup> d<sup>-1</sup> (Lana et al., 2011) and ~6 μmol m<sup>-2</sup> d<sup>-1</sup> (Wang et al., 2020b). The estimate of Lana et al. (2011) is 3 times higher than

the PPAO measurements, however that of Wang et al. (2020b) is in good agreement. Waterside concentrations of ~7 and ~3 nM were predicted (~~69-52~~ % higher and ~~27-43~~ % lower than the L4 concentration) by Lana et al. (2011) and Wang et al. (2020b) respectively. The high concentrations in the Lana et al. (2011) climatology likely correspond to phytoplankton blooms in the Apr–May period, which did not occur at the time of our measurements nor appear as intensely in the estimations of Wang et al. (2020b). This could be partly related to the interannual variability in plankton dynamics and hence seawater DMS cycling.

All the models and climatologies discussed here (except for Wang et al. (2020b)) used the Nightingale et al. (2000)  $k_w$  parameterisation, which leads to ~26, ~~~26-14~~ and ~~~14-26~~ % higher  $k_w$  values for ~~DMS~~, acetone, acetaldehyde and DMS~~acetaldehyde~~, respectively, than that of Yang et al. (2011a) over this campaign because of the solubility dependence in bubble-mediated gas exchange (Yang et al., 2011a; Bell et al., 2017). Substituting the Nightingale et al. (2000)  $k_w$  parameterisation with the Yang et al. (2011a)  $k_w$  would generally reduce the flux magnitude in these models; estimated as ~5, ~6 and ~19 % for acetone, acetaldehyde and DMS, respectively, ~~(using the COAREG 3.5 parameterisation of  $k_a$  unchanged and from COAREG 3.5).~~

~~Although isoprene EC fluxes were unresolvable, the estimated open water open water TL flux was similar to a NECS April estimate of  $-0.8 \mu\text{mol m}^{-2} \text{d}^{-1}$  (Palmer and Shaw, 2005) using MODIS satellite observations of chlorophyll a. A mean direct flux ( $0.7 \mu\text{mol m}^{-2} \text{d}^{-1}$ ) was observed in the Western North Atlantic Ocean (Kim et al., 2017); they estimated seawater concentrations were ~78 % lower than measured here but experienced much higher wind speed.~~

### 3.5 Significance of fluxes

Our measured open water open water acetone and acetaldehyde air–sea fluxes exhibit stronger deposition flux than estimated by global modelling. ~~Here we evaluate; new atmospheric~~ the lifetimes of these gases within the atmospheric marine boundary layer (MBL) ~~MBL are estimated to discuss the significance of these fluxes.~~ Atmospheric ~~The MBL~~ lifetimes ( $\tau_x$ ) are calculated similarly to a 0D model;  $\tau_x = C_x / (C_x L_x + D_x / h)$ , where  $C_x$  represents concentration (here, the mean of the PPAO observations, Table 2),  $L_x$  represents chemical loss,  $D_x$  represents the net deposition flux (here, the mean of the PPAO EC observations, Table 1) and  $h$  represents marine boundary layer (MBL) height. We assume  $h$  to be 500 m and VOCs to be homogeneously mixed within the MBL upwind of the PPAO, and  $L_x$  is  $2.7 \times 10^{-7} \text{ s}^{-1}$  for acetone and  $2.1 \times 10^{-5} \text{ s}^{-1}$  for acetaldehyde.  $L_x$  is calculated from literature OH reaction rates ( $1.7 \times 10^{-13}$  and  $1.5 \times 10^{-11} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$ , respectively; Atkinson and Arey, 2003), April surface  $50^\circ \text{ N}$  zonal average OH concentration ( $\sim 10^6 \text{ molec. cm}^{-3}$ ; Horowitz et al., 2003) and summer surface photolysis rates for acetone ( $\sim 10^{-7} \text{ s}^{-1}$ ; Blitz et al., 2004) and acetaldehyde ( $\sim 5.5 \times 10^{-6} \text{ s}^{-1}$ ; Warneck and Moortgat, 2012).

From this simple model and the values described above, the atmospheric ~~MBL~~ lifetimes of acetone and acetaldehyde are calculated to be 2.0 and 0.5 d respectively. The deposition to the ocean above accounts for 95 and 8 % of all losses for MBL acetone and acetaldehyde respectively, which highlights the importance of acetone air–sea exchange. The calculated PPAO open water open water acetone lifetime is much shorter than the 11–18 d estimates for the mean troposphere (Marandino et al., 2005; Fischer et al., 2012; Khan et al., 2015; Wang et al., 2020a), where surface deposition is less substantial (16–56 %; Jacob et al., 2002; Fischer et al., 2012). The acetaldehyde lifetime is in good agreement with literature values (0.1–0.8 d; Millet et al., 2010; Wang et al., 2019) because photochemistry is the main loss mechanism for MBL acetaldehyde rather than deposition. ~~because and~~ the Apr surface  $50^\circ \text{ N}$  zonal average OH concentration is similar to the tropospheric mean ( $1.1 \times 10^6 \text{ molec. cm}^{-3}$ ; Li et al., 2018) ~~and because photochemistry is the main loss mechanism for MBL acetaldehyde rather than deposition.~~

The noticeable increase in acetone and acetaldehyde deposition fluxes within the Plymouth Sound compared to the open water open water sector suggests that deposition can represent a large loss term for atmospheric VOCs in the coastal atmosphere. Importantly, ~~E~~ enhanced coastal deposition fluxes of acetone and acetaldehyde are not well captured in global models. The atmospheric boundary layer lifetimes during periods of offshore winds are calculated using the same approach as

before, resulting in 1.7 d for acetone and 0.5 d for acetaldehyde. Here deposition to the ocean accounts for 96 and 17 % of losses, respectively. The estimated lifetimes in the Plymouth Sound sector are similar to the open\_ocean even though the deposition fluxes were stronger because the increased deposition fluxes correspond to a greater atmospheric burden. As air masses from the Plymouth Sound (or further inland to the NE) are advected offshore over the coastal seas, the mixing ratios of carbonyls tend to decrease due to combination of deposition to the sea, chemical losses, and mixing with marine air. If additional x-y dimensional constraints are added to the Plymouth Sound OD box (e.g. 4 km N-S, 1 km E-W), dilution becomes the dominant an important loss term.

#### 4 Conclusions

This paper presents measurements of air-sea fluxes of acetone, acetaldehyde, and DMS at the coastal PPAO in the SW UK. A ~~quadrupole~~-PTR-~~quadrupole~~-MS (2.2 Hz) was used to simultaneously resolve the fluxes of these gases with the eddy covariance (EC)-method. Comparisons between the ~~open water~~ sector and a ~~terrestrially influenced estuary dominated~~ sector ~~with land influence~~ show stronger deposition fluxes of acetone and acetaldehyde from the ~~estuary sector~~ latter because of higher atmospheric carbonyl concentrations from that wind direction. Emission fluxes of DMS from the ~~terrestrially influenced estuary~~ sector are weaker than from the ~~open water~~ sector, likely because of lower wind speeds and lower seawater concentrations due to ~~riverine estuarine~~ influence/dilution.

Bulk fluxes computed from seawater and atmospheric concentrations using the ~~two-layer~~ (TL) approach agree reasonably well with our EC measurements of ~~open water~~ DMS, acetone, ~~and~~ acetaldehyde ~~and~~ DMS fluxes. The derived air-sea transfer velocities of ~~acetone, acetaldehyde and DMS, acetone, and acetaldehyde~~ are largely consistent with previous estimates over the open ocean in the mean. Along with previous measurements of air-sea CO<sub>2</sub> exchange, these data suggest that wind, rather than processes such as bottom driven turbulence, is the dominant control for gas exchange at this site for the ~~open water~~ sector (depth of ~20 m).

While the DMS flux at PPAO was in reasonable agreement with recent climatological ~~estimates~~, our measured ~~open water~~ acetone fluxes show stronger deposition compared to climatological and model estimates for the NE Atlantic Ocean coastal shelf. This is likely because the models/climatologies do not fully capture the spatial ~~and~~ temporal variability in the distribution of atmospheric and seawater acetone concentrations. The PPAO acetaldehyde flux (net sink in the mean) is in the opposite direction to global models that suggest that the northern hemisphere oceans ~~is to be~~ a net acetaldehyde source. This is mostly because models predict higher seawater acetaldehyde concentrations and lower atmospheric acetaldehyde mixing ratios at this coastal location.

Future experiments at PPAO should strive to increase the temporal and spatial resolutions of the  $C_w$  observations in order to better match up with the EC fluxes. The accuracies in the air/water concentration measurements can be further improved by using blanks that have the same CO<sub>2</sub> and humidity levels as the samples, especially for acetaldehyde. The use of a higher resolution PTR-time-of-flight-MS at  $\geq 10$  Hz would improve the precision of the flux measurement and reduce the magnitude of the high frequency flux loss. Simultaneous methanol fluxes would enable the derivation of the airside gas transfer velocity ( $k_a$ ) and benefit the lag correlation calculation (methanol flux is typically large; see Yang et al. (2014a)). Important future work would involve measuring EC fluxes,  $C_w$  and  $C_a$  at high resolution from the anthropogenically/terrestrially dominated coast through to the continental shelf edge and open ocean on a ship in order to develop our understanding of the impact of coastal seas on local atmospheric chemistry and quality. Such transect measurements would help to further constrain global models for improved estimations at the coast.

## Appendix A: Additional eddy covariance method details

Table A1 shows the PTR-MS measurement parameters used, which largely follows from Yang et al. (2013c). Importantly, a high drift voltage was used to limit the formation of hydrated clusters ( $H_3O^+ + nH_2O \rightarrow H_3O^+(H_2O)_n$ ) caused by the high humidity of marine air. These hydrated clusters have different reaction kinetics with VOCs and can affect the instrumental sensitivity. Increasing the drift voltage leads to a lower cluster formation (~~unstable high energy collision product~~), but also causes a greater degree of isoprene fragmentation (Schwarz et al., 2009; Wohl et al., 2019). Isoprene fragmentation was corrected using gas phase calibration.

Table A2 shows the quality control criteria used to filter 10 min flux data segments, developed from EC measurements of CO<sub>2</sub> and CH<sub>4</sub> fluxes at the PPAO (Yang et al., 2016a). Standard deviation in wind direction was used to ensure fairly constant wind direction and minimise influence from areas outside the specified NE and SW wind sectors. The remaining controls check for the stationary criteria required for EC measurements, remove periods with obvious flow distortion, or remove periods with substantial rain influence on the sonic anemometer.

Random uncertainty in the EC flux depends on variability (instrument noise + ambient variability) in the VOC measurement and the wind measurement. Precision (or random error) in the EC technique can be approximated as the standard deviation of 'null' fluxes (i.e. covariance data with a lag time between vertical wind and concentration measurements that is much greater than the integral time scale (Spirig et al., 2005)). Fluxes are resolvable when they are higher than 3x the measurement noise. An artificial lag time of 300 s was chosen for this calculation here. The 3x standard deviation was calculated from hourly averages over a 5 h window of stable ~~open-water~~ open water wind speed and direction as 10.1, 6.8, 4.7 and 4.7  $\mu\text{mol m}^{-2} \text{d}^{-1}$ , for acetaldehyde, acetone, DMS and isoprene respectively. Our LOD estimates are 140-250% of the LODs from a previous deployment of this PTR-MS at sea (Yang et al., 2014b), mainly due to the greater ambient variability in VOC mixing ratio at this coastal environment. Averaging hourly data to 3 h reduces random error by a factor of  $\sqrt{3}$ , thus the 3 h LOD are 5.8, 4.0, 2.7 and 2.7  $\mu\text{mol m}^{-2} \text{d}^{-1}$ , for acetaldehyde, acetone, DMS and isoprene respectively.

## Appendix B: Two-layer flux calculation additional information

Table A3 shows the solubility and solubility temperature dependence (Burkholder et al., 2015) and molar volume data (Johnson, 2010) used in the calculations of salinity dependent solubility and Schmidt number. In the diffusion coefficient calculation, the original association factor of solvent (2.6) was used as recommended by Saltzman et al. (Saltzman et al., 1993) for DMS.

During the period with satisfactory hourly blanking (27/04/18–03/05/18), the mean ( $\pm$ SD) L4 water temperature and salinity were  $10.6 \pm 0.8$  °C and  $34.9 \pm 0.1$  PSU (~~n=3~~). These were taken from the surface measurement of a weekly CTD profile collected on the RV *Plymouth Quest*. Mean atmospheric pressure was  $1006 \pm 4$  mB, wind speed at 10 m height was  $9.93 \pm 3.76$  m s<sup>-1</sup>, and friction velocity (also from EC) was  $0.364 \pm 0.166$  m s<sup>-1</sup>. Table A4 shows the available VOC concentration data just before, during and after the campaign. Simultaneous measurements and calculated fluxes were made using concentration data from as close as possible to the period with satisfactory blanking (\*).

## Appendix C: Wind sector determination

Figure S2 shows the double wind rotation tilt angle correction and drag coefficient ( $C_D = (u_* / U)^2$ ) for 10-min segments of wind data averaged into wind direction bins. The 10 min data were filtered ( $U > 3$  m s<sup>-1</sup>) to reduce the influence of the flux footprint overlapping with the narrow strip of land upwind under low wind speed conditions (Yang et al., 2019a). The majority of winds had a positive tilt angle due the rotation of ocean (horizontal) winds hitting the headland. However winds from 260–360° N were negatively angled due to the raised headland behind the observatory. The 30–70° N and 180–240° N zones exhibit

530 a small, positive tilt angle and a low drag coefficient, as might be expected for the open ocean (Kara et al., 2007; Edson et al., 2013). We define these as our two air–water exchange sectors.

### Author contribution

DP, TB and MY designed and set up the experiment at the Penlee Point Atmospheric Observatory. MY provided eddy covariance calculation code and DP undertook the analysis. FH and CW provided the seawater concentrations. DP prepared  
535 the manuscript with contribution from all authors.

### Competing interests

The authors declare that they have no conflict of interest.

### Acknowledgements

This work was supported by an EnvEast NERC DTP grant (NE/L002582/1). The first author would like to thank fellow  
540 authors and staff at PML for assistance in transporting equipment to the observatory. Thanks to Dr Rachael Beale (PML) for her L4 time series and AMT19 data that enabled a quantitative discussion and analysis with our resolved fluxes. [A further thank you to the anonymous referees who gave their time to provide helpful comments that improved the presentation and discussion of the work.](#) Data from the Western Channel Observatory is found at <http://www.westernchannelobservatory.org.uk/>. This is contribution 10 from the Penlee Point Atmospheric Observatory.

### 545 References

- Andreae, M. O., Ferek, R. J., Bermond, F., Byrd, K. P., Engstrom, R. T., Hardin, S., Houmere, P. D., LeMarrec, F., Raemdonck, H. and Chatfield, R. B.: Dimethyl sulfide in the marine atmosphere, *J. Geophys. Res.*, 90(D7), 12891, doi:10.1029/JD090iD07p12891, 1985.
- 550 Archer, S. D., Cummings, D. G., Llewellyn, C. A. and Fishwick, J. R.: Phytoplankton taxa, irradiance and nutrient availability determine the seasonal cycle of DMSP in temperate shelf seas, *Mar. Ecol. Prog. Ser.*, 394, 111–124, doi:10.3354/meps08284, 2009.
- Arnold, S. R., Spracklen, D. V., Williams, J., Yassaa, N., Sciare, J., Bonsang, B., Gros, V., Peeken, I., Lewis, A. C., Alvaïn, S. and Moulin, C.: Evaluation of the global oceanic isoprene source and its impacts on marine organic carbon aerosol, *Atmos. Chem. Phys.*, 9(4), 1253–1262, doi:10.5194/acp-9-1253-2009, 2009.
- 555 Atkinson, R.: Atmospheric chemistry of VOCs and NO<sub>x</sub>, *Atmos. Environ.*, 34(12–14), 2063–2101, doi:10.1016/S1352-2310(99)00460-4, 2000.
- Atkinson, R. and Arey, J.: Atmospheric Degradation of Volatile Organic Compounds, *Chem. Rev.*, 103(12), 4605–4638, doi:10.1021/cr0206420, 2003.
- 560 Baker, A. R., Turner, S. M., Broadgate, W. J., Thompson, A., McFiggans, G. B., Vesperini, O., Nightingal, P. D., Liss, P. S. and Jickells, T. D.: Distribution and sea-air fluxes of biogenic trace gases in the eastern Atlantic Ocean, *Global Biogeochem. Cycles*, 14(3), 871–886, doi:10.1029/1999GB001219, 2000.
- Bauer, J. E., Cai, W. J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S. and Regnier, P. A. G.: The changing carbon cycle of the coastal ocean, *Nature*, 504(7478), 61–70, doi:10.1038/nature12857, 2013.
- 565 Beale, R., Dixon, J. L., Arnold, S. R., Liss, P. S. and Nightingale, P. D.: Methanol, acetaldehyde, and acetone in the surface waters of the Atlantic Ocean, *J. Geophys. Res. Ocean.*, 118(10), 5412–5425, doi:10.1002/jgrc.20322, 2013.
- Beale, R., Dixon, J. L., Smyth, T. J. and Nightingale, P. D.: Annual study of oxygenated volatile organic compounds in UK shelf waters, *Mar. Chem.*, 171, 96–106, doi:10.1016/j.marchem.2015.02.013, 2015.
- 570 Bell, T. G., de Bruyn, W. J., Miller, S. D., Ward, B., Christensen, K. and Saltzman, E. S.: Air-sea dimethylsulfide (DMS) gas transfer in the North Atlantic: Evidence for limited interfacial gas exchange at high wind speed, *Atmos. Chem. Phys.*, 13(21), 11073–11087, doi:10.5194/acp-13-11073-2013, 2013.

- Bell, T. G., Landwehr, S., Miller, S. D., De Bruyn, W. J., Callaghan, A. H., Scanlon, B., Ward, B., Yang, M. and Saltzman, E. S.: Estimation of bubble-mediated air-sea gas exchange from concurrent DMS and CO<sub>2</sub> transfer velocities at intermediate-high wind speeds, *Atmos. Chem. Phys.*, 17(14), 9019–9033, doi:10.5194/acp-17-9019-2017, 2017.
- 575 Blando, J. D. and Turpin, B. J.: Secondary organic aerosol formation in cloud and fog droplets: A literature evaluation of plausibility, *Atmos. Environ.*, 34(10), 1623–1632, doi:10.1016/S1352-2310(99)00392-1, 2000.
- Blitz, M. A., Heard, D. E., Pilling, M. J., Arnold, S. R. and Chipperfield, M. P.: Pressure and temperature-dependent quantum yields for the photodissociation of acetone between 279 and 327.5 nm, *Geophys. Res. Lett.*, 31(6), doi:10.1029/2003gl018793, 2004.
- Blomquist, B. W., Fairall, C. W., Huebert, B. J., Kieber, D. J. and Westby, G. R.: DMS sea-air transfer velocity: Direct measurements by eddy covariance and parameterization based on the NOAA/COARE gas transfer model, *Geophys. Res. Lett.*, 33(7), 2–5, doi:10.1029/2006GL025735, 2006.
- 580 Blomquist, B. W., Huebert, B. J., Fairall, C. W. and Faloon, I. C.: Determining the sea-air flux of dimethylsulfide by eddy correlation using mass spectrometry, *Atmos. Meas. Tech.*, 3(1), 1–20, doi:10.5194/amt-3-1-2010, 2010.
- Blomquist, B. W., Brumer, S. E., Fairall, C. W., Huebert, B. J., Zappa, C. J., Brooks, I. M., Yang, M., Bariteau, L., Prytherch, J., Hare, J. E., Czerski, H., Matei, A. and Pascal, R. W.: Wind Speed and Sea State Dependencies of Air-Sea Gas Transfer: Results From the High Wind Speed Gas Exchange Study (HiWinGS), *J. Geophys. Res. Ocean.*, 122(10), 8034–8062, doi:10.1002/2017JC013181, 2017.
- 585 Booge, D., Marandino, C. A., Schlundt, C., Palmer, P. I., Schlundt, M., Atlas, E. L., Bracher, A., Saltzman, E. S. and Wallace, D. W. R.: Can simple models predict large-scale surface ocean isoprene concentrations?, *Atmos. Chem. Phys.*, 16(18), 11807–11821, doi:10.5194/acp-16-11807-2016, 2016.
- 590 Booge, D., Schlundt, C., Bracher, A., Endres, S., Zäncker, B. and Marandino, C. A.: Marine isoprene production and consumption in the mixed layer of the surface ocean – a field study over two oceanic regions, *Biogeosciences*, 15(2), 649–667, doi:10.5194/bg-15-649-2018, 2018.
- Borges, A. V., Vanderborcht, J. P., Schiettecatte, L. S., Gazeau, F., Ferrón-Smith, S., Delille, B. and Frankignoulle, M.: 595 Variability of the gas transfer velocity of CO<sub>2</sub> in a macrotidal estuary (the Scheldt), *Estuaries*, 27(4), 593–603, doi:10.1007/BF02907647, 2004.
- Borges, A. V., Delille, B. and Frankignoulle, M.: Budgeting sinks and sources of CO<sub>2</sub> in the coastal ocean: Diversity of ecosystem counts, *Geophys. Res. Lett.*, 32(14), 1–4, doi:10.1029/2005GL023053, 2005.
- 600 Broadgate, W. J., Liss, P. S. and Penkett, S. A.: Seasonal emissions of isoprene and other reactive hydrocarbon gases from the ocean, *Geophys. Res. Lett.*, 24(21), 2675–2678, doi:10.1029/97GL02736, 1997.
- Broadgate, W. J., Malin, G., Küpper, F. C., Thompson, A. and Liss, P. S.: Isoprene and other non-methane hydrocarbons from seaweeds: A source of reactive hydrocarbons to the atmosphere, *Mar. Chem.*, 88(1–2), 61–73, doi:10.1016/j.marchem.2004.03.002, 2004.
- 605 de Bruyn, W. J., Clark, C. D., Pagel, L. and Singh, H.: Loss rates of acetone in filtered and unfiltered coastal seawater, *Mar. Chem.*, 150, 39–44, doi:10.1016/j.marchem.2013.01.003, 2013.
- de Bruyn, W. J., Clark, C. D., Senstad, M., Barashy, O. and Hok, S.: The biological degradation of acetaldehyde in coastal seawater, *Mar. Chem.*, 192, 13–21, doi:10.1016/j.marchem.2017.02.008, 2017.
- Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M. and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 18, JPL Publ. [online] Available from: <https://jpldataeval.jpl.nasa.gov/>, 2015.
- 610 Carslaw, N., Bell, N., Lewis, A. C., McQuaid, J. B. and Pilling, M. J.: A detailed case study of isoprene chemistry during the EASE96 Mace Head campaign, *Atmos. Environ.*, 34(18), 2827–2836, doi:10.1016/S1352-2310(00)00088-1, 2000.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O. and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326(6114), 655–661, doi:10.1038/326655a0, 1987.
- 615 Ciuraru, R., Fine, L., Pinxteren, M. Van, D’Anna, B., Herrmann, H. and George, C.: Unravelling New Processes at Interfaces: Photochemical Isoprene Production at the Sea Surface, *Environ. Sci. Technol.*, 49(22), 13199–13205, doi:10.1021/acs.est.5b02388, 2015.
- Cloern, J. E., Foster, S. Q. and Kleckner, A. E.: Phytoplankton primary production in the world’s estuarine-coastal ecosystems, *Biogeosciences*, 11(9), 2477–2501, doi:10.5194/bg-11-2477-2014, 2014.
- 620 Dani, K. G. S. and Loreto, F.: Trade-Off Between Dimethyl Sulfide and Isoprene Emissions from Marine Phytoplankton, *Trends Plant Sci.*, 22(5), 361–372, doi:10.1016/j.tplants.2017.01.006, 2017.
- Dixon, J. L., Beale, R. and Nightingale, P. D.: Production of methanol, acetaldehyde, and acetone in the Atlantic Ocean, *Geophys. Res. Lett.*, 40(17), 4700–4705, doi:10.1002/grl.50922, 2013.

- 625 Dixon, J. L., Beale, R., Sargeant, S. L., Tarran, G. A. and Nightingale, P. D.: Microbial acetone oxidation in coastal seawater, *Front. Microbiol.*, 5(MAY), 1–9, doi:10.3389/fmicb.2014.00243, 2014.
- Edson, J. B., Jampana, V., Weller, R. A., Bigorre, S. P., Plueddemann, A. J., Fairall, C. W., Miller, S. D., Mahrt, L., Vickers, D. and Hersbach, H.: On the exchange of momentum over the open ocean, *J. Phys. Oceanogr.*, 43(8), 1589–1610, doi:10.1175/JPO-D-12-0173.1, 2013.
- 630 Exton, D. A., Suggett, D. J., McGenity, T. J. and Steinke, M.: Chlorophyll-normalized isoprene production in laboratory cultures of marine microalgae and implications for global models, *Limnol. Oceanogr.*, 58(4), 1301–1311, doi:10.4319/lo.2013.58.4.1301, 2013.
- Fischer, E. V., Jacob, D. J., Millet, D. B., Yantosca, R. M. and Mao, J.: The role of the ocean in the global atmospheric budget of acetone, *Geophys. Res. Lett.*, 39(1), 3–7, doi:10.1029/2011GL050086, 2012.
- 635 Guenther, A. B., Karl, T. R., Harley, P., Wiedinmyer, C., Palmer, P. I. and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6(11), 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- 640 Hackenberg, S. C., Andrews, S. J., Ains, R., Arnold, S. R., Bouman, H. A., Brewin, R. J. W., Chance, R. J., Cummings, D., Dall’Olmo, G., Lewis, A. C., Minaeian, J. K., Reifel, K. M., Small, A., Tarran, G. A., Tilstone, G. H. and Carpenter, L. J.: Potential controls of isoprene in the surface ocean, *Global Biogeochem. Cycles*, 31(4), 644–662, doi:10.1002/2016GB005531, 2017.
- Halsey, K. H., Giovannoni, S. J., Graus, M., Zhao, Y., Landry, Z., Thrash, J. C., Vergin, K. L. and De Gouw, J. A.: Biological cycling of volatile organic carbon by phytoplankton and bacterioplankton, *Limnol. Oceanogr.*, 62(6), 2650–2661, doi:10.1002/lno.10596, 2017.
- 645 Heald, C. L., Goldstein, A. H., Allan, J. D., Aiken, A. C., Apel, E. C., Atlas, E. L., Baker, A. K., Bates, T. S., Beyersdorf, A. J., Blake, D. R., Campos, T., Coe, H., Crouse, J. D., DeCarlo, P. F., De Gouw, J. A., Dunlea, E. J., Flocke, F. M., Fried, A., Goldan, P. D., Griffin, R. J., Herndon, S. C., Holloway, J. S., Holzinger, R., Jimenez, J. L., Junkermann, W., Kuster, W. C., Lewis, A. C., Meinardi, S., Millet, D. B., Onasch, T., Polidori, A., Quinn, P. K., Riemer, D. D., Roberts, J. M., Salcedo, D., Sive, B., Swanson, A. L., Talbot, R., Warneke, C., Weber, R. J., Weibring, P., Wennberg, P. O., Worsnop, D. R., Wittig, A. E., Zhang, R., Zheng, J. and Zheng, W.: Total observed organic carbon (TOOC) in the atmosphere: A synthesis of North American observations, *Atmos. Chem. Phys.*, 8(7), 2007–2025, doi:10.5194/acp-8-2007-2008, 2008.
- 650 Henze, D. K. and Seinfeld, J. H.: Global secondary organic aerosol from isoprene oxidation, *Geophys. Res. Lett.*, 33(9), 6–9, doi:10.1029/2006GL025976, 2006.
- Hopkins, F. E. and Archer, S. D.: Consistent increase in dimethyl sulfide (DMS) in response to high CO<sub>2</sub> in five shipboard bioassays from contrasting NW European waters, *Biogeosciences*, 11(18), 4925–4940, doi:10.5194/bg-11-4925-2014, 2014.
- 655 Horowitz, L. W., Walters, S., Mauzerall, D. L., Emmons, L. K., Rasch, P. J., Granier, C., Tie, X., Lamarque, J. F., Schultz, M. G., Tyndall, G. S., Orlando, J. J. and Brasseur, G. P.: A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2, *J. Geophys. Res. Atmos.*, 108(24), doi:10.1029/2002jd002853, 2003.
- 660 Huebert, B. J., Blomquist, B. W., Yang, M., Archer, S. D., Nightingale, P. D., Yelland, M. J., Stephens, J., Pascal, R. W. and Moat, B. I.: Linearity of DMS transfer coefficient with both friction velocity and wind speed in the moderate wind speed range, *Geophys. Res. Lett.*, 37(1), 0–4, doi:10.1029/2009GL041203, 2010.
- Hyson, P., Garratt, J. R. and Francey, R. J.: Algebraic and Electronic Corrections of Measured uw Covariance in the Lower Atmosphere, *J. Appl. Meteorol.*, 16(1), 43–47, doi:10.1175/1520-0450(1977)016<0043:AAECOM>2.0.CO;2, 1977.
- Jacob, D. J., Field, B. D., Jin, E. M., Bey, I., Li, Q., Logan, J. A., Yantosca, R. M. and Singh, H. B.: Atmospheric budget of acetone, *J. Geophys. Res. Atmos.*, 107(D10), doi:10.1029/2001JD000694, 2002.
- 665 Johnson, M. T.: A numerical scheme to calculate temperature and salinity dependent air-water transfer velocities for any gas, *Ocean Sci.*, 6(4), 913–932, doi:10.5194/os-6-913-2010, 2010.
- Kaimal, J., Wyngaard, J., Izumi, Y. and Cote, O.: Spectral characteristics of surface-layer turbulence, *Q. J. R. Meteorol. Soc.*, 098(417), 563–589, doi:10.1256/smsqj.41706, 1972.
- 670 Kara, A. B., Metzger, E. J. and Bourassa, M. A.: Ocean current and wave effects on wind stress drag coefficient over the global ocean, *Geophys. Res. Lett.*, 34(1), 2–5, doi:10.1029/2006GL027849, 2007.
- Kesselmeier, J. and Staudt, M.: Biogenic Volatile Organic Compound (VOC): An Overview on Emissions, Physiology and Ecology, *J. Atmos. Chem.*, 33(1), 23–88, doi:10.1023/A:1006127516791, 1999.
- 675 Khan, M. A. H., Cooke, M. C., Utembe, S. R., Archibald, A. T., Maxwell, P., Morris, W. C., Xiao, P., Derwent, R. G., Jenkin, M. E., Percival, C. J., Walsh, R. C., Young, T. D. S., Simmonds, P. G., Nickless, G., O’Doherty, S. and Shallcross, D. E.: A study of global atmospheric budget and distribution of acetone using global atmospheric model STOCHEM-CRI, *Atmos. Environ.*, 112, 269–277, doi:10.1016/j.atmosenv.2015.04.056, 2015.
- Kieber, R. J., Zhou, X. and Mopper, K.: Formation of carbonyl compounds from UV-induced photodegradation of humic substances in natural waters: Fate of riverine carbon in the sea, *Limnol. Oceanogr.*, 35(7), 1503–1515,




doi:10.4319/lo.1990.35.7.1503, 1990.

- 680 Kiene, R. P., Linn, L. J. and Bruton, J. A.: New and important roles for DMSP in marine microbial communities, *J. Sea Res.*, 43(3–4), 209–224, doi:10.1016/S1385-1101(00)00023-X, 2000.
- Kim, M. J., Novak, G. A., Zoerb, M. C., Yang, M., Blomquist, B. W., Huebert, B. J., Cappa, C. D. and Bertram, T. H.: Air-Sea exchange of biogenic volatile organic compounds and the impact on aerosol particle size distributions, *Geophys. Res. Lett.*, 44(8), 3887–3896, doi:10.1002/2017GL072975, 2017.
- 685 Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E. and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cycles*, 25(1), 1–17, doi:10.1029/2010GB003850, 2011.
- Lewis, A. C., Carpenter, L. J. and Pilling, M. J.: Nonmethane hydrocarbons in Southern Ocean boundary layer air, *J. Geophys. Res. Atmos.*, 106(D5), 4987–4994, doi:10.1029/2000JD900634, 2001.
- 690 Lewis, A. C., Hopkins, J. R., Carpenter, L. J., Stanton, J., Read, K. A. and Pilling, M. J.: Sources and sinks of acetone, methanol, and acetaldehyde in North Atlantic marine air, *Atmos. Chem. Phys.*, 5(7), 1963–1974, doi:10.5194/acp-5-1963-2005, 2005.
- Li, M., Karu, E., Brenninkmeijer, C. A. M., Fischer, H., Lelieveld, J. and Williams, J.: Tropospheric OH and stratospheric OH and Cl concentrations determined from CH<sub>4</sub>, CH<sub>3</sub>Cl, and SF<sub>6</sub> measurements, *Clim. Atmos. Sci.*, 1(1), 29, doi:10.1038/s41612-018-0041-9, 2018.
- 695 Liss, P. S. and Slater, P. G.: Flux of Gases across the Air-Sea Interface, *Nature*, 247(5438), 181–184, doi:10.1038/247181a0, 1974.
- Loades, D. C., Yang, M., Bell, T. G., Vaughan, A. R., Pound, R. J., Metzger, S., Lee, J. D. and Carpenter, L. J.: Ozone deposition to a coastal sea: comparison of eddy covariance observations with reactive air-sea exchange models, *Atmos. Meas. Tech.*, 13(12), 6915–6931, doi:10.5194/amt-13-6915-2020, 2020.
- 700 Mackay, D. and Yeun, A. T. K.: Mass Transfer Coefficient Correlations for Volatilization of Organic Solutes from Water, *Environ. Sci. Technol.*, 17(4), 211–217, doi:10.1021/es00110a006, 1983.
- Marandino, C. A., de Bruyn, W. J., Miller, S. D., Prather, M. J. and Saltzman, E. S.: Oceanic uptake and the global atmospheric acetone budget, *Geophys. Res. Lett.*, 32(15), doi:10.1029/2005GL023285, 2005.
- 705 Millet, D. B., Guenther, A. B., Siegel, D. A., Nelson, N. B., Singh, H. B., De Gouw, J. A., Warneke, C., Williams, J., Eerdekens, G., Sinha, V., Karl, T. R., Flocke, F., Apel, E. C., Riemer, D. D., Palmer, P. I. and Barkley, M.: Global atmospheric budget of acetaldehyde: 3-D model analysis and constraints from in-situ and satellite observations, *Atmos. Chem. Phys.*, 10(7), 3405–3425, doi:10.5194/acp-10-3405-2010, 2010.
- Moore, R. M., Oram, D. E. and Penkett, S. A.: Production of isoprene by marine phytoplankton cultures, *Geophys. Res. Lett.*, 21(23), 2507–2510, doi:10.1029/94GL02363, 1994.
- 710 Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J. and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cycles*, 14(1), 373–387, doi:10.1029/1999GB900091, 2000.
- Palmer, P. I. and Shaw, S. L.: Quantifying global marine isoprene fluxes using MODIS chlorophyll observations, *Geophys. Res. Lett.*, 32(9), 1–5, doi:10.1029/2005GL022592, 2005.
- 715 Quinn, P. K., Bates, T. S., Coffman, D. J., Upchurch, L., Johnson, J. E., Moore, R., Ziemba, L., Bell, T. G., Saltzman, E. S., Graff, J. and Behrenfeld, M. J.: Seasonal Variations in Western North Atlantic Remote Marine Aerosol Properties, *J. Geophys. Res. Atmos.*, 124(24), 14240–14261, doi:10.1029/2019JD031740, 2019.
- Rolph, G., Stein, A. and Stunder, B.: Real-time Environmental Applications and Display sYstem: READY, *Environ. Model. Softw.*, 95, 210–228, doi:10.1016/j.envsoft.2017.06.025, 2017.
- 720 Royer, S. J., Galí, M., Mahajan, A. S., Ross, O. N., Pérez, G. L., Saltzman, E. S. and Simó, R.: A high-resolution time-depth view of dimethylsulphide cycling in the surface sea, *Sci. Rep.*, 6(November 2015), 1–13, doi:10.1038/srep32325, 2016.
- Saltzman, E. S., King, D. B., Holmen, K. and Leck, C.: Experimental determination of the diffusion coefficient of dimethylsulfide in water, *J. Geophys. Res.*, 98(C9), 481–486, doi:10.1029/93jc01858, 1993.
- 725 Sander, R.: Compilation of Henry's law constants (version 4.0) for water as solvent, *Atmos. Chem. Phys.*, 15(8), 4399–4981, doi:10.5194/acp-15-4399-2015, 2015.
- Schlundt, C., Tegtmeier, S., Lennartz, S. T., Bracher, A., Cheah, W., Krüger, K., Quack, B. and Marandino, C. A.: Oxygenated volatile organic carbon in the western Pacific convective center: Ocean cycling, air-sea gas exchange and atmospheric transport, *Atmos. Chem. Phys.*, 17(17), 10837–10854, doi:10.5194/acp-17-10837-2017, 2017.
- 730 Schwarz, K., Filipiak, W. and Amann, A.: Determining concentration patterns of volatile compounds in exhaled breath by PTR-MS, *J. Breath Res.*, 3(2), 027002, doi:10.1088/1752-7155/3/2/027002, 2009.


- Shaw, S. L., Chisholm, S. W. and Prinn, R. G.: Isoprene production by *Prochlorococcus*, a marine cyanobacterium, and other phytoplankton, *Mar. Chem.*, 80(4), 227–245, doi:10.1016/S0304-4203(02)00101-9, 2003.
- 735 Spirig, C., Neftel, A., Ammann, C., Dommen, J., Grabner, W., Thielmann, A., Schaub, A., Beauchamp, J. L., Wisthaler, A. and Hansel, A.: Eddy covariance flux measurements of biogenic VOCs during ECHO 2003 using proton transfer reaction mass spectrometry, *Atmos. Chem. Phys.*, 5(2), 465–481, doi:10.5194/acp-5-465-2005, 2005.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Am. Meteorol. Soc.*, 96(12), 2059–2077, doi:10.1175/BAMS-D-14-00110.1, 2015.
- 740 Tanner, C. B. and Thurtell, G. W.: Anemometer measurements of Reynolds stress and heat transport in the atmospheric surface layer., 1969.
- Uncles, R. J., Stephens, J. A. and Harris, C.: Physical processes in a coupled bay-estuary coastal system: Whitsand Bay and Plymouth Sound, *Prog. Oceanogr.*, 137, 360–384, doi:10.1016/j.pocean.2015.04.019, 2015.
- Upstill-Goddard, R. C.: Air-sea gas exchange in the coastal zone, *Estuar. Coast. Shelf Sci.*, 70(3), 388–404, doi:10.1016/j.ecss.2006.05.043, 2006.
- 745 Veres, P. R., Andrew Neuman, J., Bertram, T. H., Assaf, E., Wolfe, G. M., Williamson, C. J., Weinzierl, B., Tilmes, S., Thompson, C. R., Thames, A. B., Schroder, J. C., Saiz-Lopez, A., Rollins, A. W., Roberts, J. M., Price, D., Peischl, J., Nault, B. A., Møller, K. H., Miller, D. O., Meinardi, S., Li, Q., Lamarque, J. F., Kupc, A., Kjaergaard, H. G., Kinnison, D., Jimenez, J. L., Jernigan, C. M., Hornbrook, R. S., Hills, A., Dollner, M., Day, D. A., Cuevas, C. A., Campuzano-Jost, P., Burkholder, J. B., Paul Bui, T., Brune, W. H., Brown, S. S., Brock, C. A., Bourgeois, I., Blake, D. R., Apel, E. C. and Ryerson, T. B.: Global airborne sampling reveals a previously unobserved dimethyl sulfide oxidation mechanism in the marine atmosphere, *Proc. Natl. Acad. Sci. U. S. A.*, 117(9), 4505–4510, doi:10.1073/pnas.1919344117, 2020.
- 750 Wang, S., Hornbrook, R. S., Hills, A., Emmons, L. K., Tilmes, S., Lamarque, J. F., Jimenez, J. L., Campuzano-Jost, P., Nault, B. A., Crouse, J. D., Wennberg, P. O., Kim, M., Allen, H. M., Ryerson, T. B., Thompson, C. R., Peischl, J., Moore, F., Nance, D., Hall, B., Elkins, J., Tanner, D., Huey, L. G., Hall, S. R., Ullmann, K., Orlando, J. J., Tyndall, G. S., Flocke, F. M., Ray, E., Hanisco, T. F., Wolfe, G. M., St. Clair, J., Commane, R., Daube, B., Barletta, B., Blake, D. R., Weinzierl, B., Dollner, M., Conley, A., Vitt, F., Wofsy, S. C., Riemer, D. D. and Apel, E. C.: Atmospheric Acetaldehyde: Importance of Air-Sea Exchange and a Missing Source in the Remote Troposphere, *Geophys. Res. Lett.*, 46(10), 5601–5613, doi:10.1029/2019GL082034, 2019.
- 755 Wang, S., Apel, E. C., Schwantes, R., Bates, K., Jacob, D., Fischer, E., Hornbrook, R. S., Hills, A., Emmons, L., Pan, L., Honomichl, S., Tilmes, S., Lamarque, J., Yang, M., Marandino, C., Saltzman, E., Bruyn, W., Kameyama, S., Tanimoto, H., Omori, Y., Hall, S., Ullmann, K., Ryerson, T., Thompson, C., Peischl, J., Daube, B., Commane, R., McKain, K., Sweeney, C., Thames, A., Miller, D., Brune, W. H., Diskin, G., DiGangi, J. and Wofsy, S.: Global Atmospheric Budget of Acetone: Air-Sea Exchange and the Contribution to Hydroxyl Radicals, *J. Geophys. Res. Atmos.*, 2, 1–23, doi:10.1029/2020JD032553, 2020a.
- 760 Wang, W.-L., Song, G., Primeau, F., Saltzman, E., Bell, T. G. and Moore, J. K.: Global ocean dimethyl sulfide climatology estimated from observations and an artificial neural network, *Biogeosciences Discuss.*, 1–26, doi:10.1002/essoar.10502127.1, 2020b.
- Warneck, P. and Moortgat, G. K.: Quantum yields and photodissociation coefficients of acetaldehyde in the troposphere, *Atmos. Environ.*, 62, 153–163, doi:10.1016/j.atmosenv.2012.08.024, 2012.
- 770 Warneke, C., De Gouw, J. A., Kuster, W. C., Goldan, P. D. and Fall, R.: Validation of atmospheric VOC measurements by proton-transfer-reaction mass spectrometry using a gas-chromatographic pre-separation method, *Environ. Sci. Technol.*, 37(11), 2494–2501, doi:10.1021/es026266i, 2003.
- Williams, J., Holzinger, R., Gros, V., Xu, X., Atlas, E. L. and Wallace, D. W. R.: Measurements of organic species in air and seawater from the tropical Atlantic, *Geophys. Res. Lett.*, 31(23), 1–5, doi:10.1029/2004GL020012, 2004.
- 775 Wofsy, S. C., Afshar, S., Allen, H. M., Apel, E. C., Asher, E. C., Barletta, B., Bent, J., Bian, H., Biggs, B. C., Blake, D. R., Blake, N., Bourgeois, I., Brock, C. A., Brune, W. H., Budney, J. W., Bui, T. P., Butler, A., Campuzano-Jost, P., Chang, C. S., Chin, M., Commane, R., Correa, G., Crouse, J. D., Cullis, P. D., Daube, B. C., Day, D. A., Dean-Day, J. M., Dibb, J. E., DiGangi, J. P., Diskin, G. S., Dollner, M., Elkins, J. W., Erdesz, F., Fiore, A. M., Flynn, C. M., Froyd, K. D., Gesler, D. W., Hall, S. R., Hanisco, T. F., Hannun, R. A., Hills, A. J., Hints, E. J., Hoffman, A., Hornbrook, R. S., Huey, L. G., Hughes, S., Jimenez, J. L., Johnson, B. J., Katich, J. M., Keeling, R. F., Kim, M. J., Kupc, A., Lait, L. R., Lamarque, J.-F., Liu, J., McKain, K., McLaughlin, R. J., Meinardi, S., Miller, D. O., Montzka, S. A., Moore, F. L., Morgan, E. J., Murphy, D. M., Murray, L. T., Nault, B. A., Neuman, J. A., Newman, P. A., Nicely, J. M., Pan, X., Paplawsky, W., Peischl, J., Prather, M. J., Price, D. J., Ray, E. A., Reeves, J. M., Richardson, M., Rollins, A. W., Rosenlof, K. H., Ryerson, T. B., Scheuer, E., Schill, G. P., Schroder, J. C., Schwarz, J. P., St. Clair, J. M., Steenrod, S. D., Stephens, B. B., Strode, S. A., Sweeney, C., Tanner, D., Teng, A. P.,
- 780 Thames, A. B., Thompson, C. R., Ullmann, K., Veres, P. R., Vieznor, N., Wagner, N. L., Watt, A., Weber, R., Weinzierl, B., et al.: ATom: Merged Atmospheric Chemistry, Trace Gases, and Aerosols, , doi:10.3334/ORNLDAAAC/1581, 2018.
- Wohl, C., Capelle, D., Jones, A., Sturges, W. T., Nightingale, P. D., Else, B. G. T. and Yang, M.: Segmented flow coil equilibrator coupled to a proton-transfer-reaction mass spectrometer for measurements of a broad range of volatile organic

- compounds in seawater, *Ocean Sci.*, 15(4), 925–940, doi:10.5194/os-15-925-2019, 2019.
- 790 Wohl, C., Brown, I. J., Kitidis, V., Jones, A. E., Sturges, W. T., Nightingale, P. D. and Yang, M.: Underway seawater and atmospheric measurements of volatile organic compounds in the Southern Ocean, *Biogeosciences*, 17(9), 2593–2619, doi:10.5194/bg-17-2593-2020, 2020.
- Yang, M. and Fleming, Z. L.: Estimation of atmospheric total organic carbon (TOC) - Paving the path towards carbon budget closure, *Atmos. Chem. Phys.*, 19(1), 459–471, doi:10.5194/acp-19-459-2019, 2019.
- 795 Yang, M., Blomquist, B. W., Fairall, C. W., Archer, S. D. and Huebert, B. J.: Air-sea exchange of dimethylsulfide in the Southern Ocean: Measurements from so GasEx compared to temperate and tropical regions, *J. Geophys. Res. Ocean.*, 116(8), 1–17, doi:10.1029/2010JC006526, 2011a.
- Yang, M., Huebert, B. J., Blomquist, B. W., Howell, S. G., Shank, L. M., McNaughton, C. S., Clarke, A. D., Hawkins, L. N., Russell, L. M., Covert, D. S., Coffman, D. J., Bates, T. S., Quinn, P. K., Zagorac, N., Bandy, A. R., de Szoeke, S. P., 800 Zuidema, P. D., Tucker, S. C., Brewer, W. A., Benedict, K. B. and Collett, J. L.: Atmospheric sulfur cycling in the southeastern Pacific – longitudinal distribution, vertical profile, and diel variability observed during VOCALS-REx, *Atmos. Chem. Phys.*, 11(10), 5079–5097, doi:10.5194/acp-11-5079-2011, 2011b.
- Yang, M., Nightingale, P. D., Beale, R., Liss, P. S., Blomquist, B. W. and Fairall, C. W.: Atmospheric deposition of methanol over the Atlantic Ocean, *Proc. Natl. Acad. Sci.*, 110(50), 20034–20039, doi:10.1073/pnas.1317840110, 2013a.
- 805 Yang, M., Archer, S. D., Blomquist, B. W., Ho, D. T., Lance, V. P. and Torres, R. J.: Lagrangian evolution of DMS during the Southern Ocean gas exchange experiment: The effects of vertical mixing and biological community shift, *J. Geophys. Res. Ocean.*, 118(12), 6774–6790, doi:10.1002/2013JC009329, 2013b.
- Yang, M., Beale, R., Smyth, T. J. and Blomquist, B. W.: Measurements of OVOC fluxes by eddy covariance using a proton-transfer-reaction mass spectrometer - method development at a coastal site, *Atmos. Chem. Phys.*, 13(13), 6165–6184, 810 doi:10.5194/acp-13-6165-2013, 2013c.
- Yang, M., Blomquist, B. W. and Nightingale, P. D.: Air-sea exchange of methanol and acetone during HiWinGS: Estimation of air phase, water phase gas transfer velocities, *J. Geophys. Res. Ocean.*, 119(10), 7308–7323, doi:10.1002/2014JC010227, 2014a.
- Yang, M., Beale, R., Liss, P. S., Johnson, M. T., Blomquist, B. W. and Nightingale, P. D.: Air-sea fluxes of oxygenated 815 volatile organic compounds across the Atlantic Ocean, *Atmos. Chem. Phys.*, 14(14), 7499–7517, doi:10.5194/acp-14-7499-2014, 2014b.
- Yang, M., Bell, T. G., Hopkins, F. E., Kitidis, V., Cazenave, P. W., Nightingale, P. D., Yelland, M. J., Pascal, R. W., Prytherch, J., Brooks, I. M. and Smyth, T. J.: Air-sea fluxes of CO<sub>2</sub> and CH<sub>4</sub> from the penlee point atmospheric observatory on the south-west coast of the UK, *Atmos. Chem. Phys.*, 16(9), 5745–5761, doi:10.5194/acp-16-5745-2016, 2016a.
- 820 Yang, M., Bell, T. G., Blomquist, B. W., Fairall, C. W., Brooks, I. M. and Nightingale, P. D.: Air-sea transfer of gas phase controlled compounds, *IOP Conf. Ser. Earth Environ. Sci.*, 35(1), doi:10.1088/1755-1315/35/1/012011, 2016b.
- Yang, M., Bell, T. G., Hopkins, F. E. and Smyth, T. J.: Attribution of atmospheric sulfur dioxide over the English Channel to dimethyl sulfide and changing ship emissions, *Atmos. Chem. Phys.*, 16(8), 4771–4783, doi:10.5194/acp-16-4771-2016, 2016c.
- 825 Yang, M., Prytherch, J., Kozlova, E., Yelland, M. J., Parenkat Mony, D. and Bell, T. G.: Comparison of two closed-path cavity-based spectrometers for measuring air-water CO<sub>2</sub> and CH<sub>4</sub> fluxes by eddy covariance, *Atmos. Meas. Tech.*, 9(11), 5509–5522, doi:10.5194/amt-9-5509-2016, 2016d.
- Yang, M., Bell, T. G., Brown, I. J., Fishwick, J. R., Kitidis, V., Nightingale, P. D., Rees, A. P. and Smyth, T. J.: Insights from year-long measurements of air–water CH<sub>4</sub> and CO<sub>2</sub> exchange in a coastal environment, *Biogeosciences*, 16(5), 961– 830 978, doi:10.5194/bg-16-961-2019, 2019a.
- Yang, M., Norris, S. J., Bell, T. G. and Brooks, I. M.: Sea spray fluxes from the southwest coast of the United Kingdom-Dependence on wind speed and wave height, *Atmos. Chem. Phys.*, 19(24), 15271–15284, doi:10.5194/acp-19-15271-2019, 2019b.
- Zhou, S., Gonzalez, L., Leithead, A., Finewax, Z., Thalman, R., Vlasenko, A., Vagle, S., Miller, L. A., Li, S. M., Bureekul, 835 S., Furutani, H., Uematsu, M., Volkamer, R. and Abbatt, J. P. D.: Formation of gas-phase carbonyls from heterogeneous oxidation of polyunsaturated fatty acids at the air-water interface and of the sea surface microlayer, *Atmos. Chem. Phys.*, 14(3), 1371–1384, doi:10.5194/acp-14-1371-2014, 2014.
- Zhou, X. and Mopper, K.: Photochemical production of low-molecular-weight carbonyl compounds in seawater and surface microlayer and their air-sea exchange, *Mar. Chem.*, 56(3–4), 201–213, doi:10.1016/S0304-4203(96)00076-X, 1997.
- 840 Zhu, Y. and Kieber, D. J.: Wavelength- and Temperature-Dependent Apparent Quantum Yields for Photochemical Production of Carbonyl Compounds in the North Pacific Ocean, *Environ. Sci. Technol.*, 52(4), 1929–1939, doi:10.1021/acs.est.7b05462, 2018.



845

Figure 1: Schematic of the tubing set-up used to draw atmospheric sample to the gas instrument (PTR-MS). Dashed lines represent the set-up of the platinum catalyst before the synthetic gas blank was added. UT represents union-tee; a three-way static connection.



850

Figure 2: Mean flux cospectra for acetone and DMS in the SW and NE wind sectors (Dashed/broken lines represent PPAO observations). Solid lines represent the idealised spectral fits (Kaimal et al., 1972), where measurement height and wind speed are specified according to the measurement conditions.




Figure 3: Time series of 3 h averages of a) wind speed ( $U_{10}$ ) and wind direction, b) acetaldehyde flux, c) acetone flux and d) DMS fluxes for the [open-water](#) (180–240° N) and Plymouth Sound (30–70° N) wind sectors that met quality control criteria. Error bars are 1 standard error. [The blue and green shaded zone lines](#) in a subplot a represent the two wind sectors. [The blue grey shaded zones lines](#) in subfigures subplots b–d represent limits of detection for 3 h averages; 5.8, 4.0 and 2.7  $\mu\text{mol m}^{-2} \text{d}^{-1}$  for acetaldehyde, acetone, DMS respectively.

855






Figure 4: Histogram of a) wind speed ( $U_{10}$ ) and b) acetone and c) DMS atmospheric gas phase mixing ratio over the entire campaign. Histograms are normalised for probability density separately and then to the same relative maximum occurrence. The [DMS-gas mixing ratio calculations here](#) values use an extrapolated [background-blanks here](#), which has introduce some uncertainty, [h-H](#), therefore DMS concentrations are likely lower, however the two wind sectors use the same background and are can still be comparable.

860



|865 Figure 5: A comparison between EC and bulk TL fluxes (hourly) for the [open-water/open water](#) sector during the dates (27/04/18–03/05/18). The error on EC measurements is one SE, whilst the error on TL is the propagated error of one SD in concentration and solubility.



870 | **Figure 6: A comparison between a) DMS, b) acetone and c) acetaldehyde EC and TL fluxes (hourly) for the [open-water/open water](#) sector during the dates 27/04/18–03/05/18. Wind speed ( $U_{10}$ ) is provided in a). The error on EC measurements is one SE, whilst the error on TL is the propagated error of one SD in concentration and solubility.**






Figure 7: Calculated transfer velocities (hourly) for a) DMS ( $K_{w,600}$ ), b) acetone ( $K_a$ ), and c) acetaldehyde ( $K_a$ ) in the [open-water/open water](#) footprint of the Penlee Point Atmospheric Observatory during the dates 27/04/18–03/05/18. **Error bars on the experimental data (black circles) are propagated uncertainty from the eddy covariance measurement. The black poly fit lines are fit to the experimental data, with the dashed black lines representing 95 % confidence in the poly fit. The red circle and blue triangle lines in subplot A are theoretical transfer velocities using the  $k_w$  parameterisations of Yang et al. (2011a) and Nightingale et al. (2000) respectively. The black-red circle lines in subplots b and c are theoretical transfer velocities calculated from the  $k_w$  parameterisation of Yang et al. (2011a) and  $k_a$  from COAREG 3.5 model (shown as blue triangles).**

Table 1: Eddy covariance flux measurements over the entire campaign. Errors indicate standard errors (SE) with  $n=135$  and  $n=40$  hour-bins for the [open-water/open water](#) and Plymouth Sound sectors respectively. Also included are the quartiles (qrt).

| Compound     | <a href="#">Open-water/Open water</a> ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) |        |            |           | Plymouth Sound ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) |        |             |           |
|--------------|--|--------|------------|-----------|---|--------|-------------|-----------|
|              | 25 % qrt.  | Median | Mean       | 75 % qrt. | 25 % qrt.   | Median | Mean        | 75 % qrt. |
| Acetaldehyde | -7.44  | -1.23  | -1.55±1.14 | 4.93      | -10.86  | -4.45  | -4.45±1.73  | 1.84      |
| Acetone      | -12.15   | -6.27  | -8.01±0.77 | -2.19     | -17.52  | -9.47  | -12.93±1.37 | -2.19     |
| DMS          | 0.79   | 4.05   | 4.67±0.56  | 8.26      | -0.96   | 1.16   | 1.75±0.80   | 4.80      |
| Isoprene     | -3.54  | 0.60   | 1.71±0.73  | 5.82      | -8.84   | -0.24  | -1.95±1.65  | 3.56      |

Table 2: Saturations and two-layer bulk fluxes of VOC in [open-water/open water](#) (SW) and Plymouth sound (NE) water (27/04/18–03/05/18) along with the eddy covariance measurements for this period. Errors for water concentration and air mixing ratio (measurement) and saturation (propagation) indicate one standard deviation. Errors for flux indicate one standard error.

| Compound     | Open water       |                         |                |  |  | Plymouth Sound          |  |
|--------------|------------------|-------------------------|----------------|--|--|-------------------------|--|
|              | Water conc. (nM) | Air mixing ratio (ppbv) | Saturation (%) | TL flux ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) | EC flux ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) | Air mixing ratio (ppbv) | EC flux ( $\mu\text{mol m}^{-2} \text{d}^{-1}$ ) |
| Acetaldehyde | 7.1±1.4          | 0.51±0.07               | 44±6           | -6.76±0.59                                       | -3.94±1.23                                       | 0.75±0.04               | -7.32±2.86                                       |
| Acetone      | 5.3±2.0          | 0.82±0.16               | 15±3           | -18.98±2.24                                      | -14.93±2.09                                      | 1.09±0.09               | -8.21±4.00                                       |
| DMS          | 3.6±0.8          | 0.20±0.09               | 1434±569       | 9.41±1.03  | 9.61±1.55  | 0.015±0.006             | 2.55±2.36  |
| Isoprene     | 0.09±0.01        | 0.18±0.05               | 971±221        | 0.30±0.05  | 1.65±1.89  | 0.28±0.03               | 9.27±3.01  |

885 **Table A1: PTR-MS method used for measurements.**

| Parameter  | Value |
|--|-------|
| Drift chamber pressure (mBar)                      | 2.3   |
| Drift chamber voltage (V)                          | 690   |
| Drift chamber temperature ( $^{\circ}\text{C}$ )   | 80    |
| H <sub>2</sub> O mass flow (mL min <sup>-1</sup> ) | 5     |
| Source/proportional valve (%)                      | 49    |
| Inlet flow rate (mL min <sup>-1</sup> )            | 150   |
| Inlet temperature ( $^{\circ}\text{C}$ )           | 80    |

**Table A2: Flux segment quality control criteria.**

| Parameter   | Value         |
|---|---------------|
| Wind direction, $\sigma$ ( $^{\circ}$ )             | < 10          |
| Momentum flux (kg m <sup>-2</sup> s <sup>-1</sup> ) | < 0           |
| Tilt angle ( $^{\circ}$ )                           | < 15          |
| $\sigma_w/u_*$                                      | 1.2 < x < 1.9 |
| $\sigma_u^2/U$ (m s <sup>-1</sup> )                 | < 0.35        |

**Table A3: Fresh water solubility ( $H^{cp}$ ), solubility temperature dependence ( $\frac{d \ln(H^{cp})}{d(1/T)}$ ) and molar volume ( $V_b$ ) Chemical data used in calculations of seawater solubility and Schmidt number at ambient temperature and salinity.  $H^{cp}$  (and  $d \ln(H^{cp})/d(1/T)$ ) for freshwater are updated from Sander (2015) and Burkholder et al. (2015).**

| Compound     | $H^{cp}$ (mol L <sup>-1</sup> atm <sup>-1</sup> ) | $\frac{d \ln(H^{cp})}{d(1/T)}$ (K) | $V_b$ (cm <sup>3</sup> mol <sup>-1</sup> ) |
|--------------|---|------------------------------------|--|
| Acetaldehyde | 13.4  | 5775                               | 56   |
| Acetone      | 29.0  | 5170                               | 77.6                                       |
| DMS          | 0.541   | 3520                               | 77   |
| Isoprene     | 0.0345  | 4400                               | 105  |

890 **Table A4: Seawater concentrations of VOCs at the L4 marine station at 2 m depth. \* denotes concentration used for two-layer calculation.**

| Date     | Acetaldehyde (nM) | Acetone (nM) | DMS (nM) | Isoprene (nM) | Measurement method |
|----------|-------------------|--------------|----------|---------------|--------------------|
| 26/03/18 | 8.09*             | 3.93*        | 1.11     | 0.09*         | SFCE-PTRMS         |
| 10/04/18 | -                 | -            | 2.56     | -             | GC                 |
| 19/04/18 | -                 | -            | 10.34    | -             | GC                 |
| 23/04/18 | -                 | -            | 3.03*    | -             | GC                 |
| 30/04/18 | -                 | -            | 4.13*    | -             | GC                 |
| 21/05/18 | -                 | -            | 10.69    | -             | GC                 |
| 30/05/18 | 6.16*             | 6.71*        | 4.99     | 0.10*         | SFCE-PTRMS         |