

# South America 2020 Regional Smoke Plume: Intercomparison with previous years, impact on solar radiation and the role of Pantanal biomass burning season

Nilton Évora do Rosário<sup>1</sup>, Elisa Thomé Sena<sup>2</sup>, and Marcia Akemi Yamasoe<sup>3</sup>

<sup>1</sup>Departamento de Ciências Ambientais, Universidade Federal de São Paulo, Diadema, Brazil

<sup>2</sup>Departamento Multidisciplinar, Universidade Federal de São Paulo, Osasco, Brazil

<sup>3</sup>Departamento de Ciências Atmosféricas, Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, São Paulo, Brazil

Correspondence: Nilton M. Évora do Rosário ([nrosario@unifesp.br](mailto:nrosario@unifesp.br))

**Abstract.** The 2020 biomass burning season in Brazil was marked by an atypical amount of fire across the Pantanal biome, which led to high levels of smoke within the biome and downwind areas. The present study analyzes fire counts and smoke over Pantanal in 2020, comparing this particular year's data with those from the previous seventeen years (2003-2019). Taking as references the most polluted years in this period, the regional smoke plume and its impact on surface solar radiation were also evaluated. In 2020, the regional smoke plume core covered an area of ~2.6 million km<sup>2</sup> at the peak of the burning season, an area well above that of the previous six years, but smaller than areas observed in a more remote past, as in 2007 and 2010 (> 5.0 million km<sup>2</sup>). The smoke loading was lower (mean AOD550 nm ~0.7) than that of 2007 and 2010 (mean AOD550 nm ~1.0). The plume radiation absorption efficiency, when compared with the previous year's plumes, did not present significant differences. Regarding the Pantanal burning season, it revealed some atypical features. Fire counts were up to 3.0 times higher than for the years from 2003 to 2019. Smoke loading over Pantanal, which is typically a fraction of that over Amazonia, in 2020 was higher than over Amazonia, an indication that local smoke surpassed smoke advection from upwind regions. The observed intraseasonal variability of smoke over Pantanal revealed to be largely driven by the nature of the burned areas in the biome. From September on, there was a significant increase of fire count in conservation and indigenous areas, where higher biomass density is present, which would explain the larger smoke plumes over Pantanal, even during October when the fire count was reduced. In October, the biome was covered by a thick smoke layer, which resulted in a mean deficit of surface solar radiation up to 200 Wm<sup>-2</sup>. Despite Pantanal biomes' massive burning in 2020, the regional smoke plume was not far from its climatological features. Nevertheless, Pantanal 2020 burning season represents the worst combination of a climate extreme applied to a fire-prone environment, coupled with inadequately enforced environmental regulations, from which there is much to be learned.

**Comentado [NR1]:** Adjusted to reply to the reviewer Query 01 (Please, see point-by point reply)

**Comentado [NR2]:** Adjusted to reply to the reviewer Query 02 (Please, see point-by point reply)

**Comentado [NR3]:** Adjusted in the conclusions to reply to the reviewer Query 18 (Please, see point-by point reply)

## 32 1 Introduction

34 In South America, the Regional Smoke Plume (RSP) produced during the biomass burning season is the most important  
signature of anthropogenic activities from the point of view of injection of pollutants into the atmosphere at continental scale  
(Prins and Menzel, 1992, Artaxo et al., 1998, Freitas et al., 2004, Longo et al., 2007). Its geographical dimension and pollutants  
36 loading, along with its climate and air quality effects, have been comprehensively studied during the last decades (Freitas et  
al., 2004, Ignotti et al., 2010, Artaxo et al., 2013, Rosario et al., 2013, Chen et al., 2013, Sena et al., 2015, Moreira et al. 2017,  
38 Thornhill et al., 2018). The RSP affects the regional climate reducing the availability of solar energy at the surface and  
perturbing the cloud microphysics and atmospheric thermodynamics and chemistry (Procópio et al., 2001, Schafer et al., 2002,  
40 Yamasoe et al., 2006, Rosario et al., 2013, Moreira et al. 2017). In the context of air quality, especially near biomass burning  
areas, RSP impact on air pollution levels surpasses the most polluted urban areas of the continent (Ignotti et al., 2010, Sena et  
42 al., 2013, Rosario et al., 2013). Systematically, along the years, the southern portion of the Amazon rainforest and the eastern  
part of the Cerrado ecosystem have been the major sources of pollutants to the RSP. Consequently, in general, these are the  
44 most affected areas by the smoke pollution (Artaxo et al., 2013, Pereira et al., 2016), in particular, when compared with the  
remaining Brazilian biomes, Pantanal, Caatinga, Mata Atlantica and Pampas (Figure 1). From the mid-2000s to the earliest  
46 years of the 2010 decade, following the trends observed in deforestation and fire counts in Amazonia and Cerrado, the RSP  
loading and dimension were significantly reduced (Reddington et al. 2015). However, in recent years, due to the increase in  
48 deforestation rates and occurrence of major biomass burning events, such as those occurred during 2020 in Pantanal and related  
to the massive regional smoke plume in 2019, that contribute to darkened cities in the southeastern region of Brazil, there is a  
50 need to perform a historical contextualization of these recent events. Therefore, one can realize the advance achieved in the  
task of bringing down the RSP loading and also to highlight the risks of jeopardizing such achievement with the current  
52 environmental governance scenario in Brazil. The 2020 biomass burning in Brazil, amid the COVID-19 pandemic, has been  
claimed to be one of the worst in recent years, an evaluation largely driven by the extension of fire count and burned area  
54 across Pantanal biome (WWF, 2020, FSP, 2020, BBC, 2020, NYT, 2020, Le Monde, 2020, FSP, 2021). Despite a significant  
number of news and several literatures produced focusing on the 2020 biomass burning season (Libonati et al., 2020, Marengo  
56 et al., 2021, Pletsch et al., 2021), there is still a lack of clarity about what was effectively exceptional during this Brazil  
particular biomass burning season. From an integrated perspective of fire counts and RSP loading, and focusing on the biomes  
58 most affected by the RSP, Amazonia, Cerrado and Pantanal, this manuscript sets out to analyze how the 2020 biomass burning  
season compares with previous biomass burning seasons. To what extent fire count in Amazonia, Cerrado and Pantanal biomes  
60 during the 2020 biomass burning season and the produced RSP surpassed the previous years? Did Pantanal biome's exceptional  
2020 biomass burning season resemble an exceptional RSP? These are some of the questions that the present analysis sought  
62 to address in order to contribute to a comprehensive contextualization of Amazonia, Cerrado and Pantanal 2020 biomass  
burning season. Additionally, we also explore the 2020 RSP loading and its impact on surface solar radiation when compared  
64 with the RSPs from previous highly polluted years. From here on, the article is organized as follows: Section 2 presents a brief

**Comentado [NR4]:** Adjusted to reply to the reviewer Query 03  
(Please, see point-by point reply)

description of the study region, a summary of the data used and the methods adopted; Section 3 presents the results divided in  
66 2 sub-sections: Sub-section 3.1 analyses the intraseasonal and interannual variability of fire counts, aerosol optical depth  
(AOD) at 550 nm for the three biomes, Amazon tropical rainforest, Cerrado ecosystem, and Pantanal, for the period from 2003  
68 to 2020. Sub-section 3.2 focuses on September and October of the most polluted years of the considered period, providing an  
interannual analysis of the spatial distribution of smoke loading, wind field and the anomaly of downward solar radiation at  
70 the surface under cloudless conditions. In Section 4, the main conclusions are summarized.

## 2 Study Region, Data and Methods

72 This study focused on the three Brazilian biomes most affected by the regional smoke plume: Amazon tropical forest (or  
Amazonia), Cerrado ecosystem and Pantanal wetlands (Figure 1). The Pampa grasslands biome, in the south of the country,  
74 and the rainforest of Mata Atlantica, distributed along the Brazilian coast, from the northeast to the southern states, are less  
exposed to the RSP. They are eventually affected by the occurrence of transport of smoke toward southeast and southern  
76 Brazil. Caatinga biome, located in the northeast of Brazil and composed of shrubland and dry forests, is in general upwind of  
Amazonia and Cerrado, therefore it is less affected by the RSP. According to Pivello (2011), the Amazon tropical forest is a  
78 fire-sensitive ecosystem and cannot tolerate fire, with consequent mortality of trees after repeated fires. Although Pantanal and  
Cerrado are considered fire-dependent biomes and, therefore, are more adapted to fire occurrence, human-driven fire frequency  
80 increased significantly in Brazil, which can lead to land degradation and biodiversity loss (Pivello, 2011). Despite its smaller  
domain, when compared to Amazonia and Cerrado, Pantanal consists of one of the world's largest freshwater wetland  
82 ecosystems, characterized by highly complex hydrological regime that support two major river systems, the Cuiabá and the  
Paraguay rivers, and an abundance of vegetation and animal life (Alho, 2008, Alho et al., 2019). Large scale biomass burning  
84 within its domain and the RSP affect the local and regional climate equilibrium and also pose a direct threat to the health of  
Pantanal's singular biodiversity and population. MODIS (Moderate Resolution Imaging Spectroradiometer) Enhanced  
86 Vegetation Index (EVI, Kidan 2021) was applied in land use characterization across Pantanal during the 2020 burning season,  
specially to highlight conservation and indigenous areas, which tend to present higher biomass density.  
88 To analyze the RSP spatial dimension and loading, monthly mean aerosol optical depth at 550 nm, from 2003 to 2020, taken  
from MODIS atmosphere Level-3 products from Aqua satellite (MYD08, Platnick et al., 2017) was analyzed. The RSP border  
90 definition was somehow arbitrary, since the dispersion of the smoke is a continuous process and it is impossible to track the  
effective ending of its influence. To provide a perspective on the plume spatial dimension and loading at its core, isolines of  
92 AOD at 550 nm of 0.5 were used as reference to evaluate the plume dimension. The value of 0.5 was found convenient since  
it is above typical monthly average values of AOD at 550 nm observed over major urban areas of South America, which  
94 prevents a misidentification related to these urban plumes. Based on this threshold to define the main area of the plume, the  
RSP loading reference was taken as the mean AOD at 550 nm within the delimited area. The mean AOD at 550 nm for each  
96 year was also calculated for the domains of the biomes Amazonia, Cerrado and Pantanal. AERONET (Holben et al., 1998)

**Comentado [NR5]:** Adjusted to reply to the reviewer Query 04  
(Please, see point-by point reply)

**Comentado [NR6]:** Adjusted to reply to the reviewer Query 05  
(Please, see point-by point reply)

**Comentado [NR7]:** Adjust to reply to the reviewer Query 07  
(Please, see point-by point reply)

**Comentado [NR8]:** Adjust to reply to the reviewer Query 08  
(Please, see point-by point reply)

**Comentado [NR9]:** Adjusted to reply to the reviewer Query 06  
(Please, see point-by point reply)

aerosol optical depth at 550 nm, estimated using Angström exponent at channels 440 nm and 675 nm and AOD at 440 nm, single scattering albedo at 440 nm (Dubovik and King, 2000) and aerosol direct radiative forcing at the surface (Garcia et al., 2012), level 1.5, from three sites located in different parts of Brazil and one in Bolivia, were included in the analysis. Such data were used to help identify differences in the aerosol intrinsic optical properties from different years and locations within the RSP influence area. Since no level 2.0 data are available for the most recent period, we compared AERONET data from levels 1.5 and 2.0 from previous years in Figure S.1 (Supplement), to demonstrate the good quality of level 1.5 data as well.

To better contextualize the 2020 biomass burning season, we also analyzed the interannual variability (2003 to 2020) of the fire counts in each of the considered biomes. Table 1 presents the variables used and their respective applications, data sources and references. Fire counts were obtained from the Brazilian Space Agency (Instituto Nacional de Pesquisas Espaciais - INPE). These are important data to explore the spatial dynamic of smoke emission sources and, to some extent, to analyze the distribution of illegal biomass burning, which has been the main driver of the regional smoke plume in Brazilian biomes. The fire counts used are based on Moderate Resolution Spectroradiometer (MODIS) data and two algorithms. In summary, according to Morissette et al. (2005), the algorithms use empirically derived thresholds based on digital numbers (DNs) at channels 20 (at around 3.7  $\mu\text{m}$ ) and 9 (around 440 nm). In the daytime algorithm, pixels are classified as “fire” if two conditions are satisfied: DN higher than 3000 at channel 20 and lower than 3300 at channel 9. The nighttime algorithm requires only one condition: DN higher than 3000 at channel 20.

To estimate the direct effect of the aerosol plume on the reduction of surface solar radiation (SSR), instantaneous retrievals of CERES (Cloud and the Earth’s Radiant Energy System) Single Scanner Footprint - Level 3, during Aqua overpasses were used. This variable is estimated by a combination of CERES SW upward irradiance at the top of the atmosphere retrievals, ancillary meteorological data, surface, aerosol, gases and cloud properties. These inputs are used in the Langley parameterized shortwave algorithm (LPSA) (Kratz et al., 2020), to estimate shortwave downward irradiance at the surface from 0.2 to 5.0  $\mu\text{m}$ . The output is gridded in 1° by 1° latitude/longitude resolution. Cloud-free scenes, when cloud fraction is less than 0.1% in a given footprint (NASA, 2018), in the presence of aerosols were selected to account only for the aerosol direct effect on radiation. The aerosol model used in CERES surface irradiance calculations is very briefly described in Kratz et al. (2020). It uses near-real time daily AODs from the Model for Atmospheric Transport and Chemistry (MATCH) (Collins et al., 2001, Rasch et al., 1997). The Optical Properties of Aerosols and Clouds (OPAC) Global Aerosol Data Set (GADS) was used to obtain the intrinsic aerosol properties, such as single scattering albedo and asymmetry parameter (Hess et al., 1998).

Wind circulation at 850 hPa plays a determinant role on the transport of smoke over South America (Freitas et al. 2004) and, therefore, on the regional smoke plume structure, especially for areas downwind of the main biomass burning areas, namely, south of Amazon rainforest and the western portion of Cerrado ecosystem. Due to its location, Pantanal biome is subject to smoke transport from Amazon Forest and Cerrado. The wind dataset to perform this analysis was taken from MERRA-2 (Modern-Era Retrospective analysis for Research and Applications, version 2) reanalysis (Gelaro, et al., 2017). The interannual variability of the fire counts and AOD were analyzed, based on monthly mean values (July, August, September and October), and from the perspective of the selected biomes. Subsequently, the interannual variability of single scattering albedo and mean

**Comentado [NR10]:** Figure A.1 in previous version included as Appendix, now was considered as Supplement and was saved and uploaded independently as supplement file

**Comentado [NR11]:** Adjust to reply to the reviewer Query 09 (Please, see point-by point reply)

**Comentado [NR12]:** Adjust to reply to the reviewer Query 10 (Please, see point-by point reply)

**Comentado [NR13]:** Adjust to reply to the reviewer Query 11 (Please, see point-by point reply)

radiative forcing were analyzed. Those variables were also based on monthly mean values from AERONET stations located in different positions of the regional plume. September and October were selected to carry out a more comprehensive geographical analysis of the features and impact on SSR of the 2020 regional smoke plume. Targeting a comparison analysis, RSPs of the most polluted years in the period analyzed were included in this analysis. The radiative impact of these plumes, based on the anomaly in downward SSR, was also compared.

**Comentado [NR14]:** Adjust to reply to the reviewer Query 12  
(Please, see point-by point reply)

### 3 Results

#### 3.1 Fire count and smoke aerosol loading over Amazon, Cerrado and Pantanal: intraseasonal and interannual variability

This section addresses the interannual and the intraseasonal variability of monthly fire counts and mean smoke aerosol loading for the three selected biomes (Amazon, Cerrado and Pantanal) during the biomass burning season of the last 18 years, from 2003 to 2020. Figure 2 shows, for each biome and multiple years, the intraseasonal evolution (July, August, September and October) of fire counts and smoke aerosol loading. The typical intra seasonality of the biomass burning season is observed for most of all the 18 years as follows: from the transition month of July, the number of fire counts increased significantly in the following months, in general, reaching its peak in September. After that, the number of fire counts decreased, but still remained high, in October, until the next wet season. The burned area variability, not shown here, closely follows the variability of fire counts. For the Amazonia biome, in terms of fire count and mean smoke aerosol loading, 2020 biomass burning season closely followed the general features of the last nine years (2011 – 2020). Regarding the peak of smoke aerosols over Amazonia, in the last 10 years, September 2017 overpowered all Septembers, including that of 2020. Looking further back in time, the Amazonia biomass burning seasons of the years of 2004, 2005, 2007 and 2010 were the most fire active and polluted years, with mean AOD at 550 nm over the biome reaching a value close to one, in some case almost twice as that observed in 2020. A similar feature can be stated for fire count. Regarding the intraseasonal behavior, for all highly polluted biomass burning seasons, September stood out as the most polluted month in Amazonia. For the Cerrado ecosystem, while fire count during the 2020 biomass burning season was not significantly different from the previous nine years (2011 – 2020) biomass burning seasons, smoke aerosol loading was exceptionally high in October, but with precedent when one looks further in the past (ex. 2007). Similar to Amazonia, during the mid-2000's Cerrado presented its more fire active and polluted years (2004, 2005, 2007) in the period here analyzed, and including 2010. However, when one looks at the Pantanal biome biomass burning seasons time series, 2020 was indeed an exceptional season in terms of fire count, with almost three and four times the fire counts typically observed in August and September of the last decades, respectively. Those figures surpass the fire counts of the years of 2005 and 2007, two of the most fire active biomass burning seasons in the last two decades in Pantanal. The 2020 high fire counts scenario for Pantanal is also seen in aerosol mean loading over the biome during September, with twice the values observed in recent years (2011 - 2020), but not unprecedented in the history of Pantanal. For the years of 2004, 2005, 2007 and 2010, Pantanal smoke aerosol loading was as high as, or even higher than, that observed in the 2020 peak. In 2007,

during September, the highest aerosol loading was observed within the period analyzed. Unlike 2020, in the polluted years of 2000s and the year 2010, large smoke loading was also observed upwind of Pantanal, in the southern of Amazon Forest and western part of Cerrado, as can be seen in the Hovmoller diagram of AOD at 550 nm (Figure 3). This indicates the relevant contribution of smoke advection from these biomes to the high aerosol loading observed over Pantanal, especially during 2010 when local fire counts were relatively low. From 2011 to 2019, both fire counts and smoke levels in Pantanal were well below the observed values during the previous polluted years, even when high levels of smoke were observed in the upwind regions, for instance in 2017 (Figure 3). In that year, the core of the transport toward the south was over Paraguay. Regarding the high levels of smoke in the northern part of the Amazon basin at the end of the year 2015 (Figure 3), and which did not have a significant influence on Pantanal region, it is worth mentioning that it occurred during an El Nino event, when this part of the Amazon typically experiences drought scenarios and higher incidence of fire counts. Indeed, according to Marengo et al. (2017), the Amazon onset of the rainy season in 2015 occurred later than normal, and the region was characterized by drought in 2016. As in 2015 (Marengo et al., 2017) and 2020 (Marengo et al., 2021), drought conditions also were present, either across central or north region of Brazil, during the polluted years of 2005, 2007 and 2010 (Jimenez et al., 2018, Libonati et al., 2021), corroborating the critical role of climate variability combined with human factors to the exacerbation of biomass burning severity in Brazil. Regarding the years characterized by less polluted scenarios in Pantanal (2011-2019), they were consistently associated with relatively low local fire counts. This scenario changed during the 2020 biomass burning season, when the total fire counts at the peak of the biomass burning season in Pantanal surpassed all the previous analyzed years (Figure 2). According to Marengo et al. (2021), since the beginning of the fire activity monitoring in Brazil, in 1998, the largest number of fires over Pantanal was detected in 2005 and, in 2020, it was 76% higher. As mentioned, the level of aerosol loading over the biome at the peak of the 2020 burning season was similar to the early polluted years (Figure 2), in the middle of 2000s. However, opposite to the former years, smoke loading in Pantanal in 2020 peaked in October, when the smoke loading upwind of Pantanal, mainly in the Amazon biome, was atypically much lower than that over Pantanal (Figure 3). This suggests that Pantanal itself was the main source of smoke in its domain, and with advection from outer regions playing a secondary role. October followed also a highly polluted September, when there was an explosion in fire counts across Pantanal. The increase in smoke over Pantanal, in mid-September, occurred simultaneously with the peak of smoke in Amazonia, suggesting that at this time the smoke plume over Pantanal had also received contribution from biomass burning emission from Amazonia (Figure 3). While the aerosol loading observed during the 2020 biomass burning over Pantanal was not unique from historical perspective, there are relevant differences between 2020 and previous polluted years. Regarding the intraseasonal variability of smoke aerosol loading, October 2020 in Pantanal can be evaluated as exceptional when compared with October from the previous years of the time series. An interesting aspect worth emphasizing for October 2020 is that Pantanal presented much lower fire counts compared to August and September (Figure 3). However, August's higher fire count in Pantanal did not translate into a high level of smoke over the wetland biome. Further analysis was done to clarify these aspects, focusing on the correlation between smoke loading over Pantanal and local fire count (Figure 4a) and on the correlation between smoke over Pantanal and over Amazonia (Figure 4b). The scatter plot of smoke loading over Amazonia versus smoke over Pantanal shows

**Comentado [NR15]:** Adjust to reply to the reviewer Query 13 (Please, see point-by point reply)

**Comentado [NR16]:** Figure 4 from previous version was splitted in two different figures in this new version (Figure 4 and Figure 5). This was done in order to assure the best contextualization and quality for the plots. The text in which they are discussed was not modified, we only adjusted the references of the next figures so we can keep the consistency of the discussion.

198 that the smoke loading over Pantanal has a stronger relationship with smoke over Amazon than with fire counts within the  
200 biome. This suggests that, in general, smoke over Pantanal is more affected by advection than by local fires. However, 2020  
202 stands out as an outlier. Typically, mean AOD at 550 nm over Pantanal domain is similar or a fraction of that over Amazonia  
204 domain. That was not the case for September and October of 2020, when mean AOD at 550 nm over Pantanal was much higher  
206 than over Amazonia, an indication that locally produced smoke played a major role on the amount of smoke over the Pantanal  
208 atmosphere column.

To analyze why in August 2020 the higher fire count did not translate in higher smoke amount, compared with September and  
204 October 2020, the monthly distribution of fire count across Pantanal (Figure 5a) was analyzed as a function of land used  
(conservation units and indigenous areas) and Enhanced Vegetation Index (EVI). The analysis (Figures 5) shows that, despite  
206 the large fire count occurrence, August presented a reduced number of fires within conservation areas (where higher biomass  
208 density is present). However, during September and October, there was a significant increase of fire number within those areas,  
which could explain the larger aerosol emissions and, consequently, higher AOD values over Pantanal compared to August.

A possible shift in the mean composition of the regional smoke aerosol plume optical properties (mainly light absorption  
210 efficiency) during the 2020 biomass burning season was also explored via retrievals of single scattering albedo (SSA) from  
AERONET stations. Since there are no AERONET stations operating in Pantanal, the analysis of SSA was carried out from  
212 stations in the neighbourhood of Pantanal (Cuiaba) and somewhere else across Amazonia and at Santa Cruz, in Bolivia. The  
results obtained did not show significant change between mean SSA for 2020 biomass burning season and from previous years  
214 (Figure 6). Referring to Cuiaba, the closest AERONET site to Pantanal, the mean SSA on 2020 was within the site typical  
variability, that also was the case of Rio Branco, Alta Floresta and Santa Cruz sites. Multiyear instantaneous aerosol radiative  
216 forcing versus AOD at 550 nm from these AERONET stations, and as function of SSA, were also analyzed (Figure 7) by  
comparing 2020 with the previous years (2003 to 2019). It is possible to observe that not only AOD but also SSA affects the  
218 downward solar irradiance at the surface. However, no exceptional difference can be noticed in 2020 data when compared  
with the entire dataset. It is worth mentioning that the AERONET data from quality level 2.0 for 2020 is not yet available, so  
220 level 1.5 was used. Therefore, future and further analysis on this matter is highly recommended in order to evaluate the posed  
hypotheses, although as observed in Figure A.1, there is very good agreement between levels 1.5 and 2.0 SSA retrievals.

222

### 3.2 Regional Smoke Plume and Surface Solar Radiation anomaly: Spatial and interannual analysis

224 Climatologically, September is the peak of the biomass burning season over a large area of South America, and of Brazil in  
particular. Therefore, September was selected to explore the spatial distribution, loading and impact on the downward solar  
226 radiation at the surface of the regional smoke plume for 2020 and the most polluted years in the time series analyzed, namely  
2004, 2005, 2007, 2010, 2017. Considering the particularity of smoke over Pantanal in October of 2020, that month was also  
228 added to this analysis of the RSP feature and impact on SSR. Figure 8 presents the interannual variability of the geographical  
distribution of the RSP for September and October for the most polluted years. Although it was one of the most polluted in the

230 last five recent years, the 2020 regional plume could not be considered as an exceptional plume when compared, for instance,  
232 with the historical RSP of the years 2007, 2010 and 2005. Among the years analyzed, 2007 stands as the top polluted, for both  
234 September and October. However, during October 2020, it is possible to visualize that the peak of the smoke over the continent  
236 was centered in Pantanal (Figure 8b), corroborating the previous discussion that 2020 biomass burning exceptionality was  
238 restricted to the Pantanal biome. The area of the RSP core, considered as the extension of the smoke plume delimited by the  
isoline of AOD at 550 nm of 0.5 (area) and its respective mean intensity (AOD at 550 nm level) are also identified. The 2007  
RSP presented the largest (5740811 km<sup>2</sup>) and most polluted domain (mean AOD at 550 nm > 1.0), a characteristic that extended  
to the month of October. The area and AOD at 550 nm figures during September for the 2020 RSP were, respectively, 2450641  
km<sup>2</sup> and 0.68.

Due to the wind circulation pattern in the region, in general, the smoke from fires in the Amazon Forest and Cerrado ecosystem  
are transported westward reaching other South-American countries until it reaches the Andes Mountain range barrier (Freitas  
et al., 2004), where the wind circulation becomes predominantly meridional and southwards with a variable zonal component  
that can fluctuate the transport of smoke between the southeast and the southern portions of Brazil. Going southwards, the  
plume can reach the Pantanal region, among other locations, but the smoke usually gets diluted during this transport process.  
Related to the flow of the regional smoke plume towards the population centers of the southeastern coast, a stronger feature of  
2020 regional smoke plume, the analysis of previous years evidenced that this has also been seen in the past. For example, in  
2004 and 2005 the monthly flow patterns were also towards the highly populated centers in the southeast of Brazil.

Figure 9 also shows that the anomaly of solar radiation in cloud-free conditions occurred over the regions more severely  
affected by the aerosol plume, consistently with the spatial distribution of AOD, as expected. Again, in 2020, the solar energy  
reduction was surpassed by previous top polluted years, specially 2007, when SSR anomaly over a large portion of the  
continent, including the western portion of Pantanal, reached values up to -300 Wm<sup>-2</sup>. It is also possible to see that in 2005,  
as occurred in 2020, the southeast portion of Brazil experienced a larger SSR reduction as a result of the persistent transport  
of smoke toward that region. As suggested by the AOD field, available solar energy reaching the ground in October 2020 over  
Pantanal was significantly reduced (up to 200 Wm<sup>-2</sup>).

#### 254 4 Conclusion

The 2020 biomass burning season in Brazil attracted unprecedented attention from national and international media as well as  
the general society. Pantanal biome was a hotspot in this entire discussion. The wetland biome's role in the Regional Smoke  
Plume (RSP) has been marginal throughout the years. However, with the explosion of fire counts across Pantanal in 2020,  
there was a question about the role of Pantanal's smoke emissions to the RSP during the 2020 biomass burning season. In this  
study we analyzed to what extent the RSP produced during the 2020 biomass burning season and its impact on Solar Surface  
Radiation under cloudless scenarios differs from previous years, from 2003 to 2019, in particular the highly polluted former  
RSPs. Additionally, we analyzed the interannual and intraseasonal variability of fire counts and aerosol loading with emphasis

**Comentado [NR17]:** Adjust to reply to the reviewer Query 14  
(Please, see point-by point reply)



262 on the biomes Amazonia tropical rainforest, Cerrado and Pantanal. In the last eighteen years, from the point of view of the  
264 Amazon Forest and Cerrado ecosystem, 2020 can be considered as an ordinary biomass burning season, far from the most fire  
active and polluted years of 2010 and 2007. For these years, 2010 and 2007, Amazonia and the western portion of Cerrado  
266 experienced their largest fire counts and aerosol loading within the analyzed period. However, for Pantanal, under certain  
aspects, 2020 was a very particular year. Not exactly due to the smoke loading over the biome, since in September of 2007 the  
268 biome experienced a higher smoke loading, but due to the relative contribution of the local fire when compared to advection  
from upwind areas and the amount of smoke observed in October. The analysis revealed that, typically, Pantanal smoke loading  
270 has a stronger relationship with smoke over Amazon than with fire counts within the biome, suggesting that AOD over Pantanal  
is more affected by advection than by local fires. However, in 2020 that was not the case, local fire dominated. Usually, smoke  
loading over Pantanal is a fraction of that over Amazonia domain, but for September and October of 2020, mean smoke loading  
272 over Pantanal was much higher than over Amazonia, an indication that smoke produced locally played an atypical role to the  
smoke level over Pantanal. In the 2020 biomass burning season, fire counts in Pantanal were 3.4 times higher than the mean  
274 value from 2003 to 2020. As important as the amount of fire, the nature of the areas being burned within Pantanal revealed to  
determine the amount of smoke produced during September and October. From September on, there was a significant increase  
276 of fire counts within conservation and indigenous areas, where higher biomass density is present, which would explain the  
large smoke plumes over Pantanal, even during October when the number of fires were significantly reduced. The entire biome  
278 was continuously covered by a thick layer of smoke from west to east and from south to north for almost one month and a half,  
which resulted in a monthly (October) mean deficit of solar radiation at the surface up to 200 Wm<sup>-2</sup>. The impact of this  
280 reduction of incoming solar radiation on biological and surface-atmosphere interactions processes and are yet to be evaluated.  
Additionally, considering the plume transport towards the highly productive central and southeast regions of Brazil, the  
282 country's capacity of production of renewable energy based on solar radiation is also affected. In a period when cloud cover  
is less frequent and the generation of hydro-electric power, the main energy source in Brazil, decreases due to the lack of  
284 precipitation, revealing a need for a comprehensive governance.

In conclusion, from RSP loading and area perspective and when compared with previous years (2003-2019), 2020 biomass  
286 burning season could not be identified as an exceptional year. However, when focusing on the Pantanal domain, there are three  
aspects that differ from typical features, the fire count observed in September, the smoke loading over the Pantanal biome  
288 during October and the relative contribution of local fire to the local smoke loading when compared to the advection of smoke  
from Amazonia domain. Pantanal biomass burning has received less attention throughout the years, when compared with  
290 Amazonia and Cerrado biomes. However, Pantanal is a critical piece in the regional hydrological cycle, with strong connection  
with the two larger biomes. Pantanal surface-atmosphere interaction processes' role in the support of local and regional  
292 biodiversity is indisputable. Although it is too early to infer a possible pattern change in Pantanal biomass burning features,  
studies focusing on the 2020 atypical biomass burning season may shed light on potential scenarios that the region may  
294 experience in the future. Because of the likelihood of an increase in drought frequency and intensity, policy makers will need  
to take measures to mitigate future scenarios similar to the one shown here in 2020. Current knowledge on the Pantanal biome

**Comentado [NR18]:** Adjust to reply to the reviewer Query 15  
(Please, see point-by point reply)

**Comentado [NR19]:** Adjust to reply to the reviewer Query 16  
(Please, see point-by point reply)

**Comentado [NR20]:** Adjust to reply to the reviewer Query 17  
(Please, see point-by point reply)

296 fire emission dynamics and smoke plume radiative properties and impacts is far limited compared to the Amazon Forest and Cerrado ecosystem.

298 Although the 2020 regional biomass burning season was not among the most polluted in the history of biomass burning in Brazil, the Pantanal 2020 biomass burning season represents the worst combination of a climate extreme applied to a fire-prone environment, coupled with inadequately enforced environmental regulations (Marengo et al. 2021, Libonati et al., 2020, Vale et al., 2021). According to Marengo et al. (2021), the years of 2019 and 2020 were characterized by the worst drought in 50 years in Pantanal. Vale et al. (2021) pointed out a large reduction in environmental fines during the pandemic. This was caused by a decrease in fine enforcement, not due to a decrease in environmental violations, as an increase in Amazonian deforestation was observed. The slashing of funding for environmental protection and climate actions during the pandemic years has compounded the environmental harm imposed by legislative acts aimed at degrading environmental protection. There are still open questions about the specific behavior of mankind intervention in Pantanal in 2020, a lack of studies of human causes and responses to fires in the Pantanal has been recognized as a challenge to a full comprehension of what happened (Libonati et al., 2020). One also has to point out that there is a need to advance scientific knowledge on the Pantanal climate process in the context of both local environmental degradation and global climate change scenarios. This evaluation has as basis the notable lack of regular monitoring and comprehensive experimental campaigns focusing on the biome and its complex interdependence relationship with the neighborhood biomes, Amazonia and Cerrado and downwind regions.

312  
314 Data availability. All the dataset (AERONET, MODIS, CERES, MERRA-2, INPE fire counts) used in this study are publicly available and can be downloaded from their respective sites provided in Table 1 of this manuscript.

### 316 Supplement

318 The supplement related to this article is available online at: <XXXXXXXXXXXX>.

320 Author contributions. N. E. R., E. S. T and M. A. Y. designed and performed the research, analyzed the data, and wrote the paper.

322 Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was supported by FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) projects 2016/18438-0 and 2018/16048-6 and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), processes number 313005/2018-4, [311984/2021-5](#) and Universal 421870/2018-4. We also acknowledge the AERONET, MERRA-2, Programa Queimadas-INPE, CERES and MODIS mission scientists, Principal Investigators and associated NASA personnel for the production of the data used in this research effort. We would like to also thank the reviewers of this manuscript, Dr Lorraine Remer and an anonymous reviewer, for their thoughtful comments and efforts towards improving this manuscript.

**Comentado [NR21]:** Adjust to reply to the reviewer Query 18 (Please, see point-by point reply)

**Comentado [NR22]:** Adjust to reply to the reviewer Query 19 (Please, see point-by point reply)

**Comentado [NR23]:** This was included to consider the Supplement file that presents and intercomparison between AERONET datasets from level 1.5 and 2.0. In the previous version is was considered as an Appendix

**Comentado [NR24]:** We added this to say thank you to the reviewer and recognize they important role in the improvement of this manuscript.

## References

- 330 Abatzoglou JT, Williams AP (2016) Impact of anthropogenic climate change on wildfire across western US forests. *Proc Natl Acad Sci USA* 113:11770–11775.
- 332 Alho, C.J.R. Biodiversity of the Pantanal: Response to seasonal flooding regime and to environmental degradation. *Brazilian Journal of Biology*, 2008, 68(4 Suppl.), 957-966. <http://dx.doi.org/10.1590/S1519-69842008000500005>. PMID:19197468.
- 334 Alho, C.J.R., Mamede, S.B., Benites, M., Andrade, B.S. and Sepúlveda, J.J.O. Threats to the biodiversity of the Brazilian Pantanal due to land use and occupation. São Paulo: Ambiente & Sociedade, 2019, 22p
- 336 Artaxo, P., Fernandes, E. T., Martins, J. V., Yamasoe, M. A., Hobbs, P. V., Maenhaut, W., Longo, K. M., Castanho, A. 1998. Large Scale Aerosol Source Apportionment in Amazonia. *Journal of Geophysical Research*, 103 (D24): 31.837-31.848.
- 338 Artaxo, P., Rizzo, L. V., Brito, J. F., Barbosa, H. M. J., Arana, A., Sena, E. T., Cirino, G. G., Bastos, W., Martin, S. T., Andreae, M. O. Atmospheric aerosols in Amazonia and land use change: from natural biogenic to biomass burning conditions. *Faraday Discussions* 165, 203-235, 2013.
- 340 BBC, 2020. Amazon fires: Year-on-year numbers doubled in October. (Available at <https://www.bbc.com/news/world-latin-america-54779877>). Accessed on: April 3rd 2021
- 342 Braghieri, R. K., Yamasoe, M. A., Rosário, N. M. É., Rocha, H. R., Nogueira, J. S. and Araújo, A. C. Characterization of the radiative impact of aerosols on CO<sub>2</sub> and energy fluxes in the Amazon deforestation arch using artificial neural networks. *Atmos. Chem. Phys.*, 20, 3439–3458, doi: 10.5194/acp-20-3439-2020, 2020.
- 344 Chen, Yang, Douglas C. Morton, Yufang Jin, G. James Collatz, Prasad S. Kasibhatla, Guido R. van der Werf, Ruth S. DeFries, James T. Randerson. Long-term trends and interannual variability of forest, savanna and agricultural fires in South America, *Carbon Management*, 4:6, 617-638, DOI: 10.4155/cmt.13.61, 2013.
- 348 Collins, W. D., P. J. Rasch, B. E. Eaton, B. V. Khatatov, J.-F. Lamarque, and C. S. Zender. Simulating aerosols using a chemical transport model with assimilation of satellite aerosol retrievals: Methodology for INDOEX. *J. Geophys. Res.*, 106, 7313–7336, <https://doi.org/10.1029/2000JD900507>, 2001
- 350 Draxler, R.R., Hess, G.D., 1998. An overview of the Hysplit 4 modelling system for trajectories, dispersion and deposition. *Aust. Meteorol. Mag.* 47, 295–308
- 352 Du, J.; Kimball, J.S.; Reichle, R.H.; Jones, L.A.; Watts, J.D.; Kim, Y. Global Satellite Retrievals of the Near-Surface Atmospheric Vapor Pressure Deficit from AMSR-E and AMSR2. *Remote Sens.* 2018, 10, 1175. <https://doi.org/10.3390/rs10081175>
- 356 Dubovik, O. and King, M. D.: A flexible inversion algorithm for retrieval of aerosol optical properties from sun and sky radiance measurements, *J. Geophys. Res.*, 105, 20673–20696, 2000.
- 358 Escobar. H. Amazon fires clearly linked to deforestation, scientists say. *Science* 30 Aug 2019 Vol. 365, Issue 6456, pp. 853
- 360 DOI: 10.1126/sci-ence.365.6456.853

Fearnside, P.M., Righi, C.A., de Alencastro Graça, P.M.L., Keizer, E.W., Cerri, C.C., Nogueira, E.M. and Barbosa, R.I., 2009. Biomass and greenhouse-gas emissions from land-use change in Brazil's Amazonian "arc of deforestation": The states of Mato Grosso and Rondônia. *Forest Ecology and Management*, 258(9), pp.1968-1978.

Freitas, S.R.; Longo, K. M.; Silva Dias, M. A.F.; Silva Dias, P. L.; Chatfield, R.; Prins, E.; Artaxo, P.; Recuero, F. 2004. Monitoring the transport of biomass burning emissions in South America. *Environmental Fluid Mechanics*, 5(1): 135-167, DOI: 10.1007/s10652-005-0243-7.

FSP, 2020. Incontrolável, fogo já consumiu 26,5% do Pantanal, mostram satélites. (Available at <https://www1.folha.uol.com.br/ambiente/2020/10/incontrolavel-fogo-ja-consumiu-265-do-pantanal-mostram-satelites.shtml>). Accessed on: April 3rd 2021

FSP, 2021. About 40% of the Pantanal of Mato Grosso Was Burnt in 2020. (Available at <https://www1.folha.uol.com.br/internacional/en/scienceandhealth/2021/01/about-40-of-the-pantanal-of-mato-grosso-was-burnt-in-2020.shtml>). Accessed on: April 6th 2021

Garcia, O. E., Díaz, J. P., Expósito, F. J., Díaz, A. M., Dubovik, O., Derimian, Y., Dubuisson, P. and Roger, J.-C. Shortwave radiative forcing and efficiency of key aerosol types using AERONET data. *Atmos. Chem. Phys.* 12, 5129–5145, doi:10.5194/acp-12-5129-2012, 2012. Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Clim.*, 30, 5419–5454, <https://doi.org/10.1175/jcli-d-16-0758.1>, 2017.

Global Modeling and Assimilation Office (GMAO) (2015), MERRA-2 tavgM\_2d\_slv\_Nx: 2d, Monthly mean, Time-Averaged, Single-Level, Assimilation, Single-Level Diagnostics V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: [Data Access Date], 10.5067/AP1B0BA5PD2

Hess, M., P. Koepke, and I. Schult (1998). Optical Properties of Aerosols and Clouds: The software package OPAC. *Bull. Amer. Meteor. Soc.*, 79, 831–844, [https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2).

Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A. (1998). AERONET – A federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16.

Ignotti, Eliane, Valente, Joaquim Gonçalves, Longo, Karla Maria, Freitas, Saulo Ribeiro, Hacon, Sandra de Souza, Artaxo Netto, Paulo. (2010). Impact on human health of particulate matter emitted from burnings in the Brazilian Amazon region. *Revista de Saúde Pública*, 44(1), 121-130. <https://doi.org/10.1590/S0034-89102010000100013>

Didan, K. (2021), MOD13C2 MODIS/Terra Vegetation Indices Monthly L3 Global 0.05Deg CMG V061. NASA EOSDIS Land Processes DAAC. <https://doi.org/10.5067/MODIS/MOD13C2.061>; obtained from the Land Processes Distributed Active

394 Archive Center (LP DAAC), located at the U.S. Geological Survey (USGS) Earth Resources Observation and Science (EROS)  
Center (lpdaac.usgs.gov) [last access: January 26, 2022],

396 King, M. D., S. Platnick, W. P. Menzel, S. A. Ackerman, and P. A. Hubanks (2013). Spatial and temporal distribution of clouds  
observed by MODIS onboard the Terra and Aqua satellites. *IEEE Trans. Geosci. Remote Sens.*, 51, 3826–3852.

398 Kratz, D. P., S. K. Gupta, A. C. Wilber, and V. E. Sthcote. (2020). Validation of the CERES Edition-4A Surface-Only Flux  
Algorithms, *J. Appl. Meteor. Climatol.*, 59(2), 281-295, doi: 10.1175/JAMC-D-19-0068.1

400 Jimenez, J. C.; Libonati, R.; Peres, L. F. Droughts over Amazonia in 2005, 2010 and 2015: a cloud cover perspective. *Frontiers  
in The Earth Science*, 2018.

402 Le Monde (2020). Le Pantanal, au Brésil, paradis de biodiversité ravagé par les flammes. (Available at  
[https://www.lemonde.fr/planete/article/2020/09/29/le-pantanal-paradis-de-biodiversite-ravage-par-les-  
404 flammes\\_6054087\\_3244.html](https://www.lemonde.fr/planete/article/2020/09/29/le-pantanal-paradis-de-biodiversite-ravage-par-les-<br/>404 flammes_6054087_3244.html)). Accessed on: April 3rd 2021

Levy, R., Remer, L., Mattoo, S., Vermote, E., and Kaufman, Y. J. (2007). Second-generation operational algorithm: Retrieval  
406 of aerosol properties over land from inversion of Moderate Resolution Imaging Spectroradiometer spectral reflectance, *J.  
Geophys. Res.-Atmos.*, 112(D13), D13211, 10.1029/2006JD007811.

408 Libonati, R., Dacamara, C., Setzer, A., Morelli, F., Melchiori, A. (2015). An Algorithm for Burned Area Detection in the  
Brazilian Cerrado Using 4  $\mu\text{m}$  MODIS Imagery. *Remote Sensing* 7, 15782-15803.

410 Libonati R., Da Camara C. C., Peres L. F., Sander de Carvalho L. A. and Garcia L. C. (2020). Rescue Brazil's burning Pantanal  
wetlands *Nature* 588 217–9

412 Libonati, R., Pereira, J.M.C., Da Camara, C.C. et al. Twenty-first century droughts have not increasingly exacerbated fire  
season severity in the Brazilian Amazon. *Sci Rep* 11, 4400 (2021). <https://doi.org/10.1038/s41598-021-82158-8>

414 Longo, K., Freitas, S. R., Andreae, M.O., Yokelson, R., and Artaxo, P.: Biomass burning in Amazonia: emissions, long range  
transport of smoke and Its regional and remote Impacts, in: *Amazonia and Global Change*, by the American Geophysical Union  
416 Press, edited by: Gash, J. and Keller, M., Mercedes Bustamante, Pedro Silva Dias, 2009

Marengo, J.A., Cunha, A.P., Cuartas, L.A., Deusdará Leal, K.R., Broedel, E., Seluchi, M.E., Michelin, C.M., De Praga Baião,  
418 C.F., Chuchón ngulo, E., Almeida, E.K. and Kazmierczak, M.L. (2021). Extreme Drought in the Brazilian Pantanal in 2019–  
2020: Characterization, Causes, and Impacts. *Frontiers in Water*, 3, p.13.

420 Marengo, J. A., Fisch, G. F., Alves, L. M., Sousa, N. V., Fu, R., Zhuang, Y., et al. (2017). Meteorological context of the onset  
and end of the rainy season in Central Amazonia during the 2014-15 Go-Amazon Experiment. *Atmos. Chem. Phys.* 17, 7671–  
422 7681. doi: 10.5194/acp-2017-22

Moreira, D. S., Longo, K. M., Freitas, S. R., Yamasoe, M. A., Mercado, L. M., Rosário, N. E., Gloor, E., Viana, R. S. M.,  
424 Miller, J. B., Gatti, L. V., Wiedemann, K. T., Domingues, L. K. G., and Correia, C. C. S. (2017). Modeling the radiative effects  
of biomass burning aerosols on carbon fluxes in the Amazon region. *Atmos. Chem. Phys.* 17, 14785-14810, doi: 10.5194/acp-  
426 17-14785-2017.

Morissette, J. T., Giglio, L., Csaszar, I., Setzer, A., Schroeder, W., Morton, D., and Justice, C. O. (2005). Validation of MODIS  
428 Active Fire Detection Products Derived from Two Algorithms. *Earth Interactions* 9, 1-25. NYT, 2020. The World's Largest  
Tropical Wetland Has Become an Inferno. (Available at [https://www.nytimes.com/interactive/2020/10/13/climate/pantanal-  
430 brazil-fires.html](https://www.nytimes.com/interactive/2020/10/13/climate/pantanal-brazil-fires.html)). Accessed on: April 3rd 2021

NASA (2018). CERES Time-Interpolated TOA Fluxes, Clouds and Aerosols.  
432 [https://ceres.larc.nasa.gov/documents/DPC/DPC\\_current/pdfs/DPC\\_SSF1deg-Day\\_R5V1.pdf](https://ceres.larc.nasa.gov/documents/DPC/DPC_current/pdfs/DPC_SSF1deg-Day_R5V1.pdf)

Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, Saulo R; Cardozo, F. S., Kaiser, J. W., Wooster, M. J. (2016).  
434 Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment.  
*Atmospheric Chemistry and Physics (Online)*, v. 16, p. 6961-6975.

Pivello, V. R. (2011). The use of fire in the Cerrado and Amazonian Rainforests of Brazil: Past and Present. *Fire Ecology* 7(1),  
436 24-30. doi: 10.4996/fireecology.0701024.

Platnick, S., et al., 2017. MODIS Atmosphere L3 Daily Product. NASA MODIS Adaptive Processing System, Goddard Space  
438 Flight Center, USA: dx.doi.org/10.5067/MODIS/MYD08\_D3.061 [last accessed: Jun 29, 2022]

Pletsch Majs, Silva Junior Chl, Penha Tv, Körting Ts, Silva Mes, Pereira G, Anderson Lo & Aragão Leoc (2021). The 2020  
440 Brazilian Pantanal fires. *An Acad Bras Cienc* 93: e20210077. DOI 10.1590/0001-3765202120210077

Prins, E. and Menzel, W. (1992). Geostationary satellite detection of biomass burning in South America, *Int. J. Remote Sens.*,  
442 13, 2783–2799.

Reddington, C., Butt, E., Ridley, D. et al. Air quality and human health improvements from reductions in deforestation-related  
444 fire in Brazil. *Nature Geosci* 8, 768–771 (2015). <https://doi.org/10.1038/ngeo2535>

Rasch, P. J., N. M. Mahowald, and B. E. Eaton. (1997). Representations of transport, convection, and the hydrologic cycle in  
446 chemical transport models: Implications for the modeling of short-lived and soluble species. *J. Geophys. Res.*, 102, 28 127–  
448 28 138, <https://doi.org/10.1029/97JD02087>.

Reid, J.S., Hobbs, P.V., Ferek, R.J., Blake, D.R., Martins, J.V., Dunlap, M.R., Liousse, C. (1998). Physical, chemical, and  
450 optical properties of regional hazes dominated by smoke in Brazil. *J. Geophys. Res.* 103, 32059–32080.

Reid, J. S., T. F. Eck, S. A. Christopher, R. Koppmann, O. Dubovik, D. P. Eleuterio, B. N. Holben, E. A. Reid, and J. Zhang  
452 (2005). A review of biomass burning emissions part III: Intensive optical properties of biomass burning particles, *Atmos.*  
*Chem. Phys.*, 5, 827–849, doi:10.5194/acp-5-827-2005.

Rosário, N. E., M. A. Yamasoe, H. Brindley, T. F. Eck, and J. Schafer (2011), Downwelling solar irradiance in the biomass  
454 burning region of the southern Amazon: Dependence on aerosol intensive optical properties and role of water vapor, *J.*  
*Geophys. Res.*, 116, D18304, doi:10.1029/2011JD015956.

Sedano, F. and Randerson, J. T. (2014). Multi-scale influence of vapor pressure deficit on fire ignition and spread in boreal  
456 forest ecosystems, *Biogeosciences*, 11, 3739–3755, <https://doi.org/10.5194/bg-11-3739-2014>.

Sena, E.T., Artaxo, P. and Correia, A.L. (2013). Spatial variability of the direct radiative forcing of biomass burning aerosols  
460 and the effects of land use change in Amazonia. *Atmospheric Chemistry and Physics*, 13(3), pp.1261-1275.

462 Sena, E.T. and Artaxo, P. (2015). A novel methodology for large-scale daily assessment of the direct radiative forcing of  
463 smoke aerosols. *Atmospheric Chemistry and Physics*, 15(10), pp.5471-5483.

464 Schafer, J. S., Eck, T. F., Holben, B. N., Artaxo, P., Yamasoe, M. A. and Procopio, A. S. (2002). Observed reductions of total  
465 solar irradiance by biomass-burning aerosols in the Brazilian Amazon and Zambian Savanna. *Geophys. Res. Lett.* 29(17),  
1823, doi:10.1029/2001GL014309.

466 Thornhill, G. D., Ryder, C. L., Highwood, E. J., Shaffrey, L. C., and Johnson, B. T. (2018). The effect of South American  
467 biomass burning aerosol emissions on the regional climate, *Atmos. Chem. Phys.*, 18, 5321–5342, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-18-5321-2018)  
468 [18-5321-2018](https://doi.org/10.5194/acp-18-5321-2018).

469 Vale MM, Berenguer E, Menezes MA de, Castro EBV de, Siqueira LP de, Portela R de CQ. (2021). The COVID-19 pandemic  
470 as an opportunity to weaken environmental protection in Brazil. *Biological Conservation* 255 1-5. Available from:  
<https://doi.org/10.1016/j.biocon.2021.108994>


472 Yamasoe, M. A.; Artaxo, P.; Miguel, A. H.; Allen, A. G. (2000). Chemical composition of aerosol particles from direct  
473 emissions of biomass burning in the Amazon Basin: water-soluble species and trace elements. *Atmospheric Environment*, 34:  
474 1.641-1.653.

475 Yamasoe, M. A., von Randow, C., Manzi, A. O., Schafer, J. S., Eck, T. F., and Holben, B. N.: Effect of smoke and clouds on  
476 the transmissivity of photosynthetically active radiation inside the canopy, *Atmos. Chem. Phys.*, 6, 1645–1656,  
<https://doi.org/10.5194/acp-6-1645-2006>, 2006.

478 WWF, 2020. Fires in the Pantanal grow more than 200% and break a record, (available at  
<https://www.wwf.org.br/informacoes/english/?76914/Fires-in-the-Pantanal-grow-more-than-200-and-break-a-record>).

480 Accessed on: April 3rd 2021

482




484

486

488

Figure 1: Spatial distribution of the Brazilian biomes of Amazon tropical forest, Caatinga, Cerrado, Pantanal, Mata Atlântica and Pampa. Locations of the AERONET stations considered in this study are also depicted. The area confined by the dashed red box represents a transect defined to study north-south smoke variability and transport taking Pantanal west and east borders as reference.




490

Figure 2: Monthly (from July to October) interannual evolution of fire counts (black), aerosol optical depth at 550 nm (grey) for the years from 2003 to 2020 for (a) Pantanal, (b) Amazonia, and (c) Cerrado biomes.

**Comentado [NR25]:** All the figures in this new version were adjusted to use colour schemes accessible to persons with colour vision deficiencies.





492

494 **Figure 3: Hovmöller diagram of mean AOD at 550 nm from Amazon Forest latitudes to Pantanal downwind regions considering the**  
 496 **average between Pantanal borders at west (58.5°W) and east (54.96°W) longitudes (see Figure 1). The magenta line represents the**  
**north and southern limits of Pantanal. Each year's label is defined at its beginning, January first, that is also the case of the months**  
**of July and October, which are used to highlight the biomass burning season period.**


498

500

502

504

506



508

510 **Figure 4:** Multiyear analysis (2003-2020) of (a) monthly mean aerosol optical Depth at 550 nm (AOD@550nm) as function of fire  
 512 count for Pantanal biome; b) monthly mean aerosol optical Depth at 550 nm (AOD@550nm) over Pantanal as function of monthly  
 mean AOD@550nm over Amazonia. August, September and October of the year 2020 are red-highlighted.

514


516

518

520

522

**Comentado [NR26]:** As explained before, Figure 4 from previous version was splitted in two different figures in this new version (Figure 4 and Figure 5). This was done in order to assure the best contextualization and quality for the plots. The texts in which they are discussed were not modified, we only adjusted the references of the next figures so the consistency of the discussion could be kept




524 **Figure 5:** a) Fire count distribution over Pantanal for August, September and October of 2020 using as background the Enhanced  
 526 Vegetation Index (EVI); b) Monthly (August, September and October) distribution of fire count over Pantanal and over conservation  
 and indigenous areas located within the biome.

528


530

532




534

536 **Figure 6:** Multiyear box plots of the monthly single scattering albedo at 440 nm (SSA@440nm) from 2003 to 2020 for four  
 538 AERONET sites: Alta Floresta, Cuiabá, Rio Branco and Santa Cruz. In each box, the black line in the center represents the median,  
 and the lower and upper limits are the first and the third quartiles, respectively. The vertical lines extending from the box represent  
 the spread of instantaneous SSA@440 nm with the length being 1.5 times the interquartile range.




540

542 **Figure 7:** Aerosol direct instantaneous radiative forcing as function of AOD at 550 nm at AERONET sites highlighting 2020 (black) against historical values (2003 – 2019).



544

546 **Figure 8:** Spatial distribution of the mean aerosol optical depth at 550 nm (AOD<sub>550nm</sub>) and wind pattern at 850 hPa for September  
 548 (a) and October (right) of different years. The area limited by the black solid line represents the regions with AOD<sub>550nm</sub> higher than  
 0.5, and its dimension (Area) and mean AOD<sub>550nm</sub> are also presented. The biomes' borders are represented by the colors blue (Amazon Forest), red (Cerrado ecosystem) and green (Pantanal).



550 **Figure 9:** Spatial distribution of the mean surface solar irradiance under cloudless conditions and in the presence of aerosol for the  
 552 months of September and October of different years. The biome's borders are represented by the colors blue (Amazon Forest), red  
 (Cerrado ecosystem) and magenta (Pantanal).

**Table 1: Set of variables used in this study, their respective applications, sources and references.**

Variables	Application	Data sources	Reference
Aerosol Optical Depth at 500 nm (AERONET) and at 550 nm (MODIS – Aqua)	Smoke plume loading and dimension	AERONET (Level, 1.5) <a href="https://aeronet.gsfc.nasa.gov/">https://aeronet.gsfc.nasa.gov/</a> MODIS Atmosphere Level 3 (L3) gridded products (Daily and Monthly) from the Aqua platform (MYD08D 3/M 3) <a href="https://giovanni.gsfc.nasa.gov/giovanni/">https://giovanni.gsfc.nasa.gov/giovanni/</a> Accessed on Feb16, 2021.	Holben et al. (1998); King et al., 2013; Levy et al., 2013; Platnick et al., 2017
Aerosol Single Scattering Albedo	Smoke plume absorption efficiency	AERONET (Level 1.5) <a href="https://aeronet.gsfc.nasa.gov/">https://aeronet.gsfc.nasa.gov/</a>	Dubovik and King (2000)
Aerosol radiative forcing	Radiative impact	AERONET (Level 1.5) <a href="https://aeronet.gsfc.nasa.gov/">https://aeronet.gsfc.nasa.gov/</a>	Garcia et al. (2012)
Wind components at 850 hPa	Circulation pattern	MERRA 2- Reanalysis atmospheric variables. <a href="https://disc.gsfc.nasa.gov/datasets/M2TMNXSLV_5.12.4">https://disc.gsfc.nasa.gov/datasets/M2TMNXSLV_5.12.4</a> Accessed on November 11, 2021.	Gelaro, et al. (2017); GMAO (2015)
Fire counts	Fire activity	<a href="http://queimadas.dgi.inpe.br/queimadas/portal/static/estatisticases_tados/">http://queimadas.dgi.inpe.br/queimadas/portal/static/estatisticases_tados/</a> – Accessed on December23, 2020.	Morisette et al. (2005)
Enhanced Vegetation Index (EVI)	Land use	<a href="https://doi.org/10.5067/MODIS/MOD13C2.061">https://doi.org/10.5067/MODIS/MOD13C2.061</a> Accessed on May, 20, 2022	Didan (2021)

554

556


558

560

562

564

### Supplement



566 **Figure S.1:** Scatterplot of level 2.0 versus level 1.5 for AOD at 500 nm, SSA at 440 nm and radiative forcing at the surface from  
AERONET. Red line indicates the best line fit and the coefficients of the linear regression are also presented.

568