

# **The Effect of Forced Change and Unforced Variability on Heat Waves, Temperature Extremes, and Associated Population Risk in a CO<sub>2</sub>- Warmed World**

**Jangho Lee, Jeffrey C. Mast, and Andrew E. Dessler**

Department of Atmospheric Sciences, Texas A&M University, College Station, TX, USA

Corresponding author: Andrew Dessler ([adessler@tamu.edu](mailto:adessler@tamu.edu))

## **Key Points**

- Unforced variability of the climate system, primarily ENSO, plays a key role in the occurrence of extreme events in a warming world.
- Uncertainty of unforced variability becomes smaller as one looks at larger regions or at a global perspective.
- Increases of heat wave indices are significant between 1.5°C and 2.0°C of warming and the risk of facing extreme heat events is higher in low GDP regions.

**Abstract**

This study investigates the impact of global warming on heat and humidity extremes by analyzing 6-hourly output from 28 members of the Max Planck Institute Grand Ensemble driven by forcing from a 1%/year CO<sub>2</sub> increase. We find that unforced variability drives large changes in regional exposure to extremes in different ensemble members, and these variations are mostly associated with ENSO variability. However, while the unforced variability of the climate can alter the occurrence of extremes regionally, variability within the ensemble decreases significantly as one looks at larger regions or at a global population perspective. This means that, for metrics of extreme heat and humidity analyzed here, forced variability of the climate is more important than the unforced variability at global scales. Lastly, we found that most heat wave metrics will increase significantly between 1.5°C and 2.0°C, and that low GDP regions shows significant higher risks of facing extreme heat events compared to high GDP regions. Considering the limited economic adaptability of population to heat extremes, this reinforces the idea that the most severe impacts of climate change may fall mostly on those least capable to adapt.

## 1. Introduction

The long-term goal of the 2015 Paris agreement is to keep the increase in global temperature well below 2°C above pre-industrial levels, while pursuing efforts and limit the warming to 1.5°C. Given that no one lives in the global average, however, understanding how these global average thresholds translate into regional occurrences of extreme heat and humidity is of great value (Harrington et al., 2018). Previous studies have reported that regional extreme heat events will not only be more frequent, but also more extreme in a warmer world. This was discussed in various assessment and reports such as US National Climate assessment and those by IPCC (Melillo et al., 2014; Wuebbles et al., 2017; Hoegh-Guldberg et al., 2018; Masson-Delmotte et al., 2018) and it is expected to have significant impacts on human society and health. More importantly, previous studies have analyzed the risk (Quinn et al., 2014; Sun et al., 2014; Lundgren et al., 2013), exposure (Dahl et al., 2019; Ruddell et al., 2009; Liu et al., 2017; Luber and McGeehin, 2008), vulnerability (Chow et al., 2012; Wilhelmi and Hayden, 2010) and susceptibility (Arbuthnott et al., 2016) of population in the current and warmer climates.

Many criteria and indices have been used to assess extreme heat, such as the absolute increase of maximum temperature from the reference period (Wobus et al., 2018), risk ratio of population's exposure to heat (Kharin et al., 2018), and heat wave magnitude index (Russo et al., 2017). In this study, we utilize four locally defined heat wave indices from Fischer and Schär (2010) and Perkins et al. (2012) of duration, frequency, amplitude, and mean. We also focus on consecutive-day extremes, which are known to cause more harm than single-day events (Baldwin et al., 2019; Simolo et al., 2011; Tan et al., 2010). In addition, because the combined effect of temperature and humidity is known to affect human health by reducing the body's

ability to cool itself through perspiration, wet-bulb temperature is frequently analyzed (Kang and Eltahir, 2018) and we will do so here.

Climate extremes are always a combination of long-term forced climate change acting in concert with unforced variability (Deser et al., 2012). Thus, characterizing and quantifying both long-term change due to external forcing and the unforced variability of the climate system is crucial in assessing the future risk of extreme events. There have been numerous studies that link dominant modes of unforced variability to extreme events. Temperature connections with El Niño Southern Oscillation (ENSO) (Thirumalai et al., 2017; Meehl et al., 2007), the Pacific Decadal Oscillation (PDO) (Birk et al., 2010), the Atlantic Multidecadal Oscillation (AMO) (Zhang et al., 2020; Mann et al., 2021) have been investigated in the previous studies. The effect of climate extremes on different populations depends on numerous factors, including the level of economic development, with impacts of heat extremes being more severe in less economically developed countries (Diffenbaugh and Burke, 2019; Harrington et al., 2016; King and Harrington, 2018). For example, as temperatures go up, increased energy demand to cool buildings will be required (Parkes et al., 2019; Sivak, 2009) in metropolitan area. But this requires resources to both install air conditioning and then operate it. The greater impacts of extreme heat in economically less developed region in a warmer climate has been discussed in multiple studies (Marcotullio et al., 2021; Russo et al., 2019).

In this paper, a single-model initial-condition ensemble of 28 simulations of a global climate model (GCM) are used to quantify heat and humidity extremes in a warmer world. We use population data to look at population risk for mortality events in daytime (Mora et al., 2017) and nighttime (Chen and Lu, 2014). We also utilize per capita gross domestic product (GDP per capita) data to investigate how climate change impacts extreme heat events on different levels of

economic status. To quantify the impact on energy demand, we also quantify changes in cooling degree days and warming degree days.

## **2. Data**

### *2.1. MPI-GE ensembles*

Simulation data in this study come from an ensemble of runs of the Max-Planck Institute Earth System Model collectively known as the MPI Grand Ensemble (MPI-GE) project (Maher et al., 2019). Each of the 28 ensemble members branches from different points of a 2000-year pre-industrial control run and are integrated for 150 years, forced by CO<sub>2</sub> concentration increasing at 1% per year (hereafter, 1% runs). Because the radiative forcing scales as the log of CO<sub>2</sub> concentration, the 1% runs feature radiative forcing that increases approximately linearly in time. We analyze 6-hourly output with  $1.875^{\circ} \times 1.875^{\circ}$  spatial resolution, which is the original resolution of the model output, for land areas between 60°N and 60°S. Our analysis will focus on 2-meter temperature (hereafter, t2m) and 2-meter dew point temperature (d2m), from which 2-meter relative humidity (rh) and wet-bulb temperature (w2m) are calculated using the methods of Davies-Jones (2008).

Unforced variability in the climate system generates uncertainties in the projection of the climate by impacting the dynamic component of the climate, especially for extreme events (Kay et al., 2015; Thompson et al., 2015). One way to analyze the impact of unforced variability in climate system is to use an initial-condition ensemble. Each members of initial-condition ensemble are generated by perturbing the initial conditions of single climate model. This perturbation will then propagate to generate different sequence of climate, such as ENSO, PDO,

etc. (Deser et al., 2012; Kay et al., 2015). In this paper, we use the ensemble to allow us to estimate the impact of unforced variability on temperature extremes.

Since the model used only considers CO<sub>2</sub> forcing without aerosols, and it represents a continuously warming climate, one might question if the model simulation accurately represents the real climate. To judge the fidelity of the simulations, we compare 15 years (2003-2017) of ERA-Interim reanalysis data (Dee et al., 2011) from the European Centre for Medium Range forecast (ECMWF) with 15 years of the MPI-GE 1% ensemble which have the same ensemble- and global-average temperatures (years 39-53); in the rest of the paper, we will refer to these as the reference periods. In both data sets, we then calculate 90<sup>th</sup> percentile and mean t2m and w2m for each grid points. This calculation was done for each member of the model ensemble. For each of the 4 values (90<sup>th</sup> percentile t2m/w2m and mean t2m/w2m), we determine if the values from the reanalysis fall into the spread of 28 ensemble members of the 1% runs. For each grid point, if the reanalysis value falls within the ensemble spread, we mask out the grid point; if not, we plot how far the reanalysis value is from the closest member of the 1% ensemble (Figure 1).

Generally, the 1% runs overpredicts t2m and w2m in Northern hemisphere, and underpredicts in Southern hemisphere, except for India. This difference is consistent with the fact that the 1% models do not contain aerosol forcing, which should lead to biases of the sign seen in Fig. 1. The w2m shows larger area of differences than t2m, which suggests that there are larger biases in the dew point, which is needed in the calculation (Davies-Jones, 2008). The area-weighted averages of these differences are -0.08°C, -0.03°C, -0.04°C, and -0.11°C globally for 90<sup>th</sup> percentile t2m, mean t2m, 90<sup>th</sup> percentile w2m, and mean w2m respectively, which means that the model is, on average, underpredicting land temperature. Breaking down to Northern and Southern hemisphere, the bias is 0.20°C, 0.21°C, 0.15°C, 0.14°C in NH and -0.64°C, -0.54°C, -

0.36°C, and -0.44°C, confirming that the model is overpredicting temperature in NH land and underpredicting in SH land.

To quantify the impact of the biases in Fig. 1 on the occurrence of heat extremes, we will perform sensitivity tests on the calculations by adding to each grid point of each member of the ensemble the average differences between the ensemble average t2m and w2m and the reanalysis. By evaluating how much our results change, we come up with an estimate of the impact of model biases on our results. As we will show later, these biases have little impact on the results of the paper.

## *2.2. Global population and GDP per capita data*

Global population data from the NASA Socioeconomic Data and Applications Center (SEDAC, 2018) are used to weight the heat wave indices by population. The data represent the population in year 2015 at 30'' × 30'' spatial resolution, and we re-gridded to the 1.875° × 1.875° grid of the MPI model by summing the values in grid boxes surrounding the MPI grid centers. In our population-weighted calculations, we assume that the relative distribution of population remains fixed into the future.

Gridded GDP per capita data (Kummu, 2019) over 1990-2015 are used to estimate the risk of heat extreme events for different levels of wealth. These data are re-gridded from the original 5'' × 5'' spatial resolution to the MPI model's resolution of 1.875° × 1.875° by averaging the GDP inside the grid box. When doing this average, per capita GDP was weighted by population and also averaged over the 1990-2015 period. We assume that the relative percentile of GDP per capita for each grid point is fixed into the future, so changes in climate risk are due to exposure to warmer climate extremes, not changes in relative per capita wealth.

153

154 **3. Method of analysis**155 *3.1. Global warming*

156 Global warming is defined as the global and annual average temperature increase  
 157 compared to the average of first 5 years of the 1% run. We find that ensemble- and global-  
 158 average t2m reaches 1.5°C, 2°C, 3°C and 4°C occur in years 59, 76, 108, and 133 years,  
 159 respectively, and reaches 4.6°C at the end of the 150-year run. The increase of global average  
 160 temperature is nearly linear for both t2m and w2m, consistent with a linear ramping of the  
 161 forcing.

162 The focus on the paper will be on heat extremes at 1.5°C, 2°C and 3°C. The 1.5°C and  
 163 2°C thresholds are the limits described in the Paris Agreement, while 3°C is the warming we are  
 164 presently on track for (Hausfather and Peters, 2020).

165

166 *3.2. Heat wave indices*

167 Identification of heat waves is done in several steps. First, for each grid point, we smooth  
 168 a daily maximum temperature (determined from 6-hourly temperatures) using a 15-day moving  
 169 window for the first 5 years of 1% runs, which is the period before significant warming has  
 170 occurred. Then, the 90<sup>th</sup> percentile of smoothed daily maximum temperature for the first 5 years  
 171 was calculated at each grid point (Fischer and Schär, 2010). This value is used as a threshold for  
 172 the heat waves at that grid point. Then we calculate the heat wave days, defined as days that  
 173 exceed the threshold for three or more consecutive days (Baldwin et al., 2019).

174 We then define four indices to represent the characteristics of these heat waves. To  
 175 determine the occurrence of events, heat wave duration (HWD; longest heat wave of the year)



and heat wave frequency (HWF; total number of heat wave days in a year) are calculated. From an intensity perspective, heat wave amplitude (HWA; maximum temperature during heat wave days during a year) and heat wave mean (HWM; mean temperature during heat wave days in a year) are selected. These indices are also calculated in an analogous fashion for wet bulb temperature (w2m), since wet-bulb temperature is arguably more relevant for human health (Heo et al., 2019; Morris et al., 2019; Buzan and Huber, 2020). These indices are summarized in Table 1.

### *3.3. Deadly days and tropical nights*

Heat wave thresholds are different for each grid point because they are based on pre-industrial temperatures at that grid point. Combined with regional differences in the ability to adapt, this means that heat waves in different regions may have different implications for human society. We therefore also count the number of days each year with daily maximum w2m above 26°C, which we refer to as “deadly days”. We note that other values could be chosen, with higher values occurring less frequently but having more significant impacts. This value is based on the analysis of Mora et al. (2017), who demonstrated that w2m of about 24°C is the threshold which fatalities from heat-related illness occur. However, since we find that there are some regions that already experience over 9 months of 24°C w2m events per year, we increase this threshold to 26°C in our analysis. We could have chosen higher w2m values, but any choice in this range is associated with negative impacts, so we have chosen a value near the bottom of the range where mortality occurs in order to maximize the signal in the model runs.

A warm nighttime minimum temperature can be as important as a high maximum temperature for human health and mortality (Argaud et al., 2007; Patz et al., 2005), so we define “tropical nights” as a daily minimum  $t_{2m}$  over  $25^{\circ}\text{C}$  (Lelieveld et al., 2012).

### *3.4. Cooling degree days and heating degree days*

To assess the economic and energy impact of heat extremes, cooling degree days (CDD) and heating degree days (HDD) are calculated. CDD and HDD are metrics of the energy demand to cool and heat buildings. For each grid point, annual CDD is calculated by subtracting  $18^{\circ}\text{C}$  from the daily average temperature and summing only the positive values over the year. HDD is the absolute value of the sum of the negative values. Previous studies reported that CDD and HDD are closely related to energy consumption (Sailor and Muñoz, 1997).

## **4. Results**

### *4.1. Impact of unforced variability of climate on regional heat extremes*

To investigate the impact of unforced variability on more regional heat extremes, we take the 15 largest cities by population (Fig. 2a) and determine the number of deadly days and tropical nights over time by averaging the  $3\times 3$  grid points surrounding the city, only including the land grid points. Figure 2b-d depicts the ensemble averaged number of deadly days and tropical nights, as well as the spread between the ensemble members. The error bars in Figure b-d show the highest and lowest values of the extremes.

This difference within the ensemble is the result of unforced variability. For all 15 cities, average spread in the number of deadly days at  $1.5^{\circ}\text{C}$ ,  $2.0^{\circ}\text{C}$ ,  $3.0^{\circ}\text{C}$ , and  $4.0^{\circ}\text{C}$  of global warming between the ensemble members with maximum and minimum numbers are 14.3, 15.1,

20.6, and 21.9 days per year. For tropical nights, the spreads are 29.3, 27.7, 29.1, and 26.7 days per year. So, on average, unforced variability can change the number of extreme days and nights by a few weeks per year. There is no significant variance of ensemble spread between the cities except for cities with very low ensemble-averaged values (e.g., Mexico City at 1.5°C warming) or very high values (e.g., tropical nights in Manila at 4.0°C warming). However, for the cities that do not see large increase in extreme temperatures (e.g., New York City), this represents a very large fraction of the predicted change of extremes, while for cities that experience much larger increase (e.g., Manila), it represents a smaller percentage.

As discussed in Section 2.1, we examine the sensitivity of our results to potential biases of the model by recalculating the deadly days and tropical nights using model data after adding in the bias estimated by comparison to the reanalysis. The average difference of deadly days in the sensitivity test (absolute difference) at 1.5°C, 2.0°C, 3.0°C, and 4.0°C warming is 2.1, 2.5, 5.5, and 7.6 days per year when averaged over 15 cities. The standard deviation of difference calculated between the cities is 2.5, 3.4, 6.7, and 9.7 days at each level of warming. For tropical nights, sensitivity test produced differences of 3.6, 3.6, 5.3, and 3.5 days per year at each level of warming, with standard deviations within the ensemble of 3.6, 4.9, 6.9, and 1.8 days. Thus, model biases are unlikely to have a large impact on our results.

Previous work has attempted to distinguish the origin and mechanisms of unforced variability of temperature and temperature extremes (Meehl et al., 2007; Zhang et al., 2020; Birk et al., 2010). To probe the statistical modes of variability affecting this ensemble spread and to identify the underlying physical mechanisms, empirical orthogonal function (EOF) analysis (North, 1984) was performed on the detrended and normalized time series of deadly days and tropical nights for the 15 cities. For each city, the 28 ensemble members are concatenated

together (total of  $28 \times 150$  years) in order for all ensemble to share the same EOF. In this way, we aim to find the dominant drivers of unforced variability that impacts heat extremes in the largest cities around the world.

The first three EOF patterns for each city are plotted in Fig. 3 as bars. The first EOF mode of deadly days shows large values for Delhi, Shanghai, Dhaka, and Karachi, while cities in other regions show lower values. The second and third EOFs for deadly days show more variability between the cities. The first EOF for tropical nights (Fig. 3d) show large positive values for cities in the India-Pakistan region, with other cities showing smaller magnitude changes. The second EOF shows large negative values in Cairo, Istanbul, and Manila, while the third EOF for tropical nights shows more variability between the cities.

The PC time series are projected onto detrended annual sea surface temperature (SST) anomalies. This allows us to investigate how heat extreme events in 15 major cities are associated with global modes of unforced variability. Maps of correlation coefficients are also plotted in Fig. 3. Characteristic patterns for ENSO (Trenberth, 2020), PDO (Deser and Trenberth, 2016), and AMO (Trenberth and Zhang, 2021) are calculated for each ensemble using all 150-year of SSTs, and the pattern is averaged over ensembles to come up with a single ENSO, PDO, and AMO SST pattern for the ensemble. Then, those patterns are compared with the PC projection on SST to see how PC projected SST resembles the patterns of unforced variability. Correlation coefficients between the standard climate indices and PC projected SST is shown on lower panel of Fig. 3 as numbers. All of the projections of deadly day PCs and projections of the first two modes of tropical nights shows patterns similar to El Niño-Southern Oscillation (ENSO) and Pacific Decadal Oscillation (PDO).

Power spectra of the PCs are calculated individually for each ensemble member, and then the ensemble average is plotted Figure 4. Overall, the spectra of the deadly day PCs look very much like the spectrum for ENSO, and it notably does not have the ~20-year peak of the PDO spectrum. This tells us that, in this model at least, the variability in the occurrence of deadly days in these large cities is strongly regulated by ENSO. This may be a consequence of the fact that these large cities are mostly located near ocean and at lower latitudes. The third deadly day PC has lower correlations with ENSO or PDO index, so it is harder to draw firm conclusions about the mechanism behind it. Also, higher modes of EOFs are unlikely to refer to a single mode of climate due to the orthogonality constraints between each mode. The tropical night PCs also show peaks at ENSO periods (Fig. 4b) suggesting that, like deadly days, tropical night variability is controlled by ENSO.

#### *4.2. Cluster analysis and population risk of heat wave indices*

We calculate HWD, HWF, HWA, and HWM for both t2m and w2m each year at each grid point, which generates eight different 150-year time series for each of the 28 ensemble members. Each time series at each grid point is regressed vs. time, yielding a slope and the intercept for each time series in all 28 ensemble members. The 16 variables (8 [heat wave indices]  $\times$  2 [slope, intercept]) are then utilized as a predictor variable for K-means clustering (Likas et al., 2003) to categorize the spatial variation of heat waves using the Euclidean distance of its predictor variables (16 variables). With slope and intercept, we can characterize the heat indices of each grid point with response to CO<sub>2</sub> forcing (slope) and climatology (intercept). The number of clusters in this study is set to 6, using the elbow method (Syakur et al., 2018). When using 5 clusters, we find that two clusters (the light and dark blue regions in Figure 5a) merge,

and when using 7 clusters, we find that one cluster (the dark blue region in Figure 5a) divides into two separate clusters.

Figure 5a shows the cluster value that most ensembles assigned to each grid point and it shows distinct geographical characteristics, as summarized in Table 2 (the result of clustering shows little difference between individual ensemble members). As might be expected from how we calculated the 16 variables for clustering, each cluster shows a different evolution of heat extremes in warmer world (Figure 6). Although the warming signal is largest in the polar regions (Figure 5b), the largest increases of HWD and HWF are found at lower latitudes (in cluster 1 and 2 on Figure 6a-d). This is mostly due to low variability in these regions compared to polar regions, making it easier for a trend to exceed the heatwave threshold.

These results are insensitive to potential model biases. Sensitivity tests show that adding the bias to the model changes HWD, HWF, deadly days, and tropical nights, by less than 5% for all metric and clusters. For HWA and HWM, the difference caused by adding the bias was less than 1°C for all metric and clusters, suggesting that the impact of model biases is small in this analysis.

For HWA and HWM, the rate of increase is similar for all clusters, with increases of  $HWA_{t2m}$  and  $HWA_{w2m}$  of 1.45°C per degree of global average warming and 0.85°C per degree of global average warming, respectively, and  $HWM_{t2m}$  and  $HWM_{w2m}$  of 0.66°C per degree of global average warming and 0.47°C per degree of global average warming, respectively (Figure 6e-h). The exception is  $HWA_{t2m}$  in cluster 6. The large increase of  $HWA_{t2m}$  in this region is connected to the strong global warming signal in high latitudes that has been predicted for decades and now observed (Stouffer and Manabe, 2017).

Turning to deadly days (Fig. 6i), we find a substantial increase occurs in cluster 1 after 2.0°C of warming; this is important because it gives additional support for the Paris Agreement’s aspirational goal of limiting global warming to 2.0°C. Almost all increases in deadly days are in low latitudes (cluster 1, 2, and 3). For tropical nights, low latitudes and deserts (cluster 4) contribute most of the increase. Figure 6 also shows the spread in within the ensemble for each metric and cluster. We find that the spread for a cluster is generally small compared to the change over time as well as the difference between the clusters.

We also generated indices weighted by global population. Heat wave indices for the 95<sup>th</sup> percentile of population (meaning 5% of the population is exposed to higher values), 90<sup>th</sup> percentile of population, and median of the population are depicted in Figure 7. Figure 7a shows that with 3°C of warming, 5% of the Earth’s population will experience heat waves lasting 122 days (standard deviation between ensemble members:  $1\sigma = 17$  days), 10% of the population will experience heat waves of 94 days ( $1\sigma = 7$  days), and half of the population will experience heat waves around 50 days ( $1\sigma = 4$  days). These are large increases over present-day values of 50, 42, and 21 days. The average of the standard deviation between the ensemble members (calculated every year and then averaged), are 10.6, 6.2 and 3.7 days for the 95<sup>th</sup>, 90<sup>th</sup> percentile and median, respectively. This is significantly smaller than values from the analyses of cities in Figure 2, where the unforced variability makes larger differences in the occurrence of heat waves.

The rate of increase of  $\text{HWF}_{\text{w2m}}$  in Fig. 7d shows a rapid increase until global average warming reaches about 2.5°C. Given that the planet has already warmed about 1°C above pre-industrial, this suggests that the world should presently be experiencing a rapid increase of wet-bulb extreme frequency, particularly in the tropics. This is related to the increased slope in

Figure 6, in which cluster 1 and 2's values of  $HWD_{w2m}$  and  $HWF_{w2m}$  increase rapidly until 3.0°C and 2.0°C of global warming. At warmer temperatures,  $HWD_{w2m}$  and  $HWF_{w2m}$  reach a plateau, since values over 300 days per year means there is little room for additional increase. For  $HWA_{t2m/w2m}$  and  $HWM_{t2m/w2m}$ , the increase is mostly linear. Also note that, at 3°C of global warming, the 90<sup>th</sup> percentile of population weighted  $HWA_{w2m}$  reaches over 29°C, which while not immediately fatal to humans may nevertheless indicate great difficulty for even a developed society to adapt to.

Currently, 10% of the total population faces more than 45 deadly days and 181 tropical nights per year. This grows to 65 and 195 days, respectively, at 1.5°C warming. With 2°C of global warming, 10% of the population will face about 3 months of deadly days and 7 months of tropical nights every year, and this increase to 4 months and 8 months in 3°C of warming. Also, with 3°C of global warming, 5% of the population will be in an environment where 8 months and 10 months in a year is a deadly days and tropical night. Our sensitivity tests suggest that model bias generates less than 5% differences for  $HWD$ ,  $HWF$ , deadly days, and tropical nights for all metrics and percentile of population at every level of global warming, except when the metrics are near-zero. Potential model biases also generate small differences in  $HWA$  and  $HWM$ , with less than 1°C difference in all metrics for every period. Furthermore, with 3°C of global warming, the minimum ensemble member of deadly days is above the maximum ensemble of the present-day reference (0.87°C) for all population percentiles (5%, 10%, and 50%). This occurs at 2°C for tropical nights. Details of ensemble spread are also shown in Table 3.

It is notable that, although there is a large spread between the ensemble members in each city (Figure 2), the spread in the clusters (Figure 6) and population-weighted metrics (Figure 7) is not as large. This emphasizes that the effect of unforced variability might be large at small



scales but, as the region expands, the impact of unforced variability decreases. This is also found in Table 3, where in each case, the standard deviation between ensembles is less than 20% of the average, except in a few cases. This indicates that unforced variability will generally play a minor role in determining global exposure to temperature above thresholds, although different people may be affected in different realizations of unforced variability.

In addition, with 1.5°C of global warming, the lowest ensemble of the 90<sup>th</sup> percentile of HWD<sub>t2m</sub>, HWD<sub>w2m</sub>, and HWF<sub>t2m</sub> exceeds the highest ensemble of the same metric in the current climate (red lines in Figure 7). With 2°C of warming, the minimum ensemble of HWD<sub>t2m/w2m</sub>, HWF<sub>t2m/w2m</sub>, HWM<sub>w2m</sub>, and tropical nights exceed the maximum ensemble of the current climate, and with 2.5°C of warming, the minimum ensemble of all metrics exceeds the maximum ensemble of the same metric in the current climate. Thus, this model predicts that the occurrence of extremes will soon be able to exceed values likely possible in our present climate for these metrics.

#### 4.3. Analysis on GDP per capita

It is well-known that not everyone is equally vulnerable to extreme weather, with rich, relatively more developed communities having more resources to deal with extreme events than poorer communities. In that context, global gridded GDP per capita is used to calculate average risk at each level of wealth. The ensemble-average result is depicted in Figure 8, which shows the absolute number of deadly days and tropical nights as well as the increase in number of deadly days and tropical nights that each level of economic level experience relative to the reference period warming of 0.87°C. This plot assumes that the relative distribution of population and GDP remains fixed through time. Our sensitivity tests show that the model bias

yields small differences in the results, with less than 5% difference in both the absolute number of extreme events as well as the changes in extremes.

For each level of warming, we find that the lower GDP regions will experience not only higher absolute numbers of extreme temperature days but also the largest increases. For deadly days, the increase is largest between 10<sup>th</sup> to 40<sup>th</sup> percentile of GDP, and for tropical nights, the increase is largest below the 30<sup>th</sup> percentile of GDP. The regions that contribute the most for the low GDP percentiles are Southeast Asia, including Myanmar, Laos, and Cambodia, and Tropical Africa, including Republic of the Congo, Kenya, Uganda, Ethiopia, and Sudan, which are in clusters 1 and 2 in our cluster analysis (Figure 5). The maximum difference of heat wave days between the ensembles is less than 25% for all GDP and global warming levels.

#### *4.4. Energy demand on large cities*

Annual CDD and HDD have been calculated for the 15 cities in section 4.1. Both CDD and HDD are calculated by averaging the CDD and HDD values of 3×3 grid points surrounding each city, including only land grid points. CDD and HDD values are then averaged for 5 years after global warming reaches each levels of threshold. Fig. 9 shows the percent change of CDD and HDD at 1.5°C, 2.0°C, 3.0°C, and 4.0°C relative to the reference period CDD and HDD values. This was done for each city, and for each ensemble member. At 1.5°C, 2.0°C, 3.0°C, and 4.0°C warming, CDDs in the 15 cities increase by an average of 9%, 22%, 54% and 70%. Our sensitivity tests show that the application of the average model bias yields changes of less than 1% in these numbers. This suggests an enormous increase in energy required for cooling.

In contrast, average energy demand on cold days (HDD) decreases by 21%, 36%, 59%, and 65% in cities considered, compared to present day, partially offsetting the increase in energy

required for cooling. Manila shows 0% change in HDD for all period, since Manila does not experience HDD days in present or future periods. Sensitivity tests also show less than a 1% difference in HDD change due to model biases.

## 5. Conclusion

In this study, we found that extreme heat events will become more frequent and severe in a warming world. We find that both forced and unforced variability play a key role in extreme heat events, highlighting the necessity of considering both contributions to extreme heat. We also look at population weighted, and GDP sorted statistics of extreme heat in warmer world.

Our results show that ENSO is the dominant mode of unforced variability impacting the occurrence of extreme heat and humidity events and that events tend to be synchronous in the world's largest 15 cities. But while the impact of unforced variability might be significant regionally and temporarily, it becomes less important when one looks at larger aggregate regions.

Looking at global population-weighted statistics, we found that with 1.5°C of global average warming, over 10% of population will face heat waves of 45°C temperature, and 28°C wet bulb temperatures. And 5% of the population will face more than 105 days of deadly days and 232 tropical nights per year. With 3°C of warming, which we are currently on track for, 10% of the population will experience over 132 days of deadly days and over 232 days of tropical nights per year. And 10% of population will face 47°C temperature and 30°C wet bulb temperature. Given these two metrics have important implications for human mortality, such increases may have significant impact on human health globally.

Sorting heat and humidity events by wealth, we confirm that increasing frequency and severity of extreme events will fall mostly on the poorer people. To further investigate some economic impacts of increasing heat extremes, cooling degree days (CDD) and heating degree days (HDD) are calculated for the world's 15 largest cities. Energy demand for cooling (CDD) increases by average of 9% on 1.5°C and 54% on 3.0°C of warming, while energy demand for heating (HDD) decreases by 21% and 59%. Since CDD is known to have a piecewise linear relationship with the energy consumption, with slope increasing with higher CDD (De Rosa et al., 2014; Shin and Do, 2016), increasing CDD in a warmer world could be one of the factors driving increased economic inequity from global warming related heat extremes, due to relative high cost and need for energy in poorest countries.

Uncertainties in this analysis include our use of gridded 6-hourly climate model output. More detailed analysis could be done with climate simulations with higher temporal and spatial resolution. The model has biases relative to measurements, potentially due to the fact that there are no aerosols in the forcing, which is another source of uncertainty. This was tested by adding the difference between the ensemble average and the reanalysis data to the model fields and recomputing the heat wave indices. In general, the impact of this bias was not important. In future analyses, this could be better resolved with use of multi-model ensembles or detailed bias-correction of the model.

Another uncertainty is that our runs are continuously warming, and it is possible that an equilibrium world at any given temperature may experience different occurrence of extremes than in the runs in this paper. Additionally, since an increasing proportion of the population is expected to live in dense metropolitan areas, there is also the possibility that actual heat and humidity extremes that populations experience could be more severe than the gridded data due to

local phenomena such as the urban heat island effect (Murata et al., 2012). Statistical or dynamical downscaling could be used for a more detailed analysis (Dibike and Coulibaly, 2006; Wood et al., 2004). Also, this study could gain further insights by considering changing population and socioeconomic distribution in the future. Overall, however, none of these things are expected to change the broad conclusions of this study that global warming will lead to increased exposure to extremes in heat and humidity.

#### **Author contribution**

Conceptualization: J.L., J.M, and A.D. Data curation: J.L. and A.D. Formal analysis: J.L. and J.M. Funding acquisition: A.D. Investigation: J.L. and J.M. Methodology: J.L. Project administration: A.D. Resources: A.D. Software: J.L. Supervision: A.D. Visualization: J.L. Writing: J.L. and A.D.

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### **Acknowledgments**

This work was supported by NSF grants AGS-1661861 and AGS-1841308, both to Texas A&M University. The authors declare that there is no conflict of interest regarding the publication of this article.

## References

- Arbuthnott, K., Hajat, S., Heaviside, C., and Vardoulakis, S.: Changes in population susceptibility to heat and cold over time: assessing adaptation to climate change, *Environmental Health*, 15, 73-93, 2016.
- Argaud, L., Ferry, T., Le, Q. H., Marfisi, A., Ciorba, D., Achache, P., Ducluzeau, R., and Robert, D.: Short- and long-term outcomes of heatstroke following the 2003 heat wave in Lyon, France, *Archives of Internal Medicine*, 167, 2177-2183, DOI 10.1001/archinte.167.20.10170147, 2007.
- Baldwin, J. W., Dessy, J. B., Vecchi, G. A., and Oppenheimer, M.: Temporally Compound Heat Wave Events and Global Warming: An Emerging Hazard, *Earth's Future*, 7, 411-427, 10.1029/2018ef000989, 2019.
- Birk, K., Lupo, A. R., Guinan, P., and Barbieri, C.: The interannual variability of midwestern temperatures and precipitation as related to the ENSO and PDO, *Atmosfera*, 23, 95-128, 2010.
- Buzan, J. R., and Huber, M.: Moist heat stress on a hotter Earth, *Annual Review of Earth and Planetary Sciences*, 48, 2020.
- Chen, R. D., and Lu, R. Y.: Dry Tropical Nights and Wet Extreme Heat in Beijing: Atypical Configurations between High Temperature and Humidity, *Monthly Weather Review*, 142, 1792-1802, 10.1175/Mwr-D-13-00289.1, 2014.
- Chow, W. T., Chuang, W.-C., and Gober, P.: Vulnerability to extreme heat in metropolitan Phoenix: spatial, temporal, and demographic dimensions, *The Professional Geographer*, 64, 286-302, 2012.
- Dahl, K., Licker, R., Abatzoglou, J. T., and Declet-Barreto, J.: Increased frequency of and population exposure to extreme heat index days in the United States during the 21st century, *Environmental Research Communications*, 1, 075002, 2019.
- Davies-Jones, R.: An efficient and accurate method for computing the wet-bulb temperature along pseudoadiabats, *Monthly Weather Review*, 136, 2764-2785, 2008.
- De Rosa, M., Bianco, V., Scarpa, F., and Tagliafico, L. A.: Heating and cooling building energy demand evaluation; a simplified model and a modified degree days approach, *Applied energy*, 128, 217-229, 2014.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, I., Kallberg, P., Kohler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553-597, 10.1002/qj.828, 2011.
- Deser, C., Phillips, A., Bourdette, V., and Teng, H. Y.: Uncertainty in climate change projections: the role of internal variability, *Climate Dynamics*, 38, 527-546, 10.1007/s00382-010-0977-x, 2012.
- Deser, C., and Trenberth, K. E.: The Climate Data Guide: Pacific Decadal Oscillation (PDO): Definition and Indices, 2016.
- Dibike, Y. B., and Coulibaly, P.: Temporal neural networks for downscaling climate variability and extremes, *Neural Networks*, 19, 135-144, 10.1016/j.neunet.2006.01.003, 2006.
- Diffenbaugh, N. S., and Burke, M.: Global warming has increased global economic inequality, *Proceedings of the National Academy of Sciences*, 116, 9808-9813, 2019.

- Fischer, E. M., and Schär, C.: Consistent geographical patterns of changes in high-impact European heatwaves, *Nature Geoscience*, 3, 398-403, 2010.
- Harrington, L. J., Frame, D. J., Fischer, E. M., Hawkins, E., Joshi, M., and Jones, C. D.: Poorest countries experience earlier anthropogenic emergence of daily temperature extremes, *Environmental Research Letters*, 11, 055007, 2016.
- Harrington, L. J., Frame, D., King, A. D., and Otto, F. E.: How uneven are changes to impact-relevant climate hazards in a 1.5° C world and beyond?, *Geophysical Research Letters*, 45, 6672-6680, 2018.
- Hausfather, Z., and Peters, G. P.: Emissions—the ‘business as usual’ story is misleading. Nature Publishing Group, 2020.
- Heo, S., Bell, M. L., and Lee, J. T.: Comparison of health risks by heat wave definition: Applicability of wet-bulb globe temperature for heat wave criteria, *Environmental Research*, 168, 158-170, 10.1016/j.envres.2018.09.032, 2019.
- Hoegh-Guldberg, O., Jacob, D., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F., and Guiot, J.: Impacts of 1.5 C global warming on natural and human systems, *Global warming of 1.5° C. An IPCC Special Report*, 2018.
- Kang, S., and Eltahir, E. A. B.: North China Plain threatened by deadly heatwaves due to climate change and irrigation, *Nat Commun*, 9, 2894, 10.1038/s41467-018-05252-y, 2018.
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J. F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M.: THE COMMUNITY EARTH SYSTEM MODEL (CESM) LARGE ENSEMBLE PROJECT A Community Resource for Studying Climate Change in the Presence of Internal Climate Variability, *Bulletin of the American Meteorological Society*, 96, 1333-1349, 10.1175/Bams-D-13-00255.1, 2015.
- Kharin, V. V., Flato, G. M., Zhang, X., Gillett, N. P., Zwiers, F., and Anderson, K. J.: Risks from Climate Extremes Change Differently from 1.5 degrees C to 2.0 degrees C Depending on Rarity, *Earths Future*, 6, 704-715, 10.1002/2018ef000813, 2018.
- King, A. D., and Harrington, L. J.: The inequality of climate change from 1.5 to 2 C of global warming, *Geophysical Research Letters*, 45, 5030-5033, 2018.
- Kummu, M. T., Maija; Guillaume, Joseph H. A.: Data from: Gridded global datasets for Gross Domestic Product and Human Development Index over 1990-2015 Dryad, Dataset, <https://doi.org/10.5061/dryad.dk1j0>. 2019.
- Lelieveld, J., Hadjinicolaou, P., Kostopoulou, E., Chenoweth, J., El Maayar, M., Giannakopoulos, C., Hannides, C., Lange, M. A., Tanarhte, M., Tyrlis, E., and Xoplaki, E.: Climate change and impacts in the Eastern Mediterranean and the Middle East, *Climatic Change*, 114, 667-687, 10.1007/s10584-012-0418-4, 2012.
- Likas, A., Vlassis, N., and Verbeek, J. J.: The global k-means clustering algorithm, *Pattern Recognition*, 36, 451-461, 2003.
- Liu, Z., Anderson, B., Yan, K., Dong, W., Liao, H., and Shi, P.: Global and regional changes in exposure to extreme heat and the relative contributions of climate and population change, *Scientific Reports*, 7, 1-9, 2017.
- Luber, G., and McGeehin, M.: Climate change and extreme heat events, *American journal of preventive medicine*, 35, 429-435, 2008.

553 Lundgren, K., Kuklane, K., Gao, C., and Holmer, I.: Effects of heat stress on working populations  
 554 when facing climate change, *Industrial health*, 51, 3-15, 2013.  
 555 Maher, N., Milinski, S., Suarez-Gutierrez, L., Botzet, M., Dobrynin, M., Kornblueh, L., Kroger, J.,  
 556 Takano, Y., Ghosh, R., Hedemann, C., Li, C., Li, H. M., Manzini, E., Notz, D., Putrasahan, D., Boysen,  
 557 L., Claussen, M., Ilyina, T., Olonscheck, D., Raddatz, T., Stevens, B., and Marotzke, J.: The Max  
 558 Planck Institute Grand Ensemble: Enabling the Exploration of Climate System Variability, *Journal*  
 559 *of Advances in Modeling Earth Systems*, 11, 2050-2069, 10.1029/2019ms001639, 2019.  
 560 Mann, M. E., Steinman, B. A., Brouillette, D. J., and Miller, S. K.: Multidecadal climate oscillations  
 561 during the past millennium driven by volcanic forcing, *Science*, 371, 1014-1019, 2021.  
 562 Marcotullio, P. J., Keßler, C., and Fekete, B. M.: The future urban heat-wave challenge in Africa:  
 563 Exploratory analysis, *Global Environmental Change*, 66, 102190, 2021.  
 564 Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A.,  
 565 Moufouma-Okia, W., Péan, C., and Pidcock, R.: Global warming of 1.5 C, An IPCC Special Report  
 566 on the impacts of global warming of, 1, 2018.  
 567 Meehl, G. A., Tebaldi, C., Teng, H., and Peterson, T. C.: Current and future US weather extremes  
 568 and El Niño, *Geophysical Research Letters*, 34, 2007.  
 569 Melillo, J. M., Richmond, T., and Yohe, G.: Climate change impacts in the United States, Third  
 570 national climate assessment, 52, 2014.  
 571 Mora, C., Dousset, B., Caldwell, I. R., Powell, F. E., Geronimo, R. C., Bielecki, C. R., Counsell, C. W.,  
 572 Dietrich, B. S., Johnston, E. T., Louis, L. V., Lucas, M. P., McKenzie, M. M., Shea, A. G., Tseng, H.,  
 573 Giambelluca, T., Leon, L. R., Hawkins, E., and Trauernicht, C.: Global risk of deadly heat, *Nature*  
 574 *Climate Change*, 7, 501-+, 10.1038/Nclimate3322, 2017.  
 575 Morris, C. E., Gonzales, R. G., Hodgson, M. J., and Tustin, A. W.: Actual and simulated weather  
 576 data to evaluate wet bulb globe temperature and heat index as alerts for occupational heat-  
 577 related illness, *Journal of Occupational and Environmental Hygiene*, 16, 54-65,  
 578 10.1080/15459624.2018.1532574, 2019.  
 579 Murata, A., Nakano, M., Kanada, S., Kurihara, K., and Sasaki, H.: Summertime temperature  
 580 extremes over Japan in the late 21st century projected by a high-resolution regional climate  
 581 model, *Journal of the Meteorological Society of Japan. Ser. II*, 90, 101-122, 2012.  
 582 North, G. R.: Empirical orthogonal functions and normal modes, *Journal of Atmospheric Sciences*,  
 583 41, 879-887, 1984.  
 584 Parkes, B., Cronin, J., Dessens, O., and Sultan, B.: Climate change in Africa: costs of mitigating  
 585 heat stress, *Climatic Change*, 154, 461-476, 2019.  
 586 Patz, J. A., Campbell-Lendrum, D., Holloway, T., and Foley, J. A.: Impact of regional climate change  
 587 on human health, *Nature*, 438, 310-317, 10.1038/nature04188, 2005.  
 588 Perkins, S., Alexander, L., and Nairn, J.: Increasing frequency, intensity and duration of observed  
 589 global heatwaves and warm spells, *Geophysical Research Letters*, 39, 2012.  
 590 Quinn, A., Tamerius, J. D., Perzanowski, M., Jacobson, J. S., Goldstein, I., Acosta, L., and Shaman,  
 591 J.: Predicting indoor heat exposure risk during extreme heat events, *Science of the total*  
 592 *environment*, 490, 686-693, 2014.  
 593 Ruddell, D. M., Harlan, S. L., Grossman-Clarke, S., and Buyantuyev, A.: Risk and exposure to  
 594 extreme heat in microclimates of Phoenix, AZ, in: *Geospatial techniques in urban hazard and*  
 595 *disaster analysis*, Springer, 179-202, 2009.




- 596 Russo, S., Sillmann, J., and Sterl, A.: Humid heat waves at different warming levels, Scientific  
597 Reports, 7, ARTN 7477
- 598 10.1038/s41598-017-07536-7, 2017.
- 599 Russo, S., Sillmann, J., Sippel, S., Barcikowska, M. J., Ghisetti, C., Smid, M., and O'Neill, B.: Half a  
600 degree and rapid socioeconomic development matter for heatwave risk, Nature communications,  
601 10, 1-9, 2019.
- 602 Sailor, D. J., and Muñoz, J. R.: Sensitivity of electricity and natural gas consumption to climate in  
603 the USA—Methodology and results for eight states, Energy, 22, 987-998, 1997.
- 604 SEDAC: Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11.  
605 Palisades, NY: NASA 2018.
- 606 Shin, M., and Do, S. L.: Prediction of cooling energy use in buildings using an enthalpy-based  
607 cooling degree days method in a hot and humid climate, Energy and Buildings, 110, 57-70, 2016.
- 608 Simolo, C., Brunetti, M., Maugeri, M., and Nanni, T.: Evolution of extreme temperatures in a  
609 warming climate, Geophysical Research Letters, 38, ArtN L16701
- 610 10.1029/2011gl048437, 2011.
- 611 Sivak, M.: Potential energy demand for cooling in the 50 largest metropolitan areas of the world:  
612 Implications for developing countries, Energy Policy, 37, 1382-1384, 2009.
- 613 Stouffer, R. J., and Manabe, S.: Assessing temperature pattern projections made in 1989, Nature  
614 Climate Change, 7, 163-165, 2017.
- 615 Sun, Y., Zhang, X., Zwiers, F. W., Song, L., Wan, H., Hu, T., Yin, H., and Ren, G.: Rapid increase in  
616 the risk of extreme summer heat in Eastern China, Nature Climate Change, 4, 1082-1085, 2014.
- 617 Syakur, M. A., Khotimah, B. K., Rochman, E. M. S., and Satoto, B. D.: Integration K-Means  
618 Clustering Method and Elbow Method For Identification of The Best Customer Profile Cluster,  
619 2nd International Conference on Vocational Education and Electrical Engineering (Icvee), 336,  
620 Unsp 012017
- 621 10.1088/1757-899x/336/1/012017, 2018.
- 622 Tan, J. G., Zheng, Y. F., Tang, X., Guo, C. Y., Li, L. P., Song, G. X., Zhen, X. R., Yuan, D., Kalkstein, A.  
623 J., Li, F. R., and Chen, H.: The urban heat island and its impact on heat waves and human health  
624 in Shanghai, International Journal of Biometeorology, 54, 75-84, 10.1007/s00484-009-0256-x,  
625 2010.
- 626 Thirumalai, K., DiNezio, P. N., Okumura, Y., and Deser, C.: Extreme temperatures in Southeast  
627 Asia caused by El Nino and worsened by global warming, Nat Commun, 8, 15531,  
628 10.1038/ncomms15531, 2017.
- 629 Thompson, D. W. J., Barnes, E. A., Deser, C., Foust, W. E., and Phillips, A. S.: Quantifying the Role  
630 of Internal Climate Variability in Future Climate Trends, Journal of Climate, 28, 6443-6456,  
631 10.1175/Jcli-D-14-00830.1, 2015.
- 632 Trenberth, K. E.: The Climate Data Guide: Nino SST Indices (Nino 1+2, 3, 3.4, 4; ONI and TNI). 2020.
- 633 Trenberth, K. E., and Zhang, R.: The Climate Data Guide: Atlantic Multi-decadal Oscillation (AMO),  
634 2021.
- 635 Wilhelmi, O. V., and Hayden, M. H.: Connecting people and place: a new framework for reducing  
636 urban vulnerability to extreme heat, Environmental Research Letters, 5, 014021, 2010.

- 637 Wobus, C., Zarakas, C., Malek, P., Sanderson, B., Crimmins, A., Kolian, M., Sarofim, M., and  
638 Weaver, C. P.: Reframing Future Risks of Extreme Heat in the United States, *Earths Future*, 6,  
639 1323-1335, [10.1029/2018ef000943](https://doi.org/10.1029/2018ef000943), 2018.
- 640 Wood, A. W., Leung, L. R., Sridhar, V., and Lettenmaier, D. P.: Hydrologic implications of  
641 dynamical and statistical approaches to downscaling climate model outputs, *Climatic Change*, 62,  
642 189-216, DOI [10.1023/B:CLIM.0000013685.99609.9e](https://doi.org/10.1023/B:CLIM.0000013685.99609.9e), 2004.
- 643 Wuebbles, D. J., Fahey, D. W., and Hibbard, K. A.: Climate science special report: fourth national  
644 climate assessment, volume I, 2017.
- 645 Zhang, G., Zeng, G., Li, C., and Yang, X.: Impact of PDO and AMO on interdecadal variability in  
646 extreme high temperatures in North China over the most recent 40-year period, *Climate*  
647 *Dynamics*, 54, 3003-3020, 2020.

648

649



650

651

**Figure 1.** Difference of 1% CO<sub>2</sub> runs compared with ERA-Interim in same level of global

652

warming (0.87°C). The grid points where ERA-Interim falls within the ensemble spread of 1%

653

runs are masked with gray, while other grid points show the difference between the nearest

654

ensemble member and ERA-Interim for (a) 90<sup>th</sup> percentile of 15-year daily average t2m, (b)


655

mean of 15-year daily average t2m, (c) 90<sup>th</sup> percentile of 15-year daily average w2m, and (d)


656

mean of 15-year daily average w2m.


657



**Figure 2.** (a) Location of 15 largest cities in the world and the number of annual heat extremes at (b) 1.5, (c) 2.0, (d) 3.0, and (e) 4.0°C of global warming. Orange (purple) bars represent the ensemble average annual number of deadly days (tropical nights), averaged 5 years after each level of warming is exceeded. Number of heat extreme days are calculated by averaging 3×3 land-only grid covering the selected city. Error bars represent the values of maximum and minimum ensemble members.




**Figure 3.** First three EOFs of annual values of deadly days (a, b, c) and tropical nights (d, e, f) in the world's 15 largest cities. For each panel, the bar graph shows the EOF pattern of the number of heat extreme days per year. Contour plots shows the SST pattern associated with the EOF mode, obtained by projecting each mode of PC onto SST anomalies. Ensemble members are averaged to yield the SST pattern. Pattern correlation with major modes of climate variability (ENSO, PDO, AMO) are also shown, as discussed in the text.



**Figure 4.** Frequency power spectrum of ENSO, PDO, and PC of first three EOF modes for (a) deadly days and (b) tropical nights. ENSO is calculated with the Niño 3.4 Index, and PDO is calculated as a leading EOF of SST anomaly in North Pacific basin. Monthly SST data is used for both ENSO and PDO, and then each index is averaged over the year to have consistency with deadly days and tropical nights.

680



681

682

**Figure 5.** (a) Clustered regions via K-means clustering. Characteristics of each cluster are listed

683

in Table 2. (b) Zonal average of temperature increases at the time of 0.87°C (our reference

684

period), 1.5°C, 2°C, and 4°C of global warming compared to pre-industrial baseline in the 1%


685

runs. Temperatures are averaged over a 5-year period after each warming threshold is exceed in

686

the model.


687



**Figure 6.** Evolution of each index averaged over each cluster. Colors are consistent with Figure 5 and Table 2. Values of each metric are calculated by averaging the grid points that belongs to each cluster. This was done for each ensemble member and then the ensemble average is plotted. Vertical lines with dots show the maximum and minimum of 28 ensemble members at each threshold of warming to represent the spread between the ensemble members.




694




695

696 **Figure 7.** Changes of population-weighted heat wave indices as a function of global average  
 697 warming. Each line denotes one ensemble member for different percentiles of population.

698



**Figure 8.** Increase in (a) deadly days and (b) tropical nights compared to the reference period (0.87°C warming), binned by percentile of GDP per capita at selected levels of warming compared to reference climate (calculated by subtracting reference values, shown as heatmap), averaged over the population within the GDP percentile (for example, averaged over population in 0~10 percentile of GDP), and over all ensemble members for 5-year window after each level of warming first occurs. Green text inside the heatmap represent the absolute number of deadly days and tropical nights in each level of warming.



**Figure 9.** Change (in percentage) of ensemble averaged cooling degree days (CDD; red) and heating degree days (HDD; blue) compared to the reference climate (0.87°C) in the 1% CO<sub>2</sub> experiments at the time they reach the global mean temperature thresholds of (a) 1.5°C, (b) 2.0°C, (c) 3.0°C, and (d) 4.0°C, respectively. Error bars represent the standard deviation of CDD and HDD values between the ensemble members.

715 **Table 1.** Explanation of heat wave indices used in this study.

Acronym	Index	Definition	Units
HWD <sub>t2m/w2m</sub>	Heat wave duration	Length of longest period of consecutive heat wave days in a year	# days
HWF <sub>t2m/w2m</sub>	Heat wave frequency	Total number of heat wave days in a year	# days
HWA <sub>t2m/w2m</sub>	Heat wave amplitude	Maximum temperature over all heat wave days in a year	°C
HWM <sub>t2m/w2m</sub>	Heat wave mean	Average temperature over all heat wave days in a year	°C
Deadly Days	Deadly Days	Daily maximum wet-bulb temperature over 26°C	# days
Tropical Nights	Tropical Nights	Daily minimum temperature over 25°C	# days
CDD	Cooling degree days	Sum of positive values after removing 18°C from daily average temperature	°C days
HDD	Heating degree days	Absolute value of sum of negative values after removing 18°C from daily average temperature	°C days

717 **Table 2.** Percentage area and major regions belonging to each cluster. Clusters are identified  
 718 only for the global land areas.

Cluster	Color	Area percentage (%)	Major regions	Cluster name
1	Maroon	2.95	Indonesia, Malaysia, Cameroon, Gabon Northern South	Tropical West Pacific
2	Orange	12.34	America, Central Africa	Tropical Africa and America
3	Pink	22.70	India, Southeast Asia, Eastern South America, Southeast U.S.	Sub-Tropical Asia and America
4	Green	21.55	Northern Africa, Middle East, Australia	Deserts
5	Sky blue	7.69	Himalayas, Andes	Mountain Range
6	Blue	32.75	Canada, Northwest U.S., Russia	Sub-Polar Region

719

**Table 3.** Number of deadly days each percentile of global population faces with reference period (0.87°C), 1.5°C, 2°C, 3°C, and 4°C global warming from the pre-industrial condition. Standard deviations between the ensembles ( $1\sigma$ ) are also shown.

		Global Warming				
Population		0.87°C	1.5°C	2.0°C	3.0°C	4.0°C
Deadly Days	95 <sup>th</sup> p.	85 ( $\pm 7$ )	105 ( $\pm 10$ )	125 ( $\pm 7$ )	161 ( $\pm 12$ )	229 ( $\pm 15$ )
	90 <sup>th</sup> p.	45 ( $\pm 5$ )	65 ( $\pm 10$ )	86 ( $\pm 8$ )	132 ( $\pm 12$ )	198 ( $\pm 12$ )
	50 <sup>th</sup> p.	0.3 ( $\pm 0.1$ )	1.5 ( $\pm 1.3$ )	5 ( $\pm 2$ )	23 ( $\pm 4$ )	63 ( $\pm 5$ )
Tropical Nights	95 <sup>th</sup> p.	211 ( $\pm 11$ )	232 ( $\pm 14$ )	253 ( $\pm 13$ )	306 ( $\pm 17$ )	358 ( $\pm 3$ )
	90 <sup>th</sup> p.	280 ( $\pm 7$ )	195 ( $\pm 9$ )	205 ( $\pm 9$ )	232 ( $\pm 12$ )	277 ( $\pm 14$ )
	50 <sup>th</sup> p.	15 ( $\pm 4$ )	27 ( $\pm 7$ )	41 ( $\pm 6$ )	71 ( $\pm 6$ )	102 ( $\pm 4$ )