

Regional impacts of black carbon morphologies on shortwave aerosol-radiation interactions: A comparative study between the US and China

Jie Luo^{1,2}, Zhengqiang Li¹, Chenchong Zhang³, Qixing Zhang^{2,*}, Yongming Zhang², Ying Zhang¹, Gabriele Curci^{5,6}, and Rajan K. Chakrabarty^{3,4}

¹State Environment Protection Key Laboratory of Satellite Remote Sensing, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100101, China

²State Key Laboratory of Fire Science, University of Science and Technology of China, Hefei, Anhui 230026, China

³Center for Aerosol Science and Engineering, Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, St. Louis, MO 63130, USA

⁴McDonnell Center for the Space Sciences, Washington University in St. Louis, St. Louis, MO 63130, USA

⁵Department of Physical and Chemical Sciences, University of L'Aquila, L'Aquila, Italy

⁶Center of Excellence in Telesensing of Environment and Model Prediction of Severe Events (CETEMPS), University of L'Aquila, L'Aquila (AQ), Italy

Correspondence: Qixing Zhang (qixing@ustc.edu.cn)

Abstract. Black carbon (BC) is one of the dominant absorbing aerosol species in the atmosphere. It normally has complex fractal-like structures due to the aggregation process during combustion. A wide range of aerosol-radiation interactions (ARI) of BC has been reported throughout experimental and modeling studies. One reason for the large discrepancies among multiple studies is the application of the over-simplified spherical morphology for BC in ARI estimates. In current climate models, the Mie theory is commonly used to calculate the optical properties of spherical BC aerosols. Here, we employ a regional chemical transport model coupled with a radiative transfer code that utilizes the non-spherical BC optical simulations to re-evaluate the effects of particles' morphologies on BC shortwave ARI, and the wavelength range of 0.3 – 4.0 μm was considered. Anthropogenic activities and wildfires are two major sources of BC emissions. Therefore, we choose the typical polluted area in eastern China which is dominated by anthropogenic emissions, and the fire region in the northwest US which is dominated by fire emissions in this study. A one-month simulation in eastern China and a seven-days simulation in the fire region in the northwest US were performed. The fractal BC model generally presents a larger clear-sky ARI compared to the spherical BC model. Assuming BC particles are externally mixed with other aerosols, the relative differences in the time-averaged clear-sky ARI between the fractal model with a fractal dimension (D_f) of 1.8 and the spherical model are 12.1% – 20.6% and 10.5% – 14.9% for typical polluted urban cities in China and fire sites in the northwest US, respectively. Furthermore, the regional-mean clear-sky ARI is also significantly affected by the BC morphology, and relative differences of 17.1% and 38.7% between the fractal model with a D_f of 1.8 and the spherical model were observed in eastern China and the northwest US, respectively. However, the existence of clouds would weaken the BC morphological effects. The time-averaged all-sky ARI relative differences between the fractal model with a D_f of 1.8 and the spherical model are 4.9% – 6.4% and 9.0% – 11.3% in typical urban polluted cities and typical fire sites, respectively. Besides, for the regional-mean all-sky ARI, the relative

20 differences between the fractal model and the spherical model are less than 7.3% and 16.8% in the polluted urban area in China and the fire region in the US, respectively. The results imply that current climate modeling may significantly underestimate the BC ARI uncertainties as the morphological effects on BC ARI are ignored in most climate models.

1 Introduction

Black carbon (BC), as the main absorbing aerosol in the atmosphere, exerts a positive radiative forcing, and lofts smoke
25 plumes (Buseck and Buseck, 2000; Streets et al., 2006; Moosmüller et al., 2009). However, there are still large uncertainties in evaluating the BC radiative effects. An important cause of the discrepancy is BC's complex morphology. BC aerosols are assumed as spheres, and the optical properties are calculated using the Mie theory in most climate and atmospheric chemical transport models, such as Community Earth System Model (CESM) (Danabasoglu et al., 2020), the Model for Interdisciplinary Research on Climate (MIROC-SPRINTARS) (Takemura et al., 2005, 2009), Weather Research and Forecasting coupled to
30 Chemistry (WRF-Chem) (Grell et al., 2005; Fast et al., 2006), and GEOS-Chem. However, many studies have shown that BC particles, especially those nascent ones, have fractal-like structures. The spherical assumption for BC can lead to a large deviation from the field measurement data and non-spherical simulated results (Chakrabarty et al., 2007; Luo et al., 2018c; He et al., 2015; Liu and Mishchenko, 2005; Luo et al., 2018d; Mishchenko et al., 2016a; Luo et al., 2021b). Based on the sampled BC images, researchers found that the shape of uncoated BC aggregates can be fitted well by a fractal law with
35 monomer number (N_s), mean monomer radius (R), fractal prefactor (k_0), fractal dimension (D_f) and the radius of gyration (R_g) (Mishchenko et al., 2002; Sorensen, 2011; Luo et al., 2021a):

$$N_s = k_0 \left(\frac{R_g}{R} \right)^{D_f} \quad (1)$$

Previous studies have shown that aggregated particle models are more realistic to reproduce the optical measurement results (Kahnert, 2010a; Luo et al., 2019, 2018b). Some studies have used the fractal BC models to investigate the radiative properties of BC (Wu et al., 2015; Liu et al., 2015a; Liu and Mishchenko, 2005; Yin and Liu, 2010; Teng et al., 2019; Luo et al., 2018a;
40 Kahnert, 2010a). The direct radiative effect (DRE) was widely used to evaluate the climate effects of aerosol (Bond et al., 2013; Saleh et al., 2015). IPCC (2014) suggested using the new terminology of the aerosol-radiation interactions (ARI) instead of DRE. Thus, in this work, we use the terminology of ARI to replace DRE. However, an extremely limited number of studies have evaluated the ARI of non-spherical BC in regional or global climate models. Kahnert (2010b) has made efforts to simulate the radiative properties of freshly emitted BC using the Multiple-scale Atmospheric Transport and CHemistry (MATCH) model.
45 That study assumed a fixed solar zenith angle (SZA) and restricted the modeling region in Western Europe. Expanding the modeling range to regions with different emission characteristics is important to understand the effects of BC sources on ARI.

A global mean BC all-sky ARI of $+0.6 \text{ W m}^{-2}$ has been reported by IPCC (2014). However, the BC ARI values estimated based on in-situ optical measurements in some regions are much larger than the rest. BC emissions in China roughly account for one-fourth of its global anthropogenic emission budget (Streets et al., 2001). Eastern China, a typical polluted region, is
50 dominated by anthropogenic emissions (Zhang et al., 2009; Li et al., 2017a). Therefore, BC ARI in eastern China has gained increasing interest.

Besides the anthropogenic sources, wildfires significantly contribute to the regional BC emissions. The extremely-high BC concentrations can be found in those fire sites. Recent studies have shown that BC emitted from fire sites can also loft the surrounding atmospheric aerosols to the upper troposphere and lower stratosphere (Yu et al., 2019). Thus, the investigation of BC ARI in these regions is important to understand the detailed plume dynamics and the warming effects of BC. Northwest US, as one of the regions where the wildfire is most frequent, has also been investigated in addition to eastern China.

In this work, we employed WRF-Chem to simulate the aerosol mass concentrations. Note here that WRF-Chem assumes aerosols to be spherical. Therefore, the radiative parameters of fractal BC aggregates were calculated offline using an optical module, Flexible Aerosol Optical Depth (FlexAOD) (Curci, 2012). We calculated the ARI at the top of the atmosphere (TOA) using a radiative transfer model, libRadtran (Mayer and Kylling, 2005), after the particles' optical properties were calculated.

2 Method

2.1 Aerosol distribution simulation

In this work, WRF-Chem version 4.1.3 was used to simulate the transport of atmospheric species. Two areas were selected to represent the BC sources with different emission characteristics. Eastern China, a major polluted region in the world, represents the typical polluted urban region. It consists of 115 east-west grids and 105 south-north grids centered at 112.00°E, 35.00°N with a grid resolution of 18 km. The northwest US, one of the regions where the wildfire is most frequent in the world, was also investigated in this work. The fire region consists of 120 east-west grids and 120 south-north grids centered at 121.48°W, 39.89°N with a grid resolution of 4 km. The schematics of the two regions are shown in Figure S1. Both regions have 33 vertical layers above the ground, with a top pressure of 50 hPa.

We used the Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1) to compute the biogenic emissions over two regions (Guenther et al., 1994, 2006). The anthropogenic inventory for eastern China in the year 2016 was compiled by the Multi-resolution Emission Inventory for China (MEIC) (Li et al., 2014; Liu et al., 2015b). We used the MIX anthropogenic inventory for the region outside China (Li et al., 2017b). The Regional Acid Deposition Model version 2 (RADM2) atmospheric chemical mechanism (Stockwell et al.) and the Model Aerosol Dynamics for Europe with the Secondary ORGanic Aerosol Model (MADE/SORGAM) were applied in the simulation of eastern China (Seinfeld et al., 2001; Ackermann et al., 1998). Fast-J photolysis scheme (Wild et al., 2000) was used to simulate the photolysis rates. The physical scheme options in WRF-Chem are shown in Table S1. The simulations in eastern China started at 0:00 UTC on November 31st, 2016, and ended at 00:00 UTC on January 1st, 2017. The data from 0:00 UTC on December 1st, 2016 to 00:00 UTC on January 1st, 2017 was used for analysis.

The Emission Database for Global Atmospheric Research for Hemispheric Transport of Air Pollution (EDGAR-HTAP) version 2 emission inventory for 2010 was used in the northwest US. MOSAIC aerosol scheme (Zaveri and Peters, 1999; Zaveri et al., 2008) and the Carbon Bond Mechanism Z (CBM-Z) photochemical mechanism (Zaveri and Peters, 1999) were used in the fire region simulation. The Fire emission was provided by the Fire INventory from NCAR (FINN) (Wiedinmyer et al., 2011). Note here that EDGAR-HTAP anthropogenic inventory and FINN were provided for the MOZART chemical mecha-

nism, so we mapped the emission for the Model for Ozone and Related chemical Tracers (MOZART) chemical mechanism to the CBM-Z chemical mechanism based on the study of Emmons et al. (2010). For both simulations in eastern China and the northwest US, the National Center for Environmental Prediction (NCEP) Global Forecast System (GFS) Final Analysis (FNL) with a horizontal grid spacing of 0.25° and 6 h intervals was used to provide the meteorological initial and boundary conditions. The chemical initial and boundary conditions were obtained from the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4). The simulations in the northwest US started at 0:00 UTC on August 5th, 2016, and ended at 00:00 UTC on August 21st, and the data from 0:00 UTC on August 14th to 00:00 UTC on August 21st was used for analysis.

2.2 The morphology of BC

In this work, we only consider externally mixed BC aerosols, which are commonly represented by fractal structures. D_f is a key parameter to describe the compactness of fractal BC (Wang et al., 2017; Yuan et al., 2019). D_f increases from approximately 1.8 to 3 when the BC morphology can vary from a chain-like structure to a spherical structure. The freshly emitted BC generally exhibits a fluffy structure with a D_f of approximately 1.8 (Heinson et al., 2010, 2017). The laboratory measurements also showed that the freshly emitted BC generally presents a small D_f . Chakrabarty et al. (2006) have shown that D_f of BC emitted from wildland fuels generally exhibits a range of 1.67 – 1.83. A D_f range of 1.6 – 1.9 was observed for BC produced from diesel combustion (Wentzel et al., 2003). China et al. (2013b) indicated that the BC freshly emitted from wildfire generally exhibits D_f with a range of 1.74 – 1.92. However, a more compact structure was commonly observed for BC in the atmosphere with the particle aging (Li et al., 2003; Adachi et al., 2014; Chen et al., 2016; Adachi et al., 2010). A D_f range of 2.2 – 2.4 was observed in the study of Adachi et al. (2010). The fractal structures with a larger D_f are widely used to describe the BC with more compact structures (Adachi et al., 2010). Chakrabarty et al. (2006) further showed that the D_f of aged BC can reach up to 2.6. To represent both fluffy and compact BC, D_f s of 1.8, 2.2, and 2.6 were considered in this study. While the fractal prefactor k_0 was also measured in a wide range in the atmosphere, its impact on the optical properties was relatively small. We assumed a fixed k_0 of 1.2 in this work. The typical morphologies of fractal BC are shown in Figure 1.

The volume-mean particle radius was commonly used to describe the size of non-spherical BC. Previous studies have observed a range of approximately 8 – 57 nm for BC monomer radius (Eggersdorfer and Pratsinis, 2012; Mikhailov et al., 2006; KÖylü and Faeth, 1992; Lee et al., 2002), while Kahnert and Kanngießner (2020) further showed that the typical range is approximately 10 – 25 nm. In this work, we assumed a fixed monomer radius of 20 nm. We considered an N_s range of 1 – 1000 to represent BC with different sizes. The volume-mean particle radius (r_p) can be calculated using:

$$r_p = RN_s^{1/3} \quad (2)$$

Note here that BC can also internally mix with other compositions, and the morphology can become more complex (Wang et al., 2021c, 2017). However, we mainly focus on the freshly emitted BC, and only consider externally mixed BC. Further investigations would be performed for more complex internally mixed BC in the future.

2.3 The refractive index and size distribution of BC

BC refractive index shows spectral dependence (Chang and Charalampopoulos, 1990), while it doesn't vary largely with wavelengths in the short wavelength range (Liu et al., 2018; Lack and Cappa, 2010; Bond and Bergstrom, 2006). The suggested BC refractive index values by Bond and Bergstrom (2006) were commonly used. In this work, the median value of $1.85 + 0.71i$ was used, as it was widely used in many regional and global climate models (e.g. WRF-Chem). The size distribution of BC also suffers large uncertainties from different fuels and conditions. The size distribution of BC is commonly fitted by a lognormal size distribution with a geometric mean radius (r_g), and a geometric standard deviation (σ_g) (Schwarz et al., 2008; Mishchenko et al., 2016b):

$$n(r_p) = \frac{N_0}{\sqrt{2\pi}r_p \ln(\sigma_g)} \exp \left[- \left(\frac{\ln(r_p) - \ln(r_g)}{\sqrt{2\ln(\sigma_g)}} \right)^2 \right] \quad (3)$$

where $n(r_p)$ is the probability density distribution of particle number concentrations; r_p is the volume-mean particle radius, N_0 is the number concentration, which can be calculated by the mass concentration obtained from WRF-Chem by assuming BC mass density, r_g , and σ_g . The details about the calculation of N_0 are shown in Curci (2012).

BC geometric mean radius of $0.05 - 0.06 \mu\text{m}$ is frequently observed by instruments and widely assumed in numerical studies (Alexander et al., 2008; Coz and Leck, 2011; Reddington et al., 2013; Liu et al., 2018). In this work, BC geometric mean radius was assumed to be $0.05 \mu\text{m}$, and σ_g was assumed to be 1.6. We used the volume-equivalent radius of to characterize the particle size of factual BC. The density of BC was assumed to be 1.8 gm^{-3} based on the suggested values by Bond and Bergstrom (2006).

2.4 BC radiative properties

In this work, BC radiative properties were calculated using the Multiple Sphere T-matrix Method (MSTM) (Mackowski and Mishchenko, 2011, 1996). The MSTM can efficiently calculate the optical properties of spheres without intersecting surfaces. The MSTM has high computational efficiency because it theoretically calculates the optical properties of randomly oriented particles without numerically averaging them over different particle orientations. The MSTM can directly calculate the extinction efficiency (Q_{ext}), scattering efficiency (Q_{sca}), and phase function (P) with the refractive index, wavelength, and the input shapefile. Then, the extinction cross-section (C_{ext}) and scattering cross-section (C_{sca}) were obtained using:

$$C_{ext} = Q_{ext} \pi r_p^2 \quad (4)$$

$$C_{sca} = Q_{sca} \pi r_p^2 \quad (5)$$

Extinction coefficient (b_{ext}), Scattering coefficient (b_{sca}) and bulk phase function $\langle P \rangle$ were calculated using the following equations:

$$b_{ext} = \int_{r_{min}}^{r_{max}} C_{ext}(r_p) n(r_p) dr \quad (6)$$

$$b_{sca} = \int_{r_{min}}^{r_{max}} C_{sca}(r_p)n(r_p)dr \quad (7)$$

$$\langle P(\theta) \rangle = \frac{\int_{r_{min}}^{r_{max}} C_{sca}(r_p)P(\theta, r_p)n(r_p)dr}{b_{sca}} \quad (8)$$

145 In climate modeling, instead of using the phase function, the Legendre expansion coefficients were commonly used:

$$\langle P(\theta) \rangle = \sum_{S=1}^{S_{max}} \alpha_S P_S(\cos\theta) \quad (9)$$

where θ is the scattering angle, P_S are generalized spherical functions, α_S are the Legendre expansion coefficients, and S_{max} is the order of truncation of the Legendre expansion coefficients.

In this work, we used the pmom tool which is available in libRadtran software for calculating the Legendre expansion
150 coefficients. With the inputs of the aerosol bulk phase function and the desired number of Legendre expansion coefficients, the pmom tool can calculate the Legendre expansion coefficients.

The radiative properties of fractal BC and spherical BC were calculated at 300 nm – 4000 nm wavelengths. The step size
of $\Delta\lambda = 50$ nm was chosen when λ is less than 1000 nm, while $\Delta\lambda = 200$ nm was selected for $1000 \text{ nm} \leq \lambda \leq 2000$ nm, and
 $\Delta\lambda = 400$ nm when $\lambda \geq 2000$ nm. We created look-up tables for b_{ext} , b_{sca} , and the Legendre expansion coefficients of phase
155 functions for each σ_g and r_g . Thus, the optical properties of BC can be obtained by interpolating the look-up tables.

2.5 Flexible Aerosol Optical Depth (FlexAOD)

The aerosol mass concentrations from WRF-Chem were inputted into an optical software Flexible Aerosol Optical Depth
(FlexAOD) (Curci, 2012; Curci et al., 2015) to calculate the aerosol radiative properties. FlexAOD is an optical post-processing
tool for the atmospheric chemistry-transport model, and it started as a tool for the GEOS-Chem model. We have made some
160 modifications to FlexAOD to make it accommodate the WRF-Chem outputs. We first mapped the aerosols from WRF-Chem
into 5 categories: BC, organic carbon(OC), inorganic salt (INS), sea salt (SA), and dust (DST). The mapping details are shown
in Tables S2 – S3. After the WRF-Chem species were mapped, the size distribution, refractive indices, and hygroscopic growth
factors were then assigned.

FlexAOD firstly reads the aerosol mass concentrations from WRF-Chem, then converts them to aerosol volume concentra-
165 tions based on the assigned mass densities. Based on the assigned normalized size distributions, we can calculate the number
concentration (N_0) of each aerosol. FlexAOD pre-calculates the optical properties of each type of aerosol by assuming N_0
= 1 with the assumed size distributions. The total scattering/extinction coefficients can be obtained by multiplying the pre-
calculated scattering/extinction cross-sections with the number concentrations. The phase function of each type of aerosol is
identical to the pre-calculated phase function by assuming $N_0 = 1$, then the total phase function was calculated according to
170 the number concentration of each aerosol. In FlexAOD, aerosol shapes were assumed to be spherical and the corresponding

optical properties of each aerosol species were calculated using the Mie code provided by Mishchenko et al. (1999). The bulk optical properties were then calculated by combining an assembly of aerosols.

BC optical properties were overwritten using the look-up tables created in Section 2.4. As described in Section 2.4, we have created look-up tables for non-spherical BC with different size distributions. Thus, if the size distribution of BC is assigned, the optical properties of BC with a normalized size distribution can be determined by interpolating the look-up tables. Once the number concentration is calculated, we can determine the total optical properties.

Apart from BC, the physical properties of other chemical species were also specified. We used the OC refractive indices suggested by Highwood (2009). The density of OC varies under different conditions. The density of the oxidized organic aerosol was reported to be approximately 1.3 g cm^{-3} (Cross et al., 2007), while Nakao et al. (2013) reported that the density of OC with lower oxidation was approximately $1 - 1.2 \text{ g cm}^{-3}$. For freshly formed OC, $0.9 - 1.1 \text{ g cm}^{-3}$ was used by Liu et al. (2017b). In this work, the density of OC was assumed to be 1.2 g m^{-3} . OC size is also commonly fitted by a lognormal size distribution. In the study of He et al. (2016) and Dentener et al. (2006), $r_g = 0.03 \mu\text{m}$ and $r_g = 0.075 \mu\text{m}$ were assumed for hydrophobic and hydrophilic OC, respectively. In this study, all OC was assumed to be hydrophilic, and we assumed a r_g of $0.075 \mu\text{m}$ and a σ_g of 1.6 for OC. The refractive indices of dust were identical to those used in The Goddard Chemistry Aerosol Radiation and Transport (GOCART) Model (Chin et al., 2002). For dust, the gamma distribution was assumed: (Martin et al., 2003; Curci, 2012):

$$n(r_p) = N_0 r_p^{(1-3b)/b} \exp\left(-\frac{r_p}{ab}\right) \quad (10)$$

where a and b are two parameters for the distribution, and b is in the range of $0 - 0.5$.

The refractive indices of other chemical species were adapted from the Optical Properties of Aerosols and Clouds (OPAC) package (Hess et al., 1998). The physical properties are displayed in Table S4. Similar to the study of Curci et al. (2019), the hygroscopic growth factors of different aerosols were taken from the OPAC package (Hess et al., 1998). Note here that many internally mixed particles exist in the atmosphere, while in this study we mainly aim to study the morphological effects of freshly emitted particles, and more complex particles may be investigated in the future. Effective refractive indices were calculated using the volume mixing method for hydrophilic particles.

The total column single-scattering albedo (SSA) and Aerosol Optical Depth (AOD) were calculated by FlexAOD, and Absorption Aerosol Optical Depth (AAOD) was calculated by:

$$\text{AAOD} = \text{AOD}(1 - \text{SSA}) \quad (11)$$

2.6 ARI modeling

The optical properties (extinction coefficient, SSA, asymmetric factor (ASY)) calculated using FlexAOD at each WRF-Chem grid were inputted into a radiative transfer model, libRadtran (Mayer and Kylling, 2005), to calculate the radiative fluxes at the top-of-the-atmosphere (TOA). The radiative transfer equation was solved by DIScrete Ordinate Radiative Transfer (DISORT) radiative transfer equation solver (Stamnes et al., 1988; Buras et al., 2011). The libRadtran can select a standard atmosphere background and determine the solar zenith angle (SZA) based on the longitude, latitude, and UTC time. The surface albedo

information was obtained from NASA Earth Observations (NEO). The radiative transfer calculations were performed for each
205 hour and then were averaged over one day. In this work, ARI of BC aerosol were calculated using following equations:

$$\text{ARI} = \text{FLUX}_{\text{With BC}} - \text{FLUX}_{\text{Without BC}} \quad (12)$$

$$\text{FLUX} = F^{\downarrow} - F^{\uparrow} \quad (13)$$

where F^{\downarrow} represents downward radiative flux and F^{\uparrow} represents upward radiative flux. In this work, we just considered the ARI
210 at the TOA. Only shortwave ARI was considered, and the wavelength is in the range of 0.3 – 4.0 μm .

In this work, both the clear-sky ARI and all-sky ARI were calculated. The daily-mean cloud optical thickness, cloud effective
radius, and cloud cover data from the MOderate Resolution Imaging Spectroradiometer (MODIS) products were used for the
all-sky ARI calculations. The regional-mean BC ARI was also calculated. The aerosol optical properties and cloud properties
were firstly averaged, then the radiative transfer calculations were performed for only one day (the median day), and similar
215 methods were also used in previous studies (eg. Saleh et al. (2015); Tuccella et al. (2020a)).

3 Results

3.1 Impacts of BC morphology on AOD and AAOD

To verify the modeling performance of the aerosol concentrations, we compared the simulated $\text{PM}_{2.5}$ concentrations with
observations at some monitoring sites, and the results are shown in Figure 2. In the figures, the left column represents the typical
220 cities in eastern China, and the right column represents the sites in the northwest US. The calculated $\text{PM}_{2.5}$ concentrations in
eastern China are generally consistent with the observations. Even though the simulated $\text{PM}_{2.5}$ concentrations in the fire region
are a little higher than the observations, the deviations are not large, and the general trends are consistent. Therefore, it is
reasonable to represent the atmospheric aerosol concentrations using WRF-Chem modeling.

In this study, we selected three fire sites to evaluate the morphological effects on the BC ARI. The positions of the selected
225 sites are shown in Table S5, and they represent the fire sites with high aerosol concentrations. As shown in Figure 3, the temporal
BC concentrations at fire sites can even exceed approximately 400 $\mu\text{g}/\text{m}^3$ when the fire occurs, while the BC concentrations
are extremely low in other days. As shown in Figure 3, even though the maximum BC concentrations in the polluted urban
cities are much smaller than the fire sites, the mean BC concentrations can reach approximately 12 $\mu\text{g}/\text{m}^3$. The simulated BC
concentrations generally agree with the measurements of Zhang et al. (2012) for the urban region, where BC concentrations
230 were observed to be approximately 4 – 12.7 $\mu\text{g}/\text{m}^3$. We further compare the calculated AOD and AAOD with observations
from AErosol RObotic NETwork (AERONET). The AERONET data of Beijing is available among all selected cities in Eastern
China. As shown in Figure 4, the calculated AAOD and AOD can generally represent the observations.

Figure 5 shows the AOD of BC with different morphologies, where BC AOD was calculated by the difference between
AOD with BC and that without BC. Zhuang et al. (2019) indicated that the time-averaged BC AOD in Beijing, Hefei, and

235 Taihu was approximately 0.05 – 0.06. Our simulations are generally in agreement with their results, and the simulated BC AOD varies from approximately 0.01 to approximately 0.12 in the selected urban cities. In Beijing and Tianjin, BC AOD can reach approximately 0.12, while in Shanghai and Nanjing, the maximum BC AOD is approximately 0.07 and 0.1, respectively. The maximum BC AOD can reach 0.5 – 0.9 in typical fire sites. From Figure 5, we can also see that BC AOD calculated using the spherical model is relatively higher than those using fractal aggregate models, which is consistent with the findings of Liu and Mishchenko (2005). As shown in Figure 6, in the urban area a spherical assumption for BC lead to an overestimation of less than 0.03 for AOD compared to the fractal model with a D_f of 1.8, while the overestimation can reach approximately 0.15 in fire sites. The overestimation accounts for a large proportion of BC AOD, which can exceed 20% of the total BC AOD.

Figure 7 shows the calculated BC AAOD using different BC models, where BC AAOD was calculated by the difference between AAOD with BC and that without BC. . Our modeling results show that a more compact structure may lead to a smaller AAOD, and this is consistent with the findings of Liu and Mishchenko (2005) for single BC particles. The reason is that a more compact structure blocks the light from transmitting into the inner part of the particle, and a smaller absorption efficiency was observed (Kahnert and Devasthale, 2011). Shin et al. (2019a) showed that most BC AAOD at 440 nm in Asia is within the range of 0 – 0.12. Our simulated BC AAOD at 450 nm is generally consistent with their findings. At fire sites, BC AAOD at 450 nm wavelength can reach approximately 0.7. As shown in Figure 8, the spherical model underestimates AAOD by approximately 0.016 compared to the fractal model with a D_f of 1.8 in typical polluted cities, while the AAOD underestimation using the spherical model can reach approximately 0.04 in fire sites. In general, the AAOD underestimation using the spherical model is approximately 8% of the total BC AAOD.

Figure 9 shows the time-averaged BC AOD at 550 nm and AAOD at 450 nm as a percentage of the total AOD and AAOD, respectively. In typical China polluted cities, BC AOD at 550 nm accounts for approximately 4.6% – 7% of the total AOD, while BC AOD in fire sites can account for larger than 10% of the total AOD. At 450 nm, in both polluted urban and fire sites, the fractions of BC AAOD are close, which is approximately 30%. This means that the relative proportions of BC and OC in polluted urban sites are close to those of fire sites.

BC morphologies also have significant impacts on the BC AOD and AAOD fractions. As BC morphologies change from a fractal dimension of 1.8 to a spherical structure, BC AOD fraction increases from 4.6% – 5.5% to 5.8% – 6.9% in typical polluted urban cities and from 9.0% – 10.3% to 11.1% – 12.7% in typical fire sites, respectively. The relative differences between the fractal model and the spherical model ((Fractal – Sphere)/Sphere) can be above 25%. For BC AAOD fraction, the values can vary in the range of 25.0% – 33.2% in typical polluted urban cities and the range of 25.4% – 30.0% in typical fire sites, respectively. The AAOD fraction relative differences between the fractal model with a D_f of 1.8 and the spherical model are approximately 10%.

265 3.2 Impacts of BC morphology on SSA

SSA, as the ratio of scattering to extinction, is widely used to infer aerosol types. Figure 10 shows the comparison of SSA using different BC models. Shin et al. (2019b) showed that the mean SSA at 440 nm was approximately 0.89 – 0.92 in Beijing using AERONET data, and Shaheen et al. (2019) demonstrated that the mean SSA at 440 nm was approximately 0.897 in

winter Beijing. At 450 nm, the calculated SSA in typical urban cities is within the range of a 0.86 – 0.92, which is generally
 270 consistent with the observations in previous studies. While in the fire region, SSA varies in a wider range. At 450 nm, SSA at
 the selected fire sites can vary in the range of 0.75 – 0.9, which is a little smaller than that in polluted urban areas due to large
 portions of carbonaceous aerosols in the fire region.

With more compact structures, SSA presents a larger value, which is consistent with the findings of Kahnert and Devasthale
 (2011). However, the effects of BC morphologies on total SSA at 450 nm are not obvious due to the small percentage of BC in
 275 the atmosphere. As shown in Figure 11, the overestimations of the spherical BC model for SSA at 450 nm are generally within
 0.005 at typical polluted cities and within 0.012 at fire sites, which is less than 1% of the total SSA. However, these values may
 have relatively larger impacts when evaluating the climate effect of BC, as BC commonly presents a relatively small value of
 0.2 – 0.4 (Kahnert and Devasthale, 2011).

3.3 Impacts of BC morphology on ARI

BC clear-sky ARI varies in different regions and the reported BC ARI varies in previous studies. Zhuang et al. (2018) esti-
 mated BC clear-sky ARI to be $+1.85 \text{ Wm}^{-2}$ in East Asia. Much larger BC clear-sky ARI during December ($+15 \text{ Wm}^{-2}$) and
 November ($+8 \text{ Wm}^{-2}$) over Ahmedabad and Gurushikhar, respectively, were reported by Rajesh and Ramachandran (2018).
 Zhuang et al. (2019) showed that clear-sky ARI averaged over East Asia were $+0.02$ to $+1.34 \text{ W/m}^2$ in summer in eastern Asia.
 Lu et al. (2020) showed daily-mean BC clear-sky ARI were within the range of $+1.37 - 4.89 \text{ Wm}^{-2}$ in Beijing. Our calculated
 285 daily-mean BC ARI in winter generally agrees with those previous studies. The daily-mean clear-sky BC ARI at the TOA
 in typical sites using different BC models is presented in Figure 12. In winter, BC clear-sky ARI at typically polluted cities
 varies in the range of approximately $+0.5 - 5.0 \text{ Wm}^{-2}$. While in large fire sites, the daily mean BC clear-sky ARI exceeds
 $+8.0 \text{ Wm}^{-2}$. Generally, with a more compact structure, BC presents a smaller clear-sky ARI at the TOA. The reason can be
 explained from the following aspects. Fractal BC can more efficiently absorb than spherical BC, while the total scattering is
 290 not significantly modified. Thus, the fractal BC leads to a larger positive clear-sky ARI.

Table 1. Time-averaged BC Clear-sky ARI at different sites (Wm^{-2}).

Location	$D_f=1.8$	$D_f=2.2$	$D_f=2.6$	Sphere
Beijing	+1.76	+1.68	+1.63	+1.57
Shanghai	+1.52	+1.45	+1.38	+1.26
Tianjin	+2.00	+1.91	+1.86	+1.77
Nanjing	+1.99	+1.90	+1.83	+1.70
Fire Loc1	+5.39	+5.16	+5.00	+4.69
Fire Loc2	+8.60	+8.27	+8.12	+7.78
Fire Loc3	+5.61	+5.38	+5.22	+4.91

Table 1 shows the time-averaged BC clear-sky ARI (averaging BC clear-sky ARI over the simulation period) in typical
 polluted urban sites and fire sites. Using the spherical model, the time-averaged clear-sky ARI is $+1.26 - +1.77 \text{ Wm}^{-2}$ in

Table 2. Time-averaged BC All-sky ARI at different sites (Wm^{-2}).

Location	$D_f=1.8$	$D_f=2.2$	$D_f=2.6$	Sphere
Beijing	+1.77	+1.70	+1.68	+1.67
Shanghai	+2.26	+2.18	+2.15	+2.15
Tianjin	+2.00	+1.93	+1.90	+1.88
Nanjing	+2.79	+2.69	+2.66	+2.66
Fire Loc1	+5.14	+4.93	+4.81	+4.62
Fire Loc2	+7.38	+7.11	+6.99	+6.77
Fire Loc3	+5.22	+5.02	+4.89	+4.69

typical polluted urban cities, while it increases to $+1.52 - +2.00 \text{ Wm}^{-2}$ using a fractal aggregate model with a D_f of 1.8. The relative differences between the fractal model with a D_f of 1.8 and spherical model $((\text{Fractal} - \text{Sphere})/\text{Sphere})$ can reach approximately 12.1% – 20.6% in typical urban cities. In fire sites, when modifying BC structure from a sphere to a fractal aggregate with a D_f of 1.8, the time-averaged BC clear-sky ARI increases from $+4.69 - +7.78 \text{ Wm}^{-2}$, to $+5.39 - +8.60 \text{ Wm}^{-2}$, and the relative differences between the two models are 10.5% – 14.9%. Lu et al. (2020) showed that BC shapes can introduce approximately 5% relative uncertainties in eastern China using different measured BC profiles. However, our results show that much larger uncertainties can be introduced from BC morphologies. The reason is that $D_f = 2.8$ was assumed for BC aggregates in the study of Lu et al. (2020), which are close to spherical shape. This D_f value is larger than the observed D_f . Besides, due to different solar zenith angles, our results show the ARI uncertainties caused by BC morphologies may vary in different regions. Therefore, the BC morphological effects on the BC ARI should be carefully considered in different regions.

IPCC (2014) has reported a global mean BC all-sky ARI of $+0.6 \text{ W m}^{-2}$, while Wang et al. (2014) estimated a smaller all-sky ARI of $+0.13 \text{ W m}^{-2}$ based on the constraints from the mass and absorption observations. Tuccella et al. (2020b) further showed that the global mean BC all-sky ARI is in the range of $+0.13$ and $+0.25 \text{ W m}^{-2}$. However, the BC all-sky ARI in some specific regions is relatively large. Based on in-situ measurements, Lamb et al. (2018) have estimated the mean column all-sky BC ARI to be $+0.48$ to $+2.01 \text{ Wm}^{-2}$ over South Korea. In Beijing, Sun et al. (2022) found that the mean BC all-sky ARI decreased from $+3.36 \text{ W m}^{-2}$ in 2012 to $+1.09 \text{ W m}^{-2}$ in 2020. The daily-mean all-sky ARI in typical urban polluted cities in eastern China and fire sites in the northwest US are shown in Figure 13. The daily-mean all-sky ARI estimated in this study generally varies in the range of approximately $+0.2 - +4.5 \text{ Wm}^{-2}$ in typical polluted cities. Table 2 shows the time-averaged BC all-sky ARI (averaging BC all-sky ARI over the simulation period) in different sites. The time-averaged all-sky ARI are $+1.67 - +2.79 \text{ Wm}^{-2}$ in urban cities, and it is relatively high compared to the regional-mean all-sky ARI due to the high BC emissions. Our estimated BC all-sky ARI is generally in the range reported by Sun et al. (2022).

The differences of all-sky ARI between the fractal model and spherical model are smaller than those of clear-sky ARI in typical polluted cities in eastern China. The relative differences in the time-averaged all-sky ARI between fractal model and the spherical model are below 6.5% in typical polluted cities. The all-sky ARI at fire sites in the northwest US is smaller than

the clear-sky ARI. The time-averaged all-sky ARI are $+4.62 - +7.38 \text{ Wm}^{-2}$ in typical fire sites. The relative differences in the time-averaged all-sky ARI between the fractal model with a D_f of 1.8 and the spherical model are 9.0% – 11.3% in typical
320 fire sites, which is relatively smaller than the differences for clear-sky ARI.

The regional-mean ARI are shown in Figures 14 – 15. As shown in Figure 14, the BC clear-sky ARI of exceeding $+3.0 \text{ W/m}^2$ is observed in eastern China. With a spherical BC model, the regional-mean clear-sky ARI in eastern China is estimated as $+1.35 \text{ Wm}^{-2}$, and it agrees well with the reported regional-mean clear-sky ARI of $+1.34 \text{ Wm}^{-2}$ in East Asia by Zhuang et al. (2019). BC morphologies also have a non-ignorable impact on the BC ARI. The regional-mean ARI deviations between
325 the fractal model and spherical model are $+0.23$, $+0.15$, and $+0.1 \text{ Wm}^{-2}$ when D_f is 1.8, 2.2, and 2.6, respectively, and the relative differences are 17.1%, 11.1%, and 7.4%, respectively.

The BC morphologies have a relatively small impact on the all-sky ARI. As shown in the lower panels of Figure 14, the all-sky ARI is generally larger than the clear-sky ARI, while the deviations between the fractal model and spherical model are smaller compared to those for clear-sky. The BC all-sky ARI can exceed $+4.0 \text{ W/m}^2$ in eastern China. A relatively larger
330 regional-mean all-sky ARI is observed than clear-sky ARI, which is $+1.79 \text{ Wm}^{-2}$ in eastern China when using the spherical model. The regional-mean all-sky ARI differences between the fractal model and spherical model are 0.13, 0.06, 0.03 Wm^{-2} when D_f is 1.8, 2.2, and 2.6, respectively. The relative differences for all-sky ARI between the two models are less than 7.3%.

The regional-mean clear-sky ARI in the fire region in the northwest US is shown in Figure 15. The clear-sky ARI in the fire sites is obviously larger than the other sites. The BC clear-sky ARI of exceeding $+5.0 \text{ W/m}^2$ is observed, and the regional-mean
335 clear-sky ARI is $+0.93 \text{ W/m}^2$. The differences of regional-mean clear-sky ARI between the fractal model and the spherical model in the fire region in the northwest US are more substantial than those in eastern China, which reach approximately $+0.36$, $+0.29$, and $+0.19 \text{ W/m}^2$ when D_f is 1.8, 2.2, and 2.6, respectively. The relative differences between the fractal model and spherical model are 38.7%, 31.2%, and 20.4% when D_f is 1.8, 2.2, and 2.6, respectively. Smaller all-sky ARI than clear-sky ARI are observed at the sites where fire occurs, while the regional-mean all-sky ARI in the fire region in the northwest US
340 is generally larger than clear-sky ARI, which is $+1.67 \text{ W/m}^2$. The differences between the two models are $+0.28$, $+0.19$, and $+0.11 \text{ W/m}^2$ when D_f is 1.8, 2.2, and 2.6, respectively, and the relative differences are 16.8%, 11.4%, and 6.6%.

4 Discussion

In current climate models, such as CESM, MIROC-SPRINTARS, and WRF-Chem, the Mie theory was commonly used to calculate the optical properties of BC aerosols. However, fractal-like BC aerosols were often observed in the atmosphere. In
345 this work, we found that the effects of BC morphology are spatially-dependent. Compared to the spherical BC model, the fractal BC model generally presents a larger clear-sky ARI, which may lead to the underestimations of BC ARI in the climate models. The relative differences in the time-averaged clear-sky ARI are 12.1% – 20.6% and 10.5% – 14.9% in typical polluted urban cities and fire sites, respectively. Furthermore, the regional-mean clear-sky ARI is also significantly affected by the BC morphology, and relative differences of 17.1% and 38.7% between the fractal model were observed in eastern China and in the
350 northwest US, respectively, while the existence of cloud would weaken the BC morphological effects. The results imply that

current climate modeling may significantly underestimate the BC ARI uncertainties as the morphological effects on BC ARI are ignored in most climate models.

355 However, this work is by no means exhaustive. This work assumed that BC aerosols are externally mixed with other chemical components, while BC aerosols are often internally mixed with other components, such as organic aerosols, sulfate, etc (China et al., 2013a; Adachi et al., 2010; Wang et al., 2021b). BC absorption can be significantly enhanced by the "lensing Effect" even if BC aerosols are internally mixed with non-absorbing materials, which may lead to larger BC ARI (Chung et al., 2012; Liu et al., 2017a). Previous studies have shown that the morphologies of internally mixed BC would significantly affect its absorption enhancement (Luo et al., 2019; Wang et al., 2021a; Luo et al., 2021c), so lead to larger uncertainties in the estimation of BC ARI. Thus, the sensitivities of BC morphologies on the ARI estimated in this work may be smaller than those
360 in real cases.

Furthermore, we found that the spherical assumption generally underestimates the clear-sky ARI for externally mixed BC, while opposite phenomenon may be found for internally mixed BC. A core-shell spherical morphology was widely used to represent the internally mixed BC. However, many partially coated BC aerosols exist in the atmosphere, while the core-shell spherical BC model commonly assumes the BC is fully embedded in a coating shell. The core-shell morphology may
365 overestimate the absorption of partially coated BC (Wang et al., 2021a; Zhang et al., 2018), so overestimate the ARI. Thus, the ARI of internally mixed BC with complex morphologies should be further investigated in the future.

5 Summary and Conclusions

The current climate modeling commonly assumes a spherical morphology for BC, while the fractal structure is more realistic than the spherical morphology for externally mixed BC. In this work, we used the fractal model to re-evaluate the BC ARI
370 in a typical polluted urban area in eastern China and a fire region in the northwest US. We found that BC morphologies have non-ignorable impacts on the aerosol optical properties. At 550 nm wavelength, the spherical BC model can overestimate the AOD up to 0.03 and 0.15 in typical polluted cities in China and fire sites in US, respectively. The overestimations roughly account for 20% of the total BC AOD. Besides, the spherical BC model underestimates BC AAOD at 450 nm up to 0.016 and 0.04 at typical polluted cities in China and fire sites in the US, respectively, compared to the fractal model with a D_f of 1.8.
375 The underestimations account for approximately 8% of the total BC AAOD.

Both the morphological effects on clear-sky and all-sky ARI are evaluated. With a spherical BC model, the estimated time-averaged clear-sky ARI is generally in the range of $+1.26 - +1.77 \text{ W/m}^2$ in typical urban polluted cities in eastern China, while this range increases to approximately $+1.52 - +2.00 \text{ W/m}^2$ when using the fractal model with a D_f of 1.8. The clear-sky ARI relative differences between the two models are approximately 12.1% – 20.6% in typical urban polluted cities. In fire sites,
380 when modifying BC structure from a sphere to a fractal aggregate, the time-averaged BC clear-sky ARI increases from $+4.69 - +7.78 \text{ Wm}^{-2}$ to $+5.39 - +8.60 \text{ Wm}^{-2}$ in typical fire sites. The relative differences between the two models are approximately 10.5% – 14.9% in typical fire sites. The existence of clouds weaken the effects of BC morphologies on the ARI. The all-sky

ARI relative differences between the fractal model and the spherical model are approximately 4.9% – 6.4% and 9.0% – 11.3% in typical urban polluted cities and typical fire sites, respectively, which is relatively smaller than those of clear-sky ARI.

385 The impacts of BC morphologies on the regional-mean ARI were also evaluated. The regional-mean clear-sky ARI was estimated as $+1.35 \text{ Wm}^{-2}$ and $+0.93 \text{ Wm}^{-2}$ in the polluted urban area and the fire region, respectively, using the spherical BC model. The regional-mean clear-sky ARI differences between the fractal model and the spherical model are approximately $+0.23$ and $+0.36 \text{ Wm}^{-2}$ in these two regions, respectively, and the relative differences between the two models are approximately 17.1% and 38.7%, respectively. The all-sky ARI differences between the fractal model and the spherical model are relatively smaller. The relative differences in the regional-mean all-sky ARI between the fractal model and the spherical model are less than 7.3% and 16.8% in these two regions, respectively. Thus, the effects of BC morphologies on the ARI should be
390 carefully considered in different regions.

Acknowledgements. This work was financially supported by the National Natural Science Foundation of China (Grant No. 41925019), the National Key Research and Development Plan under Grant No. 2020YFC1511600, and the Fundamental Research Funds for the Central Universities under Grant No. WK2320000052. RKC and CZ acknowledge support from the US National Science Foundation (AGS-1926817) and the NASA ACCDAM program (NNH20ZDA001N). CZ would like to acknowledge partial support received from the McDonnell International Scholars Academy at Washington University in St. Louis.).
395

References

- Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S., and Shankar, U.: Modal aerosol dynamics model for Europe: Development and first applications, *Atmospheric environment*, 32, 2981–2999, 1998.
- 400 Adachi, K., Chung, S. H., and Buseck, P. R.: Shapes of soot aerosol particles and implications for their effects on climate, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/https://doi.org/10.1029/2009JD012868>, 2010.
- Adachi, K., Zaizen, Y., Kajino, M., and Igarashi, Y.: Mixing state of regionally transported soot particles and the coating effect on their size and shape at a mountain site in Japan, *Journal of Geophysical Research: Atmospheres*, 119, 5386–5396, <https://doi.org/https://doi.org/10.1002/2013JD020880>, 2014.
- 405 Alexander, D. T., Crozier, P. A., and Anderson, J. R.: Brown carbon spheres in East Asian outflow and their optical properties, *Science*, 321, 833–836, 2008.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol science and technology*, 40, 27–67, 2006.
- 410 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Karcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research-Atmospheres*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- 415 Buras, R., Dowling, T., and Emde, C.: New secondary-scattering correction in DISORT with increased efficiency for forward scattering, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112, 2028–2034, 2011.
- Buseck, S. E. and Buseck, P. R.: Absorbing phenomena (vol 288, pg 989, 2000), *Science*, 289, 58–58, <GotoISI>://WOS:000088090400023, 2000.
- Chakrabarty, R. K., Moosmuller, H., Garro, M. A., Arnott, W. P., Walker, J., Susott, R. A., Babbitt, R. E., Wold, C. E., Lincoln, E. N., and 420 Hao, W. M.: Emissions from the laboratory combustion of wildland fuels: Particle morphology and size, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/https://doi.org/10.1029/2005JD006659>, 2006.
- Chakrabarty, R. K., Moosmüller, H., Arnott, W. P., Garro, M. A., Slowik, J. G., Cross, E. S., Han, J.-H., Davidovits, P., Onasch, T. B., and Worsnop, D. R.: Light scattering and absorption by fractal-like carbonaceous chain aggregates: Comparison of theories and experiment, *Applied optics*, 46, 6990–7006, 2007.
- 425 Chang, H.-c. and Charalampopoulos, T.: Determination of the wavelength dependence of refractive indices of flame soot, *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, 430, 577–591, 1990.
- Chen, C., Fan, X., Shaltout, T., Qiu, C., Ma, Y., Goldman, A., and Khalizov, A. F.: An unexpected restructuring of combustion soot aggregates by subnanometer coatings of polycyclic aromatic hydrocarbons, *Geophysical Research Letters*, 43, 11,080–11,088, <https://doi.org/https://doi.org/10.1002/2016GL070877>, 2016.
- 430 Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: "Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements", *Journal of the Atmospheric Sciences*, 59, 461–483, [https://doi.org/10.1175/1520-0469\(2002\)059<0461:TAOTFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0461:TAOTFT>2.0.CO;2), 2002.
- China, S., Mazzoleni, C., Gorkowski, K., Aiken, A. C., and Dubey, M. K.: Morphology and mixing state of individual freshly emitted wildfire carbonaceous particles, *Nature Communications*, 4, 2122, <https://doi.org/10.1038/ncomms3122>, 2013a.

- 435 China, S., Mazzoleni, C., Gorkowski, K., Aiken, A. C., and Dubey, M. K.: Morphology and mixing state of individual freshly emitted wildfire carbonaceous particles, *Nature communications*, 4, 1–7, 2013b.
- Chung, C., Lee, K., and Müller, D.: Effect of internal mixture on black carbon radiative forcing, *Tellus B: Chemical and Physical Meteorology*, 64, 10925, <https://doi.org/10.3402/tellusb.v64i0.10925>, 2012.
- Coz, E. and Leck, C.: Morphology and state of mixture of atmospheric soot aggregates during the winter season over Southern Asia—a
440 quantitative approach, *Tellus B: Chemical and Physical Meteorology*, 63, 107–116, 2011.
- Cross, E. S., Slowik, J. G., Davidovits, P., Allan, J. D., Worsnop, D. R., Jayne, J. T., Lewis, D. K., Canagaratna, M., and Onasch, T. B.: Laboratory and ambient particle density determinations using light scattering in conjunction with aerosol mass spectrometry, *Aerosol Science and Technology*, 41, 343–359, 2007.
- Curci, G.: FlexAOD: A chemistry-transport model post-processing tool for a flexible calculation of aerosol optical properties, in: Proceedings
445 of the 9th International Symposium on Tropospheric Profiling, pp. 978–90, 2012.
- Curci, G., Hogrefe, C., Bianconi, R., Im, U., Balzarini, A., Baró, R., Brunner, D., Forkel, R., Giordano, L., Hirtl, M., et al.: Uncertainties of simulated aerosol optical properties induced by assumptions on aerosol physical and chemical properties: An AQMEII-2 perspective, *Atmospheric Environment*, 115, 541–552, 2015.
- Curci, G., Alyuz, U., Barò, R., Bianconi, R., Bieser, J., Christensen, J. H., Colette, A., Farrow, A., Francis, X., Jiménez-Guerrero, P., Im, U.,
450 Liu, P., Manders, A., Palacios-Peña, L., Prank, M., Pozzoli, L., Sokhi, R., Solazzo, E., Tuccella, P., Unal, A., Vivanco, M. G., Hogrefe, C., and Galmarini, S.: Modelling black carbon absorption of solar radiation: combining external and internal mixing assumptions, *Atmospheric Chemistry and Physics*, 19, 181–204, <https://doi.org/10.5194/acp-19-181-2019>, 2019.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb,
455 W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Versteijn, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/https://doi.org/10.1029/2019MS001916>, e2019MS001916 2019MS001916, 2020.
- 460 Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J., Ito, A., et al.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmospheric Chemistry and Physics*, 6, 4321–4344, 2006.
- Eggersdorfer, M. L. and Pratsinis, S. E.: The structure of agglomerates consisting of polydisperse particles, *Aerosol Science and Technology*, 46, 347–353, 2012.
- 465 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of
470 ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/https://doi.org/10.1029/2005JD006721>, 2006.

- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled "online" chemistry within the WRF model, *Atmospheric Environment*, 39, 6957–6975, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- 475 Guenther, A., Zimmerman, P., and Wildermuth, M.: Natural volatile organic compound emission rate estimates for US woodland landscapes, *Atmospheric Environment*, 28, 1197–1210, 1994.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, 2006.
- He, C., Liou, K.-N., Takano, Y., Zhang, R., Levy Zamora, M., Yang, P., Li, Q., and Leung, L. R.: Variation of the radiative properties during black carbon aging: theoretical and experimental intercomparison, *Atmospheric Chemistry and Physics*, 15, 11 967–11 980, 480 <https://doi.org/10.5194/acp-15-11967-2015>, 2015.
- He, C., Li, Q., Liou, K.-N., Qi, L., Tao, S., and Schwarz, J. P.: Microphysics-based black carbon aging in a global CTM: constraints from HIPPO observations and implications for global black carbon budget, *Atmos. Chem. Phys.*, 16, 3077–3098, 2016.
- Heinson, W. R., Sorensen, C. M., and Chakrabarti, A.: Does Shape Anisotropy Control the Fractal Dimension in Diffusion-Limited Cluster-Cluster Aggregation?, *Aerosol Science and Technology*, 44, i–iv, <https://doi.org/10.1080/02786826.2010.516032>, 2010.
- 485 Heinson, W. R., Liu, P., and Chakrabarty, R. K.: Fractal scaling of coated soot aggregates, *Aerosol Science and Technology*, 51, 12–19, <https://doi.org/10.1080/02786826.2016.1249786>, 2017.
- Hess, M., Koepke, P., and Schult, I.: "Optical Properties of Aerosols and Clouds: The Software Package OPAC", *Bulletin of the American Meteorological Society*, 79, 831–844, [https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2), 1998.
- Highwood, E. J.: Suggested Refractive Indices and Aerosol Size Parameters for Use in Radiative Effect Calculations and Satellite Retrievals, ADIENT/APPRaise CP2 Technical Report, DRAFT V2 (5 August 2009), <http://www.met.rdg.ac.uk/~adient/refractiveindices.html>, last 490 access: 3 January 2019, 2009.
- IPCC: Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment report of the Intergovernmental Panel on Climate Change, Cambridge university press, 2014.
- Kahnert, M.: On the Discrepancy between Modeled and Measured Mass Absorption Cross Sections of Light Absorbing Carbon Aerosols, 495 *Aerosol Science and Technology*, 44, 453–460, <GotoISI>://WOS:000277436300006, 2010a.
- Kahnert, M.: Modelling the optical and radiative properties of freshly emitted light absorbing carbon within an atmospheric chemical transport model, *Atmospheric Chemistry and Physics*, 10, 1403–1416, <GotoISI>://WOS:000274410000036, 2010b.
- Kahnert, M. and Devasthale, A.: Black carbon fractal morphology and short-wave radiative impact: a modelling study, *Atmospheric Chemistry and Physics*, 11, 11 745–11 759, <https://doi.org/10.5194/acp-11-11745-2011>, 2011.
- 500 Kahnert, M. and Kanngießer, F.: Modelling optical properties of atmospheric black carbon aerosols, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 244, 106 849, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2020.106849>, 2020.
- Köylü, Ü. and Faeth, G.: Structure of overfire soot in buoyant turbulent diffusion flames at long residence times, *Combustion and Flame*, 89, 140–156, [https://doi.org/https://doi.org/10.1016/0010-2180\(92\)90024-J](https://doi.org/https://doi.org/10.1016/0010-2180(92)90024-J), 1992.
- Lack, D. and Cappa, C.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength 505 dependence of black carbon, *Atmospheric Chemistry and Physics*, 10, 4207–4220, 2010.
- Lamb, K. D., Perring, A. E., Samset, B., Peterson, D., Davis, S., Anderson, B. E., Beyersdorf, A., Blake, D. R., Campuzano-Jost, P., Corr, C. A., Diskin, G. S., Kondo, Y., Moteki, N., Nault, B. A., Oh, J., Park, M., Pusede, S. E., Simpson, I. J., Thornhill, K. L., Wisthaler, A., and Schwarz, J. P.: Estimating Source Region Influences on Black Carbon Abundance, Microphysics, and Radiative Effect Observed Over South Korea, *Journal of Geophysical Research: Atmospheres*, 123, 13,527–13,548, <https://doi.org/10.1029/2018JD029257>, 2018.

- 510 Lee, K. O., Cole, R., Sekar, R., Choi, M. Y., Kang, J. S., Bae, C. S., and Shin, H. D.: Morphological investigation of the microstructure, dimensions, and fractal geometry of diesel particulates, *Proceedings of the Combustion Institute*, 29, 647–653, [https://doi.org/https://doi.org/10.1016/S1540-7489\(02\)80083-9](https://doi.org/https://doi.org/10.1016/S1540-7489(02)80083-9), proceedings of the Combustion Institute, 2002.
- Li, J., Anderson, J. R., and Buseck, P. R.: TEM study of aerosol particles from clean and polluted marine boundary layers over the North Atlantic, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/https://doi.org/10.1029/2002JD002106>, 2003.
- 515 Li, M., Zhang, Q., Streets, D., He, K., Cheng, Y., Emmons, L., Huo, H., Kang, S., Lu, Z., Shao, M., et al.: Mapping Asian anthropogenic emissions of non-methane volatile organic compounds to multiple chemical mechanisms, *Atmos. Chem. Phys.*, 14, 5617–5638, 2014.
- Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmospheric Chemistry and Physics*, 17, 935–963, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-17-935-2017)
- 520 17-935-2017, 2017a.
- Li, M., Zhang, Q., Kurokawa, J.-i., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., et al.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmospheric Chemistry and Physics (Online)*, 17, 2017b.
- Liu, C., Yin, Y., Hu, F., Jin, H., and Sorensen, C. M.: The Effects of Monomer Size Distribution on the Radiative Properties of Black Carbon Aggregates, *Aerosol Science and Technology*, 49, 928–940, <https://doi.org/10.1080/02786826.2015.1085953>, 2015a.
- 525 Liu, C., Chung, C. E., Yin, Y., and Schnaiter, M.: The absorption Ångström exponent of black carbon: from numerical aspects, *Atmospheric Chemistry and Physics*, 18, 6259–6273, 2018.
- Liu, D., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C., Kong, S., Williams, P., Ting, Y.-C., Haslett, S., Taylor, J., Flynn, M. J., Morgan, W., McFiggans, G., Coe, H., and Allan, J.: Black-carbon absorption enhancement in the atmosphere
- 530 determined by particle mixing state, *Nature Geoscience*, 10, 184–188, <https://doi.org/10.1038/ngeo2901>, 2017a.
- Liu, D. T., Whitehead, J., Alfarra, M. R., Reyes-Villegas, E., Spracklen, D. V., Reddington, C. L., Kong, S. F., Williams, P. I., Ting, Y. C., Haslett, S., Taylor, J. W., Flynn, M. J., Morgan, W. T., McFiggans, G., Coe, H., and Allan, J. D.: Black-carbon absorption enhancement in the atmosphere determined by particle mixing state, *Nature Geoscience*, 10, 184–U132, <GotoISI>://WOS:000395791400009, 2017b.
- Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., and He, K.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, *Atmospheric Chemistry and Physics*, 15, 13 299–13 317, 2015b.
- 535 Liu, L. and Mishchenko, M. I.: Effects of aggregation on scattering and radiative properties of soot aerosols, *Journal of Geophysical Research-Atmospheres*, 110, <GotoISI>://WOS:000229988800009, 2005.
- Lu, Q., Liu, C., Zhao, D., Zeng, C., Li, J., Lu, C., Wang, J., and Zhu, B.: "Atmospheric heating rate due to black carbon aerosols: Uncertainties and impact factors", *Atmospheric Research*, 240, 104 891, <https://doi.org/https://doi.org/10.1016/j.atmosres.2020.104891>, 2020.
- 540 Luo, J., Zhang, Y., Wang, F., Wang, J., and Zhang, Q.: Applying machine learning to estimate the optical properties of black carbon fractal aggregates, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 215, 1 – 8, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2018.05.002>, 2018a.
- Luo, J., Zhang, Y., Wang, F., and Zhang, Q.: Effects of brown coatings on the absorption enhancement of black carbon: a numerical investigation, *Atmospheric Chemistry and Physics*, 18, 16 897–16 914, <https://doi.org/10.5194/acp-18-16897-2018>, 2018b.
- 545 Luo, J., Zhang, Y., and Zhang, Q.: A model study of aggregates composed of spherical soot monomers with an acentric carbon shell, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 205, 184 – 195, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2017.10.024>, 2018c.

- Luo, J., Zhang, Y. M., Zhang, Q. X., Wang, F., Liu, J., and Wang, J. J.: Sensitivity analysis of morphology on radiative properties of soot aerosols, *Optics Express*, 26, A420–A432, <GotoISI>://WOS:000432457600005, 2018d.
- 550 Luo, J., Zhang, Q., Luo, J., Liu, J., Huo, Y., and Zhang, Y.: Optical Modeling of Black Carbon With Different Coating Materials: The Effect of Coating Configurations, *Journal of Geophysical Research: Atmospheres*, 124, 13 230–13 253, <https://doi.org/10.1029/2019JD031701>, 2019.
- Luo, J., Zhang, Q., Zhang, C., Zhang, Y., and Chakrabarty, R. K.: The fractal characteristics of atmospheric coated soot: Implication for morphological analysis, *Journal of Aerosol Science*, 157, 105 804, <https://doi.org/https://doi.org/10.1016/j.jaerosci.2021.105804>, 2021a.
- 555 Luo, J., Zhang, Q., Zhang, Y., and Li, Z.: Radiative Properties of Non-spherical Black Carbon Aerosols, pp. 69–124, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-030-87683-8_3, 2021b.
- Luo, J., Zhang, Y., and Zhang, Q.: Effects of black carbon morphology on brown carbon absorption estimation: from numerical aspects, *Geoscientific Model Development*, 14, 2113–2126, <https://doi.org/10.5194/gmd-14-2113-2021>, 2021c.
- Mackowski, D. W. and Mishchenko, M. I.: Calculation of the T matrix and the scattering matrix for ensembles of spheres, *Journal of the Optical Society of America a-Optics Image Science and Vision*, 13, 2266–2278, <GotoISI>://WOS:A1996VP77000015, 1996.
- 560 Mackowski, D. W. and Mishchenko, M. I.: A multiple sphere T-matrix Fortran code for use on parallel computer clusters, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112, 2182–2192, <GotoISI>://WOS:000294518300013, 2011.
- Martin, R. V., Jacob, D. J., Yantosca, R. M., Chin, M., and Ginoux, P.: Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Mayer, B. and Kylling, A.: The libRadtran software package for radiative transfer calculations-description and examples of use, *Atmospheric Chemistry and Physics*, 5, 1855–1877, 2005.
- 565 Mikhailov, E., Vlasenko, S., Podgorny, I., Ramanathan, V., and Corrigan, C.: Optical properties of soot–water drop agglomerates: An experimental study, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Mishchenko, M. I., Dlugach, J. M., Yanovitskij, E. G., and Zakharova, N. T.: Bidirectional reflectance of flat, optically thick particulate layers: an efficient radiative transfer solution and applications to snow and soil surfaces, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 63, 409–432, 1999.
- 570 Mishchenko, M. I., Travis, L. D., and Lacis, A. A.: Scattering, absorption, and emission of light by small particles, Cambridge university press, 2002.
- Mishchenko, M. I., Dlugach, J. M., and Liu, L.: Linear depolarization of lidar returns by aged smoke particles, *Applied optics*, 55, 9968–9973, 2016a.
- 575 Mishchenko, M. I., Dlugach, J. M., Yurkin, M. A., Bi, L., Cairns, B., Liu, L., Panetta, R. L., Travis, L. D., Yang, P., and Zakharova, N. T.: "First-principles modeling of electromagnetic scattering by discrete and discretely heterogeneous random media", *Physics Reports*, 632, 1 – 75, <https://doi.org/https://doi.org/10.1016/j.physrep.2016.04.002>, 2016b.
- Moosmüller, H., Chakrabarty, R., and Arnott, W.: Aerosol light absorption and its measurement: A review, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 110, 844–878, 2009.
- 580 Nakao, S., Tang, P., Tang, X., Clark, C. H., Qi, L., Seo, E., Asa-Awuku, A., and Cocker III, D.: Density and elemental ratios of secondary organic aerosol: Application of a density prediction method, *Atmospheric environment*, 68, 273–277, 2013.
- Rajesh, T. and Ramachandran, S.: Black carbon aerosols over urban and high altitude remote regions: Characteristics and radiative implications, *Atmospheric environment*, 194, 110–122, 2018.

- 585 Reddington, C. L., McMeeking, G., Mann, G. W., Coe, H., Frontoso, M. G., Liu, D., Flynn, M., Spracklen, D. V., and Carslaw, K. S.: The mass and number size distributions of black carbon aerosol over Europe, *Atmospheric Chemistry and Physics*, 13, 4917–4939, 2013.
- Saleh, R., Marks, M., Heo, J., Adams, P. J., Donahue, N. M., and Robinson, A. L.: Contribution of brown carbon and lensing to the direct radiative effect of carbonaceous aerosols from biomass and biofuel burning emissions, *Journal of Geophysical Research: Atmospheres*, 120, 10,285–10,296, <https://doi.org/https://doi.org/10.1002/2015JD023697>, 2015.
- 590 Schwarz, J., Gao, R., Spackman, J., Watts, L., Thomson, D., Fahey, D., Ryerson, T., Peischl, J., Holloway, J., Trainer, M., et al.: Measurement of the mixing state, mass, and optical size of individual black carbon particles in urban and biomass burning emissions, *Geophysical Research Letters*, 35, 2008.
- Seinfeld, J. H., Erdakos, G. B., Asher, W. E., and Pankow, J. F.: Modeling the formation of secondary organic aerosol (SOA). 2. The predicted effects of relative humidity on aerosol formation in the α -pinene-, β -pinene-, sabinene-, Δ 3-carene-, and cyclohexene-ozone systems, *Environmental Science & Technology*, 35, 1806–1817, 2001.
- 595 Shaheen, K., Shah, Z., Suo, H. L., Liu, M., Ma, L., Alam, K., Gul, A., Cui, J., Li, C. Y., Wang, Y., Khan, S. A., and Khan, S. B.: Aerosol clustering in an urban environment of Beijing during (2005–2017), *Atmospheric Environment*, 213, 534–547, <https://doi.org/10.1016/j.atmosenv.2019.06.027>, 2019.
- Shin, S.-K., Tesche, M., Müller, D., and Noh, Y.: Technical note: Absorption aerosol optical depth components from AERONET observations of mixed dust plumes, *Atmospheric Measurement Techniques*, 12, 607–618, <https://doi.org/10.5194/amt-12-607-2019>, 2019a.
- 600 Shin, S.-K., Tesche, M., Noh, Y., and Müller, D.: Aerosol-type classification based on AERONET version 3 inversion products, *Atmospheric Measurement Techniques*, 12, 3789–3803, <https://doi.org/10.5194/amt-12-3789-2019>, 2019b.
- Sorensen, C.: The mobility of fractal aggregates: a review, *Aerosol Science and Technology*, 45, 765–779, 2011.
- Stamnes, K., Tsay, S.-C., Wiscombe, W., and Jayaweera, K.: Numerically stable algorithm for discrete-ordinate-method radiative transfer in multiple scattering and emitting layered media, *Applied optics*, 27, 2502–2509, 1988.
- 605 Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling, *Journal of Geophysical Research: Atmospheres*, 95, 16 343–16 367, <https://doi.org/10.1029/JD095iD10p16343>.
- Streets, D. G., Gupta, S., Waldhoff, S. T., Wang, M. Q., Bond, T. C., and Yiyun, B.: "Black carbon emissions in China", *Atmospheric Environment*, 35, 4281 – 4296, [https://doi.org/https://doi.org/10.1016/S1352-2310\(01\)00179-0](https://doi.org/https://doi.org/10.1016/S1352-2310(01)00179-0), 2001.
- 610 Streets, D. G., Wu, Y., and Chin, M.: Two-decadal aerosol trends as a likely explanation of the global dimming/brightening transition, *Geophysical Research Letters*, 33, <GotoISI>://WOS:000239597000006, 2006.
- Sun, J., Wang, Z., Zhou, W., Xie, C., Wu, C., Chen, C., Han, T., Wang, Q., Li, Z., Li, J., Fu, P., Wang, Z., and Sun, Y.: Measurement report: Long-term changes in black carbon and aerosol optical properties from 2012 to 2020 in Beijing, China, *Atmospheric Chemistry and Physics*, 22, 561–575, <https://doi.org/10.5194/acp-22-561-2022>, 2022.
- 615 Takemura, T., Nozawa, T., Emori, S., Nakajima, T. Y., and Nakajima, T.: Simulation of climate response to aerosol direct and indirect effects with aerosol transport-radiation model, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2004JD005029>, 2005.
- Takemura, T., Egashira, M., Matsuzawa, K., Ichijo, H., O'ishi, R., and Abe-Ouchi, A.: A simulation of the global distribution and radiative forcing of soil dust aerosols at the Last Glacial Maximum, *Atmospheric Chemistry and Physics*, 9, 3061–3073, <https://doi.org/10.5194/acp-9-3061-2009>, 2009.
- 620

- Teng, S., Liu, C., Schnaiter, M., Chakrabarty, R. K., and Liu, F.: Accounting for the effects of nonideal minor structures on the optical properties of black carbon aerosols, *Atmospheric Chemistry and Physics*, 19, 2917–2931, <https://doi.org/10.5194/acp-19-2917-2019>, 2019.
- 625 Tuccella, P., Curci, G., Pitari, G., Lee, S., and Jo, D. S.: Direct Radiative Effect of Absorbing Aerosols: Sensitivity to Mixing State, Brown Carbon, and Soil Dust Refractive Index and Shape, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030967, <https://doi.org/https://doi.org/10.1029/2019JD030967>, e2019JD030967 2019JD030967, 2020a.
- Tuccella, P., Curci, G., Pitari, G., Lee, S., and Jo, D. S.: Direct Radiative Effect of Absorbing Aerosols: Sensitivity to Mixing State, Brown Carbon, and Soil Dust Refractive Index and Shape, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030967, <https://doi.org/https://doi.org/10.1029/2019JD030967>, e2019JD030967 2019JD030967, 2020b.
- 630 Wang, X., Heald, C. L., Ridley, D. A., Schwarz, J. P., Spackman, J. R., Perring, A. E., Coe, H., Liu, D., and Clarke, A. D.: Exploiting simultaneous observational constraints on mass and absorption to estimate the global direct radiative forcing of black carbon and brown carbon, *Atmospheric Chemistry and Physics*, 14, 10989–11010, <https://doi.org/10.5194/acp-14-10989-2014>, 2014.
- Wang, Y., Liu, F., He, C., Bi, L., Cheng, T., Wang, Z., Zhang, H., Zhang, X., Shi, Z., and Li, W.: Fractal dimensions and mixing structures of soot particles during atmospheric processing, *Environmental Science & Technology Letters*, 4, 487–493, 2017.
- 635 Wang, Y., Li, W., Huang, J., Liu, L., Pang, Y., He, C., Liu, F., Liu, D., Bi, L., Zhang, X., and Shi, Z.: Nonlinear Enhancement of Radiative Absorption by Black Carbon in Response to Particle Mixing Structure, *Geophysical Research Letters*, 48, e2021GL096437, <https://doi.org/https://doi.org/10.1029/2021GL096437>, e2021GL096437 2021GL096437, 2021a.
- Wang, Y., Pang, Y., Huang, J., Bi, L., Che, H., Zhang, X., and Li, W.: Constructing Shapes and Mixing Structures of Black Carbon Particles With Applications to Optical Calculations, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034620, <https://doi.org/https://doi.org/10.1029/2021JD034620>, e2021JD034620 2021JD034620, 2021b.
- 640 Wang, Y., Pang, Y., Huang, J., Bi, L., Che, H., Zhang, X., and Li, W.: Constructing shapes and mixing structures of black carbon particles with applications to optical calculations, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034620, 2021c.
- Wentzel, M., Gorzawski, H., Naumann, K.-H., Saathoff, H., and Weinbruch, S.: Transmission electron microscopical and aerosol dynamical characterization of soot aerosols, *Journal of Aerosol Science*, 34, 1347–1370, [https://doi.org/https://doi.org/10.1016/S0021-8502\(03\)00360-4](https://doi.org/https://doi.org/10.1016/S0021-8502(03)00360-4), intercomparison of Soot Measurement Techniques, 2003.
- 645 Wiedinmyer, C., Akagi, S., Yokelson, R. J., Emmons, L., Al-Saadi, J., Orlando, J., and Soja, A.: The Fire INventory from NCAR (FINN): A high resolution global model to estimate the emissions from open burning, *Geoscientific Model Development*, 4, 625, 2011.
- Wild, O., Zhu, X., and Prather, M. J.: Fast-J: Accurate simulation of in-and below-cloud photolysis in tropospheric chemical models, *Journal of Atmospheric Chemistry*, 37, 245–282, 2000.
- 650 Wu, Y., Cheng, T. H., Zheng, L. J., and Chen, H.: A Study of Optical Properties of Soot Aggregates Composed of Poly-Disperse Monomers Using the Superposition T-Matrix Method, *Aerosol Science and Technology*, 49, 941–949, <GotoISI>://WOS:000366400100005, 2015.
- Yin, J. Y. and Liu, L. H.: Influence of complex component and particle polydispersity on radiative properties of soot aggregate in atmosphere, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 111, 2115–2126, <GotoISI>://WOS:000280887400009, 2010.
- Yu, P. F., Toon, O. B., Bardeen, C. G., Zhu, Y. Q., Rosenlof, K. H., Portmann, R. W., Thornberry, T. D., Gao, R. S., Davis, S. M., Wolf, E. T., de Gouw, J., Peterson, D. A., Fromm, M. D., and Robock, A.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume, *Science*, 365, 587–590, <GotoISI>://WOS:000483195200039, 2019.
- 655 Yuan, Q., Xu, J., Wang, Y., Zhang, X., Pang, Y., Liu, L., Bi, L., Kang, S., and Li, W.: Mixing state and fractal dimension of soot particles at a remote site in the southeastern Tibetan plateau, *Environmental science & technology*, 53, 8227–8234, 2019.

- Zaveri, R. A. and Peters, L. K.: A new lumped structure photochemical mechanism for large-scale applications, *Journal of Geophysical Research: Atmospheres*, 104, 30 387–30 415, <https://doi.org/10.1029/1999JD900876>, 1999.
- 660 Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008782>, 2008.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmospheric Chemistry and Physics*, 9, 5131–5153, <https://doi.org/10.5194/acp-9-5131-2009>, 2009.
- 665 Zhang, X., Mao, M., Yin, Y., and Wang, B.: Numerical Investigation on Absorption Enhancement of Black Carbon Aerosols Partially Coated With Nonabsorbing Organics, *Journal of Geophysical Research: Atmospheres*, 123, 1297–1308, <https://doi.org/https://doi.org/10.1002/2017JD027833>, 2018.
- Zhang, X. Y., Wang, Y. Q., Niu, T., Zhang, X. C., Gong, S. L., Zhang, Y. M., and Sun, J. Y.: Atmospheric aerosol compositions in China: spatial/temporal variability, chemical signature, regional haze distribution and comparisons with global aerosols, *Atmospheric Chemistry and Physics*, 12, 779–799, <https://doi.org/10.5194/acp-12-779-2012>, 2012.
- 670 Zhuang, B., Li, S., Wang, T., Liu, J., Chen, H., Chen, P., Li, M., and Xie, M.: Interaction between the black carbon aerosol warming effect and East Asian monsoon using RegCM4, *Journal of Climate*, 31, 9367–9388, 2018.
- Zhuang, B. L., Chen, H. M., Li, S., Wang, T. J., Liu, J., Zhang, L. J., Liu, H. N., Xie, M., Chen, P. L., Li, M. M., and Zhao, M.: The direct effects of black carbon aerosols from different source sectors in East Asia in summer, *Climate Dynamics*, 53, 5293–5310, <https://doi.org/10.1007/s00382-019-04863-5>, 2019.
- 675




Figure 1. Typical morphologies of fractal BC.




Figure 2. Comparison of measured and calculated PM_{2.5} concentrations.




Figure 3. The time series of BC concentrations in typical northwest US and eastern China.




Figure 4. Comparison of measured and calculated AOD and AAOD in Beijing, where $\lambda=500$ nm for AOD, $\lambda=440$ nm and 450 nm for measured and calculated AAOD, respectively, and the observations were obtained from AERONET.




Figure 5. The comparison of BC AOD for different BC morphologies, $\lambda=550$ nm.




Figure 6. The AOD difference between fractal aggregate models and the spherical model, $\lambda=550$ nm.




Figure 7. The comparison of BC AAOD for different BC morphologies, $\lambda=450$ nm.




Figure 8. The AAOD difference between fractal aggregate models and the spherical model, $\lambda = 450$ nm.




Figure 9. BC AOD (550 nm) and AAOD(450 nm) as a percentage of total AOD and AAOD.




Figure 10. The comparison of SSA using different BC models, $\lambda=450$ nm.




Figure 11. The SSA differences between fractal aggregate models and the spherical model, $\lambda=450$ nm.




Figure 12. The clear-sky BC ARI at the typical urban polluted cities in eastern China and fire sites in the northwest US calculated using different BC models.




Figure 13. The all-sky BC ARI at the typical urban polluted cities in eastern China and fire sites in the northwest US calculated using different BC models..




Figure 14. The regional-mean BC ARI at the TOA in eastern China for different particle models.




Figure 15. Similar to Figure 14, but for northwest US.

Code availability. The WRF-Chem is publicly available from <https://ruc.noaa.gov/wrf/wrf-chem/>; MSTM can be download from <https://www.eng.auburn.edu/~ckwski/scatcodes/>; The libRadtran is available from <http://www.libradtran.org/doku.php>; The FlexAOD can be requested from Prof. Curci (<http://pumpkin.aquila.infn.it/flexaod/>).

Data availability. The look-up tables calculated in this work can be obtained from https://figshare.com/articles/dataset/Look_up_tables_zip/13096241.
680 The PM_{2.5} data in China was obtained from <https://www.aqistudy.cn/historydata/>, and PM_{2.5} data in the northwest US can be found from <https://www.epa.gov/outdoor-air-quality-data/download-daily-data>.

Author contributions. JL and QXZ conceived the presented idea. JL developed the models, performed the computations, and wrote the paper. ZQL, CZ, YMZ, RKC, YZ verified the simulation methods and results. QXZ revised the paper and supervised the findings of this work. GC developed the FlexAOD model. All authors discussed the results and contributed to the final paper.

685 *Competing interests.* The authors declare that they have no conflict of interest.