

Effects of ozone-vegetation interactions on meteorology and air quality in China using a two-way coupled land-atmosphere model

5 Jiachen Zhu¹, Amos P.K. Tai², Steve Hung Lam Yim^{3,4}

¹Department of Geography and Resource Management, The Chinese University of Hong Kong, Sha Tin, N.T., Hong Kong, China

10 ²Earth System Science Programme, The Chinese University of Hong Kong, Sha Tin, N.T., Hong Kong, China

³Asian School of the Environment, Nanyang Technological University, 50 Nanyang Avenue, 639798 Singapore

⁴Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore

15 *Correspondence to:* Steve H.L. Yim (yimsteve@gmail.com) & Amos P.K. Tai (amostai@cuhk.edu.hk)

Abstract. Tropospheric ozone (O₃) is one of the most important air pollutants in China and is projected to continue to increase in the near future. O₃ and vegetation closely interact with each other and such interactions may not only affect plant physiology (e.g., stomatal conductance and photosynthesis) but also influence the overlying meteorology and air quality through modifying leaf stomatal behaviors. Previous studies have highlighted China as a hotspot in terms of O₃ pollution and O₃ damage to vegetation. Yet, few studies have investigated the effects of O₃-vegetation interactions on meteorology and air quality in China, especially in the light of recent severe O₃ pollution. In this study, a two-way coupled land-atmosphere model was applied to simulate O₃ damage to vegetation and the subsequent effects on meteorology and air quality in China. Our results reveal that O₃ causes up to 16% enhancement in stomatal resistance, whereby large increases are found in Henan, Hebei and Shandong provinces. O₃ damage causes more than 0.6 μmol CO₂ m⁻² s⁻¹ reductions in photosynthesis rate, and at least 0.4 and 0.8 g C m⁻² day⁻¹ decrease in leaf area index (LAI) and gross primary production (GPP), respectively, and hotspot areas appear in the northeastern and southern China. The associated reduction in transpiration causes a 5–30 W m⁻² decrease (increase) in latent heat (sensible heat) flux, which induces a 3% reduction in surface relative humidity, 0.2–0.8 K increase in surface air temperature, and 40–120 m increase in boundary layer height in China. We also found that the meteorological changes further induce a 2–6 ppb increase in O₃ concentration in northern and south-central China mainly due to enhanced isoprene emission following increased air temperature, demonstrating that O₃-vegetation interactions can lead to strong positive feedback that can amplify O₃ pollution in China. Our findings emphasize the importance of considering the effects of O₃ damage and O₃-vegetation interactions in air quality simulations, with ramifications for both air quality and forest management.

40

1. Introduction

Tropospheric ozone (O₃) is a secondary air pollutant, which is mainly formed from the photochemical oxidation of carbon monoxide (CO), methane (CH₄) and non-methane volatile organic compounds (VOCs) by hydroxyl radicals (OH) in the presence of nitrogen oxides (NO_x = NO + NO₂). O₃ is known as the third most important greenhouse gas with an estimated radiative forcing of 0.41 W m⁻² for the period of 1750–2010 (IPCC, 2013; Stevenson et al., 2013). As an air pollutant, O₃ is also shown to be harmful to not only human health but also vegetation and crop health (Anenberg et al., 2010; Cohen et al., 2017). Various field experiments and numerical modeling studies have already demonstrated that O₃ can not only reduce gross primary production (GPP) of natural vegetation as well as crop yields (Ainsworth et al., 2012; Lombardozzi et al., 2012; Tai et al., 2014; Feng et al., 2015; Yue et al., 2017; Li et al., 2018), but also decrease transpiration (Arnold et al., 2018), decrease runoff (Li et al., 2016) on larger scales and therefore affect the global carbon and water cycle (Lombardozzi et al., 2015).

Vegetation can in turn modulate O₃ concentration through influencing the sources and sinks of O₃. Dry deposition of O₃ onto vegetation is a major sink for O₃, mainly via stomatal uptake. Stomata are the pores on plant leaves; they control water exiting and carbon entering the leaf interior and hence influence the water and carbon exchange between the land and atmosphere. When vegetation is exposed to enhanced O₃ levels, cellular and tissue damage can result in a decrease in photosynthesis rate, thus altering CO₂ assimilation. Stomata conductance may decrease subsequently in response to O₃ exposure, thus reducing the dry-depositional sink of O₃ (Sadiq et al., 2017; Zhou et al., 2018), but some studies also suggest that O₃ exposure can cause stomata to respond more sluggishly to changing environmental conditions, such as drought, with complex overall effects on stomatal behaviors and dry deposition (e.g., Huntingford et al., 2018). Moreover, recent studies showed reduced dry deposition velocities of O₃ by drought-stressed vegetation, which affects surface O₃ trends and extremes (Huang et al., 2016; Lin et al., 2019; Lin et al., 2020). Vegetation also affects the sources of O₃; the most abundant biogenic VOC (BVOC) species emitted by vegetation is isoprene (C₅H₈), which is a major precursor for O₃ formation in polluted, high-NO_x environments, but removes O₃ by ozonolysis or by sequestering NO_x in more pristine, low-NO_x regions (Hollaway et al., 2017). Isoprene production is known to be highly coupled with photosynthesis and by extension to stomatal conductance (Arneeth et al., 2007). Moreover, transpiration, which is modulated by stomatal behaviors, significantly regulates surface meteorology including water vapor content and air temperature, which further influence the production and loss of O₃. Therefore, through influencing plant ecophysiology (e.g., photosynthesis and stomata behaviors), O₃-vegetation interactions can modulate boundary-layer meteorology, climate, and may further affect O₃ air quality via a series of feedback mechanisms. It is therefore essential to fully understand the O₃-vegetation interactions and the following climatic and biospheric impacts especially in areas with high O₃ concentrations and vegetation density.

In many land surface and biospheric models, such as Noah-Multi Parameterization (Noah-MP) or Community Land Model (CLM), the Farquhar-Ball-Berry model (FBB, Farquhar et al., 1980; Ball et al., 1987) is commonly used to simulate stomatal conductance and photosynthetic rate. In the FBB model, the calculation of stomata conductance is based on the calculation of photosynthesis, which makes them tightly coupled with each other. Therefore, in several land surface models that consider O₃ damage effect on vegetation, the photosynthetic rate is modified first and the stomatal conductance is modified subsequently, which means stomata conductance and photosynthesis will change collinearly under

chronic O₃ exposure (Sitch et al., 2007; Yue and Unger, 2014). However, field experiments have shown that, under chronic O₃ exposure, stomata conductance decreases with a smaller magnitude than photosynthetic rate does, which makes the simulations of stomata conductance and photosynthetic rate as well as the following water and carbon cycles in the above models less accurate (Lombardozi et al., 90 2012). Modifying stomata conductance and photosynthesis separately in land surface models is therefore more reasonable. Lombardozi et al (2012) modified the stomata conductance and photosynthetic rate separately based on the cumulative uptake of O₃ into leaves and has shown a better representation of plant responses to O₃ exposure. Efforts have been made to investigate the effects of O₃ exposure on land biosphere based on the above O₃ damage schemes. For example, based on an off-line process-based 95 vegetation model, Yue and Unger (2014) found that O₃ damage decrease GPP by 4–8% on average in the eastern US and leads to significant decreases of 11–17% in east coast hot spots. Using the offline CLM model, Lombardozi et al. (2015) estimated that the present O₃ exposure reduces GPP and transpiration globally by 8–12% and 2.0–2.4%, respectively.

100 Several modeling studies conducted so far have demonstrated the importance of considering the interactions and feedbacks between atmosphere and biosphere. By dynamically coupling O₃ and LAI but without considering the meteorological feedbacks of O₃-vegetation interactions to O₃, Zhou et al. (2018) found that O₃-induced damage on LAI can lead to changes in O₃ concentrations by –1.8 to +3 ppb in boreal summer. By considering the interactions between atmospheric chemistry with biosphere in a two- 105 way coupling model, Lei et al. (2020) quantified the damaging effects of O₃ on vegetation and found a global reduction of annual GPP by 1.5–3.6 %, with regional extremes of 10.9–14.1 % in the eastern US and eastern China. Based on the CESM model with fully interactive atmospheric chemistry, biogeochemical and biogeophysical cycles, Sadiq et al. (2017) estimated that surface O₃ is 4–6 ppb higher in Europe, North America and China in simulations with O₃-vegetation coupling comparing the surface 110 O₃ concentrations without O₃-vegetation coupling. Based on modified WRF-Chem model, Li et al (2016, 2018) investigated the effect of O₃ exposure on hydroclimate and crop productivity in the US, and highlighted O₃ damage effects on meteorological fields and surface energy balance as well as the crop yields, but the feedbacks of changing meteorology onto surface O₃ were not investigated. Arnold et al (2018) examined the global climate response to O₃ exposure and found O₃ damage on vegetation can 115 induce widespread surface warming and changes in clouds, which could be critical on regional scales. Although the interactions between O₃ and vegetation are critical to our environment, adequate representation of O₃-vegetation interactions is still missing in most atmospheric models used for climate and atmospheric chemistry simulations, at least in part due to incomplete coupling capacities with land surface or biospheric model components at high resolutions, and in part due to limited observations to 120 optimize O₃ damage schemes for wider regional applicability.

With the rapid urbanization and industrialization in the recent decades, China has experienced increasingly severe O₃ pollution, which is expected to continue to worsen in the near future. O₃ concentration in China has been observed to exceed ambient air quality standard by 100–200% (Wang 125 et al., 2017) with the maximum 8-hour mean concentration of O₃ (MDA8 O₃) increasing by 4.6% per year from 2015 to 2017 (Silver et al., 2018). Lu et al. (2018) showed that urban surface O₃ in China during 2013–2017 was significantly higher than that in other regions around the world, and thus vegetation exposure to O₃ is also higher in China. Li et al. (2018) also revealed the increasing trend of O₃ in megacity clusters of China during 2013–2017, which is closely related with meteorology,

130 anthropogenic emissions and PM_{2.5} concentrations. Global-scale studies have highlighted China as a
hotspot of O₃ pollution and damage to vegetation compared with other regions (Sadiq et al., 2017; Arnold
et al., 2018; Lei et al., 2020). However, a comprehensive study of how O₃ affects meteorology and air
quality through O₃-vegetation interactions in China at high spatial resolutions, especially under severe
O₃ pollution, is still limited but highly needed. Moreover, there have been limited studies focusing on the
135 feedbacks of O₃-vegetation coupling on O₃ concentration itself, especially in China, which is one of the
main scopes of our study.

This study, therefore, first adopted and implemented a semi-mechanistic O₃ damage scheme in a widely
used regional atmosphere-land modeling framework and hence used it to simulate and assess the impacts
140 of O₃-vegetation interactions on boundary-layer meteorology and air quality in China at a high spatial
resolution. Specifically, O₃-induced damage to vegetation, changes in meteorology in China due to O₃-
vegetation coupling, and the subsequent feedback effects onto O₃ concentration itself are examined,
which is crucial to fully understand the O₃-vegetation interactions and the following impacts on climate,
biosphere, and air quality in areas with both high O₃ concentrations and high vegetation coverage.

145

2. Methods

2.1 WRF-Chem Model Setup

The Weather Research and Forecasting (WRF) model is a state-of-the-art mesoscale nonhydrostatic
150 meteorological model. An atmospheric chemistry module that includes various gas-phase chemistry and
aerosol mechanisms has been implemented into and fully coupled with WRF to create the WRF-Chem
model (Grell et al., 2005; Fast et al., 2006). In WRF-Chem, both the air quality and meteorological
components use the same transport scheme, model grid, subgrid-scale transport physics and time step.
WRF-Chem has been widely used in previous air quality studies (e.g., Li et al., 2016; Li et al., 2018; Liu
155 et al., 2018; Liu et al., 2020). In this study, we applied our revised WRF-Chem model based on version
3.8.1 to simulate meteorological fields and O₃ concentration over China. Simulations are conducted from
24 May to 1 September every year from 2014 to 2017 and the days in May were discarded as spin-up.
For the land surface component within WRF, we used Noah-MP, which will be described in the next
subsection.

160

The model domain was configured at a horizontal resolution of 27 km on the Lambert Conformal
projection, centered at 37°N, 108.1°E and covering the whole China. The model has 26 vertical layers,
with the lowest layer at 0.17 km and the highest layer at 17.67 km. The meteorological initial and
boundary conditions are provided by the 6-hourly Final Operational Global Analysis (FNL) dataset at a
165 horizontal resolution of 1°×1°. The chemical initial and boundary conditions were generated from the
Model for Ozone and Related Chemical Tracer version 4 (MOZART-4), which is available at a horizontal
resolution of 1.9°×2.5° with 56 vertical layers (Emmons et al., 2010).

Anthropogenic emissions were from the Multi-resolution Emission Inventory for China (MEIC)
170 compiled at a spatial resolution of 27 km and a 1-hourly temporal resolution suitable for our research
domain. Biogenic emissions were calculated online by the Model of Emissions of Gases and Aerosol
from Nature (MEGAN) (Guenther et al., 2006). Biomass burning emissions were extracted from the Fire
Inventory from NCAR (FINN) version 1.5 datasets (Wiedinmyer et al., 2010). Dust emissions were

175 generated online by the Goddard Global Ozone Chemistry Aerosol Radiation and Transport model
(GOCART; Ginoux et al., 2001). Gas-phase chemistry was simulated with second generation Regional
Acid Deposition Model (RADM2; Stockwell et al., 1990) mechanism, and the Modal Aerosol Dynamics
Model for Europe (MADE; Ackermann et al., 1998), which is coupled with the Secondary Organic
Aerosol Model (SORGAM; Schell et al., 2001) for aerosol treatment. Detailed physics schemes used in
the simulations are shown in Table S1.

180

2.2 Description of Noah-MP model

Noah-MP is a land surface model that uses multiple options for key land-atmosphere interaction
processes (Niu et al., 2011). Noah-MP contains a separate vegetation canopy defined by a canopy top
and bottom, crown radius, and leaves with prescribed dimensions, orientation, density, and radiometric
185 properties. The canopy employs a two-stream radiation transfer approach along with shading effects
necessary to achieve proper surface energy and water transfer processes (Dickinson, 1983). Noah-MP is
capable of distinguishing between C_3 and C_4 photosynthesis pathways and defines vegetation-specific
parameters for plant photosynthesis and respiration.

190

Noah-MP is available for prognostic vegetation growth that combines a Ball-Berry photosynthesis-based
stomatal resistance (Farquhar et al., 1980; Ball et al., 1987) that allocates carbon to various parts of
vegetation (leaf, stem, wood and root) and soil carbon pools (fast and slow). GPP, leaf area index (LAI)
and canopy height are then predicted downstream from photosynthesis. Noah-MP also considers the
195 photosynthesis of sunlit and shaded leaves separately, whereby sunlit leaves are more limited by CO_2
concentration while shaded leaves are more constrained by insolation, which may thus have different
responses to O_3 damage. The dynamic LAI and canopy height calculation will further affect surface
energy fluxes, which will then affect the boundary-layer meteorology when coupling with the atmosphere
model in WRF-Chem. The land use types and the vegetation parameters are based on the U.S. Geological
200 Survey (USGS) embedded in Noah-MP. Fig. 1 shows the spatial distribution of vegetation fraction of
dominant vegetation types in China. The distribution of main vegetation groups (broadleaf, needleleaf,
crop and grass) that have different sensitivities to O_3 damage following Lombardozzi et al. (2015) are
shown in Fig. 1.

205 In this study, the O_3 concentration simulated by the chemical module of the WRF-Chem model was also
dynamically passed onto the Noah-MP land surface model at every time step to modify the
photosynthesis and stomatal conductance due to O_3 damage. The land surface variables simulated by
Noah-MP were also dynamically passed back onto the atmospheric components, thus allowing
immediate, two-way feedback effects onto meteorological fields, O_3 and other atmospheric chemical
210 constituents. In this way, land surface processes, atmospheric dynamics, and atmospheric chemistry in
the WRF-Chem model were fully coupled.

2.3 O_3 damage parameterization

215 In Noah-MP, the Farquhar model (Farquhar et al., 1980) was used to calculate photosynthetic rate,
whereas Ball-Berry model was used to calculate stomatal conductance (Ball et al., 1987). The

photosynthesis rate, A ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), is calculated separately for sunlit and shaded leaves and is limited by either one of three limiting factors and can be calculated as

$$A = \min(W_c, W_j, W_e) I_{gs} \quad (1)$$

where W_c is the Rubisco-limited photosynthesis rate, W_j is the light-limited photosynthesis rate, and W_e is the export-limited photosynthesis rate. I_{gs} is the growing season index with values ranging from 0 to 1. Stomatal conductance (g_s) is computed based on the photosynthesis rate from the Farquhar model as

$$g_s = \frac{1}{r_s} = m \frac{A e_s}{c_s e_i} P_{atm} + b \quad (2)$$

where g_s is the leaf stomatal conductance ($\mu\text{mol m}^{-2} \text{ s}^{-1}$); r_s is the leaf stomatal resistance ($\text{s m}^2 \mu\text{mol}^{-1}$); m is an empirical parameter that relates stomatal conductance and photosynthesis with values ranging from 5 to 9; A is the photosynthesis rate as described above; c_s is the CO_2 partial pressure at the leaf surface (Pa); e_s is the vapor pressure at the leaf surface (Pa); e_i is the saturation vapor pressure inside the leaf (Pa); P_{atm} is the atmospheric pressure (Pa); and b is the minimum stomatal conductance.

As mentioned above, following Lombardozzi et al. (2015), an O_3 damage scheme was implemented in Noah-MP embedded in WRF-Chem model version 3.8.1. The photosynthesis rate and stomatal conductance are modified independently using two sets of O_3 impact factors, $F_{p\text{O}_3}$ and $F_{c\text{O}_3}$, respectively, which are then multiplied to the initial A and g_s calculated by the Farquhar-Ball-Berry model, respectively. Lombardozzi et al. (2012) found that independently modifying stomatal conductance and photosynthesis can improve the model prediction of plant response to O_3 damage. The two damage factors are calculated based on the cumulative uptake of O_3 (CUO), which integrates the O_3 flux inside leaves through the stomata throughout the growing season. The CUO (mmol m^{-2}) is calculated as

$$\text{CUO} = 10^{-6} \sum \frac{[\text{O}_3]}{k_{\text{O}_3} r_s + r_a + r_b} \Delta t \quad (3)$$

Where $[\text{O}_3]$ is the surface O_3 concentration (nmol m^{-3}); $k_{\text{O}_3} = 1.61$ is the ratio of leaf resistance to O_3 to leaf resistance to water (Uddling et al., 2012); r_s is the stomatal resistance, r_a is the aerodynamic resistance and r_b is the boundary-layer resistance (s m^{-1}); Δt is the model time step (s). CUO is only accumulated when LAI is larger than 0.4 and O_3 flux is larger than a threshold value of $0.8 \text{ nmol O}_3 \text{ m}^{-2} \text{ s}^{-1}$ to consider the detoxification effect of plants to O_3 damage.

The two damage factors have linear relationships with CUO and can be calculated as follows:

$$F_{p\text{O}_3} = a_p \times \text{CUO} + b_p \quad (4)$$

$$F_{c\text{O}_3} = a_c \times \text{CUO} + b_c \quad (5)$$

where $F_{p\text{O}_3}$ is the O_3 damage factor for photosynthesis and $F_{c\text{O}_3}$ is the O_3 damage factor for stomatal conductance; a_p , b_p , a_c , and b_c are empirical slopes and intercepts of three different plant groups (broadleaf trees, needleleaf trees, and grasses or crops) from Lombardozzi et al. (2015). The values of

260 these slopes and intercepts are shown in Table 1. The original photosynthesis and stomatal conductance are then multiplied with the two damage factors, respectively to get the modified photosynthesis and stomatal conductance under O₃ exposure.

2.3 Model Experiments and Evaluation

265 Two sets of experiments were conducted in this study. We performed a control simulation (simu_withoutO₃) without O₃ damage on vegetation and a production simulation (simu_withO₃) with O₃ damage on vegetation. Detailed information of the experiments is shown in Table 2. In the simu_withO₃ experiment, the O₃ concentration simulated by the chemical module of the model is dynamically passed onto the land surface model at every time step to modify the photosynthesis and stomatal conductance.
270 The differences between the two sets of experiments including vegetation physiology, meteorological fields and O₃ concentration can thus be attributed to O₃-vegetation interactions. In this work, each simulation was conducted from 24 May to 1 September every year from 2014 to 2017 and the days in May were discarded as spin-up. For each simulation in the four years, anthropogenic emissions were kept at 2014 levels, while meteorological fields were changing every year. The 4-year June-July-August (JJA) averaged results were analyzed and compared. JJA was selected because of the most severe O₃ pollution in this season and because it is within the active growing season of the plants.
275

The simulated meteorological variables and air pollutant concentrations were evaluated using available in-situ observations in China. The daily meteorological observations including temperature at 2 meter (T_{2m}), relative humidity at 2 meter (RH_{2m}), and wind speed at 10 meter (WS_{10m}) above displacement height were from the National Meteorological Information Center. There are 698 stations in the study domain. The air pollutant observations were provided by the China National Environmental Monitoring Center (CNEMC) network, which offers hourly concentrations of particulate matter with an aerodynamic diameter of less than 2.5 μm (PM_{2.5}) and 10 μm (PM₁₀), carbon monoxide (CO), O₃, sulfur dioxide (SO₂) and nitrogen dioxide (NO₂). The locations of meteorological stations and the sites of CNEMC network are shown in Figure 2. The statistical parameters including mean values (Mean) of observations and simulated variables, their standard deviations (SD), indices of agreement (IOA), mean biases (MB), and correlation coefficients (CORR) were computed to evaluate the model performance in this study.
285

290 3. Results

3.1 Model evaluation

Table 3 shows the city-averaged evaluation results of meteorological variables from the modified model.
295 The information of the major cities used for evaluation is shown in Table S4. From Table 3, we can find that T_{2m} is underestimated with MB values ranging from -1.00 °C in 2017 to -0.70 °C in 2014. The IOA and CORR are generally higher than 0.8, indicating that the model could reasonably simulate the variations of T_{2m}. Unlike temperature, relative humidity is overestimated by the model simulations with MB values ranging from 4.38 in year 2014 to 7.33 in year 2016, but the CORR values with observations are still high (CORR > 0.7). Wind speed is also overestimated by more than 0.38 m s⁻¹, which might be caused by the underestimation of terrain height as reported in other WRF modeling studies (Brunner et al., 2015; Liu et al., 2020). The detailed evaluation results for each city and for seven major geographic
300

regions of China are shown in Table S5-S10. The classification of the geographic regions is shown in Fig. S2. As shown in these tables, the model can reasonably capture the spatial distribution of these meteorological variables. For example, the larger values of T_{2m} and RH_{2m} in cities from southern China compared with the cities in northern China (Table 4) can be reasonably simulated. We also found that the model simulations have better performance in northeastern China, central China and southern China in terms of IOA and CORR as shown in these tables (Table 4).

Table 5 shows the city-averaged evaluation results of six air pollutants simulated from the modified model. The information of the major cities used for air pollutant evaluation is shown in Table S11. From Table 5, positive MB values for O_3 , $PM_{2.5}$, SO_2 , NO_2 , and negative MB values for CO are found. The overestimation of O_3 by WRF-Chem was also reported by Hu et al. (2016) and Gao et al (2020). For PM_{10} , both positive and negative MB values are found for different years. The results indicate general overestimation by the model of most air pollutants except for CO. The underestimation of CO can be explained by either O_3 chemistry, which points to the problem related to low titration, or in the underestimation of dry deposition by the model, which is also affected by the modification of the model. The IOA of air pollutant concentration ranges from 0.36 (SO_2) to 0.63 (O_3). The correlation coefficient of air pollutants ranges from 0.14 (PM_{10}) to 0.66 (O_3). Detailed evaluation results for each city and major geographic regions of China are shown in Tables S9–S14 and Table 6. In terms of the evaluation for O_3 , the model has better performance in northeastern China, eastern and southern China, which may suffer the most severe O_3 damage. Our results are generally consistent with the evaluation results of CMAQ simulation over China by Liu et al. (2020). MBs of SO_2 , NO_2 and CO are consistent in both magnitude and sign with Liu et al. (2020), while the MBs of PM and O_3 are larger than Liu et al. (2020). Correlation coefficients of air pollutants are also of similar magnitude with Liu et al. (2020), showing that our model results can well capture the temporal variations of air pollutants. We also compared the evaluation results between the original model and the modified model, as shown in Table S2 and Table S3 in the supplement and Table 3 and Table 5 here. We found no obvious differences in the evaluation results between the original model results and the revised model results. It should be noted that this study might not be able to and was not meant to improve model accuracy, but our modified model is able to capture O_3 -vegetation interactions without worsening model performance. Overall, there are systematic biases in simulated variables especially the air pollutant concentrations, but the spatial distribution of both meteorological variables and air pollutant concentrations are reasonably simulated by the model, lending trust to the use of the model for sensitivity studies to examine the effects of O_3 -vegetation interactions on the atmospheric environment.

3.2 Responses of vegetation to O_3 damage

O_3 can adversely affect photosynthesis rate and stomatal conductance and therefore interfere with vegetation growth, productivity and transpiration. To understand the O_3 -induced damage on vegetation physiology, the spatial distribution and changes in stomatal resistance (RS), photosynthesis rate (PSN), LAI, GPP, and transpiration rate (TR) during 2014–2017 summer (June-July-August) were analyzed.

Figure 3a and 3d display the spatial distribution of sunlit stomatal resistance (RSSUN) and shaded stomatal resistance (RSSHA) from the `simu_withoutO3` experiment, respectively. The absolute and relative changes in RSSUN (RSSHA) between `simu_withO3` and `simu_withoutO3` experiments are shown

in the middle and the right panel of Fig. 3, separately. In general, simulated stomatal resistance in eastern China is larger than that in western China. Both RSSUN and the RSSHA are enhanced in response to O₃ damage to vegetation. The maximum increases in RSSUN and RSSHA can be up to $1.0 \times 10^3 \text{ s m}^{-1}$, which is equivalent to a ~16% increase compared to the `simu_withoutO3` simulation. Comparing the changes in RSSUN vs. RSSHA, the changes in RSSHA are larger than that in RSSUN, reflecting the larger sensitivity of shaded leaves to O₃ damage (Kinose et al., 2017). Northern China experiences larger changes in stomatal resistance generally, especially in Henan, Hebei, and Shandong provinces, where the changes in stomatal resistance are twice as much as the changes in stomatal resistance over other regions.

The spatial distribution of 2014–2017 JJA mean PSN, LAI and GPP from the `simu_withoutO3` simulations and their changes induced by O₃ damage are presented in Fig. 4. From Fig. 4a, we find that the PSN values are generally higher in eastern China compared with western China with the largest values of up to $\sim 7 \mu\text{mol CO}_2^{-1} \text{ m}^{-2} \text{ s}^{-1}$. Similar spatial distribution and hotspot areas can also be observed for LAI (Fig. 4d) and GPP (Fig. 4g), with LAI and GPP values in hotspot areas up to 3.6 and 10 $\text{g C m}^{-2} \text{ day}^{-1}$, respectively. We also find that Henan, Hebei, Shanxi and Shandong provinces have smaller values in PSN, LAI and GPP when compared with other provinces in eastern China.

With O₃ damage, PSN decreases in general, with absolute changes in PSN ranging from 0.6 to 3.6 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Fig. 4b), representing 20–40% reductions in PSN. For northeastern and southern China, where the original PSN values are large, ~20% reductions in PSN are found (Fig. 4c). In western China where the dominant vegetation type is grassland and the original PSN values are small, more than 40% of PSN is reduced due to O₃ damage (Fig. 4c). In response to the PSN reductions, LAI and GPP also decrease. More than 0.4 reductions in LAI are found in central and northern China (Fig. 4e), corresponding to more than 20% reductions in LAI; in other regions, 5–15% reductions in LAI are observed. More than 0.8 $\text{g C m}^{-2} \text{ day}^{-1}$ reductions in GPP are found generally in China. Similar to Fig. 3c, we find that GPP decreases by ~20% in northeastern and southern China and decreases by more than 40% in other regions (Fig. 4i). Based on offline models without considering atmosphere-biosphere coupling, O₃ damage was found to decrease GPP at most by 11–17% in the East Coast hotspots of the US (Yue and Unger, 2014). Using the offline CLM model, Lombardozzi et al. (2015) estimated that the present O₃ exposure reduces GPP globally by 8–12%. Based on RegCM-CHEM4 regional climate model coupled with YIBs terrestrial biosphere model, Xie et al. (2019) revealed that O₃ damage induces a significant reduction ($12.1 \pm 4.4\%$) in the GPP, up to 35% in summer over China (Table S15). Comparing our results with previous studies, our results are broadly consistent with Xie et al. (2019) but the magnitude is larger than the studies conducted by Yue and Unger (2014) and Lombardozzi et al. (2015). Differences or uncertainties may arise from the different model settings. It appears that offline models as used by Yue and Unger (2014) and Lombardozzi et al. (2015) generally found smaller damage than studies with two-way coupling between the atmosphere and biosphere as used by Xie et al. (2019) and our work; this could be due to the existence of positive biosphere-atmosphere feedbacks that potentially worsen O₃ damage, as will be discussed in subsequent sections. Different O₃ damage schemes employed in the models may also be a source of differences, although we note that both this work and Lombardozzi et al. (2015) used the same scheme, so the differences appear to arise more likely from the effect of coupling and other model settings than from the schemes alone.

390 The spatial distribution of dominant vegetation types in China are shown in Fig. 1, where we can see that
the croplands dominant in eastern China and especially in southern China suffer the greatest GPP
reductions, indicating that crop yields in China would also be heavily affected by O₃ damage.

Figure 5 depicts the spatial distribution of transpiration rate (TR) of vegetation and the changes in
395 transpiration rate induced by O₃ damage. TR values are higher in eastern China where there is larger
vegetation coverage (Fig. 5a). As shown in Fig. 5b, TR decreases by 0.2–1.0 mm day⁻¹ generally in
eastern China with large reductions in northern China, especially in Henan, Shandong, Anhui and Jiangsu
provinces. In terms of relative changes, TR decreases by ~12% in northeastern and southern China, while
400 more than 24% reductions are found in other regions. Transpiration is affected by the changes in both
RS and LAI. With O₃ damage, both the increases in RS (Fig. 3c and Fig. 3f) and decreases in LAI (Fig.
4f) cause TR to decrease, as shown in Fig. 5b and 5c. Comparing the changes in RS (Fig. 3c and Fig. 3f),
LAI (Fig. 4f) and TR (Fig. 5c), we can find that the distribution of changes in TR is more consistent with
that of RS, reflecting the dominance of RS in controlling TR.

405 3.3 Changes in meteorology due to O₃-vegetation coupling

Through interacting with vegetation, O₃ has the potential to further affect the meteorological environment
in China via modifying, e.g., surface heat fluxes, temperature, humidity, and boundary layer height. The
distribution of meteorological variables from simulations with and without O₃ damage is thus compared
410 and analyzed in this section.

Figure 6 shows the spatial distribution of latent heat (LH) flux and sensible heat (SH) flux, and the
changes in LH and SH due to O₃-vegetation coupling. With O₃ included in the model simulations, the
LH flux decreases by more than 4 W m⁻² (Fig. 6b) on average following the decreases in transpiration
415 rate. Hotspot areas are found in Henan, Shandong, Anhui and Jiangsu provinces, where reductions in LH
can be up to 30 W m⁻². Meanwhile, 5–30 W m⁻² increases in SH flux are observed in central and northern
China (Fig. 6d). With O₃-vegetation coupling, more than 20% reductions in LH flux are found in central
and northern China (Fig. 6c), 20% increment in SH flux are found in similar regions (Fig. 6f), indicating
that O₃ damage shifts the energy balance toward more net radiation being dissipated by SH flux than LH
420 flux, with ramifications for surface temperature.

Figure 7 shows the distribution and the changes in surface relative humidity, temperature and planetary
boundary layer height (PBLH) in response to O₃ damage. Reductions in transpiration rate can directly
cause reductions in relative humidity. As shown in Fig. 7b, relative humidity has at least 3% absolute
425 reductions. Values of relative humidity decrease more in northern China than in southern China. Similar
to the changes in TR (Fig. 5b), larger reductions in relative humidity (3–9%) are found over Henan,
Hebei, Shandong, Anhui provinces. The decreases in LH flux and increases in SH flux following the
changes in transpiration rate drive the increases in temperature and contribute to PBLH growth. As
presented in Fig. 7e and Fig. 7h, the distribution and hotspot areas of the changes in temperature and
430 PBLH are similar to those in relative humidity. Generally, northern China has larger increases of
temperature and PBLH compared with other regions. Generally, temperature increases by 0.2–0.8 K and
PBLH increases by 40–120 m for northern China. The hotspot areas experience at least 0.6 K increases
in temperature, and 80 m increases in PBLH.

435 As shown in Table S15, our results are comparable with results from a regional simulation conducted by
Li et al. (2016), which showed that O₃ damage decreases LH flux by 10–27 W m⁻² and O₃ damage
increases temperature by 0.6 °C–2.0 °C in the US. However, in their study, Li et al. (2016) assumed that
O₃ damage to plants happens when O₃ concentration is over a threshold of 20 ppb to imitate a weaker
440 detoxifying effect of plants, instead of the 40 ppb threshold that was commonly used in previous studies.
Considering the severe O₃ air pollution in China, we resorted to use the more universal O₃ threshold used
by previous studies (Lombardozi et al., 2015; Sadiq et al., 2017; Zhou et al., 2018) to represent a more
conventional detoxifying effect, instead of lowering the threshold value that would cause much larger
changes in the surface fluxes and meteorological fields. Using a two-way coupling model and the same
O₃ damage scheme, Arnold et al. (2018) revealed that O₃ causes less than 8 W m⁻² changes in surface
445 heat fluxes regionally, which is smaller than the changes of surface heat fluxes in our study. One possible
reason is that the simulated changes in O₃ and aerosol in Arnold et al. (2018) did not feedback onto
radiation and climate simulation or affect LAI.

3.4 O₃-vegetation feedbacks on O₃ concentrations

450 O₃-induced changes in vegetation, surface fluxes and the overlying meteorology can also constitute
important feedback effects onto O₃ concentration itself. Figure 8 shows the spatial distribution of surface
O₃ concentration. The change in surface O₃ concentration during daytime is also shown in Fig. S2. As
shown in Fig. 8a (Fig. S2), surface O₃ concentration is higher in central and northern China during
455 summer. In terms of the feedbacks on O₃ concentration, we found generally enhancements in O₃
concentration when O₃-vegetation interactions are accounted for, thus representing a positive feedback
that worsens O₃ air quality (Fig. 8b). O₃ concentration increases the most (by up to 6 %) in Hebei, Shanxi
and Henan provinces, with the maximum increment of 6 ppb. The enhancement in surface O₃
concentration from our study is at the similar magnitude with that from the study conducted by Sadiq et
460 al. (2017), in which both biogeochemical and meteorological feedbacks from O₃-vegetation interactions
to O₃ are considered. Without considering the meteorological feedbacks following the changes in
transpiration to O₃ concentrations, smaller feedbacks on surface O₃ concentrations are found by the
following studies. For instance, by incorporating O₃-LAI coupling in chemical transport model, Zhou et
al. (2018) found an O₃ feedback of -1.8 to +3 ppb globally. Another similar work conducted by Gong et
465 al. (2020) showed that O₃-induced inhibition in stomatal conductance increases surface O₃ by 2.1 ppb in
eastern China, while considering the addition effects of O₃ on isoprene emission slightly reduces surface
O₃ concentrations by influencing the precursors. Soil moisture deficit, which has been shown to reduce
stomatal uptake, if considered, will also contribute to the enhancement in O₃ concentration (Rydsgaard
et al., 2016). Together with previous findings, it is increasingly clear that meteorological feedback could
470 be an important pathway whereby O₃-vegetation interactions can further worsen O₃ air quality, almost
doubling the effect of biogeochemical feedback alone (i.e., via changes in O₃-relevant chemical fluxes
alone). It should be cautiously noted that in terms of magnitude alone the model biases in O₃ are
comparable and sometimes larger than the up to 6 ppb systematic enhancement caused by O₃ damage,
which represents be one major source of uncertainties in our study.

475 Reduced dry deposition due to stomatal closure and reduced LAI, as well as increased isoprene emission,
are all found to be the drivers for the overall positive O₃ feedback. Reductions in dry deposition velocity,

480 following closely the corresponding reductions in transpiration rate as both processes are modulated by stomatal regulation, contribute in part to the O₃ enhancement. Figure 9 shows the spatial distribution of isoprene emission and its changes due to O₃ damage. We observe general increases in isoprene emission in eastern China, mainly due to increased surface temperature (Figs. 7e and 7f) that is more than enough to offset reduced isoprene caused by reduced LAI (Figs 4e and 4f). All in all, O₃ damage on vegetation can further enhance O₃ levels via an overall positive effect, due to not only the associated reductions in dry deposition velocity, but also the reductions in transpiration, LH flux and the resulting rise in surface temperature.

4 Conclusions

490 Tropospheric O₃ is one of the most concerning air pollutants due to its global warming effects and its ability to affect human health, vegetation and crops. O₃ and vegetation closely interact with each other and such interactions may not only affect plant physiology (e.g., stomatal conductance and photosynthesis) but also influence the overlying meteorology and air quality through modifying leaf stomatal behavior, plant structure (e.g., LAI) and subsequently land-atmosphere fluxes. According to previous field experiments and modeling works, China has been recognized as one of the hotspot areas suffering from severe O₃ pollution and the resulting damage on vegetation and crops, but the feedback effects onto air quality and climate have not been fully characterized. Previous studies mainly focused on the global scale with coarse spatial resolutions, which did not fully capture the spatial distribution of O₃ damage on vegetation in China. Based on the results from global studies pointing out that China is a hotspot in terms of O₃ pollution and O₃ damage on vegetation, our model simulations performed at high spatial resolutions were capable of investigating O₃ damage effects on regional and provincial scales in China. In this study, we examined the effects of O₃-vegetation interactions on O₃ air quality and meteorology in China during 2014–2017 based on the two-way coupled WRF-Chem model simulations whereby O₃, meteorology and vegetation physiology and structure can co-evolve with each other in real time.

505 We found that in China stomatal resistance is enhanced by up to 16%, which is the direct response to O₃ damage. Northern China, especially Henan, Hebei, and Shandong provinces, is identified as a hotspot area. For photosynthesis, more than 20% reductions are observed in China. Large reductions (>2.4 μmol CO₂ m⁻² s⁻¹) are found in northeastern and southern China. Following reduced photosynthesis, LAI shows relatively small reductions (5–15%), while GPP shows more than 20% reductions (1.6 g C m⁻² day⁻¹). Changes in transpiration rate are due to both changes in stomatal resistance and changes in LAI. With the increases in stomatal resistance and decreases in LAI, transpiration decreases from 0.2 to 1.0 mm day⁻¹ in eastern China with the largest reductions occur in northern China. We also found that the distribution of changes in transpiration is consistent more with the distribution of stomatal resistance than with those of LAI, indicating the dominance of the former in contributing to the overall transpiration rate.

520 With O₃ damage, the LH fluxes decrease by more than 4 W m⁻² on average, with hotspot areas appearing in Shandong, Anhui and Jiangsu provinces, in which the decreases can be up to 30 W m⁻² following mostly the decreases in transpiration rate. SH fluxes increase in similar areas at comparable magnitudes (10–25 W m⁻²). The decreases in LH and the increases in SH cause the increases in temperature and PBLH. We found that northern China has larger decreases in relative humidity, temperature and PBLH

compared with other regions. Generally, relative humidity shows at least 4% relative reductions, temperature increases by 0.2–0.8 K, and PBLH increases by 40–120 m for northern China. This indicates that O₃-vegetation interactions will cause a shift in the energy balance toward a state where available net radiation is dissipated more by SH flux than LH flux, with ramifications for surface temperature. This represents an additional pathway whereby anthropogenic O₃ pollution can worsen warming, in addition to O₃ being a greenhouse gas itself and O₃-induced plant damage diminishing the global net carbon sink (e.g., Sitch et al., 2007; Lombardozzi et al., 2015).

O₃ induces changes in vegetation, surface fluxes and meteorology, and in turn affects its own concentration. In this study, we found that reduced dry deposition in China is mainly due to enhanced stomatal conductance, while enhanced isoprene emission is mainly due to enhanced surface temperature and the corresponding increase in O₃ concentration. O₃ concentration increases the most (up to 6%) in Hebei, Shanxi and Henan provinces, with the maximum value of 6 ppb. Our results demonstrate that O₃-vegetation interactions can lead to strong positive feedback that can amplify O₃ pollution in China, in agreement with the suggestions by previous studies focusing on a global scale (Sadiq et al., 2017; Zhou et al., 2018; Gong et al., 2020). We also found that fully considering the positive O₃-vegetation feedbacks, especially when meteorological changes are also accounted for, generates greater damage on vegetation productivity than found by studies that only considered “offline” O₃ damage on plants without feedbacks (Yue and Unger, 2014; Lombardozzi et al., 2015).

In this study, the summertime simulation period of JJA was selected due to the high O₃ pollution in this season and the overlapping with vegetation growing season to capture the severe O₃ damage on vegetation. Nevertheless, uncertainty may still arise from that our simulation period may not cover the growing season of all vegetation types and may not cover all periods that O₃ damage happens, which may represent an underestimation of the full scale of O₃ damage. Future work should be conducted for longer time periods and for all seasons, which will help us better understand O₃-vegetation interactions in China. Uncertainty may also arise from the O₃ scheme employed in this study in terms of the CUO calculation and the consideration of O₃ detoxification mechanism of different vegetation types. The calculation of CUO heavily relies on the O₃ threshold. Considering the sensitivities of different vegetation types to O₃ damage, CUO threshold should be varied with different vegetation types. However, a constant O₃ threshold was employed in our study for the whole simulation domain and for all vegetation types, which may either underestimate or overestimate the actual O₃ damage. Moreover, following the work of Lombardozzi et al. (2015), we classified all the vegetation types into only three groups, which may be too coarse to investigate O₃ damage effects on regional or local scales. For example, Zhou et al. (2018) pointed out that Lombardozzi et al. (2015) treated tropical and temperate plants equivalently, which might lead to possible biases. More studies should be conducted to derive more appropriate O₃ thresholds for CUO calculation and make them available for regional scales or for different vegetation types. Another source of uncertainty may arise from the lack of representation of the direct effect of O₃ on isoprene emission. As pointed out by Gong et al. (2020), including the effect of O₃ damage on isoprene emission may reduce O₃ concentration by influencing precursors, but increase O₃ concentration at the same time through weakening the shortwave radiative forcing of secondary organic aerosols, which would help constitute a more complete feedback mechanism between O₃ and vegetation. Moreover, uncertainties may also come from that the effect of soil moisture deficit was not considered in this study, which may underestimate the reduction in dry deposition sink of O₃. It should also be noted that keeping

the anthropogenic emission inventory fixed in 2014 levels may be another limitation because of the nonlinear chemistry involving biogenic and anthropogenic precursors. Despite these uncertainties and limitations, our study provides detailed and comprehensive results whereby O₃-vegetation impacts will adversely affect plant growth and crop production, contribute to global warming, worsen the severe O₃ air pollution in China via feedbacks, and identifies the hotspot areas in the country. Our findings clearly pinpoint the need to consider the O₃ damage effects in both air quality studies and climate change studies.

Data availability. Model output data used for analysis and plotting, and the code used for simulations can be made available upon request.

Author contributions. APKT and SHLY conceived the study. JCZ carried out the model simulations and drafted the manuscript. SHLY and APKT supervised and funded the study.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments

This work is jointly funded by the Vice-Chancellor's Discretionary Fund of The Chinese University of Hong Kong (grant no. 4930744) given to both Steve H.L. Yim and Amos P. K. Tai, by Dr. Stanley Ho Medicine Development Foundation (grant no. 8305509) given to Steve H.L. Yim, and by a Research Grants Council (Hong Kong) General Research Fund grant (grant no. 14306220) given to Amos P. K. Tai. We would like to thank Dr. Jialun Li for the sharing and guidance of WRF-Chem code implementation in this work.

References

- Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S., & Shankar, U. M. A.: Modal aerosol dynamics model for Europe: Development and first applications, *Atmos. Environ.*, 32(17), 2981-2999, [https://doi.org/10.1016/S1352-2310\(98\)00006-5](https://doi.org/10.1016/S1352-2310(98)00006-5), 1998.
- Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J., and Emberson, L. D.: The effects of tropospheric ozone on net primary productivity and implications for climate change, *Annu. Rev. Plant Biol.*, 63, 637-661, <https://doi.org/10.1146/annurevplant-042110-103829>, 2012.
- Anenberg, S. C., Horowitz, L. W., Tong, D. Q., and West, J. J.: An estimate of the global burden of anthropogenic ozone and fine particulate matter on premature human mortality using atmospheric modeling. *Environ. Health Persp.*, 118(9), 1189-1195, <https://doi.org/10.1289/ehp.0901220>, 2010.
- Arneth, A., Niinemets, Ü., Pressley, S., Bäck, J., Hari, P., Karl, T., Noe, S., Prentice, I. C., Serça, D., Hickler, T., Wolf, A., and Smith, B.: Process-based estimates of terrestrial ecosystem isoprene emissions: incorporating the effects of a direct CO₂-isoprene interaction, *Atmos. Chem. Phys.*, 7, 31-53, <https://doi.org/10.5194/acp-7-31-2007>, 2007.
- Arnold, S. R., Lombardozzi, D., Lamarque, J. F., Richardson, T., Emmons, L. K., Tilmes, S., Sitch, S. A., Folberth, G., Holloway, M. J., and Val Martin, M.: Simulated Global Climate Response to Tropospheric Ozone-Induced Changes in Plant Transpiration, *Geophys. Res. Lett.*, 45(23), 13070-13079, <https://doi.org/10.1029/2018GL079938>, 2018.

- 610 Ball, J. T., Woodrow, I. E., and Berry, J. A.: A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, *Prog. Photosynthesis*, 4, 221-224, 1987.
- Brunner, D., Savage, N., Jorba, O., Eder, B., Giordano, L., Badia, A., and Galmarini, S.: Comparative analysis of meteorological performance of coupled chemistry-273 meteorology models in the context of AQMEII phase 2, *Atmos. Environ.*, 115, 470–498. <https://doi.org/10.1016/j.atmosenv.2014.12.032>, 2015.
- 615 Cohen, A. J., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., and Feigin, V.: Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015, *The Lancet*, 389(10082), 1907-1918, [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6), 2017.
- 620 Farquhar, G. D., Caemmerer, S. V., and Berry, J. A.: A Biochemical-Model of Photosynthetic CO₂ Assimilation in Leaves of C₃ Species, *Planta*, 149, 78–90, doi:10.1007/Bf00386231, 1980.
- Dickinson, R. E.: Land surface processes and climate—Surface albedos and energy balance, *Adv. Geophys.*, 25, 305–353, [https://doi.org/10.1016/S0065-2687\(08\)60176-4](https://doi.org/10.1016/S0065-2687(08)60176-4), 1983.
- 625 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- 630 Fast J. D., Gustafson Jr. W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct forcing in an urban area using a new fully-coupled meteorology, chemistry, and aerosol model. *J. Geophys. Res.*, 111, <https://doi.org/10.1029/2005JD006721>, 2006.
- Feng, Z., Hu, E., Wang, X., Jiang, L., and Liu, X.: Ground-level O₃ pollution and its impacts on food crops in China: A review, *Environ. Pollut.*, 199, 42-48, <https://doi.org/10.1016/j.envpol.2015.01.016>, 2015.
- 635 Gao, M., Gao, J., Zhu, B., Kumar, R., Lu, X., Song, S., Zhang, Y., Jia, B., Wang, P., Beig, G., Hu, J., Ying, Q., Zhang, H., Sherman, P., and McElroy, M. B.: Ozone pollution over China and India: seasonality and sources, *Atmos. Chem. Phys.*, 20, 4399–4414, <https://doi.org/10.5194/acp-20-4399-2020>, 2020.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.-Atmos.*, 106(D17), 20255-20273, <https://doi.org/10.1029/2000JD000053>, 2001.
- 640 Gong, C., Lei, Y., Ma, Y., Yue, X., and Liao, H.: Ozone–vegetation feedback through dry deposition and isoprene emissions in a global chemistry–carbon–climate model, *Atmos. Chem. Phys.*, 20, 3841–3857, <https://doi.org/10.5194/acp-20-3841-2020>, 2020.
- 645 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model. *Atmos. Environ.*, 39(37), 6957-6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- 650 Hollaway, M. J., Arnold, S. R., Collins, W. J., Folberth, G., and Rap, A.: Sensitivity of midnineteenth century tropospheric ozone to atmospheric chemistry-vegetation interactions, *J. Geophys. Res.-Atmos.*, 122, 2452–2473, <https://doi.org/10.1002/2016JD025462>, 2017.

- Hu, J., Chen, J., Ying, Q., and Zhang, H.: One-year simulation of ozone and particulate matter in China using WRF/CMAQ modeling system, *Atmos. Chem. Phys.*, 16, 10333–10350, <https://doi.org/10.5194/acp-16-10333-2016>, 2016.
- 655 Huang, L., McDonald-Buller, E. C., McGaughey, G., Kimura, Y., and Allen, D. T.: The impact of drought on ozone dry deposition over eastern Texas. *Atmos. Environ.* 127, 176–186, 2016.
- Huntingford, C., Oliver, R. J., Mercado, L. M., and Sitch, S.: A simple theoretical model framework to describe plant stomatal "sluggishness" in response to elevated ozone concentrations, *Biogeosciences*, 15, 5415–5422, <https://doi.org/10.5194/bg-15-5415-2018>, 2018.
- 660 Kinose Y., Fukamachi Y., Okabe S., Hiroshima H., Watanabe M., and Izuta T.: Photosynthetic responses to ozone of upper and lower canopy leaves of *Fagus crenata* Blume seedlings grown under different soil nutrient conditions, *Environ. Pollut.*, 223, 213–222, <https://doi.org/10.1016/j.envpol.2017.01.014>, 2017.
- Lei, Y., Yue, X., Liao, H., Gong, C., and Zhang, L.: Implementation of Yale Interactive terrestrial Biosphere model v1. 0 into GEOS-Chem v12. 0.0: a tool for biosphere–chemistry interactions, *Geosci. Model Dev.*, 13, 1137–1153, <https://doi.org/10.5194/gmd-13-1137-2020>, 2020.
- 665 Li, J., Mahalov, A., and Hyde, P.: Simulating the impacts of chronic ozone exposure on plant conductance and photosynthesis, and on the regional hydroclimate using WRF/Chem, *Environ. Res. Lett.*, 11, 114017, <https://doi.org/10.1088/1748-9326/11/11/114017>, 2016.
- 670 Li, J., Mahalov, A., and Hyde, P.: Simulating the effects of chronic ozone exposure on hydrometeorology and crop productivity using a fully coupled crop, meteorology and air quality modeling system, *Agr. Forest Meteorol.*, 260, 287–299, <https://doi.org/10.1016/j.agrformet.2018.06.013>, 2018.
- Li, K., Jacob, D. J., Liao, H., Shen, L., Zhang, Q., and Bates, K. H.: Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China, *PNAS*, 116(2), 422–427, <https://doi.org/10.1073/pnas.1812168116>, 2019.
- 675 Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K. H., Zhang, Q., and Zhai, S.: A two-pollutant strategy for improving ozone and particulate air quality in China, *Nat. Geosci.*, 12, 906–910, <https://doi.org/10.1038/s41561-019-0464-x>, 2019.
- Lin, M., Horowitz, L. W., Xie, Y., Paulot, F., Malyshev, S., Shevliakova, E., Finco, A., Gerosa, G., Kubistin, D., and Pilegaard, K.: Vegetation feedbacks during drought exacerbate ozone air pollution extremes in Europe. *Nature Climate Change*, 10(5), 444–451, 2020.
- 680 Lin, M., Malyshev, S., Shevliakova, E., Paulot, F., Horowitz, L. W., Fares, S., Mikkelsen, T. N., and Zhang, L.: Sensitivity of ozone dry deposition to ecosystem–atmosphere interactions: A critical appraisal of observations and simulations. *Global Biogeochemical Cycles*, 33(10), 1264–1288, 2019.
- 685 Liu, Y. and Wang, T.: Worsening urban ozone pollution in China from 2013 to 2017 – Part 1: The complex and varying roles of meteorology, *Atmos. Chem. Phys.*, 20, 6305–6321, <https://doi.org/10.5194/acp-20-6305-2020>, 2020.
- Liu, Z., Yim, S. H., Wang, C., and Lau, N. C.: The impact of the aerosol direct radiative forcing on deep convection and air quality in the Pearl River Delta region, *Geophys. Res. Lett.*, 45(9), 4410–4418, <https://doi.org/10.1029/2018GL077517>, 2018.
- 690 Liu, Z., Ming, Y., Zhao, C., Lau, N. C., Guo, J., Bollasina, M., and Yim, S. H. L.: Contribution of local and remote anthropogenic aerosols to a record-breaking torrential rainfall event in Guangdong Province, China, *Atmos. Chem. Phys.*, 20, 223–241, <https://doi.org/10.5194/acp-20-223-2020>, 2020.
- Lombardozzi, D., Levis, S., Bonan, G., and Sparks, J. P.: Predicting photosynthesis and transpiration responses to ozone: decoupling modeled photosynthesis and stomatal conductance, *Biogeosciences*, 9, 3113–3130, <https://doi.org/10.5194/bg-9-3113-2012>, 2012.
- 695

- Lombardozi, D., Sparks, J. P., and Bonan, G.: Integrating O₃ influences on terrestrial processes: photosynthetic and stomatal response data available for regional and global modeling, *Biogeosciences*, 10, 6815–6831, <https://doi.org/10.5194/bg-10-6815-2013>, 2013.
- 700 Lombardozi, D., Levis, S., Bonan, G., Hess, P. G., and Sparks, J. P.: The influence of chronic ozone exposure on global carbon and water cycles, *J. Climate*, 28, 292–305, <https://doi.org/10.1175/JCLI-D-14-00223.1>, 2015.
- Lu, X., Hong, J., Zhang, L., Cooper, O. R., Schultz, M. G., Xu, X., Wang, T., Gao, M., Zhao, Y. H., and Zhang, Y.: Severe surface ozone pollution in China: A global perspective, *Environ. Sci. Technol. Lett.* 2018, 5, 8, 487–494, <https://doi.org/10.1021/acs.estlett.8b00366>, 2018.
- 705 Niu, G. Y., Yang, Z. L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D., Rosero, E., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements, *J. Geophys. Res.-Atmos.*, 116(D12), <https://doi.org/10.1029/2010JD015139>, 2011.
- 710 Rydsaa, J. H., Stordal, F., Gerosa, G., Finco, A., and Hodnebrog, O. Evaluating stomatal ozone fluxes in WRF-Chem: Comparing ozone uptake in Mediterranean ecosystems. *Atmos. Environ.* 143, 237–248, 2016.
- Sadiq, M., Tai, A. P. K., Lombardozi, D., and Val Martin, M.: Effects of ozone–vegetation coupling on surface ozone air quality via biogeochemical and meteorological feedbacks, *Atmos. Chem. Phys.*, 17, 3055–3066, <https://doi.org/10.5194/acp-17-3055-2017>, 2017.
- 715 Schell, B., Ackermann, I. J., Hass, H., Binkowski, F. S., and Ebel, A.: Modeling the formation of secondary organic aerosol within a comprehensive air quality model system., *J. Geophys. Res.-Atmos.*, 106(D22), 28275–28293, <https://doi.org/10.1029/2001JD000384>, 2001.
- Silver, B., Reddington, C. L., Arnold, S. R., and Spracklen, D. V.: Substantial changes in air pollution across China during 2015–2017, *Environ. Res. Lett.* 13 114012, <https://orcid.org/0000-0003-0395-0637>, 2018.
- 720 Sitch, S., Cox, P. M., Collins, W. J., and Huntingford, C.: Indirect radiative forcing of climate change through ozone effects on the land-carbon sink, *Nature*, 448, 791–794, <https://doi.org/10.1038/nature06059>, 2007.
- 725 Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsoren, S. B., Myhre, G., Berntsen, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L., Lee, Y. H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K. W., Wild, O., and Archibald, A.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 3063–3085, <https://doi.org/10.5194/acp-13-3063-2013>, 2013
- 730 Stockwell, W. R., Middleton, P., Chang, J. S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling, *J. Geophys. Res.-Atmos.*, 95(D10), 16343–16367, <https://doi.org/10.1029/JD095iD10p16343>, 1990.
- 735 Tai, A. P., Martin, M. V., and Heald, C. L.: Threat to future global food security from climate change and ozone air pollution. *Nature Clim. Change*, 4, 817–821, <https://doi.org/10.1038/nclimate2317>, 2014.
- Uddling, J., Matyssek, R., Pettersson, J. B., and Wieser, G.: To what extent do molecular collisions arising from water vapour efflux impede stomatal O₃ influx?, *Environ. Pollut.*, 170, 39–42, <https://doi.org/10.1016/j.envpol.2012.05.028>, 2012.
- 740

Wang, W. N., Cheng, T. H., Gu, X. F., Chen, H., Guo, H., Wang, Y., Bao, F. W., Shi, S. Y., Xu, B. R., Meng, C., and Meng, C.: Assessing spatial and temporal patterns of observed ground-level ozone in China. *Sci. Rep.*, 7, 1-12, <https://doi.org/10.1038/s41598-017-03929-w>, 2017.

745 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.

Xie, X., Wang, T., Yue, X., Li, S., Zhuang, B., Wang, M., and Yang, X.: Numerical modeling of ozone damage to plants and its effects on atmospheric CO₂ in China. *Atmos. Environ.*, 217, 116970, <https://doi.org/10.1016/j.atmosenv.2019.116970>, 2019.

750 Yue, X. and Unger, N.: Ozone vegetation damage effects on gross primary productivity in the United States, *Atmos. Chem. Phys.*, 14, 9137–9153, <https://doi.org/10.5194/acp-14-9137-2014>, 2014.

Yue, X., Unger, N., Harper, K., Xia, X., Liao, H., Zhu, T., Xiao, J., Feng, Z., and Li, J.: Ozone and haze pollution weakens net primary productivity in China, *Atmos. Chem. Phys.*, 17, 6073–6089, <https://doi.org/10.5194/acp-17-6073-2017>, 2017.

755 Zhou, S. S., Tai, A. P. K., Sun, S., Sadiq, M., Heald, C. L., and Geddes, J. A.: Coupling between surface ozone and leaf area index in a chemical transport model: strength of feedback and implications for ozone air quality and vegetation health, *Atmos. Chem. Phys.*, 18, 14133–14148, <https://doi.org/10.5194/acp-18-14133-2018>, 2018.

760

765

770

775

780

Table 1. Slopes (per mmol m^{-2}) and intercepts (unitless) used for O_3 damage factors in Eqs. (4) and (5), following Lombardozzi et al. (2015).

790

| | Photosynthesis | | Conductance | |
|-------------------|-----------------|---------------------|-----------------|---------------------|
| | Slope (a_p) | Intercept (b_p) | Slope (a_c) | Intercept (b_c) |
| Broadleaf | 0.0000 | 0.8752 | 0.0000 | 0.9125 |
| Needleleaf | 0.0000 | 0.8390 | 0.0048 | 0.7823 |
| Grasses and crops | -0.0009 | 0.8021 | 0.0000 | 0.7511 |

Table 2. Description of the two sets of model experiments.

| Experiment name | Year | Anthropogenic Emission | Meteorological ICs and BCs |
|---------------------------|---------------|------------------------|----------------------------|
| simu_without O_3 | 2014–2017 JJA | Year 2014 | FNL |
| simu_with O_3 | 2014–2017 JJA | Year 2014 | FNL |

795

Table 3. Evaluation results for the temperature at 2 meter (T_{2m}), relative humidity at 2 meter (RH_{2m}) and wind speed at 10 meter (WS_{10m}) for different years in China. Mean_obs (Mean_simu) is the mean value of observation (model simulation); SD_obs (SD_simu) is the standard deviation of the observation (model simulation); IOA is the index of agreement; CORR is the correlation coefficient; MB is the mean bias.

800

| | Year | Mean_obs | SD_obs | Mean_simu | SD_simu | IOA | CORR | MB |
|--|------|----------|--------|-----------|---------|------|------|-------|
| T_{2m} ($^{\circ}\text{C}$) | 2014 | 25.41 | 2.61 | 24.71 | 2.27 | 0.86 | 0.87 | -0.70 |
| | 2015 | 25.41 | 2.56 | 24.67 | 2.24 | 0.86 | 0.89 | -0.74 |
| | 2016 | 26.35 | 2.82 | 25.44 | 2.61 | 0.85 | 0.85 | -0.91 |
| | 2017 | 26.29 | 3.17 | 25.28 | 3.16 | 0.81 | 0.78 | -1.00 |
| RH_{2m} (%) | 2014 | 74.77 | 10.22 | 79.14 | 8.96 | 0.67 | 0.71 | 4.38 |
| | 2015 | 73.34 | 10.75 | 80.50 | 8.73 | 0.68 | 0.75 | 7.16 |
| | 2016 | 74.14 | 10.81 | 81.47 | 10.10 | 0.70 | 0.73 | 7.33 |
| | 2017 | 73.24 | 11.65 | 79.89 | 9.62 | 0.68 | 0.69 | 6.63 |
| WS_{10m} (m s^{-1}) | 2014 | 1.84 | 0.66 | 2.22 | 1.16 | 0.54 | 0.40 | 0.38 |
| | 2015 | 2.00 | 0.74 | 2.48 | 1.35 | 0.55 | 0.44 | 0.48 |
| | 2016 | 1.99 | 0.70 | 2.47 | 1.32 | 0.54 | 0.45 | 0.48 |
| | 2017 | 2.02 | 0.72 | 2.51 | 1.42 | 0.53 | 0.45 | 0.50 |

805

810

Table 4. Evaluation results of temperature at 2 meter (T_{2m}), relative humidity at 2 meter (RH_{2m}) and wind speed at 10 meter (WS_{10m}) in 7 major geographic regions from the implemented model. NE is northeast China, NC is north China, CC is central China, EC is east China, SC is south China, SW indicates southwest China, and NW is northwest China. Mean_obs (Mean_simu) is the mean value of observations (model simulations); SD_obs (SD_simu) is the standard deviation of the observations (model simulations); IOA is the index of agreement; CORR is the correlation coefficient; MB is the mean bias.

| | Region | Mean_obs | SD_obs | Mean_simu | SD_simu | IOA | CORR | MB |
|---|--------|----------|--------|-----------|---------|------|------|-------|
| T_{2m} (°C) | NEC | 23.01 | 3.05 | 22.73 | 3.01 | 0.94 | 0.91 | -0.28 |
| | NC | 24.94 | 2.76 | 25.84 | 2.84 | 0.86 | 0.88 | 0.88 |
| | CC | 27.62 | 3.05 | 26.87 | 2.75 | 0.92 | 0.88 | -0.75 |
| | EC | 27.33 | 2.99 | 26.46 | 2.58 | 0.90 | 0.89 | -0.87 |
| | SC | 28.60 | 1.49 | 28.61 | 1.30 | 0.75 | 0.64 | 0.01 |
| | SWC | 23.20 | 2.32 | 21.61 | 2.14 | 0.77 | 0.80 | -1.58 |
| | NWC | 20.20 | 2.87 | 18.55 | 3.01 | 0.77 | 0.89 | -1.65 |
| RH_{2m} (%) | NEC | 71.7 | 11.49 | 71.98 | 14.00 | 0.85 | 0.79 | 0.93 |
| | NC | 63.25 | 13.94 | 57.01 | 14.08 | 0.79 | 0.75 | -6.24 |
| | CC | 79.23 | 10.11 | 88.29 | 8.61 | 0.70 | 0.71 | 9.06 |
| | EC | 78.93 | 9.99 | 88.80 | 8.31 | 0.69 | 0.79 | 9.87 |
| | SC | 81.26 | 6.54 | 88.41 | 5.66 | 0.62 | 0.60 | 7.14 |
| | SWC | 78.92 | 9.11 | 93.34 | 5.13 | 0.52 | 0.64 | 13.40 |
| | NWC | 57.93 | 13.34 | 58.48 | 14.10 | 0.75 | 0.76 | 0.55 |
| WS_{10m} ($m\ s^{-1}$) | NEC | 2.22 | 0.93 | 3.08 | 1.80 | 0.62 | 0.62 | 0.86 |
| | NC | 2.06 | 0.72 | 2.45 | 1.29 | 0.57 | 0.48 | 0.38 |
| | CC | 2.06 | 0.81 | 2.38 | 1.41 | 0.61 | 0.51 | 0.33 |
| | EC | 2.18 | 0.76 | 2.85 | 1.54 | 0.59 | 0.55 | 0.67 |
| | SC | 2.02 | 0.76 | 2.81 | 1.51 | 0.52 | 0.43 | 0.80 |
| | SWC | 2.16 | 0.76 | 2.54 | 1.40 | 0.57 | 0.51 | 0.37 |
| | NWC | 1.46 | 0.50 | 2.91 | 1.38 | 0.30 | 0.23 | 1.45 |

815

820

825

Table 5. Evaluation results for the air pollutants in China. Mean_obs (Mean_simu) is the mean value of observation (model simulation); SD_obs (SD_simu) is the standard deviation of the observation (model simulation); IOA is the index of agreement; CORR is the correlation coefficient; MB is the mean bias.

| | Year | Mean_obs | SD_obs | Mean_simu | SD_simu | IOA | CORR | MB |
|---|------|----------|--------|-----------|---------|------|------|-------|
| O₃ (ppb) | 2014 | 29.79 | 9.95 | 51.49 | 18.60 | 0.48 | 0.57 | 22.13 |
| | 2015 | 32.04 | 10.16 | 48.98 | 18.27 | 0.54 | 0.55 | 16.95 |
| | 2016 | 33.28 | 10.59 | 48.47 | 18.18 | 0.56 | 0.58 | 15.14 |
| | 2017 | 35.74 | 11.71 | 49.50 | 19.61 | 0.63 | 0.66 | 13.82 |
| PM_{2.5} ($\mu\text{g m}^{-3}$) | 2014 | 46.30 | 21.52 | 63.28 | 27.15 | 0.52 | 0.33 | 18.61 |
| | 2015 | 38.52 | 17.30 | 55.56 | 24.85 | 0.55 | 0.42 | 16.66 |
| | 2016 | 31.86 | 13.96 | 56.70 | 25.69 | 0.47 | 0.40 | 24.54 |
| | 2017 | 28.82 | 12.23 | 56.34 | 25.70 | 0.40 | 0.30 | 27.65 |
| PM₁₀ ($\mu\text{g m}^{-3}$) | 2014 | 80.79 | 31.62 | 71.74 | 28.65 | 0.47 | 0.22 | -7.51 |
| | 2015 | 72.03 | 29.74 | 63.83 | 26.29 | 0.50 | 0.26 | -8.93 |
| | 2016 | 59.68 | 22.21 | 65.01 | 27.29 | 0.49 | 0.24 | 4.65 |
| | 2017 | 57.83 | 22.18 | 64.78 | 27.25 | 0.41 | 0.14 | 6.95 |
| SO₂ (ppb) | 2014 | 6.11 | 2.36 | 8.41 | 3.22 | 0.48 | 0.41 | 2.36 |
| | 2015 | 4.78 | 1.89 | 8.39 | 3.26 | 0.44 | 0.45 | 3.64 |
| | 2016 | 4.17 | 1.57 | 8.08 | 3.16 | 0.41 | 0.36 | 3.92 |
| | 2017 | 3.83 | 1.33 | 8.58 | 3.52 | 0.36 | 0.42 | 4.78 |
| NO₂ (ppb) | 2014 | 17.20 | 4.51 | 17.23 | 4.63 | 0.41 | 0.26 | 0.06 |
| | 2015 | 16.01 | 4.47 | 17.37 | 4.98 | 0.43 | 0.31 | 1.43 |
| | 2016 | 15.29 | 4.29 | 17.35 | 5.11 | 0.43 | 0.31 | 2.06 |
| | 2017 | 15.83 | 4.37 | 17.84 | 5.12 | 0.43 | 0.32 | 2.02 |
| CO (ppm) | 2014 | 0.76 | 0.19 | 0.44 | 0.11 | 0.48 | 0.42 | -0.32 |
| | 2015 | 0.67 | 0.15 | 0.45 | 0.11 | 0.49 | 0.42 | -0.22 |
| | 2016 | 0.65 | 0.14 | 0.45 | 0.11 | 0.50 | 0.45 | -0.20 |
| | 2017 | 0.64 | 0.12 | 0.46 | 0.11 | 0.47 | 0.38 | -0.18 |

830

835

840 **Table 6.** Evaluation results of air pollutants in 7 major geographic regions simulated by the implemented
model. NE is northeast China, NC is north China, CC is central China, EC is east China, SC is south
China, SW indicates southwest China, and NW is northwest China. Mean_obs (Mean_simu) is the mean
value of observations (model simulations); SD_obs (SD_simu) is the standard deviation of the
observations (model simulations); IOA is the index of agreement; CORR is the correlation coefficient;
845 MB is the mean bias.

| | Region | Mean_obs | SD_obs | Mean_simu | SD_simu | IOA | CORR | MB |
|---|--------|----------|--------|-----------|---------|------|-------|--------|
| O₃ (ppb) | NEC | 32.49 | 10.54 | 44.54 | 15.70 | 0.64 | 0.64 | 11.47 |
| | NC | 38.56 | 10.81 | 70.59 | 25.11 | 0.40 | 0.55 | 32.02 |
| | CC | 31.68 | 11.20 | 57.13 | 18.17 | 0.47 | 0.60 | 26.82 |
| | EC | 29.67 | 10.82 | 40.53 | 18.99 | 0.60 | 0.61 | 11.21 |
| | SC | 19.90 | 8.40 | 34.21 | 14.68 | 0.53 | 0.64 | 14.90 |
| | SWC | 24.27 | 9.12 | 42.07 | 13.19 | 0.47 | 0.50 | 18.80 |
| | NWC | 26.63 | 8.12 | 51.65 | 13.70 | 0.34 | 0.42 | 25.58 |
| PM_{2.5} (µg m⁻³) | NEC | 42.66 | 25.15 | 43.33 | 19.28 | 0.57 | 0.39 | -0.83 |
| | NC | 61.60 | 28.28 | 66.83 | 27.81 | 0.68 | 0.52 | 7.03 |
| | CC | 52.11 | 30.55 | 94.27 | 39.40 | 0.35 | 0.11 | 45.50 |
| | EC | 52.87 | 25.71 | 87.37 | 38.97 | 0.50 | 0.39 | 36.63 |
| | SC | 22.58 | 10.16 | 28.62 | 15.17 | 0.67 | 0.57 | 6.92 |
| | SWC | 32.82 | 12.22 | 76.69 | 33.49 | 0.27 | 0.08 | 47.55 |
| | NWC | 45.27 | 14.54 | 42.80 | 14.07 | 0.39 | 0.03 | -1.45 |
| PM₁₀ (µg m⁻³) | NEC | 79.68 | 36.48 | 48.99 | 20.64 | 0.49 | 0.32 | -32.25 |
| | NC | 111.17 | 39.69 | 74.29 | 29.33 | 0.54 | 0.38 | -35.90 |
| | CC | 84.80 | 41.01 | 107.65 | 41.51 | 0.37 | 0.05 | 26.30 |
| | EC | 78.16 | 35.64 | 99.51 | 40.90 | 0.54 | 0.34 | 23.64 |
| | SC | 43.64 | 15.72 | 34.11 | 16.14 | 0.58 | 0.47 | -8.54 |
| | SWC | 58.84 | 20.15 | 87.07 | 35.49 | 0.31 | -0.07 | 32.17 |
| | NWC | 88.54 | 28.17 | 47.77 | 14.72 | 0.35 | -0.13 | -39.68 |
| SO₂ (ppb) | NEC | 4.91 | 1.95 | 5.10 | 2.55 | 0.60 | 0.42 | 0.27 |
| | NC | 8.69 | 3.52 | 8.12 | 3.20 | 0.54 | 0.40 | -0.57 |
| | CC | 7.34 | 2.09 | 14.56 | 5.45 | 0.36 | 0.47 | 7.23 |
| | EC | 5.39 | 2.24 | 7.86 | 3.19 | 0.57 | 0.53 | 2.40 |
| | SC | 3.50 | 0.89 | 4.15 | 1.52 | 0.42 | 0.50 | 0.71 |
| | SWC | 4.74 | 1.93 | 15.71 | 5.12 | 0.31 | 0.13 | 11.42 |
| | NWC | 6.65 | 2.90 | 4.31 | 1.50 | 0.46 | 0.34 | -2.28 |
| NO₂ (ppb) | NEC | 19.51 | 4.84 | 14.07 | 5.27 | 0.41 | 0.11 | -5.66 |
| | NC | 19.57 | 5.13 | 14.05 | 4.14 | 0.48 | 0.27 | -5.61 |
| | CC | 16.75 | 4.32 | 19.70 | 5.57 | 0.38 | 0.32 | 2.83 |
| | EC | 16.24 | 4.78 | 28.83 | 6.88 | 0.40 | 0.39 | 12.65 |
| | SC | 13.23 | 3.48 | 14.02 | 3.96 | 0.38 | 0.29 | 1.01 |
| | SWC | 17.30 | 3.70 | 20.02 | 4.58 | 0.34 | 0.12 | 3.11 |
| | NWC | 16.93 | 4.77 | 8.92 | 2.11 | 0.41 | 0.19 | -7.98 |
| CO (ppm) | NEC | 0.64 | 0.17 | 0.38 | 0.12 | 0.48 | 0.61 | -0.27 |
| | NC | 0.90 | 0.25 | 0.47 | 0.13 | 0.47 | 0.41 | -0.42 |
| | CC | 0.81 | 0.17 | 0.58 | 0.14 | 0.49 | 0.45 | -0.22 |
| | EC | 0.66 | 0.16 | 0.56 | 0.14 | 0.63 | 0.54 | -0.09 |
| | SC | 0.66 | 0.11 | 0.32 | 0.08 | 0.36 | 0.41 | -0.34 |
| | SWC | 0.69 | 0.14 | 0.49 | 0.11 | 0.49 | 0.29 | -0.19 |
| | NWC | 0.89 | 0.25 | 0.25 | 0.04 | 0.35 | 0.18 | -0.63 |




Figure 1. The vegetation fraction of (a) broadleaf, (b) needleleaf, (c) cropland, (d) grass, (e) others, and (f) dominant vegetation types.

850




Figure 2. Site locations of air quality monitoring sites (blue dots) and the meteorological monitoring sites (pink dots) with the underlying is the terrain height (m).

855




Figure 3. Spatial distribution of mean stomatal resistance in JJA of 2014–2017 for (a) sunlit leaves (RSSUN) and (d) shaded leaves (RSSHA) from the *simu_withoutO₃* experiment. Absolute changes in (b) RSSUN and (e) RSSHA caused by O₃ damage. Relative changes in (c) RSSUN and (f) RSSHA caused by O₃ damage. Absolute changes are the RSSUN (RSSHA) from *simu_withO₃* minus RSSUN (RSSHA) from *simu_withoutO₃*. Relative changes are calculated by absolute changes over the RSSUN (RSSHA) from *simu_withoutO₃*.

860

865





Figure 4. Spatial distribution of 2014–2017 JJA mean **(a)** photosynthesis rate (PSN), **(d)** leaf area index (LAI), and **(g)** gross primary productivity (GPP) from the simu_withoutO₃ experiment; absolute changes in **(b)** PSN, **(e)** LAI and **(h)** GPP caused by O₃ damage; and relative changes in **(c)** PSN, **(f)** LAI and **(i)** GPP caused by O₃ damage. Absolute changes are the results from simu_withO₃ minus results from simu_withoutO₃. Relative changes are calculated from the absolute changes over the results from simu_withoutO₃.


870

875

880



885 **Figure 5.** Spatial distribution of 2014–2017 JJA mean **(a)** transpiration rate (TR), and **(b)** absolute changes and **(c)** relative changes in TR caused by O₃ damage.



890 **Figure 6.** Spatial distribution of mean **(a)** latent heat flux (LH) and **(d)** sensitive heat flux (SH) from the simu_withoutO₃ experiment; absolute changes in **(b)** LH flux and **(e)** SH flux in JJA of 2014–2017 caused by O₃ damage; and relative changes in **(c)** LH flux and **(f)** SH flux caused by O₃ damage. Absolute changes are the LH (SH) flux from simu_withO₃ minus LH (SH) flux simu_withoutO₃. Relative changes are calculated by absolute changes over LH (SH) flux from simu_withoutO₃.

895




Figure 7. Spatial distribution of mean **(a)** 2-m relative humidity, **(d)** 2-m temperature at, and **(g)** planetary boundary layer height (PBLH) in JJA of 2014–2017 from the `simu_withoutO3` experiment; absolute changes in **(b)** RH_{2m} , **(e)** T_{2m} and **(h)** PBLH caused by O_3 damage; and relative changes in **(c)** RH_{2m} , **(f)** T_{2m} and **(i)** PBLH caused by O_3 damage. Absolute changes are the results from `simu_withO3` minus results from `simu_withoutO3`. Relative changes are calculated by absolute changes over the results from `simu_withoutO3`.

900

905

910




Figure 8. Same as Fig 5 but for surface O_3 concentration.

915




Figure 9. Same as Fig 5 but for isoprene emission.

920