

# Mass and Density of Individual Frozen Hydrometeors

Karlie N. Rees<sup>1</sup>, Dhiraj K. Singh<sup>2</sup>, Eric R. Pardyjak<sup>2</sup>, and Timothy J. Garrett<sup>1</sup>

<sup>1</sup>Department of Atmospheric Sciences, Salt Lake City, UT, USA

<sup>2</sup>Department of Mechanical Engineering, Salt Lake City, UT, USA

**Correspondence:** Tim Garrett (tim.garrett@utah.edu)

**Abstract.** A new precipitation sensor, the Differential Emissivity Imaging Disdrometer (DEID), is used to provide the first continuous measurements of the mass, diameter, and density of individual hydrometeors. The DEID consists of an infrared camera pointed at a heated aluminum plate. It exploits the contrasting thermal emissivity of water and metal to determine individual particle mass by assuming that energy is conserved during the transfer of heat from the plate to the particle during evaporation. Particle density is determined from a combination of particle mass and morphology. A Multi-Angle Snowflake Camera (MASC) was deployed alongside the DEID to provide refined imagery of particle size and shape. Broad consistency is found between derived mass-diameter and density-diameter relationships and those obtained in prior studies. However, DEID measurements show a generally weaker dependence with size for hydrometeor density and a stronger dependence for aggregate snowflake mass.

10 *Copyright statement.* TEXT

## 1 Introduction

Predictions of precipitation amount, location, and duration have been shown to be especially sensitive to parameterized expressions for how fast a hydrometeor falls (Rutledge and Hobbs, 1984; Reisner et al., 1998; Hong et al., 2004; Fovell and Su, 2007; Lin et al., 2010; Liu et al., 2011; Iguchi et al., 2012; Thériault et al., 2012), affecting forecasts of hurricane trajectories (Fovell and Su, 2007) and storm lifetimes (Garvert et al., 2005; Colle et al., 2005; Milbrandt et al., 2010). From the perspective of fluid dynamics, fallspeed can be related to the mass and density of precipitation particles (Böhm, 1989). Observationally, one of the most frequently cited datasets that lies at the heart of current bulk microphysical parameterizations (e.g. Reisner et al., 1998; Hong et al., 2004; Tao et al., 2003) comprises just 376 snowflakes captured and photographed in the Cascade mountain range. Individual hydrometeors were melted on a sheet of plastic film, from which relationships were obtained between hydrometeor mass, fallspeed, and diameter as a function of particle habit, and in the case of graupel, density (Locatelli and Hobbs, 1974). Despite its limited scope, later numerical studies (Böhm, 1989; Khvorostyanov and Curry, 2002; Heymsfield and Westbrook, 2010; Kubicek and Wang, 2012) and ground-based disdrometer measurements (Kruger and Krajewski, 2002; Barthazy et al., 2004; Yuter et al., 2006; Newman et al., 2009) have lent general support to this reference data set, although concerns remain

about geographic and temporal specificity, measurement limitations (Yuter et al., 2006; Battaglia et al., 2010), and the ability  
25 to quantify the extent of riming (Barthazy and Schefold, 2006; Brandes et al., 2008).

Particle density measurements have proven more difficult to ascertain as they require, in addition to mass, an estimate of  
particle volume. This may be reasonably obtained for quasi-spherical particles such as lump graupel (Locatelli and Hobbs,  
1974), but the task is considerably more challenging when snow particles are formed from ice crystal aggregation. A possible  
approach is to infer density from fallen snow using column measurements (Conger and McClung, 2009), capacitance probes  
30 (Dent et al., 1998), or a combination of a camera for snow depth and an electric scale for snow mass (Muramoto et al., 1995).  
Muramoto et al. (1995) and Brandes et al. (2007) measured the bulk mass of snowflakes using a weighing gauge, from which  
the bulk volume was determined with 2D camera imagery of individual snowflakes. Tiira et al. (2016) determined a volume  
flux weighted snow density for a population of snowflakes using particle size distribution, fallspeed, and a weighing gauge to  
estimate the mass, and 2D camera imagery to determine effective diameter and volume. However, snow undergoes compaction  
35 and melting on the ground, so the relationship to individual particle density in the air is approximate (Brun et al., 1992). To  
obtain individual snowflake particle density, Magono and Nakamura (1965) collected individual wet and dry snowflakes on a  
piece of dyed filter paper, from which the outline of the flake was manually measured. Individual volume was inferred from the  
major and minor axes of the outline of snow, and the mass from the outline of the melted snowflake. Holroyd (1971) also made  
measurements of the major and minor axes of powder snow and dendrites from (Magono and Nakamura, 1965). A limitation  
40 of these datasets is that they were necessarily small given the manual nature of the effort.


Here, we present continuous measurements of individual masses of frozen hydrometeors using a new instrument, the Differential  
Emissivity Imaging Disdrometer (DEID). DEID data are combined with photographic imagery obtained using a Multi  
Angle Snowflake Camera (MASC) (Garrett et al., 2012) to obtain estimates of particle density.

## 2 Methods

45 All measurements described in this study were acquired at a meteorological measurement tower placed at the mouth of Red  
Butte Canyon (40.76857, -111.82614) in Salt Lake City, Utah at 1547 m elevation in January and February 2020. More details  
about the site and measurement campaign are provided by Singh et al. (2021).

### 2.1 Particle mass from thermal imaging

The Differential Emissivity Imaging Disdrometer (DEID) (Singh et al., 2021) consists of a thermal camera operating at a  
50 frequency between 2 and 12 Hz pointed at an aluminum plate placed atop a hotplate maintained at a self-sustained temperature  
of 85°C. The camera distinguishes hydrometeors as they melt and evaporate as white regions on a black background (Figure  
1) due to the contrasting infrared emissivities of water ( $\epsilon \approx 0.96$ ) and aluminum ( $\epsilon \approx 0.03$ ). A small strip of polyimide tape  
( $\epsilon \approx 0.95$ ) is applied to the aluminum plate to provide a reference temperature, as well as a pixel to length dimension conversion  
based on the tape's known width. The effective collection area of the DEID was  $A \approx 7 \text{ cm} \times 5 \text{ cm}$  and the per pixel resolution



**Figure 1.** Thermal camera imagery from the DEID, showing regions of high emissivity water (white) on a low emissivity aluminum plate (black).

55 of imaged particles  $p$  was  $190 \mu\text{m}$ . Processing of the thermal camera imagery yields the hydrometeor area on the plate, the temperature difference between the water and the plate, and the evaporation time.

Individual particle mass is determined using the DEID by employing the assumption that the heat gained by a hydrometeor is equivalent to the heat lost by the plate during the process of melting and evaporation. The heat balance equation is:

$$c_p \Delta T \int dM + L_{eqv} \int dM = \int_0^t \frac{K}{H} A(t) (T_p - T_w(t)) dt \quad (1)$$


60 where  $c_p$  is the specific heat capacity of water at constant pressure,  $\Delta T$  is the difference in temperature between 0 and time,  $t$ ,  $M$  is the mass of the hydrometeor, and  $L_{eqv}$  is the equivalent latent heat required for the conversion of the hydrometeor to gas. For liquid precipitation  $L_{eqv} = L_v$  where  $L_v$  is the latent heat of vaporization of water. For solid precipitation,  $L_{eqv} = L_v + L_f$  where  $L_f$  is the latent heat fusion for water.  $K$  is the thermal conductivity of the aluminum plate,  $H$  is the plate thickness,  $A(t)$  is the cross-sectional area of the water droplet at time  $t$ ,  $T_p$  is the temperature of the hotplate, and  $T_w(t)$  is the temperature of  
65 the water at time  $t$ . Taking

$$\int dM = K_d \int_0^t A(t) (T_p - T_w(t)) dt$$

a single calibrated constant  $K_d$  for the plate was determined experimentally by applying known masses of water from a micropipette to the plate. (See Appendix A for the derivation of  $K_d$ ). The heat balance equation was found in a laboratory setting to be highly insensitive to ambient winds, temperature, and humidity (Singh et al., 2021).

70 The temperature difference between the plate and water on the plate ( $\Delta T = T_p - T_w(t)$ ) can be determined using the mean pixel intensity of the particle in the thermal camera imagery (see Appendix B). The mass calculation then simplifies to:

$$M = K_d \Delta T \int A(t) dt \quad (2)$$



**Figure 2.** Evaporation volumetric profiles on the DEID plate in a space of imaged area and time  $V_t = \int A(t)dt$  for graupel.

Three-dimensional regions integrated over particle area and evaporation time were constructed for each particle during the melting process defined by  $V_t = \int A(t)dt$  so that from Eq. 2,  $M = K_d V_t \Delta T$ . An example of a one-minute time interval with 3D volumetric regions representing each individual particle is shown in Figure 2 for graupel. Additional 3D representations for other snowflake types are shown in Appendix B.


## 2.2 Hydrometeor photography

The Multi-Angle Snowflake Camera (MASC) (Garrett et al., 2012) houses three high-speed visible spectrum cameras arranged concentrically with a separation angle of  $36^\circ$ . As a particle falls through the instrument collection aperture, two vertically spaced infrared detectors simultaneously trigger the cameras to take three simultaneous photographs of the particle from the side and to measure the fallspeed between the two detectors. The MASC software processes the imagery to calculate the surface area, geometric cross-section, perimeter, orientation, aspect ratio, complexity, flatness, and whether the particle is a raindrop (Shkurko et al., 2016). Images taken by the Multi-Angle Snowflake Camera (MASC) are used in conjunction with the DEID to confirm precipitation type and refine density measurements. A mosaic of MASC images from February 5 and 6, 2020 is shown in Figure 3. Additional snowflake imagery catalogued from each storm is shown in Appendix C.


## 2.3 Particle volume and density

Immediately following the arrival of a hydrometeor on the plate, the particle cross-sectional area increases rapidly as the hydrometeor adjusts from the ambient air temperature to the temperature of the plate, melting in the process. It reaches a maximum cross-sectional area  $A_{\max}$  before shrinking during evaporation. A particle may be melted at the moment that it reaches  $A_{\max}$ . Nonetheless, due to surface tension, particles that are initially frozen tend to maintain their shape following melting so that  $A_{\max}$  is approximately representative of the frozen cross-sectional area in air. In calibration, Singh et al. (2021) found that snowflakes undergo only a 5% change in  $D_{\text{eff}}$  during the melting process.

February 5-6, 2020



**Figure 3.** A selection of MASC snowflake images obtained between February 5 and February 6, 2020.



**Figure 4.** Illustration of the diameter and volume calculations by the DEID (left), maximum and minimum theoretical bounds of volume calculations of spheroids (middle) and the corresponding volume adjustment using MASC-derived aspect ratio measurements (right).

To obtain an estimate of particle density from particle mass, a spherical volume  $V_S$  for each hydrometeor can be estimated from an effective diameter  $D_{eff}$  derived from the  $A_{max}$ .  $V_S = \frac{\pi}{6} D_{eff}^3$  where  $D_{eff} = \sqrt{\frac{4}{\pi} A_{max}}$  (Figure 4, left). This geometric definition is consistent with that taken by Locatelli and Hobbs (1974), who prescribed  $D_{eff}$  as “the diameter of the smallest circle into which the aggregate as photographed will fit without changing its density.” Unless otherwise specified, all instances of “diameter” in the text refer to  $D_{eff}$ . An alternative diameter metric is  $D_{max}$  as defined by the circumscribed diameter from the maximum horizontal dimension of the hydrometeor as it lies on the hotplate. The measured relationship between these two metrics is described in Section 3.2.

For a more precise estimate of particle volume, side-viewing MASC imagery of hydrometeors can be used to determine the average aspect ratios of hydrometeors measured over a one-minute time interval. In general, by approximating hydrometeors as an ellipsoid, the volume of a snowflake is  $V_M = \frac{\pi}{6} D_{max} D_{min} D_v$  where  $D_{max}$  is the longest dimension as seen by the DEID,  $D_{min}$  is the shortest, and  $D_v$  is bounded by the two. For example, if the hydrometeor shape is an oblate spheroid, then  $D_v = D_{max}$ , and if a prolate spheroid then  $D_v = D_{min}$  (Figure 4, middle). As a snowflake falls onto the plate, the maximum dimension is expected to lie flat with respect to the plate.  $D_{eff}$  is expected to lie between the maximum and minimum dimensions of the snowflake, so for the more general case of an ellipsoid, a reasonable assumption is that  $D_v \simeq D_{eff}$ .

The issue here is that the DEID is unlikely to provide a measure of the minimum dimension. However, it can be reasonably inferred from side views of hydrometeors provided by the multiple concentric images captured by the MASC. For a given snowflake, the minimum of the aspect ratios for each hydrometeor seen by the multiple MASC cameras as captured within a one minute interval and subsequently averaged  $\alpha_{\min}$ , can reveal a characteristic minimum dimension in the vertical direction for the time period that is otherwise invisible to the DEID (Figure 4, right). Taking  $D_{\min} = \alpha_{\min} D_{\max}$ , the MASC-adjusted volume is then

$$V_M = \frac{\pi}{6} D_{\max}^2 D_{\text{eff}} \alpha_{\min} \quad (3)$$

The spherical and mass-adjusted density of individual hydrometeors can then be calculated from the mass  $M$  and volume as  $\rho_S = M/V_S$  and  $\rho_M = M/V_M$  respectively.


### 3 Results

Over the course of five storms that took place in Salt Lake City, Utah on January 14, 2020, January 17, 2020, January 26, 2020, February 2-3, 2020, and February 5-6, 2020, the DEID detected 132,459 individual hydrometeors. Of those hydrometeors, 104,812 were snowflakes, and the remainder were either rain or a rain-snow mix. A filtering algorithm rejected small hydrometeors with fewer than three contiguous pixels of data in all three dimensions (See Appendix B), leaving a total dataset of 109,316 hydrometeors, of which 86,285 were snowflakes and the remainder rain. Of those snowflakes, density estimates for 43,649 were obtained using corresponding MASC imagery. The density of rain of course is known to be  $1000 \text{ kg m}^{-3}$  and would provide a valuable reference point for this study. Unfortunately, its measurements cannot be addressed using the techniques described here as raindrops do not preserve their area after impact on the plate.

#### 3.1 Time series

Figure 5 shows the mean and the maximum effective diameters of the hydrometeors considered for each one-minute sampling interval. The largest observed maximum diameters occurred during January 14 and January 26, when large aggregates were the primary snow type (See Appendix C for corresponding snowflake imagery). During such periods characterized primarily by aggregate snowflakes, a larger difference was observed between the mean and maximum diameter observed. Periods characterized primarily by graupel, including early January 14 and late January 17, exhibited smaller differences.

Adopting either the spherical volume or the MASC-adjusted volume, the probability density functions for  $\rho_S$  and  $\rho_M$  separated by storm and combined for all storms are shown in Figure 6. Derived density values are approximately log-normally distributed largely ranging between  $10$  and  $100 \text{ kg m}^{-3}$ . The calculated densities differ by approximately a factor of two depending on the calculation method used. For all storms, the mean spherical density was  $38 \text{ kg m}^{-3}$  and the logarithmically-weighted mean was  $35 \text{ kg m}^{-3}$ , while the respective MASC-adjusted density values were  $90 \text{ kg m}^{-3}$  and  $70 \text{ kg m}^{-3}$ . The relative absence in the distribution of high density values derived using the spherical approximation suggests an underestimate given that Locatelli and Hobbs (1974) found lump graupel to have densities ranging as high as  $450 \text{ kg m}^{-3}$  and dense graupel



**Figure 5.** One-minute averaged maximum and mean effective diameters of measured hydrometeors

has also been described by others (Magono and Nakamura, 1965; Holroyd, 1971; Muramoto et al., 1995; Fabry and Szyrmer, 1999; Brandes et al., 2007; Tiira et al., 2016). For example, for the period January 14 1243-1259 MST when MASC imagery showed graupel predominated during a period with temperatures near the melting point, the spherical calculation yielded an average density of  $88 \text{ kg m}^{-3}$  versus  $131 \text{ kg m}^{-3}$  for the MASC related estimates. The respective logarithmically-weighted mean values were 87 and  $121 \text{ kg m}^{-3}$ .


### 3.2 Diameter and aspect ratio

In the following sections, mass-diameter and density-diameter relationships are expressed with respect to  $D_{\text{eff}}$ . By way of reference, for the 43,649 snowflakes included in this study that are categorized by type,  $D_{\text{max}}$  can be correlated with  $D_{\text{eff}}$  through the power law relationship  $D_{\text{max}} = aD_{\text{eff}}^b$  with values for  $a$  and  $b$  summarized in Table 1. In general, the two quantities are highly correlated, and the relationship is nearly linear. For the ensemble taken as a whole, the relationship is  $D_{\text{max}} = 1.16D_{\text{eff}}^{1.04}$  with a square correlation coefficient of  $R^2 = 0.94$ . The average measured aspect ratios seen by the MASC for aggregates, densely rimed particles, and graupel were 0.64 0.65, and 0.82 respectively.

### 150 3.3 Mass-Diameter relationships

The mass-diameter relationship for all snowflakes is shown in Figure 7. The prefactor observed in the mass-diameter relationship is 0.018, which is approximately consistent with the range of values described by Locatelli and Hobbs (1974) for the densely rimed snowflakes (Table 2), comprising 88% of the snowflakes observed in our study. The exponent 2.52, however,






**Figure 6.** Linear probability density functions of snow density measurements using the spherical volume assumption (gray) and the MASC-adjusted volume (blue) separated into individual storms (left) and for all snowflakes (right). Mean values and  $\pm 2\sigma$  values are represented by white horizontal lines and the boundary between shading and white space under the tails of the curves.

**Table 1.** Relationships between hydrometeor circumscribed and effective diameter.

Type	$D_{\max} = aD_{\text{eff}}^b$		$R^2$	N
	a	b		
All	1.16	1.04	0.94	43,649
Graupel	1.04	1.05	0.94	34
Densely Rimed	1.16	1.04	0.94	38,562
Aggregates	1.16	1.06	0.90	5,053




**Figure 7.** Mass-diameter relationship for all 86,285 snowflakes. Contours shown are for the 25th, 50th, 75th, 90th, and 95th percentiles.

generally exceeds those values obtained by Locatelli and Hobbs (1974) for graupel-like (2.1 to 2.4), densely rimed (2.1 to 2.3),  
 155 and aggregated (1.4 to 1.9) snow.

Snowflakes were selected manually through visual classification using MASC imagery for time periods at least three minutes  
 long where no other snow types were present, defined as aggregates (N=5,053), graupel (N=34), and densely rimed (N=38,562).  
 The densely rimed category includes all snowflakes not categorized as graupel or aggregates following Garrett and Yuter  
 (2014). It also includes densely rimed aggregates and partially melted aggregates. Figures 3 and C1-C4 show mostly planar  
 160 type crystals present, but aggregated needles are frequently seen as well. Graupel occurred frequently, although there was only  
 one period 16 minutes long that did not exhibit any other type of snow. Mass-diameter relationships  $M=aD_{eff}^b$  for graupel,  
 densely rimed snow, and aggregates are compared with those found by (Locatelli and Hobbs, 1974) in Figure 8 and Table  
 2. Values of  $a$  and  $b$  for graupel determined by the DEID ( $a=0.047$  and  $b=2.73$ ) lie within the ranges observed by Locatelli  
 and Hobbs (1974)  $0.042 < a < 0.140$  and  $2.6 < b < 3.0$ . The quantity of densely rimed and aggregate snowflakes collected by  
 165 the DEID was roughly five orders more numerous than those described by Locatelli and Hobbs (1974). For densely rimed  
 snowflakes, the prefactor  $a$  lies within the range observed by Locatelli and Hobbs (1974) and the exponent  $b$  is within 11%.  
 Aggregate snowflakes differ by 12% in the exponent and by 38% in the prefactor. In contrast to the findings of Erfani and  
 Mitchell (2017) which state that particle riming changes the prefactor  $a$  but not  $b$ , here both  $a$  and  $b$  decrease with increased  
 riming.

170 Figure 9 illustrates the sensitivity of mass-diameter relationship parameters to particle type and ambient air temperature.  
 Garrett and Yuter (2014) employed the MASC-derived complexity parameter,  $\chi = P(1 + \langle \sigma \rangle) / (2r)$  where  $P$  is snowflake




**Figure 8.** Mass-Diameter relationships from the DEID (solid lines) sorted by type according to MASC imagery and compared with fits from Locatelli and Hobbs (1974), dashed lines. Contours are for 50th, 75th, 90th, and 95th percentile bounds.

perimeter,  $r$  is snowflake radius, and  $\langle \sigma \rangle$  is intensity variability, to classify snowflakes into three categories: graupel ( $\chi < 1.35$ ), densely rimed ( $1.35 \leq \chi \leq 1.75$ ), and aggregates ( $\chi > 1.75$ ). Threshold values of 1.3 and 1.8 have also been previously used (Garrett et al., 2015). Here we adopt threshold values of 1.3 and 1.7. Graupel-like snow is included in the graupel category; the two datapoints with  $\chi < 1.3$  include 9,506 snowflakes, which are unlikely to be solely graupel due to one-minute averaging of  $\chi$  and the presence of other snow types observed during graupel events. This category has values of  $b$  consistent with the exponent values of between 2.1 and 2.4 found by Locatelli and Hobbs (1974). The densely rimed category however has exponent values between 2.5 and 2.7, higher than those seen by Locatelli and Hobbs (1974), although the difference may be influenced by presence of a large number of partially melted snowflakes that bring the exponent closer to 3. Overall, smaller values of  $b$  are obtained as  $\chi$  exceeds 1.7 and snowflakes transition into aggregates. Notably, the value of  $b$  is never lower than 2.

Partially melted snowflakes are excluded favoring more aggregate-type snowflakes by restricting analysis to particles that fell when the ambient air temperature was  $< -3^\circ\text{C}$ , as represented by blue lines in Figure 9. There is a clear sensitivity in the mass-diameter relationships to ambient air temperature. For all snowflakes that occurred when the ambient air temperature was  $< 0^\circ\text{C}$  ( $N=30,651$ ), the values of  $a$  and  $b$  are 0.017 and 2.33 respectively with  $R^2=0.85$ . For all snowflakes that fell when the ambient air temperature was  $< -3^\circ\text{C}$  ( $N=4,630$ ), the corresponding values are 0.015 and 2.12 with  $R^2=0.84$ .

### 3.4 Density-Diameter relationship

Density-diameter relationships for all snowflakes in this study are shown in Figure 10 and a comparison of density-diameter relationships from this work and prior studies is shown in Figure 11. Using the spherical volume approximation (Figure 10,



**Figure 9.** Exponent (gray) and prefactor (black) values in the Mass-Diameter relationship as binned by one-minute-average particle complexity and filtered by ambient temperature (blue line)


**Table 2.** Mass-Diameter relationship comparison by type

Type	Locatelli & Hobbs (1974)				DEID			
	$M=aD^b$		$R^2$	N	$M=aD^b$		$R^2$	N
	a	b			a	b		
Graupel	0.042 - 0.140	2.6 - 3.0	0.91 - 0.98	17 - 58	0.047	2.73	0.92	34
Graupel-like Snow	0.021 - 0.059	2.1 - 2.4	0.72 - 0.91	17 - 31				
Densely Rimed	0.015 - 0.033	2.1 - 2.3	0.78 - 0.92	9 - 13	0.018	2.57	0.91	38,562
Aggregates	0.037 - 0.073	1.4 - 1.9	0.78 - 0.91	19 - 27	0.014	2.17	0.85	5,053

Mass-Diameter relationship  $M=aD^b$  where mass is in milligrams and diameter is in millimeters

190 left), the measured values for density are rarely greater than  $100 \text{ kg m}^{-3}$ , suggesting a possible underestimate. Using a similar assumption, Muramoto et al. (1995) observed similarly low values of density (Figure 11). The density-diameter relationships from other studies shown in Figure 11 include both dry and wet snowflakes and ice particles with densities extending into the 200-300  $\text{kg m}^{-3}$  range for the lowest diameters observed.

195 More refined density calculations supplemented by MASC data are shown in Figure 10, right, and include high values near the density of bulk water, as would be expected for wet snowflakes that have partially melted before reaching the hotplate. The prefactors  $a$  are 35.02 and 74.25 for the spherical and MASC-adjusted density respectively and the exponents  $b$  are -0.48 and -0.57. The values of  $a$  and  $b$  were also obtained when filtering snowflakes by temperature to exclude partially melted snowflakes ( $T < 0^\circ\text{C}$ ) and to reflect primarily aggregate snowflakes ( $T < -3^\circ\text{C}$ ) and are shown in Table 3.



**Figure 10.** Density-diameter relationships for spherical (left) and MASC-adjusted (right) volume for all snowflakes. Contours shown are for the 25th, 50th, 75th, 90th, and 95th percentiles.


**Table 3.** Density-Diameter relationship comparison by ambient air temperature.

Density Method	Ambient T (°C)	$\rho=aD^b$		$R^2$	N
		a	b		
Spherical	All	35.05	-0.48	0.24	86,285
Spherical	< 0	34.87	-0.56	0.27	72,944
Spherical	< -3	28.12	-0.91	0.50	5,042
MASC	All	74.25	-0.57	0.17	43,649
MASC	< 0	67.27	-0.91	0.27	30,651
MASC	< -3	53.64	-1.28	0.35	4,630

## 4 Discussion

200 A notable difference between the mass-diameter relationships from our study versus those described previously by Locatelli and Hobbs (1974) lies in fitted exponents for both densely rimed and aggregate type snowflakes. In the earlier study these ranged from 2.1 to 2.3 for densely rimed and 1.4 to 1.9 for aggregates, while our results point to exponents of 2.57 and 2.17. An added implication of the larger exponents is that the masses of very small aggregates with diameters less than 1 mm are generally smaller than have previously been reported (Figure 8). While the MASC-adjusted density-diameter relationships align

205 closely with several previous studies (Figure 11) for particle sizes larger than approximately 5 mm diameter, much lower values of density tend to be observed at smaller particle sizes. Where the exponents described in previous studies (Holroyd, 1971; Fabry and Szyrmer, 1999; Heymsfield, 2003; Brandes et al., 2007; Tiira et al., 2016) are close to -1, those from our study are



**Figure 11.** Density-diameter relationships from previous studies and those obtained here using the spherical density method (magenta), and MASC-adjusted density method (blue).

approximately -0.5, suggesting a weaker dependence of density on particle size than is generally assumed, especially at ambient air temperatures near 0°C. However, by excluding higher temperatures near freezing to omit partially melted snowflakes, the exponents more closely approach -1. The average density of all hydrometeors using the spherical volume assumption is 38 kg m<sup>-3</sup> and 90 kg m<sup>-3</sup> using the MASC-adjusted volume. For comparison, the density of snowfall on the ground at a high-elevation location in the Wasatch Mountains in Utah is typically less than 100 kg m<sup>-3</sup> (Alcott and Steenburgh, 2010), a location where mode snowflake diameters lie between 1 mm and 2 mm (Garrett and Yuter, 2014).

## 5 Summary

We describe measurements of the mass and density of individual frozen hydrometeors obtained using a new instrument, the Differential Emissivity Imaging Disdrometer. Power-law mass-diameter relationships obtained by the DEID derived from 86,285 measured particles agree well with widely used relationships published by Locatelli and Hobbs (1974), which was based on a much more limited dataset. The exception is that snowflakes measured by the DEID have exponents higher by between 12% and 38%. To obtain hydrometeor density from the measured mass, estimates of volume are required. Here, a simple spherical approximation for particle volume based on the particle equivalent diameter seen by a thermal camera viewing the heated plate led to density estimates approximately a factor of two lower than those using a more refined calculation that incorporated concurrent MASC measurements of particle aspect ratio. For the subset of DEID measurements that included

coincident MASC imagery totalling 43,649 hydrometeors, the resulting density-diameter relationships suggest substantially lower densities of particles smaller than 5 mm than has been observed in most prior studies. It may be that existing bulk microphysical parameterizations in numerical weather models tend to underestimate the masses of large frozen hydrometeors while overestimating those of smaller hydrometeors. If true, any revision could have possible implications for forecasts of snow water deposition in mountain reservoirs. Future anticipated refinements to the DEID particle volume algorithm at a wider range of locations may help further refine estimates of hydrometeor density.

## Appendix A: Hotplate calibration

The constant  $K_d$ , which includes the specific heat capacity, equivalent latent heat, and the plate's thickness and thermal conductivity, was calibrated experimentally for the DEID aluminum plate (Singh et al., 2021). The plate was roughened with 600 grit (P1200) sandpaper to allow for droplet spreading and more rapid evaporation of water during calibration experiments. Water drops of known masses were evaporated on the plate to determine the calibrated coefficient. Combining the constants in Equation (1) yields,

$$(c_p \Delta T_{ev} + L_v)M = \frac{K}{H} \int_0^t A(t)(T_p - T_w(t))dt \quad (A1)$$

where  $M$  is a known mass of water. A constant is determined that includes the thermal conductivity and the thickness of the hotplate using droplets of known mass  $M = 2 \pm 0.2 \times 10^{-5}$  kg:

$$\frac{K}{H} = \frac{M(c_{p,w} \Delta T_{ev} + L_v)}{\int_0^t A(t)T_p - T_w(t)dt} \quad (A2)$$


where  $\int_0^t T_p - T_w(t)A(t)dt$  was determined from DEID measurements,  $c_{p,w} = 4.28 \times 10^3$  J K<sup>-1</sup> kg<sup>-1</sup> is the specific heat of water at constant pressure,  $\Delta T_{ev} = 100$ K, and  $L_v = 2.26 \times 10^6$  J kg<sup>-1</sup> is the latent heat of vaporization. Determined through 10 trials,  $K/H = 4.1 \pm 0.1603 \times 10^3$  kg s<sup>-3</sup> K<sup>-1</sup>.

Including the latent and specific heat required to evaporate liquid water and ice, respectively, the derived values of  $K_d$  for liquid and ice are then:

$$K_{d,l} = \frac{K/H}{(c_{p,w} \Delta T_{ev} + L_v)} = \frac{4.1 \times 10^3}{2.67 \times 10^6} = 1.54 \times 10^{-3} \left[ \frac{kg}{s K m^2} \right] \quad (A3)$$

$$K_{d,i} = \frac{K/H}{(c_{p,w} \Delta T_{ev} + L_v + c_{p,i} \Delta T_m + L_f)} = \frac{4.1 \times 10^3}{3.03 \times 10^6} = 1.35 \times 10^{-3} \left[ \frac{kg}{s K m^2} \right] \quad (A4)$$

where  $c_{p,i} = 2.10 \times 10^3$  J K<sup>-1</sup> kg<sup>-1</sup> is the specific heat of ice at constant pressure and the latent heat of fusion  $L_f = 3.34 \times 10^5$  J kg<sup>-1</sup>. The equation for mass becomes  $M = K_d \int_0^t A(t)(T_p - T_w(t))dt$ .



**Figure B1.** Volumetric rendering of melting rain hydrometeors in area on the DEID plate and in time such that each isosurface represents a rain drop frozen at a point in time.

Experiments comparing mass calculations using  $\int_0^t A(t)T_p - T_w(t)dt$  and  $\Delta T \int_0^t A dt$ , where  $\Delta T = T_p - T_w(t)$  is the mean value per particle showed that the latter is a sufficient approximation and the equation for mass may then be expressed as

$$250 \quad M = K_d \Delta T \int_0^t A(t) dt \quad (A5)$$


## Appendix B: Image processing

MATLAB Image Processing software was used to extract data from the thermal camera to determine the physical properties of individual melted hydrometeors. Experimentally relevant parameters include the hotplate temperature  $T_p$ , the sampling frequency of the thermal camera  $f_s$ , the physical width of each pixel in the camera imagery  $p$ , and the sampling area of  
 255 the hotplate  $A_{\text{hot}}$ . Each thermal camera image was converted to both grayscale and binary format. 3-dimensional volumetric regions, "voxels," in a product space of hydrometeor area on the DEID plate and time during the duration of the melting of each particle were evaluated to yield the particle mass through Eq. A5, as illustrated in Figures B1-B5.


For the sake of processing, all partial voxel objects that bordered the edge of the 3D sampling area were removed for individual particle calculations. A filtering threshold was used to remove small particles with fewer than three pixel data points  
 260 in any dimension.

The temperature difference between the hotplate and water ( $\Delta T = T_p - T_w(t)$ ) is based on the mean pixel intensity of the particle during its lifetime on the hotplate. The mean pixel intensity  $I_{\text{mean}}$  is converted to  $\Delta T$  through the linear transformation ( $(T_p - T_w(t)) \approx \Delta T = T_p(255 - I_{\text{mean}})/256$ ).





**Figure B2.** As for Fig. B1 but for partly melted hydrometeors




**Figure B3.** As for Fig. B1 but for aggregate snow


The integrated cross-sectional area of the particle during its evaporation time on the hotplate is given by  $V_t = \int A(t)dt$ . The effective diameter  $D_{\text{eff}}$  is calculated from the point in time associated with the maximum recorded cross-sectional area of the particle  $A_{\text{max}}$  according to  $D_{\text{eff}} = \sqrt{\frac{4}{\pi} A_{\text{max}}}$ . The evaporation time of each particle is calculated by generating a bounding box, or the smallest box that contains the 3D region containing  $V_t$  where the circumscribed diameter  $D_{\text{max}}$  is the maximum in the area dimension and the evaporation time  $t_{\text{evap}}$  is the maximum in the time dimension.

### Appendix C: MASC snowflake imagery

MASC Imagery of frozen hydrometeors was catalogued for each snow event described in this article with representative particles shown in Figures C1 to C4




**Figure B4.** As for Fig. B1 but for densely rimed aggregate snow




**Figure B5.** As for Fig. B1 but for a mixture of densely rimed and aggregate snow.

January 14, 2020




**Figure C1.** MASC Snowflake Imagery for January 14, 2020.

January 17, 2020




**Figure C2.** MASC Snowflake Imagery for January 17, 2020.

January 26, 2020



**Figure C3.** MASC Snowflake Imagery for January 26, 2020.

February 2-3, 2020



**Figure C4.** MASc Imagery for February 2 to February 3, 2020.

*Author contributions.* TJG and ERP conceived of the project. KNR and DKS led collection and analysis of the data. KNR and TJG contributed equally to writing the manuscript with contributions from ERP.

275 *Competing interests.* The DEID is protected through patent US20210172855A1 co-authored by KNR, DKS, ERP, and TJG. TJG is a co-owner of Particle Flux Analytics, Inc. which has a licence from the University of Utah to commercialize the DEID. TJG is a member of the editorial board of this journal (senior editor).

*Acknowledgements.* This work is supported by the U.S. Department of Energy (DOE) Atmospheric System Research program award number DE-SC0016282 and National Science Foundation (NSF) Physical and Dynamic Meteorology program award number 1841870. We are grateful to David Mitchell and an anonymous reviewer for comments, and to Spencer Donovan, and Allan Reaburn and colleagues at Particle  
280 Flux Analytics for their contributions to the field program and the development of the DEID.

## References

- Alcott, T. I. and Steenburgh, W. J.: Snow-to-liquid ratio variability and prediction at a high-elevation site in Utah's Wasatch Mountains, *Wea. Forecasting*, 25, 323–337, 2010.
- Barthazy, E. and Schefold, R.: Fall velocity of snowflakes of different riming degree and crystal types, *Atmos. Res.*, 82, 391–398, 2006.
- 285 Barthazy, E., Göke, S., Schefold, R., and Högl, D.: An Optical Array Instrument for Shape and Fall Velocity Measurements of Hydrometeors, *J. Atmos. Oceanic Technol.*, 21, 1400–+, [https://doi.org/10.1175/1520-0426\(2004\)021](https://doi.org/10.1175/1520-0426(2004)021), 2004.
- Battaglia, A., Rustemeier, E., Tokay, A., Blahak, U., and Simmer, C.: PARSIVEL Snow Observations: A Critical Assessment, *J. Atmos. Oceanic Technol.*, 27, 333–344, <https://doi.org/10.1175/2009JTECHA1332.1>, 2010.
- Böhm, H. P.: A General Equation for the Terminal Fall Speed of Solid Hydrometeors., *J. Atmos. Sci.*, 46, 2419–2427, 290 [https://doi.org/10.1175/1520-0469\(1989\)046<2419:AGEFTT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1989)046<2419:AGEFTT>2.0.CO;2), 1989.
- Brandes, E. A., Ikeda, K., Zhang, G., Schönhuber, M., and Rasmussen, R. M.: A Statistical and Physical Description of Hydrometeor Distributions in Colorado Snowstorms Using a Video Disdrometer, *Journal of Applied Meteorology and Climatology*, 46, 634–650, <https://doi.org/10.1175/JAM2489.1>, 2007.
- Brandes, E. A., Ikeda, K., Thompson, G., and Schönhuber, M.: Aggregate Terminal Velocity/Temperature Relations, *J. Appl. Meteorol. Clim.*, 47, 2729–2736, <https://doi.org/10.1175/2008JAMC1869.1>, 2008.
- 295 Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *Journal of Glaciology*, 38, 13–22, <https://doi.org/10.3189/S0022143000009552>, 1992.
- Colle, B. A., Garvert, M. F., Wolfe, J. B., Mass, C. F., and Woods, C. P.: The 13 14 December 2001 IMPROVE-2 Event. Part III: Simulated Microphysical Budgets and Sensitivity Studies., *J. Atmos. Sci.*, 62, 3535–3558, <https://doi.org/10.1175/JAS3552.1>, 2005.
- 300 Conger, S. M. and McClung, D. M.: Comparison of density cutters for snow profile observations, *Journal of Glaciology*, 55, 163–169, 2009.
- Dent, J., Burrell, K., Schmidt, D., Louge, M., Adams, E., and Jazbutis, T.: Density, velocity and friction measurements in a dry-snow avalanche, *Annals of Glaciology*, 26, 247–252, 1998.
- Erfani, E. and Mitchell, D. L.: Growth of ice particle mass and projected area during riming, *Atmospheric Chemistry and Physics*, 17, 1241–1257, <https://doi.org/10.5194/acp-17-1241-2017>, <https://acp.copernicus.org/articles/17/1241/2017/>, 2017.
- 305 Fabry, F. and Szyrmer, W.: Modeling of the Melting Layer. Part II: Electromagnetic, *Journal of the Atmospheric Sciences*, 56, 3593–3600, 1999.
- Fovell, R. G. and Su, H.: Impact of cloud microphysics on hurricane track forecasts, *Geophys. Res. Lett.*, 34, L24810, <https://doi.org/10.1029/2007GL031723>, 2007.
- Garrett, T. J. and Yuter, S. E.: Observed influence of riming, temperature, and turbulence on the fallspeed of solid precipitation, *Geophysical Research Letters*, 41, 6515–6522, <https://doi.org/10.1002/2014GL061016>, 2014.
- 310 Garrett, T. J., Fallgatter, C., Shkurko, K., and Howlett, D.: Fall speed measurement and high-resolution multi-angle photography of hydrometeors in free fall, *Atmospheric Measurement Techniques*, 5, 2625–2633, <https://doi.org/10.5194/amt-5-2625-2012>, <http://www.atmos-meas-tech.net/5/2625/2012/>, 2012.
- Garrett, T. J., Yuter, S. E., Fallgatter, C., Shkurko, K., Rhodes, S. R., and Endries, J. L.: Orientations and aspect ratios of falling snow, 315 *Geophysical Research Letters*, 42, 4617–4622, <https://doi.org/10.1002/2015GL064040>, 2015.



- Garvert, M. F., Woods, C. P., Colle, B. A., Mass, C. F., Hobbs, P. V., Stoelinga, M. T., and Wolfe, J. B.: The 13–14 December 2001 IMPROVE-2 Event. Part II: Comparisons of MM5 Model Simulations of Clouds and Precipitation with Observations., *Journal of Atmospheric Sciences*, 62, 3520–3534, <https://doi.org/10.1175/JAS3551.1>, 2005.
- 320 Heymsfield, A. J.: Properties of Tropical and Midlatitude Ice Cloud Particle Ensembles. Part II: Applications for Mesoscale and Climate Models, *Journal of the Atmospheric Sciences*, 60, 2592–2611, 2003.
- Heymsfield, A. J. and Westbrook, C. D.: Advances in the Estimation of Ice Particle Fall Speeds Using Laboratory and Field Measurements, *J. Atmos. Sci.*, 67, 2469–2482, <https://doi.org/10.1175/2010JAS3379.1>, 2010.
- Holroyd, E. W.: The meso- and microscale structure of Great Lakes snowstorm bands: A synthesis of ground measurements, radar data, and satellite observations, Ph.D. thesis, State University of New York at Albany, Department of Atmospheric Science, 1971.
- 325 Hong, S., Dudhia, J., and Chen, S.: A Revised Approach to Ice Microphysical Processes for the Bulk Parameterization of Clouds and Precipitation, *Mon. Wea. Rev.*, 132, 103–120, [https://doi.org/10.1175/1520-0493\(2004\)132<0103:ARATIM>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<0103:ARATIM>2.0.CO;2), 2004.
- Iguchi, T., Matsui, T., Shi, J. J., Tao, W.-K., Khain, A. P., Hou, A., Cifelli, R., Heymsfield, A., and Tokay, A.: Numerical analysis using WRF-SBM for the cloud microphysical structures in the C3VP field campaign: Impacts of supercooled droplets and resultant riming on snow microphysics, *J. Geophys. Res.*, 117, D23 206, <https://doi.org/10.1029/2012JD018101>, 2012.
- 330 Khvorostyanov, V. I. and Curry, J. A.: Terminal Velocities of Droplets and Crystals: Power Laws with Continuous Parameters over the Size Spectrum, *J. Atmos. Sci.*, 59, 1872–1884, [https://doi.org/10.1175/1520-0469\(2002\)059<1872:TVODAC>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1872:TVODAC>2.0.CO;2), 2002.
- Kruger, A. and Krajewski, W. F.: Two-dimensional video disdrometer: A description, *J. Atmos. Oceanic. Technol.*, 19, 602–617, 2002.
- Kubicek, A. and Wang, P. K.: A numerical study of the flow fields around a typical conical graupel falling at various inclination angles, *Atmospheric Research*, 118, 15 – 26, <https://doi.org/10.1016/j.atmosres.2012.06.001>, 2012.
- 335 Lin, Y., Donner, L. J., and Colle, B. A.: Parameterization of Riming Intensity and Its Impact on Ice Fall Speed Using ARM Data, *Mon. Wea. Rev.*, 139, 1036–1047, <https://doi.org/10.1175/2010MWR3299.1>, 2010.
- Liu, C., Ikeda, K., Thompson, G., Rasmussen, R., and Dudhia, J.: High-Resolution Simulations of Wintertime Precipitation in the Colorado Headwaters Region: Sensitivity to Physics Parameterizations., *Monthly Weather Review*, 139, 3533 – 3553, 2011.
- Locatelli, J. D. and Hobbs, P. V.: Fall speeds and masses of solid precipitation particles, *J. Geophys. Res.*, 79, 2185–2197, 340 <https://doi.org/10.1029/JC079i015p02185>, 1974.
- Locatelli, J. D. and Hobbs, P. V.: Fall speeds and masses of solid precipitation particles, *Journal of Geophysical Research*, 79, 2185–2197, <https://doi.org/10.1029/JC079i015p02185>, 1974.
- Magono, C. and Nakamura, T.: Aerodynamic Studies of Falling Snowflakes, *Journal of the Meteorological Society of Japan. Ser. II*, 43, 139–147, [https://doi.org/10.2151/jmsj1965.43.3\\_139](https://doi.org/10.2151/jmsj1965.43.3_139), 1965.
- 345 Milbrandt, J. A., Yau, M. K., Mailhot, J., Bélair, S., and McTaggart-Cowan, R.: Simulation of an Orographic Precipitation Event during IMPROVE-2. Part II: Sensitivity to the Number of Moments in the Bulk Microphysics Scheme, *Mon. Wea. Rev.*, 138, 625–642, <https://doi.org/10.1175/2009MWR3121.1>, 2010.
- Muramoto, K.-i., Matsuura, K., and Shiina, T.: Measuring the Density of Snow Particles and Snowfall Rate, *Electronics & Communications in Japan, Part 3: Fundamental Electronic Science*, 78, 71–79, <https://doi.org/10.1002/ecjc.4430781107>, 1995.
- 350 Newman, A. J., Kucera, P. A., and Bliven, L. F.: Presenting the Snowflake Video Imager (SVI), *J. Atmos. Oceanic. Technol.*, 26, 167–179, <https://doi.org/10.1175/2008JTECHA1148.1>, 2009.
- Reisner, J., Rasmussen, R. M., and Bruintjes, R. T.: Explicit forecasting of supercooled liquid water in winter storms using the MM5 mesoscale model, *Q. J. Roy. Meteorol. Soc.*, 124, 1071–1107, <https://doi.org/10.1256/smsqj.54803>, 1998.

- Rutledge, S. A. and Hobbs, P. V.: The Mesoscale and Microscale Structure and Organization of Clouds and Precipitation in Midlatitude  
355 Cyclones. XII: A Diagnostic Modeling Study of Precipitation Development in Narrow Cold-Frontal Rainbands., *J. Atmos. Sci.*, 41, 2949–  
2972, [https://doi.org/10.1175/1520-0469\(1984\)041<2949:TMAMSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1984)041<2949:TMAMSA>2.0.CO;2), 1984.
- Shkurko, K., Gaustad, K., and Garrett, T. J.: Multi-Angles Snowflake Camera Value-Added Product, Tech. Rep. DOE/SC-ARM-TR-187,  
Department of Energy Atmospheric Radiation Measurement Program, 2016.
- Singh, D. K., Donovan, S., Pardyjak, E. R., and Garrett, T. J.: A differential emissivity imaging technique for measuring hydrometeor mass  
360 and type, *Atmos. Meas. Tech.*, 2021.
- Tao, W.-K., Simpson, J., Baker, D., Braun, S., Chou, M.-D., Ferrier, B., Johnson, D., Khain, A., Lang, S., Lynn, B., et al.: Microphysics,  
radiation and surface processes in the Goddard Cumulus Ensemble (GCE) model, *Meteor. Atmos. Phys.*, 82, 97–137, 2003.
- Thériault, J. M., Stewart, R. E., and Henson, W.: Impacts of terminal velocity on the trajectory of winter precipitation types, *Atmos. Res.*,  
116, 116 – 129, <https://doi.org/10.1016/j.atmosres.2012.03.008>, <ce:title>Remote Sensing of Clouds and Aerosols: Techniques and Ap-  
365 plications - Atmospheric Research</ce:title>, 2012.
- Tiira, J., Moisseev, D. N., von Lerber, A., Ori, D., Tokay, A., Bliven, L. F., and Petersen, W.: Ensemble mean density and its connection to  
other microphysical properties of falling snow as observed in Southern Finland, *Atmospheric Measurement Techniques (Online)*, 9, 2016.
- Yuter, S. E., Kingsmill, D. E., Nance, L. B., and Löffler-Mang, M.: Observations of Precipitation Size and Fall Speed Characteristics within  
Coexisting Rain and Wet Snow, *J. Appl. Meteorol. Clim.*, 45, 1450–1464, <https://doi.org/10.1175/JAM2406.1>, 2006.