

1 ***Supporting information***

2 Measurement report: Emissions of intermediate-volatility organic compounds from
3 vehicles under real-world driving conditions in an urban tunnel

4 Hua Fang^{1,2,4}, Xiaoqing Huang^{1,2,4}, Yanli Zhang^{1,2,3*}, Chenglei Pei^{1,4,5}, Zuzhao Huang⁶, Yujun Wang⁵,
5 Yanning Chen⁵, Jianhong Yan⁷, Jianqiang Zeng^{1,2,4}, Shaoxuan Xiao^{1,2,4}, Shilu Luo^{1,2,4}, Sheng Li^{1,2,4}, Jun
6 Wang^{1,2,4}, Ming Zhu^{1,2,4}, Xuwei Fu^{1,2,4}, Zhenfeng Wu^{1,2,4}, Runqi Zhang^{1,2,4}, Wei Song^{1,2}, Guohua
7 Zhang^{1,2}, Weiwei Hu^{1,2}, Mingjin Tang^{1,2}, Xiang Ding^{1,2}, Xinhui Bi^{1,2}, Xinming Wang^{1,2,3,4*}

8

9 ¹State Key Laboratory of Organic Geochemistry and Guangdong Key Laboratory of
10 Environmental Protection and Resources Utilization, Guangzhou Institute of Geochemistry,
11 Chinese Academy of Sciences, Guangzhou 510640, China

12 ²CAS Center for Excellence in Deep Earth Science, Guangzhou, 510640, China

13 ³Center for Excellence in Urban Atmospheric Environment, Institute of Urban Environment,
14 Chinese Academy of Sciences, Xiamen 361021, China

15 ⁴University of Chinese Academy of Sciences, Beijing 100049, China

16 ⁵Guangzhou Ecological and Environmental Monitoring Center of Guangdong Province,
17 Guangzhou 510060, China

18 ⁶Guangzhou Environmental Technology Center, Guangzhou 510180, China

19 ⁷Guangzhou Tunnel Development Company, Guangzhou 510133, China

20

21 *Correspondence: Dr. Xinming Wang (e-mail: wangxm@gig.ac.cn) and Dr. Yanli Zhang (e-
22 mail: Zhang_yl86@gig.ac.cn)

23 Text S1

24 Quantification of IVOCs

25 Speciated IVOCs were identified based on their retention times and mass spectra, and were
26 quantified by 18 authentic standards including C₁₂-C₂₂ n-alkanes and 8 polycyclic aromatic
27 hydrocarbons (PAHs). The pristane and phytane were quantified by the calibration curves of the n-
28 alkanes (n-C₁₇ and n-C₁₈) near them.

29 Total IVOCs mass were quantified by using the method developed by Zhao et al. (2014). Firstly, the
30 total ion chromatogram (TIC) was divided into 11 chromatogram bins based on retention times of
31 C₁₂-C₂₂ n-alkanes. The start time and end time of chromatogram bin were determined by successive
32 n-alkanes. The example was illustrated as following:

$$33 \quad t_{n, \text{ Bin-start}} = \frac{t_{n-1} + t_n}{2}$$
$$34 \quad t_{n, \text{ Bin-end}} = \frac{t_n + t_{n+1}}{2}$$

35 Where n refers to the carbon number of n-alkane which centered in the corresponding chromatogram
36 bin. The $t_{n, \text{ Bin-start}}$ and $t_{n, \text{ Bin-end}}$ respectively represent the start time and end time of chromatogram
37 bin. The t_{n-1} , t_n and t_{n+1} are the retention times of C_{n-1}, C_n and C_{n+1} n-alkanes, respectively.

38 Secondly, the IVOCs mass in each defined chromatogram bin (M_{IVOCs, B_n}) was calculated by
39 following equation:

$$40 \quad M_{\text{IVOCs}, B_n} = \frac{TA_{\text{TIC}, B_n}}{RF_{n\text{-alkanes}, C_n}} = \frac{TA_{m/z 57, B_n}}{RF_{n\text{-alkanes}, C_n}} \times \frac{1}{f_{m/z 57, \text{TIC}, B_n}}$$

41 Where TA_{TIC, B_n} is the abundance of TIC in the B_n chromatogram bin; $RF_{n\text{-alkane}, C_n}$ is the response
42 factor of the C_n n-alkane; $TA_{m/z 57, B_n}$ refers to the abundance of m/z 57 in the B_n chromatogram bin;
43 $f_{m/z 57, \text{TIC}, B_n}$ is the fraction of the m/z 57 in TIC of the B_n chromatogram bin. The total IVOCs mass
44 was the sum of the IVOCs mass calculated in each defined chromatogram bin.

45 Thirdly, the residual IVOCs (subtracting the speciated IVOCs), which was named as unresolved
46 complex mixtures IVOCs (UCM IVOCs) in previous studies (Zhao et al., 2014, 2015, 2016), were
47 further divided into unspciated branched alkanes (b-alkanes) and unspciated cyclic compounds.
48 Here, we assumed that the signal of m/z 57 all came from n-alkanes and b-alkanes. The unspciated
49 b-alkanes mass ($M_{\text{b-alkane}, B_n}$) was defined as the difference between calculated mass of m/z 57 and
50 n-alkane that falling in the same chromatogram bin (Zhao et al., 2015, 2016). Obviously, this

51 assumption would somewhat cause an overestimation of b-alkanes in that other compounds like
52 cyclic alkanes also could contribute to the abundance of m/z 57 (Zhao et al., 2014).

$$\begin{aligned} 53 \quad M_{b\text{-alkane}, B_n} &= \frac{BA_{b\text{-alkane}, B_n}}{RF_{n\text{-alkane}, C_n}} = \frac{BA_{m/z\ 57, B_n}}{RF_{n\text{-alkane}, C_n}} \times \frac{1}{f_{m/z\ 57, b\text{-alkane}}} \\ 54 \quad &= \frac{(TA_{m/z\ 57, B_n} - NA_{m/z\ 57, C_n})}{RF_{n\text{-alkane}, C_n}} \times \frac{1}{f_{m/z\ 57, b\text{-alkane}}} \end{aligned}$$

55 Where $BA_{b\text{-alkane}, B_n}$ is the abundance of b-alkanes in B_n chromatogram bin. The $BA_{m/z\ 57, B_n}$
56 represents the abundance of m/z 57 came from b-alkanes in the B_n chromatogram bin while $NA_{m/z\ 57, C_n}$
57 $NA_{m/z\ 57, C_n}$ is the abundance of m/z 57 produced by the C_n n-alkane. $TA_{m/z\ 57, B_n}$ is the total abundance of
58 m/z 57 in B_n chromatogram bin. The $f_{m/z\ 57, b\text{-alkane}}$ is the average fraction of m/z 57 in the TIC of b-
59 alkanes.

60 Lastly, the mass of unspiciated cyclic compounds were calculated as a consequence of subtracting
61 both the masses of speciated IVOCs and unspiciated b-alkanes from the total determined IVOCs
62 mass.


63 **Text S2**

64 **Calculation of fuel-based emission factor**

65 The fuel-based emission factor of IVOCs was calculated as following:


$$66 \quad \text{fuel - based EF (mg kg}^{-1}\text{)} = \frac{\text{mileage - based EF (mg km}^{-1}\text{)}}{\text{fuel density (kg L}^{-1}\text{)} \times \text{fuel efficiency (L km}^{-1}\text{)}}$$

67 Here, the 7.87 L/100 km and 7.5 L/100 km were used as the average gasoline and diesel
68 efficiency, respectively (<http://icet.org.cn/admin/upload/2014101812382577.pdf>). The fuel
69 density was 0.74 g mL⁻¹ for gasoline and 0.85 g mL⁻¹ for diesel (Zhang et al., 2016).




70

71 Figure S1. The schematic diagram of the sampling stations inside the Zhujiang tunnel.




72

73 Figure S2. The vehicle fleet compositions during the campaign.



74

75 Figure S3. Linear regression analysis of diesel fraction and fleet-average EF_{IVOCs} during the
76 campaign. The green lines represent 95% confidence intervals.




77

78 Figure S4. The comparison of distributions of total IVOCs in different volatility bins

79 determined in this study and in previous studies. The boxes represent the 75th and 25th

80 percentiles, the centerlines are the medians and squares are the averages. The whiskers represent


81 10th and 90th percentiles.



82

83 Figure S5. The Pearson correlations among IVOCs in each volatility bins (Σ represents sum of

84 IVOCs from B12 to B22).



85

86 Figure S6. The relationships of IVOCs with other primary species that concurrently measured

87 in the tunnel.

88 Table S1. The emission factors of IVOCs species (mg km⁻¹).

Species	Mean \pm 95% C.I.	Range
Dodecane	0.38 \pm 0.03	0.12–0.81
Tridecane	0.23 \pm 0.02	0.11–0.61
Tetradecane	0.22 \pm 0.02	0.10–0.53
Pentadecane	0.19 \pm 0.02	0.08–0.50
Hexadecane	0.16 \pm 0.01	0.06–0.38
Heptadecane	0.15 \pm 0.01	0.06–0.29
Pristane	0.10 \pm 0.01	0.03–0.24
Octadecane	0.13 \pm 0.01	0.05–0.25
Phytane	0.10 \pm 0.01	0.04–0.22
Nonadecane	0.10 \pm 0.01	0.03–0.24
Eicosane	0.10 \pm 0.01	0.01–0.21
Heneicosane	0.08 \pm 0.01	0.01–0.19
Docosane	0.04 \pm 0.00	0.00–0.11
Naphthalene	0.35 \pm 0.03	0.14–0.82
2-Methylnaphthalene	0.11 \pm 0.01	0.05–0.29
1-Methylnaphthalene	0.05 \pm 0.00	0.02–0.13
Acenaphthylene	0.06 \pm 0.01	0.00–0.17
Acenaphthene	0.01 \pm 0.00	0.00–0.05
Fluorene	0.00 \pm 0.00	0.00–0.01
Phenanthrene	0.01 \pm 0.00	0.00–0.05
Anthracene	0.02 \pm 0.00	0.00–0.11
Unspeciated b-alkanes B12	0.89 \pm 0.07	0.27–1.87
Unspeciated b-alkanes B13	0.58 \pm 0.04	0.19–1.03
Unspeciated b-alkanes B14	0.52 \pm 0.04	0.21–1.29
Unspeciated b-alkanes B15	0.43 \pm 0.03	0.17–1.11
Unspeciated b-alkanes B16	0.37 \pm 0.04	0.13–1.30
Unspeciated b-alkanes B17	0.34 \pm 0.03	0.15–0.81
Unspeciated b-alkanes B18	0.35 \pm 0.03	0.09–0.86
Unspeciated b-alkanes B19	0.25 \pm 0.03	0.04–0.64
Unspeciated b-alkanes B20	0.21 \pm 0.02	0.04–0.46
Unspeciated b-alkanes B21	0.19 \pm 0.02	0.04–0.61
Unspeciated b-alkanes B22	0.08 \pm 0.01	0.02–0.27
Unspeciated cyclic compounds B12	2.36 \pm 0.23	0.96–7.13
Unspeciated cyclic compounds B13	1.54 \pm 0.13	0.21–3.74
Unspeciated cyclic compounds B14	1.33 \pm 0.10	0.42–2.69
Unspeciated cyclic compounds B15	0.89 \pm 0.07	0.19–2.01
Unspeciated cyclic compounds B16	0.82 \pm 0.07	0.21–2.27
Unspeciated cyclic compounds B17	0.71 \pm 0.07	0.12–1.61
Unspeciated cyclic compounds B18	0.59 \pm 0.06	0.00–1.67
Unspeciated cyclic compounds B19	0.57 \pm 0.05	0.16–1.42
Unspeciated cyclic compounds B20	0.49 \pm 0.06	0.08–1.71
Unspeciated cyclic compounds B21	0.42 \pm 0.04	0.15–1.18
Unspeciated cyclic compounds B22	0.26 \pm 0.03	0.07–0.68
Σ Speciated IVOCs	2.59 \pm 0.14	1.43–4.37
Σ UCM IVOCs	14.19 \pm 0.79	7.41–25.60
Σ IVOCs	16.77 \pm 0.89	9.04–29.32

89 Table S2. The correlations and mass ratios of IVOCs with normal alkanes in the same volatility
 90 bin.

	B12/n-C12	B13/n-C13	B14/n-C14	B15/n-C15	B16/n-C16	B17/n-C17
Ratios	11.20±0.61	11.59±0.76	10.03±0.60	9.00±0.64	9.02±0.63	9.37±0.64
Correlations (R)	0.76**	0.51**	0.60**	0.56**	0.63**	0.53*
	B18/n-C18	B19/n-C19	B20/n-C20	B21/n-C21	B22/n-C22	
Ratios	9.90±0.73	10.15±0.83	9.71±1.62	10.73±1.97	15.74±5.38	
Correlations (R)	0.65**	0.54**	0.34	0.40*	0.42**	

91 The asterisks shown in table represent significance. **p < 0.01, *p < 0.05, no asterisks mean p > 0.05.

92 Table S3. SOA yields that used in this study.

	Species	SOA yields	Species	SOA yields
Speciated IVOCs	Dodecane	0.12	Heneicosane	0.55
	Tridecane	0.3	Docosane	0.55
	Tetradecane	0.4	Naphthalene	0.22
	Pentadecane	0.46	2-Methylnaphthalene	0.26
	Hexadecane	0.51	1-Methylnaphthalene	0.31
	Heptadecane	0.55	Acenaphthylene	0.31
	Pristane	0.46	Acenaphthene	0.31
	Octadecane	0.55	Fluorene	0.31
	Phytane	0.51	Phenanthrene	0.31
	Nonadecane	0.55	Anthracene	0.31
	Eicosane	0.55		
Unspeciate b-alkanes IVOCs	Bin12	0.05	Bin18	0.51
	Bin13	0.09	Bin19	0.55
	Bin14	0.12	Bin20	0.55
	Bin15	0.3	Bin21	0.55
	Bin16	0.4	Bin22	0.55
	Bin17	0.46		
Unspeciated cyclic compounds IVOCs	Bin12	0.12	Bin18	0.55
	Bin13	0.3	Bin19	0.55
	Bin14	0.4	Bin20	0.55
	Bin15	0.46	Bin21	0.55
	Bin16	0.51	Bin22	0.55
	Bin17	0.55		
VOCs	Benzene	0.21	n-Heptane	0.009 ^a
	Toluene	0.11	n-Octane	0.041 ^a
	Ethylbenzene	0.11	n-Nonane	0.08 ^a
	m/p-Xylene	0.06	n-Decane	0.146 ^a
	Styrene	0.11	2-Methylhexane	0.009 ^b
	o-Xylene	0.06	3-Methylhexane	0.009 ^b
	Isopropylbenzene	0.11	2,3-Dimethylpentane	0.009 ^b
	n-Propylbenzene	0.11	2,4-Dimethylpentane	0.009 ^b
	m-Ethyltoluene	0.11	2,2,4-Trimethylpentane	0.041 ^c
	p-Ethyltoluene	0.11	2,3,4-Trimethylpentane	0.041 ^c
	1,3,5-Trimethylbenzene	0.06	2-Methylheptane	0.041 ^c
	o-Ethyltoluene	0.06	3-Methylheptane	0.041 ^c
	1,2,4-Trimethylbenzene	0.06	Cyclohexane	0.04
1,2,3-Trimethylbenzene	0.06	Methylcyclohexane	0.04 ^d	

93 SOA yields of IVOCs and single-ring aromatics were from Zhao et al. (2015); ^a SOA yields are from Lim and Ziemann. (2009); ^b SOA yields are assumed the
94 same as n-Heptane; ^c SOA yields are assumed the same as n-October; ^d SOA yields are assumed the same as cyclohexane.

95 **References**

- 96 Lim, Y. B., and Ziemann, P. J.: Effects of molecular structure on aerosol yields from OH radical-
97 initiated reactions of linear, branched, and cyclic alkanes in the presence of NO_x. *Environ.*
98 *Sci. Technol.*, 43, 2328–2334, <https://doi.org/10.1021/es803389s>, 2009.
- 99 Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop,
100 A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking organic aerosols: semivolatile
101 emissions and photochemical aging. *Science* 315, 1259–1262,
102 <http://doi.org/10.1126/science.1133061>, 2007.
- 103 Zhang, Y. L., Wang, X. M., Wen, S., Herrmann, H., Yang, W. Q., Huang, X. Y., Zhang, Z.,
104 Huang, Z. H., He, Q. F., and George, C.: On-road vehicle emissions of glyoxal and
105 methylglyoxal from tunnel tests in urban Guangzhou, China. *Atmos. Environ.*, 127, 55–60,
106 <http://doi.org/10.1016/j.atmosenv.2015.12.017>, 2016.
- 107 Zhao, Y. L., Hennigan, C. J., May, A. A., Tkacik, D. S., de Gouw, J. A., Gilman, J. B., Kuster,
108 W. C., Borbon, A., and Robinson, A. L.: Intermediate-volatility organic compounds: a
109 large source of secondary organic aerosol. *Environ. Sci. Technol.*, 48, 13743–13750,
110 <https://doi.org/10.1021/es5035188>, 2014.
- 111 Zhao, Y. L., Nguyen, N. T., Presto, A. A., Hennigan, C. J., May, A. A., and Robinson, A. L.:
112 Intermediate-volatility organic compound emissions from on-road diesel vehicles:
113 chemical composition, emission factors, and estimated secondary organic aerosol
114 production. *Environ. Sci. Technol.*, 49, 11516–11526,
115 <https://doi.org/10.1021/acs.est.5b02841>, 2015.
- 116 Zhao, Y. L., Nguyen, N. T., Presto, A. A., Hennigan, C. J., May, A. A., and Robinson, A. L.:
117 Intermediate-volatility organic compound emissions from on-road gasoline vehicles and
118 small off-road gasoline engines. *Environ. Sci. Technol.*, 50, 4554–4563,
119 <https://doi.org/10.1021/acs.est.5b06247>, 2016.