



Analysis of aerosol-cloud interactions and their implications for precipitation formation using aircraft observations over the United Arab Emirates

5 Youssef Wehbe¹, Sarah A. Tessendorf², Courtney Weeks², Roelof Brientjes², Lulin Xue², Roy Rasmussen², Paul Lawson³, Sarah Woods³, and Marouane Temimi⁴

¹National Center of Meteorology, Abu Dhabi 4815, UAE

²Research Applications Laboratory, National Center for Atmospheric Research, Boulder, CO 80307, USA

³Stratton Park Engineering Company, Boulder, CO 80301, USA

⁴Department of Civil, Environmental, and Ocean Engineering, Stevens Institute of Technology, Hoboken, NJ 07030, USA

10

Correspondence to: Youssef Wehbe (ywehbe@ncms.ae)

Abstract. Aerosol and cloud microphysical measurements were collected by a research aircraft during August 2019 over the United Arab Emirates (UAE). The majority of science flights targeted summertime convection along the eastern Hajar mountains bordering Oman, while one flight sampled non-orographic clouds over the western UAE near the Saudi Arabian border. In this work, we study the evolution of growing cloud turrets from cloud base (9 °C) up to the capping inversion level (-12 °C) using coincident cloud particle imagery and particle size distributions from cloud cores under different forcing. Results demonstrate the active role of background dust and pollution as cloud condensation nuclei (CCN) with the onset of their deliquescence in the sub-cloud region. Sub-cloud aerosol sizes are shown to extend from submicron to 100 μm sizes, with higher concentrations of ultra-giant CCN (d>10 μm) from local sources closer to the Saudi border, compared to the eastern orographic region where smaller size CCN are observed. Despite the presence of ultra-giant CCN from dust and pollution in both regions, an active collision-coalescence (C-C) process is not observed within the limited depths of warm cloud (<1000 m). The state-of-the-art observations presented in this paper can be used to initialize modelling case studies to study the influence of aerosols on cloud and precipitation processes in the region and to better understand the impacts of hygroscopic cloud-seeding on these clouds.

25 1 Introduction

Aerosol particles are key components of the atmosphere and have multi-scale impacts on Earth's climate and hydrological cycle, primarily through radiative transfer and precipitation formation. Aerosol effects are grouped into direct radiative effects (Ming et al., 2005) and indirect aerosol-cloud interactions (Lohmann et al., 2010). The magnitude of uncertainty from indirect effects remains more difficult to quantify compared to that from direct effects (Solomon et al., 2007). This is primarily due to the complexity of the microphysical processes involved and their interdependencies (Morrison et al., 2005). Furthermore, the

30



contributions from thermodynamic and dynamic factors introduce additional complexities to the precipitation generation process. Hence, assessing the impact of aerosols on cloud microphysics and precipitation generation has been a longstanding research area with many processes yet to be fully understood (Morrison et al., 2020; Flossmann et al., 2019; Solomon et al., 2007; Squires, 1958; Rosenfeld, 1999). Our current knowledge is based primarily on observing the influence of background
35 aerosols on the number and size distribution of hydrometeors (Rosenfeld et al., 2008; Freud et al., 2008; Tao et al., 2007; Rosenfeld, 2000; Feingold et al., 1999). Isolating these impacts has been shown to be particularly challenging in polluted and dusty environments where the physiochemical properties of aerosols are continuously altered between large dust particles, fine particle pollution as well as complex aggregates depending on both regional and local forcing (Abuelgasim and Farahat, 2020; Filioglou et al., 2020; Semeniuk et al., 2014).

40 The role of dust aerosols as ice nucleating particles (INPs) is well established in the literature (Kanji et al., 2019; Atkinson et al., 2013; Hoose and Möhler, 2012; Prenni et al., 2009; Möhler et al., 2006). Dust INPs have been shown to prolong cloud lifetimes and enhance precipitation in relatively unpolluted continental-tropical and maritime regimes (Liu et al., 2020; Koren et al., 2005). Alternatively, when ingested into strong convective updrafts, large populations of mineral dust INPs tend to limit the growth of hydrometeors in the upper levels due to their competing effect for available water vapor (Kant et al., 2019; Min
45 et al., 2009; DeMott et al., 2003; Sassen et al., 2003). On the other hand, the role of dust as cloud condensation nuclei (CCN) in polluted desert environments with diverse aerosol physio-chemistry is less understood. The recent study by Chen et al. (2019) showed non-linear responses of precipitation to dust when it serves as both CCN and INP.

Dust introduces large particles with diameters exceeding 2 μm which could serve as giant CCN (GCCN) (Jensen and Nugent, 2017; Yin et al., 2000; Jensen and Lee, 2008). This alludes to more effective droplet collection during the collision-coalescence
50 (C-C) process. However, Rosenfeld et al. (2001) attributes the reduction in cloud droplet effective radii ($r_c < 14 \mu\text{m}$) over the Saharan desert to the presence of large concentrations of submicron CCN originating from desert mineral dust. This is shown to inhibit C-C and warm rain formation, exacerbating a reduction in precipitation over the Sahara region. A similar conclusion is derived from the modelling studies of Flossmann and Wobrock (2010). These findings contradict with previous work suggesting enhanced C-C by GCCN regardless of the fine-mode concentrations (Johnson, 1982; Takeda and Kuba, 1982).
55 Aside from size and concentration, aerosol chemistry plays an important role in determining the nucleation properties of dust particles. Laboratory deliquescence experiments show that pure insoluble mineral dust particles remain hydrophobic at 100% relative humidity (RH), while the aggregation of dust particles with soluble compounds (e.g. NaCl) results in their deliquescence at 74% RH (Wise et al., 2007). Similarly, compounds from anthropogenic pollution (sulfates) interact with mineral dust to produce internally-mixed aggregates which deliquesce at 80% RH (Semeniuk et al., 2015).

60 Located along the central portion of the dust belt, the Arabian Peninsula is largely considered a hotspot for atmospheric dust emissions (Karagulian et al., 2019; Beegum et al., 2018; Hofer et al., 2017). The UAE is situated on the southeastern coast of the Arabian Peninsula and records an annual rainfall average less than 100 mm, which is representative of the region's scarce



rainfall amounts (Wehbe et al., 2018). Aiming to augment natural rainfall over the country, the UAE has implemented an operational cloud-seeding program for the past two decades – the longest ongoing program in the region (Almazroui and Farrah, 2017). In addition to an in-depth understanding of the local meteorology, cloud-seeding programs require an accurate characterization of the background aerosol particles and their potential to serve as CCN or INPs. To this end, the UAE Unified Aerosol Experiment (UAE²) was the first comprehensive airborne assessment of aerosol characteristics in the Arabian Gulf (Reid et al., 2006). The measurements indicated an abundance of sulfate-dominant fine-mode aerosols which may have strong influences for cloud and precipitation formation. Further evaluation of the impact of background aerosols as CCN/INP on cloud microphysics and precipitation was recommended as a high priority to refine operational cloud-seeding activities over the UAE (Semeniuk et al., 2014; Breed et al., 2007).

In line with the recommendations from the UAE² work, a flight campaign was conducted in August 2019 by the Stratton Park Engineering Company (SPEC) Inc., supported by the UAE Rainfall Enhancement Program (UAEREP). The aircraft was equipped with state-of-the-art cloud physics instrumentation (see Sect. 4) to sample summertime convective clouds over the UAE (Lawson et al., 2019). Here, we use in situ aerosol and cloud microphysical measurements from the SPEC UAE campaign to assess the role of background aerosols on natural precipitation formation. Representative measurements from two separate flight cases (August 12th and 19th, 2019) are used to study dominant eastern orographic convection along the Hajar mountains (Branch et al., 2020) and the less frequent southwestern convection associated with the Arabian heat low (AHL) near the Saudi Arabian border (Steinhoff et al., 2018).

80 **2 Regional setting**

Given its geographical setting, the UAE is impacted by both mesoscale sea-land breezes and microscale orographic storms with lifetimes less than 30 minutes. Interestingly, despite its minimal contribution to the country's mean annual rainfall, the summer season records rainfall accumulations exceeding 100 mm over the eastern Hajar mountains bordering Oman (Wehbe et al., 2020; Wehbe et al., 2017). In a recent study, Branch et al. (2020) calculated back-trajectories of summertime convective events over the eastern UAE by applying thermal radiance thresholds to the European Organization for the Exploitation of Meteorological Satellites (EUMETSAT) Meteosat-7 and 10 retrievals during a 7-year period. While convection was found to be predominantly fostered along the Hajar mountain peaks (>80% of events) around noon local time, a diurnal propagation of this convection outflows to the west during the afternoon periods – coincident with the easterly sea breeze (Remiszewska et al., 2007).

90 On the other hand, rare non-orographic summertime precipitation has also been reported over the UAE. Steinhoff et al. (2018) linked such events over the southwest to increased convection and latent heat over the Arabian Sea associated with the active phases of the southwest Asian monsoon. This was shown to strengthen the circulation of the AHL, which shifted moisture transport in favour of deep convection initiation over the southwest desert inland. Kumar and Suzuki (2019) also used



95 Meteosat-10 retrievals for a satellite-based assessment of cloud climatology and seasonal variability of microphysical
properties over the UAE. They reported a high occurrence of mixed-phase convective clouds along the coast and northeastern
mountains during summer periods. Their analyses of cloud types showed higher warm cloud fractions over the mountainous
regions, while cold cloud fractions were higher over the Arabian Gulf during winter and strictly localized along the Hajar
mountains in summer. This is consistent with the radar-based analysis of Breed et al. (2007) which showed the highest
frequency of mixed-phase clouds from orographic convection along the Hajar mountains during the summer months.
100 Consequently, the August 2019 period was selected as the target period for the SPEC UAE campaign. A total of 11 science
flights (SFs) sampled frequent orographic clouds west of the Hajar foothills, in addition to one set of non-orographic clouds
over the southwest near the Saudi border. Figure 1 shows the flight tracks during the August 12th and 19th cases, referred to
hereafter as SF1 and SF4, respectively, and the composite reflectivity from the UAE National Center of Meteorology (NCM)
radar network.

105 3 Synoptic situation and thermodynamic profiles

Figure 2a-d shows the synoptic situation using the European Centre for Medium Range Weather Forecasts Reanalysis 5
(ERA5) product (Hersbach and Dee, 2016) within 1-hour before each flight – at 11:00 and 12:00 UTC for SF1 and SF4,
respectively. The mean sea-level pressure (MSLP) during SF1 (Fig. 2a) shows the two typical pressure lows impacting the
UAE during the summer season, namely, the Pakistan-India low (Bollasina and Nigam, 2011) and the Arabian Gulf low (Bitan
110 and Sa'Arani, 1992). However, the SF4 case (Fig. 2b) occurred during the less-frequent dominance of the AHL (Steinhoff et
al., 2018) over the southwestern region. The low-level atmospheric thickness (LLAT) index is used for heat low detection
(Fig. 2c,d) as proposed by Lavaysse et al. (2009) for the West African heat low. The varying threshold (red color scale) is
defined as the 90th percentile of the LLAT cumulative probability distribution function (i.e. the highest 10% values of LLAT)
computed from all the grid points (hourly scale used here). The active AHL extends from central Saudi Arabia to the southwest
115 of the UAE during SF4 (Fig. 2d).

To further assess discrepancies between the two flight cases, air mass back-trajectories were computed using the Hybrid Single
Particle Lagrangian Integrated Trajectory (HYSPPLIT) model (Stein et al., 2015). Figure 2e,f show the 1.5 and 4.5 km level
back-trajectories to two locations within the flight area and a third control location to the south for each flight date to distinguish
between regional and local air mass trajectories. During SF1, all low-level (1.5 km) trajectories (Fig. 2e) show the dominant
sea breeze convergence inland and toward the Hajar foothills, while the upper-levels (4.5 km) indicate long-range easterly
120 transport from the Indian regime. However, during SF4, more locally driven trajectories aligning with the extent of the AHL
are observed over the Empty Quarter desert area to the west.

Figure 3 shows the Skew-T profiles comprised of the aircraft-observed temperature and dew point temperature from the surface
at 970 hPa (43 °C) to 400 hPa (-15 °C) for both flight case. Co-located ERA5 profiles (Fig. 3a,b) and radiosonde observations



125 (Fig. 3c,d) from Abu Dhabi (AD) Airport (12:00 UTC) are overlaid for each case. Despite the different synoptic situations
during the SF1 and SF4 cases, similar cloud base heights and temperatures of 3.5 km and 9 °C, respectively, are observed.
These are typical values which fall within the long-term interquartile range of summertime convection (35-year analysis of
AD Airport soundings, not shown). However, much drier conditions are evident in SF4 compared to SF1 soundings above the
700-hPa level. Although the ERA5 dataset provides a high vertical resolution (~250 m), it represents the mean conditions
130 within a single ERA5 grid cell (~30 km). Hence, the mean condition as presented in the ERA5 data may not represent the
aircraft observed instance due to the high variability of the low- to mid-level atmospheric conditions in this region. Strong
upper-level temperature inversions at around -5 °C and -3 °C are observed during SF1 and SF4, respectively. The large
differences in the low-level dew point profiles between the aircraft and AD soundings is primarily due to the coastal location
of the latter. The noise in the dew point temperature profiles between the 700 and 500 hPa levels indicates the aircrafts' motion
135 in and out of clouds.

4 Dataset and methods

4.1 Aircraft instrumentation

The science flights were conducted by the SPEC Learjet 35A. The capability of the Learjet to conduct rapid maneuvers and
swift ascents/descents was key for sampling the short-lived UAE clouds while complying with local Air Traffic Control (ATC)
140 restrictions. The aircraft was equipped with state-of-the-art cloud physics instrumentation used on multiple platforms
throughout various airborne campaigns (Lawson et al., 2019). The instruments deployed for the UAE campaign included a
Condensation Particle Counter (CPC), Passive Cavity Aerosol Spectrometer Probe (PCASP), multiple optical array probes
(OAPs) and scattering probes (see Table 1). Figure 4 shows the Learjet in-flight over Al Ain and selected instruments.

The scattering probes included a standalone Fast Cloud Droplet Probe (FCDP) (Lawson et al., 2017) and Fast Forward-
145 scattering Spectrometer (FFSSP), with the latter upgraded to reduce the effects of ice shattering and processing lag time
(O'Connor, 2008). The scattering probes' retrieval and processing mechanisms are provided in Knollenberg (1981) and in the
Appendix of Lawson et al. (2017). The instrument suite also included the Hawkeye (Woods et al., 2018), a recently developed
composite system housing its own FCDP, two-dimensional stereo (2D-S) probe, and Cloud Particle Imager (CPI) (Lawson et
al., 2001). The CPI provides high resolution (2.3 μm) digital images of cloud particles for ice-phase habit identification (Woods
150 et al., 2018). The triple-coincident observations from the Hawkeye were especially useful for the mixed-phase clouds targeted
in the UAE, where a size threshold was applied on the 2D-S as a trigger for the CPI to capture ice particle habits within a large
population of cloud drops. Only the CPI component of the Hawkeye was used in this paper, given the Hawkeye-FCDP
measurements required further calibration and testing.



The OAPs included a total of three 2D-S probes (two standalone and one within the aforementioned Hawkeye) and one High
155 Volume Precipitation Spectrometer (HVPS) described in Lawson et al. (2006) and Lawson et al. (1993), respectively. A total
of five channels (three vertical and two horizontal) provided redundant measurements at 10 μm resolution (see Table 1) which
served for data quality control (see Table 1). Post-processing of the OAP imagery involved noise filtering and corrections for
particle overlapping and ice shattering as outlined in Lawson (2011). Size estimates from the two image processing methods
 M_4 and M_7 described in Lawson et al. (2017) were used for smaller (round drops) and larger (irregular ice) shapes, respectively.

160 As listed in Table 1, cloud droplet size distributions were derived using the well-established FFSSP and 2D-S instruments,
while the usage of the standalone FCDP was limited to coarse mode aerosol measurements to compliment the PCASP
accumulation mode measurements. The fine-mode PCASP measurements were quality controlled by adjusting the sample
volume to account for pressure-induced pump rate lags, especially at altitudes exceeding 4.5 km. The first two size bins (0.1–
0.12 μm) were also excluded from the total concentration calculations to avoid data contamination from excessive noise. All
165 PCASP total concentrations were filtered for out-of-cloud measurements by applying dual thresholds on the FFSSP total
number concentration (20 cm^{-3}) and hot-wire liquid water content (LWC; $0.01 \text{ g}\cdot\text{m}^{-3}$), above which is considered a cloud pass
(Korolev and Isaac, 2006). Similar to a CCN counter, but forced with higher supersaturation, the CPC provided counts of ultra-
fine mode aerosols (0.01–3 μm) through deliberate water-based condensation of intercepted particles to reach sizes detectable
by a laser counter (Wang et al., 2015; Liu et al., 2006). Finally, an Aventech Model Aircraft Integrated Meteorological
170 Measurement System (AIMMS) logged 1-Hz basic meteorological variables and air motion measurements (Beswick et al.,
2008), while a Nevzorov hot-wire probe (Korolev et al., 1998) measured the total water content and LWC – used here as the
reference value compared to LWCs registered by the scattering probes and OAPs.

4.2 Cloud penetration selection and classification

A total of 13 and 22 cloud penetrations were conducted during SF1 and SF4, respectively. All flight ascents were southward
175 out of Al Ain, climbing up to 7 km (flight altitude limit set by ATC) and penetrated clouds on the descent back over Al Ain
(SF1) or the southwest (SF4). Growing turrets were penetrated nearest to their tops to document initial ice formation (i.e. first
ice). However, as a result of the 7 km ATC limit, there are no cloud measurements at levels colder than -12°C . Penetrations
were also made around 300 m above cloud bases to measure the microphysical and dynamic properties of the updrafts. Here,
the analysis is primarily focused on penetrations from growing turrets, aiming to study early-stage cloud conditions and their
180 evolution. Penetrations from precipitating, decaying or dissipating clouds are excluded from the analysis (used only for
demonstration).

To minimize data contamination from entrainment and evaporation effects or artificial spikes in the 1-Hz acquisitions, only
measurements from cloud cores are extracted. Avoiding entrainment effects is particularly challenging in this region's dry
environment where even the most undiluted penetrations are contaminated by downdrafts. The analysed cloud cores were sub-



185 adiabatic in all cases. The Steady State (SS) and Maximum Drop Concentration (MDC) methods proposed by Tessendorf et
al. (2013) are used to manually identify the most isolated cloud core measurements within each penetration time series. Both
methods are used to identify the undiluted core and better attribute the measured spectra to the sub-cloud aerosol. As the name
suggests, the SS method is based on the assumption that LWC and droplet concentrations remain relatively constant during a
3-second window, while the MDC method averages the measurements over the 3-second window of maximum concentrations
190 to better correlate with sub-cloud aerosol measurements. Both methods are applied by manual inspection of the time series
data. We first present the background aerosol analysis, followed by a detailed microphysical analysis and inter-comparison
between the two.

5 Results and discussion

5.1 Aerosol measurements

195 Figure 5 shows the 3D flight tracks from both cases and their PCASP total concentration measurements throughout. For both
flight cases, the PCASP concentrations show a 10-fold increase on the descent over Al Ain compared to those from the ascent,
even at low altitudes (below 1500 m). This can be explained by multiple co-occurring effects. The convective outflow from
the Hajar mountains toward Al Ain and the coincident timing of the SF1 descent with the peak in diurnal sea breeze activity
(around 12:00 UTC) (Eck et al., 2008) can form a convergence zone over the region of Al Ain which is subject to orographic
200 trapping of dust along the Hajar foothills (Schwitalla et al., 2020). The effect of the orographic dust trapping is evident from
the higher PCASP descent concentrations from SF1, which is further east toward the Hajar ridge, compared to that of SF4 with
a direct descent over Al Ain (see Fig. 1). Also, the ascent-descent magnitude split in the PCASP profiles are less pronounced
for flights on non-cloud (dry) days (i.e. SF5 and SF9 – not shown here). This indicates the strong contribution from the outflow
of convection and thunderstorms during cloudy days (SF1 and SF4) coupled with the prevalent dust-laden haboob winds
205 (Miller et al., 2008). The range of penetrated cloud temperatures are shown for each flight, with high concentrations of around
1000 cm⁻³ at cloud base (9 °C) in both cases.

The vertical profiles of PCASP total number concentrations are shown in Figure 6. During the SF1 ascent, the concentrations
decrease from around 400 cm⁻³ to 100 cm⁻³ within the boundary layer up to 2 km, and increase again to 180 cm⁻³ around 4 km.
Additional variations in the change in aerosol concentration with altitude are observed up to the top 7 km level, while the
210 descent shows relatively less variability. This suggests the presence of multiple dust and aerosol layers (labelled L1-4 on Fig.
6a) on the cloud-free ascent and more mixed conditions with higher concentrations are observed on the descent with cloud
development. A well-mixed profile is observed on both the ascents and descents of SF4 as clouds were present over the
southwest and Al Ain, but the magnitude split is still evident between the two profiles. The observed stratification of dust and
aerosol layers is in line with the ground-based Lidar observations from Filioglou et al. (2020) over Al Dhaid – situated along
215 the Hajar foothills at 130 km from Al Ain – where they report four separate layers up to 6 km during August 2018. The



stratification is imposed by gravitational waves produced from the sea breeze-mountain overpasses during the afternoon period and by multiple temperature inversions (see Fig. 3) which are frequently observed during the summer months (Weston et al., 2020).

The time series of altitude (and temperature) measurements from SF1 and SF4 are shown in Figures 7a and 7c, respectively. 220 Cloud bases are sampled at around 3 km and 9 °C in both cases. The selected time series of the sub-cloud and in-cloud LWC (hot-wire), PCASP and FFSSP total number concentrations are displayed in Figures 7b and 7d. Coincident peaks in the LWC and FFSSP total concentrations indicate a cloud base penetration, while intermediate intervals are considered sub-cloud measurements. In both cases, the mean sub-cloud PCASP total number concentrations vary around 500 cm⁻³, but with higher variations (up to 1000 cm⁻³) during SF1. The mean background total concentration of 500 cm⁻³ is in close agreement with those 225 recorded over the Sahara desert by Rosenfeld et al. (2001). On the other hand, the LWC shows excessive noise during the SF4 sub-cloud passes with peaks of around 0.2 g.m⁻³ compared to SF1 (<0.05 g.m⁻³), which may be attributed to the hot-wire response to heavy dust (and haze) loading from local pollution over the southwest. This is corroborated by the more frequent FFSSP measurements during SF4 compared to SF1, suggesting the presence of large background aerosols from local pollution and dust. The CPC concentrations closely match those recorded by the PCASP during SF4 (the CPC was not operational during 230 SF1). Instances where CPC exceed PCASP concentrations within the cores (e.g. 13:53–13:55 in Fig. 7d) allude to the hygroscopic nature of the ultra-fine background aerosols over the southwest, which is consistent with the experimental results of Semeniuk et al. (2015).

Figure 8 shows the background aerosol size spectra from SF1 and SF4 merged from overlaps between the PCASP, FCDP, FFSSP and 2D-S10 measurements. A higher concentration of fine-mode aerosols between 0.1–0.2 μm is observed during SF1 235 compared to SF4 with a peak difference of approximately 5×10⁵ L⁻¹ μm⁻¹. Alternatively, higher tail concentrations from ultra-giant sizes of 20–50 μm are recorded during SF4 compared to SF1. This is in line with the previous suggestion of higher ultra-giant CCN loading from local pollution and dust aggregation over the southwest compared to the eastern regime (Semeniuk et al., 2015).

5.2 Cloud microphysics

240 The SS and MDC methods gave matching instances for the most undiluted cloud core measurements (closest to adiabatic cores) with adiabatic fractions between 0.6–0.8. Figure 9 shows the coincident imagery from the 2D-S10, Hawkeye-CPI and cockpit camera at multiple levels of growing turrets sampled during SF1. The altitude, temperature, vertical velocity range, hot-wire LWC, FFSSP total number concentration and median volume diameter (MVD) are listed in Table 2 for each level. The FFSSP and FCDP (not listed) concentrations just above cloud base (9.1 °C) are comparable at 800 and 1144 cm⁻³, 245 respectively. The maximum LWC of 0.8 g.m⁻³ is coincident with the peak updraft velocity of 3.1 m.s⁻¹ and an 8.7 μm MVD. At the next level (4.5 °C), slightly lower FFSSP (621 cm⁻³) and higher FCDP (1598 cm⁻³) concentrations are recorded with a



stronger updraft (6.9 m.s^{-1}) and higher LWC (1.2 g.m^{-3}) compared to the previous level. The marked difference between the FFSSP and FCDP concentrations at this level suggests that more than 50% of the intercepted hydrometeors are within the medium size range ($d < 6 \text{ }\mu\text{m}$) which are better captured by the FCDP. Similar values are also observed at the $3.2 \text{ }^\circ\text{C}$ level with a comparable updraft (6.2 m.s^{-1}) and LWC (1.2 g.m^{-3}). The 2D-S10 and CPI imagery for the lowest three levels (Fig. 9c–e) confirm the dominance of small drops ($d < 10 \text{ }\mu\text{m}$) with the exception of a couple of large drops captured by the 2D-S10. This is consistent with the calculated MVD values ranging from 8.9 to $10.9 \text{ }\mu\text{m}$ from the three warm levels.

In terms of ice processes in the upper portion, ice is observed by the CPI and 2D-S10 at the $-12.4 \text{ }^\circ\text{C}$ level. Very few ice particles show a habit of sector plates, as expected by nucleation at $-12 \text{ }^\circ\text{C}$ with relatively high liquid water contents of 1.1 g.m^{-3} (Pruppacher and Klett, 1980). This observation is within a decaying turret (-12.4 m.s^{-1} peak downdraft). Interestingly, at the highest sampled level ($-12.6 \text{ }^\circ\text{C}$) of a growing pileus cloud, a dominant population of liquid drops ($d < 50 \text{ }\mu\text{m}$) are observed in the absence of ice particles with a LWC of 1.4 g.m^{-3} and a strong 17.8 m.s^{-1} updraft. The calculated MVDs remain less than $20 \text{ }\mu\text{m}$ in both of these sub-freezing temperature penetrations.

Following the same format of Figure 9, the growing cloud turret penetrations at multiple levels during SF4 are shown in Figure 10. At cloud base (8.6°C), comparable concentrations from the FFSSP (541 cm^{-3}) and FCDP (605 cm^{-3}) are recorded at the peak 3.2 m.s^{-1} updraft and 0.2 g.m^{-3} LWC. The CPI and 2D-S10 capture a few $100 \text{ }\mu\text{m}$ drops which appear to be deliquesced dust and pollution aggregates, as suggested from Figure 7d. At the next slightly colder cloud base (8.3°C), more $100 \text{ }\mu\text{m}$ drops are captured by the CPI with higher total concentration and LWC of 0.5 g.m^{-3} . However, at the next $-0.3 \text{ }^\circ\text{C}$ and -3.5°C levels, small drops ($d < 50 \text{ }\mu\text{m}$) dominate the imagery similar to SF1. Additionally, the LWC remains high ($1.2\text{--}1.3 \text{ g.m}^{-3}$) for the upper sub-freezing levels where graupel and ice irregulars ($0.5\text{--}1 \text{ mm}$) are detected at -10.6 and $-11.2 \text{ }^\circ\text{C}$. The strong updraft (24.4 m.s^{-1}) may have carried a limited number of large particles aloft to serve as ice nuclei at the $-11.2 \text{ }^\circ\text{C}$ level. Hence, the larger drops observed at the lower two levels (8.6 and 8.3°C) most likely originate from melted ice that has fallen from above.

To intercompare the evolution of droplet sizes with altitude, Figure 11 shows the merged size spectra from the PCASP, FFSSP, 2D-S50 and HVPS measurements during SF1 (a) and SF4 (b) at the levels presented in Figures 9 and 10, respectively. For the 9.1°C level of SF1, a peak concentration of $10^5 \text{ L}^{-1} \text{ }\mu\text{m}^{-1}$ is shown for the sizes between $6\text{--}10 \text{ }\mu\text{m}$, which is consistent with the imagery-based analysis (Fig. 9e). For sizes larger than $10 \text{ }\mu\text{m}$, the concentrations drastically decrease to around $10 \text{ L}^{-1} \text{ }\mu\text{m}^{-1}$, while the 2D-S50 shows sizes up to $200 \text{ }\mu\text{m}$ sizes at minimal concentrations of $10^{-3} \text{ L}^{-1} \text{ }\mu\text{m}^{-1}$. At the next two $4.5 \text{ }^\circ\text{C}$ and 3.2°C levels, no broadening is observed with a similar decrease in concentrations for sizes larger than $10 \text{ }\mu\text{m}$. Similarly, the cloud base penetration of SF4 at $8.6 \text{ }^\circ\text{C}$ reveals a consistent peak in the concentrations of $6\text{--}10 \text{ }\mu\text{m}$ sizes, followed by a sharp decrease in concentrations thereafter. However, the $8.3 \text{ }^\circ\text{C}$ and $-0.3 \text{ }^\circ\text{C}$ levels show slight broadening with the peaks extending to $20 \text{ }\mu\text{m}$, which may be explained by the hygroscopic characteristics of the southwest ultra-fine background aerosols (see Fig. 7d).



However, the presence of large haze particles, turbulence and downdrafts may have also contributed to the larger sizes at this level.

280 Rosenfeld and Gutman (1994) report an effective radius larger than 14 μm needed to trigger C-C and warm rain generation, which is in line with other works (Brennguier and Chaumat, 2001). Furthermore, Pinsky et al. (2001) reports a 3% collision efficiency between collector and collected drops of 60 μm and 10 μm sizes, respectively, while the efficiency increases to 45% for collisions with collected drops of 25 μm sizes. Consequently, the concentration of intermediate sizes (10–30 μm) should be at least one order of magnitude larger than any other size range for an efficient C-C process. Given the dominance of small-sized particles with diameters less than 10 μm and the minimal concentrations of intermediate sizes (10–30 μm) from
285 measurements during both flights, an active C-C process is not achieved. A further inhibiting factor is the limited depth of warm cloud (<1000 m) which is critical for the development of C-C (Johnson, 1993).

For the sub-freezing cloud levels of SF1, the broadening of the PSD associated with the decaying turret (-12.4 °C) is primarily attributed to fallout ice (see Fig. 9b) with sizes extending to the millimeter scale. However, for the growing turret with a pure liquid phase at -12.6 °C, the broadening of the size distribution may be a result of the large vertical velocity range (-11.2–17.8
290 $\text{m}\cdot\text{s}^{-1}$) or turbulence that increases the supersaturation perturbations (Abade et al., 2018; Grabowski and Abade, 2017; Lasher-trapp et al., 2005). Dust aerosols, acting as ice nuclei, do not appear to be transported to this level, which agrees with the observations of Filioglou et al. (2020) close to the Hajar foothills. For the SF4 sub-freezing levels, broadening is observed for sizes larger than 100 μm extending up to mm-sized irregulars and graupel at -11.2 °C, but at low concentrations within a dominant population of small liquid drops (see Fig. 10a and b). Unlike the case of SF1, dust and pollution aggregates appear
295 to have served as INPs given the strong updrafts (24.4 $\text{m}\cdot\text{s}^{-1}$) at the -10.6 °C level of SF4.

Several studies show that ultra-giant nuclei ($d > 40 \mu\text{m}$) may serve as precipitation embryos when their concentration is larger than 30 m^{-3} (Bartlett, 1970). The potential sources of such nuclei can be large CCN, sulfate-dominant mineral dust (Wurzler et al., 2000), or simply water insoluble particles serving as coalescence embryos (Lasher-Trapp, 1998). The work of Hoose and Möhler (2012) suggests that dust particles may form ice particles at $T < -15^\circ\text{C}$. However, they may act as INPs at even higher
300 temperatures depending on their chemical composition, size and concentration (Harrison et al., 2016; Peckhaus et al., 2016). Reasonable K-feldspar fractions were reported by Kaufmann et al. (2016) from samples collected over Qatar, while no traces were observed in samples from Oman. This may explain the formation of ice irregulars at the sub-freezing levels of SF4 over the southwest near Qatar, given the ice-nucleating properties of K-feldspar.

5.3 Implications for hygroscopic cloud-seeding

305 The general concept of hygroscopic seeding is based on the notion of introducing large artificial (hygroscopic) particles to compete with smaller naturally occurring aerosols for available cloud LWC. Through this “competition effect”, the seeding



particles are expected to suppress the activation of smaller background aerosols, rapidly grow into larger drops and trigger C-C (Bruitjes et al., 2012; Cooper et al., 1997). Hence, the characteristics of the background aerosol population, namely their size, concentration and chemical composition are considered key precursory properties to determine, and potentially improve, the effectiveness of seeding. Hygroscopic seeding is conducted operationally in several countries and has been actively implemented over the UAE for the past two decades.

The measurements and analysis provided in this study have important implications for operational seeding activities over the UAE. Our results indicate that relatively large aerosol sizes are already present in the UAE environment - over both the eastern and southwestern region - with comparable sizes to typical hygroscopic flare particles ($d < 10 \mu\text{m}$). Furthermore, the ambient aerosols appear to be hygroscopic in nature with their deliquescence and growth to peak concentrations of $20 \mu\text{m}$ sizes above cloud base. This is more pronounced over the southwestern region where mineral aggregates are formed from sea salt and sulfate particles emitted from local oil refineries (Semeniuk et al., 2015). While the C-C process remained inactive in all observed cases, it is hypothesized that the presence of large dust and pollution aggregates causes a “natural competition effect” as reported by Tessoro et al. (2021) based on their aircraft observations from the Queensland Cloud Seeding Research Program (QCSR). Similarly, the modelling work of Segal et al. (2004) indicates a decrease in seeding effects in the presence of large background CCN due to their efficient collision. Hence, given the comparable sizes between the existing GCCN over the UAE and typical seeding particles, it is unclear if hygroscopic seeding can be effective in these clouds. Modelling studies are needed to investigate whether the concentration and hygroscopicity of the background GCCN are high enough to cause a natural competition effect. Furthermore, modelling studies can help assess the effectiveness of perhaps larger seeding particle sizes ($10\text{--}15 \mu\text{m}$) in augmenting this potentially active natural competition effect and/or in initiating C-C.

6 Conclusions

According to the most recent review on precipitation enhancement research by the World Meteorological Organization (WMO) Expert Team on Weather Modification, a more reliable assessment of the aerosol-cloud-precipitation interaction is needed, particularly using in-situ aircraft observations that can validate model results (Flossmann et al., 2019). Located in a regional dust hotspot impacted by air masses originating from five subcontinents over its coastal (west) and mountainous (east) topography, the UAE is considered a “natural laboratory” to study both mesoscale and microscale (aerosol-cloud-precipitation) processes within a limited geographical area – representative of the larger understudied Arabian Peninsula environment.

Here, we present aerosol and cloud microphysical measurements from research flights targeting two distinct summertime convective regimes in the UAE – orographic convection over the Hajar mountains bordering Oman and non-orographic convection over the southwestern Saudi border. Despite their different forcing, the thermodynamic and microphysical properties of summertime convective clouds sampled from both regimes are very similar. Sub-cloud aerosol sizes are shown to extend from 10 nm up to $100 \mu\text{m}$ sizes, with higher concentrations of larger sizes, associated with anthropogenic pollution



over the southwest, acting as ultra-giant CCN. The maximum sizes are approximately double those observed over the Sahara desert (Weinzierl et al., 2009). Despite the existence of ultra-giant CCN, no indications of a natural C-C process are observed.

340 In general, any convective with a cloud depth greater than 500 m is expected to support effective C-C growth (Johnson, 1993). However, the realistically lower LWC within the main body of a convective cloud is more sensitive to cloud base temperature and cloud depth (Warner, 1970). The sampled cloud base temperatures in the UAE (~9 °C) are just below the 10 °C temperature threshold for the onset of an active C-C process (Johnson, 1993; Pruppacher and Klett, 1980), and warm cloud depths never exceed 1000 m in all cases. The observations show a relatively low concentration of background aerosols in the 10–15 μm
345 size range which may further explain the inactivation of C-C. The results have key implications for ongoing operational cloud-seeding activities over the UAE, which currently rely on hygroscopic material with diameters less than 10 μm. Modelling studies are needed to further assess the influence of aerosols on clouds and precipitation in these clouds, as well as to study implications for hygroscopic cloud-seeding in the UAE. Model simulations can be initialized (and validated) using the observations provided in this paper to assess the potential of different seeding strategies for augmenting rainfall over this
350 water-stressed region.

Data availability

The aircraft observations are archived at the UAE National Center of Meteorology. Readers can request the dataset by contacting research@ncms.ae.

Author contribution

355 ST, CW and YW conceptualized the study. CW, SW and PL supported in data curation and visualization. YW performed the formal analysis and wrote the paper. All co-authors advised on the interpretation of results and were involved in the manuscript editing and discussion.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This material is based on work supported by the National Center of Meteorology, under the UAE Research Program for Rain Enhancement Science (UAEREP). The work is also supported by the Research Applications Laboratory (RAL) at the National



Center for Atmospheric Research (NCAR), sponsored by the National Science Foundation under Cooperative Agreement No. 1852977. The authors thank the Stratton Park Engineering Company Inc. for advising on data processing and quality control.

365 References

- Abade, G. C., Grabowski, W. W., and Pawlowska, H.: Broadening of cloud droplet spectra through eddy hopping: Turbulent entraining parcel simulations, *Journal of the Atmospheric Sciences*, 75, 3365-3379, 2018.
- Abuelgasim, A., and Farahat, A.: Effect of dust loadings, meteorological conditions, and local emissions on aerosol mixing and loading variability over highly urbanized semiarid countries: United Arab Emirates case study, *Journal of Atmospheric and Solar-Terrestrial Physics*, 199, 105215, 2020.
- 370 Almazroui, A., and Farrah, S.: The UAE seeks leading position in global rain enhancement research, *Journal of Weather Modification*, 49, 2017.
- Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., Dobbie, S., O'Sullivan, D., and Malkin, T. L.: The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds, *Nature*, 498, 355-358, 2013.
- 375 Bartlett, J.: The effect of revised collision efficiencies on the growth of cloud droplets by coalescence, *Quarterly Journal of the Royal Meteorological Society*, 96, 730-738, 1970.
- Beegum, S. N., Gherboudj, I., Chaouch, N., Temimi, M., and Ghedira, H.: Simulation and analysis of synoptic scale dust storms over the Arabian Peninsula, *Atmospheric Research*, 199, 62-81, 2018.
- Beswick, K., Gallagher, M., Webb, A., Norton, E., and Perry, F.: Application of the Aventech AIMMS20AQ airborne probe for turbulence measurements during the Convective Storm Initiation Project, *Atmospheric Chemistry & Physics*, 8, 2008.
- 380 Bitan, A., and Sa'aroni, H.: The horizontal and vertical extension of the Persian Gulf pressure trough, *International Journal of Climatology*, 12, 733-747, 1992.
- Bollasina, M., and Nigam, S.: The summertime "heat" low over Pakistan/northwestern India: evolution and origin, *Climate dynamics*, 37, 957-970, 2011.
- 385 Branch, O., Behrendt, A., Gong, Z., Schwitalla, T., and Wulfmeyer, V.: Convection Initiation over the Eastern Arabian Peninsula, *Meteorologische Zeitschrift*, 67-77, 2020.
- Breed, D., Brientjes, R., Salazar, V., and Jensen, T.: NCAR feasibility studies for weather modification programs over the past 10 years, Research Applications Laboratory (RAL), National Center for Atmospheric Research (NCAR), Boulder CO, 2007.
- Brenguier, J.-L., and Chaumat, L.: Droplet spectra broadening in cumulus clouds. Part I: Broadening in adiabatic cores, *Journal of the atmospheric sciences*, 58, 628-641, 2001.
- 390 Brientjes, R. T., Salazar, V., Semeniuk, T. A., Buseck, P., Breed, D. W., and Gunkelman, J.: Evaluation of hygroscopic cloud seeding flares, *The Journal of Weather Modification*, 44, 69-94, 2012.
- Chen, S., Xue, L., and Yau, M.-K.: Impact of hygroscopic CCN and turbulence on cloud droplet growth: A parcel-DNS approach, *Atmospheric Chemistry and Physics Discussions*, 2019, 1-17, 2019.
- 395 Cooper, W. A., Brientjes, R. T., and Mather, G. K.: Calculations pertaining to hygroscopic seeding with flares, *Journal of Applied Meteorology*, 36, 1449-1469, 1997.
- DeMott, P. J., Sassen, K., Poellot, M. R., Baumgardner, D., Rogers, D. C., Brooks, S. D., Prenni, A. J., and Kreidenweis, S. M.: African dust aerosols as atmospheric ice nuclei, *Geophysical Research Letters*, 30, 2003.
- 400 Eck, T., Holben, B., Reid, J., Sinyuk, A., Dubovik, O., Smirnov, A., Giles, D., O'Neill, N., Tsay, S. C., and Ji, Q.: Spatial and temporal variability of column-integrated aerosol optical properties in the southern Arabian Gulf and United Arab Emirates in summer, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- Feingold, G., Cotton, W. R., Kreidenweis, S. M., and Davis, J. T.: The impact of giant cloud condensation nuclei on drizzle formation in stratocumulus: Implications for cloud radiative properties, *Journal of the atmospheric sciences*, 56, 4100-4117, 1999.
- Filioglou, M., Giannakaki, E., Backman, J., Kesti, J., Hirsikko, A., Engelmann, R., O'Connor, E., Leskinen, J. T. T., Shang, X., Korhonen, H., Lihavainen, H., Romakkaniemi, S., and Komppula, M.: Optical and geometrical aerosol particle properties over the United Arab Emirates, *Atmos. Chem. Phys. Discuss.*, 2020, 1-26, 10.5194/acp-2020-133, 2020.
- 405 Flossmann, A. I., and Wobrock, W.: A review of our understanding of the aerosol-cloud interaction from the perspective of a bin resolved cloud scale modelling, *Atmospheric Research*, 97, 478-497, 2010.
- Flossmann, A. I., Manton, M., Abshaev, A., Brientjes, R., Murakami, M., Prabhakaran, T., and Yao, Z.: Review of advances in precipitation enhancement research, *Bulletin of the American Meteorological Society*, 100, 1465-1480, 2019.
- 410 Freud, E., Ström, J., Rosenfeld, D., Tunved, P., and Swietlicki, E.: Anthropogenic aerosol effects on convective cloud microphysical properties in southern Sweden, *Tellus B: Chemical and Physical Meteorology*, 60, 286-297, 2008.




- Grabowski, W. W., and Abade, G. C.: Broadening of cloud droplet spectra through eddy hopping: Turbulent adiabatic parcel simulations, *Journal of Atmospheric Sciences*, 74, 1485-1493, 2017.
- 415 Harrison, A. D., Whale, T. F., Carpenter, M. A., Holden, M., Neve, L., O'Sullivan, D., Vergara Temprado, J., and Murray, B. J.: Not all feldspars are equal: a survey of ice nucleating properties across the feldspar group of minerals, *Atmospheric Chemistry and Physics*, 16, 10927-10940, 2016.
- Hersbach, H., and Dee, D.: ERA5 reanalysis is in production, *ECMWF newsletter*, 147, 5-6, 2016.
- 420 Hofer, J., Althausen, D., Abdullaev, S. F., Makhmudov, A. N., Nazarov, B. I., Schettler, G., Engelmann, R., Baars, H., Fomba, K. W., and Müller, K.: Long-term profiling of mineral dust and pollution aerosol with multiwavelength polarization Raman lidar at the Central Asian site of Dushanbe, Tajikistan: case studies, *Atmospheric Chemistry and Physics*, 17, 14559-14577, 2017.
- Hoese, C., and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817-9854, 2012.
- 425 Jensen, J. B., and Lee, S.: Giant sea-salt aerosols and warm rain formation in marine stratocumulus, *Journal of the atmospheric sciences*, 65, 3678-3694, 2008.
- Jensen, J. B., and Nugent, A. D.: Condensational growth of drops formed on giant sea-salt aerosol particles, *Journal of the atmospheric sciences*, 74, 679-697, 2017.
- Johnson, D. B.: The role of giant and ultragiant aerosol particles in warm rain initiation, *Journal of the Atmospheric Sciences*, 39, 448-460, 1982.
- 430 Johnson, D. B.: The onset of effective coalescence growth in convective clouds, *Quarterly Journal of the Royal Meteorological Society*, 119, 925-933, 1993.
- Kanji, Z. A., Sullivan, R. C., Niemand, M., DeMott, P. J., Prenni, A. J., Chou, C., Saathoff, H., and Möhler, O.: Heterogeneous ice nucleation properties of natural desert dust particles coated with a surrogate of secondary organic aerosol, *Atmospheric Chemistry and Physics*, 19, 5091-5110, 2019.
- 435 Kant, S., Panda, J., and Gautam, R.: A seasonal analysis of aerosol-cloud-radiation interaction over Indian region during 2000–2017, *Atmospheric environment*, 201, 212-222, 2019.
- Karagulian, F., Temimi, M., Ghebreyesus, D., Weston, M., Kondapalli, N. K., Valappil, V. K., Aldababesh, A., Lyapustin, A., Chaouch, N., and Al Hammadi, F.: Analysis of a severe dust storm and its impact on air quality conditions using WRF-Chem modeling, satellite imagery, and ground observations, *Air Quality, Atmosphere & Health*, 12, 453-470, 2019.
- 440 Kaufmann, L., Marcolli, C., Hofer, J., Pinti, V., Hoyle, C. R., and Peter, T.: Ice nucleation efficiency of natural dust samples in the immersion mode, *Atmospheric Chemistry and Physics*, 16, 11177-11206, 2016.
- Knollenberg, R.: Techniques for probing cloud microstructure, *Symposium on Clouds-Their formation, optical properties, and effects*, Williamsburg, VA, 1981, 15-89,
- 445 Koren, I., Kaufman, Y. J., Rosenfeld, D., Remer, L. A., and Rudich, Y.: Aerosol invigoration and restructuring of Atlantic convective clouds, *Geophysical Research Letters*, 32, 2005.
- Korolev, A., Strapp, J., Isaac, G., and Nevzorov, A.: The Nevzorov airborne hot-wire LWC–TWC probe: Principle of operation and performance characteristics, *Journal of Atmospheric and Oceanic Technology*, 15, 1495-1510, 1998.
- Korolev, A., and Isaac, G. A.: Relative humidity in liquid, mixed-phase, and ice clouds, *Journal of the atmospheric sciences*, 63, 2865-2880, 2006.
- 450 Kumar, K. N., and Suzuki, K.: Assessment of seasonal cloud properties in the United Arab Emirates and adjoining regions from geostationary satellite data, *Remote sensing of environment*, 228, 90-104, 2019.
- Kvasov, B. I.: *Methods of shape-preserving spline approximation*, World Scientific, 2000.
- Lasher-Trapp, S. G.: Ultragiant aerosol growth by collection within a warm continental cumulus congestus, 1998.
- 455 Lasher-trapp, S. G., Cooper, W. A., and Blyth, A. M.: Broadening of droplet size distributions from entrainment and mixing in a cumulus cloud, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 131, 195-220, 2005.
- Lavaysse, C., Flamant, C., Janicot, S., Parker, D., Lafore, J.-P., Sultan, B., and Pelon, J.: Seasonal evolution of the West African heat low: a climatological perspective, *Climate Dynamics*, 33, 313-330, 2009.
- 460 Lawson, P., Gurganus, C., Woods, S., and Bruintjes, R.: Aircraft observations of cumulus microphysics ranging from the tropics to midlatitudes: Implications for a “new” secondary ice process, *Journal of the Atmospheric Sciences*, 74, 2899-2920, 2017.
- Lawson, R., Stewart, R., Strapp, J., and Isaac, G.: Airborne measurements of the origin and growth of very large snowflakes, *Geophys. Res. Lett.*, 20, 53-56, 1993.
- Lawson, R., Baker, B., Schmitt, C., and Jensen, T.: Overview of microphysical properties of summertime boundary layer clouds observed during FIRE. ACE, *J. Geophys. Res.*, 106, 14989-15014, 2001.
- 465 Lawson, R.: Effects of ice particles shattering on the 2D-S probe, *Atmospheric measurement techniques*, 4, 1361, 2011.
- Lawson, R., Woods, S., Jensen, E., Erfani, E., Gurganus, C., Gallagher, M., Connolly, P., Whiteway, J., Baran, A., and May, P.: A review of ice particle shapes in cirrus formed in situ and in anvils, *Journal of Geophysical Research: Atmospheres*, 124, 10049-10090, 2019.



- 470 Lawson, R. P., O'connor, D., Zmarzly, P., Weaver, K., Baker, B., Mo, Q., and Jonsson, H.: The 2D-S (stereo) probe: Design and preliminary tests of a new airborne, high-speed, high-resolution particle imaging probe, *Journal of Atmospheric and Oceanic Technology*, 23, 1462-1477, 2006.
- Liu, W., Kaufman, S. L., Osmondson, B. L., Sem, G. J., Quant, F. R., and Oberreit, D. R.: Water-based condensation particle counters for environmental monitoring of ultrafine particles, *Journal of the Air & Waste Management Association*, 56, 444-455, 2006.
- Liu, Y., Zhu, Q., Hua, S., Alam, K., Dai, T., and Cheng, Y.: Tibetan Plateau driven impact of Taklimakan dust on northern rainfall, *Atmospheric Environment*, 117583, 2020.
- 475 Lohmann, U., Rotstajn, L., Storelvmo, T., Jones, A., Menon, S., Quaas, J., Ekman, A., Koch, D., and Ruedy, R.: Total aerosol effect: forcing or radiative flux perturbation, *Atmospheric Chemistry and Physics*, 10, 3235-3246, 2010.
- Miller, S. D., Kuciauskas, A. P., Liu, M., Ji, Q., Reid, J. S., Breed, D. W., Walker, A. L., and Mandoos, A. A.: Haboob dust storms of the southern Arabian Peninsula, *Journal of Geophysical Research: Atmospheres*, 113, 2008.
- 480 Min, Q.-L., Li, R., Lin, B., Joseph, E., Wang, S., Hu, Y., Morris, V., and Chang, F.: Evidence of mineral dust altering cloud microphysics and precipitation, *Atmospheric Chemistry & Physics*, 9, 2009.
- Ming, Y., Ramaswamy, V., Ginoux, P. A., and Horowitz, L. H.: Direct radiative forcing of anthropogenic organic aerosol, *Journal of Geophysical Research: Atmospheres*, 110, 2005.
- Möhler, O., Field, P., Connolly, P., Benz, S., Saathoff, H., Schnaiter, M., Wagner, R., Cotton, R., Krämer, M., and Mangold, A.: Efficiency of the deposition mode ice nucleation on mineral dust particles, 2006.
- 485 Morrison, H., Shupe, M. D., Pinto, J. O., and Curry, J. A.: Possible roles of ice nucleation mode and ice nuclei depletion in the extended lifetime of Arctic mixed-phase clouds, *Geophysical research letters*, 32, 2005.
- Morrison, H., van Lier-Walqui, M., Fridlind, A. M., Grabowski, W. W., Harrington, J. Y., Hoose, C., Korolev, A., Kumjian, M. R., Milbrandt, J. A., and Pawlowska, H.: Confronting the challenge of modeling cloud and precipitation microphysics, *Journal of advances in modeling earth systems*, 12, e2019MS001689, 2020.
- 490 O'Connor, D., B. Baker, and R. P. Lawson: Upgrades to the FSSP-100 Electronics, 15th. Int. Conf. on Clouds and Precipitation, Cancun, Mexico, Universidad Nacional Autónoma de México, 2008, 13.16.
- Peckhaus, A., Kiselev, A., Hiron, T., Ebert, M., and Leisner, T.: A comparative study of K-rich and Na/Ca-rich feldspar ice-nucleating particles in a nanoliter droplet freezing assay, *Atmospheric Chemistry and Physics*, 16, 11477, 2016.
- 495 Pinsky, M., Khain, A., and Shapiro, M.: Collision efficiency of drops in a wide range of Reynolds numbers: Effects of pressure on spectrum evolution, *Journal of the atmospheric sciences*, 58, 742-764, 2001.
- Prenni, A. J., Petters, M. D., Kreidenweis, S. M., Heald, C. L., Martin, S. T., Artaxo, P., Garland, R. M., Wollny, A. G., and Pöschl, U.: Relative roles of biogenic emissions and Saharan dust as ice nuclei in the Amazon basin, *Nature Geoscience*, 2, 402-405, 2009.
- Pruppacher, H. R., and Klett, J. D.: Microphysics of clouds and precipitation, *Nature*, 284, 88-88, 1980.
- 500 Reid, J., Westphal, D., Reid, E., Walker, A., Liu, M., Miller, S., and Kuciauskas, A.: The United Arab Emirates Unified Aerosol Experiment (UAE2), Naval Research Lab Monterey CA Marine, Meteorology Div, 2006.
- Remiszewska, J., Flatau, P., Markowicz, K., Reid, E., Reid, J., and Witek, M.: Modulation of the aerosol absorption and single-scattering albedo due to synoptic scale and sea breeze circulations: United Arab Emirates experiment perspective, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- 505 Rosenfeld, D., and Gutman, G.: Retrieving microphysical properties near the tops of potential rain clouds by multispectral analysis of AVHRR data, *Atmospheric research*, 34, 259-283, 1994.
- Rosenfeld, D.: TRMM observed first direct evidence of smoke from forest fires inhibiting rainfall, *Geophysical research letters*, 26, 3105-3108, 1999.
- Rosenfeld, D.: Suppression of rain and snow by urban and industrial air pollution, *Science*, 287, 1793-1796, 2000.
- 510 Rosenfeld, D., Rudich, Y., and Lahav, R.: Desert dust suppressing precipitation: A possible desertification feedback loop, *Proceedings of the National Academy of Sciences*, 98, 5975-5980, 2001.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A., and Andreae, M. O.: Flood or drought: How do aerosols affect precipitation?, *science*, 321, 1309-1313, 2008.
- Sassen, K., DeMott, P. J., Prospero, J. M., and Poellot, M. R.: Saharan dust storms and indirect aerosol effects on clouds: CRYSTAL-FACE results, *Geophysical Research Letters*, 30, 2003.
- 515 Schwitalla, T., Branch, O., and Wulfmeyer, V.: Sensitivity study of the planetary boundary layer and microphysical schemes to the initialization of convection over the Arabian Peninsula, *Quarterly Journal of the Royal Meteorological Society*, 146, 846-869, 2020.
- Segal, Y., Khain, A., Pinsky, M., and Rosenfeld, D.: Effects of hygroscopic seeding on raindrop formation as seen from simulations using a 2000-bin spectral cloud parcel model, *Atmospheric Research*, 71, 3-34, 2004.
- 520 Semeniuk, T., Brintjes, R., Salazar, V., Breed, D., Jensen, T., and Buseck, P.: Individual aerosol particles in ambient and updraft conditions below convective cloud bases in the Oman mountain region, *Journal of Geophysical Research: Atmospheres*, 119, 2511-2528, 2014.
- Semeniuk, T., Brintjes, R., Salazar, V., Breed, D., Jensen, T., and Buseck, P.: Processing of aerosol particles within the Habshan pollution plume, *Journal of Geophysical Research: Atmospheres*, 120, 1996-2012, 2015.




- Solomon, S., Manning, M., Marquis, M., and Qin, D.: Climate change 2007-the physical science basis: Working group I contribution to the fourth assessment report of the IPCC, Cambridge university press, 2007.
- 525 Squires, P.: The microstructure and colloidal stability of warm clouds: Part I—The relation between structure and stability, *Tellus*, 10, 256-261, 1958.
- Stein, A., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 2015.
- 530 Steinhoff, D. F., Bruintjes, R., Hacker, J., Keller, T., Williams, C., Jensen, T., Al Mandous, A., and Al Yazeedi, O. A.: Influences of the Monsoon Trough and Arabian Heat Low on Summer Rainfall over the United Arab Emirates, *Monthly Weather Review*, 146, 1383-1403, 2018.
- Takeda, T., and Kuba, N.: Numerical study of the effect of CCN on the size distribution of cloud droplets, *Journal of the Meteorological Society of Japan. Ser. II*, 60, 978-993, 1982.
- 535 Tao, W. K., Li, X., Khain, A., Matsui, T., Lang, S., and Simpson, J.: Role of atmospheric aerosol concentration on deep convective precipitation: Cloud-resolving model simulations, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Tessendorf, S. A., Weeks, C. E., Axisa, D., and Bruintjes, R. T.: Aerosol characteristics observed in southeast Queensland and implications for cloud microphysics, *Journal of Geophysical Research: Atmospheres*, 118, 2858-2871, 2013.
- 540 Tessendorf, S. A., Chen, S., Weeks, C., Bruintjes, R., Rasmussen, R. M., and Xue, L.: The influence of hygroscopic flare seeding on drop size distribution over southeast Queensland, *Journal of Geophysical Research: Atmospheres*, n/a, e2020JD033771, <https://doi.org/10.1029/2020JD033771>, 2021.
- Wang, Z., Su, H., Wang, X., Ma, N., Wiedensohler, A., Pöschl, U., and Cheng, Y.: Scanning supersaturation condensation particle counter applied as a nano-CCN counter for size-resolved analysis of the hygroscopicity and chemical composition of nanoparticles, 2015.
- Warner, J.: On steady-state one-dimensional models of cumulus convection, *Journal of the Atmospheric Sciences*, 27, 1035-1040, 1970.
- 545 Wehbe, Y., Ghebreyesus, D., Temimi, M., Milewski, A., and Al Mandous, A.: Assessment of the consistency among global precipitation products over the United Arab Emirates, *Journal of Hydrology: Regional Studies*, 12, 122-135, 2017.
- Wehbe, Y., Temimi, M., Ghebreyesus, D. T., Milewski, A., Norouzi, H., and Ibrahim, E.: Consistency of precipitation products over the Arabian Peninsula and interactions with soil moisture and water storage, *Hydrological Sciences Journal*, 63, 408-425, 2018.
- Wehbe, Y., Temimi, M., and Adler, R. F.: Enhancing Precipitation Estimates Through the Fusion of Weather Radar, Satellite Retrievals, and Surface Parameters, *Remote Sensing*, 12, 1342, 2020.
- 550 Weinzierl, B., Petzold, A., Esselborn, M., Wirth, M., Rasp, K., Kandler, K., Schuetz, L., Koepke, P., and Fiebig, M.: Airborne measurements of dust layer properties, particle size distribution and mixing state of Saharan dust during SAMUM 2006, *Tellus B: Chemical and Physical Meteorology*, 61, 96-117, 2009.
- Weston, M. J., Temimi, M., Nelli, N. R., Fonseca, R. M., Thota, M. S., and Valappil, V. K.: On the Analysis of the Low-Level Double Temperature Inversion Over the United Arab Emirates: A Case Study During April 2019, *IEEE Geoscience and Remote Sensing Letters*, 2020.
- 555 Wise, M. E., Semeniuk, T. A., Bruintjes, R., Martin, S. T., Russell, L. M., and Buseck, P. R.: Hygroscopic behavior of NaCl-bearing natural aerosol particles using environmental transmission electron microscopy, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Woods, S., Lawson, R. P., Jensen, E., Bui, T., Thornberry, T., Rollins, A., Pfister, L., and Avery, M.: Microphysical properties of tropical tropopause layer cirrus, *Journal of Geophysical Research: Atmospheres*, 123, 6053-6069, 2018.
- 560 Wurzler, S., Reisin, T. G., and Levin, Z.: Modification of mineral dust particles by cloud processing and subsequent effects on drop size distributions, *Journal of Geophysical Research: Atmospheres*, 105, 4501-4512, 2000.
- Yin, Y., Levin, Z., Reisin, T. G., and Tzivion, S.: The effects of giant cloud condensation nuclei on the development of precipitation in convective clouds—A numerical study, *Atmospheric research*, 53, 91-116, 2000.




565

Figure 1. Terrain elevation (30 m, ASTER DEM), locations of the C-band radars, reflectivity (dBz) and flight tracks of the August 12th (SF1) and 19th (SF4) cases.




570 **Figure 2.** MSLP (a,b) and LLAT (c,d) with 925 hPa wind directions from ERA5 reanalysis, and HYSPLIT back-trajectories (e,f) during SF1 and SF4 cases. Red back trajectories had lower altitude end points (1500 m) while blue back trajectories had end points at 4500 m. Different line styles are used to show back-trajectories to each of the three labelled end locations.



575 **Figure 3.** Skew-T soundings from SF1 and SF4 descents (limited at 400 hPa level), co-located ERA5 grid points (a,b) and Abu Dhabi Airport (12:00 UTC) observations (c,d).




Figure 4. Learjet in-flight over Al Ain and selected instruments: HVPS, FCDP, AIMMS and Hawkeye (from left to right).




580

Figure 5. 3D flight track of SF1 and SF4 with PCASP total number concentrations (cm^{-3}). SF1 penetrations over Al Ain from cloud tops at -13 °C down to 9 °C bases (a), and SF4 penetrations over the southwest from cloud tops at -12 °C down to 9 °C bases (b) are shown.




585

Figure 6. Altitude profiles of out-of-cloud PCASP total number concentrations for SF1 (a) and SF4 (b). L1-4 designate multiple dust layers at levels of significant change in concentration gradients. The profiles are computed as the best-fit from shape-preserving spline interpolation (Kvasov, 2000) of the full 1-Hz dataset.



590

Figure 7. Flight profiles for SF1 (a) and SF4 (c) and time series of PCASP, FFSSP, CPC and hot-wire LWC measurements for sub-cloud intervals during SF1 over Al Ain (b) and SF4 over the southwest (d).



595

Figure 8. Mean sub-cloud aerosol size distributions for during SF1 (Al Ain) and SF4 (southwest).




Figure 9. SF1 coincident imagery from the 2D-S10, Hawkeye-CPI and cockpit camera. Measurements from each penetration level are listed in Table 2.





Figure 10. Same as Figure 9 but for SF4. Measurements from each penetration level are listed in Table 2.



605

Figure 11. Size distributions at multiple temperature levels of growing turrets from the PCASP (dashed), FFSSP (solid), 2D-S50 (dashed) and HVPS (solid with circle markers) measurements during SF1 (a) and SF4 (b).

610

Table 1. List of instruments used for the collection of measurements used herein.

Instrument	Purpose	Size range	Used here
CPC	Ultra-fine mode aerosol number conc.	0.01 – 3 μm	✓
PCASP	Aerosol spectra and conc.	0.1 – 3 μm	✓
FCDP	Coarse mode aerosol spectra and conc.	1.5 – 50 μm	✓
FCDP-Hawkeye	"	"	
FFSSP	Cloud particle spectra and conc.	3 – 100 μm	✓
2D-S10HV*	Cloud particle shapes and spectra	10 – 3000 μm	✓
2D-S10HV-NCAR	"	"	
2D-S10V-Hawkeye	"	"	
2D-S50H-Hawkeye	"	50 – 6400 μm	✓
CPI-Hawkeye	Cloud particle shapes (ice-phase habit)	2.3 – 2,300 μm	✓
HVPS	Precipitation particle shapes and spectra	150 μm – 2 cm	✓
Nevzorov Hot-wire	Liquid and total water content		✓
AIMMS	Basic meteorological variables		✓

(*) 10HV indicates 10 μm resolution in both the horizontal (H) and vertical (V) channels.

(*) 50H indicates 50 μm resolution in the horizontal (H) channel.



615 **Table 2.** Penetration altitude, temperature (T), vertical velocity range (Vv), FFSSP total number concentration (N-FFSSP) and MVD values from SF1 and SF4 (see Fig. 9 and 10).

Altitude (m)	T (°C)	Vv (m.s ⁻¹)	LWC (g.m ⁻³)	N-FFSSP (cm ⁻³)	MVD (µm)
SF1 penetration levels (Fig. 9a-e)					
7040	-12.6	[-11.2 ; 17.8]	1.4	442	17.4
7050	-12.4	[-12.4 ; 1.1]	1.1	223	19.2
4660	3.2	[-1.3 ; 6.2]	1.2	619	10.9
4430	4.5	[-2.9 ; 6.9]	1.2	621	9.6
3720	9.1	[-2.4 ; 3.1]	0.8	800	8.7
SF4 penetration levels (Fig. 10a-f)					
6710	-11.2	[-2.6 ; 4]	1.2	321	19.4
6700	-10.6	[-7.6 ; 24.4]	1.3	778	19.5
5400	-3.5	[-6.1 ; 4.6]	1.1	479	17.1
4910	-0.3	[-3.4 ; 11.4]	1.3	470	13.2
3620	8.3	[-2.3 ; 3.3]	0.5	753	10.6
3500	8.6	[-0.5 ; 3.2]	0.2	541	7.1