

Coupled and decoupled stratocumulus-topped boundary layers: turbulence properties

Jakub L. Nowak¹, Holger Siebert², Kai-Erik Szodry², and Szymon P. Malinowski¹

¹Institute of Geophysics, Faculty of Physics, University of Warsaw, Pasteura 5, 02-293 Warsaw, Poland

²Leibniz Institute for Tropospheric Research, Permoserstr. 15, 04318 Leipzig, Germany

Correspondence: Jakub L. Nowak (jakub.nowak@fuw.edu.pl)

Abstract. We compare turbulence properties in coupled and decoupled marine stratocumulus-topped boundary layers (STBLs) using high resolution in situ measurements performed by the helicopter-borne platform ACTOS in the region of Eastern North Atlantic.

The thermodynamically well-mixed coupled STBL was characterized by a comparable latent heat flux at the surface and in the cloud top region, and substantially smaller sensible heat flux in the entire depth. Turbulence kinetic energy (TKE) was efficiently generated by buoyancy in the cloud and at the surface, and dissipated with comparable rate across the entire depth. Structure functions and power spectra of velocity fluctuations in the inertial range were reasonably consistent with the predictions of Kolmogorov theory. The turbulence was close to isotropic.

In the decoupled STBL, decoupling was most obvious in humidity profiles. Heat fluxes and buoyant TKE production at the surface were similar to the coupled case. Around the transition level, latent heat flux decreased to zero and TKE was consumed by weak stability. In the cloud top region heat fluxes almost vanished and buoyancy production was significantly smaller than for the coupled case. TKE dissipation rate inside the decoupled STBL varied between its sublayers. Structure functions and power spectra in the inertial range deviated from Kolmogorov scaling. This was more pronounced in the cloud and subcloud layer in comparison to the surface mixed layer. The turbulence was more anisotropic than in the coupled STBL, with horizontal fluctuations dominating. The degree of anisotropy was largest in the cloud and subcloud layer of the decoupled STBL.

Integral length scales, of the order of 100 m in both cases, indicate turbulent eddies smaller than the depth of the coupled STBL or of the sublayers of the decoupled STBL. We hypothesize that turbulence produced in the cloud or close to the surface is redistributed across the entire coupled STBL but rather only inside the sublayers where it was generated in the case of the decoupled STBL. Scattered cumulus convection, developed below the stratocumulus base, may play a role in transport between those sublayers.

1 Introduction

Low-level stratocumulus clouds cover around 20 % of the Earth's surface in annual mean, more than any other cloud type. They occupy upper few hundred meters of the planetary boundary layer, preferentially in the conditions of large-scale subsidence, strong lower-tropospheric stability and moisture supply from the surface (Wood, 2012). Those are usually present in the

25 regions of subtropical and midlatitude oceans with upwelling of cold deep water. Wide-spread presence, persistence and high albedo makes marine stratocumulus important for the energy balance of the planet (Hartmann et al., 1992). Minor variations in coverage and optical thickness impact the radiation budget, therefore also model-based climate predictions (Boucher et al., 2013; Schneider et al., 2019).

The primary mechanism driving the circulation inside stratocumulus-topped boundary layer (STBL) is longwave radiative
30 cooling at the cloud top which produces convective instability. An additional source of turbulence is provided by surface buoyancy, wind shear, latent heat release in updrafts, evaporation in downdrafts or evaporative cooling associated with entrainment of dry, warm air from the free troposphere (Lilly, 1968; Stevens, 2002; Gerber et al., 2016; Mellado, 2017). Properties of the STBL are dependent on the level to which stratocumulus cloud is coupled with sea surface fluxes, in particular of latent and sensible heat (Bretherton and Wyant, 1997; Xiao et al., 2011; Zheng et al., 2018a).

35 Moderately shallow STBLs are often well mixed (Stull, 1988; Markowski and Richardson, 2010). Their typical vertical structure features an adiabatic lapse rate (dry below cloud, moist inside), a strong capping inversion at the top, near-constant concentration of moist-conserved variables (such as total water mass fraction and liquid water potential temperature) from the surface up to the inversion. However, when the circulation ceases to mix the air over entire depth, the STBL becomes decoupled, i.e. the cloud is separated from the moisture supply from the surface (Nicholls, 1984; Turton and Nicholls, 1987;
40 Wood, 2012). The radiatively driven stratocumulus layer (SCL) and the subcloud layer (SBL) in the upper part might be still mixed by negatively buoyant eddies generated at cloud top while the surface mixed layer (SML) at the bottom by positive buoyancy or shear. A stable or conditionally unstable intermediate transition layer (TSL) emerges in between. Conditional instability allows for the cumulus updrafts to penetrate through and intermittently restore the coupling (Bretherton and Wyant, 1997; De Roode and Duynkerke, 1997).

45 Decoupling can be caused either by reducing the intensity of radiatively driven circulation in relation to STBL depth or by stabilizing the subcloud layer (Zheng et al., 2018b). The first possibility might be realized with daytime shortwave radiative heating which offsets longwave cooling (Nicholls, 1984; Turton and Nicholls, 1987) or by extensive entrainment of warm and dry free-tropospheric air which deepens the STBL to such an extent that the turbulence is no longer sufficient to sustain the mixing (Bretherton and Wyant, 1997). The second possibility involves stratification of the lower part by cooling, for instance
50 due to precipitation evaporation (Caldwell et al., 2005; Dodson and Small Griswold, 2021) or advection over colder sea surface (Stevens et al., 1998).

STBL decoupling is the factor which strongly influences further evolution of cloud pattern and boundary layer structure. It constitutes an intermediate stage of transition from overcast stratocumulus into shallow cumulus convection over subtropical oceans as the air masses are advected by the trade winds towards the equator (Albrecht et al., 1995; Bretherton and Wyant,
55 1997; De Roode et al., 2016; Zheng et al., 2020). Successful representation and prediction of such transition between the two STBL regimes pose a challenge for atmospheric general circulation models (Xiao et al., 2012), in large part due to limited understanding of the interaction of various processes involved.

Previous observational studies have documented the structure of the coupled and decoupled STBLs in terms of thermodynamic and radiative features (Wood and Bretherton, 2004; Jones et al., 2011; Ghate et al., 2015; Zheng and Li, 2019) as well

60 as aerosol and cloud properties (Dong et al., 2015; Wang et al., 2016; Goren et al., 2018; Zheng et al., 2018b). On the other hand, modeling efforts provided insightful conceptual explanations of the mechanisms leading to a switch between coupled and decoupled regimes (Turton and Nicholls, 1987; Bretherton and Wyant, 1997; Stevens, 2000; Xiao et al., 2011).

Although the concept of circulation and turbulence being insufficiently strong in order to maintain the mixing throughout the entire depth plays a central role in the conventional rationale of decoupling, few works attempted to quantitatively characterize
65 small-scale (integral length scales and below) turbulence (e.g. Lambert and Durand, 1999; Dodson and Small Griswold, 2021). The major reason is the technical difficulty in measuring turbulent fluctuations of wind velocity, temperature or humidity with adequate spatial resolution and accuracy. Within the present study, we compare the properties of turbulence derived from unique helicopter-borne observations performed in coupled and decoupled STBL in the region of Eastern North Atlantic. Particular attention is given to small-scale features and deviations from the assumption of stationary homogeneous isotropic turbulence.

70 The paper is structured as follows. Section 2 introduces the measurements, including instrumentation, sampling strategy and general synoptic conditions. The selection of the two cases, coupled and decoupled STBL, is explained. Section 3 describes the stratification of the STBL in terms of thermodynamics and stability. The division into sublayers is delineated and the degree of coupling is expressed quantitatively according to literature criteria. Section 4 provides relevant details concerning derivation of turbulence parameters. Section 5 compares properties of turbulence: turbulence kinetic energy, its production and
75 dissipation rates, fluxes of sensible and latent heat, anisotropy of turbulent motions, typical length scales. Finally, the results of the comparison are summarized and discussed in the last section.

2 Measurements

2.1 Location and synoptic conditions

Observations were collected in July 2017 during the ACORES (Azores stratoCumulus measurements Of Radiation, turbulEnce
80 and aeroSols) campaign in the Eastern North Atlantic around the island of Graciosa in the Azores archipelago. Comprehensive description of the project, including weather conditions, instrumentation, sampling strategy and selected research highlights is provided by Siebert et al. (2021).

The area of the experiment is considered to be influenced by a wide range of synoptic scale meteorological conditions. Graciosa is located near the boundary of subtropics and mid-latitudes. Therefore, the impacts of both subtropical trade wind
85 system and mid-latitude cyclones are relevant. The climatology of the marine boundary layer was inferred by Rémillard et al. (2012) based on the long-term ground-based measurements of CAP-MBL project (Wood et al., 2015) utilizing the Atmospheric Radiation Measurement (ARM) facility established right next to the Graciosa airport. They reported that boundary layer decoupling and multiple cloud types (for instance cumulus under stratocumulus) are very frequent at the site throughout the year. Indeed, the range of weather conditions was observed during the ACORES, related to the location and strength of Azores high,
90 as well as occasional front passages (Siebert et al., 2021).

2.2 Instrumentation

Measurements were performed with the Airborne Cloud Turbulence Observations System (ACTOS, Siebert et al. (2006a)) and the Spectral Modular Airborne Radiation measurement sysTem - HELIcopter-borne ObservationS (SMART-HELIOS, Werner et al. (2013, 2014)). Both instrumental payloads were carried by the helicopter BO-105 as two separate external cargos on one
95 long tether: SMART-HELIOS mounted 20 m below the helicopter and ACTOS another 150 m underneath. Typical true air speed of 20 m s^{-1} and high sampling rate of individual instruments provided spatial resolution much higher than for a typical research aircraft.

For complete instrumentation of the helicopter payloads see Tables 1 and 2 in Siebert et al. (2021). In the current study we used the following ACTOS data: three-dimensional wind vector (u_e, v_e, w_e) in the Earth-fixed system and longitudinal-
100 vertical wind components (u, w) in platform-fixed system (derivation explained in sec. 4) provided by the combination of the ultrasonic anemometer-thermometer (Gill Solent HS, Siebert and Teichmann (2000)) and a high-accuracy motion package (inertial navigation system and GPS); virtual temperature T_v derived from the speed of sound measured with the same ultrasonic anemometer-thermometer (Siebert and Muschinski, 2001); temperature T and its small scale fluctuations measured by the Ultra-Fast Thermometer (UFT, Haman et al. (1997); Nowak et al. (2018)) combined with the precise calibrated PT100; specific
105 humidity q_v from the open-path infrared absorption hygrometer (LI-COR LI-7500, Lampert et al. (2018)); liquid water mass fraction q_l determined with the Particle Volume Meter (PVM-100A, Gerber et al. (1994); Wendisch et al. (2002)).

The standard deviations due to uncorrelated noise for sonic measurements are 0.02 m s^{-1} for wind and 0.02 K for virtual temperature (Siebert and Muschinski, 2001). The PT100 was calibrated prior to the campaign in a thermostated water tank using the Greisinger GMH 3750 reference thermometer which provides accuracy better than 0.05 K . The UFT was calibrated
110 for each flight separately against the PT100. For the UFT records, the standard deviation due to uncorrelated noise is 4 mK (Siebert et al., 2003). The hygrometer provides q_v with a noise floor of about 0.005 g kg^{-1} . This instrument was verified to agree well with a few hygrometers of different types and operate satisfactorily on the helicopter-towed system Helipod by Lampert et al. (2018). The PVM-100A measures q_l with the accuracy of 5% and its noise floor was estimated by Siebert et al. (2003) to be about 0.001 g kg^{-1} . The exact sensitivity depends to some extent on droplet size distribution, see Wendisch et al.
115 (2002) for details. For a more general discussion of the instrumentation on the ACTOS platform see Siebert et al. (2006a).

2.3 Data overview

Helicopter flights during ACORES were typically performed over the ocean inside the $10 \text{ by } 10 \text{ km}$ square adjacent to the northern coast of Graciosa. Specific flight path and maneuvers depended on the local cloud situation. Within the flight time of two hours, the usual strategy involved: vertical profile up to roughly 2000 m (a.s.l.), a few 10 km long horizontal legs at
120 selected levels and several steep porpoise dives around stratocumulus top. Two flights were selected for our comparative study: flight #5 on 8 July 2017 and flight #14 on 18 July 2017. The choice was dictated by stratocumulus presence, STBL stratification (considerably well-mixed in flight #5, considerably decoupled in flight #14) and a flight pattern involving substantial sampling time below SC.

Segments of two types were selected from the measurement records: profiles (PROFs) and horizontal legs (LEGs). For convenience, for each flight PROFs are ordered chronologically according to their time of execution and referred to as PROF1-PROF5 while LEGs are ordered according to their mean altitude (m above sea level). The segmentation was done manually so that the influence of sharp turns and pendulum-like motion of the payload is avoided. This resulted in the reduced length of the LEGs, between 3.5 and 12 km. LEGs were flown with a true air speed of $15\text{-}20\text{ m s}^{-1}$ and some minor displacements in vertical are unavoidable for the payload on a 170 m long rope. The mean altitudes and exact lengths are listed in Table 1. PROFs are in fact slanted with an ascent or descent rate of about $3\text{-}5\text{ m s}^{-1}$ which results in an aspect ratio of 0.15-0.25. The horizontal component of motion is necessary to avoid the downwash of the helicopter affecting wind and turbulence measurements on ACTOS. More details about measuring turbulence below a helicopter can be found in Siebert et al. (2006a); Muschinski et al. (2001).

Table 1. Mean altitude and length of the LEGs.

	Height [m]	Length [km]
Flight #5	307	5.44
	553	5.51
	819	7.93
	1079	3.94
	2018	6.25
Flight #14	143	8.11
	287	11.92
	448	7.10
	992	4.79
	2021	3.49

Flight #5 was performed in the afternoon (14:28-16:26 UTC¹) on 8 July 2017. Stratocumulus clouds emerged behind the cold front which had passed the island the day before. The cloud field was moderately thick and quite heterogeneous in structure, with some visible clearings. Satellite image from MODIS on Aqua (Fig. 1) confirms this observation showing dispersed cloud patches in the vicinity of Graciosa. The flight pattern (Fig. 2) involved: deep PROF from minimum flight level (60 m) into the free troposphere (FT), two LEGs in the FT with one close to the stratocumulus top, three LEGs in the STBL with one inside stratocumulus cloud, close to its top. Specific PROFs are indicated in the figure with different line styles which are used hereafter in the vertical profiles of various derived parameters. Altitude ranges corresponding to PROF2-PROF5 of this flight do not overlap, hence we marked them all with dashed lines.

¹On Azores the local time in summer is equivalent to UTC.




Figure 1. Satellite true color image (250 x 250 km) taken on 8 July 2017 at 15:45 UTC (i.e. during flight #5, the time given corresponds to the left swath covering most of the image) by the MODIS instrument on Aqua overpassing Azores, centered on Graciosa airport (blue circle), with overlaid helicopter operation area (red box). The image was acquired from NASA Worldview Snapshots.




Figure 2. ACTOS altitude in flight #5 with marked selected profiles and horizontal legs. PROFs are ordered chronologically, LEGs are ordered according to their altitude. Line styles of the PROFs are consistent with the figures in following sections. Altitude ranges corresponding to PROF2-PROF5 of this flight do not overlap and are all marked with dashed lines. Dots indicate the penetrations of the boundaries of the specific sublayers described in sec. 3.

Flight #14 was performed in the afternoon (15:01-17:04 UTC) on 18 July 2017, shortly after weak precipitation had been noted at the site. The sky was overcast with stratocumulus cloud of homogeneous structure. Many little cumulus clouds, probably at the initial state of formation, were reported over the ocean below stratocumulus deck. However, they were not observed to reach stratocumulus base. MODIS Aqua image (Fig. 3) shows large solid patch of stratocumulus clouds with signatures of closed-cell convection regime. The flight pattern (Fig. 4) involved: four LEGs in the STBL with one inside the stratocumulus cloud, close to its top, one LEG in the FT and a number of PROFs connecting LEG levels. In this figure and hereafter, PROF1-PROF3 are all marked with dashed lines because their altitude ranges do not overlap.




Figure 3. As in Fig. 1 but taken on 18 July 2017 at 14:43 UTC (shortly before flight #14). The image was acquired from NASA Worldview Snapshots.




Figure 4. As in Fig. 2 but for flight #14. Line styles of the PROFs are consistent with the figures in following sections. PROF1-PROF3 are all marked with dashed lines because their altitude ranges do not overlap.

3 Stratification

150 3.1 Derivation of meteorological and stability parameters

Meteorological conditions and stability parameters derived from PROFs are shown in Fig. 5 and Fig. 6 for flight #5 and #14, respectively. Liquid water potential temperature θ_l was calculated following the approximation by Betts (1973):

$$\theta_l = \theta - \frac{\theta}{T} \frac{L_v}{c_p} q_l \quad (1)$$

155 where θ denotes potential temperature, L_v latent heat of vaporization for water and c_p specific heat of dry air at constant pressure. Horizontal wind speed U and direction dd result from appropriate transformation of measured flow velocity (Edson et al., 1998). Because helicopter climb rate was not exactly constant in time and individual instruments differ in sampling rate,

data points were grouped and averaged in 10 m high altitude bins (yet separately for each PROF). To reduce the effect of random eddy penetration and improve clarity, wind profiles were additionally smoothed with five point moving average.

160 Lifting condensation level (LCL) was then derived for each height according to Bolton (1980). Such result is sensitive to gradients of thermodynamic properties in subcloud layer, signaling the degree of boundary layer coupling. To characterize static stability, the Brunt-Vaisala frequency N_b was used:

$$N_b^2 = \frac{g}{\theta_v} \frac{\partial \theta_v}{\partial z} \quad (2)$$

where θ_v is virtual potential temperature derived from speed of sound (provided by ultrasonic anemometer), g gravitational acceleration and z height above sea level. Shear rate Sh quantifies the vertical gradient of horizontal wind:

$$165 \quad Sh^2 = \left(\frac{\partial u_e}{\partial z} \right)^2 + \left(\frac{\partial v_e}{\partial z} \right)^2 \quad (3)$$

where u_e is eastward and v_e northward wind component. The derivatives were evaluated as the tangent of linear least-square fit of 10 m binned variable versus z performed inside symmetric five point windows.

3.2 Quantitative judgement of the degree of coupling

In order to objectively confirm the fact of coupling or decoupling of STBL, we employed several methods from the literature
170 (Jones et al., 2011; Wood and Bretherton, 2004; Yin and Albrecht, 2000).

The first criterion of Jones et al. (2011) involves the differences of θ_l and total water mixing ratio between the uppermost and the lowermost quarters of the boundary layer (instead of the latter quantity, we used our total water mass fraction $q_t = q_l + q_v$ which does not influence the conclusions because those two measures are approximately equal). The sounding is classified as coupled when $\Delta\theta_l = \theta_l^{top} - \theta_l^{bot} < 0.5$ K and $\Delta q_t = q_t^{bot} - q_t^{top} < 0.5$ g kg⁻¹, decoupled otherwise. The second criterion
175 of Jones et al. (2011) involves the difference between the observed cloud base height CB and the LCL corresponding to the conditions at the bottom of the boundary layer. It is classified as coupled when $\Delta z = CB - LCL^{bot} < 150$ m, decoupled otherwise. Here, we used mean conditions of the lowest leg (LEG307 for flight #5, LEG143 for flight #14) to estimate LCL^{bot} and q_l in PROFs to estimate the cloud base height.

Wood and Bretherton (2004) proposed two decoupling parameters:

$$180 \quad \alpha_\theta = \frac{\theta_l^- - \theta_l^0}{\theta_l^+ - \theta_l^0} \quad \alpha_q = \frac{q_t^- - q_t^0}{q_t^+ - q_t^0} \quad (4)$$

where superscripts +, -, 0 denote the values just above the inversion, just below the inversion and in the surface mixed layer, respectively. Wood and Bretherton (2004) calculated α_θ and α_q over subtropical Eastern Pacific at around 0 to 0.4, however no exact critical value for decoupling was determined. The higher those parameters, the more decoupled boundary layer is considered. Here, instead of finding first the SML, we apply mean values in the lower quarter of the boundary layer ($\theta_l^0 = \theta_l^{bot}$
185 and $q_t^0 = q_t^{bot}$).

Yin and Albrecht (2000) introduced a stability parameter to identify transitions in boundary layer soundings:

$$\mu = -\frac{\partial \theta}{\partial p} + \frac{\varepsilon \theta}{1 + \varepsilon r_v} \frac{\partial r_v}{\partial p} \quad (5)$$

where $\varepsilon = R_v/R_d - 1$ depends on the ratio of gas constants for water vapor R_v and dry air R_d , while r_v is water vapor mixing ratio. Their procedure detects transition anytime in the subcloud zone the value of μ exceeds by a factor of 1.3 the average $\bar{\mu}$ between 980 and 900 hPa. Here, instead of using pressure levels, we specify $\bar{\mu}$ as the boundary layer mean.

The above parameters were estimated using PROF1 of flight #5 and PROF5 of flight #14. According to the criteria of Jones et al. (2011), it is evident that flight #5 ($\Delta\theta_l = -0.51^\circ\text{C}$, $\Delta q_t = 0.13 \text{ g kg}^{-1}$, $\Delta z = -72 \text{ m}$) was performed in a coupled STBL while flight #14 ($\Delta\theta_l = 1.19^\circ\text{C}$, $\Delta q_t = 0.90 \text{ g kg}^{-1}$, $\Delta z = 216 \text{ m}$) in a decoupled STBL. Negative values of $\Delta\theta_l$ and Δz suggest instability but it might be also attributed to horizontal inhomogeneities of stratocumulus structure (sec. 2.3) in combination with slanted flight path. Consistently, the parameters of Wood and Bretherton (2004) are smaller for flight #5 ($\alpha_\theta = -0.12$, $\alpha_q = 0.04$) than for flight #14 ($\alpha_\theta = 0.26$, $\alpha_q = 0.26$). The parameter μ is plotted in panel (d) of Figs. 5 and 6. It varies significantly with height and the critical value is occasionally exceeded in both flights. This method was probably optimized for radiosoundings in different climate regime and does not seem robust in case of our data.

Following previous studies looking for differences of cloud top entrainment instability between coupled and decoupled clouds (e.g. Xiao et al., 2011), we calculated the Randall-Deardorff parameter (Randall, 1980; Deardorff, 1980):

$$\kappa = 1 + \frac{c_p}{L_v} \frac{\theta_l^+ - \theta_l^-}{q_t^+ - q_t^-}. \quad (6)$$

In both our cases ($\kappa = 0.71$ for flight #5 and $\kappa = 0.34$ for #14), it exceeds the critical value of about 0.23 indicating the possibility of buoyancy reversal resulting from mixing and evaporative cooling at cloud top.

3.3 Structure of the coupled STBL

The profiles in flight #5 exhibit a well-mixed STBL (Fig. 5). Temperature falls with height with near constant lapse rate Γ_T inside the boundary layer, followed by a sharp inversion at the top. Liquid water potential temperature is almost constant from close to the surface up to the stratocumulus top, where it features an increase of $\sim 5 \text{ K}$. Total water mass fraction behaves analogously, with a decrease of $\sim 7 \text{ g kg}^{-1}$ above cloud top. Interestingly, very dry air is located at the top of the temperature inversion. It is further capped by a layer of considerably higher q_v , however much lower than inside the boundary layer. Liquid water mass fraction in the cloud is moderate and suggest non-trivial cloud structure. There were cloud clearings penetrated as ACTOS moved along the slanted path, visible in the high rate records of q_l (not shown here). Wind velocity fluctuates in the boundary layer within $\pm 1 \text{ m s}^{-1}$ around the mean $\sim 5 \text{ m s}^{-1}$. Wind shear across the cloud top and the inversion can be noticed. Wind direction is from the NNW throughout the sampled height.

Significant differences can be observed between the PROFs in wind speed and the position of inversion. Subsequent PROFs were not performed at the same time and location, so certain variability is expected. Airborne sampling features inevitable randomness due to probing specific structures (eddies, updrafts, cloud holes etc.), thus slanted profiles do not represent mean conditions accurately.

LCL stays roughly equal from the lowest level up to the cloud base. Interestingly, it is slightly higher than the actual cloud base which might be again related to horizontal inhomogeneities in cloud structure. Brunt-Vaisala frequency indicates weak




Figure 5. Vertical structure of the coupled STBL (flight #5): (a) temperature T and liquid water potential temperature θ_l , (b) liquid water mass fraction q_l and specific humidity q_v , (c) wind speed U and direction dd , (d) lifting condensation level LCL and stability parameter μ of Yin and Albrecht (2000) with its critical level for the detection of transitions (dashed black line), (e) squared Brunt-Vaisala frequency Nb^2 and wind shear rate Sh^2 . Line styles correspond to specific profiles – consistently with Fig. 2. Color shadings denote the sublayers: entrainment interface layer (red) and stratocumulus layer (blue).

static instability in the boundary layer, stronger inside the cloud than below, and very strong stability at the capping inversion. Wind shear is more variable which can be attributed to sampling various eddies.

Based on θ_l , q_l and q_v , we manually distinguished the following sublayers: the entrainment interface layer (EIL) including the temperature inversion and the very top of the cloud, the stratocumulus layer (SCL) containing the cloud, the subcloud layer (SBL) ranging from cloud base down to the surface, and the sublayer representing free tropospheric conditions (FTL, not necessarily adjacent to the EIL top). For reference, the EIL and SCL are marked with red and blue shading in Fig. 5 and following. The heights and average properties inside the sublayers are listed in Table A1 in the appendix. The deepest profile (PROF1, solid line), was used for sublayer distinction because the specific heights may vary between the PROFs. The individual penetrations of the sublayer boundaries during other segments (PROFs and dolphin porpoises) are indicated in Fig. 2.

3.4 Structure of the decoupled STBL




Figure 6. As in Fig. 5 but for the decoupled STBL (flight #14). Line styles are consistent with Fig. 4. Color shadings denote the sublayers: entrainment interface layer (red), stratocumulus layer (blue), subcloud layer (green), transition layer (purple).

230 The profiles in flight #14 exhibit decoupled STBL (Fig. 6). Liquid water potential temperature gradually rises with height whereas specific humidity decreases step-wise. Despite the distinct q_v gradient in BL middle, its value in the lowest part and in the subcloud section is relatively stable. This suggests that both the upper and the lower STBL portion are internally mixed. The FT is quite humid, with values of q_v larger than for flight #5. The difference in θ_t at stratocumulus top is ~ 5 K while in q_t only ~ 3 g kg $^{-1}$. The stratocumulus is thicker and more abundant in liquid water than in the previous case. Wind velocity
235 varies ± 1 m s $^{-1}$ around the mean ~ 6 m s $^{-1}$. Wind direction is predominantly NW. There is significant wind shear across the inversion, with difference in U reaching ~ 4 m s $^{-1}$. LCL replicates the gradients of q_v in the middle of the boundary layer. It corresponds to the cloud base height only in the section right below the cloud which is a signature of decoupling. Brunt-Vaisala frequency indicates weak static stability throughout most of the profile, including the cloud. Its peak in the inversion layer coincides well with the maximum of Sh^2 .

240 Similarly to flight #5, we distinguished the sublayers: the FTL, the EIL, the SCL, and the SBL extending from cloud base down to the level where LCL is no longer in agreement with the observed cloud base height. In addition, two more sublayers typical for decoupled conditions were introduced: the transition layer (TSL) containing the major gradients in specific humidity and wind speed, and the surface mixed layer (SML) extending from the surface up to the bend in θ_l profile (where it begins to rise with height, c.f. Fig. 6a). Somewhat arbitrary boundary of 385 m was chosen to represent the section directly influenced by
 245 surface processes. For reference, the EIL, SCL, SBL and TSL are marked with red, blue, green and purple shading, respectively, in Fig. 6 and following. The heights and average properties inside the sublayers are listed in Table A2. PROF5 was used for sublayer identification because it covers most of the STBL depth. The individual penetrations of the sublayer boundaries during other segments are indicated in Fig. 4.

4 Methods

250 Parameters of turbulence were derived using high-resolution measurements of wind velocity, temperature and humidity. Depending on the quantity, the results were obtained for PROFs or LEGs specified in sec. 2.3. In case of PROFs, our procedure resembles the approach of Tjernstrom (1993). After timeseries of a parameter had been computed, appropriate segments were extracted and data were averaged in 10 m altitude bins (as in sec. 3.1). For LEGs, full segment was used to calculate a desired parameter. Next, each LEG was divided into 7 subsegments of equal length, overlapping by half of the length, and the
 255 very same method was applied to calculate respective quantity in each subsegment. Standard deviation among subsegments is regarded as parameter variability and shown with errorbars in plots.

The lateral channel of the ultrasonic anemometer was affected by a substantial level of artificial fluctuations (up to 1 m s^{-1} in amplitude) due to instrumental issues. The origin of this problem is under investigation. It seems to appear for true air speed above about 12 m s^{-1} which makes it relevant for most of the flight time. Therefore, we applied simplified geometrical
 260 transformation to the measured velocity vector, so that high resolution retrieval of wind velocity is possible. In comparison with the standard transformation (Lenschow, 1986), we included pitch rotation but neglected roll and yaw rotations to prevent the lateral channel from coupling with the others. The resulting vector (u, v, w) can be interpreted as wind velocity in horizontal longitudinal, horizontal lateral and vertical direction, respectively, as long as the platform is not tilted left or right (roll angle is small). This condition was satisfied throughout most of the flight time, except for major turns. For calculating turbulence
 265 properties, we selected segments with the roll angle $< 0.1 \text{ rad}$. The lateral wind v cannot be used for turbulence analysis but longitudinal u and vertical w are free from the disturbances. The modification described is not necessary to obtain mean wind profiles (U, \overline{dd}) because averaging and smoothing is applied anyway (see sec. 3.1).

Reynolds decomposition of the signals (c.f Stull, 1988)

$$x(t) = X(t) + x'(t) \tag{7}$$

270 into large scale slowly varying $X(t)$ and small scale fluctuations $x'(t)$ was realized with a simple symmetric running mean. Fluctuations $x'(t)$ were obtained by subtracting that mean from the original signal. Unless specified otherwise, the chosen

window was 50 s which corresponds to the distance of ~ 1 km. Such length is enough to penetrate at least a few large turbulent eddies typical for the atmospheric boundary layer (Malinowski et al., 2013).

4.1 Turbulence Kinetic Energy and variances

275 Variances of turbulent fluctuations $\langle u'^2 \rangle$, $\langle w'^2 \rangle$, $\langle T'^2 \rangle$, $\langle q_v'^2 \rangle$ and third moment of vertical velocity fluctuations $\langle w'^3 \rangle$ were obtained by taking the average along LEG, denoted as $\langle \rangle$. Because lateral wind fluctuations were not available, we assumed horizontal isotropy to approximate missing $\langle v'^2 \rangle$ with $\langle u'^2 \rangle$ in turbulence kinetic energy (TKE) calculation:

$$\text{TKE} = \langle u'^2 \rangle + \frac{1}{2} \langle w'^2 \rangle. \quad (8)$$

It is worth remembering that variances and TKE usually represent mostly large scales because larger eddies in turbulence
280 cascade are more energetic than smaller ones.

The accuracy of the results is severely limited by the length of the LEGs. Based on the methods of Lenschow et al. (1994), in the boundary layer the variances are subject to the systematic sampling error of about 5 % and the random sampling error of about 20 %. In the case of $\langle w'^3 \rangle$, those errors are accordingly larger (order of 10 % and 100 %, respectively, unless $\langle w'^3 \rangle$ is not very close to zero). Importantly, in the plots we provide the variability among subsegments which was found to be of the
285 same order as the total sampling error, in most cases larger than it.

4.2 TKE production and heat fluxes

Turbulence kinetic energy can be generated by buoyancy and wind shear (ignoring advection and turbulent transport). We estimated two respective terms of the TKE budget equation (Stull, 1988), buoyancy production/consumption B and shear production S , employing eddy correlation:

$$290 \quad B = \frac{g}{\langle \theta_v \rangle} \langle w' \theta_v' \rangle, \quad S = -\langle w' u' \rangle \frac{\partial u}{\partial z}. \quad (9)$$

Here, we could provide only longitudinal component of shear production because lateral wind fluctuations were not available. Correlations were computed along the LEGs. Derivatives were estimated from the PROFs covering the relevant altitude range. Inevitably, such an approach introduces some inaccuracy as the exact place and time of derivative estimation is different than for the correlation estimation. To quantify vertical transport of heat and moisture, we estimated sensible and latent heat fluxes
295 according to:

$$Q_s = \rho c_p \langle w' \theta' \rangle, \quad Q_l = \rho L_v \langle w' q_v' \rangle \quad (10)$$

where ρ is air density.

The range of scales represented in the correlations is limited by the lowest spatial resolution of the two multiplied signals. The anemometer (u , w , θ_v) resolves scales down to ~ 0.5 m (where this limit stems from the path length and spectral transfer
300 properties (Kaimal et al., 1968)), the thermometer (θ) down to ~ 2 cm, the hygrometer (q_v) down to ~ 1 m. As a result, $\langle w' \theta' \rangle$, $\langle w' u' \rangle$ and $\langle w' \theta' \rangle$ are resolved down to ~ 0.5 m while $\langle w' q_v' \rangle$ down to ~ 1 m. Those three instruments work satisfactorily

also inside clouds of moderate liquid water and droplet concentration, as our stratocumulus (Cruette et al., 2000; Siebert and Teichmann, 2000). In comparison with some other studies, the buoyancy estimation in the cloud does not include the contributions of liquid water flux $\langle w'q'_l \rangle$ and droplet sedimentation which are expected to be relatively small (considering moderate q_l) and of opposite sign, therefore partly compensate.

Similarly to variances, the accuracy of the fluxes obtained with the method of eddy correlation is limited by the length of the LEGs. In the boundary layer, the systematic sampling error was estimated for about 5-10 % while the random sampling error for about 50 % (Lenschow et al., 1994), unless the flux does not vanish. The subsegment variability (marked with errorbars in the plots) is in most cases larger than the total sampling error.

Additionally, B , Q_s and q_l at the surface were estimated with the Coupled Ocean–Atmosphere Response Experiment bulk algorithm in version 3.0 (COARE 3.0) described in Fairall et al. (2003). Sea surface temperature was taken from satellite multi-mission product provided by the Group for High Resolution Sea Surface Temperature (JPL MUR MEaSUREs Project, 2015) while all the other inputs were our measurements from the lowest point of the PROFs.

4.3 TKE dissipation rate

TKE dissipation rate ϵ was calculated invoking common assumption of homogeneous, isotropic, stationary turbulence which leads to the specific form of power spectra and structure functions (Kolmogorov, 1941). Nevertheless, theoretical assumptions are often hardly satisfied in the atmosphere, e.g. considering complex stratification, and therefore ϵ estimation from moderate-resolution (not directly resolving dissipative scales) measurements is challenging (Siebert et al., 2006b; Jen-La Plante et al., 2016; Waclawczyk et al., 2017, 2020). To account for possible anisotropy, ϵ was derived separately for longitudinal and vertical velocity fluctuations, following the methods of Siebert et al. (2006b). We also characterized the quality of estimations with additional parameters describing the deviation of experimental data from theoretical dependencies.

4.3.1 Structure function method

Second order structure function (SFC) was calculated for measured u' and w' according to the same equation:

$$D_u(r) = \langle |u'(x+r) - u'(x)|^2 \rangle \quad (11)$$

where r is distance between data points (given by true air speed) and the average is taken over positions x along the flight path. SFC was then resampled, i.e. averaged inside logarithmically equidistant bins covering the assumed inertial range $r \in [0.4, 40]$ m, with eight bins per decade (see Fig. 7). The resampling was applied in order to account for the density of data points increasing with scale in logarithmic coordinates.

Theory predicts that in the inertial range SFC has the form (Pope, 2000):

$$D(r) = C(\epsilon r)^{\frac{2}{3}} \quad (12)$$

where C is a constant, experimentally determined to $C_u \approx 2.0$ for longitudinal and $C_w \approx 2.6$ for vertical velocity component. We calculated ϵ^{sfc} by least squares fit of this relationship to the resampled SFC. Second fit was performed according to:

$$D(r) = C^* r^s \quad (13)$$

with two fitted parameters: prefactor C^* and exponent s corresponding to the slope in log-log plot. The exponent is used as a benchmark of the agreement of the SFC form with theory. Additionally, Pearson correlation coefficient R^{sfc} was computed for the resampled points. It quantifies the linearity of the experimental SFC in log-log coordinates. Consequently, s and R^{sfc} assess to some extent the reliability of derived ϵ .




Figure 7. Example of ϵ derivation with structure function method (flight #5, LEG307, vertical component). Computed SFC (Eq. (11), blue) is resampled in the assumed inertial range (yellow) to obtain logarithmically spaced points (triangles) which are used for least squares fits: one with free slope (Eq. (13), purple), one with fixed theoretical slope (Eq. (12), green).

4.3.2 Power spectrum method

Power spectral density (PSD) of u' and w' was calculated with the Welch algorithm. The window was chosen as half the length of the segment. The windows overlap by half of their length, so in turn there are three individual PSDs averaged in the Welch scheme. PSD was resampled in the assumed inertial range, analogously to SFC (see Fig. 8).

Theory predicts the following PSD form in the inertial range (Pope, 2000):

$$P(f) = C' \left(\frac{U_s}{2\pi} \right)^{\frac{2}{3}} \epsilon^{\frac{2}{3}} f^{-\frac{5}{3}} \quad (14)$$

where f is frequency and C' is a constant ($C'_u \approx 0.49$ for longitudinal and $C'_w \approx 0.65$ for vertical component). We derived ϵ^{psd} by fitting this relationship to the resampled PSD. Second fit was performed according to:

$$P(f) = C^* f^p \quad (15)$$

where fitted PSD exponent p corresponds to the slope in log-log plot. Together with Pearson correlation coefficient for the resampled points R^{psd} it measures the agreement of PSD form with theory and reliability of derived ϵ .




Figure 8. Example of ϵ derivation with power spectrum method (flight #5, LEG307, vertical component). Computed PSD (blue) is resampled in the assumed inertial range (yellow) to obtain logarithmically spaced points (triangles) which are used for least squares fits: one with free slope (Eq. (15), purple), one with fixed theoretical slope (Eq. (14), green).

4.3.3 Application of the methods

350 For PROFs, the moving window of 2 s was applied to the timeseries u' and w' . In each window, ϵ was derived separately with the two methods, together with s , R^{sfc} , p , R^{psd} . Such a solution was verified to provide sufficiently good fits and constitutes the compromise between high final spatial resolution (short window desired) and adequate representation of SFC or PSD (long window desired). Our approach follows earlier studies which determined the instantaneous dissipation rate utilizing the same type of data as ours (Siebert et al., 2006b; Katzwinkel et al., 2012). Siebert et al. (2006b) have chosen the window of 1 s based
355 on their sensitivity tests and the arguments provided by Frehlich et al. (2004) and Muschinski et al. (2004). Because we derive not only ϵ but also the slopes and correlations, we increased the window to 2 s so that the linear fit covers considerable portion of the inertial range and the sufficient number of logarithmically equidistant resampled points (see sec. 4.3.1).

In the case of LEGs, both methods were applied to the whole segment. Then, SFC and PSD were in practice averaged over relatively long horizontal distance. This approach provides an estimate of mean dissipation in contrast to local values
360 computed in short windows which might differ from the mean (Kolmogorov, 1962). Also, SFC and PSD derived on long horizontal segment are expected to follow the theoretical form more accurately which is indeed the case.

Our results (sec. 5.3) demonstrate a good agreement between the methods as long as relative variations with height are concerned. In terms of absolute values, ϵ^{psd} is usually higher than ϵ^{sfc} (around the factor of 2). In general, derived SFC

resembles its theoretical form better than PSD which is indicated by the fitted exponents and correlation coefficients. This
 365 agrees with Siebert et al. (2006b) who found the SFC method to be more robust for ϵ estimation from airborne platforms.

In order to estimate the uncertainties of the results, we used the random errors of the fitted parameters (computed with a
 standard method from least-squares fit residuals). The random error of ‘instantaneous’ (calculated in 2 s windows and serving
 for the derivation of the profiles) dissipation rate equals ~ 50 % in the boundary layer and ~ 150 % in the FT. The error of the
 LEG-derived ϵ is ~ 30 % for longitudinal component and ~ 15 % for vertical component in the boundary layer while ~ 150 %
 370 for both components in the FT. The random error of the fitted slopes is ~ 0.04 for s and ~ 0.16 for p corresponding to the
 ‘instantaneous’ estimations while ~ 0.02 in the case of both LEG-derived slopes. Notwithstanding, the given values represent
 the uncertainties due to the random errors of the fit only. The reliability of the derived dissipation rates can be also assessed by
 comparing the results of the two derivation methods, by comparing the fitted SFC and PSD slopes with their theoretical values
 or using the deviation of the computed correlation coefficients from unity.

375 4.4 Anisotropy

The assumption of isotropy might be violated in many specific situations in the atmospheric boundary layer, e.g. under strong
 buoyancy and wind shear at stratocumulus top (Malinowski et al., 2013; Jen-La Plante et al., 2016; Akinlabi et al., 2019). To
 investigate deviations from isotropy, we use anisotropy ratios A of two types, bulk and spectral, relating w -derived parameters
 to u -derived ones.

380 We define the following bulk anisotropy ratios:

$$A_2^{var} = \sqrt{\frac{\langle w'^2 \rangle}{\langle u'^2 \rangle}}, \quad A_\epsilon^{sfc} = \frac{\epsilon_w^{sfc}}{\epsilon_u^{sfc}}, \quad A_\epsilon^{psd} = \frac{\epsilon_w^{psd}}{\epsilon_u^{psd}}. \quad (16)$$

The first relates mostly to larger eddies which have dominant contribution to total variance. Isotropy is indicated by the values
 close to 1, while $A_2^{var} < 1$ and $A_2^{var} > 1$ indicate anisotropic turbulence dominated by horizontal and vertical fluctuations,
 respectively. On the other hand, A_ϵ^{sfc} and A_ϵ^{psd} regard mostly the inertial range eddies because ϵ derivation exploits SFC or
 385 PSD scaling in the inertial range. Analogously, values close to unity indicate isotropy.

The spectral anisotropy is the scale-dependent ratio of PSDs for vertical and longitudinal velocity:

$$A_P(r) = \frac{P_w(U_s/r)}{P_u(U_s/r)} \quad (17)$$

where true air speed is utilized to convert frequency into distance. A similar approach was exercised by Pedersen et al. (2018)
 who compared modeled and measured anisotropy in the region of stratocumulus top. In the inertial range, Kolmogorov theory
 390 predicts $A_P = 4/3$. Such a value of the experimentally derived $A_P(r)$ should then indicate isotropy at the particular scale r , as
 in the analysis of Siebert and Muschinski (2001). We applied the same resampling procedure as in sec. 4.3.2 to the LEG-derived
 PSDs but across the whole available range of scales (not only the inertial) and the ratio was then calculated point-by-point.

4.5 Length scales

Turbulence energy cascade is often characterized by several length scales: integral scale L , Taylor microscale λ and Kolmogorov scale η . The integral length scale corresponds to the energy-containing eddies which are involved in TKE generation. In the energy cascade, it marks the beginning of the inertial subrange where turbulent flow is considerably isotropic despite the anisotropy of large scale factors. The indefinite integral of the autocorrelation function involved in the formal definition of L cannot be evaluated experimentally due to the limited length available. We estimated the distance where the autocorrelation

$$\rho_u(r) = \frac{\langle u'(x+r)u'(x) \rangle}{\langle u'^2 \rangle} \quad (18)$$

declines by a factor of e . This method is robust enough to provide reasonable results in all our cases. The very same procedure was applied to longitudinal as well as vertical velocity to provide L_u and L_w , respectively. According to Pope (2000), under isotropic conditions $L_w = \frac{1}{2}L_u$. Such a proportion can then indicate isotropy in the relevant large eddy scale.

At the Taylor microscale, viscosity starts to substantially affect the dynamics of turbulent eddies. Under the assumption of isotropy, it can be related to velocity variance and dissipation rate. We estimated two Taylor scales, longitudinal and vertical:

$$\lambda_u = \sqrt{30\nu \frac{\langle u'^2 \rangle}{\epsilon_u^{sfc}}}, \quad \lambda_w = \sqrt{15\nu \frac{\langle w'^2 \rangle}{\epsilon_w^{sfc}}} \quad (19)$$

where ν is air viscosity for which we accounted for temperature and pressure dependence (Sutherland, 1893). In homogeneous isotropic turbulence $\lambda_w = \frac{1}{\sqrt{2}}\lambda_u$ (Pope, 2000).

Kolmogorov scale corresponds to the smallest eddies where TKE is dissipated into heat by viscosity. Following dimensional arguments of the famous similarity hypothesis, it equals:

$$\eta_u = \left(\frac{\nu^3}{\epsilon_u^{sfc}} \right)^{\frac{1}{4}}. \quad (20)$$

It was calculated separately for longitudinal (η_u) and vertical (η_w) direction with the same formula. Provided local small-scale isotropy, they should be equal. For convenience, in λ and η derivation, we used only ϵ^{sfc} and neglected ϵ^{psd} because SFC proved to resemble its theoretical form better (see sec. 5.3).

5 Observed turbulence properties

Turbulence properties in coupled and decoupled STBL are documented in a series of plots. Mean PROF-derived values inside the sublayers are listed in Tables A1 and A2.

5.1 Turbulence Kinetic Energy and variances

Figs. 9 and 10 present variances of vertical and longitudinal velocity fluctuations, TKEs, third moments of vertical velocity, variances of temperature and specific humidity in the LEGs of flight #5 and #14, respectively. Generally, the TKE inside the coupled STBL decreases with height from the middle of the SBL up to the cloud top. Despite slightly unstable stratification,




Figure 9. Statistics of turbulent fluctuations in the coupled STBL (flight #5): (a) variance of horizontal $\langle u'^2 \rangle$ and vertical velocity $\langle w'^2 \rangle$, (b) turbulence kinetic energy TKE and third moment of vertical velocity $\langle w'^3 \rangle$, (c) variance of temperature $\langle T'^2 \rangle$ and specific humidity $\langle q_v'^2 \rangle$.




Figure 10. As in Fig. 9 but for the decoupled STBL (flight #14).

the contribution from horizontal velocity variance is dominant over the vertical one. The latter reaches a minimum value below the cloud, where the buoyancy production is close to zero (compare Fig.11 in the next section).

Estimated values of the TKE are also large in the FT above the temperature inversion. This is rather an artifact due to the presence of gravity waves favored under stable conditions (the power spectra of w , u , q_v , θ_v and the cospectra of wu , wq_v and $w\theta_v$ indicate the dominant contribution of the wavelength of about 450 m). Recall that LEG1079 was flown very close to the EIL and the cloud top which often features undulated interface.

The third moment of vertical velocity is positive in the lowest LEG307, suggesting strong but localized updrafts and weak but widespread downdrafts. Higher up, it is close to zero. This results ought to be interpreted with caution because the estimation of $\langle w'^3 \rangle$ can be subject to errors due to insufficient statistics related to the small chance of penetrating infrequent but intense events (Lenschow et al., 1994; Kopec et al., 2016).

Fluctuations of temperature and humidity can be significant wherever there are spatial gradients of those quantities or in the presence of sources or sinks of heat and moisture. Such conditions occur close to the cloud top, where radiative cooling is the sink of heat and mixing between the air volumes of considerably contrasting properties occurs. Indeed, measured variances are highest in the cloud segment and decrease downward into the boundary layer where T and q_v are locally more uniform.

In the decoupled STBL, TKE level is in general lower than in the coupled case. The profiles of velocity variances across the SML resemble a typical mixed layer with shear, i.e. high TKE at the bottom and the top which is realized mostly by the contribution of horizontal velocity variance (e.g., Stull, 1988, ch. 4). The prevalence of horizontal in comparison to vertical is particularly visible for LEG448, close to the transition, where the vertical velocity variance reaches its minimum. Similarly to TKE, humidity and temperature variances exhibit a maximum at this level. T and q_v can be considered passive scalars which undergo mixing. The increased variances are caused by gradient production (Term IV in the variance budget equations in the formulation of Stull (1988), his Eqs. (4.3.2) and (4.3.3)) rather than by any diabatic sources.

Skewness of vertical velocity is slightly positive in the SML with the maximum in LEG287. At the transition and in the cloud it is close to zero with tendency towards negative values. This suggests dominant role of updrafts in the SML and downdrafts in the SCL. Altogether, the results can be interpreted as a signature of decoupling between the circulations in lower and upper parts of the boundary layer, as downdrafts originated at cloud top and updrafts originated at the surface slow down and diverge horizontally at the transition level.

5.2 TKE production and turbulent fluxes

Buoyant production of TKE is expected to be significant inside the cloud and close to the surface while the shear production is expected to be significant at the bottom and at the top of the boundary layer (Markowski and Richardson, 2010). Such a picture is in general agreement with our results for flight #5. In the coupled STBL observed there (Fig. 11), B is maximum in the LEG flown inside the cloud ($8.0 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-3}$), drops to nearly zero below the cloud and increases towards the surface, reaching $5.6 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-3}$ (estimated with COARE algorithm). S is more uniform in the boundary layer, yet subject to substantial variability among subsegments.

Sensible heat flux reaches maximum of almost 40 W m^{-2} close to the cloud top, stays small and positive in the middle of the boundary layer with the surface value of around $Q_s = 11 \text{ W m}^{-2}$ (according to COARE parameterization). Latent heat flux follows near linear decrease from $Q_l = 130 \text{ W m}^{-2}$ at the ocean surface, which is the source of moisture due to evaporation,




Figure 11. (a) TKE production by buoyancy B and shear S , (b) sensible Q_s and latent Q_l heat fluxes in the coupled STBL (flight #5). The lowest dot denotes the parameterized surface value obtained with COARE 3.0 algorithm.




Figure 12. As in Fig. 11 but in the decoupled STBL (flight #14).

to roughly zero below the cloud. At low levels in the atmosphere (at the surface and in LEG307) the contribution of moisture transport to buoyancy is of the same order as the contribution of heat transport (not shown). In the cloud top region Q_l exceeds

100 W m⁻² (subject to very large variability). It is not clear what are the contributions of radiative and evaporative cooling
 460 towards the observed heat fluxes there. LEG819 was performed close to the cloud top but neither exactly at the interface nor
 inside the EIL. Although cloud top entrainment instability parameter κ significantly exceeds the critical value (see sec. 3.2),
 which suggests the importance of evaporation, radiative cooling might still be dominant as in the study of Gerber et al. (2016).

In the decoupled STBL observed in flight #14 (Fig. 12), production terms are of the same order as in the coupled case.
 The COARE algorithm provides $B = 4.0 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-3}$, $Q_s = 6.7 \text{ W m}^{-2}$, $Q_l = 107 \text{ W m}^{-2}$ at the surface. B decreases with
 465 height turning into weak buoyancy consumption at the transition. This can be considered an important signature of decoupling.
 Above, in the cloud, B is again positive, yet significantly smaller ($2.6 \cdot 10^{-4} \text{ m}^2 \text{ s}^{-3}$) than at similar location in the coupled
 STBL. Shear production is present in the SML and at the transition as well as in the cloud top region.

Sensible heat flux in the decoupled boundary layer is relatively small reaching maximum of $\sim 10 \text{ W m}^{-2}$ at $\sim 140 \text{ m}$. Latent
 heat flux features near linear decrease with height from the maximum of $\sim 100 \text{ W m}^{-2}$ at the surface to roughly zero at the
 470 transition. In the lower part of the STBL (at the surface, in LEG143 and LEG287) the contribution of moisture transport to
 buoyancy is of the same order as the contribution of heat transport (not shown). Both sensible and latent heat fluxes observed in
 the cloud (LEG992) are small, in contrast to the coupled case. Together with rather moderate B in the cloud this suggests that
 the drivers of convection, i.e. radiative and evaporative cooling, are not as efficient in this situation which might have been one
 of the reasons why decoupling occurred. The cloud top entrainment instability parameter κ (sec. 3.2) is indeed smaller in the
 475 decoupled cloud in comparison to the coupled one which implies less efficient evaporative cooling. However, the comparison
 of radiative cooling effects between the cases requires further investigation. Another observation is that the moisture delivery
 from the ocean surface to the cloud ought to be more difficult in the decoupled STBL as Q_l vanishes at much lower height in
 relation to the cloud base than in the coupled case.

5.3 TKE dissipation rate

480 Measurements in the coupled STBL during flight #5 (Fig. 13) indicate relatively small variability of TKE dissipation rate
 throughout the boundary layer depth and substantial decrease right above the cloud top. The values fluctuate by roughly
 1 order of magnitude, between $10^{-4} \text{ m}^2 \text{ s}^{-3}$ and $10^{-3} \text{ m}^2 \text{ s}^{-3}$. Importantly, those variations do not correlate between the
 PROFs, hence they are the manifestation of some intermittency and random effects involved in airborne sampling rather than
 any systematic stratification. Among the LEGs, the highest dissipation rate was observed in the one close to the cloud top,
 485 where also substantial buoyant production of TKE was revealed (see sec. 5.2). On the other hand, continuous profiles of ϵ
 derived from PROFs do not show significant difference between the cloud and the subcloud part. It suggests that even though
 the TKE might be produced at specific places it is probably redistributed well by the circulation across the STBL before being
 dissipated by viscosity (c.f. transport analysis by Kopec et al. (2016)).

Inside the STBL, the exponents of structure function s (sec. 4.3.1) and of power spectra p (sec. 4.3.2) are close to their
 490 theoretical values ($2/3$ and $-5/3$, respectively), in striking contrast to the FT. Individual deviations occasionally reach 40 %
 in the STBL. On average, the deviations are a bit smaller inside the SCL than in the SBL (see Table A1). Typically, SFCs
 and PSDs seem to be flatter than the theory predicts (absolute values of s and p smaller than theoretical). Such behavior




Figure 13. TKE dissipation rate and inertial range scaling in the coupled STBL (flight #5): (a), (b) dissipation rate ϵ , (c), (d) fitted exponents s and p , (e), (f) correlation coefficient R . Superscripts *sfc* and *psd* denote the structure function and power spectrum methods, respectively. Subscripts *u* and *w* denote horizontal and vertical velocity components, respectively. Dissipation rates for LEG1079 which are not visible in panels (a), (b) are smaller than $10^{-5} \text{ m}^2 \text{ s}^{-3}$.

might be attributed to the non-homogeneity and non-stationarity of turbulence and different stages of its development, e.g. decay (Vassilicos, 2015). When different velocity components are concerned, SFCs and PSDs of vertical fluctuations follow
 495 Kolmogorov theory closer than the longitudinal, signaling some anisotropy in turbulence energy cascade.

Correlation coefficients R^{sfc} and R^{psd} (sec. 4.3) are close to unity in the coupled STBL. This implies both the SFC and the PSD can be considered linear in log-log coordinates in the assumed inertial range of scales. The correlation is higher for LEGs than for PROFs due to better averaging. It sharply decreases across the EIL, suggesting that in the FT the assumptions involved in the derivation of ϵ are not satisfied. Therefore, ϵ estimates above the boundary layer cannot be considered credible
 500 (Akinlabi et al., 2019). On the other hand, inside the STBL the observed forms of SFC and PSD are reasonably consistent with theoretical predictions.

Measurements in the decoupled STBL during flight #14 (Fig. 14) present lower values of ϵ and more variability with respect to height. PROF-derived results averaged across the sublayers increase from the SML up to the SCL (see Table A2). Such a trend is consistent for all derivation methods and velocity components, despite differences in the absolute values among them.
 505 The LEG-derived ϵ decreases with height, from the surface up to the transition.

Vertical profiles of the fitted exponents s and p reveal internal layering of the STBL. In contrast to the coupled case, all PROF-derived exponents deviate significantly from theoretical values. The deviations are appreciably smaller in the SML than in the SBL and the SCL, clearly demonstrating that turbulence in the upper part of decoupled STBL is further from Kolmogorov's




Figure 14. As in Fig. 13 but for the decoupled STBL (flight #14).

concepts than in the lower part. The parameters inside the SCL and the SBL are comparable, suggesting there is an efficient
 510 circulation and mixing across them. Those facts were expected, taking into account our analysis of stratification (sec. 3.4) and
 TKE production (sec. 5.2). Most probably, turbulence generated in the cloud top region is redistributed by the large eddies and
 the transport terms of the TKE balance equation (Stull, 1988) across the SCL and the SBL. Though, the properties of such
 turbulence are remarkably far from the Kolmogorov theory assuming homogeneity, isotropy and stationarity. In the light of
 this observation, the dissipation rates obtained with the methods based on the theoretical inertial range scalings can become
 515 questionable. The assumptions are better resembled by the conditions in the lowermost part of the atmosphere, albeit they are
 still distant from being exactly fulfilled. The profiles of R^{sfc} and R^{psd} are in agreement with the above hypothesis suggesting
 different character and origin of turbulence in the upper and lower part of the STBL. The absolute values are smaller than in
 the coupled case. In the SBL and the SCL the correlation is even quite poor at some particular heights.

In contrast to the PROFs, the LEG-derived exponents stay mostly close to $2/3$ or $-5/3$, accordingly, while the correlations
 520 are close to one. We suppose that the observed discrepancy results from the combination of horizontal inhomogeneity and in-
 termittency of turbulence. PROF-derived and LEG-derived parameters should not be directly compared because they represent
 small and large fluid volumes, respectively. Unfortunately, none of the horizontal segments was performed in the SBL.

5.4 Anisotropy

The coupled STBL sampled in flight #5 features bulk anisotropy ratios predominantly in the range between 0.5 and 1.0
 525 (Fig. 15). The variance anisotropy ratio is the largest (0.9) for the horizontal segment inside the cloud, close to its top where




Figure 15. Anisotropy ratios in the coupled STBL (flight #5).




Figure 16. Anisotropy ratios in the decoupled STBL (flight #14).

the turbulence is efficiently generated by buoyancy (sec. 5.2). In the SBL the values are a bit smaller. Despite substantial local fluctuations observed in A_ϵ^{sfc} and A_ϵ^{psd} , their average level can be considered constant across the boundary layer. There is very

little difference between the SBL and the SCL. The SFC-derived anisotropy ratio is relatively close to unity, suggesting near isotropic conditions. However, the PSD-derived ratio, typically around 0.6, seems to indicate the dominant role of horizontal fluctuations. The reason for such a discrepancy between the methods is not clear. It can be related to the bias in the estimation of dissipation rates between them (c.f. Waławczyk et al., 2020). Nevertheless, both anisotropy measures indicate no internal layering inside the STBL. In the FT, under static stability and weak turbulence production, horizontal motions dominate.

In the decoupled STBL investigated in flight #14, bulk anisotropy ratios are on average smaller than in the previous case (Fig. 16), signaling prevalence of horizontal fluctuations over vertical ones. A_2^{var} is the largest in the surface layer (reaching 0.72), smaller in the cloud (0.54) and close to the transition (0.41) between the two circulation systems, cloud-driven and surface-driven. Dissipation-derived anisotropy ratios imply the separation of the STBL into two parts with the border in the TSL. In the upper part, covering the SCL and the SBL, A_ϵ^{sfc} and A_ϵ^{psd} are visibly smaller than in the SML. Again, the PSD-derived rate is systematically lower than the SFC-derived, but the discrepancy is not as pronounced as in the case of flight #5. Importantly, the change at ~ 500 m correlates well with the change in the fitted SFC and PSD exponents (see sec. 5.3) as well as with the gradient of specific humidity (see sec. 3.4). This fact confirms the hypothesis involving two major circulation circuits dividing the STBL into two parts which are internally relatively well-mixed but feature turbulence of different character. In the SML, turbulence seems to be more vigorous and isotropic than in the SCL and the SBL.




Figure 17. Spectral anisotropy ratio in the coupled STBL (flight #5). The horizontal dashed line denotes the 4/3 level expected for isotropy in the inertial range.

Spectral anisotropy ratios in the coupled STBL presented in Fig. 17 are of similar form for all three LEGs inside the boundary layer, contrasting with those performed in the FT. Inside the STBL, A_P matches approximately the theoretical value of 4/3 in the range of 5-100 m, indicating isotropic properties of turbulence in the inertial subrange of the energy cascade. The anisotropy ratios gradually decreases for larger scales which are of the order of the integral lengthscale (see sec. 5.5). The scales of the size of a few hundred meters, which is close to the boundary layer depth (about 850 m), might be additionally influenced by the




Figure 18. As in Fig. 17 but for the decoupled STBL (flight #14).

proximity of the bottom and top interfaces limiting their vertical extent. On the opposite side of the spectrum (small scales), A_P can be affected by the differences in the spectral transfer functions of the sonic anemometer for different velocity components (Kaimal et al., 1968). Similar effect was briefly described by Siebert et al. (2006b). In the FT, A_P hardly reaches 1.0 because vertical excursions are damped by stability. In case of LEG1079, it is particularly small, probably because that level was very close to the strongly stable temperature inversion.

In the decoupled STBL sampled in flight #14 (Fig. 18), A_P follows similar pattern as observed in flight #5. Nonetheless, maximum values are higher, reaching up to 1.7 at the scale of 20-40 m in LEG143 and LEG992 which are the lowest and highest segment inside STBL. One may speculate those scales, featuring prevalence of vertical fluctuations, are related to the typical size of surface layer plumes and to the typical size of cloud top downdrafts manifested as cloud holes (Gerber et al., 2005). The range of scales where A_P indicates conditions close to local isotropy is narrower than in the coupled STBL. On the side of large scales, A_P falls below the theoretical 4/3 already at around 70 m for the two central LEGs and at around 50 m for the two peripheral LEGs (regarding the perspective of the STBL). This observation can be related to the integral length scales which are smaller than in flight #5 for the most part (see sec. 5.5). What is more, the depths of the two sections of the boundary layer corresponding to the supposed circulation circuits (~ 500 m) are also smaller than the total depth of the coupled STBL (~ 850 m).

5.5 Length scales

In the coupled STBL, the estimated integral scales vary around 100-150 m (Fig. 19). The longitudinal scale L_u increases, whereas the vertical L_w decreases with height. The ratio L_w/L_u decreases from about 1.3 in the lowest LEG to about 0.5 (as expected for isotropic turbulence) close to the cloud top. The variability of integral scales among the subsegments of the LEGs is extensive, reflecting poor averaging on relatively short distances which prevents accurate calculation of decorrelation length.




Figure 19. Turbulent length scales in the coupled STBL (flight #5): (a) integral scale L , (b) Taylor microscale λ , (c) Kolmogorov scale η . Subscripts u and w denote horizontal and vertical velocity components, respectively. Some of the results for LEG1079 in the FT are out of the range presented.




Figure 20. As in Fig. 19 but for the decoupled STBL (flight #14).

Estimated Taylor microscales fit into the range of 30-80 cm and decline with height from the middle to the top of the STBL. As predicted, the longitudinal λ_u are larger than the vertical λ_w . Their ratio λ_u/λ_w equals $\sqrt{2}$ (corresponding to isotropy of small-scale turbulence) only in the cloud LEG and is larger below. We may speculate that the turbulence is close to isotropic at the time and location of generation but such isotropy is broken in the process of transport. Kolmogorov microscale is almost constant across the STBL (~ 2 mm) which can be expected as it depends practically only on the dissipation rate (the viscosity changes only by a minor part in the lower atmosphere). There is also no major difference between the horizontal and vertical direction.

In the decoupled STBL, integral scales are significantly smaller in comparison to the previous case, hardly exceeding 100 m (Fig. 20). The longitudinal L_u dominates over the vertical L_w , probably due to the separation of the circulation into two circuits and weak static stability which both limit the vertical extent of eddies and promote horizontal elongation. In contrast to the coupled case, the ratio L_w/L_u equals about one half in the lowest LEG close to the surface which is, however, again the location of intensive TKE production.

Taylor microscale is mostly of the same order as in the former case. In the transition zone and in the cloud, substantial detachment between the longitudinal and the vertical can be observed. λ_u/λ_w is significantly larger than expected for isotropic turbulence. This effect is most pronounced in LEG448 close to the transition. We may speculate it might be the consequence of decaying turbulence – far from the production in the cloud and at the surface, the TKE is here dissipated and consumed by weak buoyant stability (sec. 5.2). Kolmogorov scale visibly fluctuates but on average stays close to constant across the STBL. In contrast to the coupled case, there is some difference between η_u and η_w which directly relates to $A_e^{sf_c}$ discussed in sec. 5.4.

6 Summary and discussion

Two cases of marine stratocumulus-topped boundary layer, coupled (C) and decoupled (D), have been compared in terms of stratification and turbulence properties. The observations were performed in summer in the region of Eastern North Atlantic with the use of the helicopter-borne platform ACTOS. Its moderate true air speed in combination with closely collocated fast-response instruments provides high spatial resolution measurements of turbulent fluctuations of wind velocity, temperature and humidity. Similarities and differences between the two cases can be summarized as follows.

1. Stratification

- C** Conserved variables, θ_l and q_t , feature nearly constant profiles up to the capping inversion at ~ 850 m. LCL can be considered consistent with cloud base height.
- D** Above the relatively well-mixed SML, θ_l slowly increases with height up to the capping inversion at ~ 1050 m, indicating weak stability. There is a significant gradient of q_t in the TSL. LCL is close to the observed CB in the SBL only. Decoupling of the STBL was detected according to simple thermodynamic criteria.

In both cases winds are moderate and appreciable wind shear is observed across the cloud top and the EIL.

2. TKE production

600 **C** TKE is efficiently generated by buoyancy with simultaneous importance of in-cloud and surface processes. Buoyancy production follows typical STBL profile: decreases with height from the surface upwards, vanishes or turns slightly negative below cloud base, to be again substantial inside the cloud due to latent heat release and diabatic cooling.

605 **D** TKE is generated by buoyancy at the surface and B decreases with height to zero at the SML top, turning into buoyancy consumption in the TSL. In the cloud B is weaker than at the surface, about three times smaller than for the coupled case. Buoyancy effects can be also deduced from spectral anisotropy in the uppermost and lowermost boundary layer LEGs which suggests dominance of vertical motions in scales of 10-40 m.

610 The contribution of shear to TKE production is not negligible in both cases. This result can be partly artifact because only the longitudinal term could be evaluated and due to inaccurate estimation of horizontal wind gradient involved in shear term.

3. Heat fluxes

615 In both cases latent heat flux qualitatively resembles the profile of B which is consistent with the considerable contribution of moisture transport to buoyancy in the lower part of the STBL. Q_l is large at the ocean surface and decreases to zero at similar level as the minimum of B . Sensible heat flux is positive throughout observed layers but mostly smaller than Q_l .

C Q_l and Q_s are positive and of significant magnitude close to the cloud top which can be attributed to diabatic cooling (radiative and/or evaporative).

620 **D** Q_l and Q_s are small close to the cloud top, about an order of magnitude weaker than for the coupled STBL. Additionally, Q_l vanishes at a level much lower in relation to the cloud base which might disturb moisture delivery from the ocean to the stratocumulus cloud.

4. Turbulent fluctuations

 In both cases TKE is dominated by the contribution of horizontal velocity fluctuations. Variances of temperature and humidity are significant in the regions where mixing between air volumes of different properties occurs – due to local gradients or sources/sinks, i.e. at the cloud top, at the surface and at the transition in the decoupled STBL.

625 **C** Maximum TKE is found in the middle of the SBL which together with positive $\langle w'^3 \rangle$ at this level point out the role of surface-related factors in generating convection. The vertical velocity variance suggests a profile somewhat different than the convective similarity scaling. In cloud, $\langle u'^2 \rangle$ and $\langle w'^2 \rangle$ are almost equal implying isotropic conditions.

D The SML follows the structure of a typical mixed layer with shear (c.f Stull, 1988). Updrafts are stronger than
 630 downdrafts. TKE, $\langle T'^2 \rangle$ and $\langle q_v'^2 \rangle$ are largest close to the transition. In cloud, fluctuations are relatively weak, in
 particular $\langle w'^2 \rangle$, in concordance with limited B and small heat fluxes.

5. TKE dissipation

C Derived ϵ varies weakly throughout the height, i.e. despite accidental variations no systematic layering can be
 635 observed. Although TKE is efficiently produced by buoyancy in the cloud and at the surface, it is probably re-
 distributed well across the depth before being dissipated by viscosity. The form of SFCs and PSDs is reasonably
 consistent with theoretical predictions for inertial range scaling in homogeneous, isotropic, stationary turbulence
 (Kolmogorov, 1941). Though, less steep scaling (smaller absolute values of s and p) can be found at some places
 in the SBL.

D Derived ϵ is smaller than in the coupled case and features differences between the sublayers. Despite relatively high
 640 B at the surface, similar to the coupled case, average ϵ in the SML is smaller than in the SCL. Importantly, SFCs
 and PSDs scaling in the inertial range considerably deviates from the theoretical. Such behavior is characteristic
 for decaying turbulence (less energy than expected in large scales). These deviations are more pronounced and
 more variable in the SCL and SBL in comparison with the SML, underlining different character of turbulence in
 the upper and lower part of the decoupled STBL. Probably, TKE generated in the surface region and in the cloud,
 645 respectively, is redistributed in the two circulation zones separately, without major transport through the transition.

Discrepancies between PROF-derived and LEG-derived quantities result from the contrast between local and mean
 turbulence characteristics. The observed relative tendencies are consistent among derivation methods and velocity com-
 ponents, in spite of discrepancies in the absolute values.

6. Anisotropy of turbulence

C Derived anisotropy ratios indicate that turbulence is relatively close to isotropy. This condition is met best in the
 650 cloud where significant TKE production occurs.

D The degree of anisotropy varies between the sublayers. In the uppermost part (SCL and SBL) horizontal small-
 scale velocity fluctuations dominate over the vertical. This effect is less pronounced in the SML. The change in
 anisotropy ratios in the TSL coincides with the difference in s and p right below the strong q_v gradient.

7. Length scales of turbulence

Integral length scales of the order of 100 m show that turbulent eddies are substantially smaller than the depths of STBL
 or decoupled sublayers. Thus, they can be considered small enough to be transported by larger circulations.

C In the middle SBL, w' is correlated on longer distances than u' , while the opposite holds in the SCL. This agrees
 660 with the supposed form of circulation in the boundary layer, i.e. downdrafts originated at cloud top and updrafts
 originated at the surface pair in the middle and diverge horizontally in the vicinity of top and bottom boundaries.

D Integral length scales are smaller than in the coupled case. In accordance with anisotropy ratios, L_u is larger than L_w . The same holds for Taylor microscales. The difference between λ_u and λ_w is particularly pronounced close to the transition. It seems that even smaller turbulent eddies there are elongated in horizontal.

Interestingly, $L_w/L_u \approx \frac{1}{2}$ implied by isotropy assumption holds only in the regions of intensive buoyant TKE production: in the cloud for the coupled STBL and close to the surface for the decoupled STBL. Kolmogorov scale is ~ 2 mm in both cases.

Most of our results concerning the coupled case are consistent with previous studies of stratocumulus dynamics (e.g. Nicholls and Turton, 1986; Duynkerke et al., 1995; Stevens et al., 2005; Kopec et al., 2016; Dodson and Small Griswold, 2021). In particular, the B profiles show that convection is driven both by cloud top cooling and by surface thermal instability. However, our results suggest the profile of $\langle w'^2 \rangle$ being somewhat different than the convective similarity scaling (Lenschow et al., 1980) but rather having maximum in the cloud and minimum below it, with A_2^{var} following the same behavior, similarly to Dodson and Small Griswold (2021). Together with high TKE and positive $\langle w'^3 \rangle$ in the middle SBL, this highlights the importance of surface process. It might be related to small cloud depth (relative to STBL depth) and net cooling at cloud top reduced during daytime in comparison to often considered nocturnal stratocumulus. In contrast to the works listed above, we do not clearly observe the maximum of ϵ at the top and at the bottom of the STBL, but it is rather because others applied considerable horizontal averaging in comparison to local variability captured in our PROFs.

Our observations in the decoupled STBL summarized in points 1-4 fit well into the range of conditions reported in the literature, in particular the properties of the SML. Buoyant TKE production is positive in the cloud, while there is a region of negative B around the transition (Nicholls, 1984; Nicholls and Turton, 1986; Turton and Nicholls, 1987; Durand and Bourcy, 2001). Moreover, Q_l decreases from the surface to zero at the transition and it is substantially larger than Q_s in the SML (Nicholls, 1984; Tjernström and Rogers, 1996; De Roode and Duynkerke, 1997; Lambert and Durand, 1999; Durand and Bourcy, 2001). However, Lambert and Durand (1999) dispute the nearly linear character of this decrease, suggesting rather sharp gradient right at the SML top. Comparable to Nicholls (1984), our variances $\langle T'^2 \rangle$, $\langle q_v'^2 \rangle$ are significant close to the surface and have local minimum in the middle SML where in turn $\langle w'^2 \rangle$ is relatively large. As in De Roode and Duynkerke (1997), $\langle w'^3 \rangle$ is positive in the SML and nearly zero in the SCL, although the LEGs were rather too short to ensure statistical significance of those results. On the other hand, we did not collect enough data in the SCL and SBL to judge whether they together exhibit upside-down convective scaling as in Nicholls and Turton (1986); Tjernström and Rogers (1996); De Roode and Duynkerke (1997).

The results of our comparison between coupled and decoupled STBL are in agreement with the common concept of the dominant mixing patterns in such boundary layers (e.g. Wood, 2012). Decoupling occurs when the thermally driven circulation weakens to the level that it cannot mix air throughout entire depth. Then, STBL separates into two parts: cloud driven and surface driven. Explaining the particular mechanism of decoupling operating in our case is beyond the scope of this study and would require more complete data on airmass history. Nevertheless, "deepening-warming" mechanism (Bretherton and Wyant, 1997) seems plausible. Such conclusion was reached by Kazemirad and Miller (2020) who modeled lagrangian evolution of

695 STBL on synoptic scale in the period including our measurements. Deepening-warming is typical for the region of the Eastern North Atlantic where air masses are advected over progressively warmer waters. The most important driver for this process is the increasing ratio of surface latent heat flux to net radiative cooling in the cloud. The former was indeed relatively large, the latter was probably reduced by daytime solar heating. In addition, some precipitation was reported shortly before the flight and evaporative cooling could have contributed to stabilizing the lower STBL. Finally, decoupling occurs more readily for large
700 entrainment efficiency. Derived B is weak in the cloud, much smaller than in the coupled one, which might be the result of enhanced entrainment warming offsetting radiative cooling (c.f. De Roode and Duynkerke, 1997).




Figure 21. Schematic of main processes in the coupled (left) and decoupled (right) STBL: primary circulation (yellow arrows), turbulence eddy cascade (circular arrows confined in an angle with extent proportional to inertial range scaling exponent p), TKE buoyancy production (red B letter of size proportional to strength), sensible and latent heat fluxes (purple and blue arrows, respectively, of length proportional to strength) at the surface and in the cloud top region.

The important novelty of our work are the results on small-scale turbulence (points 5-7 of the summary). As far as we know, local ϵ profile, inertial range scaling exponents and anisotropy ratios were not addressed in the context of STBL coupling before. Based on the observations, we hypothesize that turbulence is redistributed across the depth of the coupled STBL but
705 in case of the decoupled STBL primarily in the sublayer where it was generated. Therefore, specific microscopic properties – TKE dissipation rate, inertial range scaling and anisotropy – can differ between the parts of the decoupled STBL.

We consider it important to emphasize often omitted distinction between *circulation* and *turbulence*. By *circulation* we understand motions responsible for mixing across relatively deep layers, of vertical scales comparable to the boundary layer depth. They usually originate from thermally driven plumes, sinking from cloud top or rising from the surface. Circulation
710 might take form of organized structures of downdrafts and updrafts (resembling Rayleigh-Bennard convection cells). Those correspond to the peak in vertical velocity spectra, typically at ~ 1 km in STBL (Lambert et al., 1999). *Turbulence* features cascade of eddies with universal scaling properties (Kolmogorov, 1941), spanning from the integral length scale (~ 100 m in

STBL) down to the Kolmogorov scale (~ 1 mm) where TKE is dissipated by viscosity. Such turbulence can be generated by flow instabilities at specific locations (here typically close to the surface and cloud top) and distributed by circulation within
 715 STBL, alongside other constituents. Importantly, the variances and fluxes estimated in our study include contributions of both phenomena. Circulation is only partly resolved as we applied the cutoff of ~ 1 km in Reynolds decomposition due to limited length of LEGs. Similar issue was also raised by De Roode and Duynkerke (1997). The advantage of our work is a good representation of turbulence because we resolve significant portion of the inertial range. The main processes operating in the coupled and decoupled STBLs, including circulation and turbulence, are schematically illustrated in Fig. 21.



Figure 22. Cumulus clouds under stratocumulus in the decoupled STBL. Photograph was taken during PROF5 of flight #14 by the camera mounted on the bottom of ACTOS.

720 Both turbulence and circulation can contribute to vertical transport of heat and moisture which is crucial for maintaining stratocumulus cloud. In the decoupled STBL, transport by turbulence through the transition is rather limited. However, we speculate it can be efficiently realized by a small number of updrafts which are strong and moist enough to penetrate the conditionally unstable TSL (measured $\Gamma_T = -7.1 \text{ K km}^{-1}$, moist adiabatic $\Gamma_T = -4.7 \text{ K km}^{-1}$), reach their LCL and form cumulus clouds. The image of those cumuli was captured by a camera onboard ACTOS (Fig. 22). Based on the series of
 725 images from PROF5, we estimated the cloud base height ~ 660 m (equal to mean LCL in the TSL) and cloud depth ~ 100 m. None of those cumuli was penetrated by ACTOS, so it is not possible to distinguish dynamic effects responsible for their formation. Detailed analysis of vertical transport calls for high-resolution numerical simulations to be setup with the help of our results.

The onset of cumulus convection depends on the properties of the TSL which is then important for overall STBL dynamics.
 730 However, it is a challenge to conduct relevant systematic climatological analysis of TSL existence and properties due to limited number of observations. The reason is often insufficient resolution of routine radiosoundings. For instance, the layer of the strongest gradient in q_v (550-600 m) penetrated in PROF5 features the differences of $\Delta\theta = 0.4 \text{ K}$, $\Delta q_v = 1 \text{ g kg}^{-1}$ (equivalent to $\Delta RH = 8 \%$) and $\Delta LCL = 160 \text{ m}$. With the ascent rate of $\sim 5 \text{ m s}^{-1}$ and sampling interval of $\sim 2 \text{ s}$, a hygrometer with the time constant of a second and the accuracy of a single percent in RH would be desired. Moreover, TSL is not exactly flat but
 735 rather undulated as suggested by our data of LEG448. Therefore, even aircraft measurements may fail to properly capture local

conditions. This was pointed out already by (Turton and Nicholls, 1987, p. 997) who underlined the role of good observation strategy: “While cloud layer decoupling is predicted to occur quite often, the consequential modification of the horizontally averaged vertical thermodynamic structure remains fairly small. (...) Data averaged in this way will appear ‘nearly well-mixed’ whether separation has occurred or not. A more detailed analysis of individual profiles and turbulence data is necessary to
740 determine whether decoupling has taken place”.

Data availability. The whole dataset collected within the ACORES field project is planned to be archived on the PANGAEA server for public access. The data used in the present study is also available from the authors upon request.

Author contributions. All the authors participated in the instrument preparation, measurements and data postprocessing within the ACORES project where H.S. was the principal investigator. J.L.N. designed the presented analysis with advice from S.P.M and H.S. The analysis was
745 performed by J.L.N. with contributions from K.E.S. and with guidance of H.S. and S.P.M. J.L.N. wrote the manuscript with contributions from S.P.M. and H.S.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The field campaign ACORES was supported by several grants of the Deutsche Forschungsgesellschaft (DFG, with grants SI 1543/4-1, WE 1900/33-1, WE 2757/2-1, and HE 6770/2-1) and Polish National Science Centre (grant agreement 2013/08/A/ST10/00291).
750 J.L.N. acknowledges the one year grant awarded by the German Academic Exchange Service (DAAD) for his research visit to Leibniz Institute for Tropospheric Research. We acknowledge the use of imagery from the Worldview Snapshots application (<https://wvs.earthdata.nasa.gov>), part of the Earth Observing System Data and Information System (EOSDIS). The authors are also grateful to Dr. Marta Wacławczyk for discussions on the manuscript and to Katarzyna Nurowska for drawing the sketch in Fig. 21.

Appendix A: Average conditions in the sublayers

755 Average meteorological parameters and turbulence properties inside the sublayers of the atmosphere are summarized in Tables A1 and A2 for the coupled (flight #5) and decoupled (flight #14) case, respectively. The selection of the sublayers is explained in sec. 3. The average values were obtained from the data of the same PROF which served for sublayer selection, i.e. PROF1 in coupled case, PROF5 in decoupled case. Γ_T , Nb^2 and Sh^2 were calculated by estimating derivatives over sublayer depth. Other parameters were simply averaged in the relevant altitude range.

Table A1. Average conditions inside the sublayers in the case of coupled STBL (flight #5).

	SBL Subcloud layer	SCL Strato- cumulus layer	EIL Entrainment interface layer	FTL Free tropo- spheric layer
Height [m]	0 - 715	715 - 855	855 - 935	1005 - 1385
T [° C]	16.24	12.59	14.53	14.41
θ_t [° C]	17.62	17.52	20.59	23.54
Γ_T [K km ⁻¹]	-10.9	-10.1	73.9	-7.2
q_t [g kg ⁻¹]	9.53	9.43	3.19	3.89
U [m s ⁻¹]	5.3	5.0	6.5	6.8
dd [°]	337	330	329	323
LCL [m]	814	845	3363	3130
Nb ² [10 ⁻⁴ s ⁻²]	-0.4	-0.6	15.4	0.7
Sh ² [10 ⁻⁴ s ⁻²]	0.0	0.3	5.1	1.0
ϵ_w^{sfc} [10 ⁻⁴ m ² s ⁻³]	5.6	6.1	1.9	0.2
ϵ_u^{sfc} [10 ⁻⁴ m ² s ⁻³]	6.5	6.6	2.2	0.8
ϵ_w^{psd} [10 ⁻⁴ m ² s ⁻³]	5.6	5.1	1.5	0.3
ϵ_u^{psd} [10 ⁻⁴ m ² s ⁻³]	9.2	8.5	2.6	1.2
s_w	0.61	0.67	0.29	0.03
s_u	0.47	0.55	0.34	0.05
p_w	-1.53	-1.70	-1.10	-0.31
p_u	-1.25	-1.42	-1.03	-0.23
R_w^{sfc}	0.96	0.97	0.79	0.37
R_u^{sfc}	0.95	0.96	0.87	0.42
R_w^{psd}	-0.94	-0.95	-0.81	-0.49
R_u^{psd}	-0.91	-0.93	-0.81	-0.41
$\epsilon_w^{sfc}/\epsilon_u^{sfc}$	0.87	0.94	0.54	0.28
$\epsilon_w^{psd}/\epsilon_u^{psd}$	0.62	0.63	0.41	0.29
η_w^{sfc} [mm]	1.7	1.6	4.0	4.0
η_u^{sfc} [mm]	1.6	1.6	3.0	2.9

Table A2. Average conditions inside the sublayers in the case of decoupled STBL (flight #14).

	SML Surface mixed layer	TSL Transition layer	SBL Subcloud layer	SCL Strato- cumulus layer	EIL Entrainment interface layer	FTL Free tropo- spheric layer
Height [m]	0 - 385	485 - 615	615 - 865	865 - 1045	1045 - 1095	1150 - 1400
T [$^{\circ}$ C]	18.06	15.79	14.36	13.10	13.90	16.05
θ_l [$^{\circ}$ C]	18.81	19.20	19.61	20.16	22.37	26.29
Γ_T [K km $^{-1}$]	-10.1	-7.1	-7.4	-2.9	84.6	-5.2
q_t [g kg $^{-1}$]	11.65	11.08	10.75	10.68	9.65	8.48
U [m s $^{-1}$]	6.5	5.8	5.5	6.0	7.3	9.9
dd [$^{\circ}$]	314	308	314	322	322	325
LCL [m]	508	658	769	905	1328	2040
Nb 2 [10^{-4} s $^{-2}$]	0.5	0.9	0.1	1.5	28.0	1.5
Sh 2 [10^{-4} s $^{-2}$]	0.2	0.8	0.1	0.1	45.6	1.5
ϵ_w^{sfc} [10^{-4} m 2 s $^{-3}$]	0.6	1.1	1.2	1.3	1.8	0.1
ϵ_u^{sfc} [10^{-4} m 2 s $^{-3}$]	1.2	3.4	3.5	4.6	3.8	0.6
ϵ_w^{psd} [10^{-4} m 2 s $^{-3}$]	0.6	1.5	1.6	1.9	2.5	0.1
ϵ_u^{psd} [10^{-4} m 2 s $^{-3}$]	1.9	5.6	5.4	7.8	6.5	0.9
s_w	0.52	0.45	0.47	0.44	0.38	0.04
s_u	0.38	0.27	0.33	0.24	0.28	0.16
p_w	-1.39	-1.10	-1.10	-1.11	-1.05	-0.34
p_u	-1.16	-0.76	-0.78	-0.68	-0.85	-0.46
R_w^{sfc}	0.95	0.96	0.96	0.96	0.87	0.30
R_u^{sfc}	0.94	0.91	0.92	0.90	0.89	0.70
R_w^{psd}	-0.90	-0.89	-0.89	-0.89	-0.81	-0.45
R_u^{psd}	-0.89	-0.79	-0.79	-0.79	-0.83	-0.59
$\epsilon_w^{sfc}/\epsilon_u^{sfc}$	0.48	0.31	0.34	0.29	0.48	0.09
$\epsilon_w^{psd}/\epsilon_u^{psd}$	0.34	0.26	0.30	0.24	0.39	0.09
η_w^{sfc} [mm]	3.0	2.6	2.5	2.5	2.4	5.6
η_u^{sfc} [mm]	2.4	1.9	1.9	1.8	1.9	3.0

- Akinlabi, E. O., Wacławczyk, M., Mellado, J. P., and Malinowski, S. P.: Estimating turbulence kinetic energy dissipation rates in the numerically simulated stratocumulus cloud-top mixing layer: Evaluation of different methods, *Journal of the Atmospheric Sciences*, 76, 1471–1488, <https://doi.org/10.1175/JAS-D-18-0146.1>, 2019.
- Albrecht, B. A., Bretherton, C. S., Johnson, D., Schubert, W. H., and Frisch, A. S.: The Atlantic Stratocumulus Transition Experiment - AS-
765 TEX, *Bulletin - American Meteorological Society*, 76, 889–904, [https://doi.org/10.1175/1520-0477\(1995\)076<0889:TASTE>2.0.CO;2](https://doi.org/10.1175/1520-0477(1995)076<0889:TASTE>2.0.CO;2), 1995.
- Betts, A. K.: Non-precipitating cumulus convection and its parameterization, *Quarterly Journal of the Royal Meteorological Society*, 99, 178–196, <https://doi.org/10.1002/qj.49709941915>, 1973.
- Bolton, D.: The computation of equivalent potential temperature., *Monthly Weather Review*, 108, 1046–1053, [https://doi.org/10.1175/1520-
770 0493\(1980\)108<1046:TCOEPT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1980)108<1046:TCOEPT>2.0.CO;2), 1980.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.-Y.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*, edited by Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9781107415324.016>,
775 2013.
- Bretherton, C. S. and Wyant, M. C.: Moisture transport, lower-tropospheric stability, and decoupling of cloud-topped boundary layers, *Journal of the Atmospheric Sciences*, 54, 148–167, [https://doi.org/10.1175/1520-0469\(1997\)054<0148:MTL TSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<0148:MTL TSA>2.0.CO;2), 1997.
- Caldwell, P., Bretherton, C. S., and Wood, R.: Mixed-layer budget analysis of the diurnal cycle of entrainment in southeast Pacific stratocumulus, *Journal of the Atmospheric Sciences*, 62, 3775–3791, <https://doi.org/10.1175/JAS3561.1>, 2005.
- Cruette, D., Marillier, A., Dufresne, J. L., Grandpeix, J. Y., Nacass, P., and Bellec, H.: Fast temperature and true airspeed measurements with the airborne ultrasonic anemometer-thermometer (AUSAT), *Journal of Atmospheric and Oceanic Technology*, 17, 1020–1039, [https://doi.org/10.1175/1520-0426\(2000\)017<1020:FTATAM>2.0.CO;2](https://doi.org/10.1175/1520-0426(2000)017<1020:FTATAM>2.0.CO;2), 2000.
- De Roode, S. R. and Duynkerke, P. G.: Observed lagrangian transition of stratocumulus into cumulus during ASTEX: Mean state and turbulence structure, *Journal of the Atmospheric Sciences*, 54, 2157–2173, [https://doi.org/10.1175/1520-
785 0469\(1997\)054<2157:OLTOSI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<2157:OLTOSI>2.0.CO;2), 1997.
- De Roode, S. R., Sandu, I., van der Dussen, J. J., Ackerman, A. S., Blossey, P., Jarecka, D., Lock, A., Siebesma, A. P., and Stevens, B.: Large-eddy simulations of EUCLIPSE-GASS lagrangian stratocumulus-to-cumulus transitions: Mean state, turbulence, and decoupling, *Journal of the Atmospheric Sciences*, 73, 2485–2508, <https://doi.org/10.1175/JAS-D-15-0215.1>, 2016.
- Deardorff, J. W.: Cloud top entrainment instability., *Journal of the Atmospheric Sciences*, 37, 131–147, [https://doi.org/10.1175/1520-
790 0469\(1980\)037<0131:CTEI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0131:CTEI>2.0.CO;2), 1980.
- Dodson, D. S. and Small Griswold, J. D.: Turbulent and boundary layer characteristics during VOCALS-REx, *Atmospheric Chemistry and Physics*, 21, 1937–1961, <https://doi.org/10.5194/acp-21-1937-2021>, 2021.
- Dong, X., Schwantes, A. C., Xi, B., and Wu, P.: Investigation of the marine boundary layer cloud and CCN properties under coupled and decoupled conditions over the Azores, *Journal of Geophysical Research: Atmospheres*, 120, 6179–6191, <https://doi.org/10.1002/2014JD022939>, 2015.
795

Durand, P. and Bourcy, T.: Observations of the turbulence structure within two stratocumulus-topped, marine boundary layers, *Boundary-Layer Meteorology*, 99, 105–125, <https://doi.org/10.1023/A:1018999221303>, 2001.

Duynkerke, P. G., Heqing Zhang, and Jonker, P. J.: Microphysical and turbulent structure of nocturnal stratocumulus as observed during ASTEX, *Journal of the Atmospheric Sciences*, 52, 2763–2777, [https://doi.org/10.1175/1520-0469\(1995\)052<2763:MATSON>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<2763:MATSON>2.0.CO;2), 1995.

Edson, J. B., Hinton, A. A., Prada, K. E., Hare, J. E., and Fairall, C. W.: Direct covariance flux estimates from mobile platforms at sea, *Journal of Atmospheric and Oceanic Technology*, 15, 547–562, [https://doi.org/10.1175/1520-0426\(1998\)015<0547:DCFEFM>2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015<0547:DCFEFM>2.0.CO;2), 1998.

Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk parameterization of air-sea fluxes: Updates and verification for the COARE algorithm, *Journal of Climate*, 16, 571–591, [https://doi.org/10.1175/1520-0442\(2003\)016<0571:BPOASF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0571:BPOASF>2.0.CO;2), 2003.

Frehlich, R., Meillier, Y., Jensen, M. L., and Balsley, B.: A statistical description of small-scale turbulence in the low-level nocturnal jet, *Journal of the Atmospheric Sciences*, 61, 1079–1085, [https://doi.org/10.1175/1520-0469\(2004\)061<1079:ASDOST>2.0.CO;2](https://doi.org/10.1175/1520-0469(2004)061<1079:ASDOST>2.0.CO;2), 2004.

Gerber, H., Arends, B. G., and Ackerman, A. S.: New microphysics sensor for aircraft use, *Atmospheric Research*, 31, 235–252, [https://doi.org/10.1016/0169-8095\(94\)90001-9](https://doi.org/10.1016/0169-8095(94)90001-9), 1994.

Gerber, H., Frick, G., Malinowski, S. P., Brenguier, J. L., and Burnet, F.: Holes and entrainment in stratocumulus, *Journal of the Atmospheric Sciences*, 62, 443–459, <https://doi.org/10.1175/JAS-3399.1>, 2005.

Gerber, H., Malinowski, S. P., and Jonsson, H.: Evaporative and Radiative Cooling in POST Stratocumulus, *Journal of the Atmospheric Sciences*, 73, 3877–3884, <https://doi.org/10.1175/JAS-D-16-0023.1>, 2016.

Ghate, V. P., Miller, M. A., Albrecht, B. A., and Fairall, C. W.: Thermodynamic and radiative structure of stratocumulus-topped boundary layers, *Journal of the Atmospheric Sciences*, 72, 430–451, <https://doi.org/10.1175/JAS-D-13-0313.1>, 2015.

Goren, T., Rosenfeld, D., Sourdeval, O., and Quaas, J.: Satellite Observations of Precipitating Marine Stratocumulus Show Greater Cloud Fraction for Decoupled Clouds in Comparison to Coupled Clouds, *Geophysical Research Letters*, 45, 5126–5134, <https://doi.org/10.1029/2018GL078122>, 2018.

Haman, K. E., Makulski, A., Malinowski, S. P., and Busen, R.: A new ultrafast thermometer for airborne measurements in clouds, *Journal of Atmospheric and Oceanic Technology*, 14, 217–227, [https://doi.org/10.1175/1520-0426\(1997\)014<0217:ANUTFA>2.0.CO;2](https://doi.org/10.1175/1520-0426(1997)014<0217:ANUTFA>2.0.CO;2), 1997.

Hartmann, D. L., Ockert-Bell, M. E., and Michelsen, M. L.: The Effect of Cloud Type on Earth’s Energy Balance: Global Analysis, *Journal of Climate*, 5, 1281–1304, [https://doi.org/10.1175/1520-0442\(1992\)005<1281:teocto>2.0.co;2](https://doi.org/10.1175/1520-0442(1992)005<1281:teocto>2.0.co;2), 1992.

Jen-La Plante, I., Ma, Y., Nurowska, K., Gerber, H., Khelif, D., Karpinska, K., Kopec, M. K., Kumala, W., and Malinowski, S. P.: Physics of Stratocumulus Top (POST): Turbulence characteristics, *Atmospheric Chemistry and Physics*, 16, 9711–9725, <https://doi.org/10.5194/acp-16-9711-2016>, 2016.

Jones, C. R., Bretherton, C. S., and Leon, D.: Coupled vs. decoupled boundary layers in VOCALS-REx, *Atmospheric Chemistry and Physics*, 11, 7143–7153, <https://doi.org/10.5194/acp-11-7143-2011>, 2011.

JPL MUR MEaSUREs Project: GHRSSST Level 4 MUR Global Foundation Sea Surface Temperature Analysis. Ver. 4.1. PO.DAAC, <https://doi.org/10.5067/GHGMR-4FJ04>, 2015.

Kaimal, J. C., Wyngaard, J. C., and Haugen, D. A.: Deriving Power Spectra from a Three-Component Sonic Anemometer, *Journal of Applied Meteorology*, 7, 827–837, [https://doi.org/10.1175/1520-0450\(1968\)007<0827:dpsfat>2.0.co;2](https://doi.org/10.1175/1520-0450(1968)007<0827:dpsfat>2.0.co;2), 1968.

- Katzwinkel, J., Siebert, H., and Shaw, R. A.: Observation of a Self-Limiting, Shear-Induced Turbulent Inversion Layer Above Marine Stratocumulus, *Boundary-Layer Meteorology*, 145, 131–143, <https://doi.org/10.1007/s10546-011-9683-4>, <https://link.springer.com/article/10.1007/s10546-011-9683-4>, 2012.
- 835 Kazemirad, M. and Miller, M. A.: Summertime post-cold-frontal marine stratocumulus transition processes over the eastern north atlantic, *Journal of the Atmospheric Sciences*, 77, 2011–2037, <https://doi.org/10.1175/JAS-D-19-0167.1>, 2020.
- Kolmogorov, A. N.: The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers, *Dokl. Akad. Nauk SSSR*, 30, 301–304, <https://doi.org/10.1098/rspa.1991.0075>, 1941.
- Kolmogorov, A. N.: A refinement of previous hypotheses concerning the local structure of turbulence in a viscous incompressible fluid at
840 high Reynolds number, *Journal of Fluid Mechanics*, 13, 82–85, <https://doi.org/10.1017/S0022112062000518>, 1962.
- Kopec, M. K., Malinowski, S. P., and Piotrowski, Z. P.: Effects of wind shear and radiative cooling on the stratocumulus-topped boundary layer, *Quarterly Journal of the Royal Meteorological Society*, 142, 3222–3233, <https://doi.org/10.1002/qj.2903>, 2016.
- Lambert, D. and Durand, P.: The marine atmospheric boundary layer during semaphore. I: Mean vertical structure and non-axisymmetry of turbulence, *Quarterly Journal of the Royal Meteorological Society*, 125, 495–512, <https://doi.org/10.1002/qj.49712555407>, 1999.
- 845 Lambert, D., Durand, P., Thoumieux, F., Bénech, B., and Druilhet, A.: The marine atmospheric boundary layer during semaphore. II: Turbulence profiles in the mixed layer, *Quarterly Journal of the Royal Meteorological Society*, 125, 513–528, <https://doi.org/10.1002/qj.49712555408>, 1999.
- Lampert, A., Hartmann, J., Pätzold, F., Lobitz, L., Hecker, P., Kohnert, K., Larmanou, E., Serafimovich, A., and Sachs, T.: Comparison of Lyman-alpha and LI-COR infrared hygrometers for airborne measurement of turbulent fluctuations of water vapour, *Atmospheric
850 Measurement Techniques*, 11, 2523–2536, <https://doi.org/10.5194/amt-11-2523-2018>, 2018.
- Lenschow, D. H.: Aircraft Measurements in the Boundary Layer, in: *Probing the Atmospheric Boundary Layer*, pp. 39–55, American Meteorological Society, https://doi.org/10.1007/978-1-944970-14-7_5, 1986.
- Lenschow, D. H., Wyngaard, J. C., and Pennell, W. T.: Mean-field and second-moment budgets in a baroclinic, convective boundary layer., *Journal of the Atmospheric Sciences*, 37, 1313–1326, [https://doi.org/10.1175/1520-0469\(1980\)037<1313:MFASMB>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1313:MFASMB>2.0.CO;2), 1980.
- 855 Lenschow, D. H., Mann, J., and Kristensen, L.: How long is long enough when measuring fluxes and other turbulence statistics?, *Journal of Atmospheric and Oceanic Technology*, 11, 661–673, [https://doi.org/10.1175/1520-0426\(1994\)011<0661:HLILEW>2.0.CO;2](https://doi.org/10.1175/1520-0426(1994)011<0661:HLILEW>2.0.CO;2), 1994.
- Lilly, D. K.: Models of cloud-topped mixed layers under a strong inversion, *Quarterly Journal of the Royal Meteorological Society*, 94, 292–309, <https://doi.org/10.1002/qj.49709440106>, 1968.
- Malinowski, S. P., Gerber, H., Jen-La Plante, I., Kopec, M. K., Kumala, W., Nurowska, K., Chuang, P. Y., Khelif, D., and Haman, K. E.:
860 Physics of Stratocumulus Top (POST): Turbulent mixing across capping inversion, *Atmospheric Chemistry and Physics*, 13, 12 171–12 186, <https://doi.org/10.5194/acp-13-12171-2013>, 2013.
- Markowski, P. and Richardson, Y.: *Mesoscale Meteorology in Midlatitudes*, John Wiley and Sons, Ltd, Chichester, UK, <https://doi.org/10.1002/9780470682104>, 2010.
- Mellado, J. P.: Cloud-Top Entrainment in Stratocumulus Clouds, *Annual Review of Fluid Mechanics*, 49, 145–169,
865 <https://doi.org/10.1146/annurev-fluid-010816-060231>, 2017.
- Muschinski, A., Frehlich, R., Jensen, M., Hugo, R., Hoff, A., Eaton, F., and Balsley, B.: Fine-scale measurements of turbulence in the lower troposphere: An intercomparison between a kit-and balloon-borne, and a helicopter-borne measurement system, *Boundary-Layer Meteorology*, 98, 219–250, <https://doi.org/10.1023/A:1026520618624>, 2001.

- Muschinski, A., Frehlich, R. G., and Balsley, B. B.: Small-scale and large-scale intermittency in the nocturnal boundary layer and the residual layer, *Journal of Fluid Mechanics*, 515, 319–351, <https://doi.org/10.1017/S0022112004000412>, 2004.
- Nicholls, S.: The dynamics of stratocumulus: Aircraft observations and comparisons with a mixed layer model, *Quarterly Journal of the Royal Meteorological Society*, 110, 783–820, <https://doi.org/10.1002/qj.49711046603>, 1984.
- Nicholls, S. and Turton, J. D.: An observational study of the structure of stratiform cloud sheets: Part II. Entrainment, *Quarterly Journal of the Royal Meteorological Society*, 112, 461–480, <https://doi.org/10.1002/qj.49711247210>, 1986.
- Nowak, J. L., Kumala, W., Kwiatkowski, J., Kwiatkowski, K., Czyzewska, D., Karpinska, K., and Malinowski, S. P.: UltraFast Thermometer 2.0-new temperature sensor for airborne applications and its performance during ACORES 2017, in: *Geophysical Research Abstracts*, vol. 20, p. 12492, 2018.
- Pedersen, J. G., Ma, Y., Grabowski, W. W., and Malinowski, S. P.: Anisotropy of Observed and Simulated Turbulence in Marine Stratocumulus, *Journal of Advances in Modeling Earth Systems*, 10, 500–515, <https://doi.org/10.1002/2017MS001140>, 2018.
- Pope, S. B.: *Turbulent flows*, Cambridge University Press, Cambridge, <https://doi.org/10.1017/CBO9780511840531>, 2000.
- Randall, D. A.: Conditional instability of the first kind up-side down., *Journal of the Atmospheric Sciences*, 37, 125–130, [https://doi.org/10.1175/1520-0469\(1980\)037<0125:CIOTFK>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0125:CIOTFK>2.0.CO;2), 1980.
- Rémillard, J., Kollias, P., Luke, E., and Wood, R.: Marine boundary layer cloud observations in the Azores, *Journal of Climate*, 25, 7381–7398, <https://doi.org/10.1175/JCLI-D-11-00610.1>, 2012.
- Schneider, T., Kaul, C. M., and Pressel, K. G.: Possible climate transitions from breakup of stratocumulus decks under greenhouse warming, *Nature Geoscience*, 12, 164–168, <https://doi.org/10.1038/s41561-019-0310-1>, 2019.
- Siebert, H. and Muschinski, A.: Relevance of a tuning-fork effect for temperature measurements with the Gill solent HS ultrasonic anemometer-thermometer, *Journal of Atmospheric and Oceanic Technology*, 18, 1367–1376, [https://doi.org/10.1175/1520-0426\(2001\)018<1367:ROATFE>2.0.CO;2](https://doi.org/10.1175/1520-0426(2001)018<1367:ROATFE>2.0.CO;2), 2001.
- Siebert, H. and Teichmann, U.: Behaviour of an ultrasonic anemometer under cloudy conditions, *Boundary-Layer Meteorology*, 94, 165–169, <https://doi.org/10.1023/A:1002446723575>, 2000.
- Siebert, H., Wendisch, M., Conrath, T., Teichmann, U., and Heintzenberg, J.: A new tethered balloon-borne payload for fine-scale observations in the cloudy boundary layer, *Boundary-Layer Meteorology*, 106, 461–482, <https://doi.org/10.1023/A:1021242305810>, 2003.
- Siebert, H., Franke, H., Lehmann, K., Maser, R., Saw, E. W., Schell, D., Shaw, R. A., and Wendisch, M.: Probing finescale dynamics and microphysics of clouds with helicopter-borne measurements, *Bulletin of the American Meteorological Society*, 87, 1727–1738, <https://doi.org/10.1175/BAMS-87-12-1727>, 2006a.
- Siebert, H., Lehmann, K., and Wendisch, M.: Observations of small-scale turbulence and energy dissipation rates in the cloudy boundary layer, *Journal of the Atmospheric Sciences*, 63, 1451–1466, <https://doi.org/10.1175/JAS3687.1>, 2006b.
- Siebert, H., Szodry, K.-E., Egerer, U., Wehner, B., Henning, S., Chevalier, K., Lücknerath, J., Welz, O., Weinhold, K., Laueremann, F., Gottschalk, M., Ehrlich, A., Wendisch, M., Fialho, P., Roberts, G., Allwayin, N., Schum, S., Shaw, R. A., Mazzoleni, C., Mazzoleni, L., Nowak, J. L., Malinowski, S. P., Karpinska, K., Kumala, W., Czyzewska, D., Luke, E. P., Kollias, P., Wood, R., and Mellado, J. P.: Observations of Aerosol, Cloud, Turbulence, and Radiation Properties at the Top of the Marine Boundary Layer over the Eastern North Atlantic Ocean: The ACORES Campaign, *Bulletin of the American Meteorological Society*, 102, E123–E147, <https://doi.org/10.1175/bams-d-19-0191.1>, 2021.
- Stevens, B.: Cloud transitions and decoupling in shear-free stratocumulus-topped boundary layers, *Geophysical Research Letters*, 27, 2557–2560, <https://doi.org/10.1029/1999GL011257>, 2000.

- Stevens, B.: Entrainment in stratocumulus-topped mixed layers, *Quarterly Journal of the Royal Meteorological Society*, 128, 2663–2690, <https://doi.org/10.1256/qj.01.202>, 2002.
- Stevens, B., Cotton, W. R., Feingold, G., and Moeng, C. H.: Large-eddy simulations of strongly precipitating, shallow, stratocumulus-topped boundary layers, *Journal of the Atmospheric Sciences*, 55, 3616–3638, [https://doi.org/10.1175/1520-0469\(1998\)055<3616:LESOSP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<3616:LESOSP>2.0.CO;2), 1998.
- Stevens, B., Moeng, C. H., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J. C., Jiang, H., Khairoutdinov, M., Kirkpatrick, M. P., Lewellen, D. C., Lock, A., Müller, F., Stevens, D. E., Whelan, E., and Zhu, P.: Evaluation of large-eddy simulations via observations of nocturnal marine stratocumulus, *Monthly Weather Review*, 133, 1443–1462, <https://doi.org/10.1175/MWR2930.1>, 2005.
- Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Springer Netherlands, Dordrecht, <https://doi.org/10.1007/978-94-009-3027-8>, 1988.
- Sutherland, W.: The viscosity of gases and molecular force, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 36, 507–531, <https://doi.org/10.1080/14786449308620508>, 1893.
- Tjernstrom, M.: Turbulence length scales in stably stratified free shear flow analyzed from slant aircraft profiles, *Journal of Applied Meteorology*, 32, 948–963, [https://doi.org/10.1175/1520-0450\(1993\)032<0948:TLSISS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1993)032<0948:TLSISS>2.0.CO;2), 1993.
- Tjernström, M. and Rogers, D. P.: Turbulence structure in decoupled marine stratocumulus: A case study from the ASTEX field experiment, *Journal of the Atmospheric Sciences*, 53, 598–619, [https://doi.org/10.1175/1520-0469\(1996\)053<0598:TSIDMS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1996)053<0598:TSIDMS>2.0.CO;2), 1996.
- Turton, J. D. and Nicholls, S.: A Study of the Diurnal Variation of Stratocumulus Using A Multiple Mixed Layer Model, *Quarterly Journal of the Royal Meteorological Society*, 113, 969–1009, <https://doi.org/10.1002/qj.49711347712>, 1987.
- Vassilicos, J. C.: Dissipation in Turbulent Flows, *Annual Review of Fluid Mechanics*, 47, 95–114, <https://doi.org/10.1146/annurev-fluid-010814-014637>, 2015.
- Wacławczyk, M., Ma, Y.-F., Kopeć, J. M., and Malinowski, S. P.: Novel approaches to estimating turbulent kinetic energy dissipation rate from low and moderate resolution velocity fluctuation time series, *Atmos. Meas. Tech. Discuss.*, <https://doi.org/10.5194/amt-2016-401>, 2017.
- Wacławczyk, M., Gozigan, A. S., Nzotungishaka, J., Mohammadi, M., and P. Malinowski, S.: Comparison of Different Techniques to Calculate Properties of Atmospheric Turbulence from Low-Resolution Data, *Atmosphere*, 11, 199, <https://doi.org/10.3390/atmos11020199>, 2020.
- Wang, Z., Mora Ramirez, M., Dadashazar, H., MacDonald, A. B., Crosbie, E., Bates, K. H., Coggon, M. M., Craven, J. S., Lynch, P., Campbell, J. R., Azadi Aghdam, M., Woods, R. K., Jonsson, H., Flagan, R. C., Seinfeld, J. H., and Sorooshian, A.: Contrasting cloud composition between coupled and decoupled marine boundary layer clouds, *Journal of Geophysical Research: Atmospheres*, 121, 11,679–11,691, <https://doi.org/10.1002/2016JD025695>, 2016.
- Wendisch, M., Garrett, T. J., and Strapp, J. W.: Wind tunnel tests of the airborne PVM-100A response to large droplets, *Journal of Atmospheric and Oceanic Technology*, 19, 1577–1584, [https://doi.org/10.1175/1520-0426\(2002\)019<1577:WTTOTA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<1577:WTTOTA>2.0.CO;2), 2002.
- Werner, F., Siebert, H., Pilewskie, P., Schmeissner, T., Shaw, R. A., and Wendisch, M.: New airborne retrieval approach for trade wind cumulus properties under overlying cirrus, *Journal of Geophysical Research: Atmospheres*, 118, 3634–3649, <https://doi.org/10.1002/jgrd.50334>, 2013.

- Werner, F., Ditas, F., Siebert, H., Simmel, M., Wehner, B., Pilewskie, P., Schmeissner, T., Shaw, R. A., Hartmann, S., Wex, H., Roberts, G. C., and Wendisch, M.: Twomey effect observed from collocated microphysical and remote sensing measurements over shallow cumulus, *Journal of Geophysical Research: Atmospheres*, 119, 1534–1545, <https://doi.org/10.1002/2013JD020131>, 2014.
- 945 Wood, R.: Stratocumulus Clouds, *Monthly Weather Review*, 140, 2373–2423, <https://doi.org/10.1175/MWR-D-11-00121.1>, 2012.
- Wood, R. and Bretherton, C. S.: Boundary layer depth, entrainment, and decoupling in the cloud-capped subtropical and tropical marine boundary layer, *Journal of Climate*, 17, 3576–3588, [https://doi.org/10.1175/1520-0442\(2004\)017<3576:BLDEAD>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<3576:BLDEAD>2.0.CO;2), 2004.
- Wood, R., Wyant, M., Bretherton, C. S., Rémillard, J., Kollias, P., Fletcher, J., Stemmler, J., De Szoeki, S., Yuter, S., Miller, M., Mechem, D., Tselioudis, G., Chiu, J. C., Mann, J. A., O'Connor, E. J., Hogan, R. J., Dong, X., Miller, M., Ghate, V., Jefferson, A., Min, Q., Minnis, P., Palikonda, R., Albrecht, B., Luke, E., Hannay, C., and Lin, Y.: Clouds, aerosols, and precipitation in the marine boundary layer: An arm mobile facility deployment, *Bulletin of the American Meteorological Society*, 96, 419–439, <https://doi.org/10.1175/BAMS-D-13-00180.1>, 2015.
- 950 Xiao, H., Wu, C. M., and Mechoso, C. R.: Buoyancy reversal, decoupling and the transition from stratocumulus to shallow cumulus topped marine boundary layers, *Climate Dynamics*, 37, 971–984, <https://doi.org/10.1007/s00382-010-0882-3>, 2011.
- 955 Xiao, H., Wu, C. M., Mechoso, C. R., and Ma, H. Y.: A treatment for the stratocumulus-to-cumulus transition in GCMs, *Climate Dynamics*, 39, 3075–3089, <https://doi.org/10.1007/s00382-012-1342-z>, 2012.
- Yin, B. and Albrecht, B. A.: Spatial variability of atmospheric boundary layer structure over the eastern equatorial Pacific, *Journal of Climate*, 13, 1574–1592, [https://doi.org/10.1175/1520-0442\(2000\)013<1574:SVOABL>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<1574:SVOABL>2.0.CO;2), 2000.
- 960 Zheng, Y. and Li, Z.: Episodes of Warm-Air Advection Causing Cloud-Surface Decoupling During the MARCUS, *Journal of Geophysical Research: Atmospheres*, 124, 12 227–12 243, <https://doi.org/10.1029/2019JD030835>, 2019.
- Zheng, Y., Rosenfeld, D., and Li, Z.: The Relationships Between Cloud Top Radiative Cooling Rates, Surface Latent Heat Fluxes, and Cloud-Base Heights in Marine Stratocumulus, *Journal of Geophysical Research: Atmospheres*, 123, 11,678–11,690, <https://doi.org/10.1029/2018JD028579>, 2018a.
- 965 Zheng, Y., Rosenfeld, D., and Li, Z.: Estimating the Decoupling Degree of Subtropical Marine Stratocumulus Decks From Satellite, *Geophysical Research Letters*, 45, 12,560–12,568, <https://doi.org/10.1029/2018GL078382>, 2018b.
- Zheng, Y., Rosenfeld, D., and Li, Z.: A More General Paradigm for Understanding the Decoupling of Stratocumulus-Topped Boundary Layers: The Importance of Horizontal Temperature Advection, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL087697>, 2020.