

Measurement report: Receptor modelling for source identification of urban fine and coarse particulate matter using hourly elemental composition

5 Magdalena Reizer¹, Giulia Calzolari², Katarzyna Maciejewska¹, José A. G. Orza³, Silvia Nava², Franco Lucarelli², Katarzyna Juda-Rezler¹

¹ Faculty of Building Services, Hydro and Environmental Engineering, Warsaw University of Technology, Warsaw, Poland

² Department of Physics and Astronomy, University of Florence and National Institute of Nuclear Physics (INFN), Sesto Fiorentino, Italy

³ SCOLAB, Física Aplicada, Universidad Miguel Hernández, Elche, Spain

10 *Correspondence to:* Magdalena Reizer (magdalena.reizer@pw.edu.pl)

Supplementary Material




Figure S1: Hourly concentrations (ng m^{-3}) of As measured in the fine (green) and coarse (red) fractions.

15





Figure S2: Hourly concentrations (ng m^{-3}) of Ba measured in the fine (green) and coarse (red) fractions.



20 Figure S3: Hourly concentrations (ng m^{-3}) of Br measured in the fine (green) and coarse (red) fractions.

25




Figure S4: Hourly concentrations (ng m^{-3}) of Ca measured in the fine (green) and coarse (red) fractions.

30




Figure S5: Hourly concentrations (ng m^{-3}) of Cu measured in the fine (green) and coarse (red) fractions.

35




Figure S6: Hourly concentrations (ng m^{-3}) of Fe measured in the fine (green) and coarse (red) fractions.




Figure S7: Hourly concentrations (ng m^{-3}) of Mg measured in the fine (green) and coarse (red) fractions.




Figure S8: Hourly concentrations (ng m^{-3}) of Mn measured in the fine (green) and coarse (red) fractions.





Figure S9: Hourly concentrations (ng m^{-3}) of Mo measured in the fine (green) and coarse (red) fractions.

40

45



50 Figure S10: Hourly concentrations (ng m^{-3}) of Na measured in the fine (green) and coarse (red) fractions.






Figure S11: Hourly concentrations (ng m^{-3}) of Ni measured in the fine (green) and coarse (red) fractions.




55 Figure S12: Hourly concentrations (ng m^{-3}) of P measured in the fine (green) and coarse (red) fractions.



60

Figure S13: Hourly concentrations (ng m^{-3}) of Pb measured in the fine (green) and coarse (red) fractions.



65

Figure S14: Hourly concentrations (ng m^{-3}) of Rb measured in the fine (green) and coarse (red) fractions.




Figure S15: Hourly concentrations (ng m^{-3}) of S measured in the fine (green) and coarse (red) fractions.

70





Figure S16: Hourly concentrations (ng m⁻³) of Se measured in the fine (green) and coarse (red) fractions.



75 Figure S17: Hourly concentrations (ng m⁻³) of Si measured in the fine (green) and coarse (red) fractions.




Figure S18: Hourly concentrations (ng m⁻³) of Sr measured in the fine (green) and coarse (red) fractions.





Figure S19: Hourly concentrations (ng m^{-3}) of Ti measured in the fine (green) and coarse (red) fractions.



85

Figure S20: Hourly concentrations (ng m^{-3}) of V measured in the fine (green) and coarse (red) fractions.




Figure S21: Hourly concentrations (ng m^{-3}) of Y measured in the fine (green) and coarse (red) fractions.

90





Figure S22: Hourly concentrations (ng m^{-3}) of Zn measured in the fine (green) and coarse (red) fractions.



95 Figure S23: Hourly concentrations (ng m^{-3}) of Zr measured in the fine (green) and coarse (red) fractions.