

Supplement to: Modelling the size distribution of
aggregated volcanic ash and implications for operational
atmospheric dispersion modelling'

Frances Beckett¹, Eduardo Rossi², Benjamin Devenish¹, Claire Witham¹, and
Costanza Bonadonna²

¹Met Office, Exeter, UK

²University of Geneva, Switzerland




Figure S1: Sensitivity of the output aggregated GSD to the sticking efficiency parameters (a) St_{cr} , (b) q , and the physical characteristics assigned to the particles, (c) particle density (ρ_s) and (d) input GSD. Output is for 12:00 UTC on the 05/05/2010, plume height 5500 m asl.




Figure S2: Sensitivity of the output aggregated GSD to the sticking efficiency parameters (a) St_{cr} , (b) q , and the physical characteristics assigned to the particles, (c) particle density (ρ_s) and (d) input GSD. Output is for 13:00 UTC on the 06/05/2010, plume height 10000 m asl.




Figure S3: Sensitivity of the output aggregated GSD to the sticking efficiency parameters (a) St_{cr} , (b) q , and the physical characteristics assigned to the particles, (c) particle density (ρ_s) and (d) input GSD. Output is for 12:00 UTC on the 07/05/2010, plume height 5500 m asl.




Figure S4: Sensitivity of the output aggregated GSD to the entrainment coefficients (a) k_s , (b) k_n , the fraction of dry air (c) n_d and water vapour (d) n_v , the temperature of the plume at the source (e) T_0 and the source mass flux (f) Q_m . Output is for 19:00 UTC on the 04/05/2010, plume height 7000 m asl.




Figure S5: Sensitivity of the output aggregated GSD to the entrainment coefficients (a) k_s , (b) k_n , the fraction of dry air (c) n_d and water vapour (d) n_v , the temperature of the plume at the source (e) T_0 and the source mass flux (f) Q_m . Output is for 12:00 UTC on the 05/05/2010, plume height 5500 m asl.




Figure S6: Sensitivity of the output aggregated GSD to the entrainment coefficients (a) k_s , (b) k_r , the fraction of dry air (c) n_d and water vapour (d) n_v , the temperature of the plume at the source (e) T_0 and the source mass flux (f) Q_m . Output is for 13:00 UTC on the 06/05/2010, plume height 10000 m asl.




Figure S7: Sensitivity of the output aggregated GSD to the entrainment coefficients (a) k_s , (b) k_n , the fraction of dry air (c) n_d and water vapour (d) n_v , the temperature of the plume at the source (e) T_0 and the source mass flux (f) Q_m . Output is for 12:00 UTC on the 07/05/2010, plume height 5500 m asl.