



Climate impact of volcanic eruptions: the sensitivity to eruption season and latitude in MPI-ESM ensemble experiments

Zhihong Zhuo^{1, a}, Ingo Kirchner¹, Stephan Pfahl¹, Ulrich Cubasch¹

¹Institute of Meteorology, Freie Universität Berlin, Berlin, 12165, Germany

5 ^anow at: Section for Meteorology and Oceanography, Department of Geosciences, University of Oslo, Oslo, 0315, Norway

Correspondence to: Zhihong Zhuo (zhihong.zhuo@met.fu-berlin.de)

Abstract. Explosive volcanic eruptions influence near-surface temperature and precipitation especially in the monsoon regions, but the impact varies with different eruption seasons and latitudes. To study this variability, two groups of ensemble simulations are performed with volcanic eruptions in June and December at 0° representing an equatorial eruption (EQ) and
10 at 30° N and 30° S representing northern and southern hemisphere eruptions (NH and SH). Results show significant cooling especially in areas with enhanced volcanic aerosol content. Stronger cooling emerges in the northern (southern) hemisphere after the NH (SH) eruption compared to the EQ eruption. Stronger precipitation variations occur in the tropics than in the high latitudes. Summer and winter eruptions lead to similar climate impacts. The NH and the SH eruptions have reversed
15 climate impacts, especially in the South Asian monsoon regions. After the NH (SH) eruption, direct radiative effects of volcanic aerosols induce changes in the interhemispheric and land-sea thermal contrasts, which move the intertropical convergence zone southward (northward) and weaken (strengthen) the South Asian summer monsoon. This reduces
20 (increases) the moisture transport from the ocean to India, and reduces (enhances) cloud formation. The subsequent radiative feedbacks due to regional cloud cover lead to warming (cooling) in India. This emphasizes the sensitivity of regional climate impacts of volcanic eruptions to eruption latitude, which relates to the dynamical response of the climate system to radiative effects of volcanic aerosols and the subsequent regional physical feedbacks.

1 Introduction

Sulfate aerosols in the stratosphere from explosive volcanic eruptions significantly cool the surface by reflecting incoming solar radiation (Robock et al., 2000; Timmreck, 2012). This further affects the earth's hydrological cycle. Both observations and model simulations indicate a reduction of global precipitation after volcanic eruptions (Gillett et al., 2004; Iles and
25 Hegerl, 2014; Iles et al., 2013; Paik and Min, 2016; Robock and Liu, 1994; Trenberth and Dai, 2007). Precipitation was found to largely decrease in tropical areas (Colose et al., 2016; Iles and Hegerl, 2014), and also notably in summer monsoon regions (Liu et al., 2016; Man et al., 2014; Oman et al., 2006; Schneider et al., 2009; Zambri and Robock, 2016; Zhuo et al., 2020). However, Toohey et al. (2019) and Yang et al. (2019) pointed out that the spatial distribution of volcanic aerosols and associated radiative forcing affects the climate impact of volcanic eruptions. The precipitation response varies with changing



30 distributions of volcanic aerosols between the hemispheres. Inverse effects of interhemispherically asymmetric volcanic
aerosols were found in Sahelian precipitation (Haywood et al., 2013; Jacobson et al., 2020), monsoon climate (Iles and
Hegerl, 2014; Liu et al., 2016; Zhuo et al., 2014) and tropical hydroclimate in general (Colose et al., 2016; Zuo et al., 2018).
Furthermore, Stevenson et al. (2016) suggested that the eruption season has to be considered when studying the climate
35 impact of volcanic eruptions. Based on volcanic forcing reconstruction indices, most of these previous studies separated
historical eruptions into volcanic classifications with different numbers of events, eruption seasons and unequally distributed
aerosol magnitudes in different hemispheres. In order to study the sensitivity of the climate impact of volcanic eruptions to
season and location (tropical vs. Northern or Southern Hemisphere) in a systematic way, in this study we thus perform model
experiments with artificial volcanic eruptions.

Tropical eruptions have been thought to have larger climate impact than extratropical or high-latitude eruptions (Myhre et al.,
40 2013; Schneider et al., 2009). The volcanic aerosols injected into the stratosphere from a tropical eruption was believed to be
transported to both hemispheres and finally reach both poles. This generates larger areas being affected by aerosols with long
lifetimes in the stratosphere, thus causing larger and longer-lasting climate effects (Schneider et al., 2009). Based on this
hypothesis, previous reconstructions of volcanic forcing assumed a longer lifetime of aerosols injected by tropical volcanic
eruptions than by extratropical eruptions (Ammann and Naveau, 2003; Gao et al., 2008). Thus, compared to tropical
45 eruptions, model results suggested shorter cooling periods and less severe reduction of tropical precipitation after high-
latitude eruptions (Schneider et al., 2009). On the contrary, using both ice-core and tree ring-based proxy reconstruction and
model simulation, Toohey et al. (2019) found larger climate impact after extratropical eruptions than tropical eruptions. They
pointed out that the overestimated volcanic forcing from tropical eruptions results in an overestimation of their climate
impacts. Whether tropical or extratropical volcanic eruptions have larger climate impact, is still a controversial question.

50 Only few studies investigated the mechanism behind the precipitation response to volcanic eruptions (Man and Zhou, 2014;
Man et al., 2012, 2014; Oman et al., 2006; Paik and Min, 2016). Paik and Min (2016) argued that a reduced vertical motion
after volcanic eruption induces the reduction of global precipitation. The reduction of the summer monsoon precipitation was
suggested to be due to a decreased land-sea thermal contrast and the subsequent weakening of the summer monsoon (Man
and Zhou, 2014; Man et al., 2012, 2014; Oman et al., 2006). Conclusions in these studies did not consider regional
55 differences and feedback processes in the Indian monsoon region (Oman et al., 2006; Paik and Min, 2016). The inversed
climate impact after interhemispherically asymmetric volcanic aerosol injections was suggested to result from a
displacement of the ITCZ towards the warmer hemisphere with less volcanic aerosol loading (Colose et al., 2016; Haywood
et al., 2013; Iles and Hegerl, 2014; Zuo et al., 2018). Here, we study the mechanisms behind the regional climate response to
volcanic eruptions at different latitudes and in different seasons, focusing on the Indian monsoon region, with the help of a
60 series of model simulations. This study aims at answering the following questions: How are global and regional climate
impacts of volcanic eruptions affected by the eruption season and latitude? Do tropical eruptions have larger climate impact
than extratropical eruptions? What is the mechanism behind the Indian monsoon response to volcanic eruptions at different
latitudes?



65 After this introduction, we describe the methods including model description and experimental setup in section 2. In section 3, we present our results and discussion. Summary and conclusions are given in section 4.

2 Method

2.1 Model description

70 We perform simulations with the Max-Planck-Institute Earth System Model (MPI-ESM) (Giorgetta et al., 2013). The MPI-ESM is a fully coupled general circulation model with ECHAM6 as atmosphere component (Stevens et al., 2013), MPIOM as ocean component (Jungclaus et al., 2013), JSBACH for simulating the terrestrial biosphere (Reick et al., 2013; Schneek et al., 2013) and HAMOCC5 for the biogeochemistry of the ocean (Ilyina et al., 2013). The components are coupled through the exchange of energy, momentum, water and carbon dioxide using the OASIS3 coupler (Valcke, 2013). In this study, we use the low-resolution (LR) configuration of the model. In the atmospheric component, it has a horizontal resolution of T63 with 47 vertical levels extending to 0.01 hPa, thus including the stratosphere. The oceanic component has a resolution of 1.5° (near the equator) with 40 vertical levels.

75 Also because of its high simulation efficiency (Roeckner et al., 2006), the older MPI-ESM version with ECHAM5 as its atmospheric component has been extensively used to study the Asian summer monsoon (Guo et al., 2016; Man et al., 2012) and the climate response to volcanic eruptions (Man et al., 2014; Zhang et al., 2012). Studies on the impact of volcanic eruptions have also been performed with the newest model version that is also applied here (Timmreck et al., 2016; Toohey et al., 2016).

80 We use the same configuration of the MPI-ESM model as used for the historical simulation of CMIP6. The volcanic forcing used in this study is produced by the easy volcanic aerosol (EVA) module (Toohey et al., 2016). Through setting the eruption year, month, sulfate injection magnitude and hemispheric ratio in EVA, stratospheric sulfate mass injected by the volcanic eruption is transferred to mid-visible (550 nm) aerosol optical depth (AOD) and effective radius as volcanic forcing input for model simulations. Pinatubo-like eruptions are simulated because the construction of EVA relies extensively on observational constraints, especially the observational records of the 1991 Pinatubo eruption. Other external forcings for MPI-ESM are the same as in the CMIP6 historical experiment, except that the anthropogenic emissions are fixed at the 1900 level.

2.2 Experimental setup

90 In order to study the climate impacts of volcanic eruptions at different eruption latitudes and in different seasons, two groups of three different latitudinal volcanic eruptions in summer and winter are simulated (here summer and winter refer to the boreal seasons). We perform 10-member simulations for each eruption case. Following Toohey et al. (2016), 9 Tg of total sulfur injection magnitude is prescribed. The eruption latitudes are set to be 0° for the equatorial eruption (EQ case) and 30° N and 30° S for the northern and southern hemispheric eruptions (NH and SH cases), respectively. For the summer eruptions,



95 the date is set to be the same as the 1991 Pinatubo eruption on June 15, 1991; for the winter eruptions, the date is set to be December 15, 1991.

Figure 1 shows the experimental design of our simulations. An initial run is performed for the period of 1966-2012, and then we perform 23 control runs for the period of 1986-1995 without any volcanic eruption. For these 23 members, we calculate the Oceanic Niño Index (ONI) to quantify El Niño-Southern Oscillation (ENSO) variability. ONI values between -1 and 1
100 represent neutral or weak ENSO conditions. The 1991 Pinatubo eruption was accompanied by a strong El Niño event in 1991-1994. The response of ENSO to volcanic eruptions has been widely investigated (Adams et al., 2003; Khodri et al., 2017; Stevenson et al., 2016) but remains to be an unsolved question. To rule out the concurrent effect of ENSO, we pick out 10 control runs with neutral or weak ENSO in the period of 1990-1992 as a basis for our experiments. Restart files from these 10 control runs are used to initialize our three 10-member ensemble simulations for the summer (winter) EQ, NH and
105 SH eruptions in the period of 1991-1996 (1990-1996). Note that the simulated years do not correspond to the real years because of the free-running spin-up simulations.

2.3 Analysis methods

To quantify the Indian summer monsoon response, we calculate the south Asian summer monsoon index (SASMI), which is defined as the difference between the zonal wind at 850 hPa over the region 0°-20°N, 40°E-110°E and the zonal wind at 200
110 hPa over the region 0°-20°N, 40°E-110°E (Webster and Yang, 1992).

To study horizontal moisture transport, we calculate the vertically integrated moisture transport (IVT) and its divergence (IVTD). The IVT is calculated using the following equation:

$$(1/g) \int_{surface}^{model\ top} q \vec{v} dp$$

where g is the acceleration due to gravity, q is specific humidity, \vec{v} is the horizontal wind vector, and p is pressure. The
115 vertical integration of the equation is performed from the surface to the model top.

We calculate the multi-member mean (MEM) of adopted variables and the indices defined above to study the hydrological effect of the summer and winter volcanic eruptions at different latitudes. Both temporal and spatial analyses are performed to investigate the climate effects. Anomalies with respect to the MEM of the control runs without any volcanic eruption are presented. The standard deviation (SD) of the control runs is calculated to indicate the significance of the temporal analysis
120 results. Two-tailed student-t tests at the 95% and 99% confidence levels are performed for testing the significance of the spatial results. EVA and MPI-ESM model simulations of volcanic eruptions were tuned and validated largely based on observations of the Pinatubo eruption in June 1991, thus our discussions focus more on the summer eruptions.



3 Results and discussion

125 3.1 Volcanic forcing from EVA

The generated volcanic forcing is indicated by the aerosol optical depth at 550 nm (AOD_{550}). Figure 2a (2b) shows the temporal variation of global mean AOD_{550} in 1991-1996 (1990-1996) for the summer (winter) eruptions. All three eruption cases in each season lead to identical time series of global mean AOD_{550} , with a steep increase in the beginning due to the formation of sulfate aerosols after the volcanic eruptions. The highest global mean AOD_{550} of 0.12 is identical in all the cases, and is reached after several months in December 1991 (June 1991) for the summer (winter) eruptions. Figure 2c and Fig. 2d show the spatio-temporal structure of the volcanic forcing. The formation and distribution of volcanic aerosols take several months. For the EQ eruption cases, the AOD_{550} indicates that volcanic aerosols are transported to both hemispheres, associated with a stronger dispersion than in the NH and SH cases, and more aerosols are transported to the northern hemisphere in the winter case and to the southern hemisphere in the summer case. This indicates that the transport of volcanic aerosols from equatorial eruptions to high latitudes depends on the eruption season, which is related to the atmospheric circulation in the tropical stratosphere. For the NH and SH eruption cases, volcanic aerosols are mostly confined to the specific hemisphere, with large AOD_{550} in the subtropical areas between 30° and 60° latitude.

3.2 Solar radiation response

To show the primary radiative effects of the volcanic aerosols, we present the net outgoing shortwave radiation (OSR) at the top of the atmosphere (TOA) in clear-sky conditions. Figure 3 shows the time series of the global mean OSR anomaly and its zonal mean distribution. The OSR value in the MPI-ESM is negative, indicating an increased OSR after the volcanic eruptions. For all three cases of summer eruptions (Fig. 3a), the increased OSR lasts for three years until 1994. Although the global mean AOD_{550} is identical among the three cases (Fig. 1a), the OSR increases continuously after the SH and the EQ eruptions, but after the NH eruption, there is a slight decrease of OSR in the boreal winter of 1991. Similar response patterns are found for the winter eruption cases (Fig. 3b), except that the slight decrease of OSR occurs after the SH eruption case in the boreal summer of 1992, which is the austral winter of the southern hemisphere. This relates to the seasonal change of the incoming solar radiation in the different hemispheres. The zonal mean distribution of the OSR anomaly in all summer cases (Fig. 3c) and winter cases (Fig. 3d) indicates a simultaneous increase of OSR in the areas with high AOD after the eruption. This indicates that more solar radiation is reflected in regions with more volcanic aerosols. For the summer eruption cases (Fig. 3c), the shape of the OSR zonal mean distribution is similar to that of AOD_{550} in the EQ and SH cases, while OSR is reduced in the NH case in the boreal winter of 1991, resulting from the reduced incoming radiation in winter. For the winter eruption cases (Fig. 3d), similar shapes between OSR and AOD_{550} are shown in the EQ and NH cases, while a reduced OSR in the austral winter of 1991 is found in the SH case. This confirms the role of the seasonal change of the incoming solar radiation in the two hemispheres.



155 3.3 Temperature response

SR at top of the atmosphere is reflected by the volcanic aerosols. This decreases the SR reaching the surface and results in surface cooling. Figure 4 shows the time series of global mean surface temperature (T) anomaly and its zonal mean distribution. Significant cooling occurs after the volcanic eruptions. For the summer eruption cases, the largest cooling occurs in the boreal summer of 1992 (Fig. 4a) and most significantly in the northern hemisphere mid-latitudes in the NH case, in the tropics in the EQ case and in the southern hemisphere mid-latitude areas in the SH case, as indicated by the stippling in Fig. 4c. For the winter eruption cases, a significant and strong cooling is visible in the boreal summer of 1991 in the NH case (Fig. 4b), the largest cooling emerges in the northern hemisphere mid-latitude areas (Fig. 4d). Slight cooling occurs in the EQ case especially in the boreal spring of 1992 (Fig. 4b), and the abnormal cooling is pronounced in the tropics as indicated by the stippling in Fig. 4d. A similar cooling trend with a smaller magnitude is shown in the SH case (Fig. 4b), but the cooling is significant in the southern hemisphere mid-latitudes (Fig. 4d). For both the summer and winter eruptions, the largest cooling occurs several months later than the solar radiation variation but in the corresponding areas with more reflected SR by volcanic aerosols, and the response is faster in the NH cases than in the SH and EQ cases. These different responses originate from the delayed response in the ocean compared to the land, because the ocean has a larger heat capacity and content compared to the land and the NH has a larger landmass compared to the SH.

170 We further analyze the spatial distribution of the temperature response. The time series and zonal mean distribution have shown that the strongest cooling occurs in summer 1992 for the summer eruptions and in summer 1991 for the winter NH eruption and spring 1992 for the winter EQ and SH eruptions. Since we want to discuss more about the summer climate response especially in the northern hemisphere, for the spatial distribution of temperature, we focus on the JJA mean of 1992 (1991) for the summer (winter) eruptions (Fig. 5). For the NH summer eruption case (Fig. 5a), significant cooling (99% confidence level) is found for most of the land areas in the northern hemisphere, except for India and the Sahel, where positive temperature anomalies prevail. Stronger cooling occurs in the northern hemisphere than in the southern hemisphere. Over the ocean, most of the areas experience cooling (warming) in the northern (southern) hemisphere. While similar patterns with stronger magnitude are found especially in the northern hemisphere for the NH winter eruption case (Fig. 5b), different responses are shown over the southern hemispheric ocean and Antarctica. For the SH summer eruption case (Fig. 5e), significant cooling (99% confidence level) is found in the southern hemispheric land areas, but warming emerges in part of the northern hemispheric land areas. Opposite to the NH eruption case, cooling occurs in Indian and the Sahel. Over the ocean, cooling prevails in the low latitude areas while different warming and cooling emerges in different areas in mid and high latitudes. Similar response patterns are found in most of the southern hemispheric land areas in the SH winter eruption case, but warming prevails in most of the northern hemispheric land areas except for the cooling in India and the Sahel (Fig. 5f). For the EQ cases (Fig. 5c and 5d), cooling occurs in most of the land areas, but weaker in magnitude in the northern (southern) hemisphere compared to the NH (SH) eruption cases. This indicates that, compared to tropical eruptions, extratropical volcanic eruptions have stronger cooling effects in the hemisphere where the eruption occurs due to



concentration of volcanic aerosols. Furthermore, the responses in the low latitudinal land areas in the EQ summer (winter) eruption case (Fig. 5c and 5d) are similar to those in the SH summer (winter) eruption case (Fig. 5e and 5f).

190 3.4 Precipitation response

The radiative effect of volcanic aerosols, in addition to temperature variations, also leads to changes in the hydrological cycle. Time series of global mean precipitation from our experiments do not show significant changes (not shown), but there are precipitation responses on a regional level. Figure 6 shows the spatial distribution of the precipitation response to the volcanic eruption cases. Generally, stronger absolute precipitation responses occur in the tropics than in the extratropics as well as over the ocean compared to over land. For each eruption latitude, the precipitation response patterns are similar
195 well as over the ocean compared to over land. For each eruption latitude, the precipitation response patterns are similar between the summer eruptions (Fig. 6 left panel) and the winter eruptions (Fig. 6 right panel). In many tropical regions, the NH (Fig. 6a and 6b) and SH (Fig. 6e and 6f) volcanic eruptions lead to reversed precipitation responses. Precipitation response patterns in the EQ eruption cases (Fig. 6c and 6d) are close to those in the SH eruption cases (Fig. 6e and 6f). The most pronounced changes over land are found in India (black box in Fig.6), which is one of the most typical monsoon
200 regions. Precipitation is reduced in India after the NH eruptions (Fig. 6a and 6b), but strongly increased after the EQ and SH volcanic eruptions (Fig. 6c to 6f).

3.5 Mechanism of precipitation response in India

The temperature and precipitation response patterns clearly indicate a tendency towards inverse climate responses to the NH and SH volcanic eruptions, which are particularly strong in India. The following section thus aims at explaining the
205 mechanism behind the different climate effects of asymmetrically distributed volcanic aerosols in India.

3.5.1 The direct radiative effect and dynamical response

Sulfate aerosols in the stratosphere from explosive volcanic eruptions reflect solar radiation at the top of the atmosphere (Fig. 3), thus leading to significant surface cooling (Fig. 4). These cooling effects are more pronounced in the areas with more aerosols. Volcanic aerosols are asymmetrically distributed after the different latitudinal volcanic eruptions (Fig. 1). This
210 causes asymmetric cooling effects between the two hemispheres. Figure 7a shows the difference of the boreal summer mean temperature anomaly between the northern and the southern hemisphere in the three summer eruption cases. It decreases largely (slightly) in 1992 after the NH (EQ) volcanic eruptions, while it keeps close to zero after the SH volcanic eruption. This indicates a larger cooling in the northern hemisphere than in the southern hemisphere after the NH and the EQ volcanic eruption, while only a slight difference is shown between the two hemispheres after the SH volcanic eruption. Similar results
215 are found in the winter NH and EQ eruption cases, except for that in the SH case, the temperature difference between the hemispheres increases slightly, indicating a larger cooling in the southern hemisphere than in the northern hemisphere after the winter SH volcanic eruption (Fig. 7b).

The thermal contrast between the two hemispheres moves the ITCZ away from the cooler hemisphere (Broccoli et al., 2006).



As shown by the yellow line in Fig 7c. and Fig 7d, in boreal summer, large amounts of precipitation are concentrated around
220 10°N, which is the mean latitude of the ITCZ. After the NH summer volcanic eruption (Fig. 7c), the zonal mean precipitation
decreases around 10°N but increases around 0°. This indicates that the ITCZ moves southward toward the equator. The zonal
mean precipitation increases north of 10°N but decreases around the equator after the SH summer eruption, which indicates a
northward movement of the ITCZ. For the EQ summer eruption case, because the aerosols are transported to both
225 hemispheres, there is no simple displacement of the ITCZ, but a more complex change with a precipitation decrease around
10°N but increases to the north and south of 10°N. Similarly, a southward (northward) shift of the ITCZ is found after the
NH (SH) winter eruption (Fig. 7d). For the EQ winter eruption case (Fig. 7d), the zonal mean precipitation slightly increases
around 10°N, indicating a slight northward movement of the ITCZ. The movement of the ITCZ particularly affects the
climate in India.

Due to the large heat content of the ocean, the temperature response over the ocean is attenuated compared to the response
230 over land. This leads to a stronger cooling of the land compared to the ocean, which decreases the land-ocean thermal
gradient in summer and leads to a weakening of the summer monsoons (Dogar and Sato, 2019; Iles and Hegerl, 2014; Man
and Zhou, 2014; Zuo et al., 2019). Here, we quantify the Indian summer monsoon response to volcanic eruptions with the
SASMI (Webster and Yang, 1992). As shown in Fig. 8a, the SASMI decreases continuously until 1994 after the NH summer
eruption, while it increases strongly (slightly) in 1992 after the SH (EQ) summer eruption. Similarly, in Fig. 8b, a significant
235 decrease of the SASMI is found in 1991 after the NH winter eruption, while a slight increase of the SASMI occurs in 1991
after the SH and EQ winter eruptions. The SASMI keeps increasing in 1992 after the SH winter eruption. This indicates
opposite weakening and strengthening effects of the different NH and SH volcanic eruptions on the south Asian summer
monsoon: NH eruptions preferentially cool the NH land regions and tend to weaken the Indian monsoon, while SH cooling
following SH eruptions leads to a stronger monsoon circulation.

240 The altered land-ocean thermal contrast and monsoon circulation largely change the moisture transport from the ocean to
India. Since the temperature and precipitation responses, the ITCZ movement and the SASMI all show similar patterns
between the summer eruption cases and the winter eruption cases, we only use the summer eruption cases to explain the
response mechanism in India in the following. Figure 9 shows the vertically integrated moisture transport (IVT, vector) and
its divergence (IVTD, shaded) over India. In the control runs without any volcanic eruption, as shown in Fig. 9a, in the
245 western part of India, winds from the Arabian Sea bring large amounts of moisture to the land. This results in strong
precipitation along the west coast of India. In the eastern and northern parts of India, the precipitation results from the
convergence of southeasterly moisture transported from the Bay of Bengal vortex as shown by the IVTD (green shade).
After the NH volcanic eruption, the moisture transported from the Bay of Bengal to the northern part of India is largely
reduced, associated with reduced convection and upward motion, as shown by the reduced convergence (brown shade in Fig.
250 9b). On the contrary, after the EQ and the SH volcanic eruptions (Fig. 9c and 9d), more moisture is transported to the land
from both the Arabian Sea and the Bay of Bengal. Moisture convergence is reduced in the northern part of India (brown
shade), while it is strengthened in the southwest and southeast coast of India (green shade). The altered horizontal and



vertical motion of the atmospheric circulation thus change the amount and distribution of atmospheric moisture as well as the precipitation patterns over India.

255 These results show a dynamical response of the climate system to the radiative effect of volcanic aerosols. Changes in the available energy and thermal gradients between the hemispheres move the ITCZ southward (northward) after the NH (SH and EQ) volcanic eruptions (Fig. 7b), which largely affects regional precipitation in the areas along the ITCZ. The altered land-ocean thermal gradients reduce (strengthen) the monsoon circulation and thus the moisture transport and convection over the western coast of India and central India after the NH (SH and EQ) volcanic eruptions. These dynamical responses influence the regional climate in India and lead to a precipitation decrease after NH eruptions, but an increase following SH 260 eruptions (see again Fig. 6).

3.5.2 Radiative feedbacks due to changes in cloud cover

In the following, physical feedback mechanisms associated with changes in cloud cover and the corresponding radiative effects are discussed, which additionally influence the regional temperature and precipitation variations. Figure 10 shows the net surface solar radiation distribution in clear-sky and all-sky conditions. In clear-sky conditions without taking clouds into 265 consideration (Fig. 10a), the surface solar radiation is generally reduced with a zonally rather symmetric distribution. The decrease of the surface solar radiation is larger in the northern (southern) part of the area after the NH (SH) volcanic eruptions, while it decreases more uniformly after the EQ eruption. This is due to the reflection of solar radiation by the asymmetric distribution of volcanic aerosols in the two hemispheres after the NH and the SH volcanic eruptions, while being approximately balanced between the two hemispheres after the EQ eruption. However, there are some stronger and opposite 270 regional changes in the all-sky net surface solar radiation (Fig. 10b). Specifically, in India, the net surface solar radiation increases after the NH volcanic eruption, but decreases after the SH and the EQ volcanic eruptions. This indicates that in the regional climate responses to volcanic eruptions, the change of the regional cloud cover plays an important role, leading to a temperature increase (decrease) in India after the NH (the SH and the EQ) volcanic eruption (Fig. 5), which is opposite to the temperature changes in most other northern hemispheric land regions. As shown in Fig. 11, cloud cover decreases 275 significantly after the NH volcanic eruption (Fig. 11a), while it increases after the SH and the EQ volcanic eruptions (Fig. 11c and 11b). Less (more) SR is reflected with less (more) clouds, thus leading to the increase (decrease) of net surface solar radiation and temperature after the NH (the SH and the EQ) volcanic eruptions.

The dynamical response described in the previous section leads to a reduced (increased) moisture transport from the ocean to 280 the land, and that the corresponding decrease (increase) in the atmospheric water vapor content (Fig. 12) together with the reduced (strengthened) vertical motion reduce (enhance) cloud formation and precipitation in India after the NH (the SH and the EQ) volcanic eruption.



3.5.3 Summary of the response mechanism

285 Based on the above analyses, the mechanisms behind the different climate effects in India after volcanic eruptions at
different latitudes are summarized. The stratospheric aerosols injected by volcanic eruptions directly reflect large amount of
solar radiation, which decreases the net surface solar radiation. This leads to cooling in the latitude bands covered by
volcanic aerosols. It alters the thermal gradient between the northern and the southern hemisphere and between land and
ocean. These changed gradients of available energy and temperature cause subsequent dynamical responses. Specifically, the
290 altered thermal contrast between the two hemispheres moves the ITCZ towards the warmer hemisphere, and the altered land-
ocean thermal contrast changes the strength of the South Asian summer monsoon and the associated horizontal and vertical
motion of the air. This influences the regional water vapor content and the subsequent formation of clouds. Regional
temperature and precipitation are further affected by this change in moisture content and by radiative feedback processes due
to the altered cloud cover.

After the NH (the SH) volcanic eruption, the altered hemispheric thermal contrast leads to a southward (northward)
295 movement of ITCZ; the decreased land-ocean thermal contrast weakens (strengthens) the SASM and reduces (strengthens)
the horizontal and vertical motion of the air in India. This reduces (increases) the atmospheric moisture content and
decreases (increases) cloud cover, which results in a decrease (increase) of the regional precipitation. The reduced (increased)
cloud cover counteracts (strengthens) the cooling effect of volcanic aerosols. This negative (positive) feedback leads to the
increase (decrease) of the regional temperature. For the EQ eruption, evenly distributed volcanic aerosols in the two
300 hemispheres cause a relatively symmetric radiative effect in India. However, because the subsequent dynamical response is
similar to that after the SH volcanic eruption, the temperature and precipitation patterns are also similar to those after the SH
eruption. This illustrates the important roles that dynamical responses and subsequent physical feedbacks play for the
regional climate response to volcanic eruptions.

3.6 Discussion

305 Our model results indicate significant cooling and precipitation variations especially in the tropical and monsoon regions
after volcanic eruptions, in agreement with previous studies (Iles and Hegerl, 2014; Iles et al., 2013). With CESM model
simulations, Stevenson et al. (2017) showed different soil moisture responses to volcanic eruptions in April and July.
However, temperature and precipitation responses in this study do not show large difference between the summer and winter
eruptions. A different model is used, and different eruption seasons are simulated in our study compared to Stevenson et al.
310 (2017). Hence, the question on how the eruption season affects the climate impact of volcanic eruptions still needs to be
further investigated.

Tropical eruptions were widely assumed to have stronger climate impact than extratropical eruptions (Schneider et al., 2009),
until Toohey et al. (2019) pointed out that this might not be the case. Based on the atmosphere-chemistry model
MAECHAM5-HAM with fixed sea surface temperature, Toohey et al. (2019) showed a stronger hemispheric cooling after



315 extratropical explosive volcanic eruptions than after tropical eruptions. Results in this study, based on a fully coupled
atmosphere-ocean model, show stronger cooling in the relative hemisphere after the NH and SH eruptions than for the EQ
eruptions, supporting the conclusion of Toohey et al. (2019).
Results in this study agree with previous investigations regarding the different climate impacts that interhemispherically
asymmetric volcanic aerosol distributions may have in the tropics (Colose et al., 2016; Zuo et al., 2018), the monsoon
320 regions (Liu et al., 2016; Zhuo et al., 2014; Zuo et al., 2019) and the Atlantic (Yang et al., 2019). These differences are likely
caused by the movement of the ITCZ towards the warmer hemisphere with less volcanic aerosol loading (Colose et al., 2016;
Haywood et al., 2013; Iles and Hegerl, 2014) and the weakening of monsoonal circulations (Liu et al., 2016; Zuo et al.,
2018). Our study confirms these general conclusions in a quantitative way and provides detailed insights into the underlying
mechanisms based on model experiments that have been specifically designed to distinguish between NH and SH eruptions.
325 For the mechanism of the monsoon response to asymmetric volcanic aerosols, Zuo et al. (2019) suggested that the change of
the atmospheric circulation plays a dominant role in the change of precipitation, which is related to the changes of the
monsoon circulation and the cross-equator flow. Earlier, using the GISS model, Oman et al. (2005) simulated the climate
effects of high-latitude eruptions, and found that the radiative effect is larger than the dynamical effect, and the dynamical
effect mainly affects the Asian summer, as the strong cooling in northern hemisphere landmass leads to the reduction of the
330 Asian summer monsoon circulation. This study confirms that the atmospheric circulation change after volcanic eruptions
plays an important role in precipitation changes in India, because it changes the water vapor transport and the resultant
formation or depletion of clouds in different areas. In addition, regional temperature and precipitation variations are also
affected by radiative cloud feedbacks. This is in agreement with Dogar and Sato (2019), who suggested that the direct
radiative effect of tropical volcanic eruption and the associated land-sea thermal contrast result in warming and drying in the
335 Middle Eastern, African and south Asian monsoon regions, which is related to the reduction of clouds over the monsoon
region. However, the indirect circulation change was summarized to be connected with the volcanic-induced ENSO forcing
in Dogar and Sato (2019). This conclusion might be affected by large uncertainties, as only three ensemble members were
used in their study, and the simulations were conducted with an atmospheric circulation model with prescribed oceanic
boundary conditions. Our results, based on a fully coupled atmosphere-ocean model and specifically designed to be not
340 affected by ENSO anomalies at the time of the eruption, show that both the NH and the EQ volcanic eruptions favor an El
Niño-tendency in the summer of 1992, while a La Niña-tendency is found after the SH volcanic eruption (both statistically
not significant; not shown). However, after EQ volcanic eruptions, the climate response in India is similar to that after SH
eruptions (Fig. 5 and Fig. 6). This suggests that it is the direct circulation change and the subsequent physical feedback, not
so much the influence of ENSO, that dominates the regional climate impact in India.



345 4 Summary and conclusions

In order to explore how eruption latitude and eruption season affect the climate impact of volcanic eruptions, model simulations with two groups of Pinatubo-like summer and winter volcanic eruptions at 0° (EQ), 30°N (NH) and 30°S (SH) are performed based on the EVA volcanic forcing generator and the fully coupled general circulation model MPI-ESM. For each experiment case, an ensemble of 10 simulations is performed. Based on the experimental design, results in this study
350 avoid the uncertainties from unequal aerosol magnitudes among different volcanic events and the impact of concurrent ENSO events.

Results confirm that stratospheric aerosols injected by volcanic eruptions reflect incoming solar radiation and reduce the net surface solar radiation. This leads to significant surface cooling, especially in the areas covered by volcanic aerosols.

Comparing to equatorial eruptions, volcanic aerosols injected by extratropical eruptions are more concentrated in the specific
355 hemisphere. Because of this, extratropical eruptions in the northern (the southern) hemisphere (NH and SH cases) cause larger cooling over the northern (southern) hemisphere continents compared to the equatorial eruption (EQ case). The precipitation response varies regionally, with stronger responses in the tropic than in the extratropic. Summer eruptions and winter eruptions lead to similar response patterns, especially in the tropics and monsoon regions, which indicates that the eruption season plays a minor role for the climate impact of volcanic eruptions. The largest precipitation change over land
360 occurs in India, one of the most typical monsoon regions. Precipitation is reduced in the Indian monsoon region after the NH volcanic eruptions, but increases after the SH eruptions. The response patterns after the EQ eruptions are similar to those after the SH eruptions, which is likely related to the similar dynamical responses to EQ and SH volcanic eruptions.

Stratospheric volcanic aerosols directly influence the radiative budget of the climate system. This causes dynamical responses due to changes in the available energy and thermal gradients between the northern and the southern hemisphere as
365 well as between land and ocean, associated with interhemispherically asymmetric distributions of volcanic aerosols as well as the different heat capacity of land and ocean.

In India, the NH volcanic eruption leads to warming and drying. This is caused by the decreased water vapor advection and cloud cover over India, which is related to reduced monsoon circulation and convection after the volcanic eruption. The ITCZ moves southward due to the altered interhemispheric thermal contrast. The South Asian summer monsoon weakens
370 due to the decreased land-ocean thermal contrast. After the SH and EQ volcanic eruptions, cooling and wetting result from the increase of water vapor and clouds over India, connected to strengthened horizontal and vertical motions. This is associated with the northward movement of ITCZ and the strengthened South Asian summer monsoon after the volcanic eruptions.

Compared to previous studies, which are usually based on different classifications of historical volcanic eruptions with
375 different numbers of volcanic events, aerosol magnitudes, eruption latitudes and eruption seasons, this study largely reduces these uncertainties by using an idealized model setup. In this setup, the eruption latitude is the only difference between the three eruption cases, and the eruption season is the only difference between the winter and the summer eruptions. In this way,



and by performing 10-member ensemble simulations, we provide robust model results on the sensitivity of the climate effect of volcanic eruptions to eruption latitude and eruption season. These results may also support the interpretation of climate impacts of different choices of stratospheric aerosol injection in geoengineering approaches (Simpson et al., 2019).
380 In contrast to the previous assumption that tropical eruptions have stronger climate effects than extratropical eruptions, this study suggests the opposite. Further research needs to be performed to understand whether tropical or extratropical eruptions have stronger climate impact using different models and experimental setups. The simulations in this study avoid the concurrent effect of volcanic eruptions and ENSO through controlling the initial state of ENSO. Results do not show any
385 significant impact of the eruptions on ENSO variations. The impact of volcanic eruptions on ENSO and their concurrent or different impacts on climate are still unsolved questions, which deserve to be further studied.

Code availability. Post-processing and visualization of data was performed with CDO and batch scripts. The scripts are available on request from the corresponding author.

390 **Data availability.** All data required to reproduce our key results is published by Zhuo et al. (2021) at World Data Center for Climate (WDCC) at Deutsche Klimarechenzentrum (DKRZ).

Author Contributions. Z.Z. designed the study, ran the MPI-ESM experiments, analyzed the results and wrote the manuscript. I.K. and U.C. supervised and provided support for designing the study. I.K. provided support for running the experiments. I.K. and S.P. provided support for the analysis. All authors contributed to revising the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

395 **Acknowledgements.** This work is supported by China Scholarship Council (CSC). We thank Matthew Toohey for his help on the EVA module. The authors acknowledge the Deutsche Klimarechenzentrum (DKRZ, <https://www.dkrz.de/>) for the computational resources and computing facilities.

References

- 400 Adams, J. B., Mann, M. E., and Ammann, C. M.: Proxy evidence for an El Niño-like response to volcanic forcing, *Nature*, 426(6964), 274-278, doi:10.1038/nature02101, 2003.
- Ammann, C. M., and Naveau, P.: Statistical analysis of tropical explosive volcanism occurrences over the last 6 centuries, *Geophys. Res. Lett.*, 30, 1210, doi:10.1029/2002gl016388, 2003.
- Colose, C. M., LeGrande, A. N., and Vuille, M.: Hemispherically asymmetric volcanic forcing of tropical hydroclimate during the last millennium, *Earth Syst. Dynam.*, 7(3), 681-696, doi:10.5194/esd-7-681-2016, 2016.
- 405 Dogar, M. M., and Sato, T.: Regional climate response of middle eastern, African, and South Asian monsoon regions to explosive volcanism and ENSO forcing, *J. Geophys. Res.-Atmos.*, 124, 7580–7598, doi:10.1029/2019JD030358, 2019.
- Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: An improved ice core-based index for climate models, *J. Geophys. Res.*, 113, D23111, doi:10.1029/2008jd010239, 2008.




- 410 Gillett, N. P., Weaver, A. J., Zwiers, F. W., and Wehner, M. F.: Detection of volcanic influence on global precipitation, *Geophys. Res. Lett.*, 31, L12217, doi:10.1029/2004gl020044, 2004.
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., . . . Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *J. Adv. Model. Earth Syst.*, 5, 572-597, doi:10.1002/jame.20038, 2013.
- 415 Guo, Y., Cao, J., Li, H., Wang, J., and Ding, Y.: Simulation of the interface between the Indian summer monsoon and the East Asian summer monsoon: Intercomparison between MPI-ESM and ECHAM5/MPI-OM, *Adv. Atmos. Sci.*, 33(3), 294-308, doi:10.1007/s00376-015-5073-z, 2016.
- Haywood, J. M., Jones, A., Bellouin, N., and Stephenson, D.: Asymmetric forcing from stratospheric aerosols impacts Sahelian rainfall, *Nat. Clim. Change*, 3(7), 660-665, doi:10.1038/nclimate1857, 2013.
- 420 Iles, C. E., and Hegerl, G. C.: The global precipitation response to volcanic eruptions in the CMIP5 models, *Environ. Res. Lett.*, 9(10), 104012, doi:10.1088/1748-9326/9/10/104012, 2014.
- Iles, C. E., Hegerl, G. C., Schurer, A. P., and Zhang, X.: The effect of volcanic eruptions on global precipitation, *J. Geophys. Res.-Atmos.*, 118(16), 8770-8786, doi:10.1002/jgrd.50678, 2013.
- Ilyina, T., Six, K. D., Segsneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *J. Adv. Model Earth Sy.*, 5(2), 287-315, doi:10.1029/2012ms000178, 2013.
- Jacobson, T. W. P., Yang, W., Vecchi, G. A., and Horowitz, L. W.: Impact of volcanic aerosol hemispheric symmetry on Sahel rainfall, *Clim. Dynam.*, 55(7-8), 1733-1758, doi:10.1007/s00382-020-05347-7, 2020.
- 430 Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., . . . von Storch, J. S.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *J. Adv. Model Earth Sy.*, 5(2), 422-446, doi:10.1002/jame.20023, 2013
- Khodri, M., Izumo, T., Vialard, J., Janicot, S., Cassou, C., Lengaigne, M., . . . McPhaden, M. J.: Tropical explosive volcanic eruptions can trigger El Nino by cooling tropical Africa, *Nat. Commun.*, 8(1), 778, doi:10.1038/s41467-017-00755-6, 2017.
- 435 Liu, F., Chai, J., Wang, B., Liu, J., Zhang, X., and Wang, Z.: Global monsoon precipitation responses to large volcanic eruptions, *Sci Rep.*, 6, 24331, doi:10.1038/srep24331, 2016.
- Man, W., and Zhou, T.: Response of the East Asian summer monsoon to large volcanic eruptions during the last millennium, *Chin. Sci. Bull.*, 59, 4123-4129, doi:10.1007/s11434-014-0404-5, 2014.
- Man, W., Zhou, T., and Jungclaus, J. H.: Simulation of the East Asian summer monsoon during the last millennium with the MPI Earth System Model, *J. Climate*, 25(22), 7852-7866, doi:10.1175/jcli-d-11-00462.1, 2012.
- 440 Man, W., Zhou, T., and Jungclaus, J. H.: Effects of large volcanic eruptions on global summer climate and East Asian monsoon changes during the last millennium: Analysis of MPI-ESM simulations, *J. Climate*, 27(19), 7394-7409, doi:10.1175/jcli-d-13-00739.1, 2014.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and natural radiative forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 659-740, doi:10.1017/CBO9781107415324.018, 2013.
- 450 Oman, L., Robock, A., and Stenchikov, G. L.: Climatic response to high-latitude volcanic eruptions, *J. Geophys. Res.*, 110, D13103, doi:10.1029/2004jd005487, 2005.
- Oman, L., Robock, A., Stenchikov, G. L., and Thordarson, T.: High-latitude eruptions cast shadow over the African monsoon and the flow of the Nile, *Geophys. Res. Lett.*, 33(L18711), doi:10.1029/2006gl027665, 2006.
- 455 Paik, S., and Min, S.-K.: Climate responses to volcanic eruptions assessed from observations and CMIP5 multi-models, *Clim. Dynam.*, 48, 1017-1030, doi:10.1007/s00382-016-3125-4, 2016.
- Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *J. Adv. Model Earth Sy.*, 5(3), 459-482, doi:10.1002/jame.20022, 2013
- Robock, A.: Volcanic eruptions and climate, *Rev. Geophys.*, 38(2), 191-219, doi:10.1029/1998RG000054, 2000.



- 460 Robock, A., and Liu, Y.: The volcanic signal in Goddard Institute for Space Studies three-dimensional model simulations, *J. Climate*, 7, 44-55, doi:10.1175/1520-0442(1994)007<0044:TVSIGI>2.0.CO;2, 1994.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L., . . . Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Climate*, 19, 3771-3791, doi:10.1175/JCLI3824.1, 2006.
- 465 Schneck, R., Reick, C. H., and Raddatz, T.: Land contribution to natural CO₂ variability on time scales of centuries, *J. Adv. Model Earth Sy.*, 5(2), 354-365, doi:10.1002/jame.20029, 2013.
- Schneider, D. P., Ammann, C. M., Otto-Bliesner, B. L., and Kaufman, D. S.: Climate response to large, high-latitude and low-latitude volcanic eruptions in the Community Climate System Model, *J. Geophys. Res.*, 114, D15101 doi:10.1029/2008jd011222, 2009.
- 470 Simpson, I. R., Tilmes, S., Richter, J. H., Kravitz, B., MacMartin, D. G., Mills, M. J., Fasullo, J. T., Pendergrass, A. G.: The regional hydroclimate response to stratospheric sulfate geoengineering and the role of stratospheric heating, *J. Geophys. Res.-Atmos.*, 124, 12587-12616, doi:10.1029/2019JD031093, 2019.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., . . . Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, *J. Adv. Model Earth Sy.*, 5(2), 146-172, doi:10.1002/jame.20015, 2013.
- 475 Stevenson, S., Fasullo, J. T., Otto-Bliesner, B. L., Tomas, R. A., and Gao, C.: Role of eruption season in reconciling model and proxy responses to tropical volcanism, *P. Natl. Acad. Sci. USA*, 114(8), 1822-1826, doi:10.1073/pnas.1612505114, 2017.
- Stevenson, S., Otto-Bliesner, B., Fasullo, J., and Brady, E.: "El Niño like" hydroclimate responses to last millennium volcanic eruptions, *J. Climate*, 29(8), 2907-2921, doi:10.1175/jcli-d-15-0239.1, 2016.
- 480 Timmreck, C.: Modeling the climatic effects of large explosive volcanic eruptions. *WIREs Clim. Change*, 3(6), 545-564. doi:10.1002/wcc.192, 2012.
- Timmreck, C., Pohlmann, H., Illing, S., and Kadow, C.: The impact of stratospheric volcanic aerosol on decadal-scale climate predictions, *Geophys. Res. Lett.*, 43(2), 834-842, doi:10.1002/2015gl067431, 2016.
- Toohey, M., Krüger, K., Schmidt, H., Timmreck, C., Sigl, M., Stoffel, M., and Wilson, R.: Disproportionately strong climate forcing from extratropical explosive volcanic eruptions, *Nat. Geosci.*, 12(2), 100-107, doi:10.1038/s41561-018-0286-2, 2019.
- Toohey, M., Krüger, K., Sigl, M., Stordal, F., and Svensen, H.: Climatic and societal impacts of a volcanic double event at the dawn of the Middle Ages, *Climatic Change*, 136(3-4), 401-412, doi:10.1007/s10584-016-1648-7, 2016.
- Toohey, M., Stevens, B., Schmidt, H., and Timmreck, C.: Easy Volcanic Aerosol (EVA v1.0): an idealized forcing generator for climate simulations, *Geosci. Model Dev.*, 9(11), 4049-4070, doi:10.5194/gmd-9-4049-2016, 2016.
- 490 Trenberth, K. E., and Dai, A.: Effects of Mount Pinatubo volcanic eruption on the hydrological cycle as an analog of geoengineering, *Geophys. Res. Lett.*, 34(15), L15702, doi:10.1029/2007gl030524, 2007.
- Valcke, S.: The OASIS3 coupler: a European climate modelling community software, *Geosci. Model Dev.*, 6(2), 373-388, doi:10.5194/gmd-6-373-2013, 2013.
- 495 Webster, P. J., and Yang, S.: Monsoon and ENSO: Selectively interactive systems, *Q. J. Roy. Meteor. Soc.*, 118, 877-926, doi:10.1002/qj.49711850705, 1992.
- Yang, W., Vecchi, G., Fueglistaler, S., Horowitz, L. W., Luet, D. J., Muñoz, Á. G., Paynter, D., Underwood, S.: Climate impacts from large volcanic eruptions in a high-resolution climate model: The importance of forcing structure, *Geophys. Res. Lett.*, 46, 7690-7699, doi:10.1029/2019GL082367, 2019.
- 500 Zambri, B., and Robock, A.: Winter warming and summer monsoon reduction after volcanic eruptions in Coupled Model Intercomparison Project 5 (CMIP5) simulation, *Geophys. Res. Lett.*, 43, 10920-10928, doi:10.1002/2016GL070460, 2016.
- Zhang, D., Blender, R., and Fraedrich, K.: Volcanoes and ENSO in millennium simulations: global impacts and regional reconstructions in East Asia, *Theor. Appl. Climatol.*, 111(3-4), 437-454, doi:10.1007/s00704-012-0670-6, 2012.
- 505 Zhuo, Z., Gao, C., Kirchner, I., and Cubasch, U.: Impact of volcanic aerosols on the hydrology of the Asian monsoon and westerlies-dominated subregions: Comparison of proxy and multimodel ensemble means, *J. Geophys. Res.-Atmos.*, 125, e2020JD032831, doi:10.1029/2020jd032831, 2020.
- Zhuo, Z., Gao, C., and Pan, Y.: Proxy evidence for China's monsoon precipitation response to volcanic aerosols over the past seven centuries, *J. Geophys. Res.-Atmos.*, 119(11), 6638-6652, doi:10.1002/2013JD021061, 2014.




- 510 Zhuo, Z., Kirchner, I., Cubasch, U.: Different latitudinal volcanic eruption simulations in different seasons met.fu-berlin MPI-ESM-LR echam monthly. World Data Center for Climate (WDCC) at Deutsche Klimarechenzentrum (DKRZ). https://doi.org/10.26050/WDCC/LAT-SEASON-VOLCANIC_MPI-ESM-LR, 2021.
- Zuo, M., Man, W. M., Zhou, T. J., and Guo, Z.: Different impacts of northern, tropical, and southern volcanic eruptions on the tropical pacific SST in the last millennium, *J. Climate*, 31(17), 6729-6744, doi:10.1175/Jcli-D-17-0571.1, 2018.
- 515 Zuo, M., Zhou, T., and Man, W.: Hydroclimate responses over global monsoon regions following volcanic eruptions at different latitudes, *J. Climate*, 32(14), 4367-4385, doi:10.1175/jcli-d-18-0707.1, 2019.




520 Figure 1: Schematic view of the experimental design. Simulations are performed using MPI-ESM (Giorgetta et al., 2013) with volcanic forcing produced from the EVA module (Toohey et al., 2016).

525



530 Figure 2: Global mean aerosol optical depth anomaly at 550 nm for the summer (a) and winter (b) volcanic eruption cases and their zonal mean distributions (c and d).



535

Figure 3: Global mean net outgoing solar radiation (W/m^2) anomaly at top of the atmosphere in clear-sky conditions for the summer (a) and winter (b) volcanic eruption cases and their zonal mean distributions (c and d). The grey shading in (a) and (b) indicates two standard deviation of the control runs without any volcanic eruption. Dark grey and light grey stippling in (c) and (d) indicate the grid points with significant differences based on the two-tailed student's t-test at the 95% and 99% confidence level, respectively.




Figure 4: Global mean surface temperature (°C) anomaly for the summer (a) and winter (b) volcanic eruption cases and their zonal mean distributions (c and d). The grey shading in (a) and (b) indicates two standard deviations around the mean of the control runs without any volcanic eruption. Dark grey and light grey stippling in (c) and (d) indicate the grid points with significant differences based on the two-tailed student's t-test at the 95% and 99% confidence level, respectively.





Figure 5: Spatial distribution of JJA mean temperature ($^{\circ}\text{C}$) anomaly in 1992 for the summer NH (a), EQ (c), SH (e) eruption cases and in 1991 for the winter NH (b), EQ (d) and SH (f) eruption cases. The cross signs and slashes indicate the grid points with significant differences based on the two-tailed student's t-test at the 95% and 99% confidence level, respectively.



550 **Figure 6:** Spatial distribution of JJA mean precipitation (mm/day) anomaly in 1992 for the summer NH (a), EQ (c), SH (e) eruption cases and in 1991 for the winter NH (b), EQ (d) and SH (f) eruption cases. The cross signs and slashes indicate the grid points with significant differences based on the two-tailed student's t-tests at the 95% and 99% confidence level, respectively.

555




Figure 7: Difference of JJA mean temperature (°C) anomaly between the northern and the southern hemisphere after the summer eruptions in 1992 (a) and the winter eruptions in 1991 (b), and the position of the ITCZ (c and d) indicated by the JJA mean zonal mean precipitation (mm/day) anomaly between 20°N and 20°S. The grey bar in (a) and (b) indicates the two-standard deviation of the control runs.

560

565

570



575




Figure 8: South Asian summer monsoon index (m/s) anomaly for the summer eruption cases (a) and the winter eruption cases (b). The grey bar indicates one standard deviation of the control runs.

580

585

590

595

600

605

610





Figure 9: JJA mean vertically integrated moisture transport (IVT, vector, kg/m²·s) and its divergence (IVTD, shaded, kg/m²·s) for the control runs without any volcanic eruption (a), and their anomalies after the NH (b), EQ (c) and SH (d) volcanic eruptions in 1992.



620 **Figure 10: Spatial distribution of JJA mean net surface solar radiation (W/m²) anomaly in clear-sky (a) and all-sky conditions (b) after the NH (top), EQ (middle) and SH (bottom) volcanic eruptions in 1992. The vertical bars and plus signs indicate the grid points with significant differences based on the two-tailed student's t-test at the 95% and 99% confidence level, respectively.**

625



630





Figure 11: Spatial distribution of JJA mean total cloud cover (%) anomaly after the NH (a), EQ (b) and SH (c) volcanic eruptions in 1992. The vertical bars and plus signs indicate the grid points with significant differences based on the two-tailed student's t-tests at the 95% and 99% confidence level, respectively.



635

Figure 12: Spatial distribution of JJA mean vertical integrated water vapor (kg/m²) anomaly after the NH (a), EQ (b) and SH (c) summer volcanic eruptions in 1992. The vertical bars and plus signs indicate the grid points with significant differences based on two-tailed student's t-tests at the 95% and 99% confidence level, respectively.