



Measurement report: The chemical composition and temporal variability of aerosol particles at Tuktoyaktuk, Canada during the Year of Polar Prediction Special Observing Period

John MacInnis¹, Jai Prakash Chaubey¹, Crystal Weagle^{1,2}, David Atkinson³, Rachel Ying-Wen Chang¹

- 5 ¹Department of Physics and Atmospheric Science, Dalhousie University, Halifax, B3H 4R2, Canada
²Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, St. Louis, 63130-4899, United States of America
³Department of Geography, University of Victoria, Victoria, V8P 5C2, Canada

Correspondence to: Rachel Ying-Wen Chang (rachel.chang@dal.ca)

10 **Abstract.** The chemical composition, sources, and concentrations of aerosol particles vary on a seasonal basis in the Arctic. While existing research has focused on understanding the occurrence of aerosol particles during the Arctic winter and spring, less is known of their occurrence during the Arctic summer. In this study, atmospheric aerosol particle chemical composition and concentration were determined during July-September 2018 at Tuktoyaktuk, NT, Canada (69.4° N, 133.0° W) to coincide with the Year of Polar Prediction's 2nd Special Observing Period in the Arctic. The chemical composition of fine (PM_{2.5}) and
15 coarse (PM_{10-2.5}) aerosol filter samples suggests the ocean, mineral/road dust, and combustion were sources of the sampled aerosol particles. Mass concentrations of PM₂ and PM₁₀, estimated from optical particle counter measurements, remained within a similar range during the study. However, elevated mass concentrations coincided with a festival in the community of Tuktoyaktuk, suggesting local human activity was an important source of aerosol particles. Mass concentrations of PM₂, which promote negative health effects in humans, were significantly lower at Tuktoyaktuk than the national air quality standard
20 recommended by the Government of Canada. These measurements provide an important baseline to compare with future measurements associated with the assessment of aerosol chemistry and air quality in the Arctic.

1 Introduction

Aerosols are suspensions of liquid and solid particles in the atmosphere resulting from direct emissions from natural and anthropogenic sources and physical transformations, such as condensation and nucleation (Finlayson-Pitts and Pitts, 2000;
25 Seinfeld and Pandis, 2016). The concentration, size distribution, and chemical composition of aerosol particles vary significantly in the atmosphere because their sources are diverse and ephemeral in nature (Finlayson-Pitts and Pitts, 2000; Seinfeld and Pandis, 2016). The role of aerosol particles in the atmosphere is complex: they serve as short-lived climate forcers (Willis et al., 2018), provide reactive surfaces for heterogeneous chemistry (Newberg et al., 2005), and are vectors for the atmospheric transport (Wong et al., 2018) and post-depositional fate (MacInnis et al., 2019) of anthropogenic contaminants to
30 and within remote Arctic environments, respectively.



In recent years, efforts have focused on understanding the source and chemical composition of aerosol particles in the Arctic, particularly within the context of climate change (Willis et al., 2018). Aerosol particle profiles vary on a seasonal basis near the surface in the Arctic. For example, the winter-spring period is characterized by high mass concentrations of accumulation mode aerosol particles primarily from anthropogenic sources (i.e., Arctic haze), with a chemical composition dominated by sulphate (SO_4^{2-}) and black carbon (Sharma et al., 2017, 2019). In contrast, the summer is a period characterized by low mass concentrations of aerosol particles primarily associated with the Aitken mode and natural sources (Chang et al., 2011; Willis et al., 2018). Indeed, aerosol particles emitted from natural sources (e.g., marine and mineral dust aerosols) are important during the Arctic summer, however, it is anticipated that emissions from anthropogenic sources will increase as the Arctic responds to climate warming (Willis et al., 2018). An important consequence of climate warming is decreasing sea ice coverage, which promotes local warming through positive feedback albedo interactions, cloud and fog formation, the ocean-atmosphere exchange of aerosol particles and nucleation gases, and anthropogenic activities related to shipping and the exploration of minerals and fossil fuels (Willis et al., 2018; Abbatt et al., 2019). Enhancements in summertime pollution promoted by anthropogenic activity may have an effect on the role of aerosol particles as climate forcers in the Arctic. For example, while it has been suggested that cooling is an ultimate effect of aerosol particles in the Arctic atmosphere, increasing emissions of black carbon from anthropogenic activities may contribute to Arctic warming through aerosol-radiation interactions (Willis et al., 2018). Changes in the chemical composition of aerosol particles may also affect their hygroscopicity and atmospheric fate as cloud condensation nuclei, contributing to heating and cooling effects through aerosol-cloud interactions (Willis et al., 2018). The overall net impact of aerosol particles as short-lived climate forcers in the Arctic is unknown, however, increasing aerosol particle emissions are expected to have an impact on local air quality in the Arctic.

The size and chemical composition of aerosol particles has important implications for human health. In particular, aerosol particles with diameters less than or equal to $2.5 \mu\text{m}$ (referred to as $\text{PM}_{2.5}$) can accumulate in human lungs. These particles have been associated with a number of diseases in humans, such as bronchitis, asthma, and cardiovascular disease (Dominici et al., 2006; Wu et al., 2018). In response to these concerns, the World Health Organization established a guideline for $\text{PM}_{2.5}$ of $25 \mu\text{g m}^{-3}$ (24-hour average), which is suggested to be a level that minimizes risks to human health (World Health Organization, 2018). A similar guideline was established for $\text{PM}_{2.5}$ by the federal government of Canada (Canadian Ambient Air Quality Standard of $27 \mu\text{g m}^{-3}$) (Environment and Climate Change Canada, 2018), although current research explores the effects of aerosol particle exposure at lower concentrations (Brauer et al., 2019; Christidis et al., 2019; Pappin et al., 2019). For instance, several studies have noted that lower $\text{PM}_{2.5}$ mass concentrations ($5 \mu\text{g m}^{-3}$) have been associated with non-accidental death (Brauer et al., 2019; Pappin et al., 2019). Furthermore, the chemical components associated with $\text{PM}_{2.5}$ may be deleterious to human health, as it has been suggested that long-term exposure to various ions and metals in $\text{PM}_{2.5}$, including SO_4^{2-} , nitrate (NO_3^-), potassium (K), iron (Fe), zinc (Zn), and silicon (Si) were associated with annual ischemic heart disease mortality in humans (Ostro et al., 2010).

Currently, the Arctic atmosphere is characterized by low concentrations of aerosol particles in the summer (Willis et al., 2018), however, it has been predicted that the chemical composition and magnitude of aerosol particle concentrations will



65 change in the future as a result of climate change (Browse et al., 2013; Gilgen et al., 2018; Willis et al., 2018; Abbatt et al.,
2019), which may have important implications on aerosol-radiation and aerosol-cloud interactions in the Arctic (Croft et al.,
2019). In this study, the chemical composition and concentration of aerosol particles were determined in the western Canadian
Arctic during the Year of Polar Prediction's (YOPP) 2nd Special Observing Period in 2018. The YOPP was an international
collaboration led by the World Meteorological Organization designed to improve weather and sea ice forecasting and
70 environmental prediction through the implementation of intensive and modelling campaigns in polar regions
(<https://www.polarprediction.net/>). This work is a contribution to the YOPP, as the measurement of aerosol properties in the
Canadian Arctic provides a baseline for future predictions of aerosol particle concentration and composition in this region.

2 Methods and data analysis

2.1 Study area and sample collection

75 Tuktoyaktuk, NT, Canada (69.45° N, 133.04° W), located at 5 m above sea level, is a hamlet bordering the Amundsen Gulf
region of the eastern Beaufort Sea, and the latter is generally ice-covered during October-June (Herenz et al., 2018). It has a
population of 982 (2018) and is accessible primarily via Inuvik-Tuktoyaktuk Highway (an unpaved, gravel road) or the local
airport (Fig. 1). Tuktoyaktuk experiences a subarctic climate (Köppen climate classification) characterized by long, cold
winters and brief, mild summers. The average annual temperature and total annual precipitation are less than 0 °C and 300
80 mm, respectively (Herenz et al., 2018).

Fine (PM_{2.5}) and coarse (PM_{10-2.5}) aerosol filter samples were collected from Tuktoyaktuk using an AirPhoton SS4i air
sampler (Air Photon, Baltimore, USA, www.airphoton.com), as described previously (Snider et al., 2015), from 18 July to 12
September 2018. Briefly, air is drawn into the inlet of this assembly system by a vacuum pump and large particles (i.e., greater
than 10 µm) are collected on a greased impaction plate. The sampled air then passes through a porous membrane (Nuclepore®,
85 Whatman, 8 µm pore size) for the collection of PM_{10-2.5}, followed by a polytetrafluoroethylene (2 µm pore size) filter for the
collection of PM_{2.5}. This assembly can accommodate eight filter units, which were gravimetrically weighed before and after
the sampling period using a Sartorius Ultramicro Balance in a cleanroom with controlled relative humidity (32 ± 12 %) and
temperature (20.9-21.0 °C) at Dalhousie University. Due to a file writing error during the study, the sample volume could not
be determined and only the absolute mass collected on the filters is provided. We acknowledge this is a limitation of our study;
90 however, the chemical composition of these samples remains of interest, as it is relevant to our understanding of the
atmospheric fate of aerosol particles, particularly during the Arctic summer in which existing measurements are scarce (Chang
et al., 2011; Sierau et al., 2014; Tremblay et al., 2019).

Aerosol particles were sampled in situ using a GT-526S Handheld Particle Counter (Met One Instruments, Inc., Oregon,
USA) from 24 July to 13 September 2018. The instrument was calibrated by the manufacturer before the study and sampled
95 ambient air through an inlet line that was less than 1 m in length. This unit contains six channels that simultaneously measures
aerosol particles binned by diameter with lower limits of 0.3, 0.5, 1, 2, 5, and 10 µm. Sample collection was performed every

ten minutes at a flow rate of 2.8 L min^{-1} during the study period. The instruments in this study were mounted approximately 3.5 m above ground level at the Aurora College Community Centre in Tuktoyaktuk.



100

Figure 1. Location of the sampling site at the Aurora College Community Centre and the local airport in Tuktoyaktuk. Inset map shows the location of the Tuktoyaktuk in Canada and the Arctic Ocean. Map source: Toporama (Natural Resources Canada).

2.2 Laboratory analysis

105 Aerosol filter samples were extracted and analysed for ions and metals (Snider et al., 2015) at Dalhousie University in 2019. Before extraction, a ceramic blade was used to cut filters in half. One half of the filter was used for the analysis of water-soluble ions, including fluoride (F^-), chloride (Cl^-), bromide (Br^-), nitrite (NO_2^-), NO_3^- , SO_4^{2-} , phosphate (PO_4^{3-}), lithium (Li^+), sodium (Na^+), K^+ , ammonium (NH_4^+), calcium (Ca^{2+}), and magnesium (Mg^{2+}), which were extracted using 3 mL of distilled water/isopropyl alcohol (4 %) and ultrasonication. Water-soluble ions were separated by a Dionex ICS-1000 ion
110 chromatograph and analysed by conductivity detection. The other half of the filter was subjected to acid digestion using 3 mL of 10 % nitric acid (HNO_3) to extract metals, including Li, Mg, aluminum (Al), Fe, uranium (U), silver (Ag), barium (Ba), arsenic (As), Si, vanadium (V), chromium (Cr), manganese (Mn), cadmium (Cd), nickel (Ni), copper (Cu), Zn, antimony (Sb), cerium (Ce), lead (Pb), cobalt (Co), titanium (Ti), phosphorous (P), and selenium (Se). Metals were analysed by inductively coupled plasma mass spectrometry (ICP-MS, Thermo Scientific X-Series 2). Further details related to the extraction and
115 analysis of filters can be found elsewhere (Snider et al., 2015).



2.3 QA/QC

Field blanks were used to investigate contamination introduced during the sampling and transport of filters in this study. Field blanks in this study were filters that were transported to the field site, but were not deployed, and returned to the laboratory for analysis concurrently with samples. Procedural blanks were used to investigate contamination introduced during extraction.

120 These blanks consisted of the same analytical reagents used to extract filter samples, but without the aerosol filter matrix. Due to the unanticipated, brief sampling period in this study (i.e., estimated at 3 hours for each filter), the chemical mass of ions and metals in aerosol particle filter sample extracts were low and often estimated through extrapolation. We fully acknowledge that this practice introduces uncertainty into the quantitative assessment of ions in metals in aerosol particle filter samples and related analyses (Sect. 3.2). However, all reported masses are above the instrument detection limit (IDL). In addition, the

125 chemical masses of ions and metals reported in this study were subjected to screening to account for their presence in procedural and field blanks (US Environmental Protection Agency, 2016), pursuant to protocols implemented by the US EPA: (1) if the analyte is detected in all field and procedural blanks (i.e., at masses equal to or greater than the IDL), then the detection limit is defined as the mean mass of analyte in blanks plus three-times the standard deviation (SD) of masses in the

130 blanks; or (2) if some, but not all, blanks contain analyte, then the detection limit is assigned to the highest mass observed in the blanks; or (3) if the analyte is not detected in the blanks, then the detection limit is assigned to the IDL. The reported mass of ions and metals in filter samples are not blank subtracted (Supplement, Table S1).

2.4 Data analysis

The absolute mass of ions and metals on filters was calculated by multiplying extract concentrations ($\mu\text{g mL}^{-1}$ or ng mL^{-1}) by the extraction volume (3 mL) and a factor of 2 to determine the mass of ions and metals on the entire filter. A mass

135 reconstruction analysis was performed to estimate contributions by ions, metals, and particle-bound water to the total gravimetric mass, as described in the Supplement (Table S2). It was assumed that NO_3^- and SO_4^{2-} were neutralized by NH_4^+ ; however, we recognize that it is possible NO_3^- and SO_4^{2-} were associated with other species, depending on their source and atmospheric fate (e.g., NaNO_3 produced from the acid displacement reaction with NaCl and HNO_3). Similarly, there is uncertainty related to the hypothesis that the source of Cl^- in aerosol filter samples is limited to the ocean (i.e., NaCl

140 component in Table S2), particularly due to the analytical challenges associated with the quantification of Na^+ in our samples. Nevertheless, we assumed that all Cl^- in aerosol filter samples originated from marine aerosol particles for the purpose of this mass reconstruction estimate. Data from 27 August were not included in the calculation determining the average mass reconstruction profile for $\text{PM}_{2.5}$ because the chemical mass sampled was larger than the gravimetric mass. Discrepancies between the total gravimetric mass and measured chemical mass of aerosol filter samples could be attributed

145 to untargeted species (e.g., organics, other ions and metals, etc.), analytical uncertainties and/or losses of volatile species from filters during transport and laboratory analysis.



Air mass back trajectories were calculated using the Hybrid Single Particle Lagrangian Integrated Trajectory Model (National Oceanic and Atmospheric Administration, https://www.ready.noaa.gov/HYSPLIT_traj.php) to understand the source of ions and metals in aerosol filter samples. Air mass back trajectories were calculated over 48 hours using North American Regional Reanalysis (NARR) meteorology, setting the heights at the same location (end of parcel trajectory) to be 50, 200, and 400 m above ground level (Figs. S2 and S3). Available instrument log files suggest that sampling occurred once every eight days from 6:00 to 9:00 AM local time. Therefore, 9:00 AM was used as an end point for air mass back trajectory analysis.

Aerosol particle mass concentrations were estimated from measured aerosol particle number concentrations according to Eq. (1):

$$C_{\text{Mass}} = C_{\text{Num}} \cdot d_p^3 \cdot \rho_p \cdot \pi/6, \quad (1)$$

where C_{Mass} is aerosol particle mass concentration ($\mu\text{g m}^{-3}$), C_{Num} is the aerosol particle number concentration measured by the particle counter (cm^{-3}), d_p is the geometric mean diameter of aerosol particles in a given size bin (cm), and ρ_p is aerosol particle density, which was assumed to be 1.8 g cm^{-3} (Sharma et al., 2017). PM_2 and PM_{10} mass concentrations were calculated as the sum of mass concentrations in 0.3-2 μm and 0.3-10 μm size bins, respectively.

3 Results and discussion

3.1 Aerosol filter masses

The fine ($\text{PM}_{2.5}$, mean \pm SD, $15 \pm 9 \mu\text{g}$) and coarse ($\text{PM}_{10-2.5}$, $14 \pm 4 \mu\text{g}$) aerosol filter masses were similar during the study period, albeit with notable variability (Fig. 2). For instance, the masses range from 2.6-31 μg and 7.3-22 μg in $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$, respectively. Snider et al. (2016) also reported that masses of $\text{PM}_{2.5}$ (median 72 μg , lower-upper quintiles 42-131 μg) and $\text{PM}_{10-2.5}$ (median 90 μg , lower-upper quintiles 44-154 μg) were comparable in filter samples collected across a global network of sites (i.e., Surface PARTICulate mAtter Network, SPARTAN) using an AirPhoton sampler. The variability of aerosol filter mass was likely a reflection of dynamic aerosol particle sources of local and distant origin and meteorological events (e.g., changes in precipitation, wind speed and direction, Fig. S1) during the study period.

3.2 Chemical composition of aerosol filters

The detection frequencies for metals in aerosol filter samples were often low and/or variable, such as those observed for Ag, Ba, Cu, Sb, Ti, and Zn (Table S1).




Figure 2. A summary of the gravimetric mass (left panels) and chemical mass of ions (middle panels) and metals (right panels) in fine ($PM_{2.5}$) and coarse ($PM_{10-2.5}$) aerosol filter samples (top and bottom row, respectively) from Tuktoyaktuk during July-September 2018.

In contrast, Al was detected in 100 % and 75 % of $PM_{2.5}$ and $PM_{10-2.5}$ samples, respectively. Interestingly, some metals were only detected in $PM_{2.5}$, while others were only detected in $PM_{10-2.5}$. For example, Ti was only detected in $PM_{2.5}$, whereas Cu, Ag, and Ba were only detected in $PM_{10-2.5}$, suggesting the latter may have been from local sources given their shorter atmospheric lifetime due to gravitational settling. Al and Fe were the metals measured in the highest quantities in aerosol filter samples (Fig. 2).

The detection frequencies for water-soluble ions in aerosol filter samples were generally low and variable (Fig. 2). SO_4^{2-} and Cl^- were found in most $PM_{2.5}$ samples, with detection frequencies of 87 % and 62 %, respectively, whereas NO_3^- was only detected in 25 % of $PM_{2.5}$ samples. Similar to metals, several water-soluble ions were only detected in select aerosol filter samples. For example, F^- , NO_2^- , and Br^- were only detected in $PM_{10-2.5}$, suggesting a local source, whereas Cl^- , SO_4^{2-} , NH_4^+ and Ca^{2+} were only detected in $PM_{2.5}$. The highest quantities of water-soluble ions in aerosol filter samples were observed for F^- , Ca^{2+} , PO_4^{3-} , and Cl^- (Fig. 2).



In general, the chemical composition of metals in aerosol filter samples at Tuktoyaktuk was similar to other Arctic regions (Fig. 3), in that Al and Fe dominated the metal profiles, which have been linked to mineral dust emissions (Liberda et al., 2015; Ferrero et al., 2019). In contrast, contributions from Ba and Sb to the total quantity of metals at Tuktoyaktuk were greater than other Arctic regions, whereas Ti contributions from other Arctic regions were greater than those at Tuktoyaktuk (Landsberger et al., 1990; Kadko et al., 2016; Conca et al., 2019). Ion composition profiles for aerosol filter samples at Tuktoyaktuk were notably different than other Arctic regions (Leaitch et al., 2018; Ferrero et al., 2019), with the exception of the SO_4^{2-} in $\text{PM}_{2.5}$ at Tuktoyaktuk, whose composition profile was comparable to those found in aerosol particles from the Arctic Ocean and Ny-Ålesund (Fig. 3) (Ferrero et al., 2019). The ion composition profiles were dominated by Na^+ and Cl^- at Ny-Ålesund and the Arctic Ocean, and by SO_4^{2-} at Alert (Leaitch et al., 2018; Ferrero et al., 2019). Na^+ and Cl^- have been linked to marine aerosol particle emissions, while SO_4^{2-} has been linked to anthropogenic combustion and natural sources (Leaitch et al., 2018; Ferrero et al., 2019).

High detection frequencies of Al in $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$ filter samples suggest that mineral dust was a consistent source of aerosol particles at Tuktoyaktuk. It is possible that local mineral/road dust emissions were sources of aerosol particles in this region, although the mass of Al was comparable in $\text{PM}_{2.5}$ and $\text{PM}_{10-2.5}$ filter samples, which may not be an outcome expected from local mineral dust emissions if the size distribution of precursor soils is coarse (i.e., higher mass expected in $\text{PM}_{10-2.5}$). Furthermore, it has been suggested that giant ($d_p > 40 \mu\text{m}$) and saltation mode ($3 \leq d_p \leq 6 \mu\text{m}$) mineral dust aerosol particles are primarily produced from soils (Saltzman, 2013). It is noteworthy that other metals characteristic of mineral dust, such as Fe and Ti, were detected less frequently in filter samples (Table S1), suggesting they had different sources. Interestingly, the highest mass of Al, Ti, Fe, and Zn in $\text{PM}_{2.5}$ filter samples was observed on 4 September, and air masses during this period originated from north westerly directions, over the ocean. Similar air mass trajectories originating from west and north westerly directions were observed on 18, 26 July and 11-27 August (Figs. S2 and S3), and filter samples during these periods contained Ba, Ag, and Sb. Taken together, it is possible that emissions from natural or anthropogenic dust (e.g., road dust containing tire wear and mineral/soil particles) (Snider et al., 2016), and mining activities in western Canada (Government of Canada, 2018) and Alaska (Alaska Miners Association, 2020) were sources of Al, Fe, Ti, Zn, Ba, Ag, and Sb in filter samples.

It is interesting that Cl^- was only detected in $\text{PM}_{2.5}$ while Br^- was only detected in $\text{PM}_{10-2.5}$ filter samples since Cl^- and Br^- have been measured in seawater from the Canadian Arctic Archipelago (Xu et al., 2016). If the presence of Br^- in $\text{PM}_{10-2.5}$ filter samples was attributed to marine aerosol particles, then Cl^- should have also been present based on the molar composition of Cl^- and Br^- in seawater from the Canadian Arctic Archipelago (Xu et al., 2016). On the other hand, it is important to consider that the absence of Br^- in $\text{PM}_{2.5}$ filter samples could be related to analytical challenges because the mass of Br^- expected in $\text{PM}_{2.5}$ filter samples, based on the molar composition of Cl^- and Br^- in seawater from the Canadian Arctic Archipelago (Xu et al., 2016), is below the detection limit in this study.

225




Figure 3. Composition profiles of ions (top panel) and metals (bottom panel) in aerosol filter samples at Tuktoyaktuk and other Arctic regions. Average composition profiles are shown for aerosol filter samples at Tuktoyaktuk. Several elements in Landsberger et al. (1990)³ (Ca, Cl, Na), Conca et al. (2019)⁴ (Mg, K, Na, Ca), and Kadko et al. (2016)⁵ (Na, Mg) were detected in aerosol particles, however, they are not included in this figure to enable comparisons with trace metal composition profiles. Data shown for Leitch et al. (2018)⁷, Kadko et al. (2016)⁵, Ferrero et al. (2019)⁶, Landsberger et al. (1990)³, and Conca et al. (2019)⁴ correspond to sampling periods during 8 August 2014, 14 August 2011, 2011-2012, 7-18 March 1985, and 2010-2013, respectively.

It is unlikely that the absence of Cl^- in $\text{PM}_{10-2.5}$ filter samples can be explained by acid displacement reactions, considering it has been suggested that Cl^- depletion decreases with increasing aerosol particle size (Yao et al., 2003). Thus, these observations suggest that Cl^- and Br^- were from different sources. In addition to marine sources, Cl^- and Br^- can originate from biomass burning. For example, hydrochloric acid (HCl), chlorine (Cl_2), hypochlorous acid (HOCl), bromine (Br_2), and hypobromous acid (HOBr) have all been identified as products of biomass burning (Keene et al., 2006), which could have been the source of either Cl^- measured in the fine mode or Br^- measured in the coarse mode at Tuktoyaktuk.

It is challenging to identify sources of ions and metals in aerosol filter samples, especially when they are emitted from various sources in the environment (Jayarathne et al., 2014; Leitch et al., 2018; Willis et al., 2018). For instance, it is likely that ions and metals originated from continental and marine sources in this study, pursuant to air mass back trajectory analysis



during the study period (Figs. S2 and S3). One strategy that can be used to better constrain continental and marine sources is to compare known ratios of ions and metals in soils and seawater to those found in aerosol filter samples. As an example, the molar ratios of Fe/Al and Cl⁻/SO₄²⁻ in PM_{2.5} from Tuktoyaktuk are compared to those found in Icelandic soils (Đorđević et al., 2019) and seawater from the Canadian Arctic Archipelago (Xu et al., 2016), respectively. The molar ratio of Fe/Al in Icelandic soils is 0.9, and this ratio is within the range of those observed in aerosol filter samples from Tuktoyaktuk (0.4-1.4). This result further supports the hypothesis that Fe and Al could originate from mineral dust. In contrast, the molar ratio of Cl⁻/SO₄²⁻ in Arctic seawater is 27, which is higher than those in PM_{2.5} from Tuktoyaktuk (0.7-6.8). Lower molar ratios of Cl⁻/SO₄²⁻ in aerosol filter samples could be attributed to non-oceanic sources of SO₄²⁻ (i.e., natural and anthropogenic combustion sources). For example, SO₄²⁻ in PM_{2.5} at Tuktoyaktuk may have originated from natural sources, such as the biogenic emission and subsequent oxidation of dimethyl sulphide from the ocean (Bates et al., 1987) and/or emission and subsequent oxidation of sulfur dioxide (SO₂) from lignite in the Smoking Hills, NT (Radke and Hobbs, 1989). An additional source of SO₄²⁻ in PM_{2.5} at Tuktoyaktuk may include anthropogenic emissions from the combustion of fossil fuels (e.g., vehicles, aircraft, boats, etc.) (Leaitch et al., 2018; Willis et al., 2018). Other ions characteristic of combustion were also identified in aerosol filter samples from Tuktoyaktuk, such as NO₃⁻ and NH₄⁺, possibly from the emission and oxidation of nitrogen oxides (NO_x) and emissions of ammonia during fossil fuel combustion from local and long-range sources. However, ammonia emissions in the Arctic have also been associated with natural sources, such as soil (Wentworth et al., 2016) and guano (Croft et al., 2016; Wentworth et al., 2016), which could account for NH₄⁺ in aerosol filter samples at Tuktoyaktuk. It is important to consider that lower molar ratios of Cl⁻/SO₄²⁻ in PM_{2.5} filter samples from Tuktoyaktuk could also be attributed to Cl⁻ depletion from marine aerosol via acid displacement (Newberg et al., 2005). For example, it is possible that HNO₃ and sulfuric acid (H₂SO₄) displaced Cl⁻ in PM_{2.5} from Tuktoyaktuk (Laskin et al., 2012), particularly since NO₃⁻ and SO₄²⁻ were detected in aerosol filter samples that were devoid of Cl⁻ (i.e., 26 July-11 August). Although the organic composition of aerosol filter samples was not characterized in this work, it is important to consider that organic acids (Laskin et al., 2012) may have also contributed to Cl⁻ depletion in aerosol filter samples.

A mass reconstruction analysis was conducted to estimate contributions from ions and metals to the total gravimetric mass (Bari and Kindzierski, 2017). The average mass reconstruction profile for PM_{2.5} indicates that contributions from marine and mineral dust sources collectively accounted for 43 % of the total known chemical mass (i.e., NaCl and mineral dust components), while contributions from (NH₄)₂SO₄ and NH₄NO₃ collectively accounted for 31 % of the known chemical mass (Fig. 4). Lower contributions were observed for metals and particle-bound water, collectively accounting for 6 % of the total known mass in PM_{2.5} filter samples. In contrast, the average mass reconstruction profile for PM_{10-2.5} was dominated by NH₄NO₃ and unclassified inorganic ions (i.e., combined masses of F⁻, Br⁻, PO₄³⁻, NO₂⁻, Table S2), which collectively accounted for 93 % of the total chemical mass. This analysis suggests that marine, mineral dust, and combustion sources were important in this region; however, the source of PO₄³⁻ and F⁻, which were major components of the unclassified inorganic fraction of aerosol particle filter samples, could not be identified.




Figure 4. Average mass reconstruction profiles of fine and coarse aerosol filter samples at Tuktoyaktuk during July-September 280 2018. Several ions measured in this study (F^- , Br^- , PO_4^{3-} , NO_2^-) are collectively included here as unclassified inorganic components due to uncertainties associated with their sources in the environment.

The sources of these ions are interesting because PO_4^{3-} and F^- have not been previously identified in aerosol particles from the Arctic (Fig. 3). We acknowledge that it is possible that the detection of F^- could be related to laboratory contamination, 285 however, the reported masses in this study are equal to or greater than the detection limit (i.e., mean + 3SD of blank masses). It is challenging to assign sources to PO_4^{3-} and F^- because they are released from multiple sources in the environment, such as emissions from volcanos, mineral dust, combustion, and marine aerosol particles (Jayarathne et al., 2014; Vet et al., 2014). It is also challenging to assign sources to F^- because one of its gas phase precursors (i.e., hydrofluoric acid, HF), emitted during 290 smelting activity, coal combustion, and volcanic activity (Jayarathne et al., 2014) may partition to aerosol particles in the Arctic atmosphere. While there is uncertainty associated with the source of PO_4^{3-} and F^- in aerosol filter samples, the back trajectories suggest that air masses containing PO_4^{3-} travelled primarily over the ocean, whereas air masses containing F^- travelled over the ocean and land (Figs. S2 and S3). These observations suggest that marine, mineral, and combustion aerosol particles are possible sources of PO_4^{3-} and F^- at Tuktoyaktuk.

In addition to uncertainties associated with the unclassified inorganic fraction, a substantial fraction of mass on aerosol 295 filter samples could not be identified. For instance, the average chemical mass only accounted for 10 % and 12 % of the total gravimetric mass in $PM_{2.5}$ and $PM_{10-2.5}$ filter samples, respectively. Discrepancies between the total gravimetric mass and chemical mass of aerosol filter samples could be attributed to analytical uncertainties, the loss or gain of volatile species from filters after sampling (Saltzman, 2013), and/or contributions from untargeted chemical components. For example, it is possible that other inorganic ions and metals, organic material, and black carbon were components of aerosol particles at Tuktoyaktuk, 300 pursuant to the chemical composition of aerosol particles in other Arctic regions (Kadko et al., 2016; Leaitch et al., 2018;



Conca et al., 2019; Ferrero et al., 2019; Sharma et al., 2019). Despite these uncertainties, our results indicate that there are substantial, uncharacterized chemical components that could not be identified in aerosol filter samples at Tuktoyaktuk.

The summertime composition profiles at Tuktoyaktuk may provide insights into the future chemical composition of aerosol particles in the Arctic. Our analysis indicates that mineral dust, marine, and combustion sources are important during the Arctic summer, particularly in the absence of snow cover and sea ice coverage. It is expected that emissions of these aerosol particles will increase as the Arctic responds to climate warming in the future. For example, it is expected that climate warming will enhance the melting of snowpacks and sea ice within the Arctic, promoting exposed landscapes and oceans, and ship traffic (Willis et al., 2018). Our analysis also indicates that there are significant, unknown components of aerosol filter samples during the summer of 2018 at Tuktoyaktuk, which may influence the chemical properties of aerosol particles and their role in the Arctic troposphere (e.g., cloud condensation nuclei, radiative properties) (Martin et al., 2011; Herenz et al., 2018; Willis et al., 2018; Abbatt et al., 2019). For example, the unidentified components (e.g., organic components) are likely less hygroscopic than the soluble inorganics identified and would reduce the cloud condensation nuclei activity of the aerosol particles. This highlights the importance of improving understanding of the chemical composition of aerosol particles in the Arctic. It is important to emphasize that these results only provide a snapshot of the aerosol particles at Tuktoyaktuk and their representativeness is unknown.

3.3 Size distribution, temporal variability, and health implications of aerosol particles

The average number size distributions were similar throughout the study, with particle number concentrations highest in the 0.3-0.5 μm bin (Fig. S4), consistent with an earlier study in the region (Herenz et al., 2018). The mass size distributions also remained similar throughout the study, with mass concentrations dominated by the 2-5 μm aerosol particles (Fig. S4).

Total aerosol particle number concentrations (N_d) in this study were low (1-hour average, mean \pm SD, $6 \pm 8 \text{ cm}^{-3}$, median 3 cm^{-3}). These relatively low concentrations are consistent with other observations during the Arctic summer (Willis et al., 2018) and may have been attributed to enhanced aerosol particle and nucleation gas scavenging by precipitation (Croft et al., 2016). Aerosol particle mass concentrations for $\text{PM}_{2.5}$ and PM_{10} were similar in regions of the Northwest Territories, Yukon Territory, and Nunavut (Canadian Council of Ministers of the Environment, 2019) but were higher than those observed for PM_2 and PM_{10} at Tuktoyaktuk during July-September 2018 (Table 1). Although some measurements at Tuktoyaktuk are not directly comparable to those of other regions in Canada (i.e., PM_2 vs. $\text{PM}_{2.5}$), it is possible that the discrepancy in the magnitude of aerosol particle mass concentrations could be attributed to differences in regional climate (i.e., precipitation) and aerosol particle sources. For instance, it is likely that Tuktoyaktuk is influenced by marine air masses to a greater extent than other sites listed in Table 1, which are generally located inland. Differences in regional populations/human activities could further promote differences in aerosol particle concentrations (e.g., enhanced vehicle traffic). It is interesting to note in Table 1 that relatively high aerosol particle mass concentrations were observed at Norman Wells despite its low population, however, this may be attributed to major crude oil and natural gas production in that region (Canada Energy Sector, 2020).



Table 1. An overview of mass concentrations for PM₂, PM_{2.5}, and PM₁₀ at Tuktoyaktuk and other regions in Canada (Canadian Council of Ministers of the Environment, 2019) from 26 July to 13 September 2018. Population data for 2018 was retrieved from the Northwest Territories Bureau of Statistics (NWT Bureau of Statistics, 2020), the Government of Nunavut (Government of Nunavut, 2019), and the Yukon Bureau of Statistics (Yukon Bureau of Statistics, 2019).

| Location | Tuktoyaktuk, NT 69° N | | Inuvik, NT 68° N | | Norman Wells, NT 65° N | | Iqaluit, NU 64° N | | Yellowknife, NT 62° N | | Whitehorse, YT 61° N |
|----------------------|-----------------------------|------------------|------------------------|------------------|------------------------------|------------------|-------------------------|-------------------|-----------------------------|-------------------|----------------------------|
| | Population | 982 | | 3,536 | | 818 | | 38,139 | | 20,607 | |
| $\mu\text{g m}^{-3}$ | PM ₂ | PM ₁₀ | PM _{2.5} | PM ₁₀ | PM _{2.5} | PM ₁₀ | PM _{2.5} | PM _{2.5} | PM ₁₀ | PM _{2.5} | PM _{2.5} |
| Mean | 0.9 | 2.3 | 2.6 | 13 | 3.7 | 11 | 4.3 | 2.6 | 16 | 2.4 | 2.4 |
| Median | 0.6 | 1.3 | 2.0 | 6 | 3.0 | 7 | 4.0 | 0.0 | 13 | 2.0 | 2.0 |
| Max | 6.9 | 48 | 19 | 133 | 22 | 122 | 15 | 58 | 97 | 48 | 48 |

The discrepancy in the magnitude of aerosol particle mass concentrations may also be related to differences in sampling techniques. For example, one method used by the National Air Pollutant Surveillance program to determine mass concentrations is by filtration and beta attenuation (Canadian Council of Ministers of the Environment, 2019) whereas the method used here relies on aerosol particle number concentrations and estimations of aerosol particle density (Eq. 1). Despite these discrepancies, these data indicate that aerosol particle concentrations were lower during the summer of 2018 at Tuktoyaktuk than other regions in Canada.

A summary of the continuous aerosol particle mass concentrations during July-September 2018, calculated from the particle counter, is presented in Figs. 5 and 6. In general, the magnitude of mass concentrations remained within a similar range during July-September 2018, although there were notable increases in mass concentrations during 28-30 July, 3-6 August, and 22-24 August (Fig. 5). For example, the average PM₁₀ concentration was 2.3 $\mu\text{g m}^{-3}$ over the study period, whereas higher PM₁₀ concentrations occurred on 28 July (35 $\mu\text{g m}^{-3}$), 4 August (48 $\mu\text{g m}^{-3}$), and 24 August (40 $\mu\text{g m}^{-3}$). Although we are unable to confirm the source of aerosol particle emissions during these periods, it is conceivable that elevated mass concentrations could be attributed to local aerosol particle emissions from human activities at Tuktoyaktuk. For instance, elevated PM₁₀ concentrations on 28 July may have been related to human activities that proceeded during the Annual Land of the Pingos Musical Festival (e.g., barbecuing, increased vehicle traffic), which occurred during 27-29 July at Tuktoyaktuk. Similar human activities could also provide an explanation for elevated mass concentrations on 3-6 August, considering this period coincided with an extended civic holiday weekend.





Figure 5. Mass concentration profiles for $\text{PM}_{2.5}$ and PM_{10} at Tuktoyaktuk during July-September 2018. The data are presented as 1-hour averages, and each monthly period is demarcated by colour.



360

Figure 6. Diurnal variability of average aerosol particle mass concentrations (coloured bars, $\mu\text{g m}^{-3}$), wind speed (solid blue line, m s^{-1}), and temperature (dashed black line, $^{\circ}\text{C}$) at Tuktoyaktuk during 24 July to 13 September 2018. The average mass size distribution is based on 1-hour averaged mass concentration data. Hourly wind speed and temperature data from the local airport can be found in Historical Climate Archives from Environment and Climate Change Canada.



365

An investigation of the diurnal variability of aerosol particle mass concentrations suggests that maxima typically occurred in the afternoon, with the exception of 0.3-0.5 μm aerosol particles. For instance, maximum concentrations were observed at 16:00 for 0.5-1 μm aerosol particles, 17:00 for 1-2, 2-5, and 5-10 μm aerosol particles, and between 22:00 and 0:00 for 0.3-0.5 aerosol particles (Fig. 6). Elevated mass concentrations for 0.5-10 μm aerosol particles coincided with elevated
370 wind speeds and temperatures in the afternoon, suggesting daytime heating may be relevant to the occurrence of these aerosol particles (e.g., through the action of convection and the lofting mineral dust and marine aerosol particles into the atmosphere).

To assess the impact of fine aerosol particle emissions on local air quality, the mass concentration of PM_2 at Tuktoyaktuk was compared to the Canadian Ambient Air Quality Standard. The concentration of PM_2 was low at Tuktoyaktuk during July-September 2018, with 24-hour average concentrations ranging from 0.2-3 $\mu\text{g m}^{-3}$. These mass concentrations were
375 very low in Tuktoyaktuk compared to the national $\text{PM}_{2.5}$ standard (27 $\mu\text{g m}^{-3}$), suggesting PM_2 likely had minimal effects on the air quality of the community.

4 Conclusions

The chemical composition of aerosol filter samples and concentration of aerosol particles from Tuktoyaktuk were determined during July-September 2018. Although our analysis could not identify distinct sources, the results suggest that this moderately-
380 sized community in the Canadian north was influenced by a wide range of aerosol particle sources with complex processes. The observed aerosol particles were likely derived from natural sources like marine and mineral dust sources and anthropogenic sources like local road dust, distant mining activities, and the combustion of fossil fuels. Our analysis indicates that there were significant, unknown components identified in aerosol filter samples during the summer of 2018 at Tuktoyaktuk, which may influence the atmospheric fate of aerosol particles in the Arctic troposphere. While the mass concentrations of PM_2 were found
385 to be significantly lower at Tuktoyaktuk compared to the Canadian Ambient Air Quality Standard, it is likely that their concentrations will increase in the future due to climate change, which is expected to promote increases in ship and air traffic in the Arctic as well as the number of ice-free days and natural emissions from open waters. Although these measurements only represent a snapshot of the aerosol particles at Tuktoyaktuk, they can nevertheless provide insights into the chemistry and concentration of aerosol particle samples, which can be used in the future to assess aerosol particle chemistry and air quality
390 in the Canadian Arctic. Future work should focus on improving the characterization of chemical components associated with aerosol particles in order to fill current knowledge gaps related to their summertime sources and atmospheric fate in the Arctic.

Data availability. Quality-controlled aerosol filter chemistry and aerosol particle number concentrations are available through the Federated Research Data Repository (<https://doi.org/10.20383/101.0269> and <https://doi.org/10.20383/101.0278>).

395



Author contribution. RC and DA were responsible for funding acquisition and conceptualized the research. JC and DA were responsible for data acquisition. JC and CW contributed to data curation. The formal analysis and visualization of data was performed by JM. JM and RC wrote the manuscript, with editorial feedback from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

400 *Acknowledgements.* The authors thank George Hibbs for providing access to the sampling site, Nicole Chisholm for weighing filters, Brenna Walsh for helpful discussions, and Randall Martin and Jong Sung Kim for providing resources during the deployment. This work was funded by the Marine Environmental Observation, Prediction, and Response Network (MEOPAR), Polar Knowledge Canada [NST-1718-0001] and the Canada Research Chair program [CRC-2013-00056]. This is a contribution to the Year of Polar Prediction (YOPP), a flagship activity of the Polar Prediction Project (PPP), initiated by the
405 World Weather Research Programme (WWRP) of the World Meteorological Organisation (WMO). We acknowledge the WMO WWRP for its role in coordinating this international research activity.

References

Abbatt, J., Leaitch, W. R., Aliabadi, A., Bertram, A., Blanchet, J.-P., Boivin-Rioux, A., Bozem, H., Burkart, J., Chang, R., Charette, J., Chaubey, J., Christensen, R., Cirisan, A., Collins, D., Croft, B., Dionne, J., Evans, G., Fletcher, C., Galí, M.,
410 Ghahremaninezhad, R., Girard, E., Gong, W., Gosselin, M., Gourdal, M., Hanna, S., Hayashida, H., Herber, A., Hesarak, S., Hoor, P., Huang, L., Hussherr, R., Irish, V., Keita, S., Kodros, J., Köllner, F., Kolonjari, F., Kunkel, D., Ladino, L., Law, K., Lévassieur, M., Libois, Q., Liggio, J., Lizotte, M., Macdonald, K., Mahmood, R., Martin, R., Mason, R., Miller, L., Moravek, A., Mortenson, E., Mungall, E., Murphy, J., Namazi, M., Norman, A.-L., O'Neill, N., Pierce, J., Russell, L., Schneider, J., Schulz, H., Sharma, S., Si, M., Staebler, R., Steiner, N., Thomas, J., von Salzen, K., Wentzell, J., Willis, M., Wentworth, G.,
415 Xu, J.-W. and Yakobi-Hancock, J.: Overview paper: New insights into aerosol and climate in the Arctic, *Atmos. Chem. Phys.*, 19(4), 2527–2560, doi:10.5194/acp-19-2527-2019, 2019.

Alaska Miners Association. 2020. Major Mines in Alaska: www.alaskaminers.org/major-mines, last access: 16 March 2021.

Bari, M. A. and Kindzierski, W. B.: Ambient fine particulate matter (PM_{2.5}) in Canadian oil sands communities: Levels, sources and potential human health risk, *Sci. Total Environ.*, 595, 828–838, doi:10.1016/j.scitotenv.2017.04.023, 2017.

420 Bates, T. S., Charlson, R. J. and Gammon, R. H.: Evidence for the climatic role of marine biogenic sulphur, *Nature*, 329, 319–321, doi:10.1038/329319a0, 1987.

Brauer, M., Brook, J. R., Christidis, T., Chu, Y., Crouse, D. L., Erickson, A., Hystad, P., Li, C., Martin, R. V., Meng, J., Pappin, A. J., Pinault, L. L., Tjepkema, M., Van Donkelaar, A., Weichenthal, S. and Burnett, R. T.: Mortality-Air Pollution



- Associations in Low-Exposure Environments (MAPLE): Phase 1, Res. Reports Heal. Eff. Inst., 203, 2019.
- 425 Browse, J., Carslaw, K. S., Schmidt, A. and Corbett, J. J.: Impact of future Arctic shipping on high-latitude black carbon deposition, *Geophys. Res. Lett.*, 40(16), 4459–4463, doi:10.1002/grl.50876, 2013.
- Canada Energy Sector. 2020. Provincial and Territorial Energy Profiles - Northwest Territories: <https://www.cer-rec.gc.ca/en/data-analysis/energy-markets/provincial-territorial-energy-profiles/provincial-territorial-energy-profiles-northwest-territories.html>, last access: 16 March 2021.
- 430 Canadian Council of Ministers of the Environment. 2019. Ambient Air Monitoring and Quality Assurance/Quality Control Guidelines: National Air Pollution Surveillance Program, 2019.
- Chang, R. Y. W., Leck, C., Graus, M., Müller, M., Paatero, J., Burkhardt, J. F., Stohl, A., Orr, L. H., Hayden, K., Li, S. M., Hansel, A., Tjernström, M., Leaitch, W. R. and Abbatt, J. P. D.: Aerosol composition and sources in the central Arctic Ocean during ASCOS, *Atmos. Chem. Phys.*, 11, 10619–10636, doi:10.5194/acp-11-10619-2011, 2011.
- 435 Christidis, T., Erickson, A. C., Pappin, A. J., Crouse, D. L., Pinault, L. L., Weichenthal, S. A., Brook, J. R., Van Donkelaar, A., Hystad, P., Martin, R. V., Tjepkema, M., Burnett, R. T. and Brauer, M.: Low concentrations of fine particle air pollution and mortality in the Canadian Community Health Survey cohort, *Environ. Heal. A Glob. Access Sci. Source*, 18, doi:10.1186/s12940-019-0518-y, 2019.
- Conca, E., Abollino, O., Giacomino, A., Buoso, S., Traversi, R., Becagli, S., Grotti, M. and Malandrino, M.: Source identification and temporal evolution of trace elements in PM₁₀ collected near to Ny-Ålesund (Norwegian Arctic), *Atmos. Environ.*, 203, 153–165, doi:10.1016/j.atmosenv.2019.02.001, 2019.
- 440 Croft, B., Martin, R. V., Leaitch, W. R., Burkart, J., Chang, R.Y.-W., Collins, D. B., Hayes, P. L., Hodshire, A. L., Huang, L., Kodros, J. K., Moravek, A., Mungall, E. L., Murphy, J. G., Sharma, S., Tremblay, S., Wentworth, G. R., Willis, M. D., Abbatt, J. P. D. and Pierce, J. R.: Arctic marine secondary organic aerosol contributes significantly to summertime particle size distributions in the Canadian Arctic Archipelago, *Atmos. Chem. Phys.*, 19, 2787–2812, doi:10.5194/acp-19-2787-2019, 2019.
- Croft, B., Wentworth, G. R., Martin, R. V., Leaitch, W. R., Murphy, J. G., Murphy, B. N., Kodros, J. K., Abbatt, J. P. D. and Pierce, J. R.: Contribution of Arctic seabird-colony ammonia to atmospheric particles and cloud-albedo radiative effect, *Nat. Commun.*, 7, doi:10.1038/ncomms13444, 2016a.
- 450 Croft, B., Martin, R. V., Leaitch, W.R., Tunved, P., Breider, T. J., D’Andrea, S. D. and Pierce, J. R.: Processes controlling the annual cycle of Arctic aerosol number and size distributions, *Atmos. Chem. Phys.*, 16, 3665–3682, doi:10.5194/acp-16-3665-2016, 2016b.



- 455 Dominici, F., Peng, R. D., Bell, M. L., Pham, L., McDermott, A., Zeger, S. L. and Samet, J. M.: Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases, *J. Am. Med. Assoc.*, 295(10), 1127–1134, doi:10.1001/jama.295.10.1127, 2006.
- Đorđević, D., Tošić, I., Sakan, S., Petrović, S., Đuričić-Milanković, J., Finger, D. C. and Dagsson-Waldhauserová, P.: Can volcanic dust suspended from surface soil and deserts of Iceland be transferred to central Balkan similarly to African dust (Sahara)?, *Front. Earth Sci.*, 7, 142, doi:10.3389/feart.2019.00142, 2019.
- 460 Environment and Climate Change Canada. 2018. Canadian Environmental Sustainability Indicators Air Quality: www.canada.ca/en/environment-climate-change/services/environmental-indicators/air, last access: 16 March 2021.
- Ferrero, L., Sangiorgi, G., Perrone, M. G., Rizzi, C., Cataldi, M., Markuszewski, P., Pakszys, P., Makuch, P., Petelski, T., Becagli, S., Traversi, R., Bolzacchini, E. and Zielinski, T.: Chemical composition of aerosol over the Arctic ocean from summer Arctic expedition (AREX) 2011-2012 cruises: Ions, amines, elemental carbon, organic matter, polycyclic aromatic hydrocarbons, n-alkanes, metals, and rare earth elements, *Atmosphere (Basel)*, doi:10.3390/atmos10020054, 2019.
- 465 Finlayson-Pitts, B. J. and Pitts, J. N.: Particles in the Troposphere, in *Chemistry of the Upper and Lower Atmosphere*, pp. 349–420, Academic Press., 2000.
- Gilgen, A., Huang, T.K.W., Ickes, L., Neubauer, D. and Lohmann, U.: How important are future marine and shipping aerosol emissions in a warming Arctic summer and autumn?, *Atmos. Chem. Phys.*, 18, 10521–10555, doi:10.5194/acp-18-10521-2018, 2018.
- 470 Government of Canada. 2018. National Pollutant Release Inventory (NPRI) - Bulk Data: <https://open.canada.ca/data/en/dataset/40e01423-7728-429c-ac9d-2954385ccdfe>, last access: 16 March 2021.
- Government of Nunavut. 2019. Population Estimates: www.gov.nu.ca/sites/default/files/population_estimates_report_july_1_2019_0.pdf, last access: 16 March 2021.
- 475 Herenz, P., Wex, H., Henning, S., Bjerring Kristensen, T., Rubach, F., Roth, A., Borrmann, S., Bozem, H., Schulz, H. and Stratmann, F.: Measurements of aerosol and CCN properties in the Mackenzie River delta (Canadian Arctic) during spring-summer transition in May 2014, *Atmos. Chem. Phys.*, 18, 4477–4496, doi:10.5194/acp-18-4477-2018, 2018.
- Jacobi, H. W., Obleitner, F., Da Costa, S., Ginot, P., Eleftheriadis, K., Aas, W. and Zannata, M.: Deposition of ionic species and black carbon to the Arctic snowpack: Combining snow pit observations with modeling, *Atmos. Chem. Phys.*, 19, 10361–10377, doi:10.5194/acp-19-10361-2019, 2019.
- 480 Jayarathne, T., Stockwell, C. E., Yokelson, R. J., Nakao, S. and Stone, E. A.: Emissions of fine particle fluoride from biomass



- burning, *Environ. Sci. Technol.*, 48(21), 12636–12644, doi:10.1021/es502933j, 2014.
- Kadko, D., Galfond, B., Landing, W. M. and Shelley, R. U.: Determining the pathways, fate, and flux of atmospherically derived trace elements in the arctic ocean/ice system, *Mar. Chem.*, 182, 38–50, doi:10.1016/j.marchem.2016.04.006, 2016.
- Keene, W. C., Lobert, J. M., Crutzen, P. J., Maben, J. R., Scharffe, D. H., Landmann, T., Hély, C. and Brain, C.: Emissions of major gaseous and particulate species during experimental burns of southern African biomass, *J. Geophys. Res. Atmos.*, 111(D04301), 1–18, doi:10.1029/2005JD006319, 2006.
- Landsberger, S., Vermette, S. J. and Barrie, L. A.: Multielemental composition of the Arctic aerosol, *J. Geophys. Res.*, 95(D4), 3509–3515, doi:10.1029/JD095iD04p03509, 1990.
- Laskin, A., Moffet, R. C., Gilles, M. K., Fast, J. D., Zaveri, R. a., Wang, B., Nigge, P. and Shutthanandan, J.: Tropospheric chemistry of internally mixed sea salt and organic particles: Surprising reactivity of NaCl with weak organic acids, *J. Geophys. Res. Atmos.*, 117(15), 1–12, doi:10.1029/2012JD017743, 2012.
- Leaith, W. R., Russell, L. M., Liu, J., Kolonjari, F., Toom, D., Huang, L., Sharma, S., Chivulescu, A., Veber, D. and Zhang, W.: Organic functional groups in the submicron aerosol at 82.5° N, 62.5° W from 2012 to 2014, *Atmos. Chem. Phys.*, 18(5), 3269–3287, doi:10.5194/acp-18-3269-2018, 2018.
- Liberda, E. N., Tsuji, L. J. S. and Peltier, R. E.: Mining in subarctic Canada: Airborne PM_{2.5} metal concentrations in two remote First Nations communities, *Chemosphere*, 139, 452–460, doi:10.1016/j.chemosphere.2015.07.058, 2015.
- MacInnis, J. J., Lehnher, I., Muir, D. C. G., St. Pierre, K. A., St. Louis, V. L., Spencer, C. and De Silva, A. O.: Fate and transport of perfluoroalkyl substances from snowpacks into a lake in the high Arctic of Canada, *Environ. Sci. Technol.*, 53(18), 10753–10762, doi:10.1021/acs.est.9b03372, 2019.
- Martin, M., Chang, R. Y. W., Sierau, B., Sjogren, S., Swietlicki, E., Abbatt, J. P. D., Leck, C. and Lohmann, U.: Cloud condensation nuclei closure study on summer arctic aerosol, *Atmos. Chem. Phys.*, 11, 11335–11350, doi:10.5194/acp-11-11335-2011, 2011.
- Newberg, J. T., Matthew, B. M. and Anastasio, C.: Chloride and bromide depletions in sea-salt particles over the northeastern Pacific Ocean, *J. Geophys. Res. D Atmos.*, 110(D6), doi:10.1029/2004JD005446, 2005.
- NWT Bureau of Statistics. 2020. Community Data: www.statsnwt.ca/community-data/, last access: 16 March 2021.
- Ostro, B., Lipsett, M., Reynolds, P., Goldberg, D., Hertz, A., Garcia, C., Henderson, K. D. and Bernstein, L.: Long-term exposure to constituents of fine particulate air pollution and mortality: Results from the California Teachers Study, *Environ.*



- Health Perspect., 118, 363–369, doi:10.1289/ehp.0901181, 2010.
- Pappin, A. J., Christidis, T., Pinault, L. L., Crouse, D. L., Brook, J. R., Erickson, A., Hystad, P., Li, C., Martin, R. V., Meng,
510 J., Weichenthal, S., Donkelaar, A. van, Tjepkema, M., Brauer, M. and Burnett, R. T.: Examining the shape of the association
between low levels of fine particulate matter and mortality across three cycles of the canadian census health and environment
cohort, *Environ. Health Perspect.*, 127(10), doi:10.1289/EHP5204, 2019.
- Radke, L. F. and Hobbs, P. V.: Arctic hazes in summer over Greenland and the North American Arctic. III: A contribution
from the natural burning of carbonaceous materials and pyrites, *J. Atmos. Chem.*, 9, 161–167, doi:10.1007/BF00052830, 1989.
- 515 Saltzman, E. S.: Marine Aerosols, in *Surface Ocean-Lower Atmosphere Processes.*, 2013.
- Seinfeld, J. H. and Pandis, S. N.: Aerosols, in *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*,
pp. 325–359, John Wiley & Sons, Inc., 2016.
- Sharma, S., Leaitch, W.R., Huang, L., Veber, D., Kolonjari, F., Zhang, W., Hanna, S. J., Bertram, A. K. and Ogren, J. A.: An
evaluation of three methods for measuring black carbon in Alert, Canada, *Atmos. Chem. Phys.*, 17(24), 15225–15243,
520 doi:10.5194/acp-17-15225-2017, 2017.
- Sharma, S., Barrie, L. A., Magnusson, E., Brattström, G., Leaitch, W. R., Steffen, A. and Landsberger, S.: A factor and trends
analysis of multidecadal lower tropospheric observations of Arctic aerosol composition, black carbon, ozone, and mercury at
Alert, Canada, *J. Geophys. Res. Atmos.*, 124(24), 14133–14161, doi:10.1029/2019JD030844, 2019.
- Sierau, B., Chang, R. Y. W., Leck, C., Paatero, J. and Lohmann, U.: Single-particle characterization of the high-Arctic
525 summertime aerosol, *Atmos. Chem. Phys.*, 14, 7409–7430, doi:10.5194/acp-14-7409-2014, 2014.
- Snider, G., Weagle, C. L., Martin, R. V., Van Donkelaar, A., Conrad, K., Cunningham, D., Gordon, C., Zwicker, M., Akoshile,
C., Artaxo, P., Anh, N. X., Brook, J., Dong, J., Garland, R. M., Greenwald, R., Griffith, D., He, K., Holben, B. N., Kahn, R.,
Koren, I., Lagrosas, N., Lestari, P., Ma, Z., Vanderlei Martins, J., Quel, E. J., Rudich, Y., Salam, A., Tripathi, S. N., Yu, C.,
Zhang, Q., Zhang, Y., Brauer, M., Cohen, A., Gibson, M. D. and Liu, Y.: SPARTAN: A global network to evaluate and
530 enhance satellite-based estimates of ground-level particulate matter for global health applications, *Atmos. Meas. Tech.*, 8, 505–
521, doi:10.5194/amt-8-505-2015, 2015.
- Snider, G., Weagle, C. L., Murdymootoo, K. K., Ring, A., Ritchie, Y., Stone, E., Walsh, A., Akoshile, C., Anh, N. X.,
Balasubramanian, R., Brook, J., Qonitan, F. D., Dong, J., Griffith, D., He, K., Holben, B. N., Kahn, R., Lagrosas, N., Lestari,
P., Ma, Z., Misra, A., Norford, L. K., Quel, E. J., Salam, A., Schichtel, B., Segev, L., Tripathi, S., Wang, C., Yu, C., Zhang,
535 Q., Zhang, Y., Brauer, M., Cohen, A., Gibson, M. D., Liu, Y., Martins, J. V., Rudich, Y. and Martin, R. V.: Variation in global
chemical composition of PM_{2.5}: emerging results from SPARTAN, *Atmos. Chem. Phys.*, 16, 9629–9653, doi:10.5194/acp-16-



- 9629-2016, 2016.
- Thakur, R. C. and Thamban, M.: Influence of gaseous and particulate species on neutralization processes of polar aerosol and snow — A case study from Ny-Ålesund, *J. Environ. Sci. (China)*, 76, 12–25, doi:10.1016/j.jes.2018.03.002, 2019.
- 540 Tremblay, S., Picard, J. C., Bachelder, J. O., Lutsch, E., Strong, K., Fogal, P., Leaitch, W.R., Sharma, S., Kolonjari, F., Cox, C. J., Chang, R. Y. W. and Hayes, P. L.: Characterization of aerosol growth events over Ellesmere Island during the summers of 2015 and 2016, *Atmos. Chem. Phys.*, 19(8), 5589–5604, doi:10.5194/acp-19-5589-2019, 2019.
- US Environmental Protection Agency: Definition and procedure for the determination of the method detection limit—Revision 1.11., 2016.
- 545 Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C. U., Aas, W., Baker, A., Bowersox, V. C., Dentener, F., Galy-Lacaux, C., Hou, A., Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher, T., Mahowald, N. M., Nickovic, S., Rao, P. S. P. and Reid, N. W.: A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus, *Atmos. Environ.*, 93, 3–100, doi:10.1016/j.atmosenv.2013.10.060, 2014.
- Wentworth, G. R., Murphy, J. G., Croft, B., Martin, R. V., Pierce, J. R., Côté, J. S., Courchesne, I., Tremblay, J. É., Gagnon, J., Thomas, J. L., Sharma, S., Toom-Sauntry, D., Chivulescu, A., Levasseur, M. and Abbatt, J. P. D.: Ammonia in the summertime Arctic marine boundary layer: Sources, sinks, and implications, *Atmos. Chem. Phys.*, 16, 1937–1953, doi:10.5194/acp-16-1937-2016, 2016.
- 550 Willis, M. D., Leaitch, W. R. and Abbatt, J. P. D.: Processes controlling the composition and abundance of Arctic aerosol, *Rev. Geophys.*, 56(4), 621–671, doi:10.1029/2018RG000602, 2018.
- 555 Willis, M. D., Bozem, H., Kunkel, D., Lee, A. K. Y., Schulz, H., Burkart, J., Aliabadi, A. A., Herber, A. B., Leaitch, W.R. and Abbatt, J. P. D.: Aircraft-based measurements of High Arctic springtime aerosol show evidence for vertically varying sources, transport and composition, *Atmos. Chem. Phys.*, 19(1), 57–76, doi:10.5194/acp-19-57-2019, 2019.
- Wong, F., Shoeib, M., Katsoyiannis, A., Eckhardt, S., Stohl, A., Bohlin-Nizzetto, P., Li, H., Fellin, P., Su, Y. and Hung, H.: Assessing temporal trends and source regions of per- and polyfluoroalkyl substances (PFASs) in air under the Arctic Monitoring and Assessment Programme (AMAP), *Atmos. Environ.*, 172, 65–73, doi:10.1016/j.atmosenv.2017.10.028, 2018.
- 560 World Health Organization. 2018. Ambient (outdoor) air pollution: [https://www.who.int/news-room/factsheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/factsheets/detail/ambient-(outdoor)-air-quality-and-health), last access: 16 March 2021.
- Wu, J.-Z., Ge, D.-D., Zhou, L.-F., Hou, L.-Y., Zhou, Y. and Li, Q.-Y.: Effects of particulate matter on allergic respiratory diseases, *Chronic Dis. Transl. Med.*, 4(2), 95–102, doi:10.1016/j.cdtm.2018.04.001, 2018.



- 565 Xu, W., Tenuta, M. and Wang, F.: Bromide and chloride distribution across the snow-sea ice-ocean interface: A comparative study between an Arctic coastal marine site and an experimental sea ice mesocosm, *J. Geophys. Res. Ocean.*, 121(8), doi:10.1002/2015JC011409, 2016.
- Yao, X., Fang, M. and Chan, C. K.: The size dependence of chloride depletion in fine and coarse sea-salt particles, *Atmos. Environ.*, 37, 743–751, doi:10.1016/S1352-2310(02)00955-X, 2003.
- 570 Yukon Bureau of Statistics. 2019. Population Report: http://yukon.ca/sites/yukon.ca/files/ybs/populationq4_2019_0_0.pdf, last access: 16 March 2021.