

Temporary pause in the growth of atmospheric ethane and propane in 2015-2018

Hélène Angot^{1,2}, Connor Davel¹, Christine Wiedinmyer³, Gabrielle Pétron^{3,4}, Jashan Chopra¹, Jacques Hueber^{1,5}, Brendan Blanchard¹, Ilann Bourgeois^{3,6}, Isaac Vimont³, Stephen A. Montzka⁴, Ben R. Miller^{3,4}, James W. Elkins⁴, Detlev Helmig^{1,5}.

¹Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, CO, USA.

²School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne, Sion, Switzerland.

³Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA.

⁴NOAA, Global Monitoring Laboratory (GML), Earth System Research Laboratories, Boulder, CO, USA.

⁵Boulder A.I.R. LLC, Boulder, CO, USA.

⁶NOAA, Chemical Sciences Laboratory (CSL), Earth System Research Laboratories, Boulder, CO, USA.

Correspondence to: Hélène Angot (helene.angot@epfl.ch)

Supplementary Information

Table S1: 2008-2010 response factors (RF), along with the associated relative standard deviation (RSD), used to calculate mixing ratios at GEOSummit.

	2008 (Days 150-365)		2009 (Days1-149)		2009 (Days 150-201)		2009 (Days 202-251)		2009 (Days 252-302)		2009 (Days 303-365)	
	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)
ethane	7.6	5%	7.4	3%	7.7	3%	7.6	5%	7.6	5%	7.8	6%
propane	7.2	6%	6.8	2%	7.3	4%	7.1	6%	7.1	7%	7.3	7%
i-butane	6.2		5.5		6.2		6.2		6.1		6.2	
acetylene	6.1		4.5		5.8		6.0		6.6		6.4	
n-butane	6.5		6.0		6.6		6.5		6.4		6.5	
i-pentane	4.5	1%	4.2	5%	4.4	4%	4.8		4.4	1%	4.3	1%
n-pentane	4.8	1%	4.9	1%	4.6	3%	5.1	1%	4.6	4%	4.6	5%
hexane	1.3		2.8		2.1		2.5		2.0		1.9	
benzene	2.7	10%	4.5	13%	2.6	13%	3.0	9%	2.5	6%	2.4	5%
toluene	1.4	2%	2.1	16%	1.3		1.6	6%	1.4	4%	1.2	1%

	2010 (Days1-149)		2010 (Days1-49)		2010 (Days50-99)		2010 (Days100-149)		2010 (Days150-200)	
	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)
ethane	7.65	8%	7.89	8%	8.00	10%	8.48	15%	7.49	1%
propane	7.38	7%	7.44	10%	7.60	10%	7.80	8%	7.29	1%
i-butane	6.47	5%	6.25		6.56		6.67		6.79	
acetylene			6.08		6.47		6.92		6.43	
n-butane	6.82	5%	6.56		6.86		7.08		7.18	
i-pentane	5.33	15%	4.55	2%	4.91	5%	5.73	14%	6.21	5%
n-pentane	5.90	15%	4.98	8%	5.44	7%	7.19	4%	6.79	8%
hexane	4.69	50%	3.05	45%	3.16	38%	5.90	23%	6.44	31%
benzene	3.52	40%	2.65	6%	1.64	85%	4.62	40%	4.17	48%
toluene	1.71	77%	1.43	7%	0.64	141%	1.83	102%	1.87	119%

Table S2: 2012-2020 monthly response factors (RF), along with the associated relative standard deviation (RSD), calculated from the repeated analysis of two independently prepared and cross-referenced standards in use at any given time.

	January		February		March		April		May		June		July		August		September		October		November		December	
	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)	RF	RSD (%)
2012																								
ethane									7.1	0.7	7.2	0.9	7.1	1.2	7.4	2.3	7.4	2.4	7.4	0.5	7.4	0.9	7.5	0.8
propane									6.8	0.7	6.8	0.8	6.7	1.2	7.0	2.7	7.0	2.9	7.0	0.6	7.0	0.9	7.1	0.7
i-butane									6.3	0.8	6.4	1.4	6.3	1.3	6.5	2.0	6.6	2.2	6.6	0.7	6.4	1.4	6.5	1.7
acetylene									3.3	12.6	4.1	13.2	4.7	8.0	5.2	30.0	5.5	16.0	5.8	13.2	6.0	6.2	5.9	6.0
n-butane									6.8	1.4	6.8	0.6	6.7	1.2	7.2	2.5	7.1	3.8	7.1	0.7	7.2	1.5	7.0	0.6
i-pentane									6.5	1.4	6.6	1.0	6.5	1.7	6.7	2.5	6.6	2.8	6.7	0.5	6.6	1.2	6.7	0.9
n-pentane									6.9	0.8	7.1	0.7	6.9	1.3	7.0	2.2	6.9	2.7	7.0	0.5	6.9	0.9	6.9	1.1
hexane									6.3	1.2	6.3	1.9	6.3	1.5	6.6	1.2	6.3	2.3	6.3	1.5	6.3	2.5	6.3	2.1
benzene									6.9	19.1	6.7	12.5	6.5	12.8	6.7	20.6	6.6	15.6	6.7	9.5	6.5	22.1	6.6	5.4
toluene									6.7	21.1	6.8	19.4	6.6	97.5	6.1	23.2	6.2	62.8	6.4	19.6	6.1	71.4	6.1	14.4
2013																								
ethane	7.4	0.6	7.5	0.7	7.5	0.6	7.4	0.8	7.2	1.3	7.4	0.5	7.5	1.2	7.4	3.0	7.6	1.4	7.5	2.0	7.3	1.9	6.3	2.9
propane	7.0	1.8	7.1	1.5	7.1	0.8	7.1	0.7	7.0	1.3	7.1	0.7	7.1	0.8	7.1	2.8	7.2	1.7	7.2	1.9	6.8	2.3	5.8	3.1
i-butane	6.5	2.4	6.6	1.8	6.6	4.7	6.5	2.5	6.4	4.2	6.7	1.1	6.6	3.2	6.6	1.6	6.6	1.8	6.6	2.9	6.3	1.4	5.3	2.4
acetylene	6.0	1.9	6.4	3.8	6.3	3.7	6.0	5.4	5.9	1.9	6.7	0.4	5.7	5.9	6.1		6.1		6.1		6.1		5.2	
n-butane	6.9		6.9		7.3		7.1		6.8		7.0		6.9		7.0	2.0	7.1	1.7	7.1	2.3	6.8	1.7	5.8	2.6
i-pentane	6.7	2.3	6.7	0.8	6.7	2.0	6.6	1.4	6.6	0.6	6.7	0.6	6.6	1.9	6.6	2.1	6.7	1.2	6.6	1.7	6.3	2.5	5.3	2.8
n-pentane	6.8	2.1	6.9	2.6	6.9	3.6	6.9	1.9	6.7	2.0	6.8	0.1	6.9	2.5	6.9	3.3	6.9	1.2	6.8	2.9	6.6	2.3	5.5	2.9
hexane	6.3	39.4	6.3	0.0	6.3	5.3	6.3	5.2	6.6	3.4	6.6	11.1	6.7	3.8	6.6	1.5	6.7	1.4	6.5	3.4	6.2	1.3	5.2	3.3
benzene	6.6	4.3	6.7	1.4	6.6	1.6	6.6	1.7	6.6	1.3	7.0	1.2	6.8	1.7	6.7	1.1	6.8	4.5	6.6	2.6	6.4	10.2	4.9	11.1
toluene	6.0	3.9	6.4	1.3	6.2	2.4	6.1	2.8	6.1	3.1	6.2	2.6	6.2	3.3	6.2	3.3	6.2	1.2	6.0	2.9	6.5	9.6	4.7	11.4
2014																								
ethane	6.4	1.0	6.3	0.5							7.6	1.7	7.5	0.8	7.6	3.0	7.7	3.1	7.6	1.1	7.7	3.8	7.7	1.9
propane	5.9	1.5	5.9	0.6							7.3	0.6	7.1	0.9	7.2	2.7	7.5	2.4	7.4	1.2	7.3	3.9	7.4	3.6
i-butane	5.3	1.0	5.3	0.7							6.7	4.3	6.5	0.8	6.8	5.5	7.0	3.2	7.0	0.1	6.8	3.5	6.8	4.6
acetylene	5.2		5.2								7.2	2.6	8.0	7.6	7.5	1.9	7.9	7.4	7.5	5.0	7.3	4.0	7.1	4.0
n-butane	5.9	1.2	5.8	0.7							7.3	2.9	7.2	1.2	7.2	3.6	7.5	1.9	7.4	7.0	7.0	2.3	7.1	3.4
i-pentane	5.4	1.4	5.4	0.3							6.8	2.2	6.7	0.7	6.8	4.4	7.1	1.6	7.1	1.8	6.8	4.8	6.9	2.1
n-pentane	5.5	1.6	5.6	1.7							7.1	1.4	7.0	1.1	7.1	3.0	7.3	0.9	7.3	2.3	7.1	2.7	7.2	2.8
hexane	5.1	1.0	5.3	1.5							6.8	1.3	6.8	0.7	6.9	3.1	7.1	0.7	7.0	1.4	6.9	0.7	6.9	2.1
benzene	5.0	2.0	4.9	1.2							7.0	0.7	6.8	0.3	7.0	3.5	7.3	1.0	7.2	1.7	7.1	2.3	7.0	1.2
toluene	4.5	1.2	4.4	1.5							6.3	1.2	6.2	0.7	6.3	2.8					6.5	2.2	6.5	1.6
2015																								
ethane	7.7	2.6	7.7	1.8	7.7	1.0	7.8	1.1	7.7	0.8	7.6	1.0	7.5	0.5	7.7	2.0	7.7	1.1	7.6	1.0	7.8	0.6	7.7	0.8
propane	7.3	2.3	7.4	1.4	7.4	1.4	7.4	1.6	7.4	1.2	7.3	1.1	7.2	0.5	7.3	1.9	7.2	0.5	7.2	0.8	7.2	1.1	7.2	0.5
i-butane	6.7	3.4	6.8	2.4	6.8	2.1	6.8	1.7	6.9	1.7	6.9	3.1	6.6	3.2	6.9	4.7	6.3	3.4	6.6	1.0	6.5	2.2	6.6	1.6
acetylene	7.6	4.8	7.6	2.7	7.8	1.5	7.6	2.1	6.8	1.5	6.7	4.1	6.1	5.8	5.5	7.4	5.7	4.6	5.8	3.0	5.3	2.4	5.4	1.4
n-butane	7.1	3.4	7.2	3.0	7.2	3.9	7.3	0.9	7.5		7.2		7.0		7.2		6.9		7.1		7.0		7.0	
i-pentane	6.9	1.8	6.9	2.3	7.0	1.6	7.0	1.9	7.1	2.8	7.0	2.1	6.9	1.8	7.0	2.9	6.7	3.6	6.8	1.1	6.5	1.2	6.5	0.9
n-pentane	7.1	2.9	7.2	2.4	7.2	1.7	7.2	1.4	7.2	3.2	7.2	2.4	7.1	1.8	7.1	2.5	6.6	2.7	6.8	1.7	6.6	0.5	6.7	1.8
hexane	6.9	1.8	6.9	1.6	7.0	1.1	7.0	1.0	6.9	1.9	6.8	1.3	6.6	0.5	6.7	3.1	5.6	4.8	6.3	3.8	6.3	2.1	6.4	2.1
benzene	7.0	2.7	7.1	1.1	7.1	1.7	7.2	1.1	7.2	3.2	7.0	2.7	7.3	1.5	7.3	4.2	6.6	3.0	7.0	2.8	6.6	1.2	7.0	0.9
toluene	6.5	1.6	6.5	1.9	6.6	2.0	6.5	0.9	6.9	5.4	6.7	3.7	6.2	2.9	6.1	7.3	4.7	6.4	5.4	6.6	5.9	2.0	5.9	3.6
2016																								
ethane	7.7	0.8	7.7	0.5	7.6	0.9	7.6	0.9	7.8	0.7	7.4	0.7	7.4	1.2	7.3	1.3	7.2	0.6	7.2	0.5	7.2	0.9	7.2	0.9
propane	7.3	0.7	7.3	0.8	7.2	1.7	7.2	1.2	7.3	0.9	7.3	0.7	7.3	0.8	7.3	1.9	7.3	0.9	7.4	0.6	7.3	0.8	7.4	0.8
i-butane	6.5	2.0	6.7	0.8	6.6	1.4	6.6	2.0	6.6	1.2	6.4	3.3	6.5	2.0	6.6	3.2	6.7	0.8	6.7	1.0	6.6	2.4	6.7	2.4
acetylene	6.2	1.7	6.3	3.5	6.0	2.6	6.1	2.7	5.7	2.1	5.9	22.3	5.9	2.8	5.6	7.5	5.9	4.5	6.1	2.6	6.4	1.5	6.4	2.4
n-butane	7.1		7.1		6.8		7.1		7.0		6.9		7.0		7.0		7.2		7.3		7.2		7.0	
i-pentane	6.6	2.0	6.7	1.4	6.6	1.1	6.7	3.1	6.2	3.7	5.9	9.7	6.1	4.3	6.4	3.0	6.8	1.8	6.9	1.3	6.7	1.0	6.8	0.7
n-pentane	6.8	2.2	7.0	1.7	6.6	1.4	6.8	2.8	6.2	5.3	5.7	15.6	6.1	6.5	6.4	3.7	7.0	1.6	7.0	1.2	7.0	1.7	6.8	1.2
hexane	6.5	2.2	6.6	1.1	6.7	2.5	6.4	1.7	5.1	8.0	4.6	57.7	5.6	8.0	6.0	13.4	6.7	1.0	6.8	0.8	6.7	0.7	6.9	1.0
benzene	7.2	1.7	7.1	0.4	7.0	1.6	7.0	0.9	5.5	9.3	5.2	9.8	6.3	10.7	7.3	11.8	7.5	2.2	7.4	0.6	7.3	0.8	7.3	0.8
toluene	6.8	7.3	6.5	2.7	5.9	3.4	5.8	3.1	4.5	12.9	5.2	38.4	4.6	2.2	5.5	5.5	6.6	3.1	6.3	2.3	6.1	2.7	6.0	2.6




Figure S1: Mean seasonal amplitude, calculated as the relative difference between the winter maximum and the summer minimum, as a function of the lifetime of individual non-methane hydrocarbons against oxidation by the OH radical. The red line gives the fitted linear regression line with the 95 % confidence interval (grey shaded region).




Figure S2: a) Ethane, and b) propane time-series at GEOSummit from July 2008 to March 2020. Individual data points are shown in light grey, smoothed fit in dark grey, and trends in salmon. Trends inferred from discrete flask sampling are shown in turquoise. The slopes given by the Theil-Sen test for trend analysis are shown in black.




Figure S3: Scatter plot of daily averaged in-situ GC-FID measurements (x-axis) and discrete samples (y-axis) for **a)** ethane and **b)** propane. The red line gives the fitted linear regression. The slope ($\pm 95\%$ confidence interval) and correlation coefficient are given at the top. Note that these are not the same samples being compared: discrete samples are 5 minute-snapshots while in-situ GC-FID measurements are daily averages of 6-12 20 minute-samples.




Figure S4: PSCF results for ethane and propane residuals greater than the 90th percentile for April 2012-June 2019.




Figure S5: Scatter plot of ethane, propane, and benzene vs. carbon monoxide (CO) mixing ratios in the July and August 2019 biomass burning plumes observed at GEOSummit. The red line gives the fitted linear regression line with the 95 % confidence interval (grey shaded region). The slope, given at the top, gives the emission ratio (amount of compound emitted divided by that of a reference compound (CO here)).




Figure S6: Scatter plot of ethane (a, b) and propane (c, d) vs. tetrachloroethylene and hydrogen cyanide mixing ratios in the remote free troposphere during the global-scale aircraft mission ATom. Tetrachloroethylene and hydrogen cyanide are used as tracers of anthropogenic and biomass burning emissions, respectively. The red line gives the fitted linear regression line with the 95 % confidence interval (grey shaded region). The correlation coefficient is given at the top.