



Identifying the spatiotemporal variations of ozone formation regimes across China from 2005 to 2019 based on polynomial simulation and causality analysis

Ruiyuan Li¹, Miaoqing Xu¹, Manchun Li², Ziyue Chen¹, Bingbo Gao³, Na Zhao^{4,5}, Qi Yao¹

5 ¹State Key Laboratory of Remote Sensing Science, College of Global Change and Earth System Sciences, Beijing Normal University, Beijing, 100875, China

²School of Geography and Ocean Science, Nanjing University, Nanjing, 210023, China.

³College of Land Science and Technology, China Agriculture University, Beijing, 100083, China.

10 ⁴State Key Laboratory of Resources and Environment Information System, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, 100049, China

⁵University of Chinese Academy of Sciences, Beijing, 100080, China

Correspondence to: Ziyue Chen (zychen@bnu.edu.cn)

Abstract. Ozone formation regimes are closely related to the ratio of VOCs to NO_x. Different ranges of HCHO/NO₂ indicate three formation regimes, including VOCs-limited, transitional and NO_x-limited regimes. Due to the unstable interactions
15 between a diversity of precursors, the range of transitional regime, which plays a key role in identifying ozone formation regimes, remains unclear. To overcome the uncertainties from single models and the lack of reference data, we employed two models, polynomial simulation and Convergent Cross Mapping (CCM), to identify the ranges of HCHO/NO₂ across
China based on ground observations and remote sensing datasets. The ranges of transitional regime estimated by polynomial simulation and CCM were [1.0, 1.9] and [1.0, 1.8]. Since 2013, ozone formation regime has changed to the transitional and
20 NO_x-limited regime all over China, indicating ozone concentrations across China were mainly controlled by NO_x. However, despite the NO₂ concentrations, HCHO concentrations continuously exert a positive influence on ozone concentrations under transitional and NO_x-limited regimes. Under the circumstance of national NO_x-reduction policies, the increase of VOCs became the major driver for the soaring ozone pollution across China. For an effective management of ozone pollution across China, the emission-reduction of VOCs and NO_x should be equally considered.

25 1 Introduction

With the significant improvement of PM_{2.5} pollution, surface ozone has become a major airborne pollutant across China since 2017 (Li et al., 2019a; Lu et al., 2020). Due to its severe threat to public health even during a short-period exposure, ozone pollution has received growing emphasis from governments and scholars (Liu et al., 2018a; Xie et al., 2019). In the past several years, spatiotemporal distribution of ozone concentrations (Wu and Xie, 2017; Shen et al., 2019a), and the
30 influence of meteorological conditions (Chen et al., 2019b; Cheng et al., 2019; Chen et al., 2020) and anthropogenic



emissions (Cheng et al., 2018; Li et al., 2019a; Li et al., 2020) on ozone concentrations have been massively studied. However, due to the highly complicated ozone formation regime, effective ozone control remains challenging.

Different from $PM_{2.5}$, whose main precursor are NO_x , VOCs and SO_2 , the formation and decomposition of ozone are closely related to two types of precursors, VOCs and NO_x . There is a diversity of reactions between VOCs and NO_x under different meteorological conditions and concentration scenarios (Wang et al., 2017). Since VOCs and NO_x can either promote or restrict ozone production, VOCs/ NO_x is crucial for surface ozone concentrations. However, the thresholds at which VOCs/ NO_x may promote or restrict ozone production remain unclear (Jin et al., 2017; Schroeder et al., 2017). For instance, under a specific VOCs/ NO_x scenario, the reduction of NO_x may conversely increase surface ozone concentrations (Sillman et al., 1990; Kleinman, 1994). Furthermore, given the large variations of meteorological conditions and ozone level across China, the effects of VOCs/ NO_x on surface ozone concentrations also demonstrate notable spatiotemporal patterns. In this case, a comprehensive understanding of how the variations of VOCs and NO_x could influence ozone concentrations under different VOCs/ NO_x circumstances is crucial for setting effective emission-reduction policies accordingly in different regions.

To examine the complicated non-linear relationship between ozone concentrations and multiple precursors, a large body of studies has been conducted (Duncan et al., 2010; Choi et al., 2012; Pusede and Cohen, 2012; Chang et al., 2016; Jin et al., 2020). Through small-scale experiments, NO_2 and HCHO proved to be effective proxies for NO_x and VOCs (Sillman et al., 1990; Martin et al., 2004). Since NO_2 and HCHO can be monitored using remote sensing data, the two precursors have been increasingly considered in ozone-precursor sensitivity research (Jin et al., 2020; Zhang et al., 2020). Cheng et al. (2018) proved that NO_2/NO presented a good consistence with long-term ozone concentrations in Beijing. However, NO was not an easily recordable precursor based on satellite observations and not applicable in large-scale monitoring. Cheng et al. (2019) suggested that satellite retrieved HCHO/ NO_2 was strongly correlated with surface ozone concentrations in Beijing. Different HCHO/ NO_2 indicates distinct ozone formation regimes, including VOC-limited, transitional and NO_x -limited regimes. For VOC-limited (NO_x -saturated) regime, the control of VOCs emissions leads to the reduction of organic radicals (RO_2), the RO_2 - NO_x reactions and thus ozone concentrations (Milford et al., 1989). On the contrary, the decrease of NO_x promotes VOCs-CO reaction, leading to the increase of ozone concentration (Kleinman, 1994). For NO_x -limited regime, the reduction of NO_x slows down NO_2 photolysis, which products free oxygen atoms for ozone formation, and reduces ozone concentrations. The variations of VOCs exert limited influences on ozone concentrations for this regime (Kleinman, 1994). For transitional (VOCs- NO_x mixed) regime, both VOCs and NO_x impose positive influences on ozone concentrations. Since the transitional regime divides VOC-limited and NO_x -limited regimes, the estimation of the transitional regime range plays a key role to identify different ozone formation regimes.

Duncan et al. (2010) calculated the transitional regime range as [1.0, 2.0] using the Community Multiscale Air Quality Modeling System (CMAQ) model, whose uncertainties may influence the estimation accuracy (Schroeder et al., 2017). Jin et al. (2020) employed a polynomial model and calculated the transitional regime range over U.S. urban areas as [3.2, 4.1] based on decades of remote sensing and ground observation data. However, given the notable difference of meteorological



65 conditions, ozone levels and the composition of precursors across different countries, whether the transitional regime range
extracted in US is applicable to other countries remains unclear. Furthermore, the polynomial model may ignore the
complicated inner interactions between multiple precursors, meteorological factors and ozone concentrations in the
atmospheric environment (Chen et al., 2020), and may lead to large uncertainties. Consequently, ozone-precursors sensitivity,
especially the transitional regime range across China, requires further in-depth analysis.

70 To this end, this research attempts to investigate the spatiotemporal variations of ozone formation regimes across China and
identify the transitional regime range of HCHO/NO₂ based on the cross-verification of multiple models. Firstly, long-term
variations of HCHO and NO₂ across China were analyzed. Next, the datasets of HCHO, NO₂ and ozone were examined
using a polynomial model and a causality model respectively to reveal the crucial thresholds of HCHO/NO₂ that separates
the NO_x-limited, VOCs-limited, and transitional regimes. Specifically, due to the large area of China and potential spatial
75 variations in ozone formation regimes, we respectively investigated ozone formation regimes in several major regions,
including the North China Plain (NCP), Yangtze River delta (YRD), Pearl River delta (PRD), and Sichuan Basin (SCB) (The
geographical locations of four megacity clusters were shown in Figure 1), to explore the spatiotemporal variations of ozone
formation regimes. Meanwhile, we also compared the ozone formation regimes in urban and rural areas. This research sheds
useful lights for better modeling complicated ozone-precursors relationship, understanding the major drivers for enhanced
80 ozone pollution and implementing specific emission-reduction measures to mitigate ozone pollution across China.

Figure 1 inserted here.

2 Materials and methods

2.1 Data sources

In this study, OMI HCHO/NO₂ datasets were employed for exploring the spatiotemporal variations of HCHO and NO₂ in
85 China and calculating HCHO/NO₂. We connected surface ozone network data to HCHO, NO₂ and HCHO/NO₂, which
served as the input data for running third-polynomial model and Convergent Cross Mapping (CCM). MODIS land cover
product provided the spatial distribution of urban areas, which was employed for identifying urban and rural pixels.

2.1.1 OMI HCHO/NO₂

Ozone Monitoring Instrument (OMI), on board the Aura satellite, monitors global solar backscatter in the UV/Vis domain
90 (270-500 nm). OMI provides daily global observations, which crosses the equator at 13:38 (local time) (Levelt et al., 2006).
In this study, we employed daily level-3 gridded OMI HCHO product (OMHCHOd) from the Smith Astrophysical
Observatory (SAO) (González Abad et al., 2015). The HCHO vertical columns are the weighted mean values for the 0.1° ×
0.1° grid. Backscattered solar radiation, ranging from 328.5-356.5 nm, was used for fitting HCHO slant columns. Air mass
factors (AMFs) were employed for converting HCHO slant columns to vertical columns (González Abad et al., 2015). The
95 validation report suggested that the error of this product was effectively controlled within 30% over polluted areas (González



Abad et al., 2015), and validated for detecting long-term variations of HCHO columns (Zhu et al., 2017; Shen et al., 2019b). The daily level-3 gridded OMI NO₂ product (OMNO2d), provided by NASA's Goddard Space Flight Center, were utilized in this study (Bucsela et al., 2013; Lamsal et al., 2014). The spatial resolution of OMNO2d is 0.25° and each grid is generated as the weighted average of the corresponding level-2 data pixels (Krotkov et al., 2017). Differential optical absorption spectroscopy (DOAS) was employed for retrieving the NO₂ slant columns, which were successively transformed into tropospheric and stratospheric vertical columns through AMFs (Bucsela et al., 2013). The OMI NO₂ column product agrees well with other satellite products, and its overall uncertainties range from 30%-60% (Bucsela et al., 2013; Lamsal et al., 2014). To reduce uncertainties, we only selected those OMI HCHO and NO₂ data that (1) passed quality checks, (2) had a cloud coverage less than 30%, (3) had a solar zenith angle less than 60°, and (4) were not affected by row anomalies for this study (Kroon et al., 2011; Zhu et al., 2014; Krotkov et al., 2017). The May-to-September OMI HCHO and NO₂ products were acquired from NASA's Goddard Earth Sciences Data and Information Services Center (<https://disc.gsfc.nasa.gov/>).

2.1.2 Surface ozone network data

The May-to-September hourly surface ozone concentrations from 2014 to 2019, were obtained from the China Ministry of Ecology and Environment (MEE) (<https://quotsoft.net/air/>). The unit of surface ozone concentrations in this dataset is µg/m³. The network had 1633 monitoring stations, which were distributed in 330 cities across China in 2019. We used the observation data from 13:00 to 14:00 at local time to match the overpass time of OMI. This dataset has been employed in many studies to investigate the variations of surface ozone concentrations in China (Li et al., 2019a; Shen et al., 2019a; Lu et al., 2020).

2.1.3 MODIS land cover product

The annual MODIS land cover product (MCD12C1) with a spatial resolution of 0.05° from 2005 to 2019 was employed for extracting urban and rural areas. The urban and water pixels from the International Geosphere-Biosphere Program (IGBP) classification layer were employed for the following processing. The land cover product was generated based on a decision tree algorithm with boosting techniques, and its overall accuracy was about 75% (Palmer et al., 2015; Bajocco et al., 2018). MCD12C1 product was obtained from NASA's Earth System Data and Information System (<https://earthdata.nasa.gov/>).

2.1.4 Data pre-processing

Due to different spatial resolution of OMI HCHO, OMI NO₂ and MCD12C1, bilinear interpolation method was used for resampling all above-mentioned products to the same spatial size (0.25° × 0.25°). Meanwhile, we also calculated mean hourly surface ozone concentrations on the 0.25° × 0.25° grid (Figure 1).



2.2 Methods

125 Chemical transport models, such as the global chemical transport model (GEOS-Chem) (Jin et al., 2017; Li et al., 2019a) and
the Community Multiscale Air Quality Modeling System (CMAQ) (Duncan et al., 2010), have been frequently employed for
exploring the ozone sensitivity to VOCs and NO_x. However, there were large biases in estimating the range of transitional
regime based on chemical transport models (Jin et al., 2017; Jin et al., 2020), due to the uncertainties of emission inventory
and the setting of model parameters. Employing observation data alone could effectively overcome these limitations, and the
130 relationships between ozone and its precursors were fitted using linear and polynomial models (Sun et al., 2018; Jin et al.,
2020). Meanwhile, Convergent Cross Mapping (CCM) (Sugihara et al., 2012), as a robust causality analysis model, has been
widely employed for quantifying the influences of meteorological factors on surface ozone and PM_{2.5} concentrations (Chen
et al., 2018; Chen et al., 2019b; Chen et al., 2020), which is a promising tool for investigating the relationships between
ozone and its precursors. To increase the reliability of estimated range of transitional regime, both the polynomial model and
135 CCM were employed in this research. We employed the third-order polynomial model for fitting surface ozone
concentrations to the indicator of HCHO/NO₂. CCM was employed for quantifying the influences of HCHO and NO₂ on
surface ozone concentrations, and Wilcoxon test (Gehan, 1965) was used for examining whether the differences between the
causality of HCHO and NO₂ on ozone concentrations at different ranges of HCHO/NO₂ was significant. Since the algorithms
of the two models are quite different, their cross-verification provides useful reference for their reliability. Meanwhile,
140 Mann-Kendall (M-K) test (Kendall, 1948) was employed for exploring the spatiotemporal variations of HCHO, NO₂ and
ozone formation regimes in China. Furthermore, we extracted all urban and rural areas in China and compared the
differences of ozone formation regimes over these two types of areas. The workflow of the models employed in this study is
shown in Figure 2.

Figure 2 inserted here.

145 2.2.1 Estimating the transitional range of ozone formation regime using polynomial simulation

HCHO and NO₂ are considered as proxies for VOCs and NO_x, respectively. HCHO/NO₂, as an effective indicator, has been
widely employed for determining ozone formation regimes (Duncan et al., 2010; Jin and Holloway, 2015; Jin et al., 2017;
Cheng et al., 2019; Jin et al., 2020). Pusede and Cohen (2012) suggested that ozone exceedance probability (OEP) was an
effective indicator to interpret the ozone sensitivity to its precursors. The indicator is defined as the proportion of non-
150 attainment events (surface ozone concentrations exceeding 200 µg/m³) in total events at a given range of HCHO/NO₂:

$$OEP = \frac{Events_{non-attainment}}{Events_{attainment} + Events_{non-attainment}} \quad (1)$$

where $Events_{attainment}$ and $Events_{non-attainment}$ denote the attainment and non-attainment events, respectively (Pusede and Cohen,
2012; Jin et al., 2020).



In this study, we used a third-order polynomial model (Jin et al., 2020) to explore the quantitative relationships between
155 HCHO/NO₂ and ozone exceedance probability. There were 174868 paired observations of surface ozone concentrations and
HCHO/NO₂ from 2014 to 2019. The peak of fitting curve highlights the turning point of VOC-limited and NO_x-limited
regimes (Jin et al., 2020). The range of HCHO/NO₂, which corresponded to the top 10% ozone exceedance probability, was
defined as the transitional regime.

2.2.2 Estimating the transitional range of ozone formation regime using Convergent Cross Mapping

160 We also employed Convergent cross mapping (CCM) (Sugihara et al., 2012), which could reduce the influences of other
factors such as meteorological conditions (Chen et al., 2019b; Chen et al., 2020), to extract the causal influences of HCHO
and NO₂ on surface ozone concentrations. Thanks to its capability of detecting weak coupling, CCM is advantageous for
reliably comparing the influences of different meteorological factors on surface ozone concentrations (Chen et al., 2020).
Therefore, we employed CCM to compare the sensitivity of ozone to HCHO and NO₂ at different ranges of HCHO/NO₂.
165 CCM utilizes convergent maps to demonstrate the bidirectional coupling between the time series of two variables. A
convergent curve indicates that one variable imposes influences on the other variable, whilst a non-convergent curve denotes
no causality between two variables. CCM calculates cross map skill (ρ value) that explains the quantitative relationships.
Number of dimensions for the attractor reconstruction (E), time lag (τ) and number of nearest neighbors to use for prediction
(b) are required parameters for CCM. According to previous studies (Chen et al., 2020; Chen et al., 2019b), E , τ and b was
170 set as 3, 2 and 4, respectively. Since the existence of missing values imposes negative impacts on CCM results, only the
consecutive time series were retained for this research. There were 1660 observation records of HCHO time series, NO₂ time
series and corresponding surface ozone time series. CCM was implemented using “pyEDM” package in Python. Wilcoxon
test (Gehan, 1965) was used to examine whether the differences of ρ values between HCHO and NO₂ were significant at the
given HCHO/ NO₂. No significant difference was regarded as the transitional regime, while significant difference indicated
175 the VOC-limited or NO_x-limited regime.

2.2.3 Trend analysis

Mann-Kendall (M-K) (Kendall, 1948) test, which has been used in recent studies on HCHO and NO₂ (Cheng et al., 2019;
Wang et al., 2019; Zeb et al., 2019), was employed to estimate the significance of trends. M-K test is capable of processing
samples with random distributions and mitigating the effects of outliers. Z value is calculated as follow:

180

$$Z = \begin{cases} \frac{S - 1}{\sqrt{\text{Var}(S)}} (S > 0) \\ \frac{S + 1}{\sqrt{\text{Var}(S)}} (S < 0) \end{cases} \quad (2)$$

where S denotes the statistic to be tested, $\text{Var}(S)$ stands for the variance of S .



The sign and absolute value of Z indicate the direction and significance of trends, respectively. Specifically, the positive and negative values of Z indicate the upward and downward trend. 1.28, 1.64 and 2.32 are the threshold values of $|Z|$, indicating the trends of samples pass the tests at 90%, 95% and 99%, respectively.

185 2.2.4 Comparison of ozone formation regimes in urban and rural areas in China

To compare the differences of ozone formation regimes in urban and rural areas in China, the key step is to extract urban and rural pixels, respectively. Urban pixels were used for buffer analysis (Imhoff et al., 2010) to identify rural pixels. Following Peng et al. (2018), two buffers were set for urban pixels to extract candidate rural pixels (Figure 3). We set the size of each buffer as 27.75 km, which was close to the size of the $0.25^\circ \times 0.25^\circ$ grid ($27.75 \text{ km} \approx 0.25^\circ$). The first and second buffers
190 were determined as the urban fringes and candidate rural areas, respectively. Water pixels were firstly removed from candidate rural areas to avoid following uncertainties. Consequently, rural areas were regarded as buffers of 27.75-55.50 km surrounding urban areas. The use of two buffers not only assisted a complete separation of the urban and rural areas, but also minimized the uncertainties of meteorological conditions (Yao et al., 2019).

Figure 3 inserted here.

195 3 Results

3.1 Spatial and temporal variations of HCHO and NO₂

Given the national Clean Air Action implemented in 2013, we set this year as a break point to explore the spatial and temporal variations of HCHO and NO₂ in 2005-2012 and 2013-2019, respectively. Figure 4 shows the spatial distribution of HCHO in the two periods. The mean HCHO values during the period of 2005-2012 and 2013-2019 were 4.335×10^{15}
200 molec/cm^2 and $4.845 \times 10^{15} \text{ molec/cm}^2$, characterized with a 12% increase. Both periods presented an increasing trends of HCHO, and the averaged value during the two periods were $0.164 \times 10^{15} \text{ molec/cm}^2 \text{ year}^{-1}$ and $0.213 \times 10^{15} \text{ molec/cm}^2 \text{ year}^{-1}$ (Figure 5). A faster increasing trend was detected during the period of 2013-2019. The variation trend of HCHO agreed well with previous studies (Jin and Holloway, 2015; Shen et al., 2019b). We also calculated the overall linear trends of HCHO in four megacity clusters from 2005 to 2019 (Figure 6). The largest and smallest increasing trends were shown in NCP and
205 SCB, with a mean value of $0.136 \times 10^{15} \text{ molec/cm}^2 \text{ year}^{-1}$ and $0.046 \times 10^{15} \text{ molec/cm}^2 \text{ year}^{-1}$. The increasing trend of YRD and PRD were $0.066 \text{ molec/cm}^2 \text{ year}^{-1}$ and $0.058 \text{ molec/cm}^2 \text{ year}^{-1}$, respectively. Meanwhile, reversed trends were detected for NO₂ during the two periods (Figure 5), which was consistent with previous studies (Jin and Holloway, 2015; Li et al., 2019a). From 2005 to 2012, the averaged NO₂ was $2.027 \times 10^{15} \text{ molec/cm}^2$ and the annually mean increasing trend was $0.098 \times 10^{15} \text{ molec/cm}^2 \text{ year}^{-1}$. Thanks to the implementation of Clean Air Action, the averaged NO₂ were reduced to 1.900
210 $\times 10^{15} \text{ molec/cm}^2$, with a decreasing trend of $-0.029 \times 10^{15} \text{ molec/cm}^2 \text{ year}^{-1}$ from 2013 to 2019. Except for SCB, all other megacity clusters presented significant downward trends of NO₂ from 2005 to 2019. Amongst these megacity clusters, NO₂



in YRD demonstrated the largest decreasing trend of 0.104×10^{15} molec/cm² year⁻¹. NO₂ in NCP and PRD decreased by 0.010×10^{15} molec/cm² year⁻¹ and 0.092×10^{15} molec/cm² year⁻¹, respectively. A slightly increasing trend of 0.012×10^{15} molec/cm² year⁻¹ was detected in SCB (Figure 7).

215

Figure 4 inserted here.

Figure 5 inserted here.

Figure 6 inserted here.

Figure 7 inserted here.

3.2 Transitional range of ozone formation regime

220 According to HCHO/NO₂, We divided the paired observations into 200 bins for the whole country and 100 bins for these megacity clusters, and the ozone exceedance probability was calculated for each bin. The third-order polynomial was employed for fitting ozone exceedance probability to HCHO/NO₂. As shown in Figure 8a, the peak of the fitting curve was 1.4, and the vertical shaded area indicated that the transitional regime over China ranged from 1.0 to 1.9. In addition to the regime range extracted at the national scale, we also examined the range of ozone formation regimes in four major megacity
225 clusters. The range of transitional regime for NCP, YRD, PRD and SCB was [1.2, 2.1], [1.0, 1.9], [0.9, 1.8] and [1.1, 2.0] respectively, which was generally consistent with the range at the national scale. The small differences between four megacity clusters across China suggested the range of transitional regime at the national scale [1.0, 1.9] can be employed to regional or local scale research, if small-scale data and investigation was not available.

Statistical bootstrapping was used for estimating the uncertainty of the fitting model. Specifically, we iteratively extracted 50
230 randomly selected subsets from the paired observations to run the model, and the uncertainty was defined as two standard deviations from the peak of the fitting curve. The uncertainty for the third-polynomial model was 0.4, indicating a significant nonlinear relationship between ozone exceedance probability and HCHO/NO₂.

Due to the limited data used for running CCM, we set the bin size of HCHO/NO₂ as 0.2 for collecting sufficient ρ values to conduct Wilcoxon test. As shown in Figure 8b, there were no significant difference between ρ of HCHO and NO₂ when
235 HCHO/NO₂ ranged from 0.9 to 1.9, which indirectly defined the range of transitional regime. For HCHO/NO₂ < 0.9, ρ of HCHO was notably higher than that of NO₂, and this range was regarded as VOC-limited regime. Similarly, HCHO/NO₂ > 1.9 suggested the NO_x-limited regime. Through the cross verification, it was an important finding that the range of transitional ozone formation regime estimated using the third-order polynomial model and CCM was highly close, indicating the reliability of the extracted range.

240

Figure 8 inserted here.

3.3 Ozone formation regimes in China

NO₂ demonstrated a significant downward trend since 2013, while HCHO kept the increasing trend during the entire study period. Consequently, HCHO/NO₂ increased in a majority of regions across China. Specifically, the annually increasing



trend of HCHO/NO₂ in NCP, YRD and PRD was 0.035 year⁻¹, 0.023 year⁻¹ and 0.034 year⁻¹, respectively. Meanwhile, there
245 were no significant trends in SCB during this period (Figure 9). The variations of HCHO/NO₂ indicated the shrinkage of
VOC-limited regime and the expansion of transitional and NO_x-limited regimes. Since the range of transitional regime
estimated by third-order polynomial model and CCM was very close and the former included more reliable observation data,
[1.0, 1.9] was employed for identifying different ozone formation regimes. In 2005, areas with the VOC-limited regime were
concentrated in NCP, YRD and PRD. The proportions of areas with the VOC-limited regime in NCP, YRD and PRD were
250 26%, 16% and 6%, respectively. Areas with the transitional regime were mainly distributed in the marginal regions of those
megacity clusters, and scatteredly distributed in SCB. Areas with the transitional regime occupied 60%, 50%, 14% and 20%
in NCP, YRD, PRD and SCB. NO_x-limited regime dominated other areas (Figure 10a). In 2019, areas with the VOCs-
limited regime decreased significantly, and was simply found in the fringe areas of NCP and YRD. The proportion of the
VOCs-limited regime in NCP and YRD was 2% and 9%, respectively. The transitional regime was widely distributed NCP,
255 YRD and SCB, and occupied the 71 %, 56% and 36% of the total areas. The NO_x-limited regime still spread over a majority
of China (Figure 10a). We calculated the annually mean ρ of HCHO and NO₂ over those megacity clusters from 2014 to
2019 (Figure 10b). For all megacity clusters, the ρ of NO₂ was higher than HCHO, indicating that NO₂ was the dominant
factor for surface ozone concentrations. Both models suggested that NO₂ played a more important role in affecting surface
ozone concentrations than HCHO. In the past several years, NO_x-oriented emission-reduction has been conducted across
260 China, leading to the continuous decrease of NO_x concentrations. Since both VOCs and NO_x imposed positive influences on
surface ozone concentrations under the transitional and NO_x-limited ozone formation regime, the upward trend of HCHO
across China might explain recent soaring ozone concentrations across China (Shen et al., 2019a; Lu et al., 2020). It is noted
that the difference between the ρ of NO₂ and HCHO decreased notably in NCP and YRD. This may be attributed to the
following reason. NCP and YRD are the regions that received severe PM_{2.5} pollution and strict NO_x-reduction policies have
265 been conducted since 2013. With the remarkably reduced NO₂ concentrations, the variations of HCHO concentrations plays
an increasingly important role in affecting ozone concentrations in NCP and YRD. The reduction of VOCs emissions is key
for an effective management of surface ozone pollution in NCP and YRP.

Figure 9 inserted here.

Figure 10 inserted here.

270 3.4 Variations of ozone formation regimes in urban and rural areas

Previous studies suggested that the differences of ozone formation regimes existed between urban and rural areas (Tong et
al., 2017; Liu et al., 2018b; Cheng et al., 2019). We extracted HCHO and NO₂ columns in urban and rural pixels in those
megacity clusters, and calculated the annually averaged HCHO/NO₂ (Figure 11). For NCP, HCHO/NO₂ in urban areas was
higher than 1.0 since 2015, indicating a transformation from VOC-limited to transitional regime. The increase of
275 HCHO/NO₂ was attributed to the reversed variation trends of HCHO and NO₂. The rising HCHO resulted from the increase
of anthropogenic emissions and biogenic volatile organic compound (BVOC) (Shen et al., 2019b; Wang et al., 2020), while



the implementation of Clean Air Action imposed notable influences on the decrease of NO_2 (Chen et al., 2019a). HCHO/NO_2 in rural areas was in the range of [1.0, 1.9], indicating rural areas were occupied by the transitional regime from 2005 to 2019. For YRD, which was occupied by the transitional regime, no variation of ozone formation regime was found
280 in urban areas. In rural areas, HCHO/NO_2 temporally exceeded the threshold of 1.9 from 2016 to 2018, indicating the ozone formation regime changed from transitional to NO_x -limited. This phenomenon was attributed to the slight decline of HCHO, which might be attributed to the restrictions on crop residue burning in this area (Zhuang et al., 2018; Shen et al., 2019b). Due to the large differences of NO_2 concentrations, the urban and rural areas in PRD was dominated by transitional regime and NO_x -limited regime. For SCB, HCHO/NO_2 in both urban and rural areas fluctuated around the threshold value of 1.9,
285 and no significant difference between urban and rural areas was found.

Figure 11 inserted here.

4 Discussion

This research employed CCM and third-order polynomial model to estimate the transitional regime of ozone formation across China and the calculated range of HCHO/NO_2 was [0.9, 1.9] and [1.0, 1.9], respectively. Given the lack of actual
290 reference data, the close output from two different models provides a reliable reference for better understanding ozone formation regimes. Our findings were generally consistent with previous studies. For US, Duncan et al. (2010) and Choi et al. (2012) employed the OMI and GOME-2 data, whose 0.25° resolution was close to this research, and calculated the range of transitional regime as [1.0, 2.0]. The similar range of transitional regime in US and China further proved the reliability of the calculated range [1.0, 1.9] at a national scale. On the other side, the range of transitional regime can vary significantly across
295 regions (Schroeder et al., 2017; Jin et al., 2020). Sun et al. (2018) employed station-based data and calculated the range of transitional regime in Anhui Province, China as [1.3, 2.8], which was notably higher than the range across China. Jin et al. (2020) calculated the range of transitional regime in several major regions in the US using QA4ECV dataset, whose spatial resolution was 0.125° , and the output [3.2, 4.1] was much larger than the averaged range of transitional regime across US. One reason could be the severe ozone pollution in mega cities, leading to different ranges of transitional regime. Meanwhile,
300 the calculated range of transitional regime is closely related to the spatial resolution of employed HCHO and NO_2 data, and high-resolution data are more advantageous in extracting the sensitivity of ozone concentrations to precursors at the local scale (Martin et al., 2004; Jin et al., 2017; Jin et al., 2020). In addition to the locations and the spatial resolution of data sources, the uncertainties of OMI HCHO and NO_2 datasets might impose negative influences on the estimation of transitional regime range (Duncan et al., 2010; Jin et al., 2017; Schroeder et al., 2017; Jin et al., 2020). Firstly, errors exist in
305 the retrieval of HCHO and NO_2 vertical columns. Secondly, vertical mixing was not homogeneous, weakening the capability of using HCHO and NO_2 vertical columns to explore the near-surface ozone-precursors sensitivity. Therefore, future improvement of earth observation techniques and the spatiotemporal resolution of HCHO and NO_2 products can further enhance the accuracy of the estimated range of transitional regime. In general, according to the cross-verification and



310 comparison with previous studies, [1.0, 1.9] from this research is a reliable range for transitional ozone formation regime across China and can be used as an approximate criterion to follow when implementing national emission-reduction policies. On the other hand, given the potential variations of transitional regimes in different regions, when conducting small-scale research, the range of [1.0, 1.9] may be adapted accordingly based on local data.

315 Previous studies on the range of ozone formation regimes were mainly conducted using statistical models or chemical transport models. For this research, we employed both a statistical and a causality models to cross-verify the range of transitional regimes. Despite a relatively high fitting accuracy in terms of uncertainties, the findings from these studies could not be effectively compared or interpreted, due to the lack of reliable reference data. To this end, as well as numerical models, lab experiments should also be considered to extract more precise description of ozone-precursors relationship. With the rapid development of atmospheric science, smog chambers have been increasingly employed to investigate complicated interactions between multiple precursors. By setting specific meteorological conditions (e.g. temperature and humidity) and gradually adjusting the proportion of different precursors, how the proportion of NO₂ and HCHO affect ozone formation regime can be better explained in a theoretical environment. With more reliable experimental reference data, the model-based analysis on the range of transitional regime at local, regional and national scale can be further improved accordingly.

320 According to the temporal variations of OMI NO₂ concentrations across China, a notable decreasing trend was observed in three major megacity clusters, NCP, YRD and PRD. These regions were heavily polluted by PM_{2.5} and the notable decrease of NO₂ was mainly attributed to the national Clean Air Action since 2013 (Zheng et al., 2018), which aimed to reduce PM_{2.5} concentrations by cutting NO_x emissions. Conversely, HCHO concentrations during this period increased remarkably across China, due to the combined effects of anthropogenic and biogenic emissions (Shen et al., 2019b; Wang et al., 2020). The distinct temporal variations of NO₂ and HCHO led to the increase of HCHO/NO₂, and the increase of transitional areas and NO_x-limited regime areas. From 2013-2019, all these regions were dominated by the transitional or NO_x-limited regimes. 330 Attributed to the long-term variation of formation regimes, a more complicated and fragmented spatial pattern was observed across China. Consequently, for an effective control of ozone pollution, the emission-reduction of both NO_x and VOCs is required. Especially for NCP and YRD, where the NO_x has been remarkably reduced, effective approaches for controlling VOCs emissions are essential for preventing ozone pollution. This finding was consistent with previous studies (Li et al., 2019b), which recommended the simultaneous reduction of NO_x and VOCs for mitigating the composite airborne pollution in China. Admittedly, compared with NO_x-reduction, the VOCs-reduction is more complicated and the output of anthropogenic VOCs reduction is more unpredictable. In this case, reducing biogenic VOCs emissions can also be a potential solution. VOCs emitted by vegetation takes up to 50% of total VOCs in the atmospheric environment, especially in summer. The key factor that may cause enhanced biogenic emissions is summertime high temperature in summer (Chen et al., 2020). Therefore, such projects as wind corridors or contingent artificial precipitation, which can effectively reduce 340 urban heat effects, should be implemented properly to avoid summertime heat waves and successive ozone pollution (e.g. summer, 2017).



The large spatial variations of HCHO/NO₂, especially the rapid increase of transitional regime areas across China, indicates that a unified NO_x-VOCs reduction strategy is not feasible for the entire country. Instead, to effectively reduce ozone concentrations, the specific proportion of NO_x and VOCs reduction should be carefully set according to local HCHO/NO₂.
345 Meanwhile, due to the large differences in vehicle and industrial emissions (Cheng et al., 2019), the concentration of NO_x is notably higher in urban areas. Therefore, the further reduction of NO_x emissions exerts a stronger influence on ozone reduction in rural areas compared to urban areas.

5 Conclusions

To better understand the spatiotemporal variations of ozone formation regimes across China, we employed the third-order
350 polynomial model and CCM to estimate the range of transitional regime from 2005 to 2019, the results of which were [1.0, 1.9] and [0.9, 1.9], respectively. The close outputs from two distinct models proved the reliability of the extracted range. At the regional scale, we also investigated the range of transitional regime in four megacity clusters and found the range in NCP, YRD, PRD and SCB demonstrated limited differences and was generally consistent with the range at the national scale. The reverse trends of HCHO and NO₂ led to the increase of HCHO/NO₂, indicating China was dominated by the transitional and
355 NO_x-limited regimes in recent years. We also found that the ρ of NO₂ was higher than HCHO at all megacities, suggesting that the reduction of NO_x emissions would become more effective in controlling surface ozone concentrations. Meanwhile, given the rising VOCs emissions, the simultaneous reduction of NO_x and VOCs would be more effective than the sole reduction of NO_x in mitigating ozone pollution. Finally, the comparison of ozone regimes in urban and rural areas suggested that the reduction of NO_x emissions would impose stronger impacts on the control of ozone pollution in rural areas.

360 Author contributions

Ruiyuan Li: Writing – original draft, Visualization, Formal analysis. Miaoqing Xu: Data curation, Visualization, Formal analysis. Manchun Li: Conceptualization, Methodology. Ziyue Chen: Writing – original draft, Conceptualization, Formal analysis. Bingbo Gao: Data curation, Formal analysis. Na Zhao: Data curation, Visualization. Qi Yao: Data curation, Visualization.

365 Competing interests

The authors declare that they have no conflict of interest.



Acknowledgements

This research is supported by the Beijing Natural Science Foundation (Grant No. 8202031), Open Fund of the State Key Laboratory of Remote Sensing Science (Grant No. OFSLRSS201926), the Open Fund of the State Key Laboratory of Resources and Environmental Information System.

References

- Bajocco, S., Smiraglia, D., Scaglione, M., Raparelli, E., Salvati, L.: Exploring the role of land degradation on agricultural land use change dynamics. *Sci. Total Environ.*, 636, 1373-1381, doi: 10.1016/j.scitotenv.2018.04.412, 2018.
- Bucsela, E., Krotkov, N., Celarier, E., Lamsal, L., Swartz, W., Bhartia, P., Boersma, K., Veefkind, J., Gleason, J., Pickering, K.: A new stratospheric and tropospheric NO₂ retrieval algorithm for nadir-viewing satellite instruments: applications to OMI. *Atmos. Meas. Tech. Discuss.*, 6, doi: 10.5194/amt-6-2607-2013, 2013.
- Chang, C.-Y., Faust, E., Hou, X., Lee, P., Kim, H.C., Hedquist, B.C., Liao, K.-J.: Investigating ambient ozone formation regimes in neighboring cities of shale plays in the Northeast United States using photochemical modeling and satellite retrievals. *Atmos. Environ.*, 142, 152-170, doi: 10.1016/j.atmosenv.2016.06.058, 2016.
- Chen, Z., Chen, D., Kwan, M.-P., Chen, B., Gao, B., Zhuang, Y., Li, R., Xu, B.: The control of anthropogenic emissions contributed to 80% of the decrease in PM_{2.5} concentrations in Beijing from 2013 to 2017. *Atmos. Chem. Phys.*, 19, 13519-13533, doi: 10.5194/acp-19-13519-2019, 2019a.
- Chen, Z., Li, R., Chen, D., Zhuang, Y., Gao, B., Yang, L., Li, M.: Understanding the causal influence of major meteorological factors on ground ozone concentrations across China. *J. Clean Prod.*, 242, doi: 10.1016/j.jclepro.2019.118498, 2020.
- Chen, Z., Xie, X., Cai, J., Chen, D., Gao, B., He, B., Cheng, N., Xu, B.: Understanding meteorological influences on PM_{2.5} concentrations across China: a temporal and spatial perspective. *Atmos. Chem. Phys.*, 18, 5343-5358, doi: 10.5194/acp-2017-376, 2018.
- Chen, Z., Zhuang, Y., Xie, X., Chen, D., Cheng, N., Yang, L., Li, R.: Understanding long-term variations of meteorological influences on ground ozone concentrations in Beijing During 2006-2016. *Environ. Pollut.*, 245, 29-37, doi: 10.1016/j.envpol.2018.10.117, 2019b.
- Cheng, N., Chen, Z., Sun, F., Sun, R., Dong, X., Xie, X., Xu, C.: Ground ozone concentrations over Beijing from 2004 to 2015: Variation patterns, indicative precursors and effects of emission-reduction. *Environ. Pollut.*, 237, 262-274, doi: 10.1016/j.envpol.2018.02.051, 2018.
- Cheng, N., Li, R., Xu, C., Chen, Z., Chen, D., Meng, F., Cheng, B., Ma, Z., Zhuang, Y., He, B., Gao, B.: Ground ozone variations at an urban and a rural station in Beijing from 2006 to 2017: Trend, meteorological influences and formation regimes. *J. Clean Prod.*, 235, 11-20, doi: 10.1016/j.jclepro.2019.06.204, 2019.



- Choi, Y., Kim, H., Tong, D., Lee, P.: Summertime weekly cycles of observed and modeled NO_x and O_3 concentrations as a function of satellite-derived ozone production sensitivity and land use types over the Continental United States. *Atmos. Chem. Phys.*, 12, 6291-6307, doi: 10.5194/acp-12-6291-2012, 2012.
- 400 Duncan, B.N., Yoshida, Y., Olson, J.R., Sillman, S., Martin, R.V., Lamsal, L., Hu, Y., Pickering, K.E., Retscher, C., Allen, D.J., Crawford, J.H.: Application of OMI observations to a space-based indicator of NO_x and VOC controls on surface ozone formation. *Atmos. Environ.*, 44, 2213-2223, doi: 10.1016/j.atmosenv.2010.03.010, 2010.
- Gehan, E.A.: A generalized Wilcoxon test for comparing arbitrarily singly-censored samples. *Biometrika*, 52, 203-224, doi: 10.1093/biomet/52.1-2.203, 1965.
- 405 González Abad, G., Liu, X., Chance, K., Wang, H., Kurosu, T., Suleiman, R.: Updated Smithsonian Astrophysical Observatory Ozone Monitoring Instrument (SAO OMI) formaldehyde retrieval. *Atmos. Meas. Tech.*, 8, 19-32, doi: 10.5194/amt-8-19-2015, 2015.
- Imhoff, M.L., Zhang, P., Wolfe, R.E., Bounoua, L.: Remote sensing of the urban heat island effect across biomes in the continental USA. *Remote Sens. Environ.*, 114, 504-513, doi: 10.1016/j.rse.2009.10.008, 2010.
- 410 Jin, X., Fiore, A., Boersma, K.F., De Smedt, I., Valin, L.: Inferring changes in summertime surface ozone- NO_x -VOC chemistry over U.S. urban areas from two decades of satellite and ground-based observations. *Environ. Sci. Technol.*, 54, 11, 6518-6529, doi: 10.1021/acs.est.9b07785, 2020.
- Jin, X., Fiore, A.M., Murray, L.T., Valin, L.C., Lamsal, L.N., Duncan, B., Boersma, K.F., De Smedt, I., Abad, G.G., Chance, K., Tonnesen, G.S.: Evaluating a Space-Based Indicator of Surface Ozone- NO_x -VOC Sensitivity Over Midlatitude Source Regions and Application to Decadal Trends. *J. Geophys. Res.-Atmos.*, 122, 10-461, doi: 10.1002/2017JD026720, 2017.
- 415 Jin, X., Holloway, T.: Spatial and temporal variability of ozone sensitivity over China observed from the Ozone Monitoring Instrument. *J. Geophys. Res.-Atmos.*, 120, 7229-7246, doi: 10.1002/2015JD023250, 2015.
- Kendall, M.G.: Rank correlation methods. doi: 10.1111/j.2044-8295.1934.tb00727.x, 1948.
- 420 Kleinman, L.I.: Low and high NO_x tropospheric photochemistry. *J. Geophys. Res.-Atmos.*, 99, 16831-16838, doi: 10.1029/94JD01028, 1994.
- Kroon, M., De Haan, J., Veefkind, J., Froidevaux, L., Wang, R., Kivi, R., Hakkarainen, J.: Validation of operational ozone profiles from the Ozone Monitoring Instrument. *J. Geophys. Res.-Atmos.*, 116, doi: 10.1029/2010JD015100, 2011.
- Krotkov, N.A., Lamsal, L.N., Celarier, E.A., Swartz, W.H., Marchenko, S.V., Bucsel, E.J., Chan, K.L., Wenig, M., Zara, M.: The version 3 OMI NO_2 standard product. *Atmos. Meas. Tech.*, 3133-3149, doi: 10.5194/amt-10-3133-2017, 2017.
- 425 Lamsal, L., Krotkov, N., Celarier, E., Swartz, W., Pickering, K., Bucsel, E., Gleason, J., Martin, R., Philip, S., Irie, H.: Evaluation of OMI operational standard NO_2 column retrievals using in situ and surface-based NO_2 observations. *Atmos. Chem. Phys.*, 14, 11587, doi: 10.5194/acp-14-11587-2014, 2014.
- Levelt, P.F., van den Oord, G.H., Dobber, M.R., Malkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J.O., Saari, H.: The ozone monitoring instrument. *IEEE Trans. Geosci. Remote Sensing* 44, 1093-1101, doi: 10.1109/TGRS.2006.872333, 2006.
- 430



- Li, K., Jacob, D.J., Liao, H., Shen, L., Zhang, Q., Bates, K.H.: Anthropogenic drivers of 2013–2017 trends in summer surface ozone in China. *Proc. Natl. Acad. Sci. U.S.A.*, 116, 422–427, doi: 10.1073/pnas.1812168116, 2019a.
- Li, K., Jacob, D.J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K.H., Zhang, Q., Zhai, S.: A two-pollutant strategy for
435 improving ozone and particulate air quality in China. *Nat. Geosci.*, 12, 906–910, doi: 10.1038/s41561-019-0464-x, 2019b.
- Li, K., Jacob, D.J., Shen, L., Lu, X., De Smedt, I., Liao, H.: 2013–2019 increases of surface ozone pollution in China: anthropogenic and meteorological influences. *Atmos. Chem. Phys.*, doi: 10.5194/acp-2020-298, 2020.
- Liu, H., Liu, S., Xue, B., Lv, Z., Meng, Z., Yang, X., Xue, T., Yu, Q., He, K.: Ground-level ozone pollution and its health impacts in China. *Atmos. Environ.*, 173, 223–230, doi: 10.1016/j.atmosenv.2017.11.014, 2018a.
- 440 Liu, Y., Li, L., An, J., Huang, L., Yan, R., Huang, C., Wang, H., Wang, Q., Wang, M., Zhang, W.: Estimation of biogenic VOC emissions and its impact on ozone formation over the Yangtze River Delta region, China. *Atmos. Environ.* 186, 113–128, doi: 10.1016/j.atmosenv.2018.05.027, 2018b.
- Lu, X., Zhang, L., Wang, X., Gao, M., Li, K., Zhang, Y., Yue, X., Zhang, Y.: Rapid Increases in warm-season surface ozone and resulting health impact in China Since 2013. *Environ. Sci. Technol. Lett.*, doi: 10.1021/acs.estlett.0c00171, 2020.
- 445 Martin, R.V., Fiore, A.M., Van Donkelaar, A.: Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions. *Geophys. Res. Lett.*, 31, doi: 10.1029/2004GL019416, 2004.
- Milford, J.B., Russell, A.G., McRae, G.J.: A new approach to photochemical pollution control: Implications of spatial patterns in pollutant responses to reductions in nitrogen oxides and reactive organic gas emissions. *Environ. Sci. Technol.*, 23, 1290–1301, doi: 10.1021/es00068a017, 1989.
- 450 Palmer, S.C., Odermatt, D., Hunter, P., Brockmann, C., Presing, M., Balzter, H., Tóth, V.: Satellite remote sensing of phytoplankton phenology in Lake Balaton using 10 years of MERIS observations. *Remote Sens. Environ.*, 158, 441–452, doi: 10.1016/j.rse.2014.11.021, 2015.
- Peng, J., Ma, J., Liu, Q., Liu, Y., Hu, Y., Li, Y., Yue, Y.: Spatial-temporal change of land surface temperature across 285 cities in China: An urban-rural contrast perspective. *Sci. Total Environ.*, 635, 487–497, doi: 10.1016/j.scitotenv.2018.04.105,
455 2018.
- Pusede, S.E., Cohen, R.C.: On the observed response of ozone to NO_x and VOC reactivity reductions in San Joaquin Valley California 1995–present. *Atmos. Chem. Phys.*, 12, 8323–8339, doi: 10.5194/acp-12-8323-2012, 2012.
- Schroeder, J.R., Crawford, J.H., Fried, A., Walega, J., Weinheimer, A., Wisthaler, A., Müller, M., Mikoviny, T., Chen, G., Shook, M., Blake, D.R., Tonnesen, G.S.: New insights into the column CH₂O/NO₂ ratio as an indicator of near-surface
460 ozone sensitivity. *J. Geophys. Res.-Atmos.*, 122, 8885–8907, doi: 10.1002/2017JD026781, 2017.
- Shen, L., Jacob, D.J., Liu, X., Huang, G., Li, K., Liao, H., Wang, T.: An evaluation of the ability of the Ozone Monitoring Instrument (OMI) to observe boundary layer ozone pollution across China: application to 2005–2017 ozone trends. *Atmos. Chem. Phys.*, 19, 6551–6560, doi: 10.5194/acp-19-6551-2019, 2019a.
- Shen, L., Jacob, D.J., Zhu, L., Zhang, Q., Zheng, B., Sulprizio, M.P., Li, K., De Smedt, I., González Abad, G., Cao, H., Fu,
465 T.M., Liao, H.: The 2005–2016 Trends of formaldehyde columns over China observed by satellites: Increasing



- anthropogenic emissions of volatile organic compounds and decreasing agricultural fire emissions. *Geophys. Res. Lett.*, 46, 4468-4475, doi: 10.1029/2019GL082172, 2019b.
- Sillman, S., Logan, J.A., Wofsy, S.C.: The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes. *J. Geophys. Res.-Atmos.*, 95, 1837-1851, doi: 10.1029/JD095iD02p01837, 1990.
- 470 Sugihara, G., May, R., Ye, H., Hsieh, C.-h., Deyle, E., Fogarty, M., Munch, S.: Detecting causality in complex ecosystems. *Science*, 338, 496-500, doi: 10.1126/science.1227079, 2012.
- Sun, Y., Liu, C., Palm, M., Vigouroux, C., Notholt, J., Hu, Q., Jones, N., Wang, W., Su, W., Zhang, W., Shan, C., Tian, Y., Xu, X., De Mazière, M., Zhou, M., Liu, J.: Ozone seasonal evolution and photochemical production regime in the polluted troposphere in eastern China derived from high-resolution Fourier transform spectrometry (FTS) observations. *Atmos. Chem. Phys.*, 18, 14569-14583, doi: 10.5194/acp-18-14569-2018, 2018.
- 475 Tong, L., Zhang, H., Yu, J., He, M., Xu, N., Zhang, J., Qian, F., Feng, J., Xiao, H.: Characteristics of surface ozone and nitrogen oxides at urban, suburban and rural sites in Ningbo, China. *Atmos. Res.*, 187, 57-68, doi: 10.1016/j.atmosres.2016.12.006, 2017.
- Wang, H., Wu, Q., Guenther, A.B., Yang, X., Wang, L., Xiao, T., Li, J., Feng, J., Xu, Q., Cheng, H.: Land cover change dominates decadal trends of biogenic volatile organic compound (BVOC) emission in China. *Atmos. Chem. Phys., Discuss.*, 1-29, doi: 10.5194/acp-2020-28, 2020.
- 480 Wang, T., Dai, J., Lam, K.S., Nan Poon, C., Brasseur, G.P.: Twenty-five years of lower tropospheric ozone observations in tropical East Asia: The influence of emissions and weather patterns. *Geophys. Res. Lett.*, 46, 11463-11470, doi: 10.1029/2019GL084459, 2019.
- 485 Wang, T., Xue, L., Brimblecombe, P., Lam, Y.F., Li, L., Zhang, L.: Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects. *Sci. Total Environ.*, 575, 1582-1596, doi: 10.1016/j.scitotenv.2016.10.081, 2017.
- Wu, R., Xie, S.: Spatial distribution of ozone formation in China derived from emissions of speciated volatile organic compounds. *Environ. Sci. Technol.*, 51, 2574-2583, doi: 10.1021/acs.est.6b03634, 2017.
- 490 Xie, Y., Dai, H., Zhang, Y., Wu, Y., Hanaoka, T., Masui, T.: Comparison of health and economic impacts of PM_{2.5} and ozone pollution in China. *Environ. Int.*, 130, 104881, doi: 10.1016/j.envint.2019.05.075, 2019.
- Yao, R., Wang, L., Huang, X., Gong, W., Xia, X.: Greening in rural areas increases the surface urban heat island intensity. *Geophys. Res. Lett.*, 46, 2204-2212, doi: 10.1029/2018GL081816, 2019.
- Zeb, N., Khokhar, M.F., Pozzer, A., Khan, S.A.: Exploring the temporal trends and seasonal behaviour of tropospheric trace gases over Pakistan by exploiting satellite observations. *Atmos. Environ.*, 198, 279-290, doi: 10.1016/j.atmosenv.2018.10.053, 2019.
- 495 Zhang, X., Zhao, L., Cheng, M., Chen, D.: Estimating ground-level ozone concentrations in eastern China using satellite-based precursors. *IEEE Trans. Geosci. Remote Sens.*, 1-10, doi: 10.1109/TGRS.2020.2966780, 2020.



- Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., Yan, L., Zhang, Y., Zhao, H., Zheng, Y., He, K., Zhang, Q.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions. *Atmos. Chem. Phys.*, 18, 14095-14111, doi: 10.5194/acp-2018-374, 2018.
- Zhu, L., Jacob, D.J., Mickley, L.J., Marais, E.A., Cohan, D.S., Yoshida, Y., Duncan, B.N., Abad, G.G., Chance, K.V.: Anthropogenic emissions of highly reactive volatile organic compounds in eastern Texas inferred from oversampling of satellite (OMI) measurements of HCHO columns. *Environ. Sci. Technol. Lett.*, 9, 114004, doi: 10.1088/1748-9326/9/11/114004, 2014.
- Zhu, L., Mickley, L.J., Jacob, D.J., Marais, E.A., Sheng, J., Hu, L., Abad, G.G., Chance, K.: Long-term (2005-2014) trends in formaldehyde (HCHO) columns across North America as seen by the OMI satellite instrument: Evidence of changing emissions of volatile organic compounds. *Geophys. Res. Lett.*, 44, 7079-7086, doi: 10.1002/2017GL073859, 2017.
- Zhuang, Y., Li, R., Yang, H., Chen, D., Chen, Z., Gao, B., He, B.: Understanding temporal and spatial distribution of crop residue burning in China from 2003 to 2017 using MODIS data. *Remote Sens.*, 10, 390, doi: 10.3390/rs10030390, 2018.




Figure 1: The May-to-September mean hourly surface ozone network data from 2014 to 2019. Mean hourly surface ozone concentrations were calculated on the $0.25^\circ \times 0.25^\circ$ grid. Purple, blue, green and red outlines indicate the boundaries of North China Plain (NCP), Yangtze River delta (YRD), Pearl River delta (PRD), and Sichuan Basin (SCB), respectively.




Figure 2: The workflow of the polynomial simulation and the causality analysis.





Figure 3: The geographical locations of urban area, urban fringe and rural area.



520

Figure 4: May-to-September averaged HCHO and NO₂ across China during the period of 2005-2012 and 2013-2019.





Figure 5: The linear trends of May-to-September HCHO and NO₂ across China during the period of 2005-2012 and 2013-2019.



525 **Figure 6:** The time series of HCHO columns in the four megacity clusters from 2005 to 2019. Black lines indicate the linear trend of HCHO columns. Red, white and blue areas stand for VOC-limited, transitional and NO_x-limited regimes, respectively.






Figure 7: The time series of NO₂ columns in the four megacity clusters from 2005 to 2019. Black lines indicate the linear trend of NO₂ columns. Red, white and blue areas stand for VOC-limited, transitional and NO_x-limited regimes, respectively.



(a)



(b)

530 Figure 8: (a) Fitting ozone exceedance probability to HCHO/NO₂ through third-order polynomial model. The curve indicated the
 fitting result of third-order polynomial. The vertical line denoted the maximum of curve and the shaded area represented the top
 10% ozone exceedance probability. (b) The cross map skill of HCHO and NO₂ on surface ozone (The skill of using HCHO and
 535 NO₂ for predicting surface ozone concentrations) at different ranges of HCHO/NO₂. The symbols and texts above the bars were
 the results of Wilcoxon test. ***, ** indicated the difference was significant at $p = 0.01, 0.05$ confidence level, respectively. NS
 suggested non-significant differences.






Figure 9: The time series of HCHO/NO₂ in the four megacity clusters from 2005 to 2019. Black lines indicate the linear trend of HCHO/NO₂. Red, green and blue dots stand for VOC-limited, transitional and NO_x-limited regimes, respectively.



540

Figure 10: (a) The spatial distribution of HCHO/NO₂ across China in 2005 and 2019. The boundaries of NCP, YRD, PRD and SCV are denoted with the purple, blue, yellow and red bold lines. Red, green and blue stand for VOC-limited, transitional and NO_x-limited regimes. (b) The annual mean cross map skill (ρ value) of four megacity clusters. The red and blue shadow areas indicate the standard deviations.



545

Figure 11: The temporal variations of HCHO/NO₂ from 2005 to 2019 in NCP, YRD, PRD and SCB. The two red dash lines indicate the threshold values of 1.0 and 1.9, which separate the NO_x-limited, transitional and VOC-limited regime.