

Assessing the potential efficacy of marine cloud brightening for cooling Earth using a simple heuristic model

Robert Wood¹

5 ¹Department of Atmospheric Sciences, University of Washington, Seattle, WA 98195, USA

Correspondence to: Robert Wood (robwood2@uw.edu)

Abstract. A simple heuristic model is described to assess the potential for increasing solar reflection by augmenting the aerosol population below marine low clouds, which nominally leads to increased cloud droplet concentration and albedo. The model estimates the collective impact of many point-source particle sprayers, each of which generates a plume of injected particles that affects clouds over a limited area. A look-up table derived from simulations of an explicit aerosol activation scheme is used to derive cloud droplet concentration as a function of the sub-cloud aerosol size distribution and updraft speed, and a modified version of Twomey’s formulation is used to estimate radiative forcing. Plume overlap is accounted for using a Poisson distribution assuming idealized elongated cuboid plumes that have a length driven by aerosol lifetime and wind speed, a width consistent with satellite observations of ship track broadening, and a depth equal to an assumed boundary layer depth. The model is found to perform favorably against estimates of brightening from large eddy simulation studies that explicitly model cloud responses to aerosol injections over a range of conditions. Although the heuristic model does not account for cloud condensate or coverage adjustments to aerosol, in most realistic ambient remote marine conditions these tend to augment the Twomey effect in the large eddy simulations, with the resulting being a modest underprediction of brightening in the heuristic model.

20 The heuristic model is used to evaluate the potential for global radiative forcing from marine cloud brightening as a function of the quantity, size, and lifetime of salt particles injected per sprayer and the number of sprayers deployed. Radiative forcing is sensitive to both the background aerosol size distribution in the marine boundary layer into which particles are injected, and the assumed updraft speed. Given representative values from the literature, radiative forcing sufficient to offset a doubling of carbon dioxide $\Delta F_{2\times\text{CO}_2}$ is possible but would require spraying over 50% or more of the ocean area. This is likely to require at least 10^4 sprayers to avoid major losses of particles due to near-sprayer coagulation. The optimal dry diameter of injected particles, for a given salt mass injection rate, is 30-60 nm. A major consequence is that the total salt emission rate (50-70 Tg yr⁻¹) required to offset $\Delta F_{2\times\text{CO}_2}$ is a factor of five lower than the emissions rates required to generate significant forcing in previous studies with climate models, which have mostly assumed dry diameters for injected particles in excess of 200 nm. With the lower required emissions, the salt mass loading in the marine boundary layer for $\Delta F_{2\times\text{CO}_2}$ is dominated by natural salt aerosol, with injected particles only contributing ~10%. When using particle sizes optimized for cloud brightening, the aerosol direct radiative forcing is shown to make a minimal contribution to the overall radiative forcing.

1 Introduction

35 Marine low clouds reflect solar radiation and cool Earth as a result (Hartmann and Short, 1980; Ramanathan et al., 1989). The solar radiation reflected by marine low clouds (albedo) increases with the amount of liquid water they contain and as the size of cloud droplets decreases (Stephens 1978). Twomey (1974, 1977) showed that, for a fixed liquid water path (*LWP*), cloud albedo increases with the concentration of cloud droplets (N_d). Thus anthropogenic aerosol pollution increases cloud albedo and cools climate. Four decades of subsequent research has established the “Twomey effect” as the largest contributor to the overall cooling impact of aerosols on climate (Zelinka et al., 2014; Bellouin et al., 2020).

In recent decades, evidence has mounted showing cloud macrophysical adjustments to aerosol increases. Albrecht (1989) suggested that reduced droplet sizes would lead to suppressed collision-coalescence, greater retention of water, and an augmentation of the Twomey effect. Modelling and observations do both show precipitation suppression by aerosol in warm clouds (Ackerman et al., 2004; Sorooshian et al., 2010; Terai et al., 2015), and yet observations of ship tracks (Coakley and Walsh 2003, Toll et al., 2019), pollution plumes (Toll et al., 2019; Trofimov et al., 2020), and large-scale shipping lanes (Diamond et al., 2020) reveal LWP reductions in the mean. Modeling has shown that aerosols can cause both positive and negative LWP adjustments (Ackerman et al., 2004; Wood 2007), with the sign of the change dependent on meteorological and aerosol conditions. Reduced LWP stems from increased cloud top entrainment of dry free-tropospheric air due to smaller cloud droplets and/or turbulent invigoration of the boundary layer caused by suppressed precipitation (Wang et al., 2003; Ackerman et al., 2004; Bretherton et al., 2007; Wood 2007). A recent paper by Glassmeier et al. (2021) illustrates that the sign of LWP adjustments depends not only on the meteorological conditions but also on the number of aerosol particles, causing positive adjustments when the aerosol number is small and precipitation suppression increases condensate retention. Negative adjustments are found when the aerosol number is large, due to the aforementioned entrainment drying. Studies using shipping and land-based pollution sources suggest that mean LWP decreases may offset the Twomey response to a degree that ranges from 3% (Trofimov et al., 2015) to perhaps 20% (Toll et al., 2019; Diamond et al., 2020). LWP adjustments in low clouds are poorly handled in large scale models (Malavelle et al., 2017), which almost universally show LWP increases in simulations of anthropogenic aerosol impacts (Lohmann and Feichter 2005; Isaksen et al., 2009; Bellouin et al., 2020). Global models also tend to show cloud cover increases in response to aerosol, but these appear to be small compared with the Twomey responses and LWP adjustments (Zelinka et al., 2014). Cloud cover adjustments are difficult to constrain using observations (e.g., Gryspeerd et al., 2016; Possner et al., 2018).

The high sensitivity of cloud albedo to aerosol increases led Latham (1990) to speculate that cloud albedo could potentially be increased deliberately by augmenting the number of aerosol particles ingested into them. This is commonly known as “marine cloud brightening” (MCB), and it has been an increasing focus of research as a potential climate intervention strategy for over a decade (e.g., Latham et al., 2008, 2012; Jones et al., 2009; Rasch et al., 2009; Alterskjær et al., 2012; National Research Council 2015; Ahlm et al., 2017; Stjern et al., 2018). MCB involves spraying small solution drops containing sea salt into the marine boundary layer (MBL), increasing the concentration of cloud condensation nuclei. This ideally results in a higher concentration of cloud droplets and more reflective clouds. Any large-scale deployment of MCB would involve many point source injections from seagoing vessels distributed over the ocean (Salter et al., 2008). Essentially, such a deployment can be thought of as a deliberate augmentation of the natural experiment currently being conducted by the fleet of commercial ships (~60,000) that are currently emitting aerosol and precursor gases over the world’s oceans (Eyring et al., 2010). Thus, we can draw on the study of ship tracks and shipping lanes to provide insights regarding the potential efficacy of MCB.

75 A ship track is a brightened curvilinear feature in a marine cloud deck caused by the emission of particles and their precursors from an individual ship (Conover 1966). These tracks provide dramatic evidence that cloud reflectivity can increase when particles are released into the MBL. However, ship tracks are insufficient to estimate the large-scale radiative forcing possible. The global increase in reflected shortwave radiation from discernible ship tracks has been estimated from satellite observations to be $\sim 4\text{-}6 \times 10^{-4} \text{ W m}^{-2}$ (Schreier et al. 2007), which is 2-3 orders of magnitude smaller than climate model estimates of the
80 total effect of shipping emissions of aerosol and aerosol precursors, which range from 0.06-0.6 W m^{-2} (Capaldo et al., 1999; Lauer et al., 2007; Eyring et al., 2010; Peters et al., 2012; Partanen et al., 2013). The most easily discernible ship tracks form in very shallow MBLs (Durkee et al., 2000). These “Type I” ship tracks tend to occur in MBLs with particularly low concentrations of background aerosol (Hindman et al., 1994; Ackerman et al., 1995), in which turbulent mixing is weak because drizzle depletes liquid water and precludes strong cloud top radiative cooling. A more common type of ship track
85 (“Type II”) tends to be more readily discernible using near infrared rather than visible satellite imagery (Coakley et al., 1987), highlighting the smaller droplets in the track. The MBLs in which Type II ship tracks form tend to be somewhat deeper, more well-mixed and strongly driven by cloud top cooling. Ship track albedo perturbations in these cases tend to be weaker than in Type I tracks. Large eddy simulations of deep stratocumulus-topped MBLs indicate that albedo can be increased substantially by injected aerosol emissions even when a clear track is not discernible (Possner et al., 2018). In Durkee et al. (2000) no ship
90 tracks were detected in MBLs deeper than 800 m, but Possner et al. (2020) show that over 80% of all stratocumulus-topped MBLs over the oceans are deeper than 800 m, where surface emissions can increase cloud albedo but tracks may not be easy to detect.

An alternative to observational studies of individual ship tracks is to quantify the mean radiative forcing over a heavily
95 trafficked area to assess the aggregate effect of shipping. Diamond et al. (2020) was able to discern a corridor of enhanced mean N_d in clouds above a shipping lane that traverses the SE Atlantic subtropical stratocumulus deck. In this corridor, an increase in reflected diurnal-seasonal mean shortwave radiation of 2 W m^{-2} was observed associated with an increase in N_d of $\sim 5 \text{ cm}^{-3}$, which is consistent with expectations from the Twomey effect. Cloud adjustments were found to be relatively small, with reduced cloud LWP in the shipping lane offsetting $\sim 20\%$ of the Twomey effect, and a small cloud fraction increase
100 augmenting the Twomey effect by $\sim 10\%$. Although the radiative forcing would need to be somewhat stronger for MCB to offset a significant fraction of the radiative forcing from increased greenhouse gases, the lack of major cancelling cloud adjustments points to the potential for regional albedo enhancement using MCB. In this case the aerosols (from ship emissions) were inadvertently brightening clouds; aerosols of a size and concentration that targets intentional cloud brightening would very likely have a larger impact on cloud albedo and radiative forcing.

105

Climate models demonstrate the potential for producing a globally-significant radiative forcing from MCB. These studies fall into two broad categories: (i) studies in which N_d (or droplet effective radius r_e) in some fraction of the marine low cloud population is altered to some specified value to increase cloud albedo; (ii) studies that achieve cloud albedo changes by

increasing the surface aerosol source and treating the aerosol activation process, leading to changes in N_d . The latter studies
110 involve a more complete treatment of the chain of causality that links aerosol emissions to brightening, while the former studies
can be carried out without explicit representation of aerosol-cloud interaction processes.

Seeded regions in studies with specified N_d or r_c perturbations have increased N_d to different levels: 375 cm^{-3} in Jones et al.,
(2009), 1000 cm^{-3} in Rasch et al., (2009) and Baughman et al., (2012) and both 375 and 1000 cm^{-3} in Latham et al. (2008). In
115 Bala et al. (2011), the cloud effective radius is instead decreased from 14 to $11.5 \mu\text{m}$ for all marine liquid clouds, which is
approximately equivalent to increasing N_d by 80% . Stjern et al. (2018) increase N_d by 50% in all marine low clouds. Because
cloud albedo increases scale with the ratio of perturbed (seeded) to unperturbed N_d (Sect. 2.1), these changes represent a wide
diversity in how much a seeded cloud is brightened in each study. Very different fractions of the available ocean are seeded in
different studies, ranging from 1.0 , 1.6 , 2.1 and 4.7% of the ocean area in Jones et al., (2009), 9% in Baughman et al. (2012),
120 20 , 30 , 40 and 70% in Rasch et al., (2009) and the entire ocean in Bala et al. (2011). Jones et al. (2009) achieved a forcing of
 -1 W m^{-2} despite only perturbing 4.7% of the ocean surface, but perturbed regions had extensive low clouds. Rasch et al. (2009)
went further and identified the albedo susceptibility (change in albedo upon increasing N_d to 1000 cm^{-3}) for each grid box on
a seasonal basis. The most susceptible 20 , 30 , 40 and 70% of the boxes were then used as seeding regions. The wide range in
different areas seeded and in the strength of the N_d perturbation where the seeding occurs makes it difficult to intercompare
125 the effectiveness of the seeding across studies.

Climate model studies in which an aerosol surface source is added as a proxy for deliberate spraying have also been shown to
produce globally significant radiative forcing (Ahlm et al., 2017), with values in some studies more than offsetting that from
doubling CO_2 (e.g., Alterskjaer et al., 2012). Such studies introduce several additional degrees of freedom into the experimental
130 design. A comprehensive representation of the aerosol lifecycle is needed, as well as an aerosol activation parameterization to
predict N_d as a function of the aerosol size distribution in the MBL. As studies with aerosol activation schemes and/or parcel
models have shown, N_d is sensitive primarily to the concentration of aerosol in the accumulation mode (particles with dry
diameters around 50 - 200 nm), but is also sensitive to updraft speed and to small concentrations of coarse mode aerosol, which
reduce the peak supersaturation in an updraft and lower the fraction of smaller aerosols activated (Ghan et al., 1998; McFiggans
135 et al., 2006).

An additional aerosol surface source from an MCB sprayer can, in principle, be tailored to consist of particles of a specific
diameter. Connolly et al. (2014) explored the optimal particle size given energy constraints on particle production, which
primarily scales with the mass of salt injected, and found that sodium chloride particles with a modal diameter in the range 30 -
140 90 nm are optimal. Climate model studies to date have typically introduced injected particles with modal diameters that are
several times as large as this (Alterskjaer et al., 2012; Ahlm et al., 2017), which implies that these models likely require much
larger salt mass emissions than may be required if smaller particles are injected. Only Partanen et al. (2012) has tested the

sensitivity to injecting particles with modal dry diameter of 100 nm and found the same brightening as in a base case with 200 nm diameter particles but with ~ 5 times less mass injected. Consideration of total salt mass injected is important not only from
 145 the perspective of the energy required to produce particles, but also because major increases in sodium chloride aerosol mass could potentially alter natural chemical cycles in the MBL (Horowitz et al., 2020).

This study describes a simple heuristic model that predicts the global radiative forcing from MCB using physical principles to determine the collective impact of plumes from many point source sprayers distributed over the oceans on N_d and cloud albedo.
 150 The model is designed to facilitate easy experimentation on the factors controlling MCB, including details of the unperturbed aerosol size distribution, the number concentration, size and residence time of injected particles, the number of sprayers, and the fraction of the ocean over which sprayers are deployed. Section 2 describes the heuristic model in detail, and Sect. 3 tests the model using comparisons with high resolution, small-domain large eddy simulation models into which point source injections are introduced. Section 4 uses the heuristic model to examine factors controlling global radiative forcing from MCB,
 155 and critically examines some assumptions made in previous climate model studies. Finally, Sect. 5 discusses implications of the results and suggests pathways for future study, and Sect. 6 provides conclusions.

2 Heuristic model description

2.1 Radiative forcing from aerosol-cloud interactions

Central to the model is Twomey's formulation for the susceptibility of cloud albedo α_c to an increase in N_d assuming no cloud
 160 adjustments (Twomey 1977), viz.

$$\frac{d\alpha_c}{dN_d} = \frac{\alpha_c(1-\alpha_c)}{3N_d} \quad (1)$$

Integrating Eq. (1) gives an expression for the increase in cloud albedo $\Delta\alpha_c$ caused by an increase in N_d :

$$\Delta\alpha_c = \frac{\alpha_c(1-\alpha_c)(r_N^{1/3}-1)}{1+\alpha_c(r_N^{1/3}-1)} \quad (2)$$

where $r_N = N'_d/N_d$ is the ratio of the droplet concentration in seeded vs unperturbed clouds. It is worth noting that Eq. (2) is
 165 rather insensitive to α_c , such that $\Delta\alpha_c$ varies by only $\sim 10\%$ as α_c changes from 0.3-0.7. Thus, the key sensitivity in Eq. (2) is to the value of r_N .

To estimate the top-of-atmosphere (TOA) albedo for the same cloud requires a conversion to account for the absorption and scattering of solar radiation by the atmosphere above cloud. We follow the approach by Diamond et al. (2020, Eq. 17) and
 170 multiply the cloud albedo change by an atmospheric correction factor ϕ_{atm} :

$$\phi_{\text{atm}} = \frac{\Delta\alpha_{c,\text{TOA}}}{\Delta\alpha_c} = \frac{T_{\text{FT}}^2}{(1-\alpha_{\text{FT}}\alpha_c)^2} \quad (3)$$

Where T_{FT} and α_{FT} are the transmissivity and albedo of the free troposphere only. The more variable of these two parameters is T_{FT} , which depends upon free-tropospheric water vapor. Here, we assume a value of $T_{\text{FT}} = 0.8$, consistent with values over
 175 dry regions of the Tropics and midlatitudes from the CERES-SYN product (Doelling et al., 2013). Free-tropospheric albedo is less variable and we here assume a value of $\alpha_{\text{FT}} = 0.06$ (also consistent with CERES-SYN). For typical cloud albedos α_c in the range 0.25 to 0.75, ϕ_{atm} ranges from 0.66 to 0.70, so for simplicity we herein assume $\phi_{\text{atm}} = 0.70$. We estimate TOA indirect radiative forcing as $-F_{\odot}\phi_{\text{atm}}\Delta\alpha_c$, where F_{\odot} is the mean incoming solar irradiance averaged over day and night. Here, we assume a value of F_{\odot} equal to the global mean solar irradiance $F_{\odot} = 342 \text{ W m}^{-2}$. Geographical variation in insolation is not
 180 considered.

2.2 Regions where sprayers operate

Marine cloud brightening, by definition, would only be deployed over the fraction of Earth covered by ocean. We further restrict this area to minimize the likelihood that plumes will intersect land areas. This is done by summing up the ocean area of those $10 \times 10^\circ$ latitude/longitude boxes that contain less than 10% land area. The choice of boxes 10° on a side is made
 185 because plumes are of order 1000 km in length (see Sect. 2.4). This limits the eligible fraction of Earth's surface for spraying, f_{ocean} to 0.54. We then assume that sprayers are confined to operate within some specified fraction f_{spray} ($0 < f_{\text{spray}} \leq 1$) of this eligible area. If f_{spray} is chosen to be less than unity, it is assumed that sprayed regions will be those with the highest climatological unobstructed low cloud cover. To determine the mean low cloud cover for the sprayed subregions f_{low} , climatological monthly mean low cloud fractions are determined using MODIS Terra and Aqua Level 3 liquid cloud fractions
 190 (years 2006-2010) for $10 \times 10^\circ$ boxes. As f_{spray} decreases, the fraction of the ocean sprayed has a greater coverage of low clouds. If f_{spray} is chosen to be very small, spraying would occur only in regions with the highest climatological monthly mean cloud cover (~68%). The MODIS data are well-fitted with the empirically-determined expression:

$$f_{\text{low}} = 0.32 + 0.36 \exp(-3.2f_{\text{spray}}^{0.75}) \quad (4)$$

195 2.3 Expression for global radiative forcing associated with MCB

Cloud condensate and coverage adjustments to injected aerosol are assumed to be zero, so MCB indirect radiative forcing arises only from the Twomey effect. In sprayed areas without low clouds, injected particles can exert a direct radiative forcing. The direct radiative forcing from injected aerosol in cloud-free regions between clouds is quantitatively estimated (see Sect. 2.7), but increasing direct radiative forcing is not a goal of the injection design. The global mean shortwave radiative forcing
 200 ΔF from MCB aerosol-cloud interactions is written as

$$\Delta F = -F_{\odot}f_{\text{ocean}}f_{\text{spray}}f_{\text{low}}\phi_{\text{atm}}\Delta\alpha_c \quad (5)$$

To give a “back of the envelope” assessment of the potential for MCB, we take $f_{\text{ocean}} = 0.54$, assume $f_{\text{spray}} = 1$ and use Eq. (4) to set $f_{\text{low}} = 0.33$. If cloud albedo is increased by $\Delta\alpha_c = 0.01$, then $\Delta F = -0.41 \text{ W m}^{-2}$. Alternatively, it would take a cloud albedo increase of $\Delta\alpha_c = 0.09$ to produce a radiative forcing of -3.7 W m^{-2} , which would balance the longwave radiative forcing $\Delta F_{2\times\text{CO}_2}$ from doubling CO₂. Figure 1 shows ΔF as a function of r_N for different values of f_{spray} and α_c . Using Eq. (2), if we assume $\alpha_c = 0.56$ (Bender et al. 2011 finds TOA cloud albedos of 0.35 to 0.42 for overcast stratocumulus in the major subtropical Sc decks, which must be corrected to cloud albedos with Eq. 3), then the ratio of seeded to unseeded cloud droplet concentration ($r_N = N'_d/N_d$) would need to be 3.0 to produce a forcing with a magnitude equal to $\Delta F_{2\times\text{CO}_2}$. Assuming the *entire* ocean area could be seeded ($f_{\text{ocean}} = 0.7$, $f_{\text{spray}} = 1$), we find a value of $r_N = 2.4$, which is in the range of N_d increases over the ocean (2.10-2.85) that were needed to counter CO₂ doubling in an analysis of three variants on a climate model (Slingo 1990). If only half of the eligible ocean area is seeded (i.e., $f_{\text{spray}} = 0.5$), then r_N would need to be at least 7 to counter CO₂ doubling (Fig. 1). Stjern et al. (2018) analyzed an ensemble of different climate models in which N_d for all marine low clouds is increased by 50% ($r_N=1.5$) as a proxy for MCB, and found an ensemble mean $\Delta F = -1.9 \text{ W m}^{-2}$. Based on Fig. 1, and scaling the forcing to include the entire ocean, Eq. (5) produces a very similar forcing $\Delta F = -1.8 \text{ W m}^{-2}$. This is also consistent with the models in Stjern et al. (2018) having small cloud adjustments overall, so that the overwhelming bulk of the forcing is from the Twomey effect.

2.4 Aerosol delivery and plume/track configuration

Any practical MCB deployment would be unable to produce uniform increases of N_d because seeding is necessarily discrete in nature, rather than distributed evenly. It is impractical to deploy sprayers at every point over the ocean; in practice, any deployment would likely consist of an array of floating particle injection systems distributed throughout regions where low clouds occur. To extend the heuristic model to account for this, assumptions are made about the spatiotemporal extent of the region affected by a single sprayer. Sprayers are assumed to be stationary so that air masses pass over them at the rate of the near-surface wind speed U_0 , which is taken as 7 m s^{-1} , the mean value over oceans (Archer and Jacobson 2005). Each sprayer injects sodium chloride particles continuously with a salt mass rate \dot{M}_s . Injected particles have a lognormal size distribution with geometric mean dry diameter (GMD) D_s and geometric standard deviation (GMS) S . The total number of particles sprayed per second from each sprayer \dot{N}_s is then

$$\dot{N}_s = \frac{6\dot{M}_s}{\pi\rho_s D_s^3 e^{9(\ln S)^2/2}} \quad (6)$$

where ρ_s is the density of solid sodium chloride (2160 kg m^{-3}). The volume into which particles are emitted increases with time as the plume expands to fill the depth of the MBL and widens horizontally. The timescale for vertical dispersion through the depth of the MBL is 10-20 min (Chosson et al., 2008), as evidenced by the fact that, in ship tracks, brightened clouds become evident typically 10-20 km downwind of the responsible ship. As satellite data readily show, ship tracks from commercial shipping are narrower close to the emitting ship and broaden downstream (Durkee et al., 2000). After rapid vertical

235 dispersion through the MBL, dilution primarily occurs through lateral diffusion. Entrainment of lower concentration free-tropospheric air also dilutes the plume, but at a slower rate. The lateral track broadening rate is highly variable but is parameterized using the Heffter (1965) broadening rate $K=1.85 \text{ km hr}^{-1}$ (see Fig. 7 in Durkee et al. 2000). This rate is broadly consistent with large eddy simulations of horizontal tracer spread in the cloudy MBL (Wang et al., 2011).

240 It has been proposed that a spray system to inject salt particles could derive the salt from sea water droplets (Salter et al., 2008; Cooper et al., 2014). Sea water is substantially more dilute than the equilibrium size of solution droplets at the surface (see, e.g., Hoffman and Feingold 2021), and so there is some concern that cooling from the evaporation of water from equilibrating droplets may hinder or prevent the vertical mixing of injected particles. Reducing any negative buoyancy is an engineering challenge that may be addressed by increasing the turbulent mixing of the particle-laden plume with surrounding air, and/or
245 adding some thermal energy to the particle plume. This issue is beyond the scope of this study, and we herein assume the injected particles mix readily throughout the depth of the MBL. We also note that the additional water vapor introduced into the lower MBL from the evaporating sea water is negligible compared with the natural surface evaporative water flux and thus will have no impact on the MBL moisture budget.

250 Injected particles in the model have a characteristic residence e-folding timescale τ_{res} . This residence time incorporates several processes influencing particle lifetime, including removal by coalescence scavenging, scavenging by clouds and aerosol particles and dry deposition. The value of τ_{res} varies with meteorological conditions, cloud, and precipitation properties, and is also expected to be somewhat size dependent. In regions of marine stratocumulus regions values of τ_{res} of 2-3 days are consistent with estimates of precipitation scavenging (Wood et al., 2012), and $\tau_{\text{res}} = 2$ days is used as standard.

255

After a time t , the particles injected at time $t = 0$ have moved a distance $x = U_0 t$. Considering both dilution and removal processes, given a plume width $W(t) = Kt$ and assuming dispersion through the entire MBL depth h , the injected particle concentration $N_s(t)$ at time t is:

$$N_s(t) = \frac{N_s}{U_0 h K t} e^{-t/\tau_{\text{res}}} \quad (7)$$

260

The wind speed and residence time define a length scale ($L_t = U_0 \tau_{\text{res}}$) that effectively determines the streamwise length scale over which the particle concentration is affected by spraying. The area over the Earth's surface perturbed by each sprayer A is then determined by multiplying this length scale by a characteristic track width W_t , i.e. $A_t = L_t W_t = U_0 \tau_{\text{res}} W_t$. A linearly widening plume/track will expand to a width $K\tau_{\text{res}}$ over the lifetime of the particles.

265

To estimate radiative forcing, the injected particle concentration $N_s(t)$ is added to an assumed background aerosol over the entire track area and over the depth h of the MBL. Aerosol activation to form cloud droplets is carried out for the background

aerosol and for the perturbed (background+injected) aerosol using an assumed updraft speed. Section 2.6 provides details of the activation scheme, and the aerosol physical and chemical properties. The ratio of the perturbed to background cloud N_d from the activation scheme are used in the calculation of radiative forcing (Eq. 2 and 5).

Figure 2 shows results from the model for a laterally-spreading track, along with injected particle concentration and additional reflected shortwave from cloud brightening as a function of time/distance downstream of a point-source sprayer. The heuristic model assumes an elongated cuboid plume (fixed width, height, and length) with the plume length $L_t = U_0 \tau_{\text{res}}$ and plume width taken to be the width of the linearly expanding plume at time $\tau_{\text{res}}/2$, i.e., $W_t = K \tau_{\text{res}}/2$. The (time-independent) number concentration N_{s1} of injected aerosol particles in the cuboid plume is

$$N_{s1} = \frac{\dot{N}_s}{hU_0W_t} = \frac{2\dot{N}_s}{hU_0K\tau_{\text{res}}} \quad (8)$$

It is relatively straightforward to show that the overall injected particle concentration integrated over time is the same for the laterally-spreading track (Eq. 7) and the cuboid track (Eq. 8). Although the reflected solar energy from the two tracks is not identical (Fig. 2c), the values are found to be close. A heuristic model track reflects slightly less than a spreading track for a given spray rate, with the difference growing as the magnitude of the N_d perturbation increases.

Experimentation with different spray, background aerosol and cloud configurations shows that the reflected sunlight for the cuboid track is within 5% of that for the spreading track for number spray rates $\dot{N}_s < 10^{16} \text{ s}^{-1}$, with the cuboid model track being slightly less reflective. As \dot{N}_s increases, the ratio of the additional energy reflected by the spreading track to that from the cuboid track increases steadily, reaching 1.2 for $\dot{N}_s = 5 \times 10^{16} \text{ s}^{-1}$, and 1.5 for $\dot{N}_s = 10^{17} \text{ s}^{-1}$ with the exact value dependent upon the background aerosol. As the magnitude of the aerosol number perturbation increases, an increasing fraction of the energy reflected occurs at times $t > \tau_{\text{res}}$ in the spreading plume. The albedo response in the cuboid track is relatively saturated due to the high aerosol/droplet concentrations (see Fig. 1), so the dilute but widespread aerosol in the spreading plume later on is more efficient at brightening. As we show in the discussion, coagulation losses during the high concentrations near to the sprayer are likely to be large for particle spray rates much greater than $\sim 10^{16} \text{ s}^{-1}$. The cuboid tracks are thus a sufficiently accurate representation of reality to use them in the heuristic model and simplifies the treatment of overlapping tracks.

2.5 Overlapping tracks

Given the plume dimensions for the heuristic model tracks, we estimate that the number of (non-overlapping) tracks required to cover the 54% of the ocean eligible for spraying ($\sim 1.98 \times 10^{14} \text{ m}^2$), assuming $W_t = 44 \text{ km}$, $\tau_{\text{res}} = 2 \text{ days}$, and $U_0 = 7 \text{ m s}^{-1}$ (i.e., $L_t \approx 1200 \text{ km}$; $A_t = 5.28 \times 10^{10} \text{ m}^2$) is 5300. If this number of sprayers was to be deployed either randomly or uniformly then overlapping tracks would be unavoidable because air mass trajectories are not constant in time. Monte Carlo simulations were

conducted, placing N_t randomly-oriented or aligned rectangular tracks at random over a large domain of area A . The
 300 probability $p(n)$ of n tracks overlapping in the domain is well-predicted by a Poisson distribution:

$$p_p(n) \approx \frac{\zeta^n e^{-\zeta}}{n!} \quad (9)$$

where $\zeta = N_t A_t / A$ is the mean track density, i.e., the mean number of superimposed tracks (Fig. 3). Although not shown, $p(n)$
 305 is insensitive to both the track aspect ratio (L_t/W_t) and whether the tracks are aligned with their long sides in one direction or
 are randomly oriented.

The use of the Poisson distribution makes it straightforward to account for track overlap in the heuristic model; the injected
 particle concentration N_s at any given location is an integer multiple n ($n \geq 0$) of the single-track value N_{s1} from Eq. (8):
 310

$$N_s = nN_{s1} = \frac{2\dot{N}_s n}{hU_0 K \tau_{res}} \quad (10)$$

where the probability of n is given by Eq. (9). For low mean track densities ζ , the most likely value of n is zero (Fig. 3), and
 the ratio of the standard deviation to the mean value of n is high. A Poisson distribution has equal mean and variance, so the
 315 relative spatial heterogeneity of N_s , i.e., the ratio of the standard deviation to the mean track density, decreases as $\zeta^{-1/2}$.
 Because of the concave relationship between $\Delta\alpha_c$ and N_s (see e.g., Carslaw et al., 2013), a more homogeneous distribution of
 N_s over the seeded area will yield a radiative forcing with a larger magnitude for the same mean value of N_s .

2.6 Aerosol activation and physical and chemical properties

Aerosol activation to form cloud droplets is treated using a five-dimensional look-up table derived from over 6000 numerical
 320 Lagrangian parcel model simulations (see Appendix). In comparing with those using the Abdul-Razzak and Ghan (2000)
 quasi-analytical activation scheme (henceforth ARG), we find significant differences that indicate a major underprediction of
 N_d with the ARG scheme when injected dry particle diameters are smaller than ~ 200 nm (see Appendix and also Sect. 4.2),
 and so use the look-up table to treat activation in the heuristic model.

325 The aerosol size distributions used are the same for each activation approach. The background (unperturbed) aerosol particles
 are assumed to comprise lognormal accumulation and coarse modes. Accumulation mode size values ($D_{0,acc}=175$ nm;
 $S_{0,acc}=1.5$) are taken from the synthesis of marine accumulation mode measurements by Heintzenberg (2000). Measured marine
 accumulation mode number concentrations $N_{0,acc}$ vary considerably over the ocean, and impacts of this on brightening are
 explored in Sect. 4.2. Although there is significant variability in the composition of marine CCN, studies tend to find that the

330 accumulation mode aerosol in the unpolluted MBL consists of a mixture of sulfate, sea salt, and organic species. Different assessments of the hygroscopicity parameter (κ , from Petters and Kreidenweis 2007) of CCN in the MBL provide a significant diversity of values, from values as low as 0.45 (Wex et al., 2010) to ~ 0.7 (Andreae and Rosenfeld 2008). Here, we use the mean marine value of 0.7 from the model study of Pringle et al. (2010) for the unperturbed accumulation mode. The background coarse mode is lognormal with GMD $D_{0,coarse}=615$ nm and GMS $S_{0,coarse}=1.8$ taken from summertime
 335 measurements at Graciosa Island in the Azores (Zheng et al. 2018). The presence of the coarse mode suppresses the peak supersaturation in the updraft, increasing the minimum size of the particles that are activated, reducing the activated fraction (Ghan et al. 1997). This is explored further in Sect. 4.2. Injected aerosols are sodium chloride ($\kappa = 1.2$, Petters and Kreidenweis 2007), distributed lognormally with GMD D_s and GMS S , where D_s is allowed to vary, and $S=1.6$. Table 1 provides a summary of the assumed aerosol properties used. A recent study suggests that the other inorganic species in sea salt
 340 render it slightly less hygroscopic than pure sodium chloride ($\kappa=1.1$, Zieger et al., 2017). Testing showed that the results of this study are largely insensitive to small variations in κ .

For most of the analysis presented in this study, a fixed updraft speed of $w=0.4$ m s⁻¹ is assumed in the activation scheme. This is broadly representative of updrafts in the stratocumulus-topped MBL (Nicholls and Leighton 1986; Wood 2005;
 345 Bretherton et al., 2010; Zheng et al., 2016). Sensitivity to updraft speed is explored in Sect. 4.3. For simplicity, the temperature and pressure are set to be 280 K and 925 hPa respectively, but the results are not highly sensitive to these values.

2.7 Aerosol direct radiative forcing

The heuristic model is also used to produce rough estimates of the aerosol direct radiative forcing from the injected aerosol. We assume direct forcing only in clear sky regions. In an analogous formulation to Eqn. 5, we estimate the global mean direct
 350 radiative forcing as:

$$\Delta F_{\text{direct}} = -F_{\odot} f_{\text{ocean}} f_{\text{spray}} f_{\text{clear}} E \cdot AOD_{\text{spray}} \quad (11)$$

where f_{clear} is the clear sky fraction in the regions where sprayers operate, AOD_{spray} is the aerosol optical thickness (550 nm)
 355 of injected particles, and E is the clear-sky radiative forcing efficiency. We use $E = -29$ W m⁻² AOD^{-1} , which is the average over oceans for several models in the AeroCom study of Schulz et al. (2006). In this study, direct effects are only estimated for the case where $f_{\text{spray}}=1$, i.e., sprayers operate in all eligible regions of the oceans, and so $f_{\text{clear}} = 0.32$ is taken as the complement of the total cloud cover during the daytime over the global oceans from Hahn and Warren (2007). To estimate AOD_{spray} , the injected aerosol lognormal size distribution (accounting for overlapping tracks as discussed in Sect. 2.5) is used
 360 to estimate extinction σ_{spray} at 550 nm using the Mie code of Bohren and Huffman (1998). A mean relative humidity in clear sky MBLs of 80% is used to set a hygroscopic diameter growth factor for sodium chloride of 2.0 from Tang (1996). The

assumed MBL depth $h = 1$ km (Table 1) is used to determine $AOD_{\text{spray}} = h\sigma_{\text{spray}}$. Direct forcing estimates are presented in Sect. 4.4.

3. Comparison of heuristic model with large eddy simulations

365 Several existing studies in the literature have used large eddy simulations (LES) to explore impacts of salt aerosol injections on marine low cloud microphysical and macrophysical properties and albedo. In contrast to climate models, LES explicitly resolves the turbulent dynamics responsible for aerosol distribution through the MBL including ingestion into clouds, in addition to determining cloud macrophysical responses to aerosol resulting from changes in precipitation and mixing with the free troposphere. Although it is not currently possible to run LES with domain sizes large enough to examine regional and
370 global MCB, their faithful representation of injections into domains on scales of a few tens to a few hundred km can provide important insights into the potential efficacy of MCB.

The heuristic model framework is adapted to account for the limited LES domain size to test its predictions. This also allows a quantitative intercomparison of the LES results, which is needed because there is a considerable diversity in the domain
375 sizes, spray rates and particle sizes, as well as in the unperturbed cloud states, boundary layer depths, simulation durations, and in the way in which injections are introduced into the domains, across the different LES studies to date (see Table 2). Each LES experiment consisted of an unperturbed (control) case with no particle injections, and a case with particle injections. A total of 18 different injection simulation experiments are extracted from five studies.

380 Radiative forcing driven by particle injections is estimated for the LES case studies using albedo changes given in the various papers. Unless otherwise stated, heuristic model parameters are those in Table 1. Diurnal mean insolation is assumed. Where appropriate, cloud albedo changes are corrected to the TOA using a fixed value of ϕ_{atm} (Eq. 3, Table 1) consistent with that used in the heuristic model. The heuristic model uses a fixed value for unperturbed cloud albedo (Table 1), but the MBL depth is set to the value for each of the LES cases (Table 2). In several of the cases, the sprayer passes through the model domain
385 multiple times, and in other cases the track does not extend over the entire domain. For the heuristic model predictions we use the Poisson distribution approach (Sect. 2.5) as follows. The duration of the LES experiment in each case is used to determine the track width using the track spreading rate K used in the heuristic model (Table 1), and the ship speed through the domain, is used in place of the wind speed to determine the track length by multiplying by the duration of the simulation. The track area is then computed as the product of the width and length, and this is divided by the LES domain size to obtain a mean track
390 density, which is used to obtain a Poisson distribution of overlapping tracks (Eq. 9). This distribution is used in the heuristic model. In some of the LES experiments, track density is less than unity, but in cases with relatively long durations and/or small domains, it considerably exceeds unity. Wang et al., (2011, henceforth W11) also included a simulation where the same rate

is injected uniformly over the model domain, as a comparison experiment against a point-source sprayer. This is represented in the heuristic model by assuming many (weaker) sprayers operating in the domain.

395

Unperturbed (control case) aerosol size distributions for the heuristic model comprise an accumulation mode with distribution parameters from Table 1 (which are close to those assumed in the LES studies) and concentrations are adjusted to produce unperturbed N_d values reported in the various studies with a fixed 0.4 m s^{-1} updraft (i.e., the standard value used in the heuristic model). Jenkins et al. (2013, henceforth J13), used a bin aerosol scheme rather than a modal scheme. Including an additional
400 coarse mode with modal diameter 500 nm, GSD of 1.8 and concentration of 10 cm^{-3} , which we found represents a fairly good match to the size distributions shown in J13, made less than a 10% difference in the radiative forcing predicted by the heuristic model. No coarse mode is included in the heuristic model for the other cases because none was included in the LES simulations.

Figure 4 presents results from the comparison of the LES and the heuristic model. Overall, the radiative forcing in the LES
405 correlates quite well ($r = 0.62$) with predictions from the heuristic model (Fig. 4a), but the heuristic model underestimates the magnitude of the LES forcing by $\sim 30\%$ in the median. This underestimation is greatest when the unperturbed value of N_d is low (Fig. 4b). There is little bias for cases with $N_d \sim 50 \text{ cm}^{-3}$, but there is underprediction of ~ 2 for $N_d \sim 10 \text{ cm}^{-3}$, and an overprediction of brightening for high N_d cases. The sensitivity of the heuristic model brightening bias to unperturbed N_d (Fig. 4b) is not driven by a model failure to represent the Twomey effect, as the heuristic model ability to predict domain-
410 mean perturbed N_d is excellent ($r = 0.91$, Fig 4c), with only a 25% overestimate in the median. This would lead to a small ($\sim 10\%$) overprediction of Twomey forcing magnitude. Instead, the heuristic model underprediction at low N_d occurs because particle injection into very clean MBLs often leads to increases in liquid water path (LWP), cloud cover, or both. In these cases, the Twomey effect is augmented by cloud adjustments that result in stronger brightening, and this is not represented in the heuristic model. The reasons for overprediction of brightening for high N_d cases is unknown and warrants further attention
415 using more LES studies.

Forcing normalized by the total number of particles injected helps account for the different quantities of particles injected in different studies and provides a useful metric of brightening obtained per particle injected. The LES results show a remarkably strong dependence of this on the unperturbed N_d (Fig. 4d), with a factor of 20 less brightening as unperturbed N_d increases
420 from 10 to 100 cm^{-3} . Although the heuristic model underpredicts (overpredicts) brightening in the clean (polluted) cases, there is still a strong decrease (\sim factor 10) in the brightening as unperturbed N_d increases (Fig. 4e), as anticipated from the Twomey formulation (Eq. 2). In Figs. 4d and 4e, J13 stands out as an anomaly, with a much weaker per-particle brightening compared with the other models. This appears to be because the injection rates used were greater than is needed. An experiment with the precipitating case with an injection rate reduced by a factor of 5 (cyan triangles in Figs. 4d and 4e) leads to less than a 15%
425 reduction in brightening, implying asymptotic brightening as injection rates are increased, and little benefit from the high spray rates used in most of the cases in J13.

It is instructive to compare the brightening obtained per mass of salt injected, and Fig. 4f highlights just how wide a spread there is in this quantity. The most “efficient” brightening is obtained in the Chun et al. (2021) cases with the smallest injected particles (Fig. 4f). For reasons discussed in the introduction, if a forcing can be achieved by injecting less salt mass, then this is desirable, so understanding the optimal size and concentration of injected particles to achieve a required forcing should be a focus for LES studies. These issues are explored further for the heuristic model in Sect. 4.

To synthesize the findings reported here, it should be noted that all the LES studies surveyed show some level of brightening when aerosol injections are introduced. The brightening achieved in the LES experiments, which is here expressed as an equivalent diurnal mean, ranged from ~ 1 - 100 W m^{-2} , with mean of 17 W m^{-2} and a median of 7 W m^{-2} . The median unperturbed cloud N_d (29 cm^{-3}) across all the cases here, is somewhat lower than satellite estimates of average values for low clouds over the global oceans (40 - 90 cm^{-3} , Bennartz 2007). We also used the approach of Bennartz (2007) to derive a pdf of estimated N_d for all marine low clouds from MODIS data and found a median value of 50 cm^{-3} . Thus, we might anticipate that the clouds simulated in the LES cases have a somewhat greater susceptibility to brightening than the “average” marine cloud. The brightening in the deepest MBL case here (P11) does not stand out as being anomalously weak compared with similarly clean cases, although no clear track is produced in the simulated cloud field (Possner et al., 2018). It is important to stress, however, that several low cloud systems (e.g., self-aggregated cumulus, midlatitude stratus) that contribute significantly to low cloud cover over the global oceans are not represented in the LES cases in the literature to date. The LES cases also do not provide sufficient constraints on how brightening changes with injection rate and injected particle size across the different meteorological conditions. Another consideration is that almost all the LES studies examine responses that take place within the first day after injection. As Fig. 2 suggests, a significant fraction of the reflected energy likely takes place between 1-3 days after injection. However, studies suggest that cloud adjustments to aerosol may change significantly over timescales of hours to a few days (Wood 2007; Gryspeerd et al., 2019; Glassmeier et al., 2021), resulting in changes on longer timescales that may, in some cases, offset some of the Twomey brightening. Thus, although the LES simulations here provide some validation of the heuristic model, there is a need for many more simulations to test its sensitivities under the full range of meteorological, background aerosol, and aerosol injection scenarios.

4. Global forcing estimates from the heuristic model

The heuristic model is next used to estimate the global radiative forcing for MCB under different assumptions regarding the number of sprayers and the rate and size of the injected particles. Sensitivity of the forcing to the properties of the background aerosol and updraft speed is also explored.

4.1 Sprayer number and injection rate

Figure 5 presents results as a function of the number of sprayers N_{sprayers} and the salt mass injection rate per sprayer \dot{M}_s . For this case, injected particles have $D_s=100$ nm, and spraying occurs over all eligible ocean areas ($f_{\text{ocean}}=0.54$ and $f_{\text{spray}}=1$ in Eq. 5). A background accumulation mode aerosol concentration $N_{0,\text{acc}}=100$ cm^{-3} is assumed, representative of conditions over the open oceans (Heintzenberg et al., 2000), and a representative coarse mode is included (Sect. 2.6 and Table 1). Other parameters are set to the values provided in Table 1 and discussed in Sect. 2. The assumed MBL depth of 1000 m is representative of typical conditions in which stratiform marine low clouds occur. A global mean radiative forcing magnitude of 1-4 W m^{-2} can be achieved, with forcing generally increasing as total salt mass injection rates increase from ~ 10 to ~ 60 Tg yr^{-1} (Fig. 5a). As the total injection rate increases beyond 100 Tg yr^{-1} there are somewhat diminishing returns in terms of further brightening, and $-\Delta F$ reaches ~ 5 -8 W m^{-2} for an injection rate of 1000 Tg yr^{-1} . The reduced sensitivity as more particles are injected is driven by increased competition for water vapor in the updraft, resulting in a decreasing fraction of injected aerosols activated to form droplets (Fig. 5b, dotted contours). When the injected aerosol concentration is less than a few hundred cm^{-3} , such competition for vapor is relatively modest, and the activated fraction exceeds 70%, but this reduces to only 30-40% at injection rates of 300 Tg yr^{-1} .

Given that $-\Delta F$ increases approximately as a function of total mass injection rate for mass injection rates of < 50 Tg yr^{-1} (Fig. 5a), roughly the same forcing can be achieved either with a smaller number of high throughput sprayers, or a larger number of somewhat weaker sprayers. Scenario (1) in Fig. 5 has $N_{\text{sprayers}}=12000$, each injecting 6×10^{16} particles s^{-1} , for a total mass injection rate of 69 Tg yr^{-1} , achieves the same forcing (-3.7 W m^{-2}) as scenario (2), which has $N_{\text{sprayers}}=10^5$, each injecting 6×10^{15} particles s^{-1} , for a total mass injection rate of 55 Tg yr^{-1} . As we discuss in Sect. 5.1 particle spray rates approaching 10^{17} s^{-1} will likely result in significant particle losses due to high concentrations in the near field of the spray system, and so we consider scenario (1) to be close to the upper end of the injection rates that are likely to be feasible. From this, it may reasonably be concluded that if MCB were ever to be used to achieve a radiative forcing close to that needed to offset a doubling of CO_2 , considerably more sprayers would be needed than are assumed in the estimate from Salter et al. (2008), where only ~ 4500 spray vessels were assumed. The need for a greater number of sprayers in the heuristic model is primarily because of overlapping plumes, which reduce effectiveness by introducing heterogeneity into the injected particle spatial distribution. Plume overlap is not accounted for in Salter et al. (2008), wherein each sprayer uniformly increases the particle concentration over an area of 7.7×10^{10} m^2 . Our assumed track area is $A_t = 5.28 \times 10^{10}$ m^2 (Sect. 2.5) is quite similar to this, but our plumes overlap. The effect of plume overlap is demonstrated by noting that scenario (1), with fewer sprayers than scenario (2), requires $\sim 25\%$ more injected mass to achieve the same forcing. If we set N_{sprayers} to the number that would cover the ocean if there were no overlaps (5300, Sect. 2.5), the heuristic model would require over twice as much mass to produce a forcing sufficient to offset $2x\text{CO}_2$ as in the $N_{\text{sprayers}}=10^5$ case because the track coverage (fraction of the seeded area with at least one overlapping track) in seeded areas only marginally exceeds 50% (Fig. 5b, dashed lines). Thus, almost half of the

490 eligible ocean area remains unperturbed in this case, requiring increases in N_d to offset doubled CO_2 in the perturbed clouds that are harder to achieve (see Fig. 1). Figure 6 (black curves) shows $-\Delta F$ for different values of N_{sprayers} plotted as a function of the total salt mass injection rate to further illustrate the need for a high number of sprayers to minimize the total mass of salt that needs to be injected to achieve a given forcing.

495 A key result from the heuristic model, for $D_s \sim 100$ nm, is that the forcing to offset doubled CO_2 should be achievable with a total salt mass injection rate of ~ 50 -70 Tg yr^{-1} . This is much lower than the natural sea salt flux, which studies suggest ranges from 3,000 to $>10,000$ Tg yr^{-1} (Textor et al., 2006; Grythe et al., 2014). The residence time of natural sea spray particles is considerably shorter than the lifetime ($\tau_{\text{res}}=2$ days) of the injected salt particles because sea spray particles have a much larger mean size. Thus, a perhaps more useful comparison is to examine the mass loading of the injected salt particles, which increases
500 from 0.1 to 1 $\mu\text{g m}^{-3}$ as forcing magnitude increases from 1 to 4 W m^{-2} (Fig. 5c). The coarse mode aerosol assumed in the model (Table 1) has a mass loading of 12.7 $\mu\text{g m}^{-3}$, which is broadly representative of typical salt loadings in the MBL (5-20 $\mu\text{g m}^{-3}$, Jaeglé et al. 2011). Thus, the mass loading of injected particles required to deliver a significant radiative forcing is a relatively small fraction ($\sim 10\%$) of the natural salt burden in the atmosphere. This is not the case with existing climate model studies of MCB, where much higher salt mass injection rates have been required in order to provide a globally significant
505 radiative forcing. The reasons for this are explored the next section.

It is worth comparing the results from the heuristic model with estimates of radiative forcing from commercial shipping. Total SO_2 emissions from shipping are ~ 10 Tg yr^{-1} (see Lauer et al., 2007). Assuming this is all converted into sulfate, this equates to a mass of 15 Tg $\text{SO}_4 \text{ yr}^{-1}$. Introducing injections of 15 Tg of NaCl yr^{-1} would yield a Twomey radiative forcing of
510 ~ 1.5 -2.0 W m^{-2} for $D_s=100$ nm (Fig. 6a). This represents a considerably higher efficacy (forcing per mass of solute) for MCB compared with commercial shipping. Although sea salt is slightly more hygroscopic than sulfate, it is unlikely that composition differences explain the greater efficacy. On the other hand, observations show that the modal diameter of accumulation mode particles over the oceans, which consist mostly of sulfate, tend to be closer to 200 nm diameter than to 100 nm diameter (e.g., Heintzenberg et al., 2000). Although commercial ships do emit a considerable number of small Aitken particles (e.g., Hobbs
515 et al., 2000), one would expect considerable growth into the accumulation mode over the lifetime of the emitted SO_2 . One hypothesis to explain the greater efficacy of MCB is that commercial shipping emissions results in larger accumulation mode particles. Injection rates of 15 Tg yr^{-1} of NaCl particles with $D_s=200$ nm would yield a radiative forcing of only 0.5 W m^{-2} (Fig. 6a), a value within the range of estimates of forcing from marine shipping (0.06-0.6 W m^{-2} , see introduction). Additionally, shipping emissions are much more concentrated geographically than those from our heuristic model (assuming
520 $f_{\text{spray}}=1$), which reduces efficacy. A more thorough treatment of the effects of geographical heterogeneity of sulfate from commercial shipping is beyond the scope of this study.

4.2 Impacts of variations in injected aerosol size and lifetime, and background aerosol concentrations

A key unresolved question concerns what size of injected particles produces the most effective brightening. Prior studies using
525 LES and climate models have used relatively large particles with modal dry diameters exceeding 200 nm. Although particles
of this size serve as very effective CCN, it is important to take into consideration the overall mass injection rate, which
determines the energetic requirements for particle generation and impacts on atmospheric chemistry (see introduction). For
the sprayer number and injection rate from Scenario 2 (previous section, Fig. 5) the optimal geometric mean diameter D_s of
injected particles is 30-60 nm (Fig. 7a). This optimal size range is consistent with the parcel modelling of Connolly et al.
530 (2014) and is relatively insensitive to the background accumulation mode aerosol concentration $N_{0,acc}$. For fixed mass injection
rate, injected aerosol concentration increases as the inverse third power of D_s (Eqns. 5 and 7). For large injected particles
($D_s \sim 200$ nm), most of the injected particles are activated (Fig. 7b), but each particle has a large mass, and so the overall mass
injection rate required to produce a given forcing is roughly five times higher with $D_s = 200$ nm than it is with $D_s = 100$ nm
(Fig. 6a), quantitatively consistent with the GCM sensitivity tests in Partanen et al. (2012). We find that 40% more forcing
535 can be achieved per mass injected if D_s is further decreased from 100 nm to 50 nm (Fig. 7a). With D_s in this optimal range in
terms of forcing per mass injected, although the activated fraction is quite low, the gain in the added aerosol concentration
counters this. This occurs up to a point where competition for vapor draws down supersaturation and there is a reduction in
the number of injected particles that have critical supersaturations sufficiently low to activate. When D_s is smaller than ~ 40 nm,
 N_d begins to decrease again (Fig. 7b). The saturation effect of adding very small particles is also demonstrated by the gray
540 lines in Fig. 6a, which show forcing as function of total salt mass injection rate for $D_s = 50$ nm. For low mass injection rates
(< 10 Tg yr⁻¹), these very small injected particles produce twice as much brightening as particles with $D_s = 100$ nm. At higher
rates, exceeding ~ 50 Tg yr⁻¹, brightening increases very modestly.

We find a major discrepancy between the parcel model activation used here and that estimated using the ARG parameterization.
545 For D_s larger than 100 nm, droplet concentrations from ARG are in general agreement with those from the parcel model
(Fig. 7b), but as the injected particle size decreases, ARG is unable to activate enough droplets. A significant tendency to
underpredict activation fraction has been noted in several prior studies (Ghan et al., 2011; Connolly et al., 2014), and Simpson
et al. (2014) found a systematic underprediction of peak supersaturations estimated with ARG, which we confirmed with the
parcel model (Fig. 7c). This upshot of this issue is that whereas we find that a maximum in brightening for $D_s = 40$ nm, the
550 competition for vapor in ARG is so strong that it prevents activation of almost all injected particles, and so the forcing is close
to zero (Fig. 7a). It will therefore be very important to ensure that activation schemes used in climate modelling for MCB are
sufficiently accurate to represent the unusual size distributions that would be needed for effective implementation of MCB.

Alterskjaer and Kristjánsson (2013, henceforth AK13) used a climate model with ARG as its activation scheme and found that
555 injected Aitken mode particles ($D_s = 44$ nm) produced a strong negative ΔF at the lowest injection rate used (48.2 Tg yr⁻¹), but

a positive ΔF for injection rates exceeding this. The behavior is consistent with our findings using the ARG scheme but is not consistent with the results from the parcel model, where brightening continues to increase with injected mass for particles of this size (Fig. 7a) and there is no cloud darkening (positive ΔF). AK13 also conducted a sensitivity study in which peak supersaturations were fixed at 0.2% and found that the sign of the forcing changed from weakly positive to strongly negative.

560 We find that the suppression of peak supersaturation as D_s decreases is similarly strong in the parcel model and the ARG parameterization for $D_s > 30$ nm (Fig. 7c), implying that additional competition for vapor from the injected particles is not the main reason for the reduced activation fraction in ARG. Instead, it is the general underprediction of peak supersaturation in ARG occurring at all values of D_s that is the main reason for its inability to activate small Aitken particles. Indeed, the fixed supersaturation of 0.2% in the AK13 sensitivity test is quite similar to that in the parcel model (Fig. 7c), and much higher than

565 that for the ARG parameterization, showing that the use of ARG in AK13 is leading to misleading results regarding the efficacy of injecting Aitken mode particles.

Although the optimal geometric mean diameter D_s of injected particles is 30-60 nm (Fig. 7a), there are other reasons against injecting very small Aitken-sized particles, including near-field coagulation and a higher loss rate to Brownian scavenging by cloud droplets. The former is explored in Sect. 5.1. The timescale for Brownian scavenging of injected aerosol in a cloud topped MBL scales with D_s^2 (Seinfeld and Pandis, 2003) and also decreases inversely with $N_d^{2/3}$. For realistic liquid water contents, it can be shown that under MCB (i.e., N_d of a few hundred cm^{-3}) for $D_s = 60$ nm, the timescale for particle losses to Brownian scavenging by cloud droplets is ~ 4 days but falls to only ~ 1 day for $D_s = 30$ nm.

575 There is strong sensitivity of forcing to injected particle residence time τ_{res} (Fig. 6b). A longer τ_{res} increases the area of each sprayer track in proportion to τ_{res}^2 (since both track width and length are proportional to τ_{res} , see Sect. 2.4) but the injected particle concentration over that area scales as τ_{res}^{-1} (Eqn. 7). For the scenario of many overlapping tracks ($N_{\text{sprayers}}=10^5$), the total salt mass required to produce a given forcing scales with τ_{res}^{-1} (Fig. 6b). Thus, the lifetime of injected particles is a key determinant of MCB efficacy, warranting further study.

580 Higher background droplet concentration lowers a cloud's albedo susceptibility ($d\alpha_c/dN_d$, Twomey 1977; Platnick and Twomey 1994). In addition, the increase in droplet concentration ΔN_d with injection is also reduced when $N_{0,\text{acc}}$ is higher (Fig. 7b). This occurs because peak supersaturation is reduced when the background particles are more numerous, and so a lower fraction of the injected aerosol is activated. This results in a forcing that scales more weakly with $N_{0,\text{acc}}$ than would be expected based solely on the albedo susceptibility. As $N_{0,\text{acc}}$ increases from 50 to 150 cm^{-3} , the albedo susceptibility decreases

585 by a factor of three, and yet the magnitude of the radiative forcing (e.g., for $D_s=50$ nm) decreases by less than a factor of two.

The presence of a coarse mode in the unperturbed state imposes a relatively modest decrease in the effectiveness of aerosol injections in brightening clouds. For D_s in the range 50-100 nm, the realistic background coarse mode concentration ($N_{0,coarse}=10\text{ cm}^{-3}$) used throughout this study results in a forcing that is less than 10% smaller than in the absence of a coarse mode (Fig. 8), and the effect is weaker for larger D_s . Coarse mode concentrations vary with wind speed and can reach values of $\sim 20\text{-}50\text{ cm}^{-3}$ at high wind speeds (Zheng et al., 2018). As Fig. 8 shows, it is only at concentrations well in excess of 10 cm^{-3} (mass loadings well more than $10\text{ }\mu\text{g m}^{-3}$) that there would be significant limits to brightening due to the coarse mode. It is important that MCB spray technology does not introduce a significant number of coarse mode particles, as these will reduce brightening. However, it should be noted that the coarse mode mass loadings required to produce a significant dampening of forcing are considerably more than those required to produce brightening using $\sim 100\text{ nm}$ particles.

4.3 Sensitivity to updraft speed

Updraft speed w is a key determinant of the peak supersaturation during the activation process in updrafts (Sect. 2.6). A single value ($w=0.4\text{ m s}^{-1}$) is assumed for the results shown in this study but note that the sensitivity of ΔF to w is strongly dependent upon the injected particle size (Fig. 9a), with sensitivity decreasing strongly as D_s increases. The sensitivity is related to N_d (Fig. 9b), which itself depends upon the peak supersaturation in the updraft (Fig. 9c) and the size distribution of injected and unperturbed aerosol particles. Note that the suppression in peak supersaturation for the perturbed case compared with the unperturbed case is stronger for $D_s = 100\text{ nm}$ than it is for $D_s = 200\text{ nm}$ but falls no further for $D_s = 50\text{ nm}$. Smaller and more numerous particles have a greater surface area and therefore remove vapor more rapidly, but kinetic limitations on growth rates restrict the continuation of this when D_s falls much below 100 nm . For the mass spray rates assumed here (scenario 2, see Sect. 4.1), almost all injected particles activate in updrafts exceeding 0.3 m s^{-1} when $D_s = 200\text{ nm}$ (Fig. 9b). The suppression of peak supersaturation is relatively modest in this case (Fig. 9c). For $D_s = 100\text{ nm}$ the forcing magnitude increase with w is stronger because the N_d' increase with w is stronger. However, it should be noted that for $D_s=100\text{ nm}$, the forcing magnitude only increases by 30% over the range $0.2 < w < 0.6\text{ m s}^{-1}$, indicating relatively weak sensitivity to updraft speed overall. The greatest sensitivity to w occurs for the smallest injected particles ($D_s=50\text{ nm}$) where the forcing increases by 80% over the range $0.2 < w < 0.6\text{ m s}^{-1}$. This reflects the fact that the greatest sensitivity of N_d' to increasing peak supersaturation will occur when the critical supersaturation of the modal diameter (where the greatest number of particles lies) is close to the peak supersaturation. A 50 nm diameter salt particle has a critical supersaturation of 0.25% (e.g., Petters and Kreidenweis 2007), which is similar to the peak supersaturation at $w = 0.4\text{ m s}^{-1}$ (Fig. 9c). Given that activation in real MBL clouds occurs in a spectrum of updrafts (e.g., Snider et al., 2003), this result would caution against the use of injected particles that are too small to increase N_d in the majority of clouds seeded.

4.4 Direct radiative forcing

A recent study with three climate models found that direct forcing from injected salt aerosol may compete with or even exceed the indirect forcing in magnitude (Ahlm et al., 2017). We demonstrated (Sect. 4.2) that injecting particles with sizes
620 $D_s=30-60$ nm produces the greatest brightening for a given salt mass injected, and a forcing to offset doubled CO_2 can be achieved with injection rates below 100 Tg yr^{-1} (dashed circle, Fig. 10a). These “optimal” particle sizes are much smaller than the dry modal diameters of 200, 260 and 880 nm for injected particles in the models used for the GeoMIP assessment of Ahlm et al. (2017). The magnitude of the heuristic model global direct forcing ΔF_{direct} (Sect. 2.7) is very small ($<0.5 \text{ W m}^{-2}$) for total salt mass injection rates of $<100 \text{ Tg yr}^{-1}$ (Fig 10b). Indeed, to generate $\Delta F_{\text{direct}} = -4 \text{ W m}^{-2}$ requires an order of magnitude
625 greater mass injection rate than it does to produce the same forcing from MCB (compare Fig. 10a and 10b). Partanen et al. (2012) found very small contribution of direct radiative forcing with $D_s = 100$ nm, because the same indirect forcing in this case was achieved with ~ 5 times less injected mass than their case (“5×GEO”) with $D_s=200$ nm, wherein direct forcing constituted about 30% of the forcing.

630 The dry size D_s to maximize the direct forcing, for a given mass injection rate, is ~ 110 nm (Fig. 10b), which is around twice as large as the optimal size for MCB. As we have seen (Figs. 6 and 7), producing significant cloud brightening for the D_s values in the models in Ahlm et al. (2017) requires much higher mass injection, and this leads to significant direct radiative forcing. Indeed, for $D_s=880$ nm, the brightening efficiency (Fig. 7) is so low that we would expect very little brightening for spray rates of several hundred Tg yr^{-1} , which is consistent with the very small increases in N_d for the model that injected
635 particles of this size, despite an injection rate of 590 Tg yr^{-1} . In conclusion, the results here demonstrate that *marine cloud brightening is not very effective without clouds* when consideration is given to the injection rates/sizes required to produce a significant radiative forcing from aerosol-cloud interactions.

5 Implications for future work to test marine cloud brightening

The heuristic model results presented here, together with the assessment of LES studies, have implications that may help guide
640 future work to test the concept of MCB to cool Earth. Broadly speaking, these implications fall into three categories: guidance for the engineering development of particle injection (sprayer) systems; guidance for the design of climate model simulations to evaluate the feasibility of regional and global marine cloud brightening; suggestions for future LES modelling.

5.1 Sprayer development considerations

The results presented in Sect. 4.1 suggest that to produce global radiative forcing from MCB that offsets a significant fraction
645 of the forcing from doubling CO_2 , many sprayers will be required. To keep the number to below $\sim 10^5$, particle number injection rates \dot{N}_s of $\sim 10^{16} \text{ s}^{-1}$ will be needed. Similar forcing can be achieved with fewer sprayers, but this will necessitate higher \dot{N}_s .

This implies very high particle concentrations in the near field of the spray system. Taking the spray system to be a collection of nozzles arranged over some area A_0 , spraying into an airflow v_{flow} , then the initial particle concentration N_0 in the immediate wake of the sprayer is $N_0 = \dot{N}_s/A_0 v_{\text{flow}}$. The approach in Turco and Yu (1997) is used to model the downstream particle concentration assuming a fixed coagulation kernel based on a particle diameter $d = 100$ nm and $v_{\text{flow}} = 10$ m s⁻¹. The coagulation kernel is not strongly dependent upon particle diameter for this size range (see Fig. 13.5 in Seinfeld and Pandis, 2003), so variations in injected dry size and the degree of hygroscopic swelling do not have major impacts. It is worth noting that seawater droplets ultimately yielding dry diameters in the effective range for MCB may be significantly larger and may coagulate somewhat more slowly. We consider a diluting slender plume based on the Gaussian plume dispersion which yields a plume cross sectional area (and thus volume) that evolves with time t as $V = V_0 \left(1 + \left[\frac{t}{t_{\text{dil}}}\right]^a\right)$. Here $a = r_y + r_z = 1.49$ and $t_{\text{dil}} = 1.2$ s, with $t_{\text{dil}} = v_{\text{flow}}^{-1} (A_0/\pi R_y R_z)^{1/(r_y+r_z)}$, where the plume widths at distance x downstream of the sprayer in the cross-wind and vertical directions are $\sigma_y(x) = R_y x^{r_y}$ and $\sigma_z(x) = R_z x^{r_z}$ respectively. The constants R_y , r_y , R_z and r_z are those for neutral stability conditions from Klug (1969) as reproduced in Table 18.3 of Seinfeld and Pandis (2003). Numerical simulations had to be performed because the solution does not allow for analytical integration.

Results of the coagulation-dilution calculations (Fig. 11a) indicate that there are relatively weak particle losses from coagulation until particle injection rates \dot{N}_s exceed $\sim 10^{16}$ s⁻¹, above which loss rates grow sharply. Without dilution there are large losses within the first 100 s for rates exceeding 10^{15} s⁻¹, and for rates approaching 10^{17} s⁻¹ most of the losses occur within the first 10 seconds. Dilution immediately downstream of the spray system is therefore most important for particle survival. The volume profile for these simulations is shown in Fig. 11b. We assume that particle concentration within the expanding plume is uniform, which is somewhat unrealistic, because the edges of the plume will become more dilute at a faster rate than those in the center. A multi-shelled Gaussian plume model was employed in Stuart et al. (2013) and this appears to result in somewhat weaker loss rates than we find, but the same general dependencies were found. There is a somewhat weaker dependence of the fraction of particles remaining on A_0 than might be imagined (Fig. 11c), given that A_0 determines the initial concentration of particles, and loss rates scale with N^2 . This is because it takes longer for turbulent eddies to penetrate a wider plume and mix ambient air into the plume core, so that larger A_0 is associated with a longer dilution timescale t_{dil} .

The strong dependence of coagulation losses on \dot{N}_s (Fig. 11d) indicates that as rates approach 10^{17} s⁻¹ the fraction of particles remaining decreases as rapidly as \dot{N}_s increases, which essentially means no increase in the far-field concentration as injection rates are further increased. Because this is not strongly sensitive to A_0 (Fig. 11c), this imposes a limit ($\sim 5 \times 10^{16}$ s⁻¹) on the maximum rate of particles that a ship-deployable spray system can provide to the far-field environment. Recall that the spray scenario to offset doubled CO₂ forcing with 15000 ships discussed in Sect. 4.1 requires a number injection rate that is at this upper limit of feasibility. More rapid dilution than can be provided by boundary layer turbulence may be possible in the first

few seconds downstream of the spray system if the initial flow rate can be increased, but more sophisticated fluid and aerosol
680 dynamics modelling will be required to determine the maximum far-field injection rates that a sprayer can provide.

5.2 Climate modelling

The results of this study have implications for both types of climate modelling MCB studies discussed in the introduction, namely those that fix N_d at some value in seeded regions and those that attempt to model aerosol-cloud interactions using salt aerosol injection.

685

The results presented in Sect. 4.1 demonstrate that there are diminishing returns on increasing N_d as spray rate increases (e.g., Fig. 5b). Producing cloud droplet concentrations of 1000 cm^{-3} is possible but requires mass injection rates approaching 1000 Tg yr^{-1} (compare Fig. 5a and 5b). Locally high mass injection rates would reduce the fractional area of the ocean required for spraying (see next paragraph), but it should be borne in mind that increasing N_d to 1000 cm^{-3} , as has been
690 done in some climate modelling studies (e.g., Rasch et al., 2009; Baughman et al., 2012), would increase the Brownian scavenging rate of interstitial injected aerosol and reduce the overall particle residence time τ_{res} , resulting in a reduced forcing (see Sect. 4.2, Fig. 6b). Increasing N_d from 300 to 1000 cm^{-3} reduces the Brownian scavenging timescale by a factor of over 2 (Seinfeld and Pandis 2003), suggesting that attempting to implement MCB with very high droplet concentrations may not be practical.

695

Several previous climate model studies have seeded a relatively small fraction of the ocean area. Jones et al. (2009) set $N'_d = 375 \text{ cm}^{-3}$ in three regions totalling only 4.7% of the ocean and achieved $\Delta F = -0.97 \text{ W m}^{-2}$. The unperturbed N_d is not known for this study so it is not possible to predict the Twomey forcing for this case using the heuristic model, but for $f_{\text{spray}} = 0.1$, a peak forcing of -1.4 W m^{-2} is achievable (Fig. 12) but achieving a forcing magnitude in excess of 1 W m^{-2}
700 requires a mean N_d of over 800 cm^{-3} ($r_N \sim 8$) in the seeded area. One can only conclude that either the unperturbed N_d in Jones et al. (2009) was very low, or that the model produced significant positive cloud adjustments that augmented the Twomey effect. Fig. 12 suggests that global forcing magnitudes greater than $\sim 4 \text{ W m}^{-2}$ from the Twomey effect alone are only likely to be possible if $\sim 50\%$ of the eligible ocean area ($\sim 40\%$ of the total ocean area) is seeded.

705 Climate modelling to investigate MCB by injecting surface salt sources has typically injected particles with D_s of 200 nm or greater (Alterskjaer et al. 2012, 2013; Ahlm et al., 2017). The heuristic model sensitivity to injected particle size presented in Sect. 4.2 indicates that $D_s = 200 \text{ nm}$ is inefficient (Fig. 7), requiring mass spray rates 5 times higher than for $D_s = 100 \text{ nm}$ and over an order of magnitude higher than for $D_s = 50 \text{ nm}$ (Fig. 6). This has led to mass spray rates in existing MCB studies of hundreds of Tg yr^{-1} (Ahlm et al., 2017), and these high mass spray rates produce significant direct radiative forcing (Ahlm et al., 2017). Our results suggest that smaller injected particles can yield global radiative forcing of -1 to -4 W m^{-2} with very little
710

direct radiative forcing. Providing meaningful global radiative forcing using the aerosol direct effect is an extremely inefficient use of salt particles. We therefore suggest that future climate modelling should focus on smaller injected particles, to build on the sensitivity study in Partanen et al. (2012). This may challenge some models, because a dedicated injection particle mode independent of the model's accumulation mode will be required, and accurate treatment of activation for such cases is a major issue (see Sect. 4.2 and the Appendix).

5.3 Challenges for LES modelling

One notable feature of existing LES studies designed to test the sensitivity of albedo to particle injections (Sect. 3) is that all existing simulation experiments show some degree of brightening, although some do show cloud cover or condensate adjustments that partly offset the Twomey effect. In most cases, especially very clean conditions, cloud adjustments significantly augment Twomey brightening. No LES studies in the literature show domain-wide cloud adjustments that completely offset Twomey brightening. However, predicting cloud adjustments accurately is a challenge, even for LES models; some models do not represent physical processes needed to produce the correct adjustments. Such processes include size-dependent droplet evaporation, which requires an estimation of supersaturation (many LES assume saturation adjustment; see Wang et al., 2003) and sedimentation-entrainment feedback (e.g., Bretherton et al., 2007). Tests to establish the importance of these processes for determining susceptibility to particle injections are incomplete. In addition, existing LES studies represent only a small subset of possible meteorological conditions, focusing primarily upon shallow MBLs that are probably more susceptible to aerosol injections. Studies examining the susceptibility to particle injections of deeper trade wind MBLs including aggregated shallow convective systems need to be conducted to establish the efficacy of MCB in these regions.

It should be noted that most LES cases to test MCB are of insufficient duration to examine the responses at timescales longer than the injected particle residence time τ_{res} , which suggests that longer simulations will be necessary to evaluate the true radiative forcing from injections. Further, no LES studies to date have attempted to constrain the injected particle residence time τ_{res} , which is a key determinant of MCB forcing (Sect. 4.2, Fig. 6b). The dependence of τ_{res} on precipitation, entrainment, in-cloud Brownian scavenging, and other factors, warrants exploration using LES. In addition, there are several potentially important feedbacks involving aerosol residence time that are ideal for study using LES. First, it is well understood that MBL precipitation is a major sink of CCN (Wood et al., 2012; Zheng et al., 2018; Wang et al., 2021). Residence time is expected to be shorter in precipitating MBLs, and suppression of precipitation by high N_d in seeded clouds will increase τ_{res} (Wood 2012), but increased cloud surface area will increase Brownian scavenging of injected particles, reducing τ_{res} . Precipitation suppression will also impact the background aerosol properties, including potentially increasing the concentration of coarse mode and giant CCN that may counter some of the N_d -driven precipitation suppression (Feingold et al., 1999). These feedback processes may affect aerosol residence time and MCB forcing in ways not accounted for in current analyses. Finally, deficiencies in some activation schemes used in LES are likely to be a significant issue hindering accurate representation of

the effects of injected particles smaller than 100 nm (Sect. 4.2, Appendix, and Connolly et al., 2014), so it will be important to ensure that LES models can handle the activation process faithfully.

745 6 Conclusions

This study presents a simple heuristic model to produce useful quantitative estimates of the radiative forcing from the Twomey effect driven by salt particle injections over the global oceans (marine cloud brightening, MCB). The model includes a treatment of individual sprayer plumes and their overlap, and so can be used to explore brightening as a function of the number of sprayers. Brightening is predicted using Twomey's albedo susceptibility given predicted increases in N_d from an activation
750 look-up table derived using Lagrangian parcel modeling that incorporates both background and injected aerosol particle size distributions. Parameters for the model are constrained with observations of cloud cover and N_d from satellites, along with aerosol properties from syntheses of in situ observations. The model performs reasonably well in estimating the cloud brightening from a number of large eddy simulations (LES) reported in the literature, although the LES cases tend to produce more brightening for clean unperturbed states and less for polluted states, likely because of cloud adjustments (changes in
755 cloud cover and/or liquid water path in response to aerosols) that are not included in the heuristic model. The heuristic model is then used to estimate global radiative forcing from MCB and its sensitivity to injected particle spray rates and particle sizes. The key conclusions of the work are:

- Radiative forcing to offset doubled CO₂ can be achieved with global mean salt spray rates of ~50-70 Tg yr⁻¹. This is
760 much lower than the natural sea salt flux, and much lower than spray rates used in global models, which have injected larger particles than are needed to efficiently brighten clouds. To produce this radiative forcing, a large number of sprayers (10^4 - 10^5) will be required to operate over the majority of the 54% of the Earth's surface that is over ocean and remote from land.
- Injected particles with geometric mean dry diameters of 30-60 nm are most efficient at brightening clouds for a
765 fixed mass of salt injected.
- There is no evidence for marine cloud darkening (positive radiative forcing) using the parcel model-based activation scheme, although the Abdul Razzak and Ghan (2000) parameterization incorrectly shows that this occurs for very high concentrations of small injected particles due to excessive competition for water vapor.
- Competition for vapor effectively limits the maximum possible magnitude of radiative forcing from MCB to
770 approximately 8 W m⁻² for salt spray rates less than 1000 Tg yr⁻¹, assuming all ocean regions are seeded. This assumes that negative cloud adjustments remain relatively small compared with the Twomey effect.
- Heuristic model radiative forcing estimates are mostly within a factor of 3 of those from LES, across a range of different spray and unperturbed conditions.

- 775
- Brightening in the heuristic model and the LES decreases strongly with the aerosol/droplet concentrations in the unperturbed clouds, so it is critical to better understand and model the seasonal and geographical variations in these parameters in order to identify optimal locations and times for particle injections and to predict radiative forcing.
 - For injected particles with geometric mean dry diameters of ~ 100 nm or more, there is relatively weak sensitivity to updraft speed for values larger than 0.2 m s^{-1} .
 - Direct radiative forcing from injected particles is very small for mass injection rates less than 100 Tg yr^{-1} . MCB is far less effective without clouds when consideration is given to the quantity of salt that must be injected.
- 780

Appendix: Parcel model emulator

To determine aerosol activation, an explicit Lagrangian parcel model is used to construct a five-dimensional look-up table that predicts peak supersaturation and the concentration of activated aerosol (cloud droplet concentration N_d) as a function of updraft speed w , the concentration N_s and geometric mean dry diameter c of the injected particles respectively, and the unperturbed particle concentrations of accumulation mode $N_{0,acc}$ and coarse mode $N_{0,coarse}$ particles. All other aerosol size and hygroscopicity parameters are fixed at their values in Table 1. The temperature and pressure are fixed at 280 K and 900 hPa respectively. The Lagrangian parcel model is initialized just below cloud base with a saturation ratio of 0.99 and is integrated to a height of 50 m above the saturation level. Particles are determined to be activated if they have reached a diameter of $2 \mu\text{m}$ at this height. Split timestepping is used to render the integration stable for the very small sizes of some of the injected particles. The discretization of the dry size distribution is set for each case to provide an accurate estimate of the number of activated droplets. Parcel model simulations are produced for all 6468 permutations of the values of the five input parameters shown in Table A1. The injected particle concentrations N_s are set to be a function of D_s such that the overall mass of injected particles is the same for a given value of the scaling factor (Table A1). A scaling factor of unity corresponds to an injected mode mass loading of $1.2 \mu\text{g mg}^{-1}$. For the look-up table, basic linear interpolation in five dimensions is used to determine the droplet concentration and peak supersaturation for any given set of input variables w , D_s , N_s , $N_{0,acc}$, and $N_{0,coarse}$.

785

790

795

Figure A1 provides a comparison between the parcel model results and the ARG parameterization. In general, ARG significantly underpredicts the peak supersaturation (Fig. A1 panels a, c, and e) in all cases. Despite this, there is good agreement between the parcel model and ARG droplet concentrations for $D_s > 200$ nm. For this size range, the injected particles have critical supersaturations small enough that despite the underprediction of peak supersaturation in ARG, it is sufficiently high to activate most injected particles. However, as D_s falls below 200 nm, the underprediction of peak supersaturation in ARG has increasingly severe consequences, and this is exacerbated at the highest mass loadings. For $D_s = 50$ nm, ARG N_d is

800

only around 50% of that in the parcel model, and this underprediction falls rapidly as D_s falls further. Based on these findings,
805 we conclude that the biases in ARG are too large for this parameterization to produce useful results.

Code and Data availability

All code and data used in this study are available on request from the author.

Competing interests

810 The authors declare that they have no conflict of interest.

Author contribution

The sole author Robert Wood conceived and carried out the work presented and wrote the manuscript.

Acknowledgements

Support for this work was provided by the National Oceanographic and Atmospheric Administration (NOAA Award
815 NA20OAR4320271 and an Earth's Radiation Budget program grant, NOAA CPO Climate & CI #03-01-07-001), and
from Lowercarbon, the Pritzker Innovation Fund and SilverLining through the Marine Cloud Brightening Project. LES
simulations were performed at the University of Washington by Peter Blossey and Je-Yun Chun, with assistance from Matthew
Wyant. Sarah Doherty, Phil Rasch, Kelly Wanser, Tom Ackerman, Peter Blossey, Matthew Wyant, Ehsan Erfani, Je-Yun
Chun, Armand Neukermans, Chris Bretherton, Gary Cooper, Sean Garner, Kate Murphy, Paul Connolly, and Michael
820 Diamond are thanked for discussions that have helped to frame and improve this work.

References

- Abdul-Razzak, H., & Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types. *Journal of Geophysical
Research: Atmospheres (1984–2012)*, 105(D5), 6837–6844, 2000.
- 825 Ackerman, A. S., Toon, O. B., and Hobbs, P. V.: Numerical modeling of ship tracks produced by injections of cloud
condensation nuclei into marine stratiform clouds, *J. Geophys. Res.*, 100(D4), 7121– 7133, doi:[10.1029/95JD00026](https://doi.org/10.1029/95JD00026), 1995.

- 830 Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., & Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing. *Nature*, 432(7020), 1014–1017. <https://doi.org/10.1038/nature03174>, 2004.
- Ahlm, L., A. Jones, C. W. Stjern, H. Muri, B. Kravitz and J.-E. Kristjánsson: Marine Cloud Brightening – as Effective without Clouds, *Atmos. Chem. Phys.*, 17 (21), 13071–87, doi:10.5194/acp-17-13071-2017, 2017.
- 835 Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness. *Science*, 245, 1227–1230. <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Alterskjær, K., J. E. Kristjánsson, and Ø. Seland: Sensitivity to Deliberate Sea Salt Seeding of Marine Clouds – Observations and Model Simulations, *Atmos. Chem. Phys.*, 12 (5), 2795–2807, doi:10.5194/acp-12-2795-2012, 2012.
- 840 Alterskjaer, K., & Kristjánsson, J. E.: The sign of the radiative forcing from marine cloud brightening depends on both particle size and injection amount. *Geophysical Research Letters*, 40(1), 210–215. <https://doi.org/10.1029/2012GL054286>, 2013.
- 845 Andreae, M. O., & Rosenfeld, D.: Aerosol–cloud–precipitation interactions. Part 1. The nature and sources of cloud-active aerosols. *Earth-Science Reviews*, 89(1), 13–41. <https://doi.org/10.1016/j.earscirev.2008.03.001>, 2008.
- Archer, C. L., and M. Z. Jacobson: Evaluation of global wind power, *J. Geophys. Res.*, 110, D12110, doi:10.1029/2004JD005462, 2005.
- 850 Bala, G., Caldeira, K., Nemani, R., Cao, L., Ban-Weiss, G., & Shin, H.-J.: Albedo enhancement of marine clouds to counteract global warming: Impacts on the hydrological cycle. *Climate Dynamics*, 37(5–6), 915–931. <https://doi.org/10.1007/s00382-010-0868-1>, 2011.
- 855 Baughman, E., Gnanadesikan, A., Degaetano, A., & Adcroft, A.: Investigation of the Surface and Circulation Impacts of Cloud-Brightening Geoengineering. *Journal of Climate*, 25(21), 7527–7543. <https://doi.org/10.1175/JCLI-D-11-00282.1>, 2012.
- 860 Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U.,

- Malavelle, F., Mauritsen, T., ... Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change. *Reviews of Geophysics*, 58(1), e2019RG000660. <https://doi.org/10.1029/2019RG000660>, 2020.
- 865 Bender, F. A.-M., Charlson, R. J., Ekman, A. M. L., & Leahy, L. V.: Quantification of Monthly Mean Regional-Scale Albedo of Marine Stratiform Clouds in Satellite Observations and GCMs. *Journal of Applied Meteorology and Climatology*, 50(10), 2139–2148. <https://doi.org/10.1175/JAMC-D-11-049.1>, 2011.
- Bennartz, R.: Global assessment of marine boundary layer cloud droplet number concentration from satellite, *J. Geophys. Res.*, 112, D02201, doi:10.1029/2006JD007547, 2007.
- 870 Berner, A. H., Bretherton, C. S., & Wood, R.: Large eddy simulation of ship tracks in the collapsed marine boundary layer: A case study from the Monterey area ship track experiment. *Atmos. Chem. Phys.*, 15(10), 5851–5871. <https://doi.org/10.5194/acp-15-5851-2015>, 2015.
- 875 Bohren, C. F., and D. R. Huffman: *Absorption and scattering of light by small particles*. Wiley, New York, 1998, 544 pp.
- Bretherton, C. S., Blossey, P. N., & Uchida, J.: Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo. *Geophysical Research Letters*, 34(3). <https://doi.org/10.1029/2006GL027648>, 2007.
- 880 Bretherton, C. S., Wood, R., George, R. C., Leon, D., Allen, G., & Zheng, X.: Southeast Pacific stratocumulus clouds, precipitation and boundary layer structure sampled along 20° S during VOCALS-REx. *Atmos. Chem. Phys.*, 16, 2010.
- Capaldo, K., Corbett, J. J., Kasibhatla, P., Fischbeck, P., & Pandis, S. N.: Effects of ship emissions on sulphur cycling and radiative climate forcing over the ocean. *Nature*, 400(6746), 743–746. <https://doi.org/10.1038/23438>, 1999.
- 885 Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., & Pierce, J. R.: Large contribution of natural aerosols to uncertainty in indirect forcing. *Nature*, 503(7474), 67–71. <https://doi.org/10.1038/nature12674>, 2013.
- 890 Chosson, F., Paoli, R., & Cuenot, B.: Ship plume dispersion rates in convective boundary layers for chemistry models. *Atmospheric Chemistry and Physics*, 8(16), 4841–4853. <https://doi.org/10.5194/acp-8-4841-2008>, 2008.
- Chun J.-Y., R. Wood, P. Blossey, and M. Wyant: Large eddy simulations of salt tracks in shallow marine boundary layers: sensitivity to injected particle size. Manuscript in preparation, 2021.

Coakley, J. A., Robert L. Bernstein, & Philip A. Durkee: Effect of Ship-Stack Effluents on Cloud Reflectivity. *Science*, 237, 1020–1022, 1987.

Coakley Jr, J. A., & Walsh, C. D.: Limits to the aerosol indirect radiative effect derived from observations of ship tracks. *J. Atmos. Sci.*, 59, 668–680, 2002.

Connolly P. J., McFiggans G. B., Wood R., & Tsiamis A.: Factors determining the most efficient spray distribution for marine cloud brightening. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372(2031), 20140056. <https://doi.org/10.1098/rsta.2014.0056>, 2014.

905

Conover, J. H.: Anomalous Cloud Lines. *J. Atmos. Sci.*, 23, 778–785, [https://doi.org/10.1175/1520-0469\(1966\)023<0778:ACL>2.0.CO;2](https://doi.org/10.1175/1520-0469(1966)023<0778:ACL>2.0.CO;2), 1966.

Cooper G., J. Foster, L. Galbraith, S. Jain, A. Neukermans, B. Ormond: Preliminary results for salt aerosol production intended for marine cloud brightening, using effervescent spray atomization. *Phil. Trans. R. Soc. A*. 372: 20140055. <http://dx.doi.org/10.1098/rsta.2014.0055>, 2014.

Diamond, M. S., Director, H. M., Eastman, R., Possner, A., & Wood, R.: Substantial Cloud Brightening from Shipping in Subtropical Low Clouds. *AGU Advances*, 1, e2019AV000111. <https://doi.org/10.1029/2019AV000111>, 2020.

915

Doelling, D. R., N. G. Loeb, D. F. Keyes, M. L. Nordeen, D. Morstad, C. Nguyen, B. A. Wielicki, D. F. Young, M. Sun: Geostationary Enhanced Temporal Interpolation for CERES Flux Products, *Journal of Atmospheric and Oceanic Technology*, 30(6), 1072-1090. doi: [10.1175/JTECH-D-12-00136.1](https://doi.org/10.1175/JTECH-D-12-00136.1), 2013.

Durkee, P. A., Chartier, R. E., Brown, A., Trehubenko, E. J., Rogerson, S. D., Skupniewicz, C., Nielsen, K. E., Platnick, S., & King, M. D.: Composite ship track characteristics. *J. Atmos. Sci.*, 57, 2542–2553, 2000.

Eyring, V., Isaksen, I. S. A., Berntsen, T., Collins, W. J., Corbett, J. J., Endresen, O., Grainger, R. G., Moldanova, J., Schlager, H., & Stevenson, D. S.: Transport impacts on atmosphere and climate: Shipping. *Atmospheric Environment*, 44(37), 4735–4771. <https://doi.org/10.1016/j.atmosenv.2009.04.059>, 2010.

925

Feingold, G., Cotton, W. R., Kreidenweis, S. M., & Davis, J. T., 1999: The impact of giant cloud condensation nuclei on drizzle formation in stratocumulus: Implications for cloud radiative properties. *Journal of the Atmospheric Sciences*, 56(24), 4100–4117, 1999.

930

Ghan, S. J., Guzman, G., & Abdul-Razzak, H.: Competition between sea salt and sulfate particles as cloud condensation nuclei. *Journal of the Atmospheric Sciences*, 55(22), 3340–3347. [https://doi.org/10.1175/1520-0469\(1998\)055<3340:CBSSAS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<3340:CBSSAS>2.0.CO;2), 1998.

935 Ghan, S. J., Guzman, G., & Abdul-Razzak, H.: Competition between Sea Salt and Sulfate Particles as Cloud Condensation Nuclei. *Journal of the Atmospheric Sciences*, 55(22), 3340–3347. [https://doi.org/10.1175/1520-0469\(1998\)055<3340:CBSSAS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<3340:CBSSAS>2.0.CO;2), 1998.

Ghan, S. J., Abdul-Razzak, H., Nenes, A., Ming, Y., Liu, X., Ovchinnikov, M., Shipway, B., Meskhidze, N., Xu, J., & Shi,
940 X.: Droplet nucleation: Physically-based parameterizations and comparative evaluation. *Journal of Advances in Modeling Earth Systems*, 3(4). <https://doi.org/10.1029/2011MS000074>, 2011.

Glassmeier, F., Hoffmann, F., Johnson, J. S., Yamaguchi, T., Carslaw, K. S., & Feingold, G.: Aerosol-cloud-climate cooling overestimated by ship-track data. *Science*, 371(6528), 485–489. <https://doi.org/10.1126/science.abd3980>, 2021.

945

Gryspeerdt, E., J. Quaas, and N. Bellouin: Constraining the aerosol influence on cloud fraction, *J. Geophys. Res. Atmos.*, 121, doi:10.1002/2015JD023744, 2016.

Gryspeerdt, E., Goren, T., & Smith, T. W. P.: Observing the timescales of aerosol-cloud interactions in snapshot satellite
950 images. *Atmospheric Chemistry and Physics Discussions*, 1–30. <https://doi.org/10.5194/acp-2020-1030>, 2019.

Grythe, H., Ström, J., Krejci, R., Quinn, P., & Stohl, A.: A review of sea-spray aerosol source functions using a large global set of sea salt aerosol concentration measurements. *Atmospheric Chemistry and Physics*, 14(3), 1277–1297, 2014.

955 Hahn, C. J., & Warren, S. G.: *A Gridded Climatology of Clouds over Land (1971-96) and Ocean (1954-97) from Surface Observations Worldwide*. <http://cdiac.ornl.gov/epubs/ndp/ndp026e/ndp026e.html>, 2007.

Hartmann, D. L., & Short, D. A.: On the Use of Earth Radiation Budget Statistics for Studies of Clouds and Climate. *J. Atmos. Sci.*, 37, 1233–1250. [https://doi.org/10.1175/1520-0469\(1980\)037<1233:OTUOER>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<1233:OTUOER>2.0.CO;2), 1980.

960

- Heffter, J. L.: The Variation of Horizontal Diffusion Parameters with Time for Travel Periods of One Hour or Longer. *Journal of Applied Meteorology*, 4, 153–156. [https://doi.org/10.1175/1520-0450\(1965\)004<0153:TVOHDP>2.0.CO;2](https://doi.org/10.1175/1520-0450(1965)004<0153:TVOHDP>2.0.CO;2), 1965.
- 965 Heintzenberg, J., Covert, D. C., & Van Dingenen, R.: Size distribution and chemical composition of marine aerosols: A compilation and review. *Tellus B*, 52(4), 1104–1122, 2000.
- Hindman, E. E., Porph, W. M., Hudson, J. G., & Durkee, P. A.: Ship-produced cloud lines of 13 July 1991. *Atmospheric Environment*, 28, 3393–3403. [https://doi.org/10.1016/1352-2310\(94\)00171-G](https://doi.org/10.1016/1352-2310(94)00171-G), 1994.
- 970
- Hobbs, P. V., Garrett, T. J., Ferek, R. J., Strader, S. R., Hegg, D. A., Frick, G. M., Hoppel, W. A., Gasparovic, R. F., Russell, L. M., Johnson, D. W., & others: Emissions from ships with respect to their effects on clouds. *Journal of the Atmospheric Sciences*, 57(16), 2570–2590, 2000.
- 975 Hoffmann, F., & Feingold, G.: Cloud Microphysical Implications for Marine Cloud Brightening: The Importance of the Seeded Particle Size Distribution, *Journal of the Atmospheric Sciences* (published online ahead of print 2021). Retrieved Aug 17, 2021, from <https://journals.ametsoc.org/view/journals/atsc/aop/JAS-D-21-0077.1/JAS-D-21-0077.1.xml>
- Horowitz, H. M., Holmes, C., Wright, A., Sherwen, T., Wang, X., Evans, M., Huang, J., Jaeglé, L., Chen, Q., Zhai, S., &
- 980 Alexander, B.: Effects of Sea Salt Aerosol Emissions for Marine Cloud Brightening on Atmospheric Chemistry: Implications for Radiative Forcing. *Geophysical Research Letters*, 47(4), e2019GL085838. <https://doi.org/10.1029/2019GL085838>, 2020.
- Isaksen, I. S. A., Granier, C., Myhre, G., Berntsen, T. K., Dalsøren, S. B., Gauss, M., Klimont, Z., Benestad, R., Bousquet,
- 985 P., Collins, W., Cox, T., Eyring, V., Fowler, D., Fuzzi, S., Jöckel, P., Laj, P., Lohmann, U., Maione, M., Monks, P., ... Wuebbles, D.: Atmospheric composition change: Climate–Chemistry interactions. *Atmospheric Environment*, 43, 5138–5192. <https://doi.org/10.1016/j.atmosenv.2009.08.003>, 2009.
- Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., & Lin, J.-T.: Global distribution of sea salt aerosols: New constraints
- 990 from in situ and remote sensing observations. *Atmos. Chem. Phys.*, 11(7), 3137–3157. <https://doi.org/10.5194/acp-11-3137-2011>, 2011.

- Jenkins, A. K. L., Forster, P. M., & Jackson, L. S.: The effects of timing and rate of marine cloud brightening aerosol injection on albedo changes during the diurnal cycle of marine stratocumulus clouds. *Atmospheric Chemistry and Physics*, 13(3), 1659–1673. <https://doi.org/10.5194/acp-13-1659-2013>, 2013.
- Jones A, Haywood J., and Boucher O.: Climate impacts of geoengineering marine stratocumulus clouds *J. Geophys. Res.* 114 D10106, 2009.
- 1000 Klug, W.: A method for determining diffusion conditions from synoptic observations, *Staub-Reinhalt. Luft*, 29, 14–20, 1969.
- Latham, J.: Control of global warming? *Nature*, 347, 339–340. <https://doi.org/10.1038/347339b0>, 1990.
- Latham, J., P. Rasch, C.-C. Chen, L. Kettles, A. Gadian, A. Gettelman, H. Morrison, K. Bower, and T. Choulaton: Global Temperature Stabilization via Controlled Albedo Enhancement of Low-Level Maritime Clouds, *Phil. Trans. Royal Society A*, 366, no. 1882, 3969–87. <https://doi.org/10.1098/rsta.2008.0137>, 2008.
- 1005
- Latham, J., et al.: Marine cloud brightening, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 370, 4217–4262, <https://doi.org/10.1098/rsta.2012.0086>, 2012.
- 1010
- Lauer, A., Eyring, V., Hendricks, J., Jöckel, P., & Lohmann, U.: Global model simulations of the impact of ocean-going ships on aerosols, clouds, and the radiation budget. *Atmospheric Chemistry and Physics*, 7(19), 5061–5079, 2007.
- Lohmann, U., & Feichter, J.: Global indirect aerosol effects: A review. *Atmospheric Chemistry and Physics*, 5(3), 715–737, 2005.
- 1015
- Malavelle, F. F., Haywood, J. M., Jones, A., Gettelman, A., Clarisse, L., Bauduin, S., Allan, R. P., Karset, I. H. H., Kristjánsson, J. E., Oreopoulos, L., Cho, N., Lee, D., Bellouin, N., Boucher, O., Grosvenor, D. P., Carslaw, K. S., Dhomse, S., Mann, G. W., Schmidt, A., ... Thordarson, T.: Strong constraints on aerosol–cloud interactions from volcanic eruptions. *Nature*, 546(7659), 485–491. <https://doi.org/10.1038/nature22974>, 2017.
- 1020
- McFiggans, G., Artaxo, P., Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S., Gysel, M., Laaksonen, A., Lohmann, U., Mentel, T. F., Murphy, D. M., O’Dowd, C. D., Snider, J. R., & Weingartner, E.: The effect of physical and chemical aerosol properties on warm cloud droplet activation. *Atmos. Chem. Phys.*, 6(9), 2593–2649. <https://doi.org/10.5194/acp-6-2593-2006>, 2006.
- 1025

- Medeiros, B., Nuijens, L., Antoniazzi, C., & Stevens, B.: Low-latitude boundary layer clouds as seen by CALIPSO. *Journal of Geophysical Research*, 115(D23). <https://doi.org/10.1029/2010JD014437>, 2010.
- 1030 National Research Council: *Climate Intervention: Reflecting Sunlight to Cool Earth*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/18988>, 2015.
- Nicholls, S., & Leighton, J.: An observational study of the structure of stratiform cloud sheets: Part I. Structure. *Quarterly Journal of the Royal Meteorological Society*, 112(472), 431–460, 1986.
- 1035 Partanen, A.-I., Kokkola, H., Romakkaniemi, S., Kerminen, V.-M., Lehtinen, K. E. J., Bergman, T., Arola, A., & Korhonen, H.: Direct and indirect effects of sea spray geoengineering and the role of injected particle size. *Journal of Geophysical Research: Atmospheres*, 117(D2). <https://doi.org/10.1029/2011JD016428>, 2012.
- 1040 Partanen, A. I., Laakso, A., Schmidt, A., Kokkola, H., Kuokkanen, T., Pietikäinen, J. P., et al.: Climate and air quality trade-offs in altering ship fuel sulfur content. *Atmospheric Chemistry and Physics*, 13(23), 12059–12071. <https://www.atmos-chem-phys.net/13/12059/2013/>, 2013.
- Peters, K., Stier, P., Quaas, J., and Graßl, H.: Aerosol indirect effects from shipping emissions: sensitivity studies with the global aerosol-climate model ECHAM-HAM, *Atmos. Chem. Phys.*, 12, 5985–6007, <https://doi.org/10.5194/acp-12-5985-2012>, 2012.
- 1045 Petters, M. D., & Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity. *Atmospheric Chemistry and Physics*, 7(8), 1961–1971, 2007.
- 1050 Platnick, S., & Twomey, S.: Determining the Susceptibility of Cloud Albedo to Changes in Droplet Concentration with the Advanced Very High Resolution Radiometer. *Journal of Applied Meteorology*, 33(3), 334–347. [https://doi.org/10.1175/1520-0450\(1994\)033<0334:DTSOCA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0334:DTSOCA>2.0.CO;2), 1994.
- 1055 Possner, A., Wang, H., Wood, R., Caldeira, K., and Ackerman, T. P.: The efficacy of aerosol–cloud radiative perturbations from near-surface emissions in deep open-cell stratocumuli, *Atmos. Chem. Phys.*, 18, 17475–17488, <https://doi.org/10.5194/acp-18-17475-2018>, 2018.

- 1060 Possner, A., Eastman, R., Bender, F., and Glassmeier, F.: Deconvolution of boundary layer depth and aerosol constraints on
cloud water path in subtropical stratocumulus decks, *Atmos. Chem. Phys.*, 20, 3609–3621, <https://doi.org/10.5194/acp-20-3609-2020>, 2020.
- 1065 Pringle, K. J., Tost, H., Pozzer, A., Pöschl, U., & Lelieveld, J.: Global distribution of the effective aerosol hygroscopicity
parameter for CCN activation. *Atmos. Chem. Phys.*, 10(12), 5241–5255. <https://doi.org/10.5194/acp-10-5241-2010>, 2010.
- Ramanathan, V., Cess, R. D., Harrison, E. F., Minnis, P., Barkstrom, B. R., Ahmad, E., & Hartmann, D.: Cloud-Radiative
Forcing and Climate: Results from the Earth Radiation Budget Experiment. *Science*, 243, 57–63.
<https://doi.org/10.1126/science.243.4887.57>., 1989.
- 1070 Rasch, P. J., J. Latham, and C.-C. Chen: Geoengineering by Cloud Seeding: Influence on Sea Ice and Climate System, *Env.
Res. Lett.*, 4, no. 4, 045112, <https://doi.org/10.1088/1748-9326/4/4/045112>, 2009.
- Ricou, F. P., & Spalding, D. B.: Measurements of entrainment by axisymmetrical turbulent jets. *Journal of Fluid Mechanics*,
11(1), 21–32. <https://doi.org/10.1017/S0022112061000834>, 1961.
- 1075 Salter, S., Sortino G., and Latham, J.: Sea-going hardware for the cloud albedo method of reversing global warming. *Phil.
Trans. R. Soc. A*, 366, 2989–4006, 2008.
- Schreier, M., Mannstein, H., Eyring, V., & Bovensmann, H.: Global ship track distribution and radiative forcing from 1 year
1080 of AATSR data. *Geophysical Research Letters*, 34(17). <https://doi.org/10.1029/2007GL030664>, 2007.
- Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S.,
Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., ...
Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations.
1085 *Atmos. Chem. Phys.*, 6(12), 5225–5246. <https://doi.org/10.5194/acp-6-5225-2006>, 2006.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 2nd ed., John
Wiley and Sons, Inc., Hoboken, New Jersey, 1203pp, 2003.
- 1090 Simpson, E., Connolly, P., & McFiggans, G.: An investigation into the performance of four cloud droplet activation
parameterisations. *Geosci. Model Dev.*, 7(4), 1535–1542. <https://doi.org/10.5194/gmd-7-1535-2014>, 2014.

Slingo, A.: Sensitivity of the Earth's radiation budget to changes in low clouds. *Nature*, 343(6253), 49–51.
<https://doi.org/10.1038/343049a0>, 1990.

1095

Snider, J. R., Guibert, S., Brenguier, J.-L., & Putaud, J.-P.: Aerosol activation in marine stratocumulus clouds: 2. Köhler and parcel theory closure studies. *J. Geophys. Res., Atmos.*, 108(D15). <https://doi.org/10.1029/2002JD002692>, 2003.

1100

Sorooshian, A., Feingold, G., Lebsack, M. D., Jiang, H., & Stephens, G. L.: Deconstructing the precipitation susceptibility construct: Improving methodology for aerosol-cloud precipitation studies. *Journal of Geophysical Research*, 115(D17).
<https://doi.org/10.1029/2009JD013426>, 2010.

1105

Stephens, G. L.: Radiation Profiles in Extended Water Clouds. I: Theory. *J. Atmos. Sci.*, 35, 2111–2122.
[https://doi.org/10.1175/1520-0469\(1978\)035<2111:RPIEWC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2111:RPIEWC>2.0.CO;2), 1978.

1110

Stjern, C. W., Muri, H., Ahlm, L., Boucher, O., Cole, J. N. S., Ji, D., Jones, A., Haywood, J., Kravitz, B., Lenton, A., Moore, J. C., Niemeier, U., Phipps, S. J., Schmidt, H., Watanabe, S., & Kristjánsson, J. E.: Response to marine cloud brightening in a multi-model ensemble. *Atmospheric Chemistry and Physics*, 18(2), 621–634. <https://doi.org/10.5194/acp-18-621-2018>, 2018.

1115

Stuart, G. S., Stevens, R. G., Partanen, A.-I., Jenkins, A. K. L., Korhonen, H., Forster, P. M., Spracklen, D. V., & Pierce, J. R.: Reduced efficacy of marine cloud brightening geoengineering due to in-plume aerosol coagulation: Parameterization and global implications. *Atmospheric Chemistry and Physics*, 13(20), 10385–10396. <https://doi.org/10.5194/acp-13-10385-2013>, 2013.

Tang, I. N.: Chemical and size effects of hygroscopic aerosols on light scattering coefficients. *Journal of Geophysical Research: Atmospheres*, 101(D14), 19245–19250. <https://doi.org/10.1029/96JD03003>, 1996.

1120

Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., ... Tie, X: Analysis and quantification of the diversities of aerosol life cycles within AeroCom. *Atmos. Chem. Phys.*, 6(7), 1777–1813.
<https://doi.org/10.5194/acp-6-1777-2006>, 2006:

1125

Terai, C. R., Wood, R., & Kubar, T. L.: Satellite estimates of precipitation susceptibility in low-level marine stratiform clouds: *Journal of Geophysical Research: Atmospheres*, 120(17), 8878–8889. <https://doi.org/10.1002/2015JD023319>, 2015.

- Toll, V., Christensen, M., Quaas, J., & Bellouin, N.: Weak average liquid-cloud-water response to anthropogenic aerosols. *Nature*, 572(7767), 51–55. <https://doi.org/10.1038/s41586-019-1423-9>, 2019.
- 1130 Trofimov, H., Bellouin, N., & Toll, V.: Large-Scale Industrial Cloud Perturbations Confirm Bidirectional Cloud Water Responses to Anthropogenic Aerosols. *Journal of Geophysical Research: Atmospheres*, 125(14), e2020JD032575. <https://doi.org/10.1029/2020JD032575>, 2020.
- 1135 Turco, R. P., & Yu, F.: Aerosol invariance in expanding coagulating plumes. *Geophysical Research Letters*, 24(10), 1223–1226. <https://doi.org/10.1029/97GL01092>, 1997.
- Twomey, S.: Pollution and the planetary albedo. *Atmospheric Environment*, 8(12), 1251–1256, 1974.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds. *J. Atmos. Sci.*, 34, 1149–1152. [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.
- 1140 Wang, S., Wang, Q., & Feingold, G.: Turbulence, condensation, and liquid water transport in numerically simulated nonprecipitating stratocumulus clouds. *Journal of the Atmospheric Sciences*, 60(2), 262–278, 2003.
- 1145 Wang, H., Rasch, P. J., & Feingold, G.: Manipulating marine stratocumulus cloud amount and albedo: A process-modelling study of aerosol-cloud-precipitation interactions in response to injection of cloud condensation nuclei. *Atmos. Chem. Phys.*, 11(9), 4237–4249. <https://doi.org/10.5194/acp-11-4237-2011>, 2011.
- 1150 Wang, Y., Xia, W., Liu, X., Xie, S., Lin, W., Tang, Q., Ma, H.-Y., Jiang, Y., Wang, B., & Zhang, G. J.: Disproportionate control on aerosol burden by light rain. *Nature Geoscience*, 14(2), 72–76. <https://doi.org/10.1038/s41561-020-00675-z>, 2021.
- Wex, H., McFiggans, G., Henning, S., & Stratmann, F.: Influence of the external mixing state of atmospheric aerosol on derived CCN number concentrations. *Geophysical Research Letters*, 37(10). <https://doi.org/10.1029/2010GL043337>, 2010.
- 1155 Wood, R.: Drizzle in Stratiform Boundary Layer Clouds. Part I: Vertical and Horizontal Structure. *Journal of the Atmospheric Sciences*, 62(9), 3011–3033. <https://doi.org/10.1175/JAS3529.1>, 2005.
- 1160 Wood, R.: Cancellation of Aerosol Indirect Effects in Marine Stratocumulus through Cloud Thinning. *J. Atmos. Sci.*, 64, 2657–2669. <https://doi.org/10.1175/JAS3942.1>, 2007.

- Wood, R., Irons, S., & Jonas, P. R.: How important is the spectral ripening effect in stratiform boundary layer clouds? Studies using simple trajectory analysis. *Journal of the Atmospheric Sciences*, 59(18), 2681-2693, 2002.
- 1165 Wood, R., Leon, D., Lebsock, M., Snider, J., & Clarke, A. D.: Precipitation driving of droplet concentration variability in marine low clouds: *Journal of Geophysical Research: Atmospheres*, 117(D19), <https://doi.org/10.1029/2012JD018305>, 2012.
- Xue, H., & Feingold, G.: Large-Eddy Simulations of Trade Wind Cumuli: Investigation of Aerosol Indirect Effects. *Journal of the Atmospheric Sciences*, 63(6), 1605–1622. <https://doi.org/10.1175/JAS3706.1>, 2006.
- 1170 Zelinka, M. D., T. Andrews, P. M. Forster, and K. E. Taylor: Quantifying components of aerosol-cloud-radiation interactions in climate models, *J. Geophys. Res. Atmos.*, 119, 7599–7615, doi:10.1002/2014JD021710, 2014.
- 1175 Zheng, Y., D. Rosenfeld, and Z. Li: Quantifying cloud base updraft speeds of marine stratocumulus from cloud top radiative cooling, *Geophys. Res. Lett.*, 43, 11,407–11,413, doi:10.1002/2016GL071185, 2016.
- Zheng, G., Wang, Y., Aiken, A. C., Gallo, F., Jensen, M. P., Kollias, P., Kuang, C., Luke, E., Springston, S., Uin, J., Wood, R., & Wang, J.: Marine boundary layer aerosol in the eastern North Atlantic: Seasonal variations and key controlling processes. *Atmospheric Chemistry and Physics*, 18(23), 17615–17635. <https://doi.org/10.5194/acp-18-17615-2018>, 2018.
- 1180 Zieger, P., Väisänen, O., Corbin, J. C., Partridge, D. G., Bastelberger, S., Mousavi-Fard, M., Rosati, B., Gysel, M., Krieger, U. K., Leck, C., Nenes, A., Riipinen, I., Virtanen, A., & Salter, M. E.: Revising the hygroscopicity of inorganic sea salt particles. *Nature Communications*, 8(1), 15883. <https://doi.org/10.1038/ncomms15883>, 2017.
- 1185

Table 1: Parameters used in the heuristic model and their assumed values


Symbol	Parameter	Assumed value(s)	Justification
α_c	Unperturbed cloud albedo	0.56	Bender et al. (2011). See text.
ϕ_{atm}	Atmospheric correction factor	0.70	Based on Diamond et al. (2020).
f_{ocean}	Fraction of Earth's surface covered by ocean eligible for spraying	0.54	Divide globe into $10 \times 10^\circ$ boxes (approximate sprayer plume length). Only boxes with less than 10% land eligible for spraying to minimize plumes intercepting land areas. See text.
f_{spray}	Fraction of eligible ocean areas in which sprayers operate	0.5-1.0	
f_{low}	Fraction of sprayed area covered by stratiform low clouds unobscured by high clouds	Function of f_{spray} decreasing from 0.68 for $f_{\text{spray}} < 0.1$ to 0.33 for $f_{\text{spray}} = 1$	Uses MODIS liquid cloud fraction. For $f_{\text{spray}} < 1$, sort eligible $10 \times 10^\circ$ boxes by their monthly climatological mean liquid cloud fraction. Set f_{low} equal to the mean of the cloudiest f_{spray} fraction. See Equation 4 in Sect. 2.2
F_\odot	Solar irradiance	342 W m ⁻²	Assumed day+night averaged global mean solar irradiance.
\dot{M}_s	Rate of NaCl injection by each sprayer	1-1000 kg hr ⁻¹	Variable
N_{sprayers}	Number of sprayer vessels deployed	3×10^3 - 3×10^5	Variable
D_s	Geometric mean diameter of injected NaCl particles	10-1000 nm	Variable
S	Geometric standard dev. of injected NaCl particle size	1.6	
τ_{res}	Residence time of injected particles	2 days	Based on Wood et al. (2012)
$D_{0,\text{acc}}$ $D_{0,\text{coarse}}$	Geometric mean diameter of background aerosol	175 nm (acc.) 615 nm (coarse)	Accumulation mode size values based on marine aerosol climatology of Heintzenberg et al. (2000). Coarse mode values from Zheng et al. (2018).
$S_{0,\text{acc}}$ $S_{0,\text{coarse}}$	Geometric standard dev. of background aerosol size	1.5 (acc.) 1.8 (coarse)	
$N_{0,\text{acc}}$ $N_{0,\text{coarse}}$	Number concentration of background aerosol	50-150 cm ⁻³ (acc.) 10 cm ⁻³ (coarse)	Coarse mode value from summer mean at the Graciosa Island from Zheng et al. (2018).
κ	Aerosol hygroscopicity	0.7(acc.) 1.2 (coarse, injected)	Accumulation mode from Pringle et al. (2010) Coarse mode/injected salt from Petters and Kreidenweis (2007)
w	Updraft speed for aerosol activation	0.4 m s ⁻¹	Approximate value based on numerous stratocumulus field experiments
U_0	Mean surface wind speed	7 m s ⁻¹	Mean near-surface wind over global ocean (Archer and Jacobson 2005)
h	MBL depth	1 km	Typical mean value for marine low clouds over oceans
K	Plume lateral spread rate	1.85 km h ⁻¹	Based on observed ship track spreading rate (Durkee et al., 2000)

Table 2: Large eddy simulation studies included in this study. Information included in this table focused on highlighting diversity in injected particles and domain size.


Study	<i>Wang et al. (2011)</i> [W11]	<i>Berner et al. (2015)</i> [B15]	<i>Jenkins et al. (2013)</i> [J13]	<i>Possner et al. (2018)</i> [P18]	<i>Chun et al. (2020)</i> [C21]	
Case info	DYCOMS-II RF02	Collapsed MBL (Sanko Peace)	DYCOMS-II RF02	VOCALS RF06 Deep open cell	(a) Collapsed MBL (Sanko Peace)	(b) CGILS S12 Shallow MBL
MBL depth, m	900	350	750	1500	350	700
Spray rate, # s⁻¹ [diameter, nm]	1.04×10 ¹⁶ [200]	1.8×10 ¹⁵ (clean) 1.8×10 ¹⁵ (poll.) [200]	5.6×10 ¹⁶ 1.1×10 ¹⁶ (weak) [200]	1.04×10 ¹⁶ [600]	10 ¹⁶ [50]	10 ¹⁶ [50]
Spray duration	30 hr	10.7 min	30 min	48 hr	7.6 min	7.6 min
Mass of NaCl emitted total, kg [per hour, kg]	10320 [344]	10 [57]	1105 [2210]	4.5×10 ⁵ [9400]	0.7 [5.2]	0.7 [5.2]
Simulation duration	30 hr	8 hr	5 hr	48 hr	8 hr	8 hr
Domain size, km×km [area, km²]	60×120 [7200]	51.2×12.8 [660]	9×9 [81]	180×180 [32000]	48×9.6 [460]	24×4.8 115
Number of particles emitted [cm⁻³]	174	4	1650	370	29	57
Spray details	Sprayer traverses long edge of domain several times	Sprayer passes through short edge of domain once only	Sprayer traverses domain once only	Sprayer traverses domain several times	Sprayer traverses short edge of domain once only	
Simulation experiments used	Wet and dry profiles used. Wet cases 50, 100 and 200 cm ³ CCN initially. Dry case has 100 cm ³ CCN. Use single sprayer and uniform seeding only	Basetrack (clean) and SensHiAer (polluted) case used.	Non-precipitating (NP-Ch; NP-Pa) and Precipitating (WP) cases used. Sensitivity study with weaker sprayer on WP.	Single experiment	Two simulation experiments, one with reduced sea spray aerosol	Single experiment

Table A1: Parameter values used to construct the activation look-up table

Variable	Name	Values used
w	Updraft speed	0.2, 0.4, 0.6 m s ⁻¹
D_s	Injected particle geometric mean dry diameter	15, 30, 45, 60, 80, 100, 150, 200, 300, 500, 1000 nm
N_s	Injected particle concentration	Scaling factor of [0, 0.1, 0.3, 1, 3, 10, 30] times $400(100/D_s)^3$ mg ⁻¹ , where D_s is in nm
$N_{0,acc}$	Unperturbed accumulation mode concentration	5, 10, 50, 100, 150, 200, 500 mg ⁻¹
$N_{0,coarse}$	Unperturbed coarse mode concentration	0, 3, 10, 30 mg ⁻¹




1215 **Figure 1: Global radiative forcing from marine cloud brightening ΔF as a function of the ratio of the perturbed to unperturbed (background) cloud droplet concentration $r_N = N'_d/N_d$. Curves are shown for the case where sprayers are deployed over all eligible ocean regions ($f_{\text{spray}} = 1$, black lines) and where sprayers are deployed over only 50% of these areas ($f_{\text{spray}} = 0.5$, gray lines), for unperturbed cloud albedos α_c ranging from 0.3-0.7. The fraction of the Earth's surface area eligible for seeding is $f_{\text{ocean}} = 0.54$, and the atmospheric correction factor $\phi_{\text{atm}} = 0.7$**



1220

Figure 2: (a) Plan view of a realistic laterally-spreading plume/track (top) and the track assumed in the heuristic model, as a function of time/distance downstream of the sprayer. The shading qualitatively indicates the injected particle concentration. (b) Injected aerosol concentration as a function of time for the two plume types given the spray injection information in the box. A residence time $\tau_{res} = 2$ days, and a widening rate $K = 1.85 \text{ km hr}^{-1}$ are assumed, and the model is run out to 10 days to capture the total reflected solar radiation for the spreading plume. (c) additional reflected solar radiation per meter length of track from aerosol-cloud interactions for the two plume types. The total additional reflected energy E_{tot} from the two plumes is very similar. In this case, approximately 43% of the energy reflected from the spreading plume occurs for times $t > \tau_{res}$.


1225



1230 **Figure 3: Probability density functions $p(n)$ derived from the Monte Carlo simulations of overlapping rectangular tracks for three**
values of the mean track density ζ (0.4, 3.3 and 10). For the simulations, a domain of size 4000×4000 km is modelled using a
4000×4000 array, and tracks of length 1200 km and width 44 km (see Sect. 2.5) are placed randomly to the array, assuming periodic
 1235 **boundary conditions. The long dimension of each track is randomly set to be parallel to the i or j direction of the box. Poisson**
distributions (Equation 8) are shown corresponding based on the mean track densities and represent an excellent fit to the data. The
track densities $\zeta = 0.4, 3.3$ and 10 correspond to a total number of ships, if spraying were to take place over the entire eligible ocean
region, of $\sim 3000, \sim 17000$, and ~ 50000 respectively.

1240

1245



1255

1260

Figure 4: Comparison of heuristic model and LES results. Each color indicates a different study (see Table 2 for details). For W11, open circles indicate cases with uniform seeding across the domain. (a) Brightening (radiative forcing) for the LES and heuristic model. Dashed line indicates agreement, and dotted lines represent factor of 3 differences; (b) Ratio of the brightening in the heuristic model to that in the LES, plotted against the droplet concentration in the unperturbed case N_d . The dashed line shows the linear least squares fit to the data; (c) Modeled vs LES cloud droplet concentration given the injection rates and particle size distribution employed in the model (see text); Sensitivity of the normalized forcing (expressed as Joules per injected particle) to the unperturbed N_d for (d) the LES experiments; (e) the heuristic model, with lines representing least-squares fits to the data; (f) the brightening efficiency expressed as the energy reflected over the course of the simulation experiments per mass of salt injected. For J13, the triangle indicates reduced injection rate by a factor of 5, with the arrows connecting the simulation experiments with full and reduced injection rates.






Figure 5: (a) Global mean radiative forcing ΔF (colors) and total flux of sodium chloride (dotted contours) from MCB applied to all eligible ocean areas (54% of Earth's surface) as a function of the number of sprayers N_{sprayers} and the salt mass injection rate \dot{M}_s for each sprayer; (b) increase in cloud droplet concentration ΔN_d (colors), mean fraction of aerosol activated in tracks (dotted contours), and track coverage (dashed contours); (c) injected aerosol number concentration in tracks (colors) and mean mass loading in the MBL of injected salt (dotted contours). The inset of panel (a) shows the key model parameters, with others as in Table 1. Two scenarios (1) and (2), which each produce sufficient forcing to offset doubled CO₂ are highlighted. Scenario (2) has a higher number of sprayers but a lower rate of particles injected per sprayer.



1280 **Figure 6: Radiative forcing as a function of the total (global) rate of salt mass injection, for (a) three sprayer numbers ($N_{\text{sprayers}}=5300$, 15000, and 120000), and for geometric mean spray diameters D_s of 50 nm (gray), 100 nm (black), and 200 nm (yellow); (b) forcing for $N_{\text{sprayers}}=120000$ and $D_s=100$ nm as a function of injected particle lifetime. All other parameters are the same as those used in Fig. 5.**

1285



290

Figure 7: (a) Global mean radiative forcing $-\Delta F$ for a fixed salt mass spray rate (based on Scenario 2, see legend) as a function of injected particle geometric mean diameter D_s for three unperturbed accumulation mode aerosol concentrations $N_{0,\text{acc}}=50, 100$ and 150 cm^{-3} . Black curves show the results from the parcel model and gray curves from the ARG parameterization; (b) Change in mean cloud droplet concentration ΔN_d and aerosol concentration ΔN_a in regions where sprayers are operating; (c) Peak supersaturation in the updraft. The gray shaded box indicates the most effective range of D_s .




Figure 8: Global mean radiative forcing $-\Delta F$ for a fixed salt mass spray rate (based on Scenario 2, see legend) as a function of background coarse mode aerosol concentration $N_{0,coarse}$ for three injected particle sizes $D_s=50, 100$ and 200 nm. The equivalent salt mass loading in the coarse mode is indicated by the top axis.

295





Figure 9: Impact of assumed updraft speed on (a) global mean radiative forcing $-\Delta F$; (b) cloud droplet concentration N_d ; (c) peak supersaturation during activation, for a fixed salt mass spray rate (based on Scenario 2, see legend). Results for injected particle sizes $D_s=50, 100$ and 200 nm are shown. Panels (b) and (c) also show the values for the unperturbed case.

1300

1305




1315


Figure 10: Global mean radiative forcing from (a) aerosol-cloud interactions (i.e., marine cloud brightening) ΔF ; (b) direct radiative forcing of injected aerosol, ΔF_{direct} ; (c) ratio of direct to total radiative forcing, plotted as a function of the total salt injection rate and the geometric mean diameter D_s of the injected aerosol. The number of sprayers $N_{\text{sprayers}}=100000$, and all other parameters are the same as scenario 2 (see Fig. 5 and Sect. 4.1). The three models used in Ahlm et al. (2017) are shown (see legend in panel b), although it should be noted that for these models injections were confined to the Tropical belt (30°S-30°N).

1320

1325




1330 **Figure 11: Effects of coagulation on the concentration of particles at time t downstream of a hypothetical sprayer system with cross**
sectional area $A=4 \text{ m}^2$ and flow rate of air across the sprayer $v_{\text{flow}}=10 \text{ m s}^{-1}$. Here, a single particle diameter $d=100 \text{ nm}$ is assumed,
and solutions follow Turco and Yu (1997). (a) Particle concentrations with dilution proceeding according to the dispersion rates for
neutral conditions from Klug (1969), as reproduced in Seinfeld and Pandis (2006, Table 18.3); concentrations in the absence of plume
dilution are shown for comparison (gray) indicating major losses and eventual asymptotic solution; (b) the ratio of volume V to
initial volume V_0 increases super-linearly with time with dilution timescale $t_{\text{dil}}=1.2 \text{ s}$; (c) fraction of particles remaining at $t=10,000 \text{ s}$
1335 **as a function of initial plume cross sectional area A_0 for spray rate $\dot{N}_s=10^{17} \text{ s}^{-1}$; (d) fraction of particles remaining at $t=10,000 \text{ s}$**
as a function of \dot{N}_s for $A_0=4 \text{ m}^2$. The dotted line shows the scaling such that the fraction remaining decreases at the same rate as \dot{N}_s
increases, i.e., there is no increase in far-field particle concentration with increasing sprayer output.



1340

Figure 12: Sensitivity of the radiative forcing to different fractions f_{spray} of the eligible ocean where sprayers operate. Forcing is shown as a function of the total (global) rate of salt mass injection. The mass injected per sprayer is the same as that for scenario 2 (Sect. 4.1 and Fig. 5) and the sprayer density is the same in each case, i.e., the number of sprayers is proportional to f_{spray} and is 100,000 for $f_{\text{spray}}=1$. The geometric mean spray diameters $D_s=100 \text{ nm}$ and the background $N_{0,\text{acc}}=100 \text{ cm}^{-3}$. All other parameters are the same as those used in Fig. 5.

1345



1350

Figure A1: Comparison of the parcel model and ARG peak supersaturations (panels a, c, e) and activated cloud droplet concentrations N_d (panels b, d, f) as a function of the injected particle geometric mean diameter D_s and the mass loading of injected particles. The other parameters are set as $w=0.4 \text{ m s}^{-1}$, $N_{0,\text{acc}}=100 \text{ mg}^{-1}$, $N_{0,\text{coarse}}=10 \text{ mg}^{-1}$ (base values used throughout the paper). The top row shows results from the parcel model, the center from ARG, and the bottom row is the ratio of ARG to the Parcel model.