

Supplementary materials

For

Physical and chemical properties of black carbon and organic matter from different sources using aerodynamic aerosol classification

Dawei Hu^{1,*}, M. Rami Alfarra^{1,2,#}, Kate Szpek³, Justin M. Langridge³, Michael I. Cotterell⁵, Claire Belcher⁴, Ian Rule⁴, Zixia Liu⁴, Chenjie Yu¹, Yunqi Shao¹, Aristeidis Voliotis¹, Mao Du¹, Brett Smith⁶, Greg Smallwood⁶, Prem Lobo⁶, Dantong Liu⁷, Jim M. Haywood⁴, Hugh Coe¹, James D. Allan^{1,2,*}

¹Department of Earth and Environmental Sciences, University of Manchester, UK.

²National Centre for Atmospheric Science, University of Manchester, Manchester, UK.

³Observation Based Research, Met Office, Exeter, UK

⁴College for Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, UK

⁵School of Chemistry, University of Bristol, Bristol, UK

⁶Metrology Research Centre, National Research Council Canada, Ottawa, Canada.

⁷Department of Atmospheric Sciences, School of Earth Sciences, Zhejiang University, Hangzhou, Zhejiang, China.

Currently at Qatar Environment and Energy Research Institute (QEERI), Hamad Bin Khalifa University (HBKU), Doha, Qatar.

*Correspondence to: dawei.hu@manchester.ac.uk; james.allan@manchester.ac.uk




Fig. S1. TEM images of BC particles emitted from: (a) diesel engine (hot engine condition) and (b) flame burner (condition 1).