

Hemispheric contrasts in ice formation in stratiform mixed-phase clouds: Disentangling the role of aerosol and dynamics with ground-based remote sensing

Martin Radenz¹, Johannes Bühl¹, Patric Seifert¹, Holger Baars¹, Ronny Engelmann¹, Boris Barja González², Rodanthy-Elisabeth Mamouri^{3,4}, Félix Zamorano², and Albert Ansmann¹

¹Leibniz Institute for Tropospheric Research (TROPOS), Leipzig, Germany

²Atmospheric Research Laboratory, University of Magallanes, Punta Arenas, Chile

³Department of Civil Engineering and Geomatics, Cyprus University of Technology, Limassol, Cyprus

⁴ERATOSTHENES Centre of Excellence, Limassol, Cyprus

Correspondence: Martin Radenz (radenz@tropos.de)

Abstract.

Multi-year ground-based remote-sensing datasets were acquired with the Leipzig Aerosol and Cloud Remote Observations System (LACROS) at three sites. A highly polluted central European site (Leipzig, Germany), a polluted and strongly dust-influenced eastern Mediterranean site (Limassol, Cyprus), and a clean marine site in the southern mid-latitudes (Punta Arenas, Chile) are used to contrast ice formation in shallow stratiform liquid clouds. These unique, long-term datasets in key regions of aerosol-cloud interaction provide a deeper insight into cloud microphysics. The influence of temperature, aerosol load, boundary-layer coupling, and gravity wave motion on ice formation is investigated. With respect to previous studies of regional contrasts in the properties of mixed-phase clouds our study contributes the following new aspects: (1) Sampling aerosol optical parameters as a function of temperature, the average backscatter coefficient at supercooled conditions is within a factor of 3 at all three sites. (2) Ice formation was found to be more frequent for cloud layers with cloud top temperatures above -15°C than indicated by prior lidar-only studies at all sites. A virtual lidar-detection threshold of IWC needs to be considered in order to bring radar-lidar-based studies in agreement with lidar-only studies. (3) At similar temperatures, cloud layers which are coupled to the aerosol-laden boundary layer show more intense ice formation than de-coupled clouds. (4) Liquid layers formed by gravity waves were found to bias the phase occurrence statistics below -15°C . By applying a novel gravity wave detection approach using vertical velocity observations within the liquid-dominated cloud top, wave clouds can be classified and excluded from the statistics. After considering boundary layer and gravity-wave influences, Punta Arenas shows lower fractions of ice containing clouds by 0.1 to 0.4 absolute difference at temperatures between -24 and -8°C . These differences are potentially caused by the contrast in the INP reservoir between the different sites.

1 Introduction

Clouds and aerosol are inseparably linked via complex pathways of interaction whose outcome manifests in the macroscopic properties of precipitation and radiation fields. In the first place, aerosol particles are required as cloud condensation nuclei

(CCN) on which cloud droplets can form. On the other hand, primary ice formation in the heterogeneous freezing temperature range from 0 to about -40°C requires ice nucleating particles (INP) to be present in the aerosol reservoir. The ways in which aerosol and cloud particles interact are controlled by the dynamics and thermodynamics of the atmospheric environment, with 25 temperature being the most important driver. Thermodynamic processes are considered to dominate the cloud microphysical properties over aerosol-related processes because the thermodynamics control the amount of water vapor that is available for being transferred to either the liquid or the ice phase. This dominance makes it difficult to isolate aerosol-related effects in observations of cloud properties.

Despite the dominance of dynamics and thermodynamics, observations as well as aerosol-permitting model studies suggest 30 a considerable influence of the aerosol conditions on the properties and evolution of clouds and precipitation (Seifert et al., 2012; Possner et al., 2017; Solomon et al., 2018; Zhang et al., 2018). Solomon et al. (2018) used high-resolution modelling of Arctic mixed-phase clouds to show that perturbations in the INP concentration dominate over changes in the CCN concentrations. Cloud chamber studies suggest that holding CCN constant, the ratio of ice to liquid water content in the steady state is predominantly controlled by INP concentrations (Desai et al., 2019).

35 There is also a distinct spatio-temporal variability of the performance of weather and climate model simulations, which is attributed to the insufficient representation of aerosol-cloud-dynamics interaction in the models (Fan et al., 2016; Seinfeld et al., 2016). For instance, a less accurate treatment of the radiative balance in the southern-hemispheric mid-latitudes compared to their northern-hemispheric counterparts was found (Trenberth and Fasullo, 2010; Grise et al., 2015). The atmosphere of the southern hemisphere's mid-latitudes is a unique component of the Earth's climate system. It's the stormiest (e.g., Young, 1999) 40 and one of the cloudiest places on Earth (Haynes et al., 2011; Naud et al., 2014), but process understanding of clouds in that region is still limited. The reported biases in the solar radiation budget are attributed to shallow supercooled liquid-topped clouds, which are insufficiently represented by current models (Bodas-Salcedo et al., 2014; Kay et al., 2016; Bodas-Salcedo et al., 2016; Kuma et al., 2020). With the frequency of liquid-cloud tops too low, less short-wave radiation is reflected. Hence, 45 the simulations show a too strong heating at the surface, leading to a bias in ocean heat uptake (Franklin et al., 2013; Hyder et al., 2018). There is an ongoing controversy about the reasons for the observed differences and prevailing model deficiencies, but indications are given that a combination of hemispheric contrasts in aerosol load and atmospheric dynamics plays a role.

Different causes for regional contrasts in the abundance of supercooled liquid cloud layers are proposed. On the one hand, 50 reason for the excess of supercooled liquid water in southern-hemispheric cloud systems could be the inhibition of ice formation caused by the lack of ice nucleating particles in the predominantly pristine environment of the Southern Ocean (Hamilton et al., 2014), where terrestrial sources, which are frequently considered a good source for INP, are rare or far apart (Vergara-Temprado et al., 2017). In the heterogeneous freezing regime, suitable aerosol particles are a prerequisite for the formation of ice. Missing ice formation as a sink for cloud water, the liquid phase may be sustained for long periods of time. On the other hand, dynamical 55 processes can lead to an enhancement of supercooled liquid water. Korolev (2007) showed that depending on the number and size of ice crystals, a threshold vertical velocity can be found, which allows for sufficient supersaturation to grow the ice as well as the liquid phase. Gravity waves have been suggested to play a role in the phase partitioning of Southern Ocean clouds (Alexander et al., 2017; Silber et al., 2020). Due to the orographic effects and the strong westerlies, gravity waves are a

general feature in the vicinity of all landmasses in the middle and high latitudes of the southern hemisphere (Sato et al., 2012; Alexander et al., 2016). It is however also noteworthy that stronger turbulence increases the amount of ice formed in stratiform cloud layers (Bühl et al., 2019). Indications are thus given that it is necessary to also consider turbulence in studies of ice formation.

The undetermined contributions of dynamical and aerosol effects on the observed excess of supercooled liquid in the southern hemisphere requires dedicated contrasting studies. In numerous previous studies, liquid-topped supercooled stratiform cloud layers have been proven to be suitable natural laboratories for the investigation of the relationships between aerosol properties, thermodynamics, and microphysical properties of clouds in the heterogeneous freezing regime. Temperature at which the ice formation occurs, needs to be strongly constrained, because the concentration of efficient INP increases rapidly with decreasing temperature (e.g., Kanji et al., 2017). Accordingly, the amount of ice formed also increases orders of magnitude for lower temperatures (Bühl et al., 2016). The lowest temperature in such clouds occurs on top of the liquid-dominated layer, hence the cloud top temperature (CTT) can be used to constrain the ice formation temperature. Turbulence is usually confined to the liquid-dominated cloud top (Westbrook and Illingworth, 2013; de Boer et al., 2009) and due to the limited thickness of this layer, secondary ice formation or ice multiplication are strongly constrained (Fukuta and Takahashi, 1999; Myagkov et al., 2016). Contrasting microphysical properties observed in pristine, clean regions with observations from areas with higher aerosol load allows us to advance understanding of the impact of different aerosol loads (Choi et al., 2010; Kanitz et al., 2011; Seifert et al., 2015; Tan et al., 2014). Recent studies based on the A-Train satellite constellation suggest systematically lower ice amounts in the southern mid-latitudes (Zhang et al., 2018) and a strong susceptibility to dust load (Villanueva et al., 2020). Supercooled liquid clouds are frequent over the Southern Ocean (Huang et al., 2015; Hu et al., 2010) and studies by Kanitz et al. (2011) and Choi et al. (2010) showed that—at similar temperatures—ice is formed less frequently by liquid layers in the southern hemisphere mid-latitudes, than in the northern hemisphere. The study of Kanitz et al. (2011) first used a ground-based lidar at Punta Arenas (53.1°S 70.9°W, Chile) to assess the thermodynamic phase of stratiform mixed phase clouds above the Southern Hemisphere mid-latitudes. Major caveats of this study are the limited duration of the observations during austral summer and the limitations of the lidar-only setup. Alexander and Protat (2018), using ground-based lidar and A-Train observations from Cape Grim (40.7°S 144.7°E, Australia) confirm the basic findings also for the eastern parts of the Southern Ocean. Their study, similarly to McErlich et al. (2021), emphasizes the problems in A-Train derived datasets detecting shallow clouds in the lowermost part of the atmosphere.

Recent activities include shipborne, land-based and aircraft campaigns targeting aerosols and clouds above the Southern Ocean between Australia and Antarctica (60 to 160°E). An overview is provided by McFarquhar et al. (2021). In terms of stratiform clouds, the large abundance of supercooled liquid water, occasionally down to -30°C , was confirmed. But, apart from the year-long lidar/radar dataset at Macquarie Island (54.6°S 158.9°E, Australia) the cloud observations focused on austral summer.

Using a shipborne dataset, Mace and Protat (2018) also found frequent liquid-dominated clouds with low radar reflectivities and one-third of the liquid layers only observed with lidar. Comparing the observations with a Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) dataset from Hu et al. (2010), they found an overestimation of supercooled liquid

in the satellite dataset, especially strong at temperatures above -15°C . In a follow-up study, Mace et al. (2020) refined the CALIPSO classification scheme, leading to more frequent detections of the mixed phase, especially during wintertime and in the lower latitudes of the Southern Ocean. However, no CTT-resolved phase occurrence statistics is presented. Liquid layers in deeper clouds, observed during another shipborne campaign (McFarquhar et al., 2021; Alexander et al., 2021), could only be reproduced in regional model simulations, when INP parametrization was tuned to lower concentrations (Vignon et al., 2021). Zaremba et al. (2020) investigated airborne active remote-sensing observations of Southern Ocean clouds south of Tasmania. They also found widespread liquid cloud tops at temperatures down to -30°C . By investigating the ground-based remote-sensing dataset assembled at McMurdo (77.8°S 166.7°E , Antarctica), Silber et al. (2018) found frequent long-lived liquid-topped clouds, also below -30°C .

Yet, a statistical analysis of the relationship between both, aerosol conditions, cloud vertical dynamics, and the phase partitioning in stratiform cloud layers of the southern-hemisphere mid-latitudes based on long-term observations was not established. One reason is that, despite increased activity in the recent past, ground-based remote-sensing observations of clouds and aerosol are still sparsely distributed in the Southern Ocean and at the coast of Antarctica.

The goal of this study is to analyze long-term ground-based remote-sensing observations of aerosol properties, cloud micro-physics, and atmospheric dynamics from three sites with strongly contrasting aerosol conditions in order to attribute ice formation in the heterogeneous freezing regime to atmospheric dynamics and aerosol conditions. For that contrasting approach, we utilized the recent campaigns of the Leipzig Aerosol and Cloud Remote Observations System (LACROS) at Leipzig (51.4°N 12.4°E , Germany), Limassol (34.7°N 33.0°E , Cyprus), and Punta Arenas (53.1°S 70.9°W , Chile), which provide datasets that cover the aerosol conditions of a continental northern hemispheric background site, a hot-spot of mineral dust, and the marine-dominated pristine Southern Ocean, respectively. Hence, these datasets collected with a single set of ground-based remote-sensing instrumentation provide an ideal basis for contrasting studies. The broad variety of instruments covers the decisive properties of aerosols, dynamics, clouds, and precipitation for a more comprehensive picture of aerosol-cloud interaction. The observations at Punta Arenas provide the first multi-year dataset of synergistic ground-based remote sensing observations in the western half of the Southern Ocean and allows us to contextualize prior findings.

The paper is structured as follows: The instruments and field campaigns are described first (Sec. 2.1), followed by the retrievals of aerosol properties (Sec. 2.2), the automated cloud selection algorithm (Sec. 2.4), and the characterization of vertical velocity (Sec. 2.5). The results are presented in Sec. 3, including the lidar-derived average profiles of aerosol optical properties (Sec. 3.1), and frequency of ice-formation statistics with special emphasis on instrument sensitivity (Sec. 3.2.1). Boundary layer coupling and vertical dynamics are discussed in Sec. 3.2.2 and Sec. 3.2.3, respectively. The amount and efficiency of ice production is assessed in Sec. 3.2.4. The study concludes with a discussion of the contrast identified in aerosol load and stratiform cloud properties (Sec. 4) followed by a summary and outlook (Sec. 5).

Table 1. Specifications of the LACROS instruments and measured quantities used in this study.

Instrument (Reference)	Frequency ν Wavelength λ	Measured quantity	Temporal resolution	Vertical range	Vertical resolution
Doppler cloud radar METEK MIRA-35 (Görsdorf et al., 2015)	$\nu = 35 \text{ GHz}$	Radar reflectivity factor	3.5s	150 – 13000 m	30 m
		Vertical velocity	3.5s	150 – 13000 m	30 m
		Linear depolarization ratio	3.5s	150 – 13000 m	30 m
Raman-Polarization Lidar Polly ^{XT} (Engelmann et al., 2016)	$\lambda = 355, 532, 1064 \text{ nm}$	Attenuated backscatter coeff.	30s	100 – 15000 m	7.5 m
	$\lambda = 355, 532 \text{ nm}$	Raman backscatter signal	1h	300 – 5000 m	$\sim 50 \text{ m}$
	$\lambda = 355, 532 \text{ nm}$	Linear depolarization ratio	30s	100 – 15000 m	7.5 m
Microwave radiometer RPG HATPRO-G2 (Rose et al., 2005)	$\nu = 22.24 – 31.4 \text{ GHz}$	Brightness temperatures	1s	column integral	
	$\nu = 51.0 – 58.0 \text{ GHz}$	Brightness temperatures	1s	column integral	
Doppler Lidar HALO Photonics Streamline XR (Pearson et al., 2009)	$\lambda = 1.5 \mu\text{m}$	Attenuated backscatter coeff.	2s	48 – 12000 m	48 m
		Vertical velocity	2s	48 – 12000 m	48 m
Ceilometer Jenoptik CHM15kx	$\lambda = 1064 \text{ nm}$	Attenuated backscatter coeff.	30s	15 – 15300 m	15 m
Optical disdrometer Ott Parsivel ² (Löffler-Mang and Joss, 2000)	$\lambda = 650 \text{ nm}$	Hydrometeor size distribution	30s	4 m	-
Sun photometer Cimel CE318-T (Barreto et al., 2016)	$\lambda = 340 – 1064 \text{ nm}$	Aerosol optical thickness	variable	column integral	

2 Data and Methods

This section introduces the datasets and methods used in the remainder of this study. Starting with the campaigns and instrumentation (Sec. 2.1), followed by a short description of the retrievals used (Sec. 2.2 and 2.3) and finally the methods for automated selection of shallow stratiform clouds (Sec. 2.4) and characterization of vertical motion (Sec. 2.5).

2.1 Datasets from Leipzig, Limassol, and Punta Arenas

Basis of the observational datasets is the Leipzig Aerosol and Cloud Remote Observations System (LACROS), the mobile ground-based remote-sensing supersite of the Leibniz Institute for Tropospheric Research (TROPOS), Leipzig, Germany. The instrumentation used for the synergistic approaches applied in this study comprises a MIRA-35 35 GHz scanning cloud radar, a Polly^{XT} multi-wavelength Raman polarization lidar, a Streamline XR 1.5 μm scanning Doppler lidar, a HATPRO 14-channel microwave radiometer, a 1064 nm ceilometer, an optical disdrometer, and radiation sensors. Main properties of the sensors are summarized in Table 1.

LACROS was established in 2011 and was first introduced by Bühl et al. (2013b). After the initial setup phase, the basic set of instrumentation of LACROS has been kept unchanged since the year 2014. Hence, this comparative study uses data from the




Figure 1. Overview on instrument availability and campaign durations of the LACROS deployments covered in this study. Characteristics of the three different Polly^{XT} instruments used at Leipzig and named IFT, Arielle, and LACROS are described in Engelmann et al. (2016).

6 year period since then. During that period, LACROS was, besides several 4-8 weeks short-term deployments at other sites, stationed at Leipzig, Limassol, and Punta Arenas.

Fig. 1 provides an overview on the campaigns and instrument availability. From October 2016 to March 2018 LACROS was located at Limassol for the Cyprus Clouds, Aerosol, and Rain Experiment (CyCARE). In the context of the present study, this 140 location in the eastern Mediterranean in the middle of the dust belt serves as a reference for a highly aerosol-laden environment. First results of the CyCARE campaign are described by Ansmann et al. (2019). For the second long term campaign, LACROS was deployed at Punta Arenas for the two-year Dynamics, Aerosol, Clouds, And Precipitation Observations in the Pristine Environment of the Southern Ocean (DACPPO-PESO) field campaign since November 2018. Located at 53.1°S, Punta Arenas is in the midst of the Southern Ocean, farther south than any other continental land mass or major island apart Antarctica. Under 145 prevailing westerly flow, the next land mass upwind is more than 8000 km away and significantly to the north. Hence, the free tropospheric aerosol is predominantly of marine origin with low concentrations overall. First results of the DACPO-PESO campaign regarding events of significant aerosol load (Ohneiser et al., 2020; Floutsi et al., 2021), liquid cloud microphysics (Jimenez et al., 2020) and integrated water vapor observations (Bromwich et al., 2020) are already published. Observations at Leipzig were performed before October 2016 and between the deployments to Limassol and Punta Arenas. Due to the location 150 in central Europe, the Leipzig dataset serves as a reference for continental background in the northern hemisphere.

Data coverage for each of the sites is summarized in Table 2. Gaps in the observations were usually caused by periods of short-term deployments at other sites, failure of primary instruments, or power cuts (see also Fig. 1). Coverage with the core instrumentation is generally well above 80%, only exception being Doppler lidar and Polly^{XT} at Leipzig.

2.2 Lidar-based aerosol statistics

155 As a proxy for the aerosol conditions at the three measurement sites, we used vertically resolved observations from the Polly^{XT} lidar system (Althausen et al., 2009; Engelmann et al., 2016). Quantities of interest are the aerosol backscatter coefficient, the

Table 2. Overview on Location, data availability, climate, aerosol load, and related studies for the LACROS datasets used.

Location	Leipzig, Germany	Limassol, Cyprus	Punta Arenas, Chile
	51.4°N 12.4°E	34.7°N 33.0°E	53.1°S 70.9°W
Station altitude	125 m asl	11 m asl	9 m asl
Campaign name		CyCARE	DACAPO-PESO
Duration	976 d	524 d	765 d
Cloudnet data	771 d	520 d	674 d
Doppler lidar data	513 d	523 d	717 d
Polly ^{XT} data	633 d	460 d	702 d
Climate	Northern mid-latitudes	Northern subtropics	Southern mid-latitudes
Typical aerosol load	Continental background, occasionally dust	Dust, marine, continental	Marine, occasionally continental
Related studies	Ansmann (2005) Seifert et al. (2010) Bühl et al. (2013a) Bühl et al. (2016)	Ansmann et al. (2019)	Ohneiser et al. (2020) Bromwich et al. (2020) Jimenez et al. (2020) Floutsi et al. (2021)

extinction coefficient, the particle linear depolarization ratio, and the cloud-relevant concentration of INP. Their retrieval is explained below.

Basis for the statistical analysis are profiles of particle backscatter coefficient β_p at 532 nm wavelength. The profiles are 160 computed with the Klett method (Fernald, 1984) by the PollyNET retrieval whenever atmospheric conditions are suitable (Baars et al., 2016, 2017; Yin and Baars, 2021). The PollyNET retrieval chain also ensures a homogenized analysis of the data from the three different Polly^{XT} instruments, which were utilized in the frame of this study (see Fig. 1 and Engelmann et al., 2016; Baars et al., 2016).

Profiles of the particle linear depolarization ratio (hereafter referred to as particle depolarization ratio) are only calculated 165 when the ratio of molecular backscatter coefficient to β_p is below a value of 18. Additionally, any particle depolarization ratios larger than 0.7 are masked, as they are indications for noise artifacts in the cross-polarized signal component. All profiles are then filtered with the co-located Cloudnet target classification (see Section 2.3) to exclude clouds, especially optically thin ice clouds that are only clearly classified by the cloud radar. Finally, a manual screening excluded fragments of thin liquid clouds, 170 which would otherwise artificially increase β_p . For the averages, the optical data of each retrieved profile is binned to vertical intervals of 200 m or 3 K.

The derived average optical properties can be used to estimate aerosol microphysical properties, such as concentrations of ice nucleating particles (INP). This is an important step in order to evaluate the datasets of the three sites with respect to 175 contrasts in the potential contribution of aerosol effects on heterogeneous ice-formation efficiency. Conversion from optical properties as observed by lidar to microphysical aerosol properties is based on the parametrizations described by Mamouri and Ansmann (2016). By means of this approach, the lidar-measured aerosol extinction coefficient is converted to the number and surface concentration N_{500} and S_{500} of aerosol particles larger than 500 nm in diameter. These quantities are applied

in available in-situ-based parametrizations for the retrieval of INP concentrations. Prerequisite for the retrieval is a correct aerosol typing, as different types of particles differ by orders of magnitude in their ice forming efficiency. In order to do so, the average backscatter profile is separated into the categories marine, continental, and mineral dust, based on air mass source
180 (see Appendix B and Radenz et al., 2021) and particle depolarization ratio (one-step POLIPHON; Mamouri and Ansmann, 2017). The average extinction is calculated from the profiles of β_p by assuming a typical lidar ratio of 20sr for marine, 50sr for continental, and 45sr for dust aerosol (Müller et al., 2007; Baars et al., 2017; Bohlmann et al., 2018). In a next step, the extinction coefficient is converted to N_{500} and S_{500} , using sun-photometer-based conversion factors (Mamouri and Ansmann, 2016). Then, the above mentioned INP parametrizations are applied for each aerosol class, such as DeMott et al. (2015) for
185 mineral dust, DeMott et al. (2010) for continental aerosol and McCluskey et al. (2018b, a) for marine aerosol.

2.3 Cloudnet processing

Synergies between lidar, cloud radar, microwave radiometer, and meteorological data are utilized for determination of cloud
190 macro- and microphysical properties and as basis for the stratiform cloud identification scheme. State-of-the-art routines for achieving this synergy are comprised in the Cloudnet retrieval (Illingworth et al., 2007). Cloudnet re-grids the observations to a common resolution (30s and 31.18m, determined by the vertical resolution of MIRA-35 with a pulse length of 208ns) and provides products, such as a target classification and microphysical cloud properties. Regridding is a crucial step, as each of the instruments has different temporal and vertical resolution (Tab. 1). Regular scans of the MIRA-35 cloud radar (hourly) and the Streamline Doppler lidar (twice per hour) are not used within the Cloudnet processing scheme. Profiles of temperature, pressure, and humidity are obtained from the ECMWF's IFS analysis for Punta Arenas and GDAS analysis for Leipzig and
195 Limassol. Liquid water path (LWP) and integrated water vapor (IWV) are retrieved from brightness temperature observations of the microwave radiometer in two frequency bands from 22.24 to 31.4 GHz and 51.0 to 58.0 GHz (7 channels each; Crewell and Löhnert, 2007). The statistical retrieval is based on long-term radiosonde observations (Leipzig and Limassol) and high-resolution reanalysis data (Punta Arenas). Attenuated backscatter of the ceilometer is regularly cross-calibrated with Polly^{XT} using the calibrated attenuated backscatter of PollyNET. Usually, the ceilometer data is used in the synergistic retrieval, as
200 the dataset is more robust and less prone to interruptions. Rare gaps in the observations are filled with the Polly^{XT} attenuated backscatter at 1064 nm.

2.4 Automated cloud selection and characterization

While the Cloudnet algorithm provides a pixel-by-pixel classification of cloud phase and microphysical properties, information
205 on temporal coherence and cloud evolution is not readily available. Hence, an automatic reproducible filtering algorithm for the selection of targeted stratiform, supercooled cloud systems is necessary. Based on the data cube LARDA³ (Bühl et al., 2021), the approach of Bühl et al. (2016) is implemented into an automated selection algorithm.

An example of the Cloudnet processing of measurement data and the application of the cloud selection scheme is shown in Fig. 2. Starting with a profile of the Cloudnet target classification mask (Hogan and O'Connor, 2004), consecutive pixels classified as cloud pixels (liquid droplets, ice, ice and supercooled droplets) are grouped together and defined as features. In

210 case similar types of hydrometeors were observed in matching heights, single features in neighboring timesteps are connected to coherent cloud cases. For the analysis, the cloud cases are filtered for shallow stratiform clouds, which are liquid topped and either have an ice virga or not (rectangles in Fig. 2d). An overview of the microphysical parameters sampled for each cloud case is provided in Table 3. It is assumed that if ice is formed in an liquid layer it also sediments out of the cloud. This is required, as the signal in the top layer is dominated by return from liquid droplets and Cloudnet provides no reliable 215 mixed-phase classification there.

To pinpoint potential effects of aerosol load, thermodynamic, and dynamic drivers of ice formation have to be constrained. This is especially important, as the sites are situated in different climate zones. We presume for our study that thin stratiform clouds serve as a natural laboratory with only a limited number of microphysical processes being possible.




Figure 2. Stratiform liquid-topped clouds as observed with LACROS. (a) Cloud radar reflectivity overlaid with the temperature, (b) Lidar attenuated backscatter, (c) Doppler lidar vertical velocity, and (d) Cloudnet target categorization (d) on the 28 November 2018, 00:20 to 09:00 UTC. The temperature information in (a) is obtained from ECMWF IFS model data. Red rectangles in (d) indicate the automatically detected liquid-topped shallow stratiform clouds. The inset in (c) shows the autocorrelation function of the vertical velocities in the liquid-dominated layer of each cloud. The height of the velocity sampling is indicated with dashed lines. Further details are provided in Sec. 2.5.

The cloud cases are filtered for a length of more than 20 minutes and smooth cloud top heights (standard deviation $< 150\text{ m}$) 220 to exclude convective clouds. Laboratory studies of Fukuta and Takahashi (1999) and the cloud radar observations by Myagkov et al. (2016) indicate that the average thickness of the liquid dominated cloud top layer has to be less than 350 m in order to restrict the dataset to the regime of pristine ice formation and avoid strong effects of secondary ice formation. Seeding by ice clouds above is avoided by excluding clouds with ice pixel above the liquid layer. Only cloud cases with top temperatures between -38 and 0°C are considered for the statistics. For the phase occurrence frequency statistics, a cloud is classified as ice

Table 3. Parameters sampled from the automatically identified cloud cases. A more detailed description of the automated spatio-temporal selection method is provided in Bühl et al. (2016). The IWC and ice extinction is retrieved with the parametrization of Hogan et al. (2006).

Parameter	Time series	Description
Cloud top temperature	yes	Temperature of the topmost liquid pixel
Cloud top height	yes	Geometrical height of the topmost liquid pixel
Liquid layer thickness	yes	Geometrical depth of the liquid-dominated cloud top layer
Phase classification	no	Ice pixel observed below liquid-dominated layer > 5% of duration
In-cloud vertical velocity	yes	Doppler lidar vertical velocity in the middle of the cloud top layer
LWC	yes	LWC of the liquid-dominated layer
Reflectivity virga	yes	Radar reflectivity 180 m below the base of the liquid-dominated layer
IWC in virga	yes	Reflectivity-derived IWC 180 m below the base of the liquid-dominated layer
Extinction in virga	yes	Optical extinction of ice at 180 m below the base of the liquid-dominated layer
Ice-to-liquid content ratio	yes	Ratio of IWC virga to LWC in the liquid-dominated layer

225 producing, if ice pixel were observed 180 m below the liquid-dominated layer within at least 5% of the duration. On the other hand, cloud cases are classified as liquid-only, if ice pixel are observed to sediment out of the liquid-dominated layer less than 5% of the time. This threshold is simpler, than the decision tree used by Bühl et al. (2013a), but provides similar results. The resulting fraction is weighted by the length of the cloud. Similarly to Bühl et al. (2016), the properties of the ice in the virga are taken 180 m below the liquid-dominated cloud top layer to avoid uncertainties in the liquid base estimate and sublimation
230 within the virga. The ice water content (IWC) is derived from the radar reflectivity and the temperature using the Hogan et al. (2006) retrieval. With this automated cloud selection scheme multi-year datasets can be analyzed using objective criteria, while yielding statistics similar to manual cloud selection (e.g., Seifert et al., 2010; Kanitz et al., 2011; Seifert et al., 2015).

2.5 Gravity wave detection

235 In order to pinpoint effects of aerosol and dynamics on the phase partitioning in the stratiform cloud dataset, an approach is required to assign vertical motion regimes to each cloud case. Here we focus on the temporal structure of vertical velocity, to constrain the dynamics forcing on a cloud. Usually, shallow clouds are characterized by a fully developed turbulence in the liquid-dominated cloud top (Bühl et al., 2019), where the vertical motion is driven by cloud top cooling (e.g., Shao et al., 1997; Fang et al., 2014; Simmel et al., 2015). In the turbulent layer at cloud top, up- and downdrafts alternate at horizontal scales in the order of 100 m or less.

240 However, orographic gravity waves can trigger vertical motion and associated cloud formation, as well. Microphysical processes in these wave clouds are governed by large-scale dynamics, where vigorous up- and downdrafts may appear stationary. Due to this dynamics, the mixed-phase and the ice phase are horizontally separated, with the liquid drops predominantly in the ascending branch and the ice particles in the descending branch (Heymsfield and Miloshevich, 1993; Baker and Lawson, 2006). The properties of the horizontal wind field determines the regions of the up- and downdraft in such orographic clouds.

245 Observing these clouds by stationary ground-based remote sensing might thus not sample the full horizontal extent of the cloud, which causes a misclassification of the cloud case in terms of liquid-only and ice-producing. An illustrative example of such a wave cloud is shown in appendix A

250 Also, the flow is highly laminar, opposed to the confined, fully developed turbulence found in layered mixed-phase clouds. Cloud microphysics in wave conditions cannot directly be compared to layered clouds. For clouds in the heterogeneous freezing regime, Cotton and Field (2002) found that only rapid evaporation freezing once the downdraft commences could explain their observations, where evaporation freezing can be better characterized as inside-out contact freezing of shrinking particles (Durant, 2005). A more recent study by Field et al. (2012) found that condensation and immersion freezing are needed together with deposition and evaporation freezing to explain their aircraft observations of ice formation in wave clouds.

255 On the other hand, the frequent occurrence of atmospheric gravity waves in a specific region might increase the frequency of thermodynamic conditions that favor the presence of a sustained liquid phase. As Korolev (2007) demonstrate, long-lasting steady updrafts are required in order to make the liquid phase dominating over the ice phase. Probing an observational dataset for the presence of long-lasting updrafts could therefore provide a hint on the role of atmospheric gravity waves in the occurrence of enhanced concentration of supercooled liquid.

260 Turbulence properties of the liquid layer in the clouds subject to our study can be derived from the vertical velocity observation by Doppler lidar (Bühl et al., 2019). The small size of droplets in the liquid-dominated cloud-top layer and their negligible terminal velocity makes them tracers of air motion. From the Doppler lidar observations, the vertical velocity is sampled at the average height of the center of the layer identified as liquid-containing in the Cloudnet classification. The height of this sampling is indicated as dashed lines in Fig. 2c and Fig. A1d. The temporal resolution of the resulting time series is 2s, i.e., equal to the Doppler lidar raw data. This time-series is then used to get insights into the turbulent properties of the cloud-top 265 layer. To separate both regimes in the large dataset, an characteristic parameter is needed. The autocorrelation function was found to be a pragmatic choice for such a parameter.

The autocorrelation function $\Psi(\tau)$ for a time series of vertical velocities v_t is defined as

$$\Psi(\tau) = \sum_t v_t v_{t+\tau} \quad (1)$$

270 with the temporal shift τ and the vertical velocity v at time t . To compare different cloud cases, the autocorrelation function $\Psi(\tau)$ is normalized with $\Psi(0)$. The temporal shift τ from the observations is converted into a horizontal shift or autocorrelation length l with the Cloudnet model-based horizontal wind velocity v_{hor} :

$$\Psi(l) = \Psi(\tau) v_{\text{hor}} \quad (2)$$

275 Similarly the vertical-velocity spectral power density is calculated by a Fast Fourier Transform of the vertical-velocity time series. High autocorrelation coefficients for large shifts and low power density are indications for wave-driven, low-turbulent flow. The inset in Fig. 2c shows the autocorrelation function for each of the identified cloud cases. All of them are weakly affected by gravity waves, but small-scale turbulence dominates. In contrast, the wave cloud in Fig. A1d shows high autocorrelation coefficients for longer shifts. As a characteristic value of the autocorrelation function, the shift at which the

coefficient drops below 0.8 was chosen after visually inspecting the whole dataset. When the ice-formation frequency statistics is investigated for an influence of gravity waves (Sec. 3.2.3), the shift threshold is reduced step by step. As the autocorrelation 280 function is generally decreasing, a lower characteristic value than 0.8 would result in larger shift thresholds. For shifts larger than about 500 m, random fluctuations appear for clouds with a rapid drop in autocorrelation coefficients. Thus, 0.8 is a robust choice for the characteristic value.

3 Results

In this section, the thermodynamic phase partitioning and quantitative ice mass production in ice forming shallow cloud layers 285 are presented and compared between all measurement sites under study. First, the average profiles of aerosol optical and microphysical properties are presented. Then, instrumental detection thresholds, boundary layer effects and gravity wave activity are all analyzed as potential influencing factors on the retrieved ice-formation characteristics.

3.1 Aerosol conditions at Leipzig, Limassol, and Punta Arenas

To provide a general insight into the aerosol conditions at the three sites, the average optical and microphysical aerosol properties 290 as derived from the lidar observations (Sec. 2.2) are shown in Fig. 3 and Fig. 4. The impact of aerosol on clouds is controlled more strongly by temperature than geometrical height, hence the averages are also calculated with temperature as a vertical coordinate.

3.1.1 Optical properties

The average aerosol backscatter coefficient β_p at 532 nm and the particle depolarization ratio derived from the Polly^{XT} 295 observations with the Klett method (see Sec. 2.2) are investigated in this section. The evaluation of the air mass sources is covered more quantitatively in the next section and Appendix B.

The central European site of Leipzig is characterized by predominantly continental aerosol mixed with anthropogenic pollution (Baars et al., 2016). Long-range transport of dust may occur periodically, especially during spring and autumn (Ansmann et al., 2003) as well as lofted smoke layers from wildfires (Haarig et al., 2018; Baars et al., 2021). Aerosol optical thickness 300 (AOT) is derived from sun-photometer observations. Mean AOT at 500 nm is 0.198 at TROPOS, Leipzig, between 2014 and 2018. When only the periods with co-located Polly^{XT} observations (used in this study) are considered, the AOT is 0.216. Mean β_p at 532 nm drops below $0.2 \text{ Mm}^{-1} \text{sr}^{-1}$ only above 4 km height, which corresponds to an extinction coefficient of 1.0 Mm^{-1} assuming continental aerosol conditions and a corresponding lidar ratio of 50 sr.

Limassol is characterized by a distinct dry season with no precipitation and very few clouds during the summer. Generally, 305 Limassol is frequently affected by aerosol transport from Africa, the Middle East and Europe with aerosol characteristics including dust (mineral and soil), marine (organics and sea salt) and anthropogenic pollution as well as mixtures of these (Nisantzi et al., 2015). Mean AOT at 500 nm is 0.176 during the whole observational period and 0.165 during the ‘cloudy season’ from October to May. In the following, the non-cloud season from June to September is excluded from the statistics.

The profile of mean backscatter is similar to the one at Leipzig within a factor of 1.5, whereas the median generally is higher
310 at Leipzig.

The aerosol load at Punta Arenas can be separated into two distinct layers with an aerosol-rich boundary layer and pristine conditions aloft. The free troposphere is dominated by marine aerosol from the Southern Ocean and was reported to show no changes compared to the pre-industrial conditions (Hamilton et al., 2014). Nevertheless, events of aerosol long-range transport also occur occasionally (Foth et al., 2019; Floutsi et al., 2021). The boundary layer is laden with a mixture of marine and
315 continental aerosol, as Punta Arenas is located 230km inland from the Pacific coast. Mean AOT at 500nm is 0.055 during the whole campaign, but dropping to 0.047, when excluding the period of long-range wildfire smoke transport in early 2020 (Ohneiser et al., 2020). Average boundary layer height is around 1.5km (Foth et al., 2019) with negligible β_p above 2.0km height (90% percentile of β_p at 532nm dropping below $0.2\text{Mm}^{-1}\text{sr}^{-1}$). Comparing the backscatter at Punta Arenas and Limassol, the 90% percentile at Punta Arenas is more than 30% below the mean of Limassol and Leipzig at similar heights
320 above ground. Both European sites show quite some variability, as well, with the 90% percentile β_p twice as large as the mean.

At temperatures above 5°C , β_p shows the strongest difference between the three sites (Fig. 3b), also explaining the larger AOT at Limassol and Leipzig. Between 0°C and -12°C , the mean β_p at Limassol and Punta Arenas are almost equal. This counterintuitive behavior can be explained by different temperature regimes. The -5°C isotherm at Punta Arenas varies between 1.1 and 3.3km height, whereas at Limassol it varies between 3.0 and 4.8km height and at Leipzig between 2.1 and
325 4.8km. At Punta Arenas, corresponding to the height of 2.0km, only very low values of β_p are observed at heights below a temperature of -10°C . The pronounced decrease of backscatter at Leipzig and Limassol is observed at slightly higher temperatures, which agrees with the, on average, higher boundary layer temperature there. Assuming typical lidar ratios of 50 (continental), 45 (dust), and 20sr (marine) at Leipzig, Limassol, and Punta Arenas, respectively, typical aerosol extinction coefficients can be estimated from the median β_p profile (Sec. 2.2). As shown in Fig. 3c, the extinction coefficients at Punta
330 Arenas are a factor 2-3 lower than over Limassol for temperatures below -10°C . This difference decreases to a factor of 1.5 for slightly supercooled conditions of above -10°C . Comparing Punta Arenas and Leipzig, the extinction is a factor of 3-4 higher at the latter site for all temperatures below freezing. This optical properties serve as a proxy for the background reservoir of aerosol particles that could act as cloud condensation nuclei and ice nucleating particles in the free troposphere.

The particle depolarization ratio provides hints on the aerosol particle types (Fig. 3d). At Leipzig, the depolarization ratio
335 is approximately 0.05 at all temperatures, typical for a continental aerosol with a slight contribution of mineral dust. The depolarization ratio at Limassol is bimodal, with one peak above 20°C , a second one at -14°C and a minimum at 4°C . The first peak can be ascribed to mineral dust in the boundary layer and originating from local sources. The second one at is caused by long-range transport of mineral dust. At Punta Arenas, a maximum at $+8^\circ\text{C}$ is caused by occasional events of dried sea salt aerosol at the top of the atmospheric boundary layer (Haarig et al., 2017; Bohlmann et al., 2018). Going to lower temperatures,
340 the depolarization ratio has a minimum of 0.01 at -20°C and a slight increase afterwards.




Figure 3. Profiles of average aerosol optical properties at 532 nm wavelength derived by the PollyNET retrieval using the Klett method. (a) particle backscatter coefficient over height and (b) temperature for Leipzig, Limassol, and Punta Arenas. (c) Extinction coefficient and (d) particle depolarization ratio for the same locations. For the extinction coefficient only the median values based on the typical lidar ratio at each site are shown. Variability caused by different aerosol mixtures (see Sec. 3.1.2) is denoted by shading.

3.1.2 Contrasts in INP concentrations

An important aspect for the discussion of microphysical contrasts in clouds over the three sites is how the differences in aerosol optical profiles are linked to contrasts in the INP load. An estimate of average INP concentrations covering the whole troposphere can be derived with the parametrizations described in Sec. 2.2. The decision for the aerosol typing required in the 345 INP retrieval is based on the particle depolarization ratio and air mass source estimates. The air mass source methodology is based on Radenz et al. (2021) and shown in Appendix B.

At Punta Arenas marine sources are by far the most frequent, contributing 90% to the residence time throughout the troposphere and peaking 95% at 2.7 km height. Only below 2.0 km height (or above -10°C) local terrestrial sources make up to 10% of the air mass. Above 5 km height (below -20°C) sparsely vegetated areas in Australia contribute up to 4% to the air 350 mass source. Mean particle depolarization ratios below 0.02 in the free troposphere exclude frequent presence of mineral dust (Fig. 3d). Contrarily, at Limassol air masses with a marine source contribute 50% to 80% to the mixture. But these air masses are not pristine marine, as the eastern Mediterranean is enclosed by strongly populated landmasses. Below 5 km height (above -16°C) continental Europe is second strongest source, responsible for up to 45% at 0.5 km height. In the upper troposphere, barren ground from the Sahara is the strongest terrestrial source with contributions around 15%. POLIPHON (see Sec. 2.2) 355 shows a peak at -12°C with mean dust fractions of 0.3 (90% percentile 0.87). Hence, the backscatter is divided into dust and non-dust according to the dust fraction. The non-dust portion is then split up into continental and marine, with the 40% contribution of the continent above -12°C and 20% below. The aerosol mixture at Leipzig is dominated by continental aerosol, with an average dust fraction of 0.1.




Figure 4. Average INP concentrations derived from optical properties for Leipzig (green), Limassol (orange), and Punta Arenas (blue). Shading shows the spread covered by different aerosol compositions, similar to Fig. 3c. Temperature is used as a vertical coordinate. The fraction of continental aerosol is abbreviated ‘conti.’. The used parametrizations are described in Sec. 2.2.

The derived INP concentrations are depicted in Fig. 4. Note that the INP concentration at a certain atmospheric temperature
 360 (as a proxy for height) is shown based on the optical properties, aerosol types, and the parameterization, not a freezing spectrum
 obtained from sampling a single air parcel and varying temperature.

At temperatures above -10°C average INP concentrations between 4×10^{-3} to $6 \times 10^{-2} \text{ L}^{-1}$ can be expected at all three locations. With decreasing temperatures, the concentration increases to $0.1 - 1 \text{ L}^{-1}$ at -25°C at the northern-hemispheric sites
 365 of Leipzig and Limassol. A strong increase of ice nucleating efficiency with decreasing temperature is counterbalanced by a
 decreasing aerosol concentration at lower temperatures, which are tied to greater heights. INP concentrations at Punta Arenas
 are strongly controlled by the fraction of continental aerosol in the free troposphere. Continental sources can contribute up
 to 1% between -11 and -24°C and up to 4% at lower temperatures. When only pristine marine aerosol is present, the INP
 concentration can be expected to be 3-4 orders of magnitude lower than at the two other sites. Equally low concentrations of INP
 were also found by McFarquhar et al. (2021) in the remote Southern Ocean south of Australia. As soon as very small fractions
 370 of continental aerosol are present, the INP concentration increases significantly. However, even if few percent of continental
 aerosol are assumed, throughout the heterogeneous freezing regime, INP concentrations remain a factor of 2-6 lower at Punta
 Arenas, compared to Leipzig and Limassol. Due to the absence of suitable remote-sensing or in-situ measurements, the actual
 contribution of continental aerosol to the free-tropospheric aerosol load over Punta Arenas can to date not be obtained. The
 range given in Fig. 4 is thus a solid estimate of the expectable range of possible INP concentrations. At temperatures above

375 -10°C , which refers to a height range that is frequently within the boundary layer, INP concentrations at Punta Arenas are high and within an order of magnitude to the concentrations at Leipzig and Limassol at the same temperature. Concentrations that are equally high as retrieved from the remote sensing observations above -10°C were also found in-situ at the up-wind hilltop station on Cerro Mirador in an altitude of 622m above sea level (Gong et al., in preparation).

3.2 Mixed-phase stratiform cloud properties

380 The automated stratiform cloud selection algorithm introduced in Sec. 2.4 is applied to the Cloudnet datasets of all three locations. An overview over identified clouds and temperature-resolved phase occurrence frequency is provided in Sec. 3.2.1. Afterwards, it is shown how boundary layer aerosol load (Sec. 3.2.2) and gravity waves affect the phase occurrence (Sec. 3.2.3). In Sec. 3.2.4 contrasts of ice content in the virga are briefly discussed.

3.2.1 Phase occurrence frequency and detection thresholds

385 Fig. 5 shows all the cloud cases that fulfill the selection criteria introduced in section 2.4. The time share of ice virga sedimenting out of the liquid-dominated top is given by the fraction of profiles for which ice was classified below the liquid layer. Values range from 0 (no ice virga observed at all) to 1 (ice virga observed all the time). At all three locations, the fraction of profiles (i.e., fraction of time) in which a virga originated from the liquid layer increases with decreasing temperature. Similarly, the ice-to-liquid content ratio (Bühl et al., 2016) increases for lower temperatures. Liquid only clouds (fraction of profiles where 390 no ice was produced from the liquid layer) were observed at Leipzig and Limassol down to -16°C . Whereas at Punta Arenas such clouds were observed even at temperatures as low as -38°C .




Figure 5. All stratiform clouds identified by the automated selection algorithm for (a) Leipzig, (b) Limassol, and (c) Punta Arenas by their cloud top temperature and the fraction of profiles, where ice is observed below the liquid-dominated cloud top. The 0.05 threshold for the classification is marked by a dashed line. The color gives the median ice-to-liquid content ratio. n gives the number of cloud cases in each dataset. Red squares in (c) denote the clouds also shown in Fig. 2.

First, the temperature-resolved fraction of occurrence of ice-forming clouds is depicted in Fig. 6. For the phase occurrence frequency, a cloud is classified as ice-producing, if an ice virga was observed at least during 5% of the duration of a cloud case (Sec. 2.4). Comparing the frequency of ice-containing clouds at different locations provides insights into differences of primary ice formation (Choi et al., 2010; Kanitz et al., 2011; Seifert et al., 2015; Tan et al., 2014; Zhang et al., 2018). Generally, clouds contain ice more frequently for decreasing temperature (Fig. 6, solid lines). While at Leipzig and Limassol nearly all clouds with CTTs below -16°C contained ice, at Punta Arenas a fraction of 0.4 ± 0.1 of shallow stratiform clouds at these temperatures were classified as liquid only. This behavior is discussed further in Sec. 3.2.3.

Compared to prior lidar-based studies (e.g., Kanitz et al., 2011; Choi et al., 2010; Seifert et al., 2010), the fraction of ice-containing clouds in the synergistic dataset is higher at temperatures above -10°C . At these temperatures, the amounts of ice produced and hence, the radar reflectivity and optical extinction are usually very low and stays undetected for lidar (Bühl et al., 2013a) and space-borne radars (Bühl et al., 2016). To quantify a lidar detection threshold in terms of optical extinction, the reflectivity-to-IWC and the reflectivity-to-extinction relationships by Hogan et al. (2006) are used. The only additional information needed for both parametrizations is the ambient temperature. Using these relationships, the response of the occurrence statistics to arbitrary extinction detection thresholds can be tested. As shown in Fig. 6 (dashed lines), for the lidar data used in the Cloudnet classification, an extinction threshold of 12 Mm^{-1} had to be applied in order to best match the lidar-only statistics from Punta Arenas and Leipzig presented by Kanitz et al. (2011).




Figure 6. Fraction of ice containing clouds over temperature for Leipzig, Limassol, and Punta Arenas. Total duration of the clouds in each bin is given by the numbers on top in hours. Dashed curves mark the occurrence frequency when using a lidar detection threshold of 12 Mm^{-1} .

3.2.2 Effect of boundary layer aerosol load on phase occurrence

As discussed in Sec. 3.1, the aerosol load at Punta Arenas is confined to the lowermost 2 km and the aerosol load at temperatures above -10°C is similar to Limassol. To check for possible impact of this boundary layer aerosol on the ice-formation

efficiency, the basic temperature-resolved phase occurrence frequency (Fig. 6) is split into two subsets, one containing cloud cases with bases below 2km height and one with cloud cases having bases above that threshold, in the following denoted as coupled and uncoupled clouds, respectively. The resulting ice-formation frequency is shown in Fig. 7. At any height, temperatures vary by more than 11°C (10% to 90% percentile), providing ample coverage for the height threshold. Generally, coupled 415 clouds show higher fractions of ice (absolute difference increases by 0.3 at all locations at -8°C). The temperature-resolved phase occurrence frequency for the coupled clouds show rapid increase in fraction of ice-containing clouds, reaching 1.0 at temperatures of only -15°C . The absolute difference between the fractions is less than 0.20, with the lowest fractions still being observed at Punta Arenas. Below -15°C , almost no clouds were observed at such low heights, especially at Limassol, hence no meaningful comparison can be done in this temperature interval.




Figure 7. Fraction of ice containing clouds over temperature for Leipzig, Limassol, and Punta Arenas, (a) below 2km and (b) above 2km. Total duration of the clouds in each bin is given by the numbers on top in hours.

420 Considering only uncoupled clouds (cloud base above 2km height, Fig. 7b), stronger contrasts become evident. The wave clouds discussed in the following section are still included and impact the frequency at temperatures below -18°C at Punta Arenas. The frequencies of ice-forming clouds are similar at Leipzig and Limassol, but 0.15 to 0.5 at -10°C lower at Punta Arenas. Comparing the coupled and uncoupled state at Leipzig and Limassol, ice formation is more frequent in the coupled case by 0.15 between -5 and -12°C .

425 3.2.3 Gravity wave influence on phase occurrence at low temperatures

The lower frequency of ice-containing cloud layers at temperatures below -18°C over Punta Arenas is a prominent feature of both, the lidar-radar and the lidar-only-equivalent datasets shown in Fig. 6. The associated high abundance of supercooled liquid water is frequently reported as a general phenomenon of stratiform clouds in the southern hemisphere mid and higher latitudes. Within this subsection, the reasons for the found behavior over Punta Arenas will be elaborated in more detail.

430 Frequently, the stratiform liquid-only cloud layers observed over Punta Arenas at temperatures below -18°C are embedded in orographic gravity waves. Following the wave detection methodology introduced in Sec. 2.5, Fig. 8 shows the autocorrelation and power spectra of the Doppler lidar vertical velocity for clouds classified as liquid only over Leipzig, Limassol, and Punta Arenas, respectively. These supercooled liquid-only clouds at Punta Arenas show high autocorrelation coefficients at long shifts (Fig. 8c1), whereas liquid only clouds with similar characteristics are absent at Limassol (Fig. 8b1) and Leipzig (Fig. 8a1). In
435 terms of spectral power density (Fig. 8c2), the strongly supercooled clouds at Punta Arenas show only low turbulence.




Figure 8. Vertical-velocity autocorrelation function (a1, b1, and c1) and power density (a2, b2, and c2) of the Doppler lidar vertical velocities for cloud cases classified as liquid-only. The columns are (a) Leipzig, (b) Limassol, and (c) Punta Arenas. The curves are binned to CTT intervals of 5 K with color indicating the CTT. The number of clouds for each temperature interval is given as numbers in the respective panel. Shading denotes the 10%–90% percentile range for the -5 and -30°C bin.

440 In a next step, the autocorrelation is used to filter the dataset for clouds affected by gravity waves. As described in Sec. 2.5, the length at which the autocorrelation coefficient drops below 0.8 is used as a characteristic value. Decreasing threshold value for this characteristic length will remove gravity-wave influenced clouds from the dataset. For large values of the length threshold, e.g., larger 1000 m, only clouds that are strongly forced by gravity waves will be removed, whereas going to shorter thresholds (< 500 m) will also remove weakly gravity-wave-influenced clouds. Fig. 9 shows an increase in the fraction of ice-containing clouds below -12°C with decreasing correlation-length thresholds from 30000 to 300 m. At Punta Arenas,

the fraction of ice-containing clouds increases from 0.5 to 0.85. In the temperature interval between -15 and -12°C , the fraction is 0.05 to 0.1 lower at Punta Arenas compared to Leipzig and Limassol. Hence, clouds with fully developed turbulence show similar ice-formation frequencies, independent of the location, with indications for a still slightly reduced ice-formation efficiency over Punta Arenas.

445




Figure 9. Fraction of ice containing clouds over temperature for Leipzig, Limassol, and Punta Arenas. Dash-dotted lines show the fractions for Leipzig, Limassol, and Punta Arenas with an autocorrelation coefficient smaller than 0.8 for a horizontal shift of 300 m. The fractions for cloud cases with longer autocorrelation are only shown for Punta Arenas. Total duration of the clouds in each bin is given by the numbers on top in hours.

3.2.4 Comparison of radar reflectivity factor of the ice virga

Prior studies of Zhang et al. (2018) identified a strong contrast in radar reflectivity factor between the different 30-deg latitude bands of the globe, with the southern hemisphere mid-latitudes (30 - 60°S) showing the lowest mean reflectivity of all regions. They concluded that this difference in reflectivity factor is associated to a respective difference in ice crystal mass and number concentration. In the following we provide a similar representation of regional contrasts of ice-virga reflectivity from ground-based perspective that is based on a single radar instrument. As described in Sec. 2.4, the amount of ice formed in the mixed phase layer is measured at six height bins (180 m) below the base of the liquid dominated cloud top and hence at the top of the virga (Bühl et al., 2016). Fig. 10a shows the cloud top temperature-resolved statistics of reflectivity for the three stations, which is based on the full cloud dataset, including the wave-influenced clouds (as these are also included in the study by Zhang et al., 2018). From -28°C to -16°C and again above -12°C , Punta Arenas shows the lowest reflectivity and Limassol the highest. For most temperatures, Punta Arenas is 5–8 dB below the northern hemispheric stations, only exception is the interval between -4 and -8°C , where the reflectivity for Punta Arenas and Leipzig is almost equal. At temperatures below -28°C , Limassol and Punta Arenas show equal reflectivity with values at Leipzig being slightly higher.




Figure 10. Amount of ice observed in the virga, (a) radar reflectivity factor observed at the top of the virga (180 m below the liquid-dominated layer base) and (b) ice-to-liquid-content-ratio binned by CTT. Temperature bins of 4°C.

We further investigated the properties of the ice-forming liquid-dominated cloud top layers and found that cloud thickness
460 agrees within 40 m above -30°C . The ice-to-liquid content ratio (Fig. 10b) is smaller at Punta Arenas, than at the northern
hemispheric locations, especially (factor 3) between -24 and -20°C , but also above -10°C . Hence, in these temperature
regimes, the liquid phase is less efficiently converted into ice in stratiform clouds above Punta Arenas.

4 Discussion of observed contrasts in properties of shallow clouds

In the previous section, a comprehensive overview of the aerosol conditions and stratiform cloud properties at the strongly
465 contrasting sites of Leipzig, Limassol, and Punta Arenas was presented. Several aerosol-, temperature-, surface-coupling-, as
well as dynamics-related differences have been identified and will be discussed in the following.

The analysis of the profiles of aerosol optical properties obtained from the Polly^{XT} lidar observations reveals almost similar
aerosol load at Punta Arenas and Limassol between 0 and -10°C . Leipzig shows a higher aerosol load in this temperature
interval. For lower temperatures the lowest average extinction was observed at Punta Arenas. Low values of particle depolar-
470 ization ratio indicate that non-spherical particles, such as mineral dust are completely absent at Punta Arenas below -10°C . The absence of continental aerosol species in the free troposphere is also consistent with the Southern Ocean south of Aus-
tralia (e.g., Alexander and Protat, 2019). Additionally, Uetake et al. (2020) showed for this region that INP are related to sea
spray emissions. Together with an air mass source estimate, the average profiles of optical parameters were used as an input
for INP parametrizations for a first estimate of long-term averages of INP concentrations. The lidar-based estimate of average
475 INP concentrations identifies the strongest differences in INP concentrations between -12 and -35°C . However, it is still
subject to discussion which fraction of marine particles is transported from the boundary layer to the free troposphere. This
insufficient knowledge propagates further into uncertainties in the fraction of continental aerosol, which leads to considerable

uncertainties in the estimated concentrations of INP. Bourgeois et al. (2018) argue that due to efficient removal processes, marine particles predominantly remain in the boundary layer (globally 80% of their AOT). Murphy et al. (2019) argue in a
480 similar direction, especially when sea salt is regarded a tracer of marine origin.

Opposed to prior lidar-only studies (e.g., Kanitz et al., 2011; Alexander and Protat, 2018), more than half of the clouds observed at -8°C at the three sites were found to contain ice. This is in line with the lidar detection threshold introduced by Bühl et al. (2013a), but extending their results to Limassol and Punta Arenas. This result has significance for lidar-only or
485 ceilometer datasets that are currently used for validation of climate models (e.g., Kuma et al., 2020). The occurrence of ice at these relatively warm temperatures is also in line with airborne in-situ observations over the Southern Ocean (Huang et al., 2017; D'Alessandro et al., 2019).

Ice formation at slightly supercooled conditions above -10°C was found to be equally frequent, suggesting the presence of
490 equally efficient INP at all sites. At Punta Arenas, such temperatures usually occur within the boundary layer, where presumably INP of biological origin are present (Gong et al., in preparation). Similarly high fractions were also observed for Arctic boundary layer clouds during summer and autumn (Achtert et al., 2020). Griesche et al. (2020) also found that surface coupling increased the frequency of ice formation in boundary layer clouds observed during Arctic summer.




Figure 11. Fraction of ice containing clouds over temperature for Leipzig, Limassol, and Punta Arenas when considering only fully turbulent clouds with an autocorrelation coefficient smaller than 0.8 for a horizontal shift of 300 m (see Fig. 7) and cloud bases in the free troposphere above 2 km height (see Fig. 9).

The liquid-only clouds at Punta Arenas found below -14°C are identified to be predominantly wave clouds forced by orography, likely in an early period of their lifecycle. In this kind of cloud, the liquid-dominated mixed-phase is separated horizontally from the ice virga, due to the rapid flow and weak turbulence (Heymsfield and Miloshevich, 1993). When sampled
495 by stationary observations, only parts of the clouds are covered and the phase occurrence frequency is biased. Using Doppler lidar vertical velocity observations, clouds driven by orographic wave dynamics can be identified based on their autocorrelation

function, as presented in Sec. 3.2.3. At temperatures below -14°C , approximately one-third of the cloud layers observed at Punta Arenas were influenced by gravity waves. Similar frequencies of occurrence of non-turbulent liquid layers were also found at Utqiagvik (Alaska) and McMurdo (Antarctica) deriving stability criteria from radiosoundings (Silber et al., 2020).

500 In a final step, the separation techniques for coupling and orographic waves can be combined, to assess contrasts ice frequency for free-tropospheric and fully-turbulent clouds. The resulting occurrence frequency is shown in Fig. 11. The fraction of ice-containing clouds at temperatures below -15°C is above 0.85 at all three sites. However, comparing to Leipzig and Limassol, the fraction of ice forming clouds with CTT between -25 and -15°C remains 0.1 lower at Punta Arenas. Mineral dust is known to be an efficient INP at these temperatures (Kanji et al., 2017), but was not observed at the respective temperatures at 505 Punta Arenas (Fig. 3). This difference in the non-wave cloud phase occurrence is also in agreement with satellite-based studies of Villanueva et al. (2020), who also attributed latitudinal differences in the ice occurrence to associated differences in the dust load. Fig. 11 also depicts the low frequency of ice formation in free-tropospheric fully-turbulent clouds at temperatures above -15°C . With less coupling to the near-surface aerosol reservoir, ice formation is strongly suppressed compared to Leipzig and Limassol.

510 The ice mass formed by stratiform liquid layers is lowest at Punta Arenas, as was found by investigating the radar reflectivity factor in the ice virga as a proxy. Especially between -28°C and -16°C , the observed radar reflectivity factor is up to 7 dB lower, compared to both sites in the northern hemisphere. This result is consistent with estimates from space-borne sensors covering the full Southern Ocean (Zhang et al., 2018). Contradicting, Arctic mixed-phase clouds were found to respond with lower IWC to increased loads of anthropogenic pollution (Norgren et al., 2018). As for the lower frequency of ice formation 515 discussed above, the difference in ice mass coincides with a lack of dust INPs at these temperatures. Slight differences in IWC or Z , respectively, might be explained by a slower mass growth rate, caused by a smaller vapor diffusion coefficient at higher ambient pressure (Hall and Pruppacher, 1976). When temperature and particle size are considered similar, the stratiform clouds subject to this study, experience 10% to 20% larger growth rates at Leipzig and Limassol than at Punta Arenas. Hence, the difference of a factor 3-6 larger ice mass cannot be explained solely by this effect alone.

520 Frequent occurrences of ice forming clouds above -10°C were found, which were not covered by studies based on space-borne active remote sensing. Similar to their colder counterparts, they show a lower ice amount in the virga above Punta Arenas, but with smaller difference compared to the sites in the northern hemisphere. With an average reflectivity of -36 dBZ , they are usually well below the detection limit of the CloudSat satellite in the A-Train constellation (Bühl et al., 2013a, 2016). Occurrence of such clouds in other parts of the Southern Ocean cannot be ruled out. A misclassification of supercooled drizzle 525 clouds as ice containing is unlikely, as they exceed -30 dBZ neither at cloud top, nor in the virga. From previous studies it is known that the onset of drizzle formation is usually associated to higher reflectivities, either above approximately -20 dBZ (Liu et al., 2008; Acquistapace et al., 2019) or at least above -30 dBZ (Wu et al., 2020).

5 Summary and Outlook

This study investigated contrasts in aerosol-cloud interactions in shallow supercooled stratiform clouds observed with the
530 ground-based remote sensing supersite LACROS at Leipzig, Limassol, and Punta Arenas.

Sampling the profiles of optical properties with temperature as a vertical coordinate revealed aerosol load at temperatures between -15 and 0°C being within a factor of 2 at Punta Arenas and Limassol. This finding is related to the cold and (compared to the free troposphere) aerosol laden boundary layer at Punta Arenas. At lower temperatures, the lowest β_p and extinction was observed over Punta Arenas, the highest over Leipzig. The very low particle depolarization ratio at Punta Arenas between -25
535 and -10°C , suggests the absence of mineral dust in a temperature regime, where dust is known to be an efficient INP. An estimate of INP concentrations at the respective temperatures based on the optical properties reveals differences of 1-4 orders of magnitude between Punta Arenas and the two northern hemispheric sites. In absence of abundant INPs from marine sources, the contribution of terrestrial sources causes strong variability.

The phase occurrence frequencies showed a higher fraction of ice containing clouds at weakly-supercooling temperatures
540 of above -10°C compared to prior lidar-only studies. A cloud radar with a sensitivity better than -40 dBZ is needed to sufficiently characterize low ice water contents in the virga formed by shallow stratiform clouds in this temperature regime. Coupling to the boundary layer increases the frequency of ice formation at slightly supercooling temperatures at all sites. The strongest contrasts in the ice-formation frequency between free-tropospheric and surface-coupled conditions were found for Punta Arenas. This finding is in compliance to the found contrasts in the INP profiles at the three sites and further indicates
545 that the free-tropospheric INP reservoir over the Southern Ocean is limited.

Frequent liquid only layers below -20°C at Punta Arenas were found to be associated with orographic gravity waves, causing two implications: (1) potential phase misclassification by stationary observers due to horizontal separation of ice and liquid phase; (2) sustained liquid water in updrafts, because the associated vertical velocities allow supersaturation over water; The newly developed Doppler lidar autocorrelation approach helps to address "[...]" the difficulty in characterizing gravity waves
550 from existing single-site measurements [...] as posed by Silber et al. (2020). Ice mass in the virga and the ice-to-liquid content ratio was found to be lowest at Punta Arenas, especially at temperatures where dust serves as an efficient INP at the northern hemispheric sites.

From the results of this study, two issues for further investigations arise. Firstly, importance of terrestrial sources of INP in the Southern Ocean. Observations of free-tropospheric aerosols upwind and downwind of major landmasses are required,
555 to quantify the terrestrial emission and identify regions, where the emitted INP impact cloud microphysics the strongest. Secondly, how important are gravity waves in the formation of clouds in other regions of the Southern Ocean? How frequent do gravity waves occur above the open ocean, which covers the vast majority of the Earths surface between 30 and 70°S ? Such measurements are required in order to further constrain the role of INP in the evident excess of liquid water in clouds of the southern-hemispheric mid-latitudes.

560 *Code and data availability.* The Cloudnet datasets are provided by the ACTRIS Data Centre node for cloud profiling under the following links: <https://hdl.handle.net/21.12132/2.e130b236928a4932> (Leipzig), <https://hdl.handle.net/21.12132/2.a056a828a6b94d1f> (Limassol) and <https://hdl.handle.net/21.12132/2.b6c194d7d33b448e> (Punta Arenas). In the upcoming future, the Doppler lidar and PollyNET datasets will also be available via ACTRIS. Meanwhile, they can be obtained upon request at polly@tropos.de. The source code of the cloud sniffer, the database of identified clouds and analysis scripts are available at https://github.com/martin-rdz/larda_cloud_sniffer (last accessed 27.04.2021; Radenz and Bühl, 2021) and requires a setup of pyLARDA (<https://github.com/lacros-tropos/larda>, last accessed 27.04.2021; Bühl et al., 2021). AERONET photometer observations at Leipzig, Limassol and Punta Arenas are available from the AERONET database (<http://aeronet.gsfc.nasa.gov/>).

Appendix A: Wave-cloud example

Fig. A1 shows a typical wave cloud at Punta Arenas. Updrafts in stationary or slowly propagating waves can sustain liquid layers without showing signatures of sedimenting ice crystals to the fixed observer. The ice phase is only visible downwind as soon as the droplets have evaporated. Such a wave cloud was observed at Punta Arenas on the 27 September 2019. A mid-tropospheric long-wave ridge caused a strong, zonal westerly flow at Punta Arenas. Velocity of the horizontal wind exceeded 25 m s⁻¹ at 4000 m height. Layerd clouds developed within the orographic gravity waves triggered by this flow. One of the cloud layers with a CTT between -34 and -31°C is depicted in Fig. A1. It illustrates how liquid and ice phase are horizontally separated. Embedded in a slowly propagating gravity wave, the cloud was advected over the site and the full life cycle could be observed. Based on the Doppler lidar vertical velocity (Fig. A1d), the updraft was present until 05:32 UTC. Afterwards, downward motion with increasing velocity set in. Only after the liquid droplets evaporated at 06:17 UTC, the ice particles can be clearly identified in the remote-sensing observations (Fig. A1b, c).

Appendix B: Air mass source estimate

580 The automated time–height-resolved air mass source attribution described by Radenz et al. (2021) is used to characterize air mass origin for the LACROS observations. As in the original publication, 10-day HYSPLIT ensemble backward trajectories are calculated in intervals of 3 hours and 500 m throughout the period of the deployment. For the calculation of the residence times, a reception height of 2 km is used. Additionally to the MODIS based surface cover classification and the the named geography (individually for Leipzig, Limassol, and Punta Arenas), 30°-wide bands of latitude are used to characterize meridional transport. The assignment of the surface cover classed and named geography are depicted in Fig. 2 and 3 of Radenz et al. (2021). The average residence time for each set of surface types is shown in Fig. B1. However, as for the aerosol optical properties, geometrical height is less insightful as temperature. Thus, the residence time at each 3 h profile is binned by ambient temperature, using analysis profiles. Then, the average residence time is calculated per temperature bin (Fig. B2).




Figure A1. Wave cloud observed at Punta Arenas on 27 September 2019, 05:00–06:45 UTC. (a) Lidar attenuated backscatter, (b) Cloudnet target categorization, (c) Lidar volume linear depolarization ratio and (d) Doppler lidar vertical velocity. The temperature information in (b) is obtained from ECMWF IFS model data. The red rectangle in (d) indicates the automatically detected liquid-topped shallow stratiform cloud. The inset in (d) shows the autocorrelation function of the vertical velocities in the liquid-dominated layer. The height of the velocity sampling is indicated with a dashed line.

Author contributions. MR analyzed the data and drafted the manuscript. MR, PS, JB, HB, RE, BBG, REM, and FZ conducted the campaigns 590 and operated LACROS. PS, MR, JB generated the Cloudnet dataset. HB processed the Polly^{XT} lidar data. PS, JB, and AA supervised the work and revised the manuscript. All authors jointly contributed to the paper and the scientific discussion.

Competing interests. No competing interests are present.

Acknowledgements. The authors wish to thank TROPOS, Cyprus University of Technology and University of Magallanes for their logistic and infrastructural support during the LACROS deployments. We furthermore thank Teresa Vogl, Willi Schimmel, Audrey Teisseire, Cristofer 595 Jimenez, Heike Kalesse-Los, and Roland Schrödner for keeping LACROS operational at Punta Arenas. We gratefully acknowledge the ACTRIS Cloud Remote Sensing Unit for making the Cloudnet datasets publicly available. LACROS operations were supported by the European Union (EU) Horizon 2020 (ACTRIS; grant no. 654109) and the Seventh Framework Programme (BACCHUS; grant no. 603445). Observations at Leipzig were supported by the German Bundesministerium für Bildung und Forschung funded projects “High Definition 600 Clouds and Precipitation for Climate Prediction – HD(CP)2” (grant nos. 01LK1503F, 01LK1502I, 01LK1209C, and 01LK1212C). Part of the CyCARE and DACAPO-PESO campaigns were funded by the Deutsche Forschungsgemeinschaft (DFG – German Research Foundation)




Figure B1. Mean normalized residence time profile at (a) Leipzig, (b) Limassol, and (c) Punta Arenas for a reception height of 2 km. Residence times are shown for the surface cover classification (1), named areas (2) and latitude bands (3).

project PICNICC (SE2464/1-1 and KA4162/2-1). JB acknowledges funding by DFG project COARSEMIX (398285025). REM has been financial supported by the SIROCCO project (grant no. EXCELLENCE/1216/0217) co-funded by the Republic of Cyprus and the structural funds of the EU for Cyprus through the Research and Innovation Foundation. Thanks are also given to EXCELSIOR EU H2020 Widespread Teaming program with Grant Agreement No 857510 (www.excelsior2020.eu) and the Republic of Cyprus for the support of the establishment 605 the ERATOSTHENES Centre of Excellence. Thanks are also given to EXCELSIOR EU H2020 Widespread Teaming program with Grant Agreement No 857510 (www.excelsior2020.eu) and the Republic of Cyprus for the support of the establishment the ERATOSTHENES Centre of Excellence. BBG acknowledges funding by the ANID/CONICYT/FONDECYT Iniciación 11181335. HB acknowledges funding




Figure B2. Mean normalized residence time binned by temperature at (a) Leipzig, (b) Limassol, and (c) Punta Arenas for a reception height of 2 km. Residence times are shown for the surface cover classification (1), named areas (2) and latitude bands (3). Hatching indicates bins with less than 1 % of the values.

by the AEOLUS Cal/Val activities (German Bundesministeriums für Wirtschaft und Energie; grant no. 50EE1721C). We thank AERONET-Europe for providing the calibration service of the photometer. The authors gratefully acknowledge the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport and dispersion model. We thank Timothy J. Dunkerton for serving as an editor and gratefully acknowledge the constructive comments of the two referees.

References

Achert, P., O'Connor, E. J., Brooks, I. M., Sotiropoulou, G., Shupe, M. D., Pospichal, B., Brooks, B. J., and Tjernström, M.: Properties of Arctic Liquid and Mixed-Phase Clouds from Shipborne Cloudnet Observations during ACSE 2014, *Atmospheric Chemistry and Physics*, 20, 14 983–15 002, <https://doi.org/10.5194/acp-20-14983-2020>, 2020.

615 Acquistapace, C., Löhner, U., Maahn, M., and Kollias, P.: A New Criterion to Improve Operational Drizzle Detection with Ground-Based Remote Sensing, *Journal of Atmospheric and Oceanic Technology*, 36, 781–801, <https://doi.org/10.1175/JTECH-D-18-0158.1>, 2019.

Alexander, S. P. and Protat, A.: Cloud Properties Observed From the Surface and by Satellite at the Northern Edge of the Southern Ocean, *Journal of Geophysical Research: Atmospheres*, 123, 443–456, <https://doi.org/10.1002/2017JD026552>, 2018.

620 Alexander, S. P. and Protat, A.: Vertical Profiling of Aerosols With a Combined Raman-Elastic Backscatter Lidar in the Remote Southern Ocean Marine Boundary Layer (43–66°S, 132–150°E), *Journal of Geophysical Research: Atmospheres*, 124, 12 107–12 125, <https://doi.org/10.1029/2019JD030628>, 2019.

Alexander, S. P., Sato, K., Watanabe, S., Kawatani, Y., and Murphy, D. J.: Southern Hemisphere Extratropical Gravity Wave Sources and Intermittency Revealed by a Middle-Atmosphere General Circulation Model, *Journal of the Atmospheric Sciences*, 73, 1335–1349, 625 <https://doi.org/10.1175/JAS-D-15-0149.1>, 2016.

Alexander, S. P., Orr, A., Webster, S., and Murphy, D. J.: Observations and fine-scale model simulations of gravity waves over Davis, East Antarctica (69°S, 78°E), *Journal of Geophysical Research: Atmospheres*, 122, 7355–7370, <https://doi.org/10.1002/2017JD026615>, 2017.

Alexander, S. P., McFarquhar, G. M., Marchand, R., Protat, A., Vignon, É., Mace, G. G., and Klekociuk, A. R.: Mixed-phase Clouds and Precipitation in Southern Ocean Cyclones and Cloud Systems Observed Poleward of 64°S by Ship-based Cloud Radar and Lidar, *Journal 630 of Geophysical Research: Atmospheres*, <https://doi.org/10.1029/2020JD033626>, 2021.

Althausen, D., Engelmann, R., Baars, H., Heese, B., Ansmann, A., Müller, D., and Komppula, M.: Portable Raman Lidar PollyXT for Automated Profiling of Aerosol Backscatter, Extinction, and Depolarization, *Journal of Atmospheric and Oceanic Technology*, 26, 2366–2378, <https://doi.org/10.1175/2009JTECHA1304.1>, 2009.

Ansmann, A.: Ice Formation in Saharan Dust over Central Europe Observed with Temperature/Humidity/Aerosol Raman Lidar, *Journal 635 of Geophysical Research*, 110, D18S12, <https://doi.org/10.1029/2004JD005000>, 2005.

Ansmann, A., Bösenberg, J., Chaikovsky, A., Comerón, A., Eckhardt, S., Eixmann, R., Freudenthaler, V., Ginoux, P., Komguem, L., Linné, H., Márquez, M. n. L., Matthias, V., Mattis, I., Mitev, V., Müller, D., Music, S., Nickovic, S., Pelon, J., Sauvage, L., Sobolewsky, P., Srivastava, M. K., Stohl, A., Torres, O., Vaughan, G., Wandinger, U., and Wiegner, M.: Long-range transport of Saharan dust to northern Europe: The 11–16 October 2001 outbreak observed with EARLINET, *Journal of Geophysical Research: Atmospheres*, 108, AAC 12, 640 <https://doi.org/10.1029/2003JD003757>, 2003.

Ansmann, A., Mamouri, R.-E., Bühl, J., Seifert, P., Engelmann, R., Hofer, J., Nisantzi, A., Atkinson, J. D., Kanji, Z. A., Sierau, B., Vrekoussis, M., and Sciare, J.: Ice-Nucleating Particle versus Ice Crystal Number Concentrationin Altocumulus and Cirrus Layers Embedded in Saharan Dust:A Closure Study, *Atmospheric Chemistry and Physics*, 19, 15 087–15 115, <https://doi.org/10.5194/acp-19-15087-2019>, 2019.

645 Baars, H., Kanitz, T., Engelmann, R., Althausen, D., Heese, B., Komppula, M., Preißler, J., Tesche, M., Ansmann, A., Wandinger, U., Lim, J.-H., Ahn, J. Y., Stachlewska, I. S., Amiridis, V., Marinou, E., Seifert, P., Hofer, J., Skupin, A., Schneider, F., Bohlmann, S., Foth, A., Bley, S., Pfüller, A., Giannakaki, E., Lihavainen, H., Viisanen, Y., Hooda, R. K., Pereira, S. N., Bortoli, D., Wagner, F., Mattis, I., Janicka, L., Markowicz, K. M., Achert, P., Artaxo, P., Pauliquevis, T., Souza, R. A. F., Sharma, V. P., van Zyl, P. G., Beukes, J. P., Sun, J., Rohwer,

650 E. G., Deng, R., Mamouri, R.-E., and Zamorano, F.: An overview of the first decade of Polly^{NET}: an emerging network of automated Raman-polarization lidars for continuous aerosol profiling, *Atmospheric Chemistry and Physics*, 16, 5111–5137, <https://doi.org/10.5194/acp-16-5111-2016>, 2016.

Baars, H., Seifert, P., Engelmann, R., and Wandinger, U.: Target categorization of aerosol and clouds by continuous multiwavelength-polarization lidar measurements, *Atmospheric Measurement Techniques*, 10, 3175–3201, <https://doi.org/10.5194/amt-10-3175-2017>, 2017.

655 Baars, H., Radenz, M., Floutsi, A. A., Engelmann, R., Althausen, D., Heese, B., Ansmann, A., Flament, T., Dabas, A., Trapon, D., Reitebuch, O., Bley, S., and Wandinger, U.: Californian Wildfire Smoke Over Europe: A First Example of the Aerosol Observing Capabilities of Aeolus Compared to Ground-Based Lidar, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL092194>, 2021.

Baker, B. A. and Lawson, R. P.: In Situ Observations of the Microphysical Properties of Wave, Cirrus, and Anvil Clouds. Part I: Wave Clouds, *Journal of the Atmospheric Sciences*, 63, 3160–3185, <https://doi.org/10.1175/JAS3802.1>, 2006.

660 Barreto, Á., Cuevas, E., Granados-Muñoz, M.-J., Alados-Arboledas, L., Romero, P. M., Gröbner, J., Kouremeti, N., Almansa, A. F., Stone, T., Toledano, C., Román, R., Sorokin, M., Holben, B., Canini, M., and Yela, M.: The New Sun-Sky-Lunar Cimel CE318-T Multiband Photometer – a Comprehensive Performance Evaluation, *Atmospheric Measurement Techniques*, 9, 631–654, <https://doi.org/10.5194/amt-9-631-2016>, 2016.

Bühl, J., Ansmann, A., Seifert, P., Baars, H., and Engelmann, R.: Toward a quantitative characterization of heterogeneous ice formation with lidar/radar: Comparison of CALIPSO/CloudSat with ground-based observations, *Geophysical Research Letters*, 40, 4404–4408, <https://doi.org/10.1002/grl.50792>, 2013a.

665 Bühl, J., Seifert, P., Wandinger, U., Baars, H., Kanitz, T., Schmidt, J., Myagkov, A., Engelmann, R., Skupin, A., Heese, B., Klepel, A., Althausen, D., and Ansmann, A.: LACROS: The Leipzig Aerosol and Cloud Remote Observations System, in: SPIE Remote Sensing, edited by Comeron, A., Kassianov, E. I., Schäfer, K., Stein, K., and Gonglewski, J. D., p. 889002, Dresden, Germany, <https://doi.org/10.1117/12.2030911>, 2013b.

Bühl, J., Seifert, P., Myagkov, A., and Ansmann, A.: Measuring ice- and liquid-water properties in mixed-phase cloud layers at the Leipzig Cloudnet station, *Atmospheric Chemistry and Physics*, 16, 10 609–10 620, <https://doi.org/10.5194/acp-16-10609-2016>, 2016.

Bühl, J., Seifert, P., Engelmann, R., and Ansmann, A.: Impact of vertical air motions on ice formation rate in mixed-phase cloud layers, *npj Climate and Atmospheric Science*, 2, 36, <https://doi.org/10.1038/s41612-019-0092-6>, 2019.

675 Bühl, J., Radenz, M., Schimmel, W., Vogl, T., Röttenbacher, J., and Lochmann, M.: pyLARDA v3.2, <https://doi.org/10.5281/zenodo.4721311>, 2021.

Bodas-Salcedo, A., Williams, K. D., Ringer, M. A., Beau, I., Cole, J. N. S., Dufresne, J.-L., Koshiro, T., Stevens, B., Wang, Z., and Yokohata, T.: Origins of the Solar Radiation Biases over the Southern Ocean in CMIP2 Models*, *Journal of Climate*, 27, 41–56, <https://doi.org/10.1175/JCLI-D-13-00169.1>, 2014.

680 Bodas-Salcedo, A., Hill, P., Furtado, K., Williams, K., Field, P., Manners, J., Hyder, P., and Kato, S.: Large contribution of supercooled liquid clouds to the solar radiation budget of the Southern Ocean, *Journal of Climate*, 29, 4213–4228, 2016.

Bohlmann, S., Baars, H., Radenz, M., Engelmann, R., and Macke, A.: Ship-Borne Aerosol Profiling with Lidar over the Atlantic Ocean: From Pure Marine Conditions to Complex Dust–Smoke Mixtures, *Atmospheric Chemistry and Physics*, 18, 9661–9679, <https://doi.org/10.5194/acp-18-9661-2018>, 2018.

685 Bourgeois, Q., Ekman, A. M. L., Renard, J.-B., Krejci, R., Devasthale, A., Bender, F. A.-M., Riipinen, I., Berthet, G., and Tackett, J. L.: How
much of the global aerosol optical depth is found in the boundary layer and free troposphere?, *Atmospheric Chemistry and Physics*, 18,
7709–7720, <https://doi.org/10.5194/acp-18-7709-2018>, 2018.

690 Bromwich, D. H., Werner, K., Casati, B., Powers, J. G., Gorodetskaya, I. V., Massonnet, F., Vitale, V., Heinrich, V. J., Liggett, D., Arndt, S.,
Barja, B., Bazile, E., Carpentier, S., Carrasco, J. F., Choi, T., Choi, Y., Colwell, S. R., Cordero, R. R., Gervasi, M., Haiden, T., Hirasawa,
N., Inoue, J., Jung, T., Kalesse, H., Kim, S.-J., Lazzara, M. A., Manning, K. W., Norris, K., Park, S.-J., Reid, P., Rigor, I., Rowe, P. M.,
Schmithüsen, H., Seifert, P., Sun, Q., Uttal, T., Zannoni, M., and Zou, X.: The Year of Polar Prediction in the Southern Hemisphere
(YOPP-SH), *Bulletin of the American Meteorological Society*, 101, E1653–E1676, <https://doi.org/10.1175/BAMS-D-19-0255.1>, 2020.

700 Choi, Y.-S., Lindzen, R. S., Ho, C.-H., and Kim, J.: Space observations of cold-cloud phase change, *Proceedings of the National Academy of
Sciences*, 107, 11 211–11 216, <https://doi.org/10.1073/pnas.1006241107>, 2010.

695 Cotton, R. J. and Field, P. R.: Ice nucleation characteristics of an isolated wave cloud, *Quarterly Journal of the Royal Meteorological Society*,
128, 2417–2437, <https://doi.org/10.1256/qj.01.150>, 2002.

Crewell, S. and Löhner, U.: Accuracy of Boundary Layer Temperature Profiles Retrieved With Multifrequency Multiangle Microwave
Radiometry, *IEEE Transactions on Geoscience and Remote Sensing*, 45, 2195–2201, <https://doi.org/10.1109/TGRS.2006.888434>, 2007.

D'Alessandro, J. J., Diao, M., Wu, C., Liu, X., Jensen, J. B., and Stephens, B. B.: Cloud Phase and Relative Humidity Distributions over
700 the Southern Ocean in Austral Summer Based on In Situ Observations and CAM5 Simulations, *Journal of Climate*, 32, 2781–2805,
<https://doi.org/10.1175/JCLI-D-18-0232.1>, 2019.

de Boer, G., Eloranta, E. W., and Shupe, M. D.: Arctic Mixed-Phase Stratiform Cloud Properties from Multiple Years of Surface-Based Mea-
surements at Two High-Latitude Locations, *Journal of the Atmospheric Sciences*, 66, 2874–2887, <https://doi.org/10.1175/2009JAS3029.1>,
2009.

705 DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.:
Predicting global atmospheric ice nuclei distributions and their impacts on climate, *Proceedings of the National Academy of Sciences*,
107, 11 217–11 222, <https://doi.org/10.1073/pnas.0910818107>, 2010.

710 DeMott, P. J., Prenni, A. J., McMeeking, G. R., Sullivan, R. C., Petters, M. D., Tobo, Y., Niemand, M., Möhler, O., Snider, J. R., Wang, Z.,
and Kreidenweis, S. M.: Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust
particles, *Atmospheric Chemistry and Physics*, 15, 393–409, <https://doi.org/10.5194/acp-15-393-2015>, 2015.

Desai, N., Chandrakar, K. K., Kinney, G., Cantrell, W., and Shaw, R. A.: Aerosol-Mediated Glaciation of Mixed-Phase Clouds: Steady-State
715 Laboratory Measurements, *Geophysical Research Letters*, 46, 9154–9162, <https://doi.org/10.1029/2019GL083503>, 2019.

Durant, A. J.: Evaporation Freezing by Contact Nucleation Inside-Out, *Geophysical Research Letters*, 32, L20 814,
<https://doi.org/10.1029/2005GL024175>, 2005.

720 Engelmann, R., Kanitz, T., Baars, H., Heese, B., Althausen, D., Skupin, A., Wandinger, U., Komppula, M., Stachlewska, I. S.,
Amiridis, V., Marinou, E., Mattis, I., Linné, H., and Ansmann, A.: The Automated Multiwavelength Raman Polarization and
Water-Vapor Lidar Polly^{XT}: The neXT Generation, *Atmospheric Measurement Techniques*, 9, 1767–1784,
<https://doi.org/10.5194/amt-9-1767-2016>, 2016.

Fan, J., Wang, Y., Rosenfeld, D., and Liu, X.: Review of Aerosol–Cloud Interactions: Mechanisms, Significance, and Challenges, *Journal of
725 the Atmospheric Sciences*, 73, 4221–4252, <https://doi.org/10.1175/JAS-D-16-0037.1>, 2016.

Fang, M., Albrecht, B. A., Ghate, V. P., and Kollias, P.: Turbulence in Continental Stratocumulus, Part I: External Forcings and Turbulence
Structures, *Boundary-Layer Meteorology*, 150, 341–360, <https://doi.org/10.1007/s10546-013-9873-3>, 2014.

Fernald, F. G.: Analysis of Atmospheric Lidar Observations: Some Comments, *Applied optics*, 23, 652–653, 1984.

Field, P. R., Heymsfield, A. J., Shipway, B. J., DeMott, P. J., Pratt, K. A., Rogers, D. C., Stith, J., and Prather, K. A.: Ice in Clouds Experiment—Layer Clouds. Part II: Testing Characteristics of Heterogeneous Ice Formation in Lee Wave Clouds, *Journal of the Atmospheric Sciences*, 69, 1066–1079, <https://doi.org/10.1175/JAS-D-11-026.1>, 2012.

Floutsi, A. A., Baars, H., Radenz, M., Haarig, M., Yin, Z., Seifert, P., Jimenez, C., Ansmann, A., Engelmann, R., Barja, B., Zamorano, F., and Wandinger, U.: Advection of Biomass Burning Aerosols towards the Southern Hemispheric Mid-Latitude Station of Punta Arenas as Observed with Multiwavelength Polarization Raman Lidar, *Remote Sensing*, 13, 138, <https://doi.org/10.3390/rs13010138>, 2021.

Foth, A., Kanitz, T., Engelmann, R., Baars, H., Radenz, M., Seifert, P., Barja, B., Fromm, M., Kalesse, H., and Ansmann, A.: Vertical aerosol distribution in the southern hemispheric midlatitudes as observed with lidar in Punta Arenas, Chile (53.2° S and 70.9° W), during ALPACA, *Atmospheric Chemistry and Physics*, 19, 6217–6233, <https://doi.org/10.5194/acp-19-6217-2019>, 2019.

Franklin, C. N., Sun, Z., Bi, D., Dix, M., Yan, H., and Bodas-Salcedo, A.: Evaluation of clouds in ACCESS using the satellite simulator package COSP: Global, seasonal, and regional cloud properties, *Journal of Geophysical Research: Atmospheres*, 118, 732–748, <https://doi.org/10.1029/2012JD018469>, 2013.

Fukuta, N. and Takahashi, T.: The Growth of Atmospheric Ice Crystals: A Summary of Findings in Vertical Supercooled Cloud Tunnel Studies, *Journal of the Atmospheric Sciences*, 56, 1963–1979, [https://doi.org/10.1175/1520-0469\(1999\)056<1963:TGOAIC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<1963:TGOAIC>2.0.CO;2), 1999.

Görsdorf, U., Lehmann, V., Bauer-Pfundstein, M., Peters, G., Vavrik, D., Vinogradov, V., and Volkov, V.: A 35-GHz Polarimetric Doppler Radar for Long-Term Observations of Cloud Parameters—Description of System and Data Processing, *Journal of Atmospheric and Oceanic Technology*, 32, 675–690, <https://doi.org/10.1175/JTECH-D-14-00066.1>, 2015.

Griesche, H. J., Ohneiser, K., Seifert, P., Ansmann, A., and Engelmann, R.: Contrasting Ice Formation in Arctic Clouds: Surface Coupled vs Decoupled Clouds, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–22, <https://doi.org/10.5194/acp-2020-1096>, 2020.

Grise, K. M., Polvani, L. M., and Fasullo, J. T.: Reexamining the Relationship between Climate Sensitivity and the Southern Hemisphere Radiation Budget in CMIP Models, *Journal of Climate*, 28, 9298–9312, <https://doi.org/10.1175/JCLI-D-15-0031.1>, 2015.

Haarig, M., Ansmann, A., Gasteiger, J., Kandler, K., Althausen, D., Baars, H., Radenz, M., and Farrell, D. A.: Dry versus wet marine particle optical properties: RH dependence of depolarization ratio, backscatter, and extinction from multiwavelength lidar measurements during SALTRACE, *Atmospheric Chemistry and Physics*, 17, 14 199–14 217, <https://doi.org/10.5194/acp-17-14199-2017>, 2017.

Haarig, M., Ansmann, A., Baars, H., Jimenez, C., Veselovskii, I., Engelmann, R., and Althausen, D.: Depolarization and lidar ratios at 355, 532, and 1064 nm and microphysical properties of aged tropospheric and stratospheric Canadian wildfire smoke, *Atmospheric Chemistry and Physics*, 18, 11 847–11 861, <https://doi.org/10.5194/acp-18-11847-2018>, 2018.

Hall, W. D. and Pruppacher, H. R.: The Survival of Ice Particles Falling from Cirrus Clouds in Subsaturated Air, *Journal of the Atmospheric Sciences*, 33, 1995–2006, [https://doi.org/10.1175/1520-0469\(1976\)033<1995:TSOIPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<1995:TSOIPF>2.0.CO;2), 1976.

Hamilton, D. S., Lee, L. A., Pringle, K. J., Reddington, C. L., Spracklen, D. V., and Carslaw, K. S.: Occurrence of pristine aerosol environments on a polluted planet, *Proceedings of the National Academy of Sciences*, 111, 18 466–18 471, <https://doi.org/10.1073/pnas.1415440111>, 2014.

Haynes, J. M., Jakob, C., Rossow, W. B., Tselioudis, G., and Brown, J.: Major Characteristics of Southern Ocean Cloud Regimes and Their Effects on the Energy Budget, *Journal of Climate*, 24, 5061–5080, <https://doi.org/10.1175/2011JCLI4052.1>, 2011.

Heymsfield, A. J. and Miloshevich, L. M.: Homogeneous Ice Nucleation and Supercooled Liquid Water in Orographic Wave Clouds, *Journal of the Atmospheric Sciences*, 50, 2335–2353, [https://doi.org/10.1175/1520-0469\(1993\)050<2335:HINASL>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<2335:HINASL>2.0.CO;2), 1993.

Hogan, R. J. and O'Connor, E. J.: Facilitating cloud radar and lidar algorithms: the Cloudnet Instrument Synergy/Target Categorization product, Tech. rep., University of Reading, Department of Meteorology, <http://www.met.rdg.ac.uk/~swrhgnrj/publications/categorization.pdf>, 2004.

Hogan, R. J., Mittermaier, M. P., and Illingworth, A. J.: The Retrieval of Ice Water Content from Radar Reflectivity Factor and Temperature and Its Use in Evaluating a Mesoscale Model, *Journal of Applied Meteorology and Climatology*, 45, 301–317, <https://doi.org/10.1175/jam2340.1>, 2006.

Hu, Y., Rodier, S., Xu, K.-m., Sun, W., Huang, J., Lin, B., Zhai, P., and Josset, D.: Occurrence, liquid water content, and fraction of super-cooled water clouds from combined CALIOP/IIR/MODIS measurements, *Journal of Geophysical Research: Atmospheres*, 115, D00H34, <https://doi.org/10.1029/2009JD012384>, 2010.

Huang, Y., Protat, A., Siems, S. T., and Manton, M. J.: A-Train Observations of Maritime Midlatitude Storm-Track Cloud Systems: Comparing the Southern Ocean against the North Atlantic, *Journal of Climate*, 28, 1920–1939, <https://doi.org/10.1175/JCLI-D-14-00169.1>, 2015.

Huang, Y., Chubb, T., Baumgardner, D., deHoog, M., Siems, S. T., and Manton, M. J.: Evidence for Secondary Ice Production in Southern Ocean Open Cellular Convection: Secondary Ice Production over the Southern Ocean, *Quarterly Journal of the Royal Meteorological Society*, 143, 1685–1703, <https://doi.org/10.1002/qj.3041>, 2017.

Hyder, P., Edwards, J. M., Allan, R. P., Hewitt, H. T., Bracegirdle, T. J., Gregory, J. M., Wood, R. A., Meijers, A. J. S., Mulcahy, J., Field, P., Furtado, K., Bodas-Salcedo, A., Williams, K. D., Copsey, D., Josey, S. A., Liu, C., Roberts, C. D., Sanchez, C., Ridley, J., Thorpe, L., Hardiman, S. C., Mayer, M., Berry, D. I., and Belcher, S. E.: Critical Southern Ocean Climate Model Biases Traced to Atmospheric Model Cloud Errors, *Nature Communications*, 9, 3625, <https://doi.org/10.1038/s41467-018-05634-2>, 2018.

Illingworth, A. J., Hogan, R. J., O'Connor, E., Bouniol, D., Brooks, M. E., Delanoé, J., Donovan, D. P., Eastment, J. D., Gaussiat, N., Goddard, J. W. F., Haeffelin, M., Baltink, H. K., Krasnov, O. A., Pelon, J., Piriou, J.-M., Protat, A., Russchenberg, H. W. J., Seifert, A., Tompkins, A. M., van Zadelhoff, G.-J., Vinit, F., Willén, U., Wilson, D. R., and Wrench, C. L.: Cloudnet: Continuous Evaluation of Cloud Profiles in Seven Operational Models Using Ground-Based Observations, *Bulletin of the American Meteorological Society*, 88, 883–898, <https://doi.org/10.1175/BAMS-88-6-883>, 2007.

Jimenez, C., Ansmann, A., Engelmann, R., Donovan, D., Malinka, A., Seifert, P., Wiesen, R., Radenz, M., Yin, Z., Bühl, J., Schmidt, J., Barja, B., and Wandinger, U.: The Dual-Field-of-View Polarization Lidar Technique: A New Concept in Monitoring Aerosol Effects in Liquid-Water Clouds – Case Studies, *Atmospheric Chemistry and Physics*, 20, 15 265–15 284, <https://doi.org/10.5194/acp-20-15265-2020>, 2020.

Kanitz, T., Seifert, P., Ansmann, A., Engelmann, R., Althausen, D., Casiccia, C., and Rohwer, E. G.: Contrasting the impact of aerosols at northern and southern midlatitudes on heterogeneous ice formation, *Geophysical Research Letters*, 38, L17802, <https://doi.org/10.1029/2011gl048532>, 2011.

Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of Ice Nucleating Particles, *Meteorological Monographs*, 58, 1.1–1.33, <https://doi.org/10.1175/AMSMONOGRAPH-D-16-0006.1>, 2017.

Kay, J. E., Bourdages, L., Miller, N. B., Morrison, A., Yettella, V., Chepfer, H., and Eaton, B.: Evaluating and improving cloud phase in the Community Atmosphere Model version 5 using spaceborne lidar observations, *Journal of Geophysical Research: Atmospheres*, 121, 4162–4176, <https://doi.org/10.1002/2015JD024699>, 2016.

Korolev, A.: Limitations of the Wegener–Bergeron–Findeisen Mechanism in the Evolution of Mixed-Phase Clouds, *Journal of the Atmospheric Sciences*, 64, 3372–3375, <https://doi.org/10.1175/JAS4035.1>, 2007.

Kuma, P., McDonald, A. J., Morgenstern, O., Alexander, S. P., Cassano, J. J., Garrett, S., Halla, J., Hartery, S., Harvey, M. J., Parsons, S.,
800 Plank, G., Varma, V., and Williams, J.: Evaluation of Southern Ocean cloud in the HadGEM3 general circulation model and MERRA-2
reanalysis using ship-based observations, *Atmospheric Chemistry and Physics*, 20, 6607–6630, <https://doi.org/10.5194/acp-20-6607-2020>,
2020.

Liu, Y., Geerts, B., Miller, M., Daum, P., and McGraw, R.: Threshold Radar Reflectivity for Drizzling Clouds, *Geophysical Research Letters*,
35, L03 807, <https://doi.org/10.1029/2007GL031201>, 2008.

805 Löffler-Mang, M. and Joss, J.: An Optical Disdrometer for Measuring Size and Velocity of Hydrometeors, *Journal of Atmospheric and
Oceanic Technology*, 17, 130–139, [https://doi.org/10.1175/1520-0426\(2000\)017<0130:AODFMS>2.0.CO;2](https://doi.org/10.1175/1520-0426(2000)017<0130:AODFMS>2.0.CO;2), 2000.

Mace, G. G. and Protat, A.: Clouds over the Southern Ocean as Observed from the R/V Investigator during CAPRICORN. Part I: Cloud
Occurrence and Phase Partitioning, *Journal of Applied Meteorology and Climatology*, 57, 1783–1803, [https://doi.org/10.1175/JAMC-D-17-0194.1](https://doi.org/10.1175/JAMC-D-
17-0194.1), 2018.

810 Mace, G. G., Benson, S., and Hu, Y.: On the Frequency of Occurrence of the Ice Phase in Supercooled Southern Ocean Low Clouds Derived
From CALIPSO and CloudSat, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL087554>, 2020.

Mamouri, R.-E. and Ansmann, A.: Potential of polarization lidar to provide profiles of CCN- and INP-relevant aerosol parameters, *Atmo-
spheric Chemistry and Physics*, 16, 5905–5931, <https://doi.org/10.5194/acp-16-5905-2016>, 2016.

Mamouri, R.-E. and Ansmann, A.: Potential of polarization/Raman lidar to separate fine dust, coarse dust, maritime, and anthropogenic
815 aerosol profiles, *Atmospheric Measurement Techniques*, 10, 3403–3427, <https://doi.org/10.5194/amt-10-3403-2017>, 2017.

McCluskey, C. S., Hill, T. C. J., Humphries, R. S., Rauker, A. M., Moreau, S., Strutton, P. G., Chambers, S. D., Williams, A. G., McRobert,
I., Ward, J., Keywood, M. D., Harnwell, J., Ponsonby, W., Loh, Z. M., Krummel, P. B., Protat, A., Kreidenweis, S. M., and De-
Mott, P. J.: Observations of Ice Nucleating Particles Over Southern Ocean Waters, *Geophysical Research Letters*, 45, 11,989–11,997,
<https://doi.org/10.1029/2018GL079981>, 2018a.

820 McCluskey, C. S., Ovadnevaite, J., Rinaldi, M., Atkinson, J., Belosi, F., Ceburnis, D., Marullo, S., Hill, T. C. J., Lohmann, U., Kanji,
Z. A., O'Dowd, C., Kreidenweis, S. M., and DeMott, P. J.: Marine and Terrestrial Organic Ice-Nucleating Particles in Pristine Ma-
rine to Continentally Influenced Northeast Atlantic Air Masses, *Journal of Geophysical Research: Atmospheres*, 123, 6196–6212,
<https://doi.org/10.1029/2017JD028033>, 2018b.

McErlich, C., McDonald, A., Schuddeboom, A., and Silber, I.: Comparing Satellite- and Ground-Based Observations of Cloud Occurrence
825 Over High Southern Latitudes, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020JD033607>, 2021.

McFarquhar, G. M., Bretherton, C. S., Marchand, R., Protat, A., DeMott, P. J., Alexander, S. P., Roberts, G. C., Twohy, C. H., Toohey, D.,
Siems, S., Huang, Y., Wood, R., Rauber, R. M., Lasher-Trapp, S., Jensen, J., Stith, J. L., Mace, J., Um, J., Järvinen, E., Schnaiter, M.,
Gettelman, A., Sanchez, K. J., McCluskey, C. S., Russell, L. M., McCoy, I. L., Atlas, R. L., Bardeen, C. G., Moore, K. A., Hill, T. C. J.,
Humphries, R. S., Keywood, M. D., Ristovski, Z., Cravigan, L., Schofield, R., Fairall, C., Mallet, M. D., Kreidenweis, S. M., Rainwater,
830 B., D'Alessandro, J., Wang, Y., Wu, W., Saliba, G., Levin, E. J. T., Ding, S., Lang, F., Truong, S. C. H., Wolff, C., Haggerty, J., Harvey,
M. J., Klekociuk, A. R., and McDonald, A.: Observations of Clouds, Aerosols, Precipitation, and Surface Radiation over the Southern
Ocean: An Overview of CAPRICORN, MARCUS, MICRE, and SOCRATES, *Bulletin of the American Meteorological Society*, 102,
E894–E928, <https://doi.org/10.1175/BAMS-D-20-0132.1>, 2021.

Müller, D., Ansmann, A., Mattis, I., Tesche, M., Wandinger, U., Althausen, D., and Pisani, G.: Aerosol-Type-Dependent Lidar Ratios Ob-
835 served with Raman Lidar, *Journal of Geophysical Research*, 112, D16 202, <https://doi.org/10.1029/2006JD008292>, 2007.

Murphy, D. M., Froyd, K. D., Bian, H., Brock, C. A., Dibb, J. E., DiGangi, J. P., Diskin, G., Dollner, M., Kupc, A., Scheuer, E. M., Schill, G. P., Weinzierl, B., Williamson, C. J., and Yu, P.: The distribution of sea-salt aerosol in the global troposphere, *Atmospheric Chemistry and Physics*, 19, 4093–4104, <https://doi.org/10.5194/acp-19-4093-2019>, 2019.

Myagkov, A., Seifert, P., Bauer-Pfundstein, M., and Wandinger, U.: Cloud Radar with Hybrid Mode towards Estimation of Shape and 840 Orientation of Ice Crystals, *Atmospheric Measurement Techniques*, 9, 469–489, <https://doi.org/10.5194/amt-9-469-2016>, 2016.

Naud, C. M., Booth, J. F., and Del Genio, A. D.: Evaluation of ERA-interim and MERRA cloudiness in the Southern Ocean, *J. Climate*, 27, 2109–2124, <https://doi.org/10.1175/JCLI-D-13-00432.1>, 2014.

Nisantzi, A., Mamouri, R. E., Ansmann, A., Schuster, G. L., and Hadjimitsis, D. G.: Middle East versus Saharan dust extinction-to-backscatter 845 ratios, *Atmospheric Chemistry and Physics*, 15, 7071–7084, <https://doi.org/10.5194/acp-15-7071-2015>, 2015.

Norgren, M. S., de Boer, G., and Shupe, M. D.: Observed Aerosol Suppression of Cloud Ice in Low-Level Arctic Mixed-Phase Clouds, 850 *Atmospheric Chemistry and Physics*, 18, 13 345–13 361, <https://doi.org/10.5194/acp-18-13345-2018>, 2018.

Ohneiser, K., Ansmann, A., Baars, H., Seifert, P., Barja, B., Jimenez, C., Radenz, M., Teisseire, A., Flouts, A., Haarig, M., Foth, A., Chudnovsky, A., Engelmann, R., Zamorano, F., Bühl, J., and Wandinger, U.: Smoke of extreme Australian bushfires observed in the stratosphere over Punta Arenas, Chile, in January 2020: optical thickness, lidar ratios, and depolarization ratios at 355 and 532 nm, 855 *Atmospheric Chemistry and Physics*, 20, 8003–8015, <https://doi.org/10.5194/acp-20-8003-2020>, 2020.

Pearson, G., Davies, F., and Collier, C.: An Analysis of the Performance of the UFAM Pulsed Doppler Lidar for Observing the Boundary Layer, *Journal of Atmospheric and Oceanic Technology*, 26, 240–250, <https://doi.org/10.1175/2008JTECHA1128.1>, 2009.

Possner, A., Ekman, A. M. L., and Lohmann, U.: Cloud response and feedback processes in stratiform mixed-phase clouds perturbed by ship exhaust, *Geophysical Research Letters*, <https://doi.org/10.1002/2016GL071358>, 2017.

855 Radenz, M. and Bühl, J.: larda cloud sniffer, <https://doi.org/10.5281/zenodo.5574997>, 2021.

Radenz, M., Seifert, P., Baars, H., Flouts, A. A., Yin, Z., and Bühl, J.: Automated Time–Height-Resolved Air Mass Source Attribution for Profiling Remote Sensing Applications, *Atmospheric Chemistry and Physics*, 21, 3015–3033, <https://doi.org/10.5194/acp-21-3015-2021>, 2021.

Rose, T., Crewell, S., Löhnert, U., and Simmer, C.: A Network Suitable Microwave Radiometer for Operational Monitoring of the Cloudy 860 Atmosphere, *Atmospheric Research*, 75, 183–200, <https://doi.org/10.1016/j.atmosres.2004.12.005>, 2005.

Sato, K., Tateno, S., Watanabe, S., and Kawatani, Y.: Gravity Wave Characteristics in the Southern Hemisphere Revealed by a High-Resolution Middle-Atmosphere General Circulation Model, *Journal of the Atmospheric Sciences*, 69, 1378–1396, <https://doi.org/10.1175/JAS-D-11-0101.1>, 2012.

Seifert, A., Köhler, C., and Beheng, K. D.: Aerosol-cloud-precipitation effects over Germany as simulated by a convective-scale numerical 865 weather prediction model, *Atmospheric Chemistry and Physics*, 12, 709–725, <https://doi.org/10.5194/acp-12-709-2012>, 2012.

Seifert, P., Ansmann, A., Mattis, I., Wandinger, U., Tesche, M., Engelmann, R., Müller, D., Pérez, C., and Haustein, K.: Saharan dust and heterogeneous ice formation: Eleven years of cloud observations at a central European EARLINET site, *Journal of Geophysical Research: Atmospheres*, 115, D20 201, <https://doi.org/10.1029/2009JD013222>, 2010.

Seifert, P., Kunz, C., Baars, H., Ansmann, A., Bühl, J., Senf, F., Engelmann, R., Althausen, D., and Artaxo, P.: Seasonal variability 870 of heterogeneous ice formation in stratiform clouds over the Amazon Basin, *Geophysical Research Letters*, 42, 5587–5593, <https://doi.org/10.1002/2015gl064068>, 2015.

Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V.,

875 Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving Our Fundamental Understanding of the
Role of Aerosol-cloud Interactions in the Climate System, *Proceedings of the National Academy of Sciences*, 113, 5781–5790,
<https://doi.org/10.1073/pnas.1514043113>, 2016.

880 Shao, Q., Randall, D. A., Moeng, C.-H., and Dickinson, R. E.: A Method to Determine the Amounts of Cloud-Top Radiative and Evap-
orative Cooling in a Stratocumulus-Topped Boundary Layer, *Quarterly Journal of the Royal Meteorological Society*, 123, 2187–2213,
<https://doi.org/10.1002/qj.49712354403>, 1997.

885 Silber, I., Verlinde, J., Eloranta, E. W., and Cadeddu, M.: Antarctic Cloud Macrophysical, Thermodynamic Phase, and Atmospheric Inversion
Coupling Properties at McMurdo Station: I. Principal Data Processing and Climatology, *Journal of Geophysical Research: Atmospheres*,
123, 6099–6121, <https://doi.org/10.1029/2018JD028279>, 2018.

890 Silber, I., Fridlind, A. M., Verlinde, J., Russell, L. M., and Ackerman, A. S.: Nonturbulent Liquid-Bearing Polar Clouds: Ob-
served Frequency of Occurrence and Simulated Sensitivity to Gravity Waves, *Geophysical Research Letters*, 47, e2020GL087099,
<https://doi.org/10.1029/2020GL087099>, 2020.

Simmel, M., Bühl, J., Ansmann, A., and Tegen, I.: Ice Phase in Altocumulus Clouds over Leipzig: Remote Sensing Observations and Detailed
Modeling, *Atmospheric Chemistry and Physics*, 15, 10 453–10 470, <https://doi.org/10.5194/acp-15-10453-2015>, 2015.

895 Solomon, A., de Boer, G., Creamean, J. M., McComiskey, A., Shupe, M. D., Maahn, M., and Cox, C.: The relative impact of cloud conden-
sation nuclei and ice nucleating particle concentrations on phase partitioning in Arctic mixed-phase stratocumulus clouds, *Atmospheric
Chemistry and Physics*, 18, 17 047–17 059, <https://doi.org/10.5194/acp-18-17047-2018>, 2018.

Tan, I., Storelvmo, T., and Choi, Y.-S.: Spaceborne lidar observations of the ice-nucleating potential of dust, polluted dust, and smoke aerosols
in mixed-phase clouds, *Journal of Geophysical Research: Atmospheres*, 119, 6653–6665, <https://doi.org/10.1002/2013JD021333>, 2014.

Trenberth, K. E. and Fasullo, J. T.: Simulation of Present-Day and Twenty-First-Century Energy Budgets of the Southern Oceans, *Journal of
Climate*, 23, 440–454, <https://doi.org/10.1175/2009JCLI3152.1>, 2010.

895 Uetake, J., Hill, T. C. J., Moore, K. A., DeMott, P. J., Protat, A., and Kreidenweis, S. M.: Airborne Bacteria Confirm the Pris-
tine Nature of the Southern Ocean Boundary Layer, *Proceedings of the National Academy of Sciences*, 117, 13 275–13 282,
<https://doi.org/10.1073/pnas.2000134117>, 2020.

900 Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Browse, J., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Bur-
rows, S. M., Ceburnis, D., DeMott, P. J., Mason, R. H., O'Dowd, C. D., Rinaldi, M., and Carslaw, K. S.: Contribution of feldspar
and marine organic aerosols to global ice nucleating particle concentrations, *Atmospheric Chemistry and Physics*, 17, 3637–3658,
<https://doi.org/10.5194/acp-17-3637-2017>, 2017.

Vignon, É., Alexander, S. P., DeMott, P. J., Sotiropoulou, G., Gerber, F., Hill, T. C. J., Marchand, R., Nenes, A., and Berne, A.: Challenging
and Improving the Simulation of Mid-level Mixed-phase Clouds over the High-latitude Southern Ocean, *Journal of Geophysical Research:
Atmospheres*, <https://doi.org/10.1029/2020JD033490>, 2021.

905 Villanueva, D., Heinold, B., Seifert, P., Deneke, H., Radenz, M., and Tegen, I.: The day-to-day co-variability between mineral dust and cloud
glaciation: a proxy for heterogeneous freezing, *Atmospheric Chemistry and Physics*, 20, 2177–2199, <https://doi.org/10.5194/acp-20-2177-2020>, 2020.

Westbrook, C. D. and Illingworth, A. J.: The Formation of Ice in a Long-Lived Supercooled Layer Cloud, *Quarterly Journal of the Royal
Meteorological Society*, 139, 2209–2221, <https://doi.org/10.1002/qj.2096>, 2013.

910 Wu, P., Dong, X., Xi, B., Tian, J., and Ward, D. M.: Profiles of MBL Cloud and Drizzle Microphysical Properties Retrieved From Ground-Based Observations and Validated by Aircraft In Situ Measurements Over the Azores, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2019JD032205>, 2020.

Yin, Z. and Baars, H.: PollyNET/Pollynet_Processing_Chain: Version 2.1, <https://doi.org/10.5281/zenodo.4694451>, The program has been tested in MATLAB 2014a and 2018a. And it can run on both window 10 and centos 6, which has been fully tested., 2021.

915 Young, I.: Seasonal variability of the global ocean wind and wave climate, *International Journal of Climatology*, 19, 931–950, [https://doi.org/10.1002/\(SICI\)1097-0088\(199907\)19:9<931::AID-JOC412>3.0.CO;2-O](https://doi.org/10.1002/(SICI)1097-0088(199907)19:9<931::AID-JOC412>3.0.CO;2-O), 1999.

Zaremba, T. J., Rauber, R. M., McFarquhar, G. M., Hayman, M., Finlon, J. A., and Stechman, D. M.: Phase Characterization of Cold Sector Southern Ocean Cloud Tops: Results from SOCRATES, *Journal of Geophysical Research: Atmospheres*, n/a, e2020JD033673, <https://doi.org/10.1029/2020JD033673>, e2020JD033673 2020JD033673, 2020.

920 Zhang, D., Wang, Z., Kollias, P., Vogelmann, A. M., Yang, K., and Luo, T.: Ice particle production in mid-level stratiform mixed-phase clouds observed with collocated A-Train measurements, *Atmospheric Chemistry and Physics*, 18, 4317–4327, <https://doi.org/10.5194/acp-18-4317-2018>, 2018.