

The author's responses to the reviewer's comments and the revised manuscript are merged below.

Response to Reviewer #2

Overall, the addition of the PCA results greatly strengthens the results of the manuscript and I find the significant revisions by the authors have ameliorated the gravest of my earlier concerns. Despite these greatly improved state of the study, I have several minor technical comments and suggested typographical edits that I would like to see addressed to clarify language in the paper before it is accepted. Many of the extended new passages seem rather hastily worded. I recommend the authors have a native English speaker or editing service review newly incorporated text.

We appreciate your time for carefully reviewing our manuscript. We would like to thank you for the constructive comments and suggestions, which encourage and help us to improve the manuscript. The manuscript has been revised accordingly, and Dr. Timothy Logan has done a thorough grammar check for the manuscript. In the response below, your comments are provided in black text and our responses are provided in blue text.

Response:

MINOR COMMENTS (L refers to line number)

L31-32: What do you mean by the phrase “underneath CCN and moisture sources?” Do you mean surface sources? Or sources underneath the cloud that aren't the surface?

This phrase is changed to ‘...and the below-cloud CCN and moisture sources’ in the revised manuscript.

L53-55: You need to rephrase the sentence beginning “The enhance Nc conversion...” as it is very difficult to understand. When you say “Nc conversion” do you mean activation of CCN or autoconversion? In addition, what do you mean by “intrusions of CCN?” Intrusions from where? Advected aerosol, stronger sources, from the free troposphere?

This phrase is changed to ‘...enhanced activation of CCN and the cloud droplet condensational growth induced by the higher below-cloud CCN loading’ in the revised manuscript.

L180: Do you mean northeast Atlantic instead of southeast?

Thanks for capturing. This term is changed to ‘northeast Atlantic’ in the revised manuscript.

Section 3.5.1: Can you spell out the implications of the moderate negative correlations between cloud properties and PC1? In my mind, greater TKE should lead to a more adiabatic PBL cloud so it is worth explicitly pointing out that adiabaticity (probably) depends more on decoupling state than turbulence. This is a somewhat counter-intuitive argument, maybe a weakness of your use of sub-cloud TKE vs. full PBL mean including the cloud layer. In addition, I think the low correlations you find between PC2 and cloud properties are likely a consequence of subsetting for single-layer stratocumulus. If you looked at *all* boundary layer clouds at ENA, you would naturally be sorting by LTS and wind direction.

Thanks for the comments and suggestions, we have expanded the discussion in the revised section 3.5.1 as follows:

These negative correlations suggest that under the higher PWV_{BL} condition, the sufficient water vapor supply allows more CCN to become cloud droplets, as previously discussed, and hence increases the cloud adiabaticity due to the dominant condensational growth process. While in the situation of relatively higher TKE_w , the decrease in the N_c and f_{ad} might be partly attributed to the association with the active in-cloud coalescence process and entrainment of dry air. However, owing to the obstacle of retrieving in-cloud TKE_w from the ground-based remote sensing, the usage of sub-cloud TKE_w in this study captures part of the relationship between turbulence and adiabaticity. Therefore, in this situation, the cloud adiabaticity might depend more on PWV_{BL} and the boundary layer decoupling state.

Additionally, the following statement is also added:

These weak correlations might likely be due to the subset of MBL single-layer stratocumulus in this study, as the previous study over the ENA found that the sensitivity of MBL cloud adiabaticity largely depends on the strength of cloud top inversion (which can be partially indicated by the increased LTS) and slightly depends on the boundary layer decoupling (Terai et al., 2019; Zheng et al., 2020).

General comment on sections 3.4-3.5: I wonder if the missing factor that could explain the D–TKE negative correlation is surface fluxes? There is likely a diurnal cycle factor as well, perhaps obtainable through examining net cloud top longwave cooling rate

Thanks for the comments, we have added the discussion in the revised section 3.4 as follows:

Note that the negative correlation between D_i and TKE_w examined here might also be partly attributed to the diurnal cycle of the turbulence, which is studied to be associated with the cloud-top longwave radiative cooling over the ENA, especially for the drizzling clouds (Ghate et al., 2021; Zheng et al., 2016). However, this study focuses on the non-precipitating clouds where the effect of drizzle on the cloud-top radiative cooling driven turbulence is minimum, and examining the cloud-top radiative cooling rate from ground-based remote sensing is beyond the scope of the current study. It would be with interest to get the accurate cloud-top radiative cooling rate using a radiative transfer model to perform further study in the future.

TYPOGRAPHICAL COMMENTS

L129: Instead of “where sits in,” try “which sits in”

It is changed to ‘which sits in’ in the revised manuscript.

L130-132: This sentence is rather colloquial in tone. I suggest removing “So that” at the beginning.

The term ‘So that...’ is removed.

L184: singular, “buoyancy generation and shear”

It is changed to ‘buoyancy generation and shear’ in the revised manuscript.

L186: “growth process” instead of “growing process”

It is changed to ‘growth process’ in the revised manuscript.

L972-977: this is a mega-long sentence and is rather awkwardly worded. Please revisit to simplify and clarify wording.

Thanks for pointing it out. This sentence is changed to ‘From the aircraft in-situ measurements during the ACE-ENA, we found that the observed vertical profile of N_c is near-constant in middle part of the cloud (even in the drizzling cloud where the collision-coalescence processes are more active), and the signal of entrainment-induced N_c depletion is shown near the cloud top (Wu et al., 2020a)’ in the revised manuscript.

L1061: “to have” instead of “to has”

It is changed to ‘to have’ in the revised manuscript.

References.

- Ghate, V. P., Cadeddu, M. P., Zheng, X. and O’Connor, E.: Turbulence in The Marine Boundary Layer and Air Motions Below Stratocumulus Clouds at the ARM Eastern North Atlantic Site, *J. Appl. Meteorol. Climatol.*, doi:10.1175/jamc-d-21-0087.1, 2021.
- Terai, C. R., Zhang, Y., Klein, S. A., Zelinka, M. D., Chiu, J. C. and Min, Q.: Mechanisms Behind the Extratropical Stratiform Low-Cloud Optical Depth Response to Temperature in ARM Site Observations, *J. Geophys. Res. Atmos.*, doi:10.1029/2018JD029359, 2019.
- Zheng, Y., Rosenfeld, D. and Li, Z.: Quantifying cloud base updraft speeds of marine stratocumulus from cloud top radiative cooling, *Geophys. Res. Lett.*, doi:10.1002/2016GL071185, 2016.
- Zheng, Y., Rosenfeld, D. and Li, Z.: A More General Paradigm for Understanding the Decoupling of Stratocumulus-Topped Boundary Layers: The Importance of Horizontal Temperature Advection, *Geophys. Res. Lett.*, doi:10.1029/2020GL087697, 2020.

Revised Version

Environmental Effects on Aerosol-Cloud Interaction in non-precipitating MBL Clouds over the Eastern North Atlantic

Xiaojian Zheng¹, Baike Xi¹, Xiquan Dong¹, Peng Wu², Timothy Logan³ and Yuan Wang^{4,5}

¹Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, AZ, USA

²Pacific Northwest National Laboratory, Richland, WA, USA

³Department of Atmospheric Sciences, Texas A&M University, College Station, TX, USA

⁴Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

⁵Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

Correspondence: Baike Xi (baikex@arizona.edu)

Abstract. Over the eastern north Atlantic (ENA) ocean, a total of 20 non-precipitating single-layer marine boundary layer (MBL) stratus and stratocumulus cloud cases are selected to investigate the impacts of the environmental variables on the aerosol-cloud interaction (ACI_r) using the ground-based measurements from the Department of Energy Atmospheric Radiation Measurement (ARM) facility at the ENA site during 2016 – 2018. The ACI_r represents the relative change of cloud-droplet effective radius r_e with respect to the relative change of cloud condensation nuclei (CCN) number concentration at 0.2% supersaturation ($N_{CCN,0.2\%}$) in the water vapor stratified environment. The ACI_r values vary from -0.01 to 0.22 with increasing sub-cloud boundary layer precipitable water vapor (PWV_{BL}) conditions, indicating that r_e is more sensitive to the CCN loading under sufficient water vapor supply, owing to the combined effect of enhanced condensational growth and coalescence processes associated with higher N_c and PWV_{BL} . The principal component analysis shows that the most pronounced pattern during

the selected cases is the co-variations of the MBL conditions characterized by the vertical component of turbulence kinetic energy (TKE_w), decoupling index (D_i), and PWV_{BL} . The environmental effects on ACI_r emerge after the data are stratified into different TKE_w regimes. The ACI_r values, under both relatively lower and higher PWV_{BL} conditions, increase more than double from the low TKE_w to high TKE_w regime. It can be explained by the fact that stronger boundary layer turbulence maintains a well-mixed MBL, strengthening the connection between cloud microphysical properties and the below-cloud CCN and moisture sources. With sufficient water vapor and low CCN loading, the active coalescence process broadens the cloud droplet size spectra, and consequently results in an enlargement of r_e . The enhanced activation of CCN and the cloud droplet condensational growth induced by the higher below-cloud CCN loading can effectively decrease r_e , which jointly presents as the increased ACI_r . This study examines the importance of environmental effects on the ACI_r assessments and provides observational constraints to future model evaluations on aerosol-cloud interactions.

1. Introduction

Clouds are one of the most important parts of the Earth's climate system. They can impact the global climate by modulating the radiative balance in the atmosphere. Moreover, the radiative effects of cloud adjustments due to aerosols remain one of the largest uncertainties in climate modeling (IPCC, 2013). Over the oceanic area, the lower troposphere is dominated by marine boundary layer (MBL) clouds. MBL clouds can persistently reflect the solar radiation by their long-lasting nature maintained by cloud-top radiative cooling, and therefore act as a major modulator of the Earth's radiative budget (Seinfeld et al., 2016). The climatic importance of MBL cloud radiative properties is primarily induced by cloud microphysical properties such as cloud-droplet number concentration (N_C) and effective radius (r_e), and has been intensively investigated by many researchers (Garrett and Zhao, 2006; Rosenfeld, 2007; Wood et al., 2015; Seinfeld et al., 2016). The ambient aerosol conditions can influence these cloud microphysical properties via the aerosol-cloud interaction (ACI). Compared to the clean regions, clouds under the regions having relatively higher below-cloud aerosol concentrations exhibited smaller cloud droplets (reduced r_e and increased N_C) and enhanced both cloud liquid water contents and optical depths (McComiskey et al., 2009; Chen et al., 2014; Wang et al., 2018). The changes of MBL cloud microphysical properties induced by aerosols have been investigated

from previous studies using in-situ measurements, ground- and satellite-based observations, and model simulations in multiple oceanic areas such as the eastern Pacific and eastern Atlantic (Twohy et al., 2005; Lu et al., 2007; Hill et al., 2009; Costantino and Bréon, 2010; Mann et al., 2014; Dong et al., 2015; Diamond et al., 2018; Yang et al., 2019; Zhao et al., 2019; Wang et al., 2020).

The assessments of ACI, particularly using ground-based remote sensing, vary in terms of the quantitative values, which represent the different cloud susceptibilities to aerosol loadings. Owing to the numerous approaches in assessing the ACI, such as the spatial and temporal scales, N_c and r_e retrieval methods, and more importantly, the different aerosol proxies used in the ACI quantification, different ACI results could be achieved. For example, the studies using total aerosol number concentration and aerosol scattering/extinction coefficients to represent the aerosol loadings would result in relatively lower ACI values (Pandithurai et al., 2009; Liu et al., 2016). This is primarily attributed to the inclusion of aerosol species with different abilities to activate, which is determined by their physicochemical properties, and thus will cause non-negligible uncertainties in capturing the information of aerosol intrusion to the cloud (Feingold et al. 2006; Logan et al., 2014). While some studies found relatively higher ACI values using cloud condensation nuclei (CCN) number concentration (N_{CCN}), presumably due to the fact that CCN represents the portion of aerosols that can be activated and possesses the potential ability to further grow into cloud droplets, this favorably yields a more straightforward assessment of ACI (McComiskey et al., 2009; Qiu et al., 2017; Zheng et al., 2020). It is noteworthy that the ACI variations have been found to have both increasing and decreasing trends in response to changing environmental water availability (Martin et al., 2004; Kim et al., 2008; McComiskey et al., 2009; Pandithurai et al., 2009; Martin et al., 2011; Liu et al., 2016; Zheng et al., 2020). Although these contradicting results have been postulated due to multiple factors such as cloud adiabaticity, condensational growth, collision coalescence, and atmospheric thermodynamics and dynamics, the underlying mechanisms in altering the ACI and causing the uncertainties in the ACI assessments remain unclear. Therefore, further studies are necessary (Fan et al., 2016; Feingold and McComiskey, 2016; Seinfeld et al., 2016).

The Eastern North Atlantic (ENA) is a remote oceanic region that features persistent but diverse subtropical MBL clouds, owing to complex meteorological influences from the semi-permanent Azores High and prevailing large-scale subsidence (Wood et al., 2015). The ENA has become a favorable region to study the aerosol indirect effects on MBL clouds under a

relatively clean environment with occasional intrusions of long-range transport of continental air mass (Logan et al., 2014; Wang et al., 2020). The atmospheric radiation measurement (ARM) program established the ENA permanent observatory site on the northern edge of Graciosa Island, Azores, in 2013, which continuously provides comprehensive measurements of the atmosphere, radiation, cloud, and aerosol from ground-based observation instruments. Owing to the location of the site, which sits in between the boundaries of mid-latitude and subtropical regimes, the ENA is under the mixed influence of diverse meteorological conditions. In terms of the aerosol influence on the cloud properties, the roles of meteorological factors on cloud formation and development are not negligible and hence are being explored in this study. The large-scale thermodynamic variables of the lower troposphere are widely used, such as the lower tropospheric stability (LTS), where the higher LTS values are found to be associated with a relatively shallow and well-mixed marine boundary layer, and are prone to stratiform cloud formations with higher cloud fractions (Klein and Hartmann, 1993; Wood, 2012; Wood and Bretherton, 2006; Yue et al., 2011; Rosenfeld et al., 2019), especially over the subtropical ocean such as the northeast Atlantic. Over the ENA site, the spatial gradient of the LTS has been studied to be associated with the contribution terms of MBL turbulence and the wind directional change (Wu et al., 2017).

In the cloud-topped MBL which is maintained by cloud-top radiative cooling, the buoyancy generation and shear contribute most to the turbulence kinetic energy (TKE) production (Nicholls, 1984; Hogan et al., 2009), where the intensity of turbulence denotes the coupling of MBL clouds to the below-cloud boundary layer. In terms of the cloud droplet growth process, especially in a clean environment with low N_{CCN} below the cloud layer, the cloud droplets at the cloud base experience rapid growth via the diffusion of water vapor, and subsequently enter the regime of active coalescence (Rosenfeld and Woodley, 2003; Martins et al., 2011). The intensive turbulence effectively modulates the cloud droplet growth by strengthening the coalescence process and the cloud cycling (Feingold et al., 1996, 1999; Pawlowska et al., 2006). In particular, the unique topography of Graciosa Island induces an island effect which could cause disturbances in the updraft and hence impact the MBL turbulence, depending on the surface wind directions (Zheng et al., 2016). The environmental effects on the MBL cloud formation and development processes and cloud microphysical properties have been widely implemented and considered in climate modeling (Medeiros and Stevens, 2011; West et al., 2014; Zhang et al., 2016). Thus, it is important to provide observational constraints on the environmental effects. The assessment of ACI from the

ground-based perspective highly relies on the sensitivities of cloud droplet number concentrations and size distribution to the changing of below-cloud CCN loadings. Hence, studying the relationship between the environmental effect and the MBL cloud microphysical responses is a nontrivial task.

In this study, we target the non-precipitating single-layer MBL stratus and stratocumulus clouds during the period between September 2016 and May 2018 and examine the role of thermodynamical and dynamical variables on ACIs. This study aims to advance the understanding of ACI by disentangling the environmental effects and providing observational constraints on quantifying the ACI when modeling aerosol effects on MBL clouds. The ground-based observations and retrievals, and the reanalysis are introduced in section 2. Section 3 describes the aerosol, cloud and meteorological properties, and the variations of cloud microphysical properties under different environmental regimes. Moreover, the ACIs under given water vapor conditions and the roles of environmental effects on ACI are discussed in Section 3. The conclusion of the key findings and the future work are presented in section 4.

2. Data and methods

2.1 Cloud and aerosol properties

The cloud boundaries at the ARM ENA site are primarily determined by the ARM Active Remotely-Sensed Cloud Locations (ARSCL) product, which is a combination of data detected by multiple active remote-sensing instruments, including the Ka-band ARM Zenith Radar (KAZR) and laser ceilometer. The KAZR has an operating frequency at 35 GHz and is sensitive in cloud detection with very minimum attenuation up to the cloud top height (Widener et al., 2012). The temporal and vertical resolutions of KAZR reflectivity are 4 seconds and 30 m, respectively. The ceilometer operates at 910 nm and its attenuated backscatter data can be converted to the cloud base height up to 7.7 km with an uncertainty of ~ 10 m (Morris, 2016). Combining both KAZR and ceilometer measurements, the cloud base (z_b) and top (z_t) heights can be identified accordingly. The single-layer low cloud is defined as having a cloud top height lower than 3 km, with no additional cloud layer in the atmosphere above (Xi et al., 2010).

The cloud microphysical properties are retrieved from a combination of ground-based observations, including KAZR, ceilometer, and microwave radiometer. The detailed retrieval methods and procedures are described in Wu et al. (2020a). The retrieved cloud microphysical

properties, both in time series and vertical profiles, have been validated using the collocated aircraft in-situ measurements during the Aerosol and Cloud Experiments in the Eastern North Atlantic field campaign (ACE-ENA). The retrieval uncertainties are estimated to be ~15% for cloud droplet effective radius (r_e), ~35% for cloud droplet number concentration (N_c), and ~30% for the cloud liquid water content (LWC) (Wu et al., 2020a). Furthermore, the cloud adiabaticity is calculated using the retrieved in-cloud vertical profile of LWC and the adiabatic LWC_{ad}. The LWC_{ad} is given by $\text{LWC}_{\text{ad}(z)} = \Gamma_{\text{ad}}(z - z_b)$, following the method in Wu et al. (2020b), where Γ_{ad} denotes the linear increase of LWC with height under an ideal adiabatic condition (Wood, 2005). The cloud adiabaticity (f_{ad}) is defined as the ratio of LWC to LWC_{ad}.

The surface CCN number concentrations (N_{CCN}) are measured by the CCN-100 (single-column) counter. Since the supersaturation (SS) levels cycle between approximately 0.10% and 1.10% within one hour, N_{CCN} under a relatively stable supersaturation level has to be carefully calculated to rule out the impact of supersaturation on N_{CCN} . This study adopts the interpolation method given by $N_{\text{CCN}} = c\text{SS}^k$ (Twomey, 1959), where parameters c and k are fitted by a power-law function for every periodic cycle. In this study, the supersaturation level of 0.2% is used because it represents typical supersaturation conditions of boundary-layer stratiform clouds (Hudson and Noble, 2013; Logan et al., 2014; Wood et al., 2015; Siebert et al., 2021), and N_{CCN} at 0.2% supersaturation (hereafter $N_{\text{CCN},0.2\%}$) is interpolated to a 5-min temporal resolution.

2.2 Environmental conditions and cloud case selections

The integrated precipitable water vapor (PWV) is obtained from a 3-channel microwave radiometer (MWR3C), which operates at three frequency channels of 23.834, 30, and 89 GHz. The uncertainty of PWV is estimated to be ~0.03 cm (Cadeddu et al., 2013). To capture the information of MBL water vapor more accurately, the sub-cloud boundary layer integrated precipitable water vapor (PWV_{BL}) is calculated using the interpolated sounding product following:

$$\text{PWV}_{\text{BL}} = \frac{1}{\rho_w} \sum (z_{i+1} - z_i) * (\rho_{v,i+1} + \rho_{v,i}) / 2, \quad (1)$$

where the ρ_w is the liquid water density and the ρ_v is the water vapor density collected from the Interpolated Sounding and Gridded Sounding Value-Added Products (Toto and Jensen, 2016),

the subscripts i and $i + 1$ represent the bottom and top of each interpolated sounding height layer. Both PWV and PWV_{BL} are temporally collocated to 5-min resolutions and plotted against each other in Fig. S1a to test the contribution of PWV_{BL} to PWV. The Pearson correlation coefficient of 0.85 shows that the PWV_{BL} are strongly positively correlated with PWV, while the distribution of the percentage ratio of PWV_{BL} to PWV (Fig. S1b) indicates that, on average, PWV_{BL} contributes to ~58% of PWV. Considering the cloud-topped MBL, the majority of cases (~74%) associate with a relatively moist boundary layer compared to the amount of water vapor in the free troposphere, where PWV_{BL} already contributed over 50% of the total column PWV. In contrast, only ~9% of cloud samples occur under a relatively dry boundary layer and moist free troposphere, where PWV_{BL} contributions are less than 40%. In general, PWV can well capture the variation of PWV_{BL} . In the rest of the study, PWV_{BL} are used, as it represents the sub-cloud boundary layer water vapor availability which is more closely related to the MBL cloud processes.

The LTS parameter is used as a proxy of large-scale thermodynamic structure and is defined as the difference between the potential temperature at 700 hPa and surface ($\theta_{700} - \theta_{sfc}$). The LTS values are calculated from European Centre for Medium-Range Weather Forecasts (ECMWF) model outputs of potential temperature, by averaging over a grid box of $0.56^\circ \times 0.56^\circ$ centered at the ENA site. To match the temporal resolutions of the other variables, the original 1-hour LTS data are downsampled to 5-min under the assumption that the large-scale forcing would not have significant changes within an hour.

The boundary layer decoupling condition is represented by the decoupling index (D_i), which is given by $D_i = (z_b - z_{LCL})/z_b$, where the z_{LCL} is the lifting condensation level calculated analytically following the method in Romps (2017), with an uncertainty of around 5 m. The surface temperature, pressure, relative humidity, and mass fraction of water vapor are used in the z_{LCL} calculation, as long as the vector-averaged wind directions (in 360° coordinate) over the ENA site are obtained from the ARM surface meteorology systems (ARM MET handbook, 2011).

As for the boundary layer dynamics, the higher-order moments of vertical velocity are widely used in different model parameterization practices, such as higher-order turbulence closure and probability density function methods (Lappen and Randall, 2001; Zhu and Zuidema, 2009; Ghate et al., 2010). The vertical velocity variance can be used to represent the turbulence

intensity in the below-cloud boundary layer (Feingold et al., 1999). In this study, the vertical component of the turbulence kinetic energy (TKE_w) is used, which is defined as:

$$TKE_w = \frac{1}{2} \overline{(w')^2},$$

(2)

where the $(w')^2$ is the variance of vertical velocity measured from the Doppler lidar standard 10-min integration, which is collected in the Doppler Lidar Vertical Velocity Statistics Value-Added Product (Newson et al., 2019). The noise correction has been applied to reduce the uncertainty of the variance to ~10% (Hogan et al., 2009; Pearson et al., 2009). In this study, the mean value of TKE_w in the sub-cloud boundary layer proportion of the Doppler lidar range is used, and the data temporal resolution is further downscaled to 5-min for temporal collocation purposes.

In this study, the non-precipitating cloud periods are determined when the KAZR reflectivity at the ceilometer-detected cloud base height range does not exceed -37 dBZ (Wu et al., 2015, 2020b), which extensively rules out the wet-scavenging depletion on below-cloud CCN (Wood, 2006) and ensures the accuracy in capturing the below-cloud CCN loadings. Both retrieved cloud microphysical properties and CCN data are available from September 2016 to May 2018 and confine this period in this study.

3. Result and Discussion

3.1 Aerosol, cloud, and meteorological properties of selected cloud cases

A total of 20 non-precipitating cloud cases are selected in this study, with the detailed time periods listed in Table 1, including 1143 samples with temporal resolutions of 5-min, which corresponds to ~95 hours. Among the selected cases, there are three, eight, five, and four cases for Spring, Summer, Fall, and Winter seasons, respectively. MBL clouds often produce precipitation in the form of drizzle (Wood 2012, Wu et al., 2015, 2020b). A recent study of the seasonal variation of the drizzling frequencies (Wu et al., 2020b) showed that the MBL clouds in the cold months (Oct-Mar) have the highest drizzling frequency of the year (~70%), while the clouds in the warm months (Apr-Sept) are found to have a lower chance of drizzling (~45%). Therefore, the selection of a non-precipitating single-layer low cloud case that lasts at least 2

hours is limited, with only 6 cases found in the cold months and 14 cases found during the warm months.

The probability distribution functions (PDFs) of the aerosol and cloud properties, and the environmental conditions for the selected cases are shown in Fig. 1. The PDF of $N_{CCN,0.2\%}$ presents a normal distribution with a mean value of 215 cm^{-3} and median value of 217 cm^{-3} . About 97% of the $N_{CCN,0.2\%}$ samples lie below 350 cm^{-3} and represents a relatively clean environment (Logan et al., 2014, 2018). A few instances of aerosol intrusions ($\sim 3\%$) with higher $N_{CCN,0.2\%}$ were likely a result of continental air mass transport from North America, Europe, and Africa (Logan et al., 2014; Wang et al., 2020). As for the cloud microphysical properties, the cloud-layer mean N_c and r_e (Fig. 1b and 1c) are also both normally distributed with median values close to the mean values. The majority of the N_c values ($\sim 91\%$) are lower than 125 cm^{-3} with a mean value of 86 cm^{-3} , and the r_e distribution peaks between 9 - 11 μm with a mean value of 10.1 μm . Both N_c and r_e values fall in the typical ranges of the non-precipitating MBL cloud characteristics over the ENA site (Dong et al., 2014; Wu et al., 2020b). The distribution of f_{ad} is slightly skewed to the left with a median value of 0.66 (Fig. 1d), indicating that the bulk of cloud samples are close to adiabatic environments, while the left tail denotes a wide range of cloud sub-adiabaticities, which allows us to investigate the role of cloud adiabaticities on the cloud microphysical variations.

For all selected cases, the LTS, which represents the large-scale thermodynamic structure, is distributed bimodally across the range from 14K to 23K with mean and median values of 19.1K in Fig. 1e. A higher LTS magnitude represents a relatively stable environment and is favorable to the formation of marine stratocumulus (Medeiros and Stevens, 2011; Gryspeerdt et al., 2016). Note that the median LTS of 19.1 K in this study is close to the separation threshold of 18.55K suggested by prior studies to distinguish the marine stratocumulus from a global assessment of marine shallow cumulus clouds (Smalley and Rapp, 2020). Therefore, leveraging the demarcation line at 19.1K may allow us to investigate the aerosol-cloud relationships under contrasting thermodynamic regimes. The PDF of D_i parameter spreads widely with a median value of 0.34 for the selected cases (Fig. 1f), which provides an opportunity to study the cloud sample behaviors under MBL conditions range from well-mixed to decoupled. Higher D_i values indicate more decoupled MBL with weaker turbulence which cannot sufficiently maintain the well-mixed MBL, while lower D_i values often associate with stronger turbulence which maintains a coupled MBL (Jones et al., 2011). As an indicator of the

below-cloud boundary layer turbulence, the TKE_w values present a gamma distribution that is highly skewed to the right (Fig. 1e), with a mean value of 0.11 and a median value of $0.08 \text{ m}^2\text{s}^{-2}$. About half of the cloud samples are observed within a relatively less turbulent environment (which is also implied by the higher half of D_i), suggesting weak connections between the cloud layer and the below-cloud boundary layer. The other half of the cloud samples, with relatively higher TKE_w values up to $0.4 \text{ m}^2/\text{s}^2$, imply tighter connections between cloud microphysical properties and below-cloud boundary layer accompanied by intensive turbulent conditions, which is favorable to enhance cloud droplet growth (Albrecht et al., 1995; Hogan et al., 2009; Ghate et al., 2010; West et al., 2014; Ghate and Cadeddu, 2019).

It is noteworthy that PWV_{BL} values exhibit a bimodal distribution with a median value of 1.2 cm (Fig. 1f). About 49% of the samples have their PWV_{BL} values in the range of 0.4 - 1.2 cm with the first peak in 0.6 - 0.8 cm, and 51% of the samples have PWV_{BL} values higher than 1.2 cm with a second peak in 1.6 - 1.8 cm, which may be due to the seasonal difference of the selected cases. Fig. S2 shows the seasonal variation of the PWV_{BL} from 2016 to 2018 when single-layered low clouds are present. The monthly PWV_{BL} values are as low as ~ 0.9 cm and remain nearly invariant from January through March, then increase to ~ 2.0 cm (doubled) in September, and decrease dramatically to the winter months. The selected cloud cases are distributed across the seasons, with $\sim 34\%$ of the samples occurring during the months with the lowest mean PWV_{BL} (Jan-Mar), while $\sim 43\%$ of the samples fall in the highest PWV_{BL} months (Jun-Sept). These two different PWV_{BL} regions will provide a great opportunity for us to further examine the ACI under relatively lower and higher water vapor conditions.

3.2 Dependent of cloud microphysical properties on CCN and PWV_{BL}

Figure 2 shows the cloud microphysical properties as a function of $N_{\text{CCN},0.2\%}$ and PWV_{BL} for the samples from 20 selected cases. As illustrated in Fig. 2a, there is a statistically significant positive correlation ($R^2=0.9$) between $\ln(N_c)$ and $\ln(N_{\text{CCN},0.2\%})$. The linear fit of $\ln(N_c)$ to $\ln(N_{\text{CCN},0.2\%})$ is then mathematically transformed to a power-law fitting function of N_c to $N_{\text{CCN},0.2\%}$, and plotted as dash lines in Fig. 2a. The power-law fitting indicates that 90.3% of the variation in binned $\ln(N_c)$ can be explained by the change in the binned $\ln(N_{\text{CCN},0.2\%})$ and further suggests that with more available below-cloud CCN, higher number concentrations are expected. The logarithmic ratio $\partial \ln(N_c) / \partial \ln(N_{\text{CCN},0.2\%})$ is computed to be 0.435 from our study. This ratio is very close to 0.48 as was shown by McComiskey et al. (2009), who also

used ground-based measurements to study the marine stratus clouds over the California coast. The logarithmic ratio (0.435) is also close to the result (0.458) of Lu et al. (2007) who used aircraft in-situ measured cloud droplet and accumulation mode aerosol number concentration for the marine stratus and stratocumulus clouds over the eastern Pacific Ocean. The ratio reflects the relative conversion efficiency of cloud droplets from the CCN, regardless of the water vapor availability. Theoretically, it has the boundaries of 0 - 1, where the lower bound means no change of N_c with N_{CCN} , and the upper bound indicates a linear relationship that every CCN would result in one cloud droplet. Our result is comparable with the previous studies targeting the MBL stratiform clouds, indicating a certain similarity of the bulk cloud microphysical responses with respect to aerosol intrusion in those types of cloud and over different marine environments, further support that the assessment in this study is valid.

The PWV_{BL} values are represented as blue circles (larger one for higher PWV_{BL}) in Fig. 2a in order to study the role of water vapor availability on the CCN- N_c conversion process. As demonstrated in Fig. 2a, the PWV_{BL} values almost mimic the increasing $N_{CCN,0.2\%}$ trend, which is also governed by the seasonal $N_{CCN,0.2\%}$ and the selected cloud cases. Fig. S3 shows the seasonal variation of $N_{CCN,0.2\%}$ from 2016 to 2018. It is noticeable that the monthly $N_{CCN,0.2\%}$ values, which mimic the monthly variation of PWV_{BL} , are much higher during warm months (May-Oct) than during cold months (Nov-Apr). This seasonal $N_{CCN,0.2\%}$ variation is also found in recent studies of MBL aerosol composition and number concentration. During the warm months, the below-cloud boundary layer is enriched by the accumulation mode of sulfate and organic particles via local generation and long-range transport induced by the semi-permanent Azores High, which are found to be hydrophilic and can be great CCN contributors (Wang et al., 2020; Zawadowicz et al., 2020; Zheng et al., 2018, 2020). Therefore, the coincidence of high $N_{CCN,0.2\%}$ and PWV_{BL} does not necessarily imply a physical relationship, but instead is the result of their similar seasonal trend. The potential co-variabilities between $N_{CCN,0.2\%}$ and PWV_{BL} , and hence the implication on the N_c variation will be further investigated in the latter section. When taking the PWV_{BL} into account, R^2 increases from 0.903 to 0.982, and this new relationship suggests that the co-variability between the binned $\ln(N_{CCN,0.2\%})$ and $\ln(PWV_{BL})$ are in a stronger correlation with the change in binned $\ln(N_c)$. Intuitively, if the CCN- N_c relationship is primarily dominated by the diffusion of water vapor, more CCN and higher PWV_{BL} should result in a continuously increasing of N_c . However, the rapid increase of N_c (37 to 92 cm^{-3}) in the first half of $N_{CCN,0.2\%}$ bins ($<250 \text{ cm}^{-3}$) does not happen in the second half

of the $N_{CCN,0.2\%}$ bins ($>250 \text{ cm}^{-3}$) where the slope of N_c increase ($96 \text{ to } 103 \text{ cm}^{-3}$) appears to be flattened for higher $N_{CCN,0.2\%}$ and PWV_{BL} bins. Furthermore, the joint power-law fitting of N_c (to $N_{CCN,0.2\%}$ and PWV_{BL}) appears to be constantly lower than the single power-law fitting of N_c (to $N_{CCN,0.2\%}$ solely) in each bin. The negative power of PWV_{BL} in this relationship suggests that PWV_{BL} might play a stabilization role in the diffusional growth process, which will be further analyzed in the following sections.

The relationship between r_e and $N_{CCN,0.2\%}$ is shown in Fig. 2b where there is no significant relationship between r_e with $N_{CCN,0.2\%}$ solely, given a near-zero slope and the low correlation coefficient (fitted line not plotted). However, after applying a multiple linear regression to the logarithmic form of r_e , $N_{CCN,0.2\%}$ and PWV_{BL} , a significant correlation among those three variables is found. The r_e is negatively correlated with $N_{CCN,0.2\%}$ and positively correlated with PWV_{BL} , and 73.7% of the variations in binned $\ln(r_e)$ can be explained by the joint changes of the binned $\ln(N_{CCN,0.2\%})$ and $\ln(PWV_{BL})$. This indicates that in the bulk part, r_e decreases with increasing $N_{CCN,0.2\%}$ and enlarges with increasing PWV_{BL} . Notice that in the lower $N_{CCN,0.2\%}$ bins ($<150 \text{ cm}^{-3}$) where the PWV_{BL} values are the lowest among all the bins ($0.76 - 0.85 \text{ cm}$), the limitation of cloud droplet growth by competing for the available water vapor is evident by the changes in N_c and r_e . For example, the $N_{CCN,0.2\%}$ changes from 47 to 128 cm^{-3} , the N_c increases from 37 to 71 cm^{-3} and r_e only increases from 9.30 to $9.74 \text{ }\mu\text{m}$. In other words, nearly tripling the CCN loading leads to roughly doubling N_c , while the r_e is only enlarged by $0.44 \text{ }\mu\text{m}$ (4.7%). In the relatively low available PWV_{BL} regime, it is clear that even with more CCN being converted into cloud droplets, the limited water vapor condition prohibits the further diffusional growth of those cloud droplets. However, in the higher $N_{CCN,0.2\%}$ bins ($>150 \text{ cm}^{-3}$) with relatively higher PWV_{BL} , the binned r_e values fluctuate and decrease with increasing CCN bins under similar PWV_{BL} (i.e., the two $N_{CCN,0.2\%}$ ranges from $200\text{-}400 \text{ cm}^{-3}$, and from $400\text{-}500 \text{ cm}^{-3}$). Since r_e essentially represents the area-weighted information of the cloud droplet size distribution (DSD), this sorting method of r_e inevitably entangles multiple cloud droplet evolution processes and environmental effects that can alter the DSD, especially under the condition of sufficient water supply. Therefore, the further assessment of the r_e responses to the $N_{CCN,0.2\%}$ loading under the constraint of water vapor should be discussed in order to untangle the impacts of different processes and environmental effects on r_e .

3.3 Aerosol-cloud interaction under different water vapor availabilities

As previously discussed above and suggested by earlier studies, the conditions of water vapor supply have a substantial impact on various processes from CCN- N_c conversion to in-cloud droplet condensational growth and coalescence processes, hence effectively altering the cloud DSD (Feingold et al., 2006; McComiskey et al., 2009; Zheng et al., 2020). Moving forward to examine how r_e responds to the changes of $N_{CCN,0.2\%}$ in the context of given water vapor availability, an index describing the aerosol-cloud interaction process is introduced as follows:

$$ACI_r = - \left. \frac{\partial \ln(r_e)}{\partial \ln(N_{CCN,0.2\%})} \right|_{PWV_{BL}}. \quad (3)$$

The ACI_r represents the relative change of r_e with respect to the relative change of $N_{CCN,0.2\%}$, where positive ACI_r denotes the decrease of r_e with increasing $N_{CCN,0.2\%}$ under binned PWV_{BL} . This assessment of ACI_r focuses on the relative sensitivity of the cloud microphysics response in the water vapor stratified environment, while previous studies used the cloud liquid water path (LWP) as the constraint (Twomey, 1977; Feingold et al., 2003; Garrett et al., 2004). LWP describes the liquid water (i.e., existing cloud droplets) physically linked to r_e and N_c which have an interdependent relationship in cloud retrieval procedures, and hence to a certain extent, share co-variabilities with cloud microphysical properties (Dong et al., 1998; Wu et al., 2020a). In this study, by using the PWV as a sorting variable, we are trying to capture the role of ambient available water vapor in the cloud droplet growth process (especially the water vapor diffusional growth), using measurement independent to the cloud retrievals. Fig. 3 shows the variation of ACI_r under different PWV_{BL} bins, and illustrates the calculation of ACI_r in three different PWV_{BL} ranges. Note that in Fig. 3a, the regressions are derived from all points (statistically significant with a confidence level of 95%). As shown in Fig. 3a, the ACI_r values range from close-to-zero values (-0.01) to 0.22, with the mean value of 0.117 ± 0.052 . The ACI_r range of this study agrees well with the previous studies of MBL cloud aerosol-cloud interactions (McComiskey et al., 2009; Pandithurai et al., 2009; Liu et al., 2016). It is noteworthy that the variation of ACI_r with PWV_{BL} suggests two different relationships under separated PWV_{BL} conditions, as discussed in the following two paragraphs.

Under the relatively lower PWV_{BL} condition (<1.2 cm), the low values of ACI_r (-0.01 - 0.057) indicate that r_e is less sensitive to $N_{CCN,0.2\%}$, and the dependence on PWV_{BL} is also insignificant as given by flat regression line (green dashed line) and low correlation coefficient of 0.38 (Fig. 3a). As discussed in section 3.2, the limited water vapor can weaken the ability of condensational growth of the cloud droplet converted from CCN, that is, the increase of CCN loading cannot be effectively reflected by a decrease in r_e . For example, a 307% increase of $N_{CCN,0.2\%}$ only leads to a 10% decrease in r_e in the PWV_{BL} range of 0.8-1.0 cm as shown in Fig. 3b. So that in this regime, even with a slight PWV_{BL} increase, the lack of a sufficient amount of large cloud droplets is favorable to the predominant condensational growth process, which effectively narrows the cloud DSD and, in turn, confines the variable range of r_e with respect to $N_{CCN,0.2\%}$ (Pawlowska et al., 2006; Zheng et al., 2020). In this situation, the ability of CCN to convert to cloud droplets as well as droplet condensational growth are limited by insufficient water vapor, rather than an influx of CCN.

However, under the relatively higher PWV_{BL} regime (>1.2 cm), the ACI_r values become more positive and express a significant increasing trend with PWV_{BL} (correlation coefficient of 0.83, blue dashed line), which indicates that r_e is more susceptible to $N_{CCN,0.2\%}$ in this regime. On the one hand, due to the sufficient water vapor supply, the enhanced condensational growth process allows more CCN to grow into cloud droplets, so that the limiting factor of the droplet growth corresponds to the changes in CCN loading. On the other hand, the increased N_c values associated with higher water vapor supply in the cloud effectively enhance the coalescence process. This results in broadening the cloud DSD and increasing the variation range of r_e in response to the changes of $N_{CCN,0.2\%}$. To test our hypothesis of active coalescence under higher water vapor conditions, Table 2 lists the occurrence frequencies of large r_e values (> 12 and $14 \mu m$) under the six high PWV_{BL} bins ($1.2 - 2.4$ cm), because this range of $12-14 \mu m$ can serve as the critical demarcation of an efficient coalescence process (Gerber, 1996; Freud and Rosenfeld, 2012; Rosenfeld et al., 2012). As listed in Table 2, for the six high PWV_{BL} bins, the occurrence frequencies of $r_e > 12 \mu m$ are 25.0%, 30.6%, 54.1%, 74.2%, 93.8%, and 97.5%, and the occurrence frequencies of $r_e > 14 \mu m$ are 1.25%, 1.77%, 7.4%, 17.7%, 31.9%, and 20.1%, respectively.

The increasing trends of large r_e occurrences mimic the trend of ACI_r and suggest that with increased PWV_{BL} , cloud droplets have a greater chance to grow via the effective coalescence process and subsequently lead to an enlargement of ACI_r . Although previous

studies have brought up the potential impacts of the cloud droplet coalescence process on ACI, it is rarely seen that the relationship among them has been discussed in detail. Here we provide possible explanations on how the enhanced coalescence process can enlarge ACI_r . Quantitatively, ACI_r is described by the logarithmic partial derivative ratio of r_e to $N_{CCN,0.2\%}$, thus a sharper decrease of r_e with respect to a given $N_{CCN,0.2\%}$ range can result in a steeper slope and in turn, larger ACI_r (i.e., a 239% increase in $N_{CCN,0.2\%}$ leads to a r_e decrease of 48% in the 2.2-2.4 cm bin in Fig. 3b). Physically, this relies on how the cloud droplet size distribution (DSD) would change with different CCN loadings. Therefore, particularly in low CCN conditions, sufficient water vapor availability will allow cloud droplets to continuously grow via diffusion of water vapor (i.e., condensational growth), and enter the active cloud-droplet coalescence regime. In contrast, the increase in cloud droplet size can effectively reduce N_c via the process of large cloud droplets collecting small droplets, and small droplets be coalesced into large droplets. Consequently, the cloud DSD becomes effectively broadened toward the large tail by the coalescence, so that r_e is enlarged. With more CCN available, the cloud DSD is narrowed by the enhanced condensational growth and regresses toward the small tail by increasing the amount of newly converted cloud droplets which result in decreased r_e . These interactions between CCNs and cloud droplets ultimately result in the broadened changeable range of r_e , and in turn, the enlarged ACI_r .

In order to investigate the theoretical implication of supersaturation conditions on the aerosol-cloud interaction observed here in the MBL stratiform clouds, the ACI_r values are calculated with respect to the surface N_{CCN} theoretically at two additional high supersaturation levels (0.5% and 1.2%), under all PWV_{BL} conditions. The results in Table 3 show that the ACI_r signals are both weak and do not have significant changes under relatively lower PWV_{BL} conditions, while the ACI_r signals tend to strengthen with the increase of supersaturation under the relatively higher PWV_{BL} . Based on Köhler theory, if the supersaturation exceeds the critical point for the given droplet, the droplet will thus experience continued growth, so theoretically the ACI should increase with the supersaturation under same aerosol number concentration. However, the observed limited water vapor cannot support this ideal droplet growth, results in weak responses of cloud droplets to aerosol intrusion. With the increase of observed water vapor, the continued growth of cloud droplets becomes more plausible, hence the high supersaturation yields larger droplets with low number of aerosols, more efficient droplet activation with a large number of aerosols, and in turns, larger ACI_r (even out of the theoretical bounds). However, considering these high supersaturation environments are unphysical in the

observed MBL cloud layers, and estimating the real supersaturation conditions using ground-based remote-sensing is beyond the scope of this study, we chose the supersaturation level of 0.2% because it represents the most typical supersaturation conditions of MBL stratiform clouds.

3.4 The co-variabilities of the meteorological factors

The environmental conditions over the ENA have been widely studied as not independent but entangled with each other (Wood et al., 2015; Zheng et al., 2016; Wu et al., 2017; Wang et al., 2021). To better understand the dependencies and the co-variabilities of the meteorological factors, a principal component analysis (PCA) is performed comprising the following variables: (1) PWV_{BL} denotes the water vapor availability within the boundary layer; (2) D_i describes the boundary layer coupling conditions; (3) TKE_w represents the strength of boundary layer turbulence; (4) $W_{dir,NS}$ reflects the surface wind directions in terms of northerly and southerly; and (5) LTS infers the large-scale thermodynamic structures. Note that the $W_{dir,NS}$ are taken as $W_{dir,NS} = abs(W_{dir} - 180^\circ)$, so that the original W_{dir} ($0-360^\circ$) can be transformed to $W_{dir,NS}$ ($0-180^\circ$) where the values smaller than 90° are close to the southerly wind, and those greater than 90° are close to the northerly wind. The $W_{dir,ns}$ are transformed as such to capture the island effects better, because the cliff is located north of the ENA site.

The input data metric of the PCA is constructed from the above five variables, thus the principal components (PCs) that explaining the variations of those dependent variables can be output from the eigenanalysis. The result shows that for the five selected meteorological factors, the proportions of the total intervariable variance explained by the PCs are 43.72%, 22.01%, 18.26%, 8.95% and 7.06%, and the eigenvalues are 2.19, 1.10, 0.91, 0.45, and 0.35, respectively. Note that the first three PCs have the highest eigenvalues and explain most ($\sim 84\%$) of the total variance, which indicates that they can capture the significant variation patterns of the selective meteorological factors.

To determine the relative contributions of the variables to PCs, all the five selected meteorological variables are projected to the first three PCs and the Pearson correlation coefficients between them are listed in Table 4. For the first PC (PC1) which accounts for the highest proportion (43.72%) of the total variance, the PC1 is strongly negatively correlated with PWV_{BL} (-0.84) and D_i (-0.73), but strongly positively correlated with TKE_w (0.69). These

results suggest that PC1 mainly represents the boundary layer conditions, and the co-variations of the boundary layer water vapor and turbulence are the most distinct environmental patterns for the selected cloud cases. The PC2 and PC3 are most correlated with LTS (0.58 and 0.65 for PC2 and PC3, respectively) and $W_{dir,NS}$ (0.60 and -0.50 for PC2 and PC3, respectively), indicating that the PC2 and PC3 mainly describe the variations in large-scale thermodynamic and the surface wind patterns, which are likely associated with the variations of the Azores High position and strength (Wood et al., 2015).

To further understand the correlations between the meteorological variables, the principal component loadings plot is constructed by projecting the variables onto PC1 and PC2 as shown in Fig. 4. Each point denotes the variable correlations with PC1 (x-coordinate) and PC2 (y-coordinate), so that each vector represents the strength and direction of the original variable influences on the pair of PCs. The angle between the two vectors represents the correlation between each other. In Fig. 4, both TKE_w and $W_{dir,NS}$ vectors are located in the same quadrant (positive in both PC1 and PC2) and close to each other with a small degree of an acute angle, which means the TKE_w are strongly correlated with the $W_{dir,NS}$. When the surface wind is coming from the north side of the island, the topographic lifting effect of the cliff would induce additional updraft over the ENA site (Zheng et al., 2016), so that the wind closer to the northerly wind (larger $W_{dir,NS}$) is more correlated with higher TKE_w . Note that TKE_w and D_i vectors are almost in an opposite direction, which denotes a strongly negative correlation between the two variables. The angles of PWV_{BL} with D_i ($\sim 45^\circ$) and TKE_w ($\sim 142^\circ$) suggest that PWV_{BL} is moderately positively correlated with D_i but negatively correlated with TKE_w . A higher D_i indicates a more decoupled MBL, where MBL is not well-mixed and separated into a radiative-driven layer and a surface flux driven layer that caps the surface moisture (Jones et al., 2011). This situation is more likely to be associated with a relatively higher PWV_{BL} and weaker TKE_w condition. Note that the negative correlation between D_i and TKE_w examined here might also be partly attributed to the diurnal cycle of the turbulence, which is studied to be associated with the cloud-top longwave radiative cooling over the ENA, especially for the drizzling clouds (Ghate et al., 2021; Zheng et al., 2016). However, this study focuses on the non-precipitating clouds where the effect of drizzle on the cloud-top radiative cooling driven turbulence is minimum, and examining the cloud-top radiative cooling rate from ground-based remote sensing is beyond the scope of the current study. It would be with interest to get the accurate cloud-top radiative cooling rate using a radiative transfer model to perform further study in the future. As for the LTS parameter, the close to 90° angle with TKE_w

suggests no correlation between them, since the LTS is mostly capturing the large-scale thermodynamical structures and is obtained from a coarser temporal resolution. Thus, the LTS does not essentially have correspondence to the strength of boundary layer turbulence and can be treated as independent to TKE_w over the ENA site. The loading plot intuitively tells us the directions and strengths of the co-variabilities of the selected meteorological variables, and sheds the light on determining the key factors that are feasible to use in examining the environmental impacts on the aerosol-cloud interactions.

3.5 Linking the meteorological factors to aerosol-cloud interaction

3.5.1 Relations of meteorological factors with aerosol and cloud properties

The PCs are, mathematically, the linear combination of the selected variables, and hence independent of each other after the PCA. Therefore, treating the aerosol and cloud properties as dependents and correlated with the PCs allows us to infer their co-variation with the meteorological factors statistically. A weakly negative correlation between $N_{CCN,0.2\%}$ and PC1 ($R_{PC1,CCN} = -0.35$) suggests that the relatively higher $N_{CCN,0.2\%}$ could be sometimes found under higher PWV_{BL} and lower TKE_w . Though the correlation is low, the plausible contributions could come from the seasonal variations of $N_{CCN,0.2\%}$ and PWV_{BL} as discussed in the previous section, and the weaker TKE_w might prevent the vertical mixing of CCN and induce higher surface $N_{CCN,0.2\%}$. On the other hand, a weakly positive correlation between $N_{CCN,0.2\%}$ and PC2 ($R_{PC2,CCN} = 0.21$) suggests that there are no fundamental relationships between CCN with thermodynamic and the surface wind direction, and they are not the key controlling factor of surface $N_{CCN,0.2\%}$ variation because the surface CCN concentration is primarily contributed by the accumulation-mode aerosols which come from the condensational growth of Aitken-mode aerosols (Zheng et al., 2018). As for the cloud properties, both N_c and f_{ad} are negatively correlated with PC1 ($R_{PC1,N_c} = -0.51$ and $R_{PC1,f_{ad}} = -0.62$, respectively), suggesting a moderate relationship between N_c , f_{ad} , and the boundary layer condition. These negative correlations suggest that under the higher PWV_{BL} condition, the sufficient water vapor supply allows more CCN to become cloud droplets, as previously discussed, and hence increases the cloud adiabaticity due to the dominant condensational growth process. While in the situation of relatively higher TKE_w , the decrease in the N_c and f_{ad} might be partly

attributed to the association with the active in-cloud coalescence process and entrainment of dry air. However, owing to the obstacle of retrieving in-cloud TKE_w from the ground-based remote sensing, the usage of sub-cloud TKE_w in this study captures part of the relationship between turbulence and adiabaticity. Therefore, in this situation, the cloud adiabaticity might depend more on PWV_{BL} and the boundary layer decoupling state. Moreover, their low correlations with PC2 ($R_{PC2,N_c} = -0.10$ and $R_{PC2,f_{ad}} = -0.17$, respectively) indicate very weak relations with the large-scale thermodynamic variables. These weak correlations might likely be due to the subset of MBL single-layer stratocumulus in this study, as the previous study over the ENA found that the sensitivity of MBL cloud adiabaticity largely depends on the strength of cloud top inversion (which can be partially indicated by the increased LTS) and slightly depends on the boundary layer decoupling (Terai et al., 2019; Zheng et al., 2020). Note that the same sign of correlations with PC1 statistically infer the similar directional co-variation of $N_{CCN,0.2\%}$, N_c , and f_{ad} to a certain extent.

To examine the physical relation between $N_{CCN,0.2\%}$, N_c and f_{ad} , the profiles of cloud r_e and LWC are plotted in normalized height from cloud base (z_b) to cloud top height (z_t) (Fig. 5), which is given by $z_n = (z - z_b) / (z_t - z_b)$. The solid lines denote the mean values, and the shaded area represents one standard deviation at each normalized height z_n . The normalized r_e increases from $\sim 8.6 \mu\text{m}$ at the cloud base toward $\sim 11 \mu\text{m}$ near the upper part of the cloud where z_n is 0.7 (Fig. 5a), through condensational growth and coalescence processes, and then decreases toward the cloud top due to cloud-top entrainment. Similar in-cloud vertical variation of r_e is also found by previous study using aircraft in-situ measurements (Zhao et al., 2018; Wu et al. 2020a). Profiles of retrieved LWC and calculated adiabatic LWC_{ad} (blue line) are presented in Fig. 5b. As demonstrated in Fig. 5b, the f_{ad} values, which is the ratio of LWC to LWC_{ad} , reach a maximum of 0.8 at the cloud base and a minimum of 0.38 at the cloud top. The shaded areas of r_e and LWC denote the range from near-adiabatic to sub-adiabatic cloud environments, where in the near-adiabatic cloud (higher f_{ad}) the cloud droplets experience adiabatic growth and LWC should be close to LWC_{ad} . In contrast, in the sub-adiabatic cloud regime, the decrease of f_{ad} is largely due to cloud-top entrainment and coalescence processes even in non-precipitating MBL clouds (Wood, 2012; Braun et al., 2018; Wu et al. 2020b). Furthermore, to understand the implication of cloud adiabaticity with respect to $\text{CCN}-N_c$ conversion, all of the f_{ad} samples are separated into two groups by the median value of the layer-mean f_{ad} (0.66) for further analysis.

Figure 6 shows N_c against the binned $N_{CCN,0.2\%}$ for the near-adiabatic regime ($f_{ad} > 0.66$) and sub-adiabatic regime ($f_{ad} < 0.66$). For the near-adiabatic regime, N_c increases from $\sim 60 \text{ cm}^{-3}$ to 119 cm^{-3} with increased $N_{CCN,0.2\%}$ and PWV_{BL} , and both $N_{CCN,0.2\%}$ and PWV_{BL} appear to play positive roles in terms of the N_c increase. The result is as expected because the process of condensational growth is predominant in the near-adiabatic clouds, that is, with increasing water vapor supply, the higher CCN loading can effectively lead to more cloud droplets. However, in the sub-adiabatic cloud regime, N_c increases with increased $N_{CCN,0.2\%}$ but possesses a negative correlation with PWV , which results in a slower increase of N_c under higher $N_{CCN,0.2\%}$ and PWV_{BL} conditions. The mean reduction of N_c in the sub-adiabatic regime is computed to be $\sim 37\%$ compared to that for the near-adiabatic clouds. As previously studied, the coalescence process contributes significantly to N_c depletion, even in a non-precipitating MBL clouds (Feingold et al., 1996; Wood, 2006). Thus, lower N_c in the sub-adiabatic regime may be partly due to the combined effect of coalescence and entrainment (Wood, 2006; Hill et al., 2009; Yum et al., 2015; Wang et al., 2020). Note that the retrieved N_c represents the cloud layer-mean information. In summary, the Wu et al. (2020a) retrieval works to separate the reflectivity into the contributions of cloud (Z_c) and drizzle. The retrieval assumes an initial guess of the representative layer-mean N_c based on the climatology over ENA sites (Dong et al., 2014), and such allows the first guess of the vertical profile of LWC based on N_c and Z_c , and then constrains the N_c and LWC using the LWP derived from MWR, and finally output r_e values (Fig. 3 in Wu et al., 2020a). Therefore, the final retrieved N_c is updated to in response to the cloud microphysical processes within this time-step. From the aircraft in-situ measurements during the ACE-ENA, we found that the observed vertical profile of N_c is near-constant in the middle part of the cloud (even in the drizzling cloud where the collision-coalescence processes are more active), and the signal of entrainment-induced N_c depletion is shown near the cloud top (Wu et al., 2020a). However, it is difficult and beyond the scope of the ground-based retrieval to compare the vertical dependency of depletion rate within one time-step. Therefore, as the retrieval currently works to represent the layer-mean information from the given time-step, the preferred method in this study is to compare N_c at different times, where in this case are the adiabatic versus sub-adiabatic conditions which hence yields different N_c that we retrieved from the ground-based snapshot perspective. From the PCA and binning analysis, the effect of cloud adiabaticities on CCN- N_c conversions may shed light on interpreting the aerosol-cloud interaction under different environmental effects.

3.5.2 The role of meteorological factors on ACI_r assessment

Since ACI_r can only be calculated by the logarithmic derivatives from a set of $N_{CCN,0.2\%}$ and r_e data within a certain regime, it will be inappropriate to linearly correlate the data with PCs directly, in both mathematical and physical perspectives. Therefore, the meteorological factors which have the strongest influence on the most explanatory PCs, namely PWV_{BL} and TKE_w are selected to be the sorting variables in assessing the environmental impacts on the ACI_r . In addition, LTS is also selected as it represents the large-scale thermodynamic factor and is independent to the boundary-layer environment conditions. The data samples are first separated into two regimes using the median values of the targeting factors, and then separated into four quadrants by the median PWV_{BL} because ACI_r is found to have significant differences under different water vapor availabilities. The ACI_r values are further calculated for all quadrants to examine whether the ACI_r can be distinguished by the targeting factors.

Combining LTS and PWV_{BL} as sorting variables, the ACI_r values for four regimes are shown in Fig. S4. The ACI_r differences between low and high PWV_{BL} regimes are still retained. In the low PWV_{BL} regime, the ACI_r values are limited to 0.016 and 0.056 for low and high LTS regimes, respectively. In the high PWV_{BL} regime, the ACI_r values are 0.150 and 0.171 for low and high LTS regimes, respectively, which is about 3-5 times greater than those in low PWV_{BL} regime. However, the ACI_r in different LTS regimes cannot be distinctly differentiated (ACI_r differences between LTS regimes are ~ 0.02 and ~ 0.04), and the main difference in ACI_r are still induced by the PWV_{BL} . Owing to the location of the ENA site where it locates near the boundary of mid-latitude and subtropical climate regimes, the MBL clouds over the ENA are found to be often under the influences of cold fronts associated with mid-latitude cyclones, where the cloud evolutions are subject to the combine effects of post-frontal and large-scale subsidence (Wood et al., 2015; Zheng et al., 2020; Wang et al., 2021). Therefore, over the ENA, although the spatial gradient of LTS is studied to be associated with the production of MBL turbulence and the change in wind direction (Wu et al., 2017), the LTS value itself is examined to have a weak impact on the aerosol-cloud interaction from this study.

The TKE_w has been found to be strongly positively correlated with $W_{dir,NS}$ and negatively correlated with D_i from the PCA, that is, the values of TKE_w already account for the co-variabilities in these variables. Therefore, treating TKE_w as the sorting variable would lead to a more physical process-orientated assessment. Accordingly, to examine the role of the

dynamical factors on ACI_r , the samples are separated into four regimes demarcated by the median values of PWV_{BL} and TKE_w (Fig. 7), and the mean values of D_i and f_{ad} in the four quadrants are also displayed in Fig. 7. The effect of PWV_{BL} on ACI_r is demonstrated by the mean ACI_r values where they are much higher in the high PWV_{BL} regime than those in the low PWV_{BL} regime no matter what the TKE_w regimes. Furthermore, the result illustrates that TKE_w does play an important role in ACI_r , because the ACI_r values in the high TKE_w regime are more than double than the values in the low TKE_w regime.

In the regimes of high TKE_w and PWV_{BL} , which are closely associated with coupled MBL ($D_i = 0.21$) and more sub-adiabatic cloud conditions ($f_{ad} = 0.52$), r_e is highly sensitive to CCN loading with the highest ACI_r of 0.259. The sufficient water vapor availability allows CCN to be converted into cloud droplets more effectively, while the relatively higher TKE_w indicates stronger turbulence in the below-cloud boundary layer and maintains a nearly well-mixed MBL. The CCN and moisture below-cloud layer are efficiently transported and mixed aloft via the ascending branch of the eddies (Nicholls, 1984; Hogan et al., 2009), hence are effectively connected to the cloud layer. Therefore, under the lower CCN loading condition, the active coalescence process (which indicated by the low f_{ad} values) results in the depletion of small cloud droplets and broadening of cloud DSD (Chandrakar et al., 2016), and in turn, leads to further enlarged r_e . However, with higher CCN intrusion into the cloud layer, the enhanced cloud droplet conversion and the subsequential condensational growth behave contradictorily to narrow the DSD (Pinsky and Khain, 2002; Pawlowska et al., 2006), which leads to decreased r_e . Therefore, the MBL clouds are distinctly susceptible to CCN loading under the environments of sufficient water vapor and strong turbulence in which the ACI_r is enlarged.

Under high PWV_{BL} but low TKE_w conditions, the mean ACI_r reduces to 0.101 (~ 39% of that under high TKE_w). The MBL is more likely decoupled where $D_i = 0.54$, which indicates that the weaker turbulence loosens the connection between the cloud layer and the underlying boundary layer. This results in a less effective conversion of CCN into cloud droplets, while the more adiabatic cloud environment ($f_{ad} = 0.75$) denotes the lack of coalescence growths and thus diminishes the r_e sensitivity to CCN. Although the constraints of insufficient water vapor on ACI_r are still evident, the ACI_r values increase from 0.008 in the low TKE_w regime to 0.024 in the high TKE_w regime. The ACI_r differences between the two TKE_w regimes attest that ACI_r strongly depends on the connection between the cloud layer and

the below-cloud boundary layer CCN and moisture, that is, stronger turbulence can enhance the susceptibility of r_e to CCN.

In this study, the relationship between turbulence and ACI is found to be valid in non-precipitating MBL clouds. Theoretically, the effect of turbulence on ACI_r would appear to be artificially amplified, if in the presence of precipitation. The intensive turbulence can enhance the coalescence process and accelerate the CCN-cloud cycling, and subsequently, the CCN depletion due to precipitation and coalescence scavenging would result in quantitatively enlarged ACI_r (Feingold et al., 1996, 1999; Duong et al., 2011; Braun et al., 2018). Though it is beyond the scope of this study, it would be of interest to perform such analysis on the aerosol-cloud-precipitation interaction using ground-based remote sensing and model simulations in a future study.

4. Summaries and Conclusions

Over the ARM-ENA site, a total of 20 non-precipitating single-layered MBL stratus and stratocumulus cloud cases have been selected in order to investigate the aerosol-cloud interaction (ACI). The distributions of CCN and cloud properties for selected cases represent the typical characteristics of non-precipitating MBL clouds in a relatively clean environment over the remote oceanic area. The diversity of boundary layer conditions and cloud adiabaticities among the selected cases enable the investigation of different environmental effects on ACI.

The overall variations of N_c with $N_{CCN,0.2\%}$ show an increasing trend, regardless of the water vapor condition, while the sufficient PWV_{BL} appears to stabilize the CCN- N_c conversion process. The water vapor limitation on cloud droplet growth is evident in the lower $N_{CCN,0.2\%}$ up to 150 cm^{-3} with low PWV_{BL} values, where a near tripling of CCN loading leads to a near doubling of N_c but only 4.7% increase in r_e . When $N_{CCN,0.2\%}$ is greater than 250 cm^{-3} and PWV_{BL} values are also relatively high, r_e appears to decrease with increasing $N_{CCN,0.2\%}$ under similar water vapor conditions. As for bulk aerosol-cloud interaction, the ACI_r values vary from -0.01 to 0.22 for different PWV_{BL} conditions where ACI_r appears to be diminished under limited water vapor availability due to limited droplet activation and condensational growth processes. While under relatively sufficient water supply conditions, r_e shows more sensitive

responses to the changes of $N_{CCN,0.2\%}$, due to the combined effect of condensational growth and coalescence processes accompanying the higher N_c and PWV_{BL} .

The theoretical diagram describing the mechanism proposed above is shown in Fig. 8. Under the relatively lower PWV_{BL} condition, the limited water vapor weakens the ability of condensational growth of the cloud droplet converted from CCN, which results in both less newly converted as well as large cloud droplets, with the lack of chance of coalescence processes under this circumstance. Therefore, the variable range of r_e versus $N_{CCN,0.2\%}$ is narrowed and presented as small ACI_r . While under the relatively higher PWV_{BL} condition, particularly in low CCN conditions, the sufficient water vapor availability allows cloud droplets growing via the condensation of water vapor, and thus enter the active cloud-droplet coalescence regime. In contrast, the increase in cloud droplet size can effectively reduce N_c via the coalescence process and the size distributions are effectively broadened toward the large tail by the coalescence, so that r_e is enlarged. Under a higher $N_{CCN,0.2\%}$ intrusion, the cloud droplet size distribution is narrowed by the enhanced condensational growth and regresses toward the small tail by increasing the amount of newly converted cloud droplets which results in decreased r_e . Combinedly, the interactions between CCNs and cloud droplet growth processes ultimately result in a broadened changeable range of r_e , and in turn, the enlarged ACI_r .

The co-variabilities among the environmental factors are examined using the multi-dimensional PCA. The variables of PWV_{BL} , D_i , TKE_w , LTS and $W_{dir,NS}$ are constructed as the input of the eigenanalysis. Results show that the first three PCs can describe the majority (~84%) of the variance among the selected variables. The most explanatory PC1 (account for 43.72% contribution) strongly correlated with PWV_{BL} , D_i (both negatively) and TKE_w (positively), and hence describe the co-variation of the boundary layer conditions. While the PC2 and PC3 (account for 22.01% and 18.26% contributions, respectively) are strongly correlated with LTS and $W_{dir,NS}$, which likely indicates the variations of the Azores High position and strength. By projecting the variables onto PC1 and PC2, the PCA loading analysis shows that TKE_w is strongly negatively correlated with D_i , which is what we expected. A decoupled MBL cloud is often separated into two layers where the lower one can cap the surface moisture, while the higher TKE_w denote sufficient turbulence that maintains the well-mixed MBL. Additionally, the island effect is also indicated by the eigenanalysis, where surface northerly wind would induce additional updraft velocity and hence disturb TKE_w , owing to the effect of the cliff north

of the ENA site. The role of cloud adiabaticities on the behaviors of CCN- N_c conversion is examined using both binning and eigenanalysis. In a near-adiabatic cloud vertical structure, the cloud droplet growth process is dominated by condensational growth, thus the N_c responses to increased $N_{CCN,0.2\%}$ and PWV_{BL} are strengthened. When the cloud layer becomes more sub-adiabatic, the effect of coalescence leads to the depletion of N_c and thus results in the lower retrieved N_c from a ground-based snapshot perspective. The competition between the condensational growth and coalescence processes strongly impacts the variations of cloud microphysics to CCN loading.

To investigate the environmental effects on ACI_r , the factors having the most influence on the explanatory PCs are selected as the sorting variables in the ACI_r assessments. The LTS sorting method cannot distinguish the ACI_r values, which means the LTS values themselves have a weak impact on ACI_r due to the MBL cloud cover over the ENA is mainly impacted by the mid-latitude cyclone systems. In contrast, the intensity of boundary layer turbulence represented by TKE_w plays a more important role in ACI_r , since the values of TKE_w already account for the co-variations of the MBL conditions, and hence leads to a physical process-orientated assessment. The ACI_r assessments in four different TKE_w and PWV_{BL} regimes show that the constraints of insufficient water vapor on the ACI_r are still evident, but in both PWV_{BL} regimes the ACI_r values increase more than double from low TKE_w to high TKE_w regimes. Noticeably, the ACI_r increases from 0.101 in the low TKE_w regime to 0.259 in the high TKE_w regime, under high PWV_{BL} conditions. The intensive below-cloud boundary layer turbulence strengthens the connection between the cloud layer and below-cloud CCN and moisture. So that with sufficient water vapor, an active coalescence leads to further enlarged r_e , particularly for low CCN loading conditions, while the enhanced N_c from condensational growth induced by increased $N_{CCN,0.2\%}$ can effectively decrease r_e . Combining these processes together, the enlarged ACI_r is presented.

In this study, the non-precipitating MBL clouds are found to be most susceptible to the below-cloud CCN loading under environments with sufficient water vapor and stronger turbulence. This study examines the importance of the environmental effects on the ACI_r assessments, and provides the observational constraints to the future model evaluations on the aerosol-cloud interactions. Future studies will be focusing on exploring the role of environmental effects on the aerosol-cloud-precipitation interactions in MBL stratocumulus through an integrative analysis of observations and model simulations.

Data availability. Data used in this study can be accessed from the DOE ARM's Data Discovery at <https://adc.arm.gov/discovery/>

Author contributions. The original idea of this study is discussed by XZ, BX, and XD. XZ performed the analyses and wrote the manuscript. XZ, BX, XD, PW, YW and TL participated in further scientific discussions and provided substantial comments and edits on the paper.

Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue “Marine aerosols, trace gases, and clouds over the North Atlantic (ACP/AMT inter-journal SI)”. It is not associated with a conference.

Acknowledgments. The ground-based measurements were obtained from the Atmospheric Radiation Measurement (ARM) Program sponsored by the U.S. Department of Energy (DOE) Office of Energy Research, Office of Health and Environmental Research, and Environmental Sciences Division. The reanalysis data were obtained from the ECMWF model output, which provides explicitly for the analysis at the ARM ENA site. The data can be downloaded from <https://adc.arm.gov/discovery/>. This work was supported by the NSF grants AGS-1700728/1700727 and AGS-2031750/2031751, and was also supported as part of the “Enabling Aerosol-cloud interactions at GLobal convection-permitting scales (EAGLES)” project (74358), funded by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Earth System Modeling program with the subcontract to the University of Arizona. The Pacific Northwest National Laboratory is operated for the Department of Energy by Battelle Memorial Institute under Contract DE-AC05-76 RL01830. And a special thanks to coeditor Dr. Hang Su and three anonymous reviewers for the constructive comments and suggestions, which helped to improve the manuscript.

References.

- Albrecht, B. A., Bretherton, C. S., Johnson, D., Schubert, W. H. and Frisch, A. S.: The Atlantic Stratocumulus Transition Experiment - ASTEX, Bull. - Am. Meteorol. Soc., doi:10.1175/1520-0477(1995)076<0889:TASTE>2.0.CO;2, 1995.
- ARM MET Handbook: ARM Surface Meteorology Systems (MET) Handbook, DOE ARM Climate Research Facility, DOE/SC-ARM/TR-086. Available at: https://www.arm.gov/publications/tech_reports/handbooks/met_handbook.pdf, last access: 21 August 2021.
- Braun, R. A., Dadashazar, H., MacDonald, A. B., Crosbie, E., Jonsson, H. H., Woods, R. K., Flagan, R. C., Seinfeld, J. H. and Sorooshian, A.: Cloud Adiabaticity and Its Relationship to Marine Stratocumulus Characteristics Over the Northeast Pacific Ocean, J. Geophys. Res. Atmos., doi:10.1029/2018JD029287, 2018.
- Cadeddu, M. P., Liljegren, J. C. and Turner, D. D.: The atmospheric radiation measurement (ARM) program network of microwave radiometers: Instrumentation, data, and retrievals, Atmos. Meas. Tech., doi:10.5194/amt-6-2359-2013, 2013.
- Chandrakar, K. K., Cantrell, W., Chang, K., Ciochetto, D., Niedermeier, D., Ovchinnikov, M., Shaw, R. A. and Yang, F.: Aerosol indirect effect from turbulence-induced broadening of cloud-droplet size distributions, Proc. Natl. Acad. Sci. U. S. A., doi:10.1073/pnas.1612686113, 2016.
- Chen, Y. C., Christensen, M. W., Stephens, G. L. and Seinfeld, J. H.: Satellite-based estimate of global aerosol-cloud radiative forcing by marine warm clouds, Nat. Geosci., doi:10.1038/ngeo2214, 2014.
- Costantino, L. and Bréon, F. M.: Analysis of aerosol-cloud interaction from multi-sensor satellite observations, Geophys. Res. Lett., doi:10.1029/2009GL041828, 2010.

- Diamond, M. S., Dobracki, A., Freitag, S., Griswold, J. D. S., Heikkila, A., Howell, S. G., Kacarab, M. E., Podolske, J. R., Saide, P. E. and Wood, R.: Time-dependent entrainment of smoke presents an observational challenge for assessing aerosol-cloud interactions over the southeast Atlantic Ocean, *Atmos. Chem. Phys.*, doi:10.5194/acp-18-14623-2018, 2018.
- Dong, X., Xi, B., Kennedy, A., Minnis, P. and Wood, R.: A 19-month record of marine aerosol-cloud-radiation properties derived from DOE ARM mobile facility deployment at the Azores. Part I: Cloud fraction and single-layered MBL cloud properties, *J. Clim.*, doi:10.1175/JCLI-D-13-00553.1, 2014.
- Dong, X., Schwantes, A. C., Xi, B. and Wu, P.: Investigation of the marine boundary layer cloud and CCN properties under coupled and decoupled conditions over the azores, *J. Geophys. Res.*, doi:10.1002/2014JD022939, 2015.
- Duong, H. T., Sorooshian, A. and Feingold, G.: Investigating potential biases in observed and modeled metrics of aerosol-cloud-precipitation interactions, *Atmos. Chem. Phys.*, doi:10.5194/acp-11-4027-2011, 2011.
- Fan, J., Wang, Y., Rosenfeld, D., Liu, X.: Review of Aerosol-Cloud Interactions: Mechanisms, Significance and Challenges, *J. Atmo. Sci.* 73(11), 4221-4252, 2016.
- Feingold, G., Kreidenweis, S. M., Stevens, B. and Cotton, W. R.: Numerical simulations of stratocumulus processing of cloud condensation nuclei through collision-coalescence, *J. Geophys. Res. Atmos.*, doi:10.1029/96jd01552, 1996.
- Feingold, G., Frisch, A. S., Stevens, B. and Cotton, W. R.: On the relationship among cloud turbulence, droplet formation and drizzle as viewed by Doppler radar, microwave radiometer and lidar, *J. Geophys. Res. Atmos.*, doi:10.1029/1999JD900482, 1999.

- Feingold, G., Furrer, R., Pilewskie, P., Remer, L. A., Min, Q. and Jonsson, H.: Aerosol indirect effect studies at Southern Great Plains during the May 2003 Intensive Operations Period, *J. Geophys. Res. Atmos.*, doi:10.1029/2004JD005648, 2006.
- Feingold, G. and McComiskey, A.: ARM's Aerosol–Cloud–Precipitation Research (Aerosol Indirect Effects), *Meteorol. Monogr.*, doi:10.1175/amsmonographs-d-15-0022.1, 2016.
- Freud, E. and Rosenfeld, D.: Linear relation between convective cloud drop number concentration and depth for rain initiation, *J. Geophys. Res. Atmos.*, doi:10.1029/2011JD016457, 2012.
- Garrett, T. J. and Zhao, C.: Increased Arctic cloud longwave emissivity associated with pollution from mid-latitudes, *Nature*, doi:10.1038/nature04636, 2006.
- Garrett, T. J., Zhao, C., Dong, X., Mace, G. G. and Hobbs, P. V.: Effects of varying aerosol regimes on low-level Arctic stratus, *Geophys. Res. Lett.*, doi:10.1029/2004GL019928, 2004.
- Gerber, H.: Microphysics of marine stratocumulus clouds with two drizzle modes, *J. Atmos. Sci.*, doi:10.1175/1520-0469(1996)053<1649:MOMSCW>2.0.CO;2, 1996.
- Ghate, V. P., Albrecht, B. A. and Kollias, P.: Vertical velocity structure of nonprecipitating continental boundary layer stratocumulus clouds, *J. Geophys. Res. Atmos.*, doi:10.1029/2009JD013091, 2010.
- Ghate, V. P. and Cadeddu, M. P.: Drizzle and Turbulence Below Closed Cellular Marine Stratocumulus Clouds, *J. Geophys. Res. Atmos.*, doi:10.1029/2018JD030141, 2019.
- Ghate, V. P., Cadeddu, M. P., Zheng, X. and O'Connor, E.: Turbulence in The Marine Boundary Layer and Air Motions Below Stratocumulus Clouds at the ARM Eastern North Atlantic Site, *J. Appl. Meteorol. Climatol.*, doi:10.1175/jamc-d-21-0087.1, 2021.

- Gryspeerd, E., Quaas, J. and Bellouin, N.: Constraining the aerosol influence on cloud fraction, *J. Geophys. Res.*, doi:10.1002/2015JD023744, 2016.
- Hill, A. A., Feingold, G. and Jiang, H.: The influence of entrainment and mixing assumption on aerosol–cloud interactions in marine stratocumulus, *J. Atmos. Sci.*, doi:10.1175/2008JAS2909.1, 2009.
- Hogan, R. J., Grant, A. L. M., Illingworth, A. J., Pearson, G. N. and O’Connor, E. J.: Vertical velocity variance and skewness in clear and cloud-topped boundary layers as revealed by Doppler lidar, *Q. J. R. Meteorol. Soc.*, doi:10.1002/qj.413, 2009.
- Hudson, J. G. and Noble, S.: CCN and Vertical Velocity Influences on Droplet Concentrations and Supersaturations in Clean and Polluted Stratus Clouds, *J. Atmos. Sci.*, doi:10.1175/jas-d-13-086.1, 2013.
- Jones, C. R., Bretherton, C. S., and Leon, D.: Coupled vs. decoupled boundary layers in VOCALS-REx, *Atmos. Chem. Phys.*, 11, 7143–7153, <https://doi.org/10.5194/acp-11-7143-2011>, 2011.
- Klein, S. A. and Hartmann, D. L.: The seasonal cycle of low stratiform clouds, *J. Clim.*, doi:10.1175/1520-0442(1993)006<1587:TSCOLS>2.0.CO;2, 1993.
- Kim, B. G., Miller, M. A., Schwartz, S. E., Liu, Y. and Min, Q.: The role of adiabaticity in the aerosol first indirect effect, *J. Geophys. Res. Atmos.*, doi:10.1029/2007JD008961, 2008.
- Liu, J., Li, Z. and Cribb, M.: Response of marine boundary layer cloud properties to aerosol perturbations associated with meteorological conditions from the 19-month AMF-Azores campaign, *J. Atmos. Sci.*, doi:10.1175/JAS-D-15-0364.1, 2016.

- Lappen, C. L. and Randall, D. A.: Toward a unified parameterization of the boundary layer and moist convection. Part I: A new type of mass-flux model, *J. Atmos. Sci.*, doi:10.1175/1520-0469(2001)058<2021:TAUPOT>2.0.CO;2, 2001.
- Logan, T., Xi, B. and Dong, X.: Aerosol properties and their influences on marine boundary layer cloud condensation nuclei at the ARM mobile facility over the Azores, *J. Geophys. Res.*, doi:10.1002/2013JD021288, 2014.
- Logan, T., Dong, X. and Xi, B.: Aerosol properties and their impacts on surface CCN at the ARM Southern Great Plains site during the 2011 Midlatitude Continental Convective Clouds Experiment, *Adv. Atmos. Sci.*, doi:10.1007/s00376-017-7033-2, 2018.
- Lu, M. L., Conant, W. C., Jonsson, H. H., Varutbangkul, V., Flagan, R. C. and Seinfeld, J. H.: The marine stratus/stratocumulus experiment (MASE): Aerosol-cloud relationships in marine stratocumulus, *J. Geophys. Res.*, doi:10.1029/2006JD007985, 2007.
- Mann, J. A., Christine Chiu, J., Hogan, R. J., O'Connor, E. J., L'Ecuyer, T. S., Stein, T. H. and Jefferson, A.: Aerosol impacts on drizzle properties in warm clouds from ARM Mobile Facility maritime and continental deployments, *J. Geophys. Res.*, doi:10.1002/2013JD021339, 2014.
- Martin, G. M., Johnson, D. W. and Spice, A.: The Measurement and Parameterization of Effective Radius of Droplets in Warm Stratocumulus Clouds, *J. Atmos. Sci.*, doi:10.1175/1520-0469(1994)051<1823:tmapoe>2.0.co;2, 1994.
- Martins, J. V., Marshak, A., Remer, L. A., Rosenfeld, D., Kaufman, Y. J., Fernandez-Borda, R., Koren, I., Correia, A. L., Zubko, V. and Artaxo, P.: Remote sensing the vertical profile of cloud droplet effective radius, thermodynamic phase, and temperature, *Atmos. Chem. Phys.*, doi:10.5194/acp-11-9485-2011, 2011.

- McComiskey, A, Feingold, G., Frisch, A. S., Turner, D. D., Miller, M., Chiu, J. C., Min, Q. and Ogren, J.: An assessment of aerosol-cloud interactions in marine stratus clouds based on surface remote sensing, *J. Geophys. Res.*, 114, D09203, doi:10.1029/2008JD011006, 2009.
- McComiskey, A. and Feingold, G.: The scale problem in quantifying aerosol indirect effects, *Atmos. Chem. Phys.*, doi:10.5194/acp-12-1031-2012, 2012.
- Medeiros, B. and Stevens, B.: Revealing differences in GCM representations of low clouds, *Clim. Dyn.*, doi:10.1007/s00382-009-0694-5, 2011.
- Morris, V. R.: Ceilometer Instrument Handbook, DOE ARM Climate Research Facility, DOE/SC-ARM-TR-020, 2016. Available at: https://www.arm.gov/publications/tech_reports/handbooks/ceil_handbook.pdf, last access: 23 April 2021.
- Newsom, R. K., Sivaraman, C., Shippert, T.R. and Riihimaki, L. D.: Doppler Lidar Vertical Velocity Statistics Value-Added Product. DOE ARM Climate Research Facility, DOE/SC-ARM/TR-149, 2019. Available at: https://www.arm.gov/publications/tech_reports/doe-sc-arm-tr-149.pdf, last access: 2 September 2021.
- Nicholls, S.: The dynamics of stratocumulus: Aircraft observations and comparisons with a mixed layer model, *Q. J. R. Meteorol. Soc.*, doi:10.1002/qj.49711046603, 1984.
- Pandithurai, G., Takamura, T., Yamaguchi, J., Miyagi, K., Takano, T., Ishizaka, Y., Dipu, S. and Shimizu, A.: Aerosol effect on cloud droplet size as monitored from surface-based remote sensing over East China Sea region, *Geophys. Res. Lett.*, doi:10.1029/2009GL038451, 2009.
- Pawlowska, H., Grabowski, W. W. and Brenguier, J. L.: Observations of the width of cloud droplet spectra in stratocumulus, *Geophys. Res. Lett.*, doi:10.1029/2006GL026841, 2006.

- Pearson, G., Davies, F. and Collier, C.: An analysis of the performance of the UFAM pulsed Doppler lidar for observing the boundary layer, *J. Atmos. Ocean. Technol.*, doi:10.1175/2008JTECHA1128.1, 2009.
- Pinsky, M. B. and Khain, A. P.: Effects of in-cloud nucleation and turbulence on droplet spectrum formation in cumulus clouds, *Q. J. R. Meteorol. Soc.*, doi:10.1256/003590002321042072, 2002.
- Qiu, Y., Zhao, C., Guo, J. and Li, J.: 8-Year ground-based observational analysis about the seasonal variation of the aerosol-cloud droplet effective radius relationship at SGP site, *Atmos. Environ.*, doi:10.1016/j.atmosenv.2017.06.002, 2017.
- Romps, D. M.: Exact expression for the lifting condensation level, *J. Atmos. Sci.*, doi:10.1175/JAS-D-17-0102.1, 2017.
- Rosenfeld, D. and Woodley, W. L.: Closing the 50-year circle: From cloud seeding to space and back to climate change through precipitation physics. Chapter 6 of “Cloud Systems, Hurricanes, and the Tropical Rainfall Measuring Mission (TRMM)”, edited by: Tao, W.-K. and Adler, R. F., *Meteor. Monogr.*, 51, 234 pp., 59–80, AMS, 2003.
- Rosenfeld, D.: Aerosol-Cloud Interactions Control of Earth Radiation and Latent Heat Release Budgets, in *Solar Variability and Planetary Climates.*, 2007.
- Rosenfeld, D., Wang, H. and Rasch, P. J.: The roles of cloud drop effective radius and LWP in determining rain properties in marine stratocumulus, *Geophys. Res. Lett.*, doi:10.1029/2012GL052028, 2012.
- Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T. and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science* (80-.), doi:10.1126/science.aav0566, 2019.

Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G. and Wood, R.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, *Proc. Natl. Acad. Sci. U. S. A.*, doi:10.1073/pnas.1514043113, 2016.

Siebert, H., Szodry, K.-E., Egerer, U., Wehner, B., Henning, S., Chevalier, K., Lückerrath, J., Welz, O., Weinhold, K., Lauermann, F., Gottschalk, M., Ehrlich, A., Wendisch, M., Fialho, P., Roberts, G., Allwayin, N., Schum, S., Shaw, R. A., Mazzoleni, C., Mazzoleni, L., Nowak, J. L., Malinowski, S. P., Karpinska, K., Kumala, W., Czyzewska, D., Luke, E. P., Kollias, P., Wood, R. and Mellado, J. P.: Observations of Aerosol, Cloud, Turbulence, and Radiation Properties at the Top of the Marine Boundary Layer over the Eastern North Atlantic Ocean: The ACORES Campaign, *Bull. Am. Meteorol. Soc.*, doi:10.1175/bams-d-19-0191.1, 2021.

Terai, C. R., Zhang, Y., Klein, S. A., Zelinka, M. D., Chiu, J. C. and Min, Q.: Mechanisms Behind the Extratropical Stratiform Low-Cloud Optical Depth Response to Temperature in ARM Site Observations, *J. Geophys. Res. Atmos.*, doi:10.1029/2018JD029359, 2019.

Thorsen, T. J. and Fu, Q.: Automated retrieval of cloud and aerosol properties from the ARM Raman Lidar. Part II: Extinction, *J. Atmos. Ocean. Technol.*, doi:10.1175/JTECH-D-14-00178.1, 2015.

Toto, T. and Jensen, M.: Interpolated Sounding and Gridded Sounding Value-Added Products. DOE ARM Climate Research Facility, DOE/SC-ARM-TR-183, 2016. Available at: https://www.arm.gov/publications/tech_reports/doe-sc-arm-tr-183.pdf, last access: 2 September 2021.

- Twohy, C. H., Petters, M. D., Snider, J. R., Stevens, B., Tahnk, W., Wetzel, M., Russell, L. and Burnet, F.: Evaluation of the aerosol indirect effect in marine stratocumulus clouds: Droplet number, size, liquid water path, and radiative impact, *J. Geophys. Res. D Atmos.*, doi:10.1029/2004JD005116, 2005.
- Twomey, S.: The nuclei of natural cloud formation part II: The supersaturation in natural clouds and the variation of cloud droplet concentration, *Geofis. Pura e Appl.*, doi:10.1007/BF01993560, 1959.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *J. Atmos. Sci.*, doi:10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2, 1977.
- Wang, Y., Jiang, J.H., Su, H., Choi, S., Huang, L., Guo, J., and Yung, Y. L.: Elucidating the Role of Anthropogenic Aerosols In Arctic Sea Ice Variations, *J. Climate* 31(1), 99-114, 2018.
- Wang, Y., Zheng, X., Dong, X., Xi, B., Wu, P., Logan, T., and Yung, Y. L.: Impacts of long-range transport of aerosols on marine-boundary-layer clouds in the eastern North Atlantic, *Atmos. Chem. Phys.*, 20, 14741–14755, <https://doi.org/10.5194/acp-20-14741-2020>, 2020.
- West, R. E. L., Stier, P., Jones, A., Johnson, C. E., Mann, G. W., Bellouin, N., Partridge, D. G. and Kipling, Z.: The importance of vertical velocity variability for estimates of the indirect aerosol effects, *Atmos. Chem. Phys.*, doi:10.5194/acp-14-6369-2014, 2014.
- Widener, K, Bharadwaj, N, and Johnson, K: Ka-Band ARM Zenith Radar (KAZR) Instrument Handbook. DOE ARM Climate Research Facility, DOE/SC-ARM/TR-106, 2012. Available at: https://www.arm.gov/publications/tech_reports/handbooks/kazr_handbook.pdf, last access: 23 April 2021.

- Wood, R.: Rate of loss of cloud droplets by coalescence in warm clouds, *J. Geophys. Res. Atmos.*, doi:10.1029/2006JD007553, 2006.
- Wood, R. and Bretherton, C. S.: On the relationship between stratiform low cloud cover and lower-tropospheric stability, *J. Clim.*, doi:10.1175/JCLI3988.1, 2006.
- Wood, R.: Stratocumulus clouds, *Mon. Weather Rev.*, doi:10.1175/MWR-D-11-00121.1, 2012.
- Wood, R., Wyant, M., Bretherton, C. S., Rémillard, J., Kollias, P., Fletcher, J., Stemmler, J., De Szoeko, S., Yuter, S., Miller, M., Mechem, D., Tselioudis, G., Chiu, J. C., Mann, J. A. L., O'Connor, E. J., Hogan, R. J., Dong, X., Miller, M., Ghate, V., Jefferson, A., Min, Q., Minnis, P., Palikonda, R., Albrecht, B., Luke, E., Hannay, C. and Lin, Y.: Clouds, aerosols, and precipitation in the marine boundary layer: An arm mobile facility deployment, *Bull. Am. Meteorol. Soc.*, doi:10.1175/BAMS-D-13-00180.1, 2015.
- Wu, P., Dong, X. and Xi, B.: Marine boundary layer drizzle properties and their impact on cloud property retrieval, *Atmos. Meas. Tech.*, doi:10.5194/amt-8-3555-2015, 2015.
- Wu, P., Dong, X., Xi, B., Liu, Y., Thieman, M. and Minnis, P.: Effects of environment forcing on marine boundary layer cloud-drizzle processes, *J. Geophys. Res.*, doi:10.1002/2016JD026326, 2017.
- Wu, P., Dong, X., Xi, B., Tian, J. and Ward, D. M.: Profiles of MBL Cloud and Drizzle Microphysical Properties Retrieved From Ground-Based Observations and Validated by Aircraft In Situ Measurements Over the Azores, *J. Geophys. Res. Atmos.*, doi:10.1029/2019JD032205, 2020a.
- Wu, P., Dong, X. and Xi, B.: A climatology of marine boundary layer cloud and drizzle properties derived from ground-based observations over the azores, *J. Clim.*, doi:10.1175/JCLI-D-20-0272.1, 2020b.

- Xi, B., Dong, X., Minnis, P. and Khaiyer, M. M.: A 10 year climatology of cloud fraction and vertical distribution derived from both surface and GOES observations over the DOE ARM SPG site, *J. Geophys. Res. Atmos.*, doi:10.1029/2009JD012800, 2010.
- Yang, Y., Zhao, C., Dong, X., Fan, G., Zhou, Y., Wang, Y., Zhao, L., Lv, F. and Yan, F.: Toward understanding the process-level impacts of aerosols on microphysical properties of shallow cumulus cloud using aircraft observations, *Atmos. Res.*, doi:10.1016/j.atmosres.2019.01.027, 2019.
- Yue, Q., Kahn, B. H., Fetzer, E. J. and Teixeira, J.: Relationship between marine boundary layer clouds and lower tropospheric stability observed by AIRS, CloudSat, and CALIOP, *J. Geophys. Res. Atmos.*, doi:10.1029/2011JD016136, 2011.
- Yum, S. S., Wang, J., Liu, Y., Senum, G., Springston, S., McGraw, R. and Yeom, J. M.: Cloud microphysical relationships and their implication on entrainment and mixing mechanism for the stratocumulus clouds measured during the VOCALS project, *J. Geophys. Res.*, doi:10.1002/2014JD022802, 2015.
- Zhang, S., Wang, M., J. Ghan, S., Ding, A., Wang, H., Zhang, K., Neubauer, D., Lohmann, U., Ferrachat, S., Takeamura, T., Gettelman, A., Morrison, H., Lee, Y., T. Shindell, D., G. Partridge, D., Stier, P., Kipling, Z. and Fu, C.: On the characteristics of aerosol indirect effect based on dynamic regimes in global climate models, *Atmos. Chem. Phys.*, doi:10.5194/acp-16-2765-2016, 2016.
- Zhao, C., Qiu, Y., Dong, X., Wang, Z., Peng, Y., Li, B., Wu, Z. and Wang, Y.: Negative Aerosol-Cloud re Relationship From Aircraft Observations Over Hebei, China, *Earth Sp. Sci.*, doi:10.1002/2017EA000346, 2018.
- Zhao, C., Zhao, L. and Dong, X.: A case study of stratus cloud properties using in situ aircraft observations over Huanghua, China, *Atmosphere (Basel)*, doi:10.3390/atmos10010019, 2019.

- Zawadowicz, M. A., Suski, K., Liu, J., Pekour, M., Fast, J., Mei, F., Sedlacek, A., Springston, S., Wang, Y., Zaveri, R. A., Wood, R., Wang, J., and Shilling, J. E.: Aircraft measurements of aerosol and trace gas chemistry in the Eastern North Atlantic, *Atmos. Chem. Phys. Discuss.* [preprint], <https://doi.org/10.5194/acp-2020-887>, in review, 2020.
- Zheng, G., Wang, Y., Aiken, A. C., Gallo, F., Jensen, M. P., Kollias, P., Kuang, C., Luke, E., Springston, S., Uin, J., Wood, R., and Wang, J.: Marine boundary layer aerosol in the eastern North Atlantic: seasonal variations and key controlling processes, *Atmos. Chem. Phys.*, 18, 17615–17635, <https://doi.org/10.5194/acp-18-17615-2018>, 2018.
- Zheng, G., Kuang, C., Uin, J., Watson, T., and Wang, J.: Large contribution of organics to condensational growth and formation of cloud condensation nuclei (CCN) in the remote marine boundary layer, *Atmos. Chem. Phys.*, 20, 12515–12525, <https://doi.org/10.5194/acp-20-12515-2020>, 2020.
- Zheng, X., Xi, B., Dong, X., Logan, T., Wang, Y. and Wu, P.: Investigation of aerosol-cloud interactions under different absorptive aerosol regimes using Atmospheric Radiation Measurement (ARM) southern Great Plains (SGP) ground-based measurements, *Atmos. Chem. Phys.*, doi:10.5194/acp-20-3483-2020, 2020.
- Zheng, Y., Rosenfeld, D. and Li, Z.: Quantifying cloud base updraft speeds of marine stratocumulus from cloud top radiative cooling, *Geophys. Res. Lett.*, doi:10.1002/2016GL071185, 2016.
- Zheng, Y., Rosenfeld, D. and Li, Z.: A More General Paradigm for Understanding the Decoupling of Stratocumulus-Topped Boundary Layers: The Importance of Horizontal Temperature Advection, *Geophys. Res. Lett.*, doi:10.1029/2020GL087697, 2020.
- Zhu, P. and Zuidema, P.: On the use of PDF schemes to parameterize sub-grid clouds, *Geophys. Res. Lett.*, doi:10.1029/2008GL036817, 2009.

Table 1. Dates and time periods of selected non-precipitating MBL cloud periods

Case No.	Start Date	Start UTC	End Date	End UTC	Valid Samples
1	20160915	2200	20160916	0020	24
2	20170219	2110	20170220	0520	87
3	20170222	0830	20170222	1200	38
4	20170605	1430	20170605	1900	54
5	20170616	1230	20170616	1510	32
6	20170617	0320	20170617	0520	24
7	20170627	0020	20170627	0250	28
8	20170630	0530	20170630	0930	42
9	20170630	1400	20170630	1700	34
10	20170706	0140	20170706	0900	62
11	20170707	0130	20170707	1000	91
12	20170910	2100	20170911	0600	94
13	20170911	1930	20170911	2150	24
14	20170912	0820	20170912	1100	32
15	20171006	2110	20171006	2320	26
16	20180130	1030	20180131	0500	152
17	20180203	1930	20180204	0500	72
18	20180324	0210	20180324	0600	46
19	20180508	0730	20180508	1110	42
20	20180513	2130	20180514	1200	139

Table 2. Occurrence frequencies of large in-cloud r_e * under relatively high PWV conditions

PWV (cm)	1.2- 1.4	1.4- 1.6	1.6- 1.8	2.8- 2.0	2.0- 2.2	2.2- 2.4
$r_e > 12 \mu\text{m}$ (%)	25.0	30.6	54.1	74.2	93.8	97.5
$r_e > 14 \mu\text{m}$ (%)	1.25	1.77	7.4	17.7	31.9	20.1

*The occurrence of large r_e is defined when the r_e is found to be larger than $12 \mu\text{m}$ or $14 \mu\text{m}$ using the retrieved in-cloud vertical profiles.

Table 3. ACI_r calculated with respect to N_{CCN} theoretically at different supersaturation levels, under all PWV_{BL} conditions

PWV_{BL} (cm)	0.4-0.6	0.6-0.8	0.8-1.0	1.0-1.2	1.2-1.4	1.4-1.6	1.6-1.8	1.8-2.0	2.0-2.2	2.2-2.4
ACI_r										
$(N_{CCN}@0.2\%SS)$	0.020	0.057	0.002	-0.014	0.108	0.076	0.145	0.151	0.221	0.175
$(N_{CCN}@0.5\%SS)$	0.023	0.057	0.0002	0.024	0.129	0.121	0.309	0.136	0.293	0.159
$(N_{CCN}@1.2\%SS)$	0.023	0.045	0.002	0.072	0.125	0.123	0.323	0.175	0.347	0.186

Table 4. The first three principal components from eigenanalysis

Eigenanalysis	PC1	PC2	PC3
Eigenvalues	2.17	1.10	0.91
Proportion of variance explained (%)	43.72	22.01	18.26
Cumulative proportion (%)	43.72	65.73	83.99
Correlations (Variables vs. PCs)	PC1	PC2	PC3
PWV _{BL}	-0.84	0.20	-0.11
D _i	-0.73	-0.48	-0.20
TKE _W	0.69	0.35	-0.44
W _{dir,ns}	0.52	0.60	-0.50
LTS	-0.43	0.58	0.65




Figure 1. Probability distribution functions (PDFs), mean, standard deviation and median values of aerosol, cloud, and meteorological properties for 20 selected non-precipitating cloud cases at the DOE ENA site during the period 2016-2018. (a) Cloud condensation nuclei (CCN) number concentration at 0.2% supersaturation ($N_{CCN,0.2\%}$); (b) cloud-droplet number concentration (N_c); (c) cloud-droplet effective radius (r_e); (d) cloud adiabaticity (f_{ad}); (e) lower tropospheric stability (LTS); (f) decoupling index (D_i); (g) mean vertical component of turbulence kinetic energy (TKE_w); and (h) sub-cloud boundary-layer precipitable water vapor (PWV_{BL}).




Figure 2. (a) N_c and (b) r_e as a function of $N_{CCN,0.2\%}$ (x-axis) and PWV (blue filled circles) for all selected samples. The larger blue circles represent relatively higher PWV values. Whiskers denote one standard deviation for each bin.




Figure 3. (a) Relationship of ACI_r (dots) to binned PWV_{BL} . Whiskers denote one standard deviation for each bin. Linear regressions are performed in relatively low PWV_{BL} regime (< 1.4 cm, green) and high PWV_{BL} regime (> 1.4 cm); and (b) illustration of ACI_r derived from r_e to $N_{CCN,0.2\%}$ in following three PWV_{BL} bins: 0.8-1.0 cm (green), 1.2-1.4 cm (purple), 2.2-2.4 cm (blue). The ACI_r represents the relative change of r_e with respect to the relative change of $N_{CCN,0.2\%}$, where positive ACI_r denotes the decrease of r_e with increased $N_{CCN,0.2\%}$ under binned PWV .




Figure 4. The projections of TKE_w (purple), $W_{dir,NS}$ (red), LTS (orange), PWV_{BL} (blue) and D_i (green) onto the first principal component (PC1) and the second principal component (PC2). The x-coordinates denote variables' correlations with PC1, and the y-coordinates denote variables' correlations with PC2.




Figure 5. Normalized in-cloud vertical profiles of retrieved (a) r_e and (b) LWC (black) and calculated adiabatic LWC_{ad} (blue) for all selected cloud cases, 0 is cloud base and 1 is cloud top. Solid dotted lines denote mean values and shaded areas denote one standard deviation at each height.




Figure 6. N_c as a function of $N_{CCN,0.2\%}$ (x-axis) and PWV (dots) for high adiabaticity f_{ad} (red) and low f_{ad} (black) regimes. The larger circles represent relatively higher PWV values. Whiskers denote one standard deviation for each bin.




Figure 7. ACI_r derived from r_e to $N_{\text{CCN},0.2\%}$ for (a) low TKE_w and (b) high TKE_w regimes. Samples in the low PWV regime are plotted in green, and samples in the high PWV regime are plotted in blue. The mean values of D_i and f_{ad} are displayed for each quadrant with the corresponding color-coded.




Figure 8. Theoretical mechanism of the responses of cloud droplet size distributions to different CCN intrusion, under relative insufficient (low PWV_{BL}) versus sufficient (high PWV_{BL}) water vapor availabilities.