

Anthropogenic Aerosol effects on Tropospheric Circulation and Sea Surface Temperature (1980-2020): Separating the role of Zonally Asymmetric Forcings

5 Chenrui Diao ¹, Yangyang Xu ^{1,*}, Shang-Ping Xie ²

¹ Department of Atmospheric Sciences, Texas A&M University, College Station, Texas 77843, USA
² Scripps Institute of Oceanography, University of California, San Diego, La Jolla, CA 92093, USA

10 *Correspondence to:* Yangyang Xu (yangyang.xu@tamu.edu)

1

Deleted: 1

Abstract. Anthropogenic Aerosols (AA) induce global and regional tropospheric circulation adjustments due to the radiative energy perturbations. The overall cooling effects of AA, which mask a portion of global warming, have been the subject of many studies but still have large uncertainty. The interhemispheric contrast in AA forcing has also been demonstrated to induce a major shift in atmospheric circulation. The zonally heterogeneous changes in AA emissions since the late 20th century, with a notable decline in the Western Hemisphere and continuous increase in the Eastern Hemisphere, received less attention. Here we utilize four sets of single-model initial-condition large-ensemble simulations with various combinations of external forcings to quantify the different radiative and circulation responses due to aerosol emissions changes during 1980-2020. In particular, we focus on the distinct climate responses to Fossil-Fuel (FF) related aerosols from Western Hemisphere (WH) versus Eastern Hemisphere (EH).

Deleted: since the pre-industrial (PI) era, to

Deleted: with

Deleted: remaining

The zonal and meridional redistribution of FF aerosols from WH to EH results in negative radiative forcing over Asia and positive radiative forcing over North America and Europe. This leads to a counter-clockwise anomaly of zonal mean stream function over the tropics (i.e. a northward shift of Hadley cell) to balance the interhemispheric difference in radiative forcing. The redistribution of FF aerosol also induces equatorward shift of the Northern Hemisphere (NH) jet stream, consistent with the thermal wind argument of surface air temperature (SAT) gradient. Furthermore, the consistent relationship between the jet stream shift and the Top-of-Atmosphere net solar flux (FSNTOA) gradient suggests that the latter can be considered as a rule-of-thumb indicator.

Deleted: counterclockwise

Deleted: and stronger

Deleted: with the gradient

Deleted: predictive metric.

Two sets of regional FF simulations (Fix_EastFF1920 and Fix_WestFF1920) are performed to separate the roles of East versus West aerosol forcings, which had clearly opposite trends in the last 40 years. We find that the WH aerosol reduction dominated the simulated warming over NH mid-to-high latitudes. The increased aerosol over the EH low-to-mid latitudes is confined more locally but also induces slight warming over the northeastern Pacific and North Atlantic. The competing role of FF forcing originating from EH and WH in shaping tropospheric circulation and surface climate response indicates the importance of both zonal and meridional distribution of aerosol forcing within the NH, and previous idealized models that only consider the zonal difference of aerosol emission may oversimplify the real aerosol forcing.

Deleted: and reveals

Deleted: dominating role

Deleted: WH forcing due to

Deleted: reduction

Deleted: NH. The Aerosol reduction over WH mid-to-high latitudes dominates

Deleted: 2

1 Introduction

The external forcings due to anthropogenic activities and internal variabilities originating from the ocean-atmosphere system together determine climate change at decadal time scales (Kirtman et al., 2013; Meehl et al., 2013). Since the Industrial Revolution, the increasing GHG emissions have been shown to be the leading cause of global warming of about 1.1 °C (as of the late 2010s; IPCC, 2018). On the other hand, the internal variation of the climate, which fluctuates at time scales ranging from years to decades, modulate the paces of global warming at a shorter decadal to a multi decadal time scale (Dai et al., 2015; Xie and Kosaka, 2017; Dong and McPahden, 2017), also with regional implications such as sea ice retreat (Ding et al., 2019).

Deleted: multidecadal

In addition to GHG forcing and internal variability, another major confounding factor affecting global climate change at decadal scales is anthropogenic aerosol forcing. Despite decades of research into this subject, quantitative understandings of the regional climate effects of Anthropogenic Aerosols (AA) remain highly uncertain. There is still limited understanding of the physical mechanisms governing the strength of AA radiative forcing, for example, due to complex aerosol-cloud interaction (Fiedler et al., 2017; Bender, 2020), the brownness of organic aerosols (Bahadur et al., 2012; Jacobson, 2012; Kodros et al., 2015), surface albedo changes due to black carbon aerosols (Xu et al., 2016; Liu et al., 2020).

Deleted: regional

Additionally, there are at least two more reasons why a robust attribution of past climate change to AA is difficult: uneven spatial distributions and fast temporal evolutions of emission/concentration/forcing. Unlike GHGs, the lifetimes of aerosols are as short as days, and thus the spatial distribution of aerosol concentration and its forcing is highly heterogeneous, which may perturb regional climate differently compared to the well-mixed GHG (Ming and Ramaswamy, 2011; Shindell et al., 2015). Lin et al. (2018) analyzed the relationship between aerosol and precipitation extremes and showed that precipitation extremes are more sensitive to aerosols than GHGs, consistent with Salzmann (2016) which examined global mean precipitation. Also, the relatively shorter lifetime means that AA concentrations respond to local emission changes quickly. Indeed, global sulfate aerosol concentration has declined following strengthened emission control measures in the developed nations in the West (Klimont et al., 2013), in contrast to the monotonic increase in GHG concentration since the industrial revolution.

Deleted: That is, unlike

Deleted: .

Moved down [1]: Wang et al.

Deleted: (2020) showed a large shift in South Hadley circulation due to AA in the 20th century.

Moved down [2]: Recently, Wang et al.

Deleted: (2020) demonstrated that the reduced aerosol emission over Europe suppresses the Eurasia wintertime extremes. The decline of AA is likely to continue in the future

Moved down [3]: (Andreae et al., 2005; Zheng et al., 2020),

Deleted: which is projected to intensify severe heat extremes (Zhao et al., 2019; Xu et al., 2020).

However, despite the subtle differences between GHGs and aerosols, other studies found similar climate responses to GHGs and aerosols. Xie et al. (2013) found that the regional ocean temperature and precipitation in response to GHGs and aerosols are similar, suggesting the importance of the spatial distribution of radiative changes. Song et al. (2021) show that increasing GHGs and decreasing aerosols in the recent decades both delay rainfall by inducing a moister atmosphere. Both the differences and similarities between GHGs- and aerosol-induced climate responses indicate the complexity and importance of the temporal and spatial distribution of AA forcings.

Deleted: 3

10

Because of the unique temporal and spatial features of AA, some have argued that the aerosol forcing can induce an external-forced “decadal variability”, which can then be imposed onto the natural variabilities, further confounding a robust attribution of observed changes at a shorter time scale. For example, several recent studies suggested the reduction in aerosol emission over Europe contributes to the Atlantic Multidecadal Variability, which is to a large part attributed to internal oceanic processes (Booth et al., 2012; Bellomo et al., 2018; Hua et al., 2019; Watanabe and Tatebe, 2019). The Atlantic Meridional Overturning Circulation (AMOC) is also argued by recent studies to be induced by AA forcing (Hassan et al., 2020; Menary et al., 2020), though with large uncertainties. Some studies focused on aerosol effects on the Pacific decadal to multidecadal variations, arguing that aerosol forcings can induce Pacific decadal variation (Allen et al., 2014; Dong et al., 2014; Hua et al., 2018), but the relative contribution of external forcing and internal variability remains unclear.

Deleted: . 2019).

15

20

25

Given the rapid temporal evolution of global aerosol emission, as well as the regional redistribution of dominant aerosol emission regions from the West to East, the question of how AA affects the regional climates needs further investment. Many previous studies that examined the aerosol geographical distribution effect on circulation, forcing and temperature (Chemke et al., 2018; Shen et al., 2018). The recent aerosol unmasking in the West can have profound implications on regional climate (Samset et al., 2018; Zhao et al., 2019; Wang et al., 2020b), Arctic sea ice (Krishnan et al., 2020), etc. Wang et al. (2015) demonstrated that the redistribution of aerosol from west to east induces a southward shift of circulation systems and the weakening of tropical circulation. Persad and Caldeira (2018) demonstrated the divergent temperature responses to the regional aerosol forcings at different latitudes based on idealized model simulations. Wang et al. (2020a) showed a large shift in South Hadley circulation due to AA in the 20th century. Recently, Wang et al. (2020b) demonstrated that the reduced aerosol emission over Europe suppresses the Eurasia wintertime extremes.

Deleted: we need to address

Deleted: decadal variations over Asia and the Pacific Ocean. Recent...

Deleted: have compared

Deleted: heterogeneous

Deleted: emission patterns and tried to explain how the climate responds to the competition of EH and WH aerosol emission. The regional warming due to the

Deleted: extremes

Deleted: 2020

Deleted: tropics

Moved (insertion) [1]

Moved (insertion) [2]

Deleted: a

30

35

Understanding the climate response to zonally asymmetric forcing is the main motivation of the present study, because an improved understanding can help shed light on other relevant problems on regional forcings, such as land-use changes (e.g. deforestation over Amazon vs. Africa), volcanic eruption (Verma et al., 2019), geoengineering solutions such as stratospheric or tropospheric aerosol injection conducted over different locations, and the potential contrast of China and India’s future emission trajectories in future decades (Samset et al., 2019; Wang et al., 2021). The FF-related aerosols are projected to further decrease in future decades (Andreae et al., 2005; Zheng et al., 2020), even for Asian regions, with more strict air quality measures in developing nations. The future decline of FF aerosol will lead to further unmasking and warming in addition to GHG-induced global warming (Xu and Xie, 2015; Lelieveld et al., 2019; Allen et al., 2020; Wang et al., 2020a) and have consequences for heat extremes (Zhao et al., 2019; Xu et al., 2020) and humidity and precipitation (Song et al., 2021).

Moved (insertion) [3]

Deleted: The FF aerosol is projected to further decrease in future decades, even for Asian regions, with more strict air quality measurements in developing nations, and thus will lead to further unmasking and warming in addition to GHG-induced global warming (Xu et al., 2015; Wang et al., 2018; Lelieveld et al., 2019; Allen et al., 2020).

Deleted: 4

4

This study leveraged a recently available large ensemble simulation using the fully coupled global climate model and conducted additional “regional” single forcing experiments to assess the aerosol impact on global climate change in the past few decades (1980-2020). The present study focuses on the zonal (WH to EH) asymmetry of aerosol forcing within the Northern Hemisphere. We aim to detail how the upward EH AA emission trend and the downward WH AA trend competes to affect tropical and mid-latitude circulation, and simultaneously, affect the North Pacific surface climate which may have played a role in determining the observed Pacific decadal variations.

Deleted: utilized

Deleted: simulation

Deleted: west

Deleted: east

Deleted: ,

The structure of this paper is the following. In Sect. 2, we provide the details of the climate model, published simulation, and our new model experiment. In Sect. 3, we present simulated responses on the global and regional radiation budget (Sect. 3.1), air temperature (Sect. 3.2), and NH tropospheric circulation (Sect. 3.3) with a focus on separating the role of WH and EH FF forcing that have clear zonal asymmetry. The importance of the latitudinal distribution of AA forcing in driving North Pacific temperature change is highlighted in Sect. 3.4. In Sect. 4, we summarize our findings and suggest scientific questions for future research.

Deleted: over WH and EH with

Deleted: Importance

Deleted: discussed

2 Methods

2.1 Climate model

The climate model used in this study is the Community Earth System Model 1 (CESM1). CESM1 is a fully coupled model developed by NCAR and community scientists (Hurrell et al., 2014) and is one of the models participating in the Coupled Model Intercomparison Project Phase 5 (CMIP5) (Meehl et al., 2016). The CESM1 has been extensively applied in a variety of climate studies including the ones focusing on external forcing and internal variability (e.g., Swart et al., 2015; Xu et al., 2015; Kay et al., 2015; Ding et al., 2019). Studies utilizing CMIP5 multi-model comparison (e.g., Samset et al., 2016; Smith et al., 2016; Lin et al., 2018) also demonstrated its capability for attribution studies on human-induced regional climate change.

Relevant to the aerosol effect focused on this study, a scheme of the three-mode aerosol model (MAM3) - Aitken, accumulation, and coarse modes (Liu et al., 2012), is used by default in CESM1 (CAM5). Aerosol concentration (including sulfates, black carbons, organic carbons) in CESM1 (CAM5) is calculated online from the historical (up to 2005) and future (RCP8.5 thereafter) emission scenarios. The cloud physics scheme allows ice supersaturation and features activation of aerosols to form cloud droplets and ice crystals and thus enables simulations of aerosol indirect effects (Morrison and Gettlemen, 2008), which was missing in the model’s predecessors.

Deleted: full

Deleted: Morris

Deleted: 5

The simulations used in this study are based on a model version of nominal 1° horizontal resolution (0.9° X 1.25°) and 30 vertical levels. All simulation outputs analyzed in this study are monthly data.

15 2.2 Existing simulations

Our study relies on two published large ensemble datasets using CESM1 (CAM5):

- a. CESM1 Large Ensemble Project (CESM1-LENS; Kay et al., 2015);
- b. CESM1 “Single Forcing” Large Ensemble Project (Deser et al., 2020).

20 The CESM1-LENS includes a 40-member ensemble of fully coupled simulations for the period of 1920-2100 with the same historical radiative forcing up to 2005 and the RCP8.5 scenario thereafter (Kay et al., 2015). Each ensemble member starts from the same simulation restart file in 1920 but with slightly different air temperatures perturbed at the level of round-off error. In this paper, we use “ALL” (i.e., all forcing considered) to represent this large ensemble. One advantage of having a large ensemble
25 simulation is that we can separate climate responses to external forcings from internal variabilities by ensemble averaging. Thus, all results in this study are based on the ensemble average.

The CESM1 “Single Forcing” Large Ensemble uses the same model setup of the CESM1-LENS but with individual external forcing fixed at the 1920 level while keeping all other external forcing evolving
30 with time into the 21st century. The “Single Forcing” Large Ensemble includes four sets of ensembles with different single forcings fixed: (1) industrial aerosols (XAERindus, 20 members, 1920-2080), (2) biomass burning aerosols (XAERbmb, 15 members, 1920-2029), (3) greenhouse gases (XGHGs, 20 members, 1920-2080), and (4) land-use/land-cover (XLULC, 5 members, 1920-2029). Here in this
35 study, we only used the first two ensembles. We changed the notation of the two ensembles to “Fix_FF1920” (“FF” stands for Fossil Fuel) and Fix_BB1920 (“BB” stands for Biomass Burning) because, in the emission inventory dataset, energy/transportation sector-related emission is also fixed, rather than the industrial activities only as the original notation implies. We also emphasize the timing (the year 1920) of leveling emission here because anthropogenic aerosol emissions in these simulations are not removed entirely but rather stay at a relatively low level (blue lines in Fig. 1 a and b).

40 By subtracting the “Fix_FF1920” or “Fix_BB1920” ensemble average results from the “ALL” ensemble average, we can obtain climate responses to the Fossil-Fuel-related aerosol forcing (FF) or Biomass-Burning aerosol forcing (BB). Note that other sets of single forcing simulations (e.g., historicalMisc cases in CMIP5 (Taylor et al., 2012), and “hist-aer” cases in CMIP6, (Gillett et al., 2016))
45 simulate only historical aerosol evolution with all other forcings fixed at pre-industrial state. The fix-aerosol method adopted here (as in Deser et al., 2020, but also in earlier studies such as Xu et al., 2015)

6

Deleted: FF1920”

Deleted:)

Deleted: 6

50 serves to estimate the aerosol effects with all other external forcings (such as GHGs) evolving in the background, arguably an advantage in experimental design to assess the actual impact of single forcing.

55 One potential issue of using the fixed single forcing approach is that we have assumed additivity when differencing “ALL” and fixed single forcing cases (i.e., Fix_FF1920 or Fix_BB1920). The additivity assumption is examined in several recent studies focusing on the nonlinear interaction between aerosols and GHGs (Deng et al., 2019) and for various climate variables, in particular for extreme precipitation (Lin et al., 2018).

2.3 New simulation

60 The existing two sets of single forcing large ensemble simulations (Fix_FF1920 and Fix_BB1920) enables a robust separation of aerosol-induced responses and a comparison of the role of FF and BB forcing. However, FF forcing features a strong zonal asymmetry starting from the 1980s (blue solid vs. dashed lines in Fig. 1 c, d), which continuously increases over the EH (dashed lines) and decreases over the WH (solid lines). The sharp contrast and competition between WH and EH in FF emission and forcing trend (driven by different air pollution policy) brings extra complexity to our attribution.

65 To gain further insights on the role of regional forcing (East vs. West), we conducted two additional sets of “regional” single forcing large ensemble simulations (10 realizations for each case) by branching from the existing run of Fix_FF1920. In the experiment of Fixed *Eastern* Fossil Fuel simulation (Fix_EastFF1920), we use the same initialization protocol as Fix_FF1920, but only fix the aerosols over the EH box (0–80°N, 60–150°E; shown as the blue dashed box in Fig. 2), where FF aerosol emissions over other regions are allowed to evolve, including the decline over North America and Europe (shown as the blue solid box in Fig. 2). The experiment of Fixed *Western* Fossil Fuel simulation (Fix_WestFF1920) is similar to the setup of Fix_EastFF1920 case except that we fix the aerosols over WH box (20°–80°N, 130°–10°W, and 30°–80°N, 10°W–40°E; shown as the blue dashed box in Fig. 2).
70 We run the two sets of simulations from 1920 through 1980 for one realization, and then expand the ensemble size to be 10 for 1980–2020. A small random perturbation of surface temperature is applied to each realization to generate ensemble spreads.

80 In Fix_EastFF1920, except for the strong **negative** forcing in the lower latitudes of East Asia, a weak **positive** forcing over Siberia due to the extension of WH aerosol reduction is also included. However, the extended positive forcing is considerably weak compared to the negative forcing and is largely constrained in the small emission domain. Therefore, the difference between ALL and Fix_EastFF1920 can be safely used to represent the climate in response to the dominant role of the negative radiative forcing from lower latitudes of Asia.

Deleted: 2020

Deleted: in review). Furthermore, as in Deser et al., (2020), we consider the simple addition of FF response and BB response as the overall climate response to anthropogenic aerosol forcing (denoted as “AA”). Nevertheless, the focus of this study is on the contrasting role of FF and BB, motivated by the apparent difference in timing (blue vs. green lines in Fig. 1c, d) and emission locations (both in zonal and meridional directions).

Deleted: cooling

Deleted: warming

Deleted: 7

7

95

Similar to how we obtain FF response, we subtract the ensemble average results of Fix_EastFF1920 and Fix_WestFF1920 from the ALL ensemble average results respectively to obtain climate responses to regional Fossil-Fuel-related aerosol forcings (EastFF and WestFF). An additivity test is conducted to evaluate whether the summation of EastFF and WestFF can roughly reproduce FF. The SO4 column burden (BURDENSO4) and surface temperature (TS) in response to FF and EastFF + WestFF (SUM hereafter) are shown in Fig. 3. The FF-induced SO4 column burden resembles the sum of the SO4 burden from SUM. The TS responses are also very similar between FF and SUM, except for the central Pacific and part of the Arctic region. The warmer patterns over the central Pacific in SUM compared to FF is possibly related to the TS responses to remote forcings beyond the two regions in consideration here (e.g., Arabian Peninsula, South America, and Africa), the residues effects of internal variability even after ensemble average due to limited ensemble sizes. Overall, the sum of two sets of regional fixed single forcing experiments well represent the major patterns of FF aerosol induced response, and thus the two new sets of simulations here are capable of separating the East versus West aerosol forcings.

00

05

10

Note that we did not conduct the analogous simulation for BB because, for NH, significant BB emission and forcing trends during 1980-2020 are only over the EH (green dashed lines in Fig. 1c and d), specifically from Northeast Asia. Thus, the existing Fix_BB1920 simulation already captures the regional contribution from the EH.

Deleted: 1 c

Deleted: (blue oval in the right panel of Fig. 2).

15

3 Results

3.1 Zonal asymmetry of anthropogenic aerosol forcing in the recent decades

Figure 1 shows the global and regional emissions of two major types of AA (sulfur aerosols and organic carbon). Globally, it is clear that the AA emission started to decline since the late 20th century (shaded area of 1980-2020 as the focused period of this study), but the decrease in aerosols mainly comes from developed countries in North America and Europe (solid lines as Western Hemisphere (WH) in Fig. 1c, d) while the developing countries in Asia (e.g., China and India) are still in the phase of increasing aerosol emission (dashed lines as Eastern Hemisphere (EH) in Fig. 1c, d).

20

25

The 1st row of Fig. 2 depicts the 40-year linear trend of the SO4 column burden between 1980 and 2020. SO4 trend, as the dominant cooling aerosol produced by FF, shows a clear heterogeneous pattern in NH, with a decrease over North America and Europe (shown as the blue solid box) and a strong increase over China and India (shown as the lower part of the blue dashed box). Note that the decrease

Deleted: blue dashed box).

Deleted: 8

8

in SO4 column burden also occurs over the mid-to-high latitudes of Asia, though with weaker trends compared to Europe and North America.

35 Unlike Sulfate, another major cooling aerosol species, primary organic matter (POM) burden shows different distributions (2nd row of Fig. 2). FF-related POM is similar to that of SO4 burden but with much weaker negative trends over North America and also a weaker positive trend over China, compared to SO4. In contrast, BB-related POM features a much stronger increasing trend over
40 northeastern Asia (40 °N–70 °N, 70 °E–150 °E), and a slight decrease over rain forests of Amazon and Congo. Combining FF and BB, the significant increasing trend of POM occurs over Asia at both low latitudes and high latitudes, while a relatively weaker decline trend can be found over Europe, Africa, and South America, again constituting a west-east zonal asymmetry. The Secondary Organic Aerosol (SOA) burden resembles the SO4 burden in both FF and BB cases but with weaker trends.

45 Aerosols with heating effects (such as Black Carbon; not shown) resemble the spatial pattern of SO4 burden shown in Fig. 2. However, the overall aerosols effect is dominated by cooling aerosols such as SO4. Thus, in this study, we only focus on the total cooling effect of aerosols, without separating the warming and cooling competition as done in several earlier studies (Xu and Xie, 2015; Lin et al., 2016; Wang et al., 2017).

50

Looking at aerosol mass burden only is insufficient to establish connections between the radiative forcing response and aerosol emissions because different aerosol species could have different radiative forcing efficiency. Thus, we further show Aerosol Optical Depth due to anthropogenic aerosol emission (AOD_AA) in the bottom panels of Fig. 2. To remove AOD induced by natural aerosols such as dust and sea salt, we derive AOD_AA following Eq. (1), in which, the AODVIS is the total AOD at the
55 550nm band, and the AODDUST 1-3 represent the dust AOD with different sizes. The background AOD (bkg_AOD) is the 100-year climatology of (AODVIS - AODDUST(1-3)) in the CESM1 pre-industrial control run, which is dominated by sea salt.

$$AOD_{AA} = AODVIS - AODDUST(1-3) - bkg.AOD, \quad (1)$$

60 As expected, the AOD_AA trend (third row in Fig. 2) in response to FF resembles the SO4 burden (first row in Fig. 2), while AOD_AA in response to BB is in close agreement with the POM. Both FF and BB AOD trends feature zonal asymmetry across the Pacific ocean, with subtle differences in terms of latitudinal distribution (increase at lower latitudes versus decrease at higher latitudes in FF. The implications of these spatial contrasts on climatic responses will be further discussed in the next section.

65

It is clear that BB shows a simple distribution without zonal competition, where a significant increase occurs over northeastern Asia. Therefore, our following discussion will only focus on the FF responses, which show subtle zonal differences. Based on the released simulation and the new regional-FF

9

Deleted:)].

Deleted: responses

Deleted: 9

simulations, we are able to separate the climate responses in response to aerosol increase over EH and aerosol reduction over WH.

In line with the zonal asymmetry of AOD_AA trends, simulated solar radiation flux also has significant zonal contrast due to aerosols' direct and indirect climate effects. The first row of Fig. 3 shows the Surface Downward Solar radiation (FSDS), broadly consistent with the patterns of the AOD_AA trend (third row of Fig. 2). Note the opposite colors, though, because a decline in AOD_AA leads to an increase in FSDS. The global surface radiative forcing shows an overall positive trend in response to the decrease in global sulfate emission, but with significant spatial heterogeneity due to the opposite regional emission trends. An increase in FSDS occurs over North America, Europe, and the northern part of Asia, consistent with reducing aerosol forcings over these regions. In contrast, the increasing aerosol emission over east Asia induces a substantial decrease in FSDS. The Net Solar Radiation at the Top-Of-Atmosphere (FSNTOA; the second row of Fig. 3), as the main metric for aerosol forcing, is also consistent with FSDS patterns, but shows more obvious responses over the ocean. Both FSDS and FSNTOA show significant trends over not only the emission domain but also over extended regions into the ocean surface. Due to WestFF, the north Atlantic region shows strong increases in solar radiations, which is consistent with the significant decrease in cloud droplet concentration (CDNUMC, third row of Fig. 3) in response to the WestFF aerosols. However, the cloud fractions (fourth row of Fig. 3) show very weak changes over the north Atlantic, which indicates the critical role of the aerosol first indirect effect over the north Atlantic. The reduction of aerosol emission over North America also leads to smaller cloud droplet concentrations over the North Pacific in the WestFF case, further contributing to a positive radiative forcing. In contrast, in response to EastFF, cloud droplet concentration shows a significant increase over the subtropical Pacific in the Northern Hemisphere, which is consistent with the weak increase in SO4 burden over this region. The larger cloud droplet concentration increases cloud albedo and amplifies the negative radiative forcing. The negative radiative forcing over the subtropical Pacific is evident in the EastFF case, but weaker in the (total) FF case due to the offset by the decreasing aerosols from North America.

The cloud fractions over this region also show an increasing trend in the FF and EastFF, but fail to pass the significance test in the FF case. Surprisingly, the eastern subtropical Pacific in the South Hemisphere also shows significant changes in TOA solar radiation and the cloud fraction, without much aerosol changes. This may possibly be explained by the slow response of sea surface temperature (SST) to the aerosol forcing, where the cloud fraction is affected by the climate adjustment due to SST or circulation changes (Xu and Xie, 2015; Wang et al., 2016; Dong et al., 2019; Kang et al., 2021). The slow responses of cloud fraction to aerosol forcing could also occur near the emission regions where SST changes more significantly; however, as discussed above, the simulated radiation changes over and near the emission regions are highly consistent with the changes in cloud droplet concentrations, indicating a dominant role of indirect aerosol forcing through microphysics perturbation. Here we mainly focused on the overall circulation changes in response to regional aerosol forcings using a fully-

Deleted: major
Deleted: , because of both
Deleted: of aerosols
Deleted:) which is largely
Deleted: net

Deleted: Different from AOD_AA changes which are concentrated over or near emission regions, FSDS changes are also seen over expanded regions due to the aerosol indirect effects, where the cloud droplet numbers and cloud lifetimes are enhanced by aerosols. This is corroborated by the Total Cloud fraction (CLDTOT, the second row of Fig. 3). CLDTOT in response to FF decreases over the North Atlantic, contributing to the significant enhancement of FSDS over NH high latitudes. The Pacific warm pool region in the and FF EastFF cases shows a strong increase in CLDTOT due to the strong local emission increase, and this leads to a strong negative trend in FSDS. Surprisingly, the Pacific warm pool also shows slight increase of CLDTOT in response to WestFF without local emission, which might be associated with the remote influence of WH forcing on the Pacific region

Deleted: Net Solar Radiation at the Top-Of-Atmosphere (FSNTOA; third row of Fig. 3), as
Deleted: main metric for

Deleted: is also

Deleted: FSDS patterns, showing positive global net radiative forcing trend. Interestingly, the

Deleted: 10

35 coupled climate model, therefore a clear separation of the slow and fast responses of clouds and climate to aerosol forcing is beyond the scope of this study.

40 The North Pacific region, a focused region of this study, shows complex competition of the two emission sources, where WestFF induces a significant decrease in cloud droplet concentration (along with increasing FSNTOA) northward of 30 °N. In contrast, EastFF leads to an opposite trend at 30 °N and south. One may expect an increase in FF aerosol over Asia would lead to a negative forcing trend over the North Pacific, as is claimed in previous studies, but actually, the simulated negative trends are confined to lower latitude regions (30 °N and south). The two sets of regional forcing simulations reveal clearly that the decline of FF aerosol over the WH mid-latitudes induces the positive radiative forcing trend at mid-high latitudes of the North Pacific, producing a weak positive FSNTOA trend. This demonstrated East-West competition is a focal point of our following analysis. In the subsequent sections, we will discuss the possible mechanisms in terms of temperature and circulation changes.

50 3.2 Simulated responses in the hemispheric average of surface air temperature

In Sect. 3.1, we demonstrate the distinct East-West pattern of the aerosol emission changes and its radiative effect. This section analyzes the simulated response in the hemispheric average of Surface Air Temperature (SAT).

55 Figure 5a shows the Northern Hemisphere (NH) mean SAT from the two large ensemble simulations. Without the FF aerosol emission in Fix_FF1920 simulation (blue line in Fig. 5a), NH-mean SAT is significantly warmer than the air temperature in the ALL simulation (black line in Fig. 5a). Large volcanic eruptions (Four major ones are shown as vertical dashed lines in Fig. 5a) also strongly affect the NH-mean SAT by causing abrupt cooling of about 0.1 to 0.3 K episodically but the cooling effects quickly recover in a few years, which means it hardly affects the multidecadal climate trend. The global-mean SAT evolutions resemble the NH result but with a weaker magnitude. The stronger response over NH is reasonable because most of the emission sources (and land regions) are located at NH and aerosol burden and radiative forcing (Fig. 2 and 3) are regionally concentrated.

65 Fig. 5b shows the climate response to FF by calculating the difference between ALL and Fix_FF1920 simulation. It indicates that the mid-20th-century aerosol cooling effect is dominated by FF aerosol (Meehl et al., 2004; Diao and Xu., in review) with a cooling trend of 0.16 K/decade over NH. Since the 1980s, as the aerosol emission started to decline over WH, the NH-mean SAT response to FF aerosol has shifted to a slightly warming trend by about 0.07 K/decade.

70

11

Deleted: competitions

Deleted: increase of FSNTOA

Deleted: a decreasing CLDTOT while

Deleted: and weaker

Deleted: . Overall, the negative trend of FSNTOA due to EastFF is offset by WH forcing and leads to a slightly positive trend over this region. By only looking at the FF response over North Pacific, it is somewhat counter-intuitive: one

Deleted: close to the emission source (blue oval in Fig. 2 bottom row left). But the

Deleted: (red oval in Fig. 2 bottom row right) dominates

Deleted: radiative forcings

Deleted: kind of

Deleted: and

Deleted: further

Deleted: later

Deleted: in the subsequent sections.

Deleted: 4a

Deleted: three

Deleted: 4a

Deleted: 4a

Deleted: 4a

Deleted: multi-decadal

Deleted: 4b

Deleted: 11

95 Because of the distinct East-West aerosol forcing asymmetry, we further examine how the aerosol
emissions would influence regional SAT differently. Fig. 5c–d shows the temperature responses over
the Eastern (80 °E–140 °W) and the Western (90 °W–30 °E) portion of NH separately. The domains are
shown as the red boxes in Fig. 4 but here we only have the NH portion to be considered to be consistent
with Fig. 5a and b. The Western-NH temperature in response to FF (Fig. 5d) is largely following local
00 emission evolution (Fig. 1), with a cooling along with the increasing emission before the 1980s and
warming along with emission reduction afterward.

Deleted: 4c

Deleted: 3

Deleted: 4a

Deleted: 4d

Notably, the Eastern-NH responses to FF are counter-intuitive: There is a warming trend of 0.06
K/decade (same as Western-NH) after the 1980s (Fig. 5c) even with continuously increasing aerosol
05 emission over this region, suggesting that Eastern-NH is sensitive to the remote influence of WH
aerosols. Indeed, the Eastern-NH cooling response is even larger than the Western-NH response during
the previous cooling period (1940–1980; -0.17 K/decade versus -0.11 K/decade). The apparent
contradiction of the EH warming in response to FF (Fig. 5c) and the local negative FF forcing (3rd row
of Fig. 2) bears important implications on tropospheric temperature and circulation changes (to be
10 further explored in Sect. 3.4). Here we argue that the larger remote response of EH temperature to
WestFF is due to the latitudinal difference in aerosol emission location: WH emission changes mainly
occur over mid-to-high latitudes (30–60°N; the first row of Fig. 2), while the EH emission changes are
mainly located over low-to-mid latitudes (southward of 40°N, the first row of Fig. 2). As a result, during
15 1980–2020, there is only a weak regional cooling over EH, due to local FF aerosols and is confined over
low latitudes of EH; but the WH decline of emission (positive forcing) dominates the mid-to-high
latitudes SAT change, including North Pacific. Detailed analysis of this subtle latitudinal contrast in
forcing will be provided in Sect. 3.4 for regional SAT over North Pacific.

Deleted: 4c

Deleted: 4c

Deleted: below

Deleted: (visualized in colored ovals in Fig. 2 bottom)

3.3 Tropospheric responses

20 Because of the complex zonal and meridional differences in aerosol emission during 1980–2020 (Sect.
3.1), and the competition between EastFF and WestFF in changing NH air temperature (Sect. 3.2),
tropospheric circulation responses could also be distinct over different regions. In this subsection, we
discuss the global and regional tropospheric circulation responses due to the evolving anthropogenic
aerosol emission, which have a major implication on mid-latitude climate (Xu and Xie, 2015; Mann et
25 al., 2017; Wang et al., 2020b).

Deleted: extreme weather

Deleted: , 2020

Previous studies have explored the tropospheric circulation responses to inter-hemispheric (meridional)
forcing gradient due to anthropogenic aerosols – more reflecting aerosols over NH compared to SH will
lead to an equatorward shift of NH Hadley circulation and NH westerly wind (e.g., Hwang et al., 2013;
30 Hilgenbrink et al., 2018). Meanwhile, recent studies also put effort into how the west-east contrast
effects of aerosol induce the circulation changes (Wang et al., 2015; Kang et al., 2021). However, from

Deleted: 12

1980 to 2020, NH anthropogenic aerosol forcing (Sect. 3.1) is highly heterogeneous, with both strong zonal contrasts and subtle latitudinal differences (Fig. 4), further compounding the forcing-response relationship (Shindell and Faluvegi, 2009; Persad and Caldeira, 2018). Next, we will analyze the aerosol-induced tropospheric responses (in terms of zonal average) both globally and regionally, for the EH and WH portions (domains as red boxes in Fig. 4a).

Deleted: colored ovals at the bottom of

Deleted: 3

Deleted: .

Deleted: 3a

Figure 6a–c shows the decadal trend of global Zonal Mean Meridional overturning Stream Function (ZMMSF) in response to FF, EastFF, and WestFF during 1980–2020. The ZMMSF in response to FF features a counter-clockwise Hadley Cell anomaly (shown in blue) over the tropics, which indicates a northward shift of the Hadley Cell into NH. The northward shift of Hadley Cell also occurs in response to WestFF, but not to EastFF, indicating that the shift of Hadley Cell is mainly due to the WestF. The global mean ZMMSF shifts in our results are consistent with previous studies (Xu and Xie, 2015; Allen and Ajoku, 2016; Amaya et al., 2018; Shen and Ming, 2018) focusing on the inter-hemispheric forcing gradient. That is, the tropical circulation always tends to move towards a warmer hemisphere with larger positive forcing.

Deleted: 5a

Deleted: contour plots in

Deleted: , and the corresponding FSDS trend gradient in the line plot beneath the contour plots

Deleted: with an increasing positive FSDS gradient (dash line in “FSDS gradient” panel).

Deleted: WestFF, which induces warming over NH mid-latitudes and thus leading to a positive interhemispheric FSDS gradient. On the other hand, the EastFF forcing induces a slight negative FSDS gradient over the NH low latitudes, which fails to drive the shift of Hadley Cell...

Deleted: et al.,

Deleted: .

Deleted: et al.,

Deleted: Are NH mid-

To further diagnose why EastFF and WestFF induce distinct changes of the Hadley Cell, we analyse the changes of zonal, column integrated meridional energy transport in response to aerosol forcings, which is shown in Fig. 7b–d. The atmospheric energy transport (AET) is calculated based on the:

$$\frac{\partial}{\partial \phi} F_a = R_{TOA} - Q, \quad (2)$$

Where ϕ is latitude, F_a is the meridional energy flux, R_{TOA} is the net radiative flux at the top-of-atmosphere (downward positive) and Q is the net downward energy flux at the surface. Q includes shortwave radiation, longwave radiation, sensible heat flux, and latent heat flux. AET is then obtained by integrating the energy flux from south to north:

Deleted: jet streams changing consistently as tropical circulation?

$$AET(\Phi) = 2\pi a^2 \int_{-\pi/2}^{\Phi} \cos \Phi' (R_{TOA} - Q) d\Phi' \quad (3)$$

Where a is the Earth radius. Similarly, the oceanic energy transport (OET) is calculated based on the surface radiative flux:

$$\frac{\partial}{\partial \phi} F_o = Q \quad (4)$$

The positive radiative forcing in NH extratropics from WestFF induces a negative AET at the equator (Fig. 7d), which leads to the northward shifts of Hadley Cell and ITCZ to balance the interhemispheric difference in radiative forcing. Previous studies demonstrated that cooling NH leads to southward shift of ITCZ (Broccoli et al., 2006; Kang et al., 2021) consistent with the findings here. On the other hand, the EastFF introduces strong negative radiative forcing in the tropics and weak positive forcing in NH extratropics, but the AET shows small trends at all latitudes compared to that due to WestFF (Fig. 7c).

Deleted: 13

Therefore, the Hadley Cell does not shift significantly in response to EastFF. The AET changes in response to FF resembles that in response to WestFF, indicating the dominant role of WestFF in shifting the Hadley Cell.

Figure 6d-f shows the global-mean zonal wind (U) trends. The FF and EastFF forcing induce slowing U on the poleward flank of the NH jet core while strengthening U on the equatorward flank (especially in the EastFF case), indicating equatorward shifts of the NH jet stream. However, in the WestFF case, U decreases on both flanks of the jet core (slightly greater on the equatorward flank) and the position of the jet core has no significant shift. This is not consistent with the shift of Hadley Cell. As a result, the shift of Hadley Cell and NH jet stream in FF case are in opposite directions, which appears to disagree with previous studies (Xu and Xie, 2015). However, based on the regional FF simulations (Fig 6b-c & e-f), we show that the jet stream and Hadley Cell in FF are controlled by different forcings during this period (negative forcing due to EastFF vs. positive forcing due to WestFF), which agrees with the argument provided by Xu and Xie (2015). The competitions between EastFF and WestFF in shaping the Hadley Cell and NH mid-latitude Jet stream further indicate the importance of the meridional location of the aerosol forcings except for the zonal difference. Previous studies (Seo et al., 2014; Kang et al., 2021) also suggest the importance of latitudinal position of the radiative forcing to the movement of tropical circulations, which is consistent with our findings here.

The latitudinal profiles under the contour plots in Fig. 6 d-f indicate the corresponding zonal mean FSNTOA gradient trend (black curves; in the unit of $W/m^2/decade/10^\circ Lat$), which seems to provide a good rule-of-thumb guidance of the expected shift of NH jet stream. The FSNTOA gradient in response to FF and EastFF show an increase of FSNTOA gradient trend over the mid-latitudes (black dashed lines), which is consistent with the equatorward shift of the NH jet stream. On the contrary, the FSNTOA gradient in WestFF shows a slight negative gradient, while the NH jet stream shows no significant shift. Note that the FSNTOA gradient is only a quick rule-of-thumb guidance of the NH jet shift, and one cannot explain the jet stream shift only based on it. More precise mechanisms of the jet stream shift driven by FF forcings will be discussed below.

Figure 8a-c shows the zonal-mean geopotential wind in the zonal direction (U_g) in EH (red dashed box in Fig. 4), which is derived from geopotential height (Z) following the geostrophic wind equation. The derived U_g patterns always resemble the simulated U pattern in EH (Fig. 8d-f), WH, and Global (not shown), revealing the strong correlation between tropospheric circulation changes and the tropospheric temperature changes (and thus the geopotential height changes). Instead of the gradient of radiative variables as in Fig. 6, here we show the latitudinal profiles of the trend of SAT gradient in response to each force. It has been previously demonstrated that the tropospheric responses to sulfate aerosol are anchored to the SST gradient (Xu and Xie, 2015).

Deleted: 5d...d-f shows the global-mean zonal wind (U) trends. The FF and EastFF forcing induce a reduction of...slowing U speed...on the poleward flank of the NH jet core and an increase of...hile strengthening U speed...n the equatorward flank of the jet core (...especially in the EastFF case), which indicate the ...indicating equatorward shifts of the NH jet stream (marked by red arrows in Fig. 5d-e)... However, in the WestFF case, the ... speed...increases on both sides...lanks of the jet core decreases (...slighter greater on the equatorward side greater than the poleward side...lank) and the position of the jet core has no significant shift. ¶

To summarize, the shift of NH jet stream in response to FF forcing is mainly controlled by the EastFF, but not WestFF. In contrast, as we discussed in the previous paragraph, the shift of the Hadley cell is mainly controlled by the WestFF...his is not consistent with the shift of Hadley Cell. As a result, the shift of Hadley Cell and NH jet stream in FF case are in opposite directions, which appears to disagree with previous studies (Xu et al....nd Xie, 2015). However, based on the regional FF simulations (Fig 5b...b-c & e-f), we show that the jet stream and Hadley Cell in FF are controlled by opposite...ifferent forcings during this period (cooling...egative forcing induced by...ue to EastFF vs. warming...ositive forcing induced by...ue to WestFF), which agrees with the argument provided by Xu et al

Deleted: 2020...021) also suggest the importance of latitudinal position of the radiative forcing to that (... [2]

Deleted: 5... d-f is...ndicate the corresponding zonal mean FSNTOA gradient trend (black curves; in the unit of $W/m^2/decade/10^\circ Lat$), which we argue provides...eems to provide a good rule-of-thumb guidance of the expected shift of NH jet stream. The FSNTOA gradient in response to FF and EastFF show an increase of FSNTOA gradient trend over the mid-latitudes (black dashed lines in Fig. 5..., which tends to drive ...s consistent with the equatorward shift of the NH jet stream. On the contrary, the FSNTOA...SNTOA gradient in WestFF slightly decreases with latitude, with...hows a slight negative gradient, while the NH jet stream shows no significant shift of NH jet stream. Actually, the FSNTOA gradients in FF and WestFF are similar although with a different latitudinal gradient (black dashed lines in Fig.5), which means... Note that purely...he FSNTOA gradient is only a quick of-thumb guidance of the NH jet shift, and one cannot perfectly indicate the shift of NH jet stream. However, considering the consistent relationship between...xplain the shift of ...et stream and the FSNTOA gradient, we argue that...hift only based on it can provide a simple and quick rule-of-thumb predictor of the expected shift of NH jet stream, and more... More precise mechanisms of the jet stream shift driven by FF forcings will be discussed later. (... [3]

Deleted: It's worth noting that in the line plots of Fig. 5, we adopt the latitudinal gradient of radiation flux trend (surface or TOA; in the unit of $W/m^2/decade/10^\circ Lat$), rather than the actual trend. If we adopt the global mean forcing trend instead, it will suggest an opposite shifting direction of moving into the warmer side of the climatological jet stream (for example Fig. 5f), and thus would not serve well as a rule-of-thumb predictor. More fundamentally, it is the latitudinal gradient of forcing (i.e. warming or cooling rate) that drives the jet stream to change, not the local forcing, because it reflects the zonal component of the geostrophic wind equation, which is demonstrated next. ¶ (... [4]

Deleted: 14

55 FF induces nearly no SAT gradient at low latitudes and positive gradients northward of 20 °N (Fig. 8a), which is similar to WestFF. In contrast, EastFF induces a negative SAT gradient southward of 35 °N. The EastFF-induced SAT shows negative gradient on the equator flank of the NH jet core (southward of 35 °N) and positive gradient on the polar flank (max at about 45 °N), indicating a great strengthening of U on the equatorward flank and weakening on the poleward flank. This is producing a net effect of an equatorward shift of NH jet. The WestFF- and FF-induced SAT does not show a negative gradient at low latitudes, and thus U on the equatorward flank of the jet core decreases (WestFF) or slightly increases (FF). The WH jet stream shifts are much weaker compared to the global-mean (Fig. 6d–f) and EH results (Fig. 8d–f) because the air temperature gradients are much weaker and thus are not shown in the figures.

65 To summarize, Fig. 8 shows that the local trend of SAT gradient well explains the weakening or strengthening of the NH jet stream following the geostrophic wind equation, while the latitudinal slope of the SAT gradient (dashed line as the linear fit in Fig. 8) indicates the shift of jet stream. A consistent governing principle emerging from Fig. 8 is that: NH jet stream always tends to shift towards the more negative portion of SAT gradient. This is consistent with Fig. 6 – NH jet stream shifts towards the more negative portion of forcing gradient. Based on Fig. 7, the AET fails to explain the shift of NH jet stream, indicating that the jet stream may be more controlled by the slow response of the aerosol forcing due to surface temperature change rather than the fast response.

75 As shown in Fig. 5, the EH still experiences a warming tendency, in response to FF, the same as WH, despite an increasing aerosol emission locally at low-to-mid latitudes. So, to further reveal the contrast between EH and WH in response to FF as well as regional FF forcings, we re-assess the identified relationship between radiative forcing gradient, temperature gradient, and the tropical circulation changes in Fig. 6–7, by extracting the regional signals from the global mean states.

80 Figure 8 (1st and 2nd rows) shows the regional ZMMSF and U changes relative to the global-mean state. The EH-Globe (EH minus global mean) and WH-Globe show opposite ZMMSF trends at low latitudes with opposite slopes of FSNTOA gradients in both the FF case and the regional FF cases, indicating the importance of cross-equatorial AET to govern the tropospheric circulation adjustment. The increasing FSNTOA gradient from south to north in NH leads to a clockwise ZMMSF trend and a poleward shift of Hadley Cell, which is consistent with the result shown in Xu and Xie (2015) and references within.

90 The relationship between the gradient of the SAT trend and the shift of the NH jet stream is re-assessed in the middle row of Fig. 9, in terms of regional anomalies over EH and WH relative to the global average. The results also support the simple relationship we identified; the NH jet stream shifts to the flank with a more negative SAT gradient because the magnitude of U trend at certain latitudes is

- Deleted: In EH,
- Deleted: above
- Deleted: 6a
- Deleted: While on the other hand
- Deleted: below
- Deleted: The sign of SAT gradient supports our previous argument that the jet stream changes are largely linked with Ug.
- Deleted: in NH has a slightly
- Deleted: below
- Deleted: a
- Deleted: SAT
- Deleted: leading to
- Deleted: zonal wind
- Deleted: of the jet core
- Deleted: (arrow in Fig. 6b), which is consistent with the results in Fig. 5.
- Deleted: the zonal wind
- Deleted: does not increase significantly
- Deleted: continue to decrease but with smaller magnitude compared with the poleward flank.
- Deleted: changes
- Deleted: in comparison with
- Deleted: 5d
- Deleted: 6d
- Deleted: 6
- Deleted: 6
- Deleted: 6
- Deleted: 5
- Deleted: always
- Deleted: 4
- Deleted: , the same as WH,
- Deleted: 5–
- Deleted: 7
- Deleted: FSDS
- Deleted: radiative gradient within NH
- Deleted: In the low-to-mid latitudes of EH (5 °N to 35 °N), (... [5])
- Deleted: FSDS
- Deleted: 7
- Deleted: in Fig. 5–6
- Deleted: 15

determined by the local gradient of air temperature trend, with a more negative gradient strengthening the westerly wind there. One Counterexample here is the WH-Globe U in response to WestFF, where the jet stream does not shift even with a slight negative SAT gradient. The air temperature trend pattern (bottom row of Fig. 9) reveals that the negative gradient of air temperature locates at 30 °N at all pressure levels, which is the latitude of the jet core. As a result, the jet core does not shift much as desired. In the low-to-mid latitudes of EH (5 °N to 35 °N), the gradient of the FSNTOA trend in response to FF is negative compared to the global-mean, which is largely consistent with the increasing aerosol emission in EH (3rd row of Fig. 2).

3.4 Surface temperature responses over the NH with a focus on North Pacific

Having demonstrated the tropical circulation changes and NH jet stream changes in Sect. 3.3, now we look at the SAT response to the regional aerosol forcings. Many previous studies have examined the relationship and mechanisms about Atlantic changes (Booth et al., 2012; Bellomo et al., 2018; Hua et al., 2019; Watanabe and Tatebe, 2019). In contrast, the aerosol effects on the Pacific ocean are comparatively less studied in the previous work (Allen et al., 2014; Dong et al., 2014; Hua et al., 2018), and the potential effects of aerosol redistribution need further discussion. Since this study focuses on the comparison and competition of East and West aerosol forcings, we are specifically interested in how the increasing Asia aerosol forcing affects the North Pacific and how that might be compensated by declining aerosol forcing from North America.

The 1st row of Fig. 10 shows the 40-year linear trends of Surface Air Temperature (SAT) over the ocean in response to FF, EastFF, and WestFF during 1980–2020, which bears a close similarity to SST (not shown). Overall, FF forcing induces significant warming over the North Pacific northward of 40 °N, which is even stronger than the North Atlantic warming but with weaker trend in TOA net radiative flux (Rtoa; second row of Fig10). The EastFF-induced SAT response, unlike FF case, shows significant cooling over the major part of the North Pacific except for the east coast region, which is mentioned by previous studies that the north Pacific cooling due to Asia aerosol emissions (Dong et al., 2014; Takahashi and Watanabe, 2016; Smith et al., 2016). On the other hand, the WestFF, with positive forcing at WH mid-to-high latitudes (30 °N–60 °N; blue oval in Fig. 10), induces a strong warming pattern not only locally at North Atlantic but more so over the North Pacific.

The North Atlantic warming, as many other studies (e.g., Acosta Navarro et al., 2017; Qin et al. 2020) pointed out, can be attributed to the reduction of aerosol emission over North America and Europe since the 1980s, which is clearly seen in TOA net energy flux (blue circle in the 2nd row of Fig. 10). A North Atlantic warming hole is also significant in the FF response (Dagan et al., 2020; Fiedler and Putrasahan 2021). Notably, the simulated warming hole is less significant in response to WestFF forcing alone. In

Deleted: 7

Deleted: Having demonstrated a coherent relationship between tropical circulation changes and that in radiation fluxes and SAT in Sect. 3.3, now we address the following apparent paradox in the FF case: relative to the global mean, EH mid-to-high latitudes have a positive SAT response despite an EH negative forcing, and similarly, WH mid-to-high latitudes have a negative T response (relative to the global mean) despite a local positive forcing (due to aerosol reduction; Fig. 3). Indeed, more than just SAT, EH mid-latitudes (45 °N–60 °N), relative to Global, also shows a warming tendency in tropospheric temperature below 250 hPa, which is contradictory to a local near zero external forcings (Fig. 7c). In contrast, when there is no WH external forcing in the Fix_EastFF1920 simulation, the EH mid-to-high latitudes show significant cooling response due to only EH aerosol forcing (Fig. 7f), which is mentioned by previous studies that demonstrate the north Pacific cooling due to Asia aerosol emissions. This suggests that the mid-to-high latitudes of EH is heavily influenced by WH external forcing, consistent with the hemispheric mean SAT trends shown in Sect. 3.2 and Fig. 4. Next, We provide further discussion of the mechanisms of tropospheric and surface air temperature adjustment in EH and WH.

Moved (insertion) [4]

Moved up [4]: 1
The 1st row of Fig.

Deleted: 8

Deleted: the

Deleted: above

Deleted: during 1980–2020

Deleted: . The warming is inconsistent with the negative FF forcing at EH low-to-mid latitudes (marked by the blue oval in the middle left panel of Fig. 8), which only produces a weak local ... [6]

Deleted: above 50 North North Atlantic shows a warming trend in response to EastFF.

Deleted: warming

Deleted: °N

Deleted: marked by the red

Deleted: the middle left panel of

Deleted: 8) induce es

Deleted: trend, inducing warming patterns

Deleted: Comparing the three sets of responses. It is clear that the North Pacific warming trend above 40 °N is more dominated ... [7]

Moved down [5]: 1

Deleted: starting

Deleted: middle row of Fig. 8). However, the TOA net energy flux over the North Pacific region does not show any stronger pos ... [8]

Deleted: . This reveals that the response of radiative forcing to WestFF cannot fully explain the North Pacific warming. So ... [9]

Deleted: 16

response to the EastFF forcing, the high latitudes of the North Atlantic show a warming trend despite insignificant local changes of TOA net energy flux (2nd row of Fig. 10).

Deleted: ? Intuitively, it seems that increasing aerosol emission over EH (notably China and India) could contribute to regional cooling. However, as shown in Sect. 2 and 3.3, there is a clear contradiction between FF aerosol emission increase and regional-mean warming over the North Pacific. Previous

The equatorial Pacific SAT response to EastFF also shows a El Niño-like pattern, but fails to pass the 95% significance test. This is contrary to some previous studies arguing that Asia aerosol leads to a La Niña-like pattern (Kaufmann et al., 2011; Smith et al., 2016; Kang et al., 2021). The WestFF induced an asymmetric SAT pattern, with cooling over South Pacific while cooling North Pacific. However, both EastFF and WestFF induce significant SAT responses over the North Pacific but not South Pacific, the question about whether aerosol induces a positive or negative phase over the Pacific is still unclear and beyond the scope of this study.

Deleted: also shared similar conclusions,

Deleted: Atlantic SST is more tightly controlled by local

Let's compare the three sets of responses. The weak cooling over the Pacific warm pool region in response to FF can be explained by the offsetting effects between EastFF and WestFF, where the WestFF-induced warming weakens the strong cooling due to EastFF. Similarly, at the North Pacific region southward of 40 °N, the extended aerosol cooling effect from East Asia is largely offset by the warming effect due to WestFF in the total FF response. A notable finding is that, at least based on the simulation here, the North Pacific warming northward of 40 °N is dominated by the positive forcing from WH mid-to-high latitudes, overwhelming the cooling from EH low-to-mid latitudes. The subtle difference in the latitudinal displacement of EH and WH forcings is playing a role here. Indeed, Fig. 4 shows that the EastFF-induced CDNUMC changes are concentrated over low-to-mid latitudes (close to the emission sources and western subtropical Pacific). In contrast, the WestFF-induced CDNUMC changes expand to a larger domain over North Pacific and North Atlantic. The simulation here indicates that mid-to-high latitude SAT is more sensitive to extratropical forcing than forcings originating from a lower latitude. This finding is consistent with previous findings that emission at higher latitudes generates stronger temperature responses (Shindell and Faluvagi, 2009; Persad and Caldeira, 2018). We also examine the BB case (not shown), which has a strong negative forcing over northeastern Asia over 50 °N, and we find that BB-induced cooling occurs over the entire North Pacific similar to WestFF-induced response. Therefore, we highlight that the latitudinal distribution of aerosol forcing is essential to the North Pacific climate responses.

Deleted: radiation change (Boo 2015; Navarro et al., 2017), while North Pacific SST can be more affected by atmospheric circulation changes ...

Deleted: by external forcing

The competition of EastFF and WestFF over the North Pacific deserves some more discussion. There is an apparent paradox in the FF case: compared to the North Atlantic, the high-latitude region of North Pacific shows stronger warming, though with much weaker Rtoa trend. This suggests that the Rtoa cannot fully explain the North Pacific warming. So how does WH forcing lead to North Pacific warming (and also in the tropospheric circulations of EH mid-to-high latitudes in Fig. 9)? We now try to discuss the mechanisms of North Pacific SAT adjustment based on the meridional energy transport.

Moved (insertion) [5]

Deleted: /or internal

Deleted: variations (Qin 2020). ¶

Deleted: 7)?

Deleted: 17

10 The trends of the zonal mean atmospheric meridional heat transport (ZMMHT) are shown in the third row of Fig. 10, where WH shows positive trends northward of 60 °N in all three cases (note that the climatology of ZMMHT over both North Pacific and North Atlantic are still positive in poleward direction, as shown in line contour). This suggests that the meridional energy transfer from the Atlantic Ocean to the Arctic is enhanced, slowing down the North Atlantic warming rate due to local aerosol positive forcing. Conversely, North Pacific ZMMHT (poleward in climatology) is weakened and thus exports less heat from North Pacific mid-latitudes to the polar region (blue color in the EH panels of Fig. 10).

Deleted: ,

Deleted: 8), over

Deleted: a

Deleted: trend above

Deleted: of Fig. 8 bottom

15 In addition to the atmospheric ZMMHT, we also show the zonal, column integrated AET response in EH versus WH (bottom row of Fig. 10). The AET results show a strong northward energy transport trend in the WH, while the equatorward energy transport trend in the EH, both of which resemble the ZMMHT trends. Moreover, it is clear that the strong WestFF positive radiative forcing induces the strong poleward energy transport trend in the WH and the AET in response to EastFF is much weaker.

Deleted: 8). This explains why the North Pacific shows a stronger warming trend even without local emission change or local forcing. Comparing WestFF and EastFF, it is clear that the WestFF forcing is the dominating driver of the equatorward meridional heat transport over North Pacific.

20 Previous studies have also shown that the surface temperature response to aerosol forcing is also strongly modulated by the oceanic energy transport (OET; Cai et al., 2006; Delworth and Dixon, 2006; Dagan et al., 2020; Menary et al., 2020; Fiedler and Putrasahan 2021; Hassan et al., 2021). Here we show that OET (blue lines in Figure 10 bottom row) can have different trends from AET or ZMMHT. In response to the WestFF (positive) forcing, OET increases in WH high latitude and decreases in EH, indicating that poleward heat transport via the ocean is strengthened over the North Atlantic, but is weakened over the North Pacific. The stronger poleward OET in WH, in addition to poleward AET in a larger magnitude, explains why the North Pacific shows a stronger warming trend without local forcing. The EastFF also induces increasing OET in WH and decreasing OET in EH, but with small magnitudes compared with WestFF-induced OET responses, which is similar to AET responses. This further suggests that the WestFF forcing at mid-to-high latitude is the dominant driver of atmospheric and oceanic poleward energy transport in NH.

25 To summarize, the simulated North Pacific warming northward of 50 °N is dominated by the WestFF positive radiative forcing at mid-to-high latitudes. The EastFF forcing at low-to-mid latitudes, which are extensively discussed by previous studies, only induces cooling trend at mid-to-low latitudes (southward of 50 °N) near the emission domain, which is similar to the positive El Niño-like pattern response, but the signal is overwhelmed by the warming tendency induced by WestFF, via both radiative forcing response and meridional heat transport from pole region. Another possible explanation for the stronger North Atlantic warming is that the mid-latitude region of EH is controlled by the WH emission reduction effects at mid-to-high latitudes via the zonal energy transport between the Atlantic and Pacific (McGregor et al., 2014), but it is not tested yet in this study.

Deleted: 2014).

Deleted: 18

4 Summary

The main findings of this paper are:

(1) The **significant** zonal contrast in aerosol emission redistribution, and to a **lesser** extent the latitudinal difference, leads to opposite local radiative forcing that has competing effects on regional climate.”

(2) In terms of hemispheric surface temperature response: the overall FF emission decreases since the 1980s, mostly driven by the WH, induces a significant warming trend over NH. Interestingly, although the FF emission over EH continuously increases, the SAT in EH still shows a warming trend as large as WH, because of the heavy influence from WH aerosol reduction.

(3) In terms of tropospheric circulation responses: in response to FF, NH shows an overall positive gradient of temperature trend (cooling low latitude, warming high latitude), inducing a **counter-clockwise** anomaly (i.e., a poleward shift) of Hadley and an equatorward shift of NH jet stream (Fig. 6). Previous studies show that the shift of mid-latitude jet stream is associated with the SST meridional gradient, and a cooling NH drives an equatorward shift of both Hadley Cell and NH jet stream (Xu and Xie, 2015; Xu et al., 2016), which appears to be inconsistent with our results. The main difference is that previous results are largely based on a sharp interhemispheric forcing gradient with a time scale of the entire 20th century (Wang et al., 2020a), or focusing on the mid-century era where global aerosol emission is on the rise.

In this study, with a focus on the period of 1980–2020, the regional FF forcing competitions within the NH is more complex given that the EH and WH show heterogeneous distributions in both zonal and meridional direction, and even opposite trends (an increase of aerosol emission over EH versus reduction over WH). The competitions between EH and WH are illustrated with two sets of separate large ensemble simulations (Fix_EastFF1920 and Fix_WestFF1920) we conducted. Our result shows that the shift of Hadley Cell is **dominated** by the WestFF forcing at mid-to-high latitudes. **WestFF induces positive radiative forcing in the NH extratropics and drives the northward shift of the Hadley Cell to balance the interhemispheric difference in radiative forcing.**

The equatorward shift of the NH jet stream is mainly driven by EastFF forcing at low-to-mid latitudes. The competitions between EastFF and WestFF in shaping the Hadley Cell and mid-latitude Jet stream

Deleted: To summarize, the simulated North Pacific warming above 50 °N is dominating by the WestFF forcing at mid-to-high latitudes. The EastFF forcing at low-to-mid latitudes, which are extensively discussed by previous studies, induces cooling trend at mid latitudes (below 50 °N) over North Pacific, which is similar to the positive PDO-like response, but the signal is overwhelmed by the warming tendency induced by WestFF, via both radiative forcing response and meridional heat transport from pole region.⁴

Deleted: In terms of aerosol forcing, the Eastern Hemisphere (EH) and the Western Hemisphere (WH) show very different anthropogenic aerosol emission trends since the 1980s, where WH started to reduce Fossil Fuel (FF) emission in comparison to the continuous increasing in both FF and BB emission over EH (Fig.

Deleted: and Fig. 2). Due to FF, positive forcing occurs at mid to high latitudes (above 30 °N) and negative forcing occurs over low latitude regions (particularly West tropical Pacific ocean and Indian ocean). Due to BB, negative forcing mainly occurs close to the emission domain (Northeastern Asia) with some extension to the North Pacific (Fig. 3).

Deleted: major

Deleted: less

Deleted: .

Deleted: counterclockwise

Deleted: 5

Deleted: 2020

Deleted: (Diao and Xu, in review).

Deleted: poleward

Deleted: controlled

Deleted: , while

Deleted: 19

20 indicate the key role of the meridional distribution of the aerosol forcings. The gradient of Top-of-
Atmosphere net solar flux (FSNTOA) provides a rule-of-thumb guidance of the expected shift of the
NH jet stream (Fig. 9). The jet stream patterns largely agree with the geostrophic wind (Ug) patterns
(Fig. 8), which are derived from the latitudinal gradient of air temperature. The NH jet stream tends to
shift to the negative gradient portion of FSNTOA.

25 (4) In terms of North Pacific temperature response, the FF forcing during 1980-2020, unlike
suggested by previous studies, induces North Pacific and pan-Pacific warming due to a competition
between EastFF and WestFF, with the latter dominating the former (Fig. 10). The dominance is due to
the latitudinal distribution of aerosol forcing within the NH. The negative forcing of EastFF, which
30 occurs at the EH tropical and subtropical region (0–40 °N), is largely confined to the emission domain.
In contrast, negative FF forcing over WH mid-to-high latitudes (northward of 30 °N), not only
introduces local warming but also imposes a heavy influence over North Pacific warming. Diagnostic
data shows that the remote contribution to North Pacific warming from WestFF is due to the
35 combination of both radiative forcing responses and meridional energy transfer anomaly from the North
Atlantic to North Pacific via the Arctic pathway.

The importance of the inter-hemispheric asymmetry of external forcing has been extensively discussed
in previous studies (e.g., Xu and Xie, 2015; Chung and Soden, 2017; Wang et al., 2020a), in which the
40 NH is usually considered as a whole. Here we further emphasize that the latitudinal difference of
external forcing within the NH is important in determining the tropospheric and surface responses.
More specifically, the EastFF induces a PDO-like pattern (negative phase) with a cooling over the
northwestern Pacific and warming over the tropical Pacific. However, when combined with the aerosol
reduction over the WH, the EastFF-driven northwestern Pacific cooling is completely offset and instead
45 exhibits a warming trend.

We provide the following discussions related to our conclusion and suggest some future research
directions.

(1) The issue of nonlinearity. The CESM1 single forcing large ensemble simulations applied in this
50 study treats the FF and BB forcing separately, which enables many of our discussions above. However,
one problem is the additivity of the two anthropogenic aerosol sources. Deser et al. (2020) sum up the
FF, BB, and GHGs responses to reconstruct the all forcing (ALL) response, and find some
inconsistencies due to the nonlinear interactions between aerosol- and GHG-induced responses. So, it is
also possible that adding FF and BB may cause some nonlinear problems. Similarly, adding EastFF and
55 WestFF responses may also cause such problems. This study is not affected by such additivity problems
because we treat each set of simulations separately, but future studies that aim to utilize the combined
AA-induced changes, should be cautious and a rigorous test on additivity would be necessary.

20

Deleted: The gradient of surface downward solar flux (FSDS) provides a rule-of-thumb guidance of the expected shift of the NH jet stream (Fig. 7). It is because the jet stream patterns largely agree with the geostrophic wind (Ug) patterns (Fig. 6), which are derived from the latitudinal gradient of air temperature. The NH jet stream always tends to shift to the negative gradient portion of FSDS.

Deleted: 8

Deleted: above

Deleted: et al.,

Deleted: et al.,

Deleted: 2020

Deleted: The

Deleted: 20

75 (2) The issue of RCP8.5 scenarios applied in CESM1. As is described in Sect. 2.1, the RCP8.5 scenarios are applied in all five sets of large ensemble simulations used in this study for 2006-2020. Studies have pointed out that the RCP dataset is different from the new SSP scenarios which contain the historical emission data up to 2015. For example, the sulfate emissions over India and China over the 2006-2020 period in the RCP8.5 scenarios are lower than observations and SSP scenarios, which could lead to an underestimate of the sulfate cooling effects locally and globally (Lin et al., 2018). Future experiments utilizing the newer GCMs and SSP scenarios would be helpful to repeat and test the current results.

80 (3) The single model large ensemble method. The single model large ensemble simulations applied in this study effectively separate the external forcing induced response from the model internal variability (Kay et al., 2015; Deser, et al., 2020). However, one limitation here is that only one GCM is utilized, so the potential systematic errors from a single model cannot be tested. Some new single-forcing large ensemble simulations based on models other than CESM1 (e.g., CanESM2 (Oudar et al., 2018); CESM2) are also coming out recently as part of the “Single Model Initial-condition Large Ensemble” (SMILE) efforts, which brings the possibility of testing “multi-model large ensemble” in future works and improving our understanding of aerosol induced climate change.

90 (4) The implication of the North Pacific response on the PDO mode and global warming rate. The increasing FF emission over EH low latitudes (without WH) induce a PDO-like SAT response over the North Pacific and tropical Pacific, suggesting the potential impact of anthropogenic forcing onto the internal variability. More future research is needed on this topic based on the current large ensemble simulations and the upcoming CESM2 simulations.

95 (5) The subtle difference of EastFF and WestFF in the north-south direction. It is clear that the redistribution of anthropogenic aerosols since the 1980s is not a pure zonal shift, but also shifts in the meridional direction. The aerosol loading shows a net decrease, especially in the mid-latitude/subpolar region. Future studies may consider both zonal and meridional shifts in SO4 loading.

00

21

Deleted: aerosol

Deleted: 21

References

- Acosta Navarro, J. C., Ekman, A. M. L., Pausata, F. S. R., Lewinschal, A., Varma, V., Seland, Ø., Gauss, M., Iversen, T., Kirkevåg, A., Riipinen, I., and Hansson, H. C.: Future Response of Temperature and Precipitation to Reduced Aerosol Emissions as Compared with Increased Greenhouse Gas Concentrations, *J. Clim.*, 30, 939–954, <https://doi.org/10.1175/JCLI-D-16-0466.1>, 2017.
- Allen, R. J. and Ajoku, O.: Future aerosol reductions and widening of the northern tropical belt, *J. Geophys. Res. Atmos.*, 121, 6765–6786, <https://doi.org/10.1002/2016JD024803>, 2016.
- Allen, R. J., Norris, J. R., and Kovilakam, M.: Influence of anthropogenic aerosols and the Pacific Decadal Oscillation on tropical belt width, *Nat. Geosci.*, 7, 270–274, <https://doi.org/10.1038/ngeo2091>, 2014.
- Allen, R. J., Turnock, S., Nabat, P., Neubauer, D., Lohmann, U., Olivíć, D., Oshima, N., Michou, M., Wu, T., Zhang, J., Takemura, T., Schulz, M., Tsigaridis, K., Bauer, S. E., Emmons, L., Horowitz, L., Naik, V., van Noije, T., Bergman, T., Lamarque, J.-F., Zanis, P., Tegen, I., Westervelt, D. M., Le Sager, P., Good, P., Shim, S., O'Connor, F., Akritidis, D., Georgoulas, A. K., Deushi, M., Sentman, L. T., John, J. G., Fujimori, S., and Collins, W. J.: Climate and air quality impacts due to mitigation of non-methane near-term climate forcers, *Atmos. Chem. Phys.*, 20, 9641–9663, <https://doi.org/10.5194/acp-20-9641-2020>, 2020.
- Amaya, D. J., Siler, N., Xie, S.-P., and Miller, A. J.: The interplay of internal and forced modes of Hadley Cell expansion: lessons from the global warming hiatus, *Clim. Dyn.*, 51, 305–319, <https://doi.org/10.1007/s00382-017-3921-5>, 2018.
- Andreae, M. O., Jones, C. D., and Cox, P. M.: Strong present-day aerosol cooling implies a hot future, *Nature*, 435, 1187–1190, <https://doi.org/10.1038/nature03671>, 2005.
- Bahadur, R., Praveen, P. S., Xu, Y., and Ramanathan, V.: Solar absorption by elemental and brown carbon determined from spectral observations, *Proc. Natl. Acad. Sci.*, 109, 17366–17371, <https://doi.org/10.1073/pnas.1205910109>, 2012.
- Bellomo, K., Murphy, L. N., Cane, M. A., Clement, A. C., and Polvani, L. M.: Historical forcings as main drivers of the Atlantic multidecadal variability in the CESM large ensemble, *Clim. Dyn.*, 50, 3687–3698, <https://doi.org/10.1007/s00382-017-3834-3>, 2018.
- Bender, F. A.-M.: Aerosol Forcing: Still Uncertain, Still Relevant, *AGU Adv.*, 1, e2019AV000128, <https://doi.org/10.1029/2019AV000128>, 2020.
- Booth, B. B. B., Dunstone, N. J., Halloran, P. R., Andrews, T., and Bellouin, N.: Aerosols implicated as a prime driver of twentieth-century North Atlantic climate variability, *Nature*, 484, 228–232, <https://doi.org/10.1038/nature10946>, 2012.
- Cai, W., Bi, D., Church, J., Cowan, T., Dix, M., and Rotstayn, L.: Pan-oceanic response to increasing anthropogenic aerosols: Impacts on the Southern Hemisphere oceanic circulation, *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2006GL027513>, 2006.

- 40 [Chemke, R. and Dagan, G.: The Effects of the Spatial Distribution of Direct Anthropogenic Aerosols Radiative Forcing on Atmospheric Circulation, J. Clim., 31, 7129–7145, https://doi.org/10.1175/JCLI-D-17-0694.1, 2018.](https://doi.org/10.1175/JCLI-D-17-0694.1)
- [Chung, E.-S. and Soden, B. J.: Hemispheric climate shifts driven by anthropogenic aerosol–cloud interactions, 10, 566–571, https://doi.org/10.1038/ngeo2988, 2017.](https://doi.org/10.1038/ngeo2988)
- 45 [Dagan, G., Stier, P., and Watson-Parris, D.: Aerosol Forcing Masks and Delays the Formation of the North Atlantic Warming Hole by Three Decades, Geophys. Res. Lett., 47, e2020GL090778, https://doi.org/https://doi.org/10.1029/2020GL090778, 2020.](https://doi.org/https://doi.org/10.1029/2020GL090778)
- Dai, A., Fyfe, J. C., Xie, S. P., and Dai, X.: Decadal modulation of global surface temperature by internal climate variability, Nat. Clim. Chang., 5, 555, https://doi.org/10.1038/nclimate2605, 2015.
- 50 [Delworth, T. L. and Dixon, K. W.: Have anthropogenic aerosols delayed a greenhouse gas-induced weakening of the North Atlantic thermohaline circulation?, Geophys. Res. Lett., 33, https://doi.org/https://doi.org/10.1029/2005GL024980, 2006.](https://doi.org/https://doi.org/10.1029/2005GL024980)
- Deng, J., Dai, A., and Xu, H.: Nonlinear Climate Responses to Increasing CO2 and Anthropogenic Aerosols Simulated by CESM1, J. Clim., 33, 281–301, https://doi.org/10.1175/JCLI-D-19-0195.1, 2019.
- 55 Deser, C., Phillips, A. S., Simpson, I. R., Rosenbloom, N., Coleman, D., Lehner, F., Pendergrass, A. G., DiNezio, P., and Stevenson, S.: Isolating the Evolving Contributions of Anthropogenic Aerosols and Greenhouse Gases: A New CESM1 Large Ensemble Community Resource, J. Clim., 33, 7835–7858, https://doi.org/10.1175/JCLI-D-20-0123.1, 2020.
- Ding, Q., Schweiger, A., L’Heureux, M., Steig, E. J., Battisti, D. S., Johnson, N. C., Blanchard-Wrigglesworth, E., Po-Chedley, S., Zhang, Q., Harnos, K., Bushuk, M., Markle, B., and Baxter, I.: Fingerprints of internal drivers of Arctic sea ice loss in observations and model simulations, Nat. Geosci., 12, 28–33, https://doi.org/10.1038/s41561-018-0256-8, 2019.
- 60 [Dong, B., Wilcox, L. J., Highwood, E. J., and Sutton, R. T.: Impacts of recent decadal changes in Asian aerosols on the East Asian summer monsoon: roles of aerosol–radiation and aerosol–cloud interactions, Clim. Dyn., 53, 3235–3256, https://doi.org/10.1007/s00382-019-04698-0, 2019.](https://doi.org/10.1007/s00382-019-04698-0)
- 65 [Dong, L. and McPhaden, M. J.: The role of external forcing and internal variability in regulating global mean surface temperatures on decadal timescales, Environ. Res. Lett., 12, 034011, https://doi.org/10.1088/1748-9326/aa5dd8, 2017.](https://doi.org/10.1088/1748-9326/aa5dd8)
- 70 [Dong, L., Zhou, T., and Chen, X.: Changes of Pacific decadal variability in the twentieth century driven by internal variability, greenhouse gases, and aerosols, Geophys. Res. Lett., https://doi.org/10.1002/2014GL062269, 2014.](https://doi.org/10.1002/2014GL062269)
- [Fiedler, S. and Putrasahan, D.: How Does the North Atlantic SST Pattern Respond to Anthropogenic Aerosols in the 1970s and 2000s?, Geophys. Res. Lett., 48, e2020GL092142, https://doi.org/https://doi.org/10.1029/2020GL092142, 2021.](https://doi.org/https://doi.org/10.1029/2020GL092142)

- 75 [Fiedler, S., Stevens, B., and Mauritsen, T.: On the sensitivity of anthropogenic aerosol forcing to model-internal variability and parameterizing a Twomey effect, J. Adv. Model. Earth Syst., 9, 1325–1341, https://doi.org/https://doi.org/10.1002/2017MS000932, 2017.](https://doi.org/https://doi.org/10.1002/2017MS000932)
- [Gillett, N. P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., Santer, B. D., Stone, D., and Tebaldi, C.: The Detection and Attribution Model Intercomparison Project \(DAMIP v1.0\) contribution to CMIP6, Geosci. Model Dev., 9, 3685–3697, https://doi.org/10.5194/gmd-9-3685-2016, 2016.](https://doi.org/https://doi.org/10.5194/gmd-9-3685-2016)
- 80 [Hassan, T., Allen, R. J., Liu, W., and Randles, C. A.: Anthropogenic aerosol forcing of the Atlantic meridional overturning circulation and the associated mechanisms in CMIP6 models, 21, 5821–5846, https://doi.org/10.5194/acp-21-5821-2021, 2021.](https://doi.org/10.5194/acp-21-5821-2021)
- Hilgenbrink, C. C. and Hartmann, D. L.: The Response of Hadley Circulation Extent to an Idealized Representation of Poleward Ocean Heat Transport in an Aquaplanet GCM, J. Clim., 31, 9753–9770, <https://doi.org/10.1175/JCLI-D-18-0324.1>, 2018.
- 85 Hua, W., Dai, A., and Qin, M.: Contributions of Internal Variability and External Forcing to the Recent Pacific Decadal Variations, Geophys. Res. Lett., 45, 7084–7092, <https://doi.org/10.1029/2018GL079033>, 2018.
- Hua, W., Dai, A., Zhou, L., Qin, M., and Chen, H.: An Externally Forced Decadal Rainfall Seesaw Pattern Over the Sahel and Southeast Amazon, Geophys. Res. Lett., 46, 923–932, <https://doi.org/10.1029/2018GL081406>, 2019.
- 90 [Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A Framework for Collaborative Research, Bull. Am. Meteorol. Soc., 94, 1339–1360, https://doi.org/10.1175/BAMS-D-12-00121.1, 2013.](https://doi.org/10.1175/BAMS-D-12-00121.1)
- IPCC: IPCC Special Report on the impacts of global warming of 1.5°C, Ipcc - Sr15, 2018.
- Jacobson, M. Z.: Investigating cloud absorption effects: Global absorption properties of black carbon, tar balls, and soil dust in clouds and aerosols, J. Geophys. Res. Atmos., 117, <https://doi.org/10.1029/2011JD017218>, 2012.
- 00 [Kang, S. M., Xie, S.-P., Deser, C., and Xiang, B.: Zonal mean and shift modes of historical climate response to evolving aerosol distribution, Sci. Bull., https://doi.org/https://doi.org/10.1016/j.scib.2021.07.013, 2021.](https://doi.org/https://doi.org/10.1016/j.scib.2021.07.013)
- Kaufmann, R. K., Kauppi, H., Mann, M. L., and Stock, J. H.: Reconciling anthropogenic climate change with observed temperature 1998–2008, Proc. Natl. Acad. Sci., 108, 11790 LP – 11793, <https://doi.org/10.1073/pnas.1102467108>, 2011.
- 05 Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J.-F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M.: The Community

- 10 Earth System Model (CESM) Large Ensemble Project: A Community Resource for Studying Climate Change in the Presence of Internal Climate Variability, *Bull. Am. Meteorol. Soc.*, 96, 1333–1349, <https://doi.org/10.1175/BAMS-D-13-00255.1>, 2015.
- Kirtman, B., Power, S. B., Adedoyin, A. J., Boer, G. J., Bojariu, R., Camilloni, I., Doblas-Reyes, F., Fiore, A. M., Kimoto, M., Meehl, G., Prather, M., Sarr, A., Schar, C., Sutton, R., van Oldenborgh, G. J.,
15 Vecchi, G., and Wang, H.-J.: Chapter 11 - Near-term climate change: Projections and predictability, in: *Climate Change 2013: The Physical Science Basis. IPCC Working Group I Contribution to AR5*, edited by: IPCC, Cambridge University Press, Cambridge, 2013.
- Klimont, Z., Smith, S. J., and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions, *Environ. Res. Lett.*, 8, 14003, <https://doi.org/10.1088/1748-9326/8/1/014003>, 2013.
- 20 Kodros, J. K., Scott, C. E., Farina, S. C., Lee, Y. H., L'Orange, C., Volckens, J., and Pierce, J. R.: Uncertainties in global aerosols and climate effects due to biofuel emissions, *Atmos. Chem. Phys.*, 15, 8577–8596, <https://doi.org/10.5194/acp-15-8577-2015>, 2015.
- Krishnan, S., Ekman, A. M. L., Hansson, H.-C., Riipinen, I., Lewinschal, A., Wilcox, L. J., and Dallafior, T.: The Roles of the Atmosphere and Ocean in Driving Arctic Warming Due to European
25 Aerosol Reductions, *Geophys. Res. Lett.*, 47, e2019GL086681, <https://doi.org/10.1029/2019GL086681>, 2020.
- [Lelieveld, J., Klingmüller, K., Pozzer, A., Burnett, R. T., Haines, A., and Ramanathan, V.: Effects of fossil fuel and total anthropogenic emission removal on public health and climate. *Proc. Natl. Acad. Sci.*, 116, 7192–7197. <https://doi.org/10.1073/pnas.1819989116>, 2019.](https://doi.org/10.1073/pnas.1819989116)
- 30 Lin, L., Gettelman, A., Xu, Y., and Fu, Q.: Simulated responses of terrestrial aridity to black carbon and sulfate aerosols, *J. Geophys. Res. Atmos.*, 121, 785–794, <https://doi.org/10.1002/2015JD024100>, 2016.
- Lin, L., Xu, Y., Wang, Z., Diao, C., Dong, W., and Xie, S.-P.: Changes in Extreme Rainfall Over India and China Attributed to Regional Aerosol-Cloud Interaction During the Late 20th Century Rapid Industrialization, *Geophys. Res. Lett.*, 45, 7857–7865, <https://doi.org/10.1029/2018GL078308>, 2018.
- 35 Liu, D., He, C., Schwarz, J. P., and Wang, X.: Lifecycle of light-absorbing carbonaceous aerosols in the atmosphere, *npj Clim. Atmos. Sci.*, 3, 40, <https://doi.org/10.1038/s41612-020-00145-8>, 2020.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward
40 a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*, 5, 709–739, <https://doi.org/10.5194/gmd-5-709-2012>, 2012.
- Mann, M. E., Rahmstorf, S., Kornhuber, K., Steinman, B. A., Miller, S. K., and Coumou, D.: Influence of Anthropogenic Climate Change on Planetary Wave Resonance and Extreme Weather Events, *Sci. Rep.*, 7, 45242, <https://doi.org/10.1038/srep45242>, 2017.
- 45

- McGregor, S., Timmermann, A., Stuecker, M. F., England, M. H., Merrifield, M., Jin, F.-F., and Chikamoto, Y.: Recent Walker circulation strengthening and Pacific cooling amplified by Atlantic warming, *Nat. Clim. Chang.*, 4, 888–892, <https://doi.org/10.1038/nclimate2330>, 2014.
- 50 Meehl, G. A., Hu, A., Arblaster, J. M., Fasullo, J., and Trenberth, K. E.: Externally Forced and Internally Generated Decadal Climate Variability Associated with the Interdecadal Pacific Oscillation, *J. Clim.*, 26, 7298–7310, <https://doi.org/10.1175/JCLI-D-12-00548.1>, 2013.
- Meehl, G. A., Washington, W. M., Ammann, C. M., Arblaster, J. M., Wigley, T. M. L., and Tebaldi, C.: Combinations of Natural and Anthropogenic Forcings in Twentieth-Century Climate, *J. Clim.*, 17, 3721–3727, [https://doi.org/10.1175/1520-0442\(2004\)017<3721:CONAAF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<3721:CONAAF>2.0.CO;2), 2004.
- 55 Meehl, G. A., Hu, A., and Teng, H.: Initialized decadal prediction for transition to positive phase of the Interdecadal Pacific Oscillation, *Nat. Commun.*, 7, <https://doi.org/10.1038/ncomms11718>, 2016.
- [Menary, M. B., Robson, J., Allan, R. P., Booth, B. B. B., Cassou, C., Gastineau, G., Gregory, J., Hodson, D., Jones, C., Mignot, J., Ringer, M., Sutton, R., Wilcox, L., and Zhang, R.: Aerosol-Forced AMOC Changes in CMIP6 Historical Simulations, *Geophys. Res. Lett.*, 47, e2020GL088166, <https://doi.org/https://doi.org/10.1029/2020GL088166>, 2020.](https://doi.org/https://doi.org/10.1029/2020GL088166)
- 60 Ming, Y. and Ramaswamy, V.: A Model Investigation of Aerosol-Induced Changes in Tropical Circulation, *J. Clim.*, 24, 5125–5133, <https://doi.org/10.1175/2011JCLI4108.1>, 2011.
- [Morrison, H. and Gettelman, A.: A New Two-Moment Bulk Stratiform Cloud Microphysics Scheme in the Community Atmosphere Model, Version 3 \(CAM3\). Part I: Description and Numerical Tests, *J. Clim.*, 21, 3642–3659, <https://doi.org/10.1175/2008JCLI2105.1>, 2008.](https://doi.org/10.1175/2008JCLI2105.1)
- 65 Oudar, T., Kushner, P. J., Fyfe, J. C., and Sigmond, M.: No Impact of Anthropogenic Aerosols on Early 21st Century Global Temperature Trends in a Large Initial-Condition Ensemble, *Geophys. Res. Lett.*, 45, 9245–9252, <https://doi.org/https://doi.org/10.1029/2018GL078841>, 2018.
- [Persad, G. G. and Caldeira, K.: Divergent global-scale temperature effects from identical aerosols emitted in different regions, *Nat. Commun.*, 9, 3289, <https://doi.org/10.1038/s41467-018-05838-6>, 2018.](https://doi.org/10.1038/s41467-018-05838-6)
- 70 Qin, M., Dai, A., and Hua, W.: Aerosol-forced multidecadal variations across all ocean basins in models and observations since 1920, *Sci. Adv.*, 6, eabb0425, <https://doi.org/10.1126/sciadv.abb0425>, 2020.
- [Salzmann, M.: Global warming without global mean precipitation increase?, *Sci. Adv.*, 2, e1501572, <https://doi.org/10.1126/sciadv.1501572>, 2016.](https://doi.org/10.1126/sciadv.1501572)
- 75 Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, Ø., Andrews, T., Faluvegi, G., Fläschner, D., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivie, D., Richardson, T., Shindell, D., Shine, K. P., Takemura, T., and Voulgarakis, A.: Fast and slow precipitation responses to individual climate forcings: A PDRMIP multimodel study, *Geophys. Res. Lett.*, 43, 2782–2791, <https://doi.org/10.1002/2016GL068064>, 2016.
- 80

Deleted: 26

85 [Samset, B. H., Stjern, C. W., Andrews, E., Kahn, R. A., Myhre, G., Schulz, M., and Schuster, G. L.: Aerosol Absorption: Progress Towards Global and Regional Constraints, *Curr. Clim. Chang. Reports*, 4, 65–83, <https://doi.org/10.1007/s40641-018-0091-4>, 2018.](#)

[Samset, B. H., Lund, M. T., Bollasina, M., Myhre, G., and Wilcox, L.: Emerging Asian aerosol patterns, *Nat. Geosci.*, 12, 582–584, <https://doi.org/10.1038/s41561-019-0424-5>, 2019.](#)

Shen, Z. and Ming, Y.: The Influence of Aerosol Absorption on the Extratropical Circulation, *J. Clim.*, 31, 5961–5975, <https://doi.org/10.1175/JCLI-D-17-0839.1>, 2018.

90 Shindell, D. T., Faluvegi, G., Rotstayn, L., and Milly, G.: Spatial patterns of radiative forcing and surface temperature response, *J. Geophys. Res. Atmos.*, 120, 5385–5403, <https://doi.org/doi:10.1002/2014JD022752>, 2015.

[Shindell, D. and Faluvegi, G.: Climate response to regional radiative forcing during the twentieth century, *Nat. Geosci.*, 2, 294–300, <https://doi.org/10.1038/ngeo473>, 2009.](#)

95 Smith, D. M., Booth, B. B. B., Dunstone, N. J., Eade, R., Hermanson, L., Jones, G. S., Scaife, A. A., Sheen, K. L., and Thompson, V.: Role of volcanic and anthropogenic aerosols in the recent global surface warming slowdown, *Nat. Clim. Chang.*, 6, 936–940, <https://doi.org/10.1038/nclimate3058>, 2016.

[Song, F., Leung, L. R., Lu, J., Dong, L., Zhou, W., Harrop, B., and Qian, Y.: Emergence of seasonal delay of tropical rainfall during 1979–2019, *Nat. Clim. Chang.*, 11, 605–612, <https://doi.org/10.1038/s41558-021-01066-x>, 2021.](#)

00 Swart, N. C., Fyfe, J. C., Hawkins, E., Kay, J. E., and Jahn, A.: Influence of internal variability on Arctic sea-ice trends, *Nat. Clim. Chang.*, 5, 86–89, <https://doi.org/10.1038/nclimate2483>, 2015.

[Takahashi, C. and Watanabe, M.: Pacific trade winds accelerated by aerosol forcing over the past two decades, *Nat. Clim. Chang.*, 6, <https://doi.org/10.1038/nclimate2996>, 2016.](#)

05 [Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *Bull. Am. Meteorol. Soc.*, 93, 485–498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.](#)

Verma, T., Saravanan, R., Chang, P., and Mahajan, S.: Tropical Pacific Ocean Dynamical Response to Short-Term Sulfate Aerosol Forcing, *J. Clim.*, 32, 8205–8221, <https://doi.org/10.1175/JCLI-D-19-0050.1>, 2019.

10 Wang, H., Xie, S.-P., and Liu, Q.: Comparison of Climate Response to Anthropogenic Aerosol versus Greenhouse Gas Forcing: Distinct Patterns, *J. Clim.*, 29, 5175–5188, <https://doi.org/10.1175/JCLI-D-16-0106.1>, 2016.

[Wang, H., Xie, S.-P., Zheng, X.-T., Kosaka, Y., Xu, Y., and Geng, Y.-F.: Dynamics of Southern Hemisphere Atmospheric Circulation Response to Anthropogenic Aerosol Forcing, *Geophys. Res. Lett.*, 47, e2020GL089919, <https://doi.org/10.1029/2020GL089919>, 2020a.](#)

15 Wang, Y., Jiang, J. H., and Su, H.: Atmospheric responses to the redistribution of anthropogenic aerosols, *J. Geophys. Res. Atmos.*, 120, 9625–9641, <https://doi.org/10.1002/2015JD023665>, 2015.

27

Deleted: 27

Wang, Y., Le, T., Chen, G., Yung, Y. L., Su, H., Seinfeld, J. H., and Jiang, J. H.: Reduced European aerosol emissions suppress winter extremes over northern Eurasia, *Nat. Clim. Chang.*, <https://doi.org/10.1038/s41558-020-0693-4>, 2020b.

Deleted: 2020

20 Wang, Z., Lin, L., Yang, M., Xu, Y., and Li, J.: Disentangling fast and slow responses of the East Asian summer monsoon to reflecting and absorbing aerosol forcings, *Atmos. Chem. Phys.*, 17, 11075–11088, <https://doi.org/10.5194/acp-17-11075-2017>, 2017.

Watanabe, M. and Tatebe, H.: Reconciling roles of sulphate aerosol forcing and internal variability in Atlantic multidecadal climate changes, *Clim. Dyn.*, 53, 4651–4665, <https://doi.org/10.1007/s00382-019-04811-3>, 2019.

25 Xie, S.-P. and Kosaka, Y.: What Caused the Global Surface Warming Hiatus of 1998–2013?, *Curr. Clim. Chang. Reports*, 3, 128–140, <https://doi.org/10.1007/s40641-017-0063-0>, 2017.

[Xie, S.-P., Lu, B., and Xiang, B.: Similar spatial patterns of climate responses to aerosol and greenhouse gas changes. *Nat. Geosci.*, 6, 828–832, <https://doi.org/10.1038/ngeo1931>, 2013.](#)

30 Xu, Y. and Xie, S.-P.: Ocean mediation of tropospheric response to reflecting and absorbing aerosols, *Atmos. Chem. Phys.*, 15, 5827–5833, <https://doi.org/10.5194/acp-15-5827-2015>, 2015.

[Xu, Y., Ramanathan, V., and Washington, W. M.: Observed high-altitude warming and snow cover retreat over Tibet and the Himalayas enhanced by black carbon aerosols. 16, 1303–1315, <https://doi.org/10.5194/acp-16-1303-2016>, 2016.](#)

Deleted:

35 Xu, Y., Lamarque, J. F., and Sanderson, B. M.: The importance of aerosol scenarios in projections of future heat extremes, *Clim. Change*, <https://doi.org/10.1007/s10584-015-1565-1>, 2018.

[Xu, Y., and X. Xu. Days with both extreme heat and extreme air pollution are becoming more common – which can't be a good thing for global health. *The Conversation*, 2020.](#)


40 Zhao, A., Bollasina, M. A., and Stevenson, D. S.: Strong Influence of Aerosol Reductions on Future Heatwaves, *Geophys. Res. Lett.*, 46, 4913–4923, <https://doi.org/10.1029/2019GL082269>, 2019.

Zheng, Y., Zhang, Q., Tong, D., Davis, S. J., and Caldeira, K.: Climate effects of China's efforts to improve its air quality, *Environ. Res. Lett.*, 15, 104052, <https://doi.org/10.1088/1748-9326/ab9e21>, 2020.

45

Deleted: 28

Figures



50

Figure 1:

(a) & (b): The global average anthropogenic emission of Sulfur and OC from 1920 to 2020 in the three sets of simulations used in this study. The “FF” represents the emission-related to Fossil Fuels. The “BB” represents Biomass Burning. Shading areas of 1980-2020 are the focused period.

55 (c) & (d): Similar to (a) & (b) but as the difference between ALL and corresponding fixed aerosol experiments to show the regional FF or BB emission. FF (blue lines) uses the left-hand side Y-axis and BB (green lines) uses the right-hand side Y-axis. Solid lines are for the “Western box” (West-box; 0-80°N, 120°W-40°E), and dashed lines are for the “Eastern box” (East-box; 0-80°N, 60°E-150°E). Boxes are shown in Fig. 2. All numbers in (c) and (d) are relative to the 1980 level to illustrate the change from 1980 to 2020.

60




Figure 2:

65 1st & 2nd row : 40-year trend of Sulfate and Primary OC (POM) column burden ($\text{kg}/\text{km}^2/\text{decade}$) in response to FF (left) from 1980 to 2020, and BB (right) emission changes. BC and SOA are not shown but the patterns are very similar to SO_4 .

3rd row: 40-year trend of AOD-AA (aerosol optical depth due to anthropogenic aerosols). Note the color scale of FF is twice of BB.

70 The blue dashed box in the upper left panel indicates the “Eastern Hemisphere” box (EH-box, $0^\circ\text{--}80^\circ\text{N}$, $60^\circ\text{--}150^\circ\text{E}$) used for the regional fixed-aerosol simulations. The solid box indicates the “Western Hemisphere” box (WH-box, $20^\circ\text{--}80^\circ\text{N}$, $130^\circ\text{--}10^\circ\text{W}$, and $30^\circ\text{--}80^\circ\text{N}$, $10^\circ\text{W}\text{--}40^\circ\text{E}$).

Deleted:),

Deleted: 30




Figure 3:

- 75 [Left column: the 40-year trend of \(top\) sulfate column burden \(kg/km²/decade\), and \(bottom\) surface air temperature in response to FF forcing from 1980 to 2020.](#)
- [Right column: Same as the left column but for the summation of EastFF and WestFF, obtained by first adding EastFF and WestFF responses together and then calculating the 40-year trend.](#)




Figure 4:

(1st row) 40-year trend in surface downward solar flux (FSDS, W/m²/decade) in response to FF (left), EastFF (mid), and WestFF (right); (2nd row) Top-of-Atmosphere net solar flux (FSNTOA, W/m²/decade); (3rd row) vertically-integrated droplet concentration (CDNUMC, %/decade); and (4th row) Total cloud fraction (CLDTOT, %/decade). The CDNUMC and CLDTOT patterns show the percentage trend relative to the 40-year climatology.

Dotted areas indicate the region where the trend passes the 95% significance test. The high latitude ocean regions with significant sea ice change (e.g. the Arctic and the Sea of Okhotsk) are masked to remove the surface albedo change effects. The dashed box in the left panel of 1st row indicates the “Eastern Hemisphere” (EH box, 80°E-140°W) used in our subsequent analysis, and the solid box indicates the “Western Hemisphere” (WH box, 90°W-30°E).

Deleted: Total cloud fraction (CLDTOT, 1/decade); and (3rd row)

Deleted:).

Deleted: 32




Figure 5:

(a) The Northern Hemisphere (NH) surface air temperature (SAT) anomalies (relative to 1920) in two sets of simulations: ALL (all forcing: historical+RCP8.5) (black line; 40-member), and Fix_FF1920 (blue line; 20-member). The four dashed lines indicate the four major volcanic eruption events (greater than category 5) during the 20th century. Shaded areas are the focused period of this study.

(b) The SAT responses to FF (blue). The thicker straight lines indicate the 40-year linear fits for 1940-1980 and 1980-2020 respectively. (c) & (d) are similar to (b) but for the EH and WH boxes (see boxes in Fig. 4).

Deleted: 4


Deleted: the three

Deleted: eastern part of NH

Deleted: western part of NH

Deleted: 3

Deleted: 33



10

Figure 6:

(a-c) The 1980-2020 trend of zonal mean meridional stream function (ZMMSF, 10^9 kg/s/decade) in response to (a) FF, (b) EastFF, and (c) WestFF. The positive values (solid lines for climatology and red shading for long-term trend) indicate clockwise circulation. The positive values indicate westerly wind. The statistically insignificant trend is masked in white. The arrows indicate the latitudinal shift of Hadley cells.

15

(d-f) Similar to (a-c) but for zonal mean zonal wind (U, m/s/decade). The latitudinal profiles below show the Top-of-Atmosphere net solar flux trend (FSNTOA as in Fig. 5c; $W/m^2/decade$). The arrows indicate the latitudinal shift direction of the NH Jetstream core.

20

The profiles below show the 40-year trend of TOA net solar flux (FSNTOA; grey curves; in units of $K/decade$), and the gradient of FSNTOA trend (black curves; in units of $W/m^2/decade/10^\circ lat$). The trend and gradient lines are smoothed using the moving average method (with 30 degrees of latitudinal range sampling window). The dashed lines are the linear fit from 0° to $60^\circ N$.

34

Deleted: 5

Moved down [6]: The arrows indicate the latitudinal shift of Hadley cells.(d-f) Similar to (a-c) but for zonal mean zonal wind (U, m/s/decade).

Moved down [7]: (d-f) Similar to (a-c) but for zonal mean zonal wind (U, m/s/decade).

Deleted: 4

The latitudinal profiles below show surface downward solar flux trend (FSDS as shown in Fig. 3; grey curves; $W/m^2/decade$), and its latitudinal gradient (FSDS_gradient; black curves; $W/m^2/decade/10^\circ lat$) smoothed in 30 degrees of latitudinal range, in which positive gradient indicates a larger FSDS trend at higher NH latitude. The dashed lines are the linear fit from 0° to $60^\circ N$.

Moved (insertion) [6]

Deleted: The latitudinal profiles below show the Top-of-Atmosphere net solar flux trend (FSNTOA as in Fig. 4c; $W/m^2/decade$), and its latitudinal gradient (FSNTOA gradient; $W/m^2/decade/10^\circ lat$).

Moved (insertion) [7]

Deleted: 34





Figure 7:

(a) The 40-year climatology of northward energy transport (Pwatt) is calculated based on ALL experiments ensemble average. (b–d) The decadal trend of northward energy transport in response to FF, EastFF, and WestFF (Pwatt/decade), which are obtained by subtracting the fixed single forcing experiments from the ALL experiment.

The dashed blue lines represent the oceanic energy transport; the solid red lines represent the atmospheric energy transport.

Moved (insertion) [8]

Deleted: 35



50 **Figure 8:**

(a–c) Similar to Fig. 6d–f but for geostrophic wind (U_g) trend (m/s/decade) derived from geopotential height (Z) during 1980–2020 in Eastern Hemisphere (EH). The climatology is shown in contour lines with an interval of 6. The positive values indicate westerly wind. The statistically insignificant trend is masked in white. The arrows indicate the latitudinal shift in each case.

55 The profiles below show the gradient of surface air temperature trend (black curves; in units of $W/m^2/decade/10^\circ lat$). The gradient lines are smoothed using the moving average method (with 30 degrees of latitudinal range, sampling window). The dashed lines are the linear fit from 0° to $60^\circ N$. Positive values indicate a larger SAT trend at higher latitude, and vice versa.

60 (d–f) Similar to the bottom row of Fig. 6 but for the U trend in the Eastern Hemisphere (EH) in response to three types of forcings.

Moved (insertion) [9]

Deleted: 6 (d

Deleted:)

Deleted: latitudinal

Deleted: (SAT)

Deleted: in


Deleted: ; in unit of $K/decade/10^\circ lat$.

Deleted: The dashed lines are the linear fit from 0° to $60^\circ N$.

Deleted: (d–f)

Deleted:).

Deleted: 36



Moved up [8]: Figure 7: 1

Figure 9:

Ist row: ZMMSF trend, Same as Fig. 6 (a–c) but showing the difference between global mean and EH mean (a–b), and the difference between global mean and WH mean (g–h). The latitudinal profiles below show the latitudinal gradient of the FSDS trend, which is also calculated as the difference between EH/WH and Globe.

2nd row: U trend, Same as Fig. 8 (d–f) but showing the difference between global mean and EH (WH). The latitudinal profiles below show the latitudinal gradient of SAT trend, which is also calculated as the difference between EH (WH) and Globe.

3rd row: Similar to the above two rows but showing tropospheric air temperature (T) trend.

37

Deleted: 5

Deleted: 6

Deleted: /

Deleted: 37

75

80




Figure 10:

1st & 2nd rows: 40-year sea surface air temperature trend (SAT) and net energy flux at TOA (R_{toa}). Arctic regions with significant sea ice changes are masked.

The blue (red) circles in the 2nd row indicate the major regions with a decline (increase) of aerosol emission.

3rd row: 40-year trends of zonal mean meridional heat transport (ZMMHT) in EH and WH. The positive values (solid lines for climatology and red shading for long-term trend) indicate poleward transfer. The arrows indicate the energy transfer responses relative to the climatology.

4th row: Similar to Fig. 7 but showing the regional northward energy transport ($P_{watt/decade}$; northward positive) in EH and WH. The dashed blue lines indicate oceanic northward energy transport (OET), while the solid red lines indicate atmospheric northward energy transport (AET).

Moved up [9]: Figure 8: [↗](#)

Deleted: trends of Sea

Deleted: (SSAT),

Deleted: that of

Deleted: including shortwave and longwave, not FSNTOA shown in Fig. 4).

Deleted: in

Deleted: significant

Deleted: of

Deleted: 38

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [1] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [2] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [2] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [3] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [4] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [4] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [4] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [4] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [4] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 14: [4] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 15: [5] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 16: [6] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 16: [7] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 16: [8] Deleted Diao, Chenrui 10/26/21 4:45:00 PM

▼
Page 16: [9] Deleted Diao, Chenrui 10/26/21 4:45:00 PM