



Aerosol radiative impact during the summer 2019 heatwave produced partly by an inter-continental Saharan dust outbreak – Part 2: Long-wave and net dust direct radiative effect

5 Michaël Sicard^{1,2}, Carmen Córdoba-Jabonero³, María-Ángeles López-Cayueta³, Albert Ansmann⁴,
Adolfo Comerón¹, María-Paz Zorzano^{5,6}, Alejandro Rodríguez-Gómez¹, Constantino Muñoz-Porcar¹

¹CommSensLab, Dept. of Signal Theory and Communications, Universitat Politècnica de Catalunya (UPC), 08034-Barcelona, Spain

²Ciències i Tecnologies de l'Espai-Centre de Recerca de l'Aeronàutica i de l'Espai/Institut d'Estudis Espacials de Catalunya (CTE-CRAE/IEEC), Universitat Politècnica de Catalunya (UPC), 08034-Barcelona, Spain

10 ³Instituto Nacional de Técnica Aeroespacial (INTA), Atmospheric Research and Instrumentation Branch, Torrejón de Ardoz, 28850-Madrid, Spain

⁴Leibniz Institute for Tropospheric Research (TROPOS), 04318-Leipzig, Germany

⁵Centro de Astrobiología (CSIC-INTA), Ctra. Ajalvir, km. 4, Torrejón de Ardoz, 28850-Madrid, Spain

⁶School of Geosciences, University of Aberdeen, Aberdeen, AB24 3FX, UK

15

Correspondence to: Michaël Sicard (msicard@tsc.upc.edu)

Abstract. This paper is the companion paper of Córdoba-Jabonero et al. (2021). It deals with the estimation of the longwave (LW) and net dust direct radiative effect (DRE) during the dust episode that occurred between 23 and 30 June, 2019, and coincided with a mega-heatwave. The analysis is performed at two European sites where polarized-Micro-Pulse Lidars ran
20 continuously to retrieve the vertical distribution of the dust optical properties: Barcelona, Spain, 23-30 June, and Leipzig, Germany, 29-30 June. The radiative effect is computed with the GAME radiative transfer model separately for the fine- and coarse-mode dust. The instantaneous and daily radiative effect and radiative efficiency (DREff) are provided for the fine-mode, coarse-mode and total dust at the surface, top of the atmosphere (TOA) and in the atmosphere. The fine-mode daily LW DRE is small (< 6 % of the shortwave (SW) component) which makes the coarse-mode LW DRE the main modulator of the
25 total dust net DRE. The coarse-mode LW DRE starts exceeding (in absolute values) the SW component in the middle of the episode which produces positive coarse-mode net DRE at both the surface and TOA. Such an unusual tendency is attributed to increasing coarse-mode size and surface temperature along the episode. This has the effect of reducing the SW cooling in Barcelona up to the point of reaching total dust net DRE positive (+0.9 W m⁻²) on one occasion at the surface and quasi-neutral (-0.6 W m⁻²) at TOA. When adding the LW component, the total dust SW radiative efficiency is reduced by a factor 1.6 at
30 both surface (on average over the episode, the total dust net DREff is -54.1 W m⁻² τ⁻¹) and TOA (-37.3 W m⁻² τ⁻¹). A sensitivity study performed on the surface temperature and the air temperature in the dust layer, both linked to the heatwave and upon which the LW DRE strongly depends, shows that the heatwave contributed to reduce the dust net cooling effect at the surface



and that it had nearly no effect at TOA. Its subsequent effect was thus to reduce the heating of the atmosphere produced by the dust particles.

35 **1 Introduction**

It now makes no doubt that extreme air temperatures have an effect on mortality (Basu and Samet, 2002) and so have mineral dust intrusions (Diaz et al., 2017). Studies of heatwaves and elevated temperatures episodes indicate that the main underlying mechanism leading to death is that stress on the cardiovascular and respiratory systems increases during periods of high air temperature (Kunst et al., 1993). During periods of mineral dust intrusions, the effect on human health is associated to the
40 biological matter and microorganisms transported in the dust plume that can be harmful to humans (Griffin et al., 2007) and/or to the increase of the particulate matter load at ground level when dust is present (Tobias et al., 2011). When both atmospheric phenomena (heatwave and mineral dust intrusion) occur simultaneously, their effects may cumulate. A place on Earth with a high population density and a high probability that both phenomena occur at the same time is southern Europe because of its proximity to the North African deserts, the largest source of mineral dust on Earth (Prospero et al., 2002).

45 A recent study from Sousa et al. (2019) quantifies the proportion of Saharan dust intrusions over southwestern Europe since 2000 which were actually accompanied by a heatwave. They find that 39 and 14 % of the Saharan dust intrusions over the western and eastern Iberian Peninsula (IP), respectively, were accompanied by a heatwave. Although dust intrusions are more frequent in the eastern (868 cases) than in the western IP (295 cases), their transport towards the eastern IP is often characterized by an accentuated SW-NE tilt in the mean position of the 580-dam geopotential height thickness, which favors
50 heatwaves in the central Mediterranean, with little incidence in the eastern IP. In their introduction Sousa et al. (2019) stress “an incomplete description of the relationship between such air masses [Saharan dust] originated in the desertic region of northwestern Africa and heatwave episodes” in the IP.

A recent event of heatwave accompanied with a Saharan dust intrusion occurred in June 2019. The event was outstanding in the sense that temperature records were broken in several places in Europe. The month of June 2019 is still the hottest June
55 ever recorded at European level (Copernicus, 2021). Average temperatures were more than 2°C above normal and if we consider the 5-day period 25-29 June the temperatures were 6 to 10°C above normal, with local differences even higher in NE Spain, France and the United Kingdom according to a detailed article issued by the Spanish state meteorological agency (AEMET) on the “June 2019 heatwave in the context of the climate crisis” (AEMET, 2019). Although quite recent, the heatwave of 25-29 June 2019, classified as mega-heatwave by some authors, has already been the subject of several studies.
60 Xu et al. (2020a; 2021) quantified the role of dynamical and thermodynamical processes in triggering this extreme event and investigated how changes of these processes observed over the last decades may have affected the occurrence probability of such extreme events. Ma et al. (2020) and Vautard et al. (2020) used climate models to quantify the role of human contribution through the anthropogenic climate change. Sousa et al. (2019) studied the relationship between heatwaves and Saharan warm air intrusions in the IP in the long-term context. Finally Córdoba-Jabonero et al. (2021) studied the shortwave (SW) dust direct
65 radiative effect during 23-30 June 2019 at two sites in Barcelona (BCN), Spain, and Leipzig (LPZ), Germany.

The present paper is the companion paper of Córdoba-Jabonero et al. (2021). It deals with the estimation of the longwave and net dust direct radiative effect during 23-30 June 2019. This second part of the study is motivated by the high temperatures caused by the heatwave that accompanied with the dust event. Indeed, high temperature, clear skies and high insolation imply high land surface temperature and thus strong longwave radiation emission. The paper is organized as follows: the description
70 and discussion of the dust microphysical and optical properties in the longwave spectral range are exposed in Section 2 (instruments, methodology and radiative effects in the shortwave spectral range can be found in the companion paper); Section 3.1 includes the results in terms of longwave and net dust direct radiative effect and radiative efficiency and Section 3.2



presents a sensitivity study on the relationship between heatwave (by means of surface temperature and air temperature in the dust layer) and coarse-mode longwave and total dust radiative effects. Conclusions are given in Section 4.

75 2 Dust radiative properties in the longwave spectral range and GAME parametrization

The longwave (LW) module of the GAME radiative transfer model (Dubuisson et al., 1996; 2004; 2005) has been used in a recently increasing number of studies (Sicard et al., 2014a; 2014b; Barragan et al., 2016; 2017; 2020; Granados-Muñoz et al., 2019a; 2019b). GAME calculates spectrally integrated, upward and downward radiative fluxes in 40 plane and homogeneous layers from 0 to 100 km with a 1-km resolution from 0 to 25 km. Spectral limits were set in terms of wavenumber from 200 to 2500 cm^{-1} (wavelength: 4 – 50 μm) at a fixed resolution of 20 cm^{-1} (115 points). GAME accounts for thermal emission, absorption and scattering, as well as their interactions, using the Discrete Ordinates Method (DISORT) (Stamnes et al., 1988). Gaseous absorption (H_2O , CO_2 , O_3 , N_2O , CO , CH_4 and N_2) is treated from the correlated k distribution (Lacis and Oinas, 1991). More details of the longwave module of GAME can be found in Sicard et al. (2014a). GAME presents the advantage of the complete representation of the longwave aerosol scattering, in addition to their absorption. The moderate spectral resolution of GAME makes it possible to account for the spectral variations of aerosol properties, especially in the infrared window. The spectral optical properties of aerosols are defined for each atmospheric layer where dust is present: the single scattering albedo (SSA) and the asymmetry factor (asyF) are assumed constant vertically; the extinction coefficient (α) varies with altitude. GAME outgoing (i.e. leaving the terrestrial atmosphere) longwave radiation (OLR) was validated through comparison with CERES (Clouds and the Earth's Radiant Energy System) OLR measurements in 11 cases of dust intrusion in Barcelona (Sicard et al., 2014a). Their results indicate a bias between simulated and measured OLR of -0.8 % and a root mean square error of 2.52 W m^{-2} .

The convention followed for the definition of the dust direct radiative effect (DRE) at either the surface (SRF) or the Top-Of-the-Atmosphere (TOA) is the one of Eq. (9) of Córdoba-Jabonero et al. (2021). The atmospheric (ATM) DRE is the difference between the TOA DRE and the SRF DRE. To simplify abbreviations the shortwave, longwave and net (shortwave + longwave) DRE are noted DRE_{SW} , DRE_{LW} and DRE_{NET} , respectively.

GAME is used to calculate the instantaneous longwave radiative effect in a hourly basis between 5 and 19 UTC over the 8 days of the event in Barcelona (23-30 June) and over 2 days in Leipzig (29 June, Episode 1, and 30 June, Episode 2; see Córdoba-Jabonero et al., 2021). In the rest of this section the dust microphysical and radiative properties are sometimes averaged over the whole event. For Barcelona this means from 23 to 30 June or from 24 to 29 June in order to avoid the beginning and the end of the dust intrusion when abrupt changes in the microphysics and radiative properties are expected. In Leipzig the dust microphysical and radiative properties are averaged over the afternoons of 29 and 30 June (when the dust was present, see Figures 5 and 6 of Córdoba-Jabonero et al., 2021), respectively noted 29J-pm and 30J-pm, corresponding to the two dust episodes described in the former reference. In all cases the averaging period is always indicated.

2.1 Dust microphysics

The dust radiative properties in the longwave spectral range were calculated using a Mie code in the range 4 – 50 μm . The input of our Mie code is the geometric median radius, r_g , and its standard deviation, σ_g , of the lognormal distribution, the particle number, and the spectrally-resolved refractive index. The code calculates the extinction coefficient normalized to that at 532 nm (the wavelength of the P-MPL systems) spectrally-resolved in the range 4 – 50 μm , α/α_{532} , the single scattering albedo and the asymmetry factor. Although the particle number is provided in input, it has no effect on our calculations since SSA and asyF are intensive parameters and the extinction is normalized to that at 532 nm, thus the dependence on the particle



number is removed. We distinguish between the dust coarse-mode (D_c) and the dust fine-mode (D_f). The total dust (DD) is the sum of both D_c and D_f components.

The spectral refractive index (real and imaginary part) is the same than in Sicard et al. (2014a) and comes from measurements of long range transport mineral dust taken in Meppen in western Germany (Volz, 1983). The table giving the refractive index as a function of the wavelength was found in Krekov (1993). The spectral variation of both real and imaginary parts of the refractive index can be seen in Figure 1 of Sicard et al. (2014a).

For each of the coarse and fine modes, the geometric median radius and the standard deviation were estimated from column-integrated AERONET retrievals. AERONET provides volume median radius, r_v , and its standard deviation, σ_v . The geometric median radius and standard deviation were calculated as follows:

$$r_g = r_v e^{-3(\ln \sigma_g)^2} \quad (1)$$

$$\sigma_g = \sigma_v \quad (2)$$

AERONET V3L2.0 data were used at both sites. 36 retrievals are available in BCN over the period 23-30 June, and 13 retrievals are available in LPZ for both days of 29 and 30 June. To have a hourly estimation of both r_g and σ_g both parameters were interpolated. Both retrievals and hourly interpolated values for r_v , σ_v and r_g are represented as a function of time in Figure 1 at both sites and for both size modes. In BCN, the mean (over 23-30 June) D_c and D_f geometric median radii (and standard deviation) are 1.262 (0.652) and 0.063 (0.600) μm , respectively; in LPZ we find for the mean of 29J-pm and 30J-pm: $r_g(D_c) = 1.399 \mu\text{m}$ ($\sigma_g(D_c) = 0.680$) and $r_g(D_f) = 0.032 \mu\text{m}$ ($\sigma_g(D_f) = 0.505$). Interestingly, the mean large dust particles in LPZ are larger than in BCN and their sizes distribute over a larger scatter interval; $r_g(D_c)$ in LPZ on 29J-pm and 30J-pm ($\sim 1.4 \mu\text{m}$) is similar to $r_g(D_c)$ in BCN on 29 and 30 June, possibly indicating a conservation of the coarse mode size distribution during transport between both sites. The small dust particles in LPZ are on average much smaller than in BCN (Figure 1b). It is known that dust aging encompasses many processes that can alter the dust chemical composition, shape and size. In particular, when the transport of mineral dust occurs over polluted regions anthropogenic inorganic acids can be absorbed by the dust surface forming hygroscopic salt compounds that coat the dust particles (Abdelkader et al., 2015; Athanasopoulou et al., 2016). Dust also favors the formation of secondary pollutants (Querol et al., 2019; Xu et al., 2020b). Both processes (acid absorption and secondary aerosol formation) can lead to dust particle growth in different size ranges. Secondary aerosol formation is enhanced in stagnant (low winds) and high humidity conditions (Xu et al., 2020b). NCEP (National Centers for Environmental Prediction) reanalysis horizontal wind in BCN (not shown) reveals strong winds ($\sim 18 \text{ m}$




Figure 1. AERONET volume median radius and standard deviation and corresponding geometric median radius in BCN (red) and LPZ (blue) for (a) the coarse mode and (b) the fine mode. Circles indicate retrievals and the dash lines represent the hourly interpolated data. The legend in the first plot applies to all plots. The black lines are a linear fit of the retrievals in BCN over the period 24-29 June.



s^{-1}) at the beginning of the episode and much more stagnant conditions ($<5 \text{ m s}^{-1}$) towards the end of the episode. All in all, such conditions might have favored the transfer of small dust particles to larger sizes and the growth of coarse-mode particles along the transport. A linear fit (black lines in Figure 1) has been applied to the BCN retrievals of r_g and σ_g over the period 24-29 June in order to avoid the beginning and the end of the dust intrusion when abrupt changes in the microphysics are expected. Coarse and fine modes have opposite tendencies: $r_g(Dc)$ increases at a rate of almost $+10 \text{ \% day}^{-1}$ and its scatter interval (σ_g) also increases ($\sim +2 \text{ \% day}^{-1}$); $r_g(Df)$ slightly decreases (at a rate of $\sim -1 \text{ \% day}^{-1}$) and its scatter interval (σ_g) also decreases ($\sim -2 \text{ \% day}^{-1}$). The coarse mode radius increase and widening reflects the possible size growth mentioned earlier that might have occurred during the dust transport. This fact is also nicely illustrated by Figure 2 in which both daily Dc and Df normalized size distributions are represented on a day-by-day basis. One sees clearly how the Dc radii increase between 24 and 29 June, while the Df radii decrease during the same period. Although the decreasing rate of the Df radius along the event is small ($<1 \text{ \% day}^{-1}$), its representation in a logarithmic scale (X axis) in Figure 2 shows clearly that it is noteworthy.

2.2 Dust radiative properties in the longwave spectral range

With the microphysics defined in the previous section, spectral dust radiative properties (α , SSA and asyF) in the longwave spectral range are calculated with the Mie code on a hourly basis at both sites. Results are presented in Figure 3 separately at BCN and LPZ and for Dc and Df. In BCN a color code is used for each curve corresponding to the number of hours past since the beginning of the period considered: 23 June at 00 UTC. In LPZ the average of 29J-pm (Episode 1) and 30J-pm (Episode 2) is shown. The effect of the increasing $r_g(Dc)$ along the dust event in BCN is visible on the Dc extinction plot (Figure 3a, top panel): α/α_{532} is smaller at the beginning of the event (blue-green curves below the mean) than at the end (red-brown curves above the mean). The main differences between Dc and Df are: the Df extinction is at least three orders of magnitude smaller than the Dc one (in LPZ it is even smaller); Df SSA converges rapidly towards 0 with increasing wavelengths and so does Df asyF. Differences between BCN and LPZ are essentially due to the differences in the dust radii at each site: the mean Dc extinction in BCN is slightly lower than in LPZ ($r_g(Dc)$ is smaller in BCN than in LPZ) and the mean Df extinction in BCN is larger than in LPZ ($r_g(Df)$ is twice larger in BCN than in LPZ). SSA and asyF are similar at both sites.





Figure 2. Day-by-day evolution of the daily mean normalized lognormal distributions in Barcelona between 24 and 29 June calculated with the daily mean (r_g , σ_g) of Dc (solid lines) and Df (dash lines).

The profiles of extinction coefficients at 532 nm for both coarse and fine modes were taken from the POLIPHON inversion performed by Córdoba-Jabonero et al. (2021). $\alpha_{532}(Dc)$ and $\alpha_{532}(Df)$ are averaged in 1-km layer-mean values from 0 to 10 km and then input in GAME. In each layer GAME calculates the LW spectral extinction by multiplying the measured α_{532} by the calculated normalized extinction, α/α_{532} . Figure 4 represents the Dc and Df dust optical depth (DOD) at 532 nm at both sites. In BCN Dc and Df DOD peak at 0.420 and 0.090, respectively. In LPZ both episodes are much more




175 moderate ($DOD(D_c)$ peaks at 0.072 and $DOD(D_f)$ at 0.045). Except on 24 and 25 June, the Dc DOD variations in BCN are smooth; for the fine mode DOD swings up and down. In LPZ, DOD outside the dust episodes is very small (< 0.006).

To get an idea of the dust stratification we plot in Figure 5 the dust layer center of mass (CoM) of both size modes and at both sites. A striking feature is the shape similarity in BCN between Dc DOD (Figure 4a) and Dc CoM (Figure 5a). Such similarity is usually not expected and probably reflects the homogeneous vertical mixing of the dust plume arriving in BCN all along the intrusion. In BCN the Dc and Df CoM are quite similar. A peak at ~ 4 km is observed at the beginning of the event and next a general temporal decreasing tendency is observed. Towards the end of the intrusion CoM is ~ 2 km. In LPZ the centers of mass of the Dc layer are similar to the CoM in BCN at the beginning of the event (23-24 June). Dc CoM of Episode 1 (3.7 – 4.5 km) is higher than that of Episode 2 (3.3 – 3.6 km) as noted by Córdoba-Jabonero et al. (2021). It is noteworthy to mention that the dust CoM in BCN falls within the climatological dust bottom and top heights, 1.5 and 3.5 km, respectively,



185

Figure 3. Spectral normalized extinction, α/α_{532} , SSA and asyF in BCN for (a) Dc and (b) Df, and in LPZ for (c) Dc and (d) Df. In BCN all hourly calculations are represented: 192 (8 days). In LPZ the average of 29J-pm (Episode 1) and 30J-pm (Episode 2) is represented. In the BCN plots the thick black line represents the mean over the whole period (23-30 June).



190 Figure 4. Hourly dust optical depth calculated from the extinction profiles for (a) the coarse mode and (b) the fine mode obtained using the POLIPHON algorithm in BCN (red) and LPZ (blue).




Figure 5. Hourly dust layer center of mass (CoM) for (a) the coarse mode and (b) the fine mode in BCN (red) and LPZ (blue).

estimated by Papayannis et al. (2008) from lidar measurements in BCN.

195 2.3 GAME parametrization

Finally the rest of input in GAME are the atmospheric profiles and the Earth surface properties. The gas absorption is parametrized from profiles of pressure, temperature and relative humidity. In BCN, radiosoundings launched twice a day (at 00 and 12 UTC) by the University of Barcelona in collaboration with the Servei Meteorològic de Catalunya, the Catalonia meteorological agency, were used. No radiosoundings are available in LPZ, thus the 6-hour profiles from the Global Data Assimilation System (GDAS) provided by the National Oceanic and Atmospheric Administration (NOAA) were used instead.

The Earth surface is assumed lambertian in GAME and its albedo in the longwave spectral range was set to a constant value of 0.017. This value corresponds to the climatological value found by Sicard et al. (2014a) in Barcelona from CERES measurements in the spectral window 8.1 – 11.8 μm and averaged over spring and summer seasons in the period June 2007 – May 2012. The same value was used at both sites.

We used the hourly land surface temperature (LST) V2 product provided globally at 5-km resolution by the Copernicus Global Land Service (<https://land.copernicus.eu/global/products/lst>). The LST V2 datasets are estimated from TOA brightness temperatures from the infrared spectral channels of a constellation of geostationary satellites, Meteosat Second Generation being the one covering Europe. Its estimation further depends on the albedo, the vegetation cover and the soil moisture. Further details about the temperature retrieval can be found in Freitas et al., (2013). The LST V2 product user manual and validation reports can be found at https://land.copernicus.eu/global/products/lst?qt-lst_characteristics=5#qt-lst_characteristics. Figure 6 shows the hourly LST in Barcelona (23-30 June) and Leipzig (29-30 June). The diurnal cycle is nicely visible at both sites on all days. In Barcelona the temperature steadily increases between 23 and 28 June. On 29 June the LST starts its decrease. A maximum LST of 45.22 $^{\circ}\text{C}$ is reached in the afternoon of 28 June. In Leipzig the LST increase is even more pronounced than in Barcelona. The maximum night/day LST difference is 32.2 $^{\circ}\text{C}$ on 29 June and the maximum LST (46.2 $^{\circ}\text{C}$) is reached in the afternoon of 30 June.




Figure 6. Hourly LST in BCN (red) and LPZ (blue). The associated uncertainty is added as a shaded area, although it is so small that it is almost imperceptible.

3 Results

The results are first discussed for the longwave spectral range and for the whole spectral range (net effect = shortwave + longwave) in terms of dust direct radiative effect (Section 3.1.1), then in terms of radiative efficiency (Section 3.1.2) and a deeper analysis is performed at BCN site (Section 3.2). All shortwave magnitudes were taken from the Part 1 paper by



Córdoba-Jabonero et al. (2021). Instantaneous (from 5 to 19 UTC) and daily averages are presented. Daily DRE_{LW} is difficult to estimate because GAME calculates longwave fluxes only during daytime hours, so nighttime DRE_{LW} estimations are not available, although they are not zero unlike in the shortwave spectral range. Here the daily DRE_{LW} has been calculated as the average of the 15 estimations available everyday (from 5 to 19 UTC). Di Sarra et al. (2011) used the same method averaging instantaneous DRE_{LW} obtained every 6 hours. Meloni et al. (2015) assumed their instantaneous DRE_{LW} retrieval constant throughout the 24 hours.

3.1 Longwave and net dust direct radiative effect

3.1.1 Dust direct radiative effect

The calculations of the instantaneous dust DRE_{LW} are presented as a function of time in both BCN and LPZ in Figure 7 separately for Dc, Df and DD and at SRF, TOA and in the atmosphere. At BCN site we also plot the temporal evolution of the Df/DD DRE_{LW} ratio along the eight days of the event. Daily values of Dc, Df and DD DRE_{LW} are reported at SRF in Table 1 and at TOA in Table 2. Both size modes produce a positive radiative effect at both SRF and TOA. Except on a couple of occasions on 25 June, DRE_{LW} at SRF is always larger than at TOA resulting in a negative atmospheric DRE_{LW} , indicator of a cooling effect of the atmosphere.

Independently of the atmospheric level where it is estimated, the Df DRE_{LW} is small: it is in general more than one order of magnitude smaller than Dc DRE_{LW} , and it is smaller at TOA than at SRF. In BCN LW Df instantaneous values do not exceed $+0.6 \text{ W m}^{-2}$ (at SRF, Figure 7) and the daily values are smaller than $+0.42 \text{ W m}^{-2}$ (Table 1). Instantaneous Df DRE_{LW} are in the range of values of Sicard et al. (2014b) also estimated for the dust fine mode: $[+0.4; +0.7 \text{ W m}^{-2}]$ and $[+0.1; +0.3 \text{ W m}^{-2}]$ at SRF and TOA, respectively. In LPZ instantaneous (daily) Df DRE_{LW} are smaller than $+0.2 (+0.08) \text{ W m}^{-2}$. In BCN the contribution of Df DRE_{LW} to the total dust DRE_{LW} is smaller than 5-6 % (Figure 7c). In average over the period 24-30 June (23 June is discarded because it is the day the dust arrived in BCN) it is 3.6 and 2.5 % at SRF and TOA, respectively, and it seems to slightly decrease with time along the event (negative slope of the fittings in Figure 7c). Hence Df DRE_{LW} is expected to have a very little effect on DRE_{NET} with respect to Dc DRE_{LW} . This is also true with respect to Df DRE_{SW} as nicely illustrated in Figure 8 which shows for Dc, Df and DD the daily LW/SW DRE ratio in absolute value at SRF, TOA and ATM. In the middle plot (Df) the LW/SW DRE ratio is not higher than 10.0 and 5.2 % at SRF and TOA, respectively. The average over the whole event is 6.7 and 3.7 % at SRF and TOA, respectively. Note that the decrease of the Df-to-DD DRE_{LW} ratio with respect to time (opposite of what was observed for DRE_{SW} , see Córdoba-Jabonero et al. (2021)) is due to a combination of various factors such as $\tau_g(Dc)$ (increases), $\tau_g(Df)$ (decreases), Dc DOD (decreases) and Df DOD (constant).

Unlike the Df component, Dc DRE_{LW} is quite significant. It is also smaller at TOA than at SRF. In BCN Dc instantaneous values reach $+11.4 \text{ W m}^{-2}$ (at SRF) and the daily values are as high as $+9.8 \text{ W m}^{-2}$. In LPZ the instantaneous and daily values (at SRF) do not exceed $+3.3$ and $+1.3 \text{ W m}^{-2}$, respectively. For comparison, Sicard et al. (2014b) found instantaneous Dc DRE_{LW} in the range $[+2.7; +10.2]$ and $[+0.5; +4.7 \text{ W m}^{-2}]$ at SRF and TOA, respectively. As far as the daily LW/SW Dc DRE ratio is concerned (Figure 8), one sees that at both SRF and TOA it increases with time along the event. It is ~ 80 % (at both SRF and TOA) on the first day of the event and it reaches 170 % at SRF (on 28 June) and 183 % at TOA (on 30 June). On the last four days of the event, the daily LW/SW Dc DRE ratio is higher than 119 %, indicating that Dc DRE_{SW} is going to be fully compensated by Dc DRE_{LW} , and that, on the daily time scale, the sign of Dc DRE_{NET} will be that of the longwave component, i.e. a positive (warming) DRE. This result is further explored in Section 3.2.

Because Df DRE_{LW} is small compared to Dc DRE_{LW} , the total dust DRE_{LW} is similar to that of the coarse mode. In BCN instantaneous values do not exceed $+12.1 \text{ W m}^{-2}$ at SRF and $+7.9 \text{ W m}^{-2}$ at TOA. The average over the whole event of the daily values are $+6.12 \text{ W m}^{-2}$ at SRF and $+3.50 \text{ W m}^{-2}$ at TOA, respectively, resulting in a LW/SW DD DRE ratio of 67 and 60 %. The fraction of the LW component compared to the SW one is thus similar at both SRF and TOA, and it is higher




than other estimations from the literature. Di Sarra et al. (2011) found daily LW/SW ratios of 49 and 35 % at SRF and TOA, respectively, and Meloni et al. (2015) of 52 and 26 %. Both works deal with desert dust measurements in the Mediterranean, i.e. above the sea. The explanation why our daily LW/SW ratios are higher than those reported in the literature probably lies in the surface temperature. Indeed the LST at solar noon used here is over +15°C higher than the sea surface temperature considered in Di Sarra et al. (2011) and Meloni et al. (2015), which contributes to enhance DRE_{LW} . In LPZ instantaneous values do not exceed +3.3 W m⁻² at SRF and +3.4 W m⁻² at TOA. The average over the period 29-30 June of the daily values are +0.96 W m⁻² at SRF and +1.04 W m⁻² at TOA, respectively. It is interesting to observe that in LPZ DRE_{LW} is higher at TOA than at SRF (in BCN the reverse occurs). As shown by Dufresne et al. (2002) and Sicard et al. (2014a) DRE_{LW} is highly dependent on the dust layer height and its variations with the latter at SRF and TOA are opposite (DRE_{LW} decreases at SRF with increasing height while it increases at TOA). The height of the coarse dust layer is much higher in LPZ than in BCN (Figure 5a), which enhances DRE_{LW} at TOA with respect to the surface. The DD DRE_{LW} does not follow systematically a constant diurnal cycle like the SW one does. On the first three days no pattern is identified, probably because of the strong variations of the DOD (see Figure 4). On 26 June DD DRE_{LW} is rather flat, while on the last four days of the event both SRF and TOA DRE_{LW} have a clear diurnal cycle with the shape of a semi cosine curve (28 June) or an inverted-V (27, 29 and 30 June) showing an increase in the morning, a maximum reach at 12 or 13 UTC and a decrease afterward. This pattern of the total dust DRE_{LW} is mostly due to the coarse mode since the fine mode DRE_{LW} is much smaller than the coarse one. The diurnal cycle of DD (or Dc) DRE_{LW} is more pronounced at TOA than at SRF. It is well known that the OLR has a marked diurnal cycle (Slingo et al., 1987) and that it is highly correlated to the surface heating and cooling due to the diurnal cycle of insolation (Chung et al., 2009). For DRE_{LW} these statements are less straightforward. When holding all parameters constant except the surface temperature, Osborne et al. (2011) showed with Saharan dust measurements performed in Mauritania and Niger in June 2007 that the shape of the diurnal cycle of DRE_{LW} mimicked that of the surface temperature. In our analysis the surface temperature is also the only variable with a diurnal cycle with the shape of a cosine curve, so we believe that the cosine or inverted-V shape of DD DRE_{LW} observed on the last four days of the dust event is related to the diurnal cycle of the surface temperature.

The instantaneous dust DRE_{NET} ($DRE_{SW} + DRE_{LW}$) is presented as a function of time in both BCN and LPZ in Figure 9 separately for Dc, Df and DD and at SRF, TOA and in the atmosphere. Daily values of Dc, Df and DD DRE_{NET} are reported at SRF in Table 1 and at TOA in Table 2. The fine mode DRE_{NET} is similar to DRE_{SW} since $DRE_{LW} \ll DRE_{SW}$. It produces a negative net radiative effect at both SRF and TOA. The daily mean values (averaged over the whole event) are -3.2 (BCN) and -1.15 (LPZ) W m⁻² at SRF and -2.5 (BCN) and -0.85 (LPZ) W m⁻² at TOA. Dc and DD instantaneous DRE_{NET} at SRF and TOA are similar to their corresponding DRE_{SW} but shifted towards positive values (Figure 9). At SRF daily Dc DRE_{NET} is negative in the first half period of the event in BCN and in LPZ. In BCN the average over the whole event is +0.2 W m⁻², i.e. that at SRF the LW component wins over the shortwave one at the event scale. At TOA daily Dc DRE_{NET} is essentially positive in BCN (except on the first three days of the event) and in LPZ. In BCN (LPZ) the average over the whole event is +0.2 (+0.5) W m⁻², i.e. that at TOA the LW component wins over the shortwave one at the event scale and at both sites. It makes no doubt that for this particular dust episode and for the coarse-mode dust particles the longwave DRE is as important as the shortwave one. Not having it taken into account would have led to large errors in the final estimation of the total dust DRE and to an overestimation of the dust cooling effect. Because Dc DRE_{SW} and DRE_{LW} nearly compensate, the dust coarse mode net contribution to the total dust DRE_{NET} is overall small (Dc $|DRE_{NET}| < 0.5$ W m⁻²), independently of the atmospheric level. Finally, as far as DD DRE_{NET} is concerned, most of the DD DRE_{NET} instantaneous values at SRF and TOA are negative and remain lower than 20 W m⁻² in absolute value. The daily means are -3.0 (BCN) and -1.40 (LPZ) W m⁻² at SRF and -2.3 (BCN) and -0.35 (LPZ) W m⁻² at TOA. At SRF the highest daily DD DRE_{NET} are -7.4 and -1.5 W m⁻² in BCN and LPZ, respectively. In one occasion in BCN (on 28 June) the daily DD DRE_{NET} at SRF takes a positive value +0.9 W m⁻², meaning that on a daily basis the dust radiative effect produces a net warming at SRF. This is quite an unusual results. Our values are



310 significantly smaller (in absolute value) than the limited number of dust, daily DRE_{NET} that can be found in the literature. For instance, Di Sarra et al. (2011) found daily DD DRE_{NET} of -43.7 (SRF) and -37.6 $W\ m^{-2}$ (TOA); Meloni et al. (2015) of -14.7 (SRF) and -15.5 $W\ m^{-2}$ (TOA); and Valenzuela et al. (2017) of $[-64; -37$ $W\ m^{-2}]$ (SRF) and $[-31; -18$ $W\ m^{-2}]$ (TOA). As said earlier, the most probable explanation why our values are smaller than the literature lies in the surface temperature which is



315 **Figure 7.** Longwave instantaneous dust direct radiative effect (DRE_{LW}) at SRF, TOA and in the atmosphere (ATM) in (a) BCN and (b) LPZ. (c) Df/DD DRE ratio of the instantaneous DRE_{LW} at the SRF and TOA in BCN; the mean values and best linear fits have been calculated between 24 and 30 June. The absolute decrease of Df/DD DRE_{LW} ratio at the SRF (TOA) is -0.2 (-0.1) $\% day^{-1}$.

320 The objective of this last paragraph is to complete for the net radiative effect the discussion about the diurnal cycle of DRE_{SW} made in the companion paper by Córdoba-Jabonero et al. (2021). Figure 10 shows the diurnal cycle of DRE_{NET} for Dc, Df and DD at SRF, TOA and in the atmosphere for 26 June. Dc and DD DRE_{SW} are also reproduced. Df DRE_{SW} is not reproduced since it is nearly equal to Df DRE_{NET} (Df DRE_{LW} is small). We find again that the diurnal cycle of the Dc and DD DRE_{NET} is singular at SRF and TOA with the shape of a “W”, showing two minima, one in the morning (around 6 UTC) and one in the afternoon (17-18 UTC), and a maximum at central hours of the day. This shape is the one of the diurnal cycle of DRE_{SW} (the diurnal cycle of DRE_{LW} is rather flat on 26 June) which is due to a combination of solar geometry and dust anisotropic scattering (Osborne et al., 2011; Osipov et al., 2015). We refer to Córdoba-Jabonero et al. (2021) for a detailed explanation of the “W” shape of the diurnal cycle of DRE_{SW} . At SRF and TOA both Dc and DD DRE_{NET} are similar to DRE_{SW} shifted towards positive values (because of the Dc LW offset). The cooling effect produced during the day by DRE_{SW}

325



Table 1: Daily DRE_{LW} and DRE_{NET} ($W m^{-2}$) and $DREff$ ($W m^{-2} \tau^{-1}$) at SRF produced by Dc, Df and DD particles in BCN and LPZ.

\bar{X} indicates daily-averages. The daily DOD at 532 nm, DOD^{532} , is also included. Df and DD DRE_{LW} and the 29-30 June mean DRE in LPZ are given with a precision of 2 digits.


June 2019	BCN										LPZ (*)		
	23	24	25	26	27	28	29	30	23-30	29	30	29-30	
DOD^{532}	\bar{D}_c	0.055	0.189	0.098	0.193	0.140	0.090	0.092	0.071	0.116	0.028	0.020	0.024
	\bar{D}_f	0.019	0.041	0.031	0.062	0.045	0.029	0.039	0.031	0.037	0.016	0.013	0.015
	$\bar{D}\bar{D}$	0.074	0.230	0.129	0.255	0.185	0.119	0.131	0.102	0.153	0.044	0.033	0.039
DRE	$LW \bar{D}_c$	+1.8	+6.9	+3.7	+7.8	+9.8	+7.5	+5.6	+4.0	+5.9	+1.3	+0.5	+0.90
	$LW \bar{D}_f$	+0.11	+0.16	+0.14	+0.35	+0.42	+0.24	+0.26	+0.14	+0.22	+0.08	+0.05	+0.06
	$LW \bar{D}\bar{D}$	+1.91	+7.06	+3.84	+8.15	+10.22	+7.74	+5.86	+4.14	+6.12	+1.38	+0.55	+0.96
	$NET \bar{D}_c$	-0.2	-2.7	-2.0	-1.7	+3.0	+3.1	+1.2	+0.6	+0.2	-0.2	-0.3	-0.25
	$NET \bar{D}_f$	-1.6	-3.6	-3.0	-5.7	-3.7	-2.2	-3.1	-2.6	-3.2	-1.3	-1.0	-1.15
	$NET \bar{D}\bar{D}$	-1.8	-6.3	-5.0	-7.4	-0.7	+0.9	-1.9	-2.0	-3.0	-1.5	-1.3	-1.40
DREff	$LW \bar{D}_c$	+39.6	+34.6	+30.5	+40.2	+64.8	+71.6	+54.6	+48.7	+44.3	+46.2	+32.8	+41.3
	$LW \bar{D}_f$	+5.0	+3.9	+3.7	+5.0	+8.1	+7.1	+5.8	+3.8	+5.3	+4.7	+4.1	+4.4
	$LW \bar{D}\bar{D}$	+28.5	+29.0	+23.9	+30.7	+50.2	+55.6	+39.8	+34.8	+34.9	+37.1	+26.0	+28.2
	$NET \bar{D}_c$	-33.8	-40.6	-55.4	-38.3	-7.7	+5.1	-12.2	-13.5	-30.8	-39.9	-62.8	-48.2
	$NET \bar{D}_f$	-118.0	-141.6	-140.2	-128.5	-116.7	-103.9	-113.3	-108.5	-124.4	-144.1	-163.5	-153.5
	$NET \bar{D}\bar{D}$	-60.8	-58.9	-76.3	-62.7	-36.0	-21.9	-42.8	-42.9	-54.1	-74.0	-101.8	-85.3

330 (*) 29 and 30 June represent the dust Episode 1 and Episode 2, respectively, as observed in LPZ.

Table 2: Same as Table 1 but at the TOA. DOD^{532} values are shown in Table 1.

June 2019	BCN										LPZ (*)		
	23	24	25	26	27	28	29	30	23-30	29	30	29-30	
DRE	$LW \bar{D}_c$	+1.0	+3.7	+3.2	+6.1	+4.6	+3.2	+2.7	+2.6	+3.4	+1.4	+0.6	+1.0
	$LW \bar{D}_f$	+0.05	+0.05	+0.06	+0.22	+0.16	+0.10	+0.08	+0.06	+0.10	+0.05	+0.03	+0.04
	$LW \bar{D}\bar{D}$	+1.05	+3.75	+3.26	+6.32	+4.76	+3.30	+2.78	+2.66	+3.50	+1.45	+0.63	+1.04
	$NET \bar{D}_c$	-0.3	-2.2	-0.3	+0.3	+0.9	+1.0	+0.6	+1.1	+0.2	+0.7	+0.3	+0.50
	$NET \bar{D}_f$	-1.3	-3.2	-2.4	-4.6	-2.9	-1.6	-2.3	-1.7	-2.5	-1.0	-0.7	-0.85
	$NET \bar{D}\bar{D}$	-1.6	-5.4	-2.7	-4.3	-2.0	-0.6	-1.7	-0.6	-2.3	-0.3	-0.4	-0.35
DREff	$LW \bar{D}_c$	+23.4	+17.7	+26.9	+30.9	+31.2	+31.8	+28.4	+31.3	+26.5	+52.9	+40.7	+48.5
	$LW \bar{D}_f$	+2.0	+1.1	+1.9	+3.2	+3.2	+2.7	+2.1	+1.8	+2.4	+3.7	+2.8	+3.3
	$LW \bar{D}\bar{D}$	+16.5	+14.6	+20.7	+23.5	+24.2	+24.8	+20.4	+22.2	+20.8	+37.1	+26.0	+32.6
	$NET \bar{D}_c$	-29.9	-32.3	-29.8	-15.6	-7.4	-0.9	+0.1	+10.8	-17.4	+1.0	-5.7	-1.4
	$NET \bar{D}_f$	-95.7	-133.1	-117.7	-99.9	-83.1	-73.1	-76.7	-66.8	-96.2	-108.2	-119.5	-113.6
	$NET \bar{D}\bar{D}$	-51.3	-49.8	-52.1	-38.6	-26.9	-18.8	-23.1	-13.2	-37.3	-106.3	-125.6	-114.2

(*) 29 and 30 June represent the dust Episode 1 and Episode 2, respectively, as observed in LPZ.



335 **Figure 8.** Daily LW/SW DRE ratio in BCN at the SRF, TOA and in the atmosphere (ATM) for (top) Dc (middle) Df and (bottom) DD.





Figure 9. Net instantaneous dust direct radiative effect (DRE_{NET}) at SRF, TOA and in the atmosphere (ATM) in (a) BCN and (b) LPZ.

340 is thus partly counterbalanced by DRE_{LW} . At TOA during the central hours (11-13 UTC), when Dc DRE_{SW} reaches its minimum, DD even produces a quasi-neutral effect ($-0.7 < DD DRE_{NET} < -0.2 \text{ W m}^{-2}$). At SRF, the instantaneous DD DRE_{NET} is negative and it is larger (in absolute value) than 12 W m^{-2} ; the daily DD DRE_{NET} (Table 1) is equal to -7.4 W m^{-2} . As expected, when considering a complete daily cycle (24 hours) instead of instantaneous values, the LW DRE plays a larger role with respect to the SW DRE because the latter is zero during the night hours. The diurnal DRE variations at TOA are larger than at SRF. As a consequence, DRE_{NET} in the atmosphere shows the shape of an inverted U for both Dc and DD. The dust (all modes) produces a heating of the atmosphere during most of the hours of the day (DRE_{NET} at TOA $>$ DRE_{NET} at SRF) and a slight cooling at dawn/dusk. The effect of DRE_{LW} on the atmospheric DRE_{NET} is rather small: the difference for Dc and DD between the atmospheric DRE_{SW} and DRE_{NET} is not greater than 2 W m^{-2} .



350 **Figure 10.** Diurnal cycle on 26 June in BCN of the dust net radiative effect (DRE_{NET}) for Dc, Df and DD (bottom) at SRF, (middle) in the atmosphere, and (top) at TOA. For Dc (red) and DD (black) the diurnal cycle of DRE_{SW} is also reported by dash lines. Df DRE_{SW} is not represented since it would overlap with Df DRE_{NET} .


3.1.2 Dust direct radiative efficiency

The dust direct radiative efficiency (DREff) is assessed on a daily basis (and at the event scale) by calculating the best linear fit forced to 0 of the scatterplot of instantaneous values of one full day (and of the whole dust episode) of DRE vs. DOD^{532} .
 355 First the longwave dust direct radiative efficiency is discussed at SRF and TOA: see Figure 11 showing all instantaneous DRE_{LW} vs. DOD^{532} at SRF and TOA separately for Dc/Df and BCN/LPZ, and Table 1 and Table 2 reporting the daily LW DREff values; and then the net direct radiative efficiency is discussed compared to the SW and LW components separately for Dc, Df and DD and at SRF, TOA and in the atmosphere: see Figure 12 showing only the linear fits (for the sake of clarity) of
 360 SW, LW and net DRE components vs. DOD^{532} , and Table 1 and Table 2 reporting the daily net DREff values. As expected Df LW DREff is small at SRF and TOA and at both sites (Df LW DREff $< +5.3 \text{ W m}^{-2} \tau^{-1}$). At the event scale at SRF Dc LW DREff in BCN ($+44.3 \text{ W m}^{-2} \tau^{-1}$) and in LPZ ($+41.3 \text{ W m}^{-2} \tau^{-1}$) are similar, while at TOA they differ significantly ($+26.5$ and $+48.5 \text{ W m}^{-2} \tau^{-1}$, respectively). There are two reasons for that difference: 1) in average the dust coarse particles are larger in LPZ than in BCN (see Section 2.1 and Figure 1) inducing larger extinction coefficients in the longwave spectral range (Figure
 365 3a and c) and 2) the center of mass of the dust plume is much higher in LPZ than in BCN (Figure 5) which produces higher LW DRE at TOA (Dufresne et al., 2002; Sicard et al., 2014a). According to Sicard et al. (2014a), when maintaining the aerosol optical depth constant, Dc DRE_{LW} does not vary much for $r_g(Dc) > 1 \mu\text{m}$. This result lets us think that the main reason why Dc LW DREff at TOA in LPZ is higher than in BCN is related with the height of the dust plume. When Df and Dc are considered together, DD LW DREff drops down (with respect to the Dc value) to $+34.9$ (BCN) $+28.2 \text{ W m}^{-2} \tau^{-1}$ (LPZ) at SRF
 370 and to $+20.8$ (BCN) and $+32.6 \text{ W m}^{-2} \tau^{-1}$ (LPZ) at TOA.

To study the effects of each particle size mode (Dc and Df) and spectral components (SW and LW) we now look at Figure 12 which has been produced with the data from BCN for the whole event (23-30 June). The central plots show DRE vs. DOD^{532} for the dust fine mode. Df LW DREff is small at all atmospheric levels which leads to similar values of Df SW and net DREff (black and red lines almost overlap in Figure 12b, e and h). This allows to conclude that the dust Df net DREff is
 375 driven basically by the SW component. Things are different for the dust coarse mode (left plots in Figure 12) for which the LW component offsets significantly the SW one. Indeed, at SRF and TOA, we observe a decrease of the net Dc radiative efficiency with respect to the SW one of a factor 2.5: -30.8 (net) vs. $-75.2 \text{ W m}^{-2} \tau^{-1}$ (SW) at SRF, and -17.4 (net) vs. $-43.9 \text{ W m}^{-2} \tau^{-1}$ (SW) at TOA. In the atmosphere (Figure 12g), the inclusion of the LW component reduces the Dc net DREff of a factor




1.6 with respect to the SW one. When both size modes are added together (DD, right plots of Figure 12), a reduction of the
 380 SW radiative efficiency is also observed when the LW component is taken into account. At SRF and TOA, we observe a decrease of the net DD radiative efficiency with respect to the SW one of a factor 1.6: -54.1 (net) vs. $-88.9 \text{ W m}^{-2} \tau^{-1}$ (SW) at SRF, and -37.3 (net) vs. $-58.0 \text{ W m}^{-2} \tau^{-1}$ (SW) at TOA. Interestingly, the ratio of the total dust SW-to-NET radiative efficiency is the same (1.6) at both SRF and TOA. It is a value significantly larger than 1 which highlights again the importance of considering the longwave component in studies focused on dust radiative effects. Finally, in the atmosphere (Figure 12i), the
 385 inclusion of the LW component reduces the DD net DREff of a factor 1.8 with respect to the SW one.



390 **Figure 11. Dust longwave direct radiative effect (DRE_{LW}) at (a) SRF and (b) TOA as a function of DOD^{532} , as shown separately for Dc (circles) and Df (crosses) at BCN (23-30 June) and LPZ (29-30 June). Corresponding DREff values (slope of the linear fitting of DRE_{LW} vs. DOD^{532}) are included in the legend (solid and dashed lines, respectively, for Dc and Df particles).**

3.2 Heatwave and dust cooling/warming effect of the Earth-Atmosphere system

How the net direct radiative effect of mineral dust is modified when the dust intrusion occurs simultaneously with a heatwave? This question is the motivation of this section which focuses on the results in BCN. We will start with some features of the
 395 June 2019 heatwave. Data provided by the Copernicus Climate Change Service show that the European-average temperature for June 2019 was higher than for any other month of June on record (Copernicus, 2021). Average temperatures were more than 2°C above normal and if we consider the 5-day period 25-29 June the temperatures were 6 to 10°C above normal, with local differences even higher (up to 18°C !) in NE Spain, France and the United Kingdom according to a detailed article issued by the Spanish state meteorological agency (AEMET) on the “June 2019 heatwave in the context of the climate crisis”
 400 (AEMET, 2019). In June 2019, the synoptic conditions were marked by the presence of an anomalous long-lasting anticyclone in the upper troposphere which advected warm air from the Sahel and Mediterranean region and enhanced incoming solar radiation and surface turbulent fluxes. According to Xu et al. (2020a) this situation is at the origin of three heatwaves over



405 **Figure 12.** Linear fitting of the dust direct radiative effect (SW, LW and net) as a function of DOD^{532} at SRF for (a) Dc, (b) Df and (c) DD; at TOA for (d) Dc, (e) Df and (f) DD; and in the atmosphere for (g) Dc, (h) Df and (i) DD, in BCN (23-30 June). Corresponding DREff values (slope of the linear fitting of DRE vs. DOD^{532}) are included next to the right axis. DREff units are $W m^{-2} \tau^{-1}$. In Dc and DD plots the arrows and number indicate the SW / NET ratio. The legend in plot (a) applies to all plots.

Europe and it is during the last one, from 25 to 29 June, that the highest temperatures were reached. Towards the end of June, this anticyclone merged with a high-pressure system located over northeastern Europe that had previously produced extreme temperatures, and extended towards the northeast as a strong subtropical ridge intensified by a low-pressure system located over the eastern Atlantic. This subtropical ridge injected very warm air and mineral dust from the Sahara region towards western Europe, and this air overheated during its transport while traveling over previously warmed land (Sousa et al., 2019; Xu et al., 2020a). Sousa et al. (2019) classified the heatwave of 25-29 June 2019 as a mega-heatwave (Barriopedro et al., 2011) because of its outstanding duration, intensity and spatial extent. Other climate studies involving multi-model methodologies have come to the conclusion that the June 2019 heatwave had been most probably triggered by anthropogenic climate change, and that such heatwaves could become more widespread, long-lasting, and severe over Europe in the future (Ma et al., 2020; Vautard et al., 2020).

As seen in the previous section, the total dust net radiative effect is modulated by the dust coarse mode LW radiative effect. Figure 13a shows on the same plot the daily Dc LW/SW DRE ratio and the daily Df and DD DRE_{NET} at SRF for the BCN site and for the period 23-30 June; Figure 13b shows the same magnitudes at TOA. In Figure 13c the daily LST is reported. A clear correlation exists between the LST and the Dc LW/SW DRE ratio at both SRF and TOA. In the same way a clear correlation exists between Dc LW/SW DRE ratio and DD DRE_{NET} at SRF and also at TOA. 26 June is a pivotal moment of the episode in terms of dust radiative effect: during that day the TOA (first) and SRF (second) Dc LW/SW DRE ratio cross the 100 % threshold (Dc $|DRE_{SW}| = Dc |DRE_{LW}|$). This affects directly DD DRE_{NET} with respect to Df DRE_{NET} : DD



$|DRE_{NET}| > Df |DRE_{NET}|$ before 26 June (Dc DRE_{NET} amplifies the dust fine mode cooling effect) and $DD |DRE_{NET}| < Df |DRE_{NET}|$ after 26 June (Dc DRE_{NET} reduces the dust fine mode cooling effect). As expected the difference between Df and DD DRE_{NET} is larger at SRF than TOA which reflects that the contribution of Dc DRE_{NET} is more significant at SRF, where most of the LW radiation comes from, than at TOA. When Dc LW/SW DRE ratio reaches values above 150 % at SRF, DD DRE_{NET} takes values close to zero (-0.7 and +0.9 W m⁻² on 27 and 28 June, respectively), i.e. the total dust net radiative effect is quasi neutral at this moment of the episode.


The aim of this last paragraph is to estimate the impact of the heatwave on the dust net radiative effect. For that, two parameters directly related to the heatwave and needed for the computation of the LW radiative effect are considered: the surface temperature and the profile of the air temperature. In the following we calculate Dc DRE_{LW} (the variations produced on Df DRE_{LW} are neglected since Df DRE_{LW} is small) in a controlled way assuming the following three scenarios:

1. Heatwave-1 (HW1): This study with a climatological LST from the 8 years 2011-2018.
2. Heatwave-2 (HW2): This study with an air temperature profile in the dust layer equal to the actual one minus 6 °C.
3. Heatwave-3 (HW3): This study with a climatological LST from the 8 years 2011-2018, and an air temperature profile in the dust layer equal to the actual one minus 6 °C (equivalent to a combination of HW1+HW2 scenarios).

The climatological LST was calculated as the average over the 8 years preceding 2019 (2011-2018). The data were provided by the Copernicus Global Land Service in the way than the hourly LST of 23 – 30 June, 2019 (see Section 2.3). Although the analysis was originally planned for a 10-year climatology, Copernicus data of the LST are available only back to 2011. In HW1 and HW3 the LST has been replaced by the 8-year climatological value. According to AEMET (2019), the temperature anomaly at 850 hPa on 28 June, 2019, in the region of Barcelona was on the order of +12 °C. This is probably the maximum of the episode (see Figure 6 and Figure 13c). As a rule of thumb the mean anomaly over the episode is estimated to be +6 °C. For comparison the difference between the maximum daily LST (32.7°C reached on 28 June, Figure 6) and the mean over the episode (28.8°C) is 3.9°C. In HW2 and HW3 6°C have been subtracted to the temperature profiles in the dust layer. The different scenarios allow to estimate how the dust radiative effect would have been modified: HW1, if the surface temperature had not been so high; HW2, if the air had not been so warm; and HW3, if the dust episode had not been accompanied by a heatwave (“normal” surface and air temperature). Figure 14a and b show the daily DD DRE_{NET} and the daily Dc LW/SW DRE ratio at SRF and TOA, respectively, for the three scenarios as well as in the heatwave conditions (this study). Figure 14c shows the daily LST (2019) and the climatological LST (2011-2018) for comparison. From the latter figure, one sees that at the beginning of the episode (23 and 24 June) the 2019 daily LST is indeed lower than the climatological one. From 24 June onwards it increases strongly. On the last three days of the episode (28, 29 and 30 June) the 2019 daily LST is approximately 4.0 – 4.5 °C above the climatological LST. In HW1 (red lines in Figure 14a and b) the use of the climatological LST has no impact on the Dc LW/SW DRE ratio at SRF (and thus on the DD DRE_{NET} either). Contrarily the impact at the TOA is quite noticeable: towards the end of the episode, the reduction of LST in HW1 yields logically a reduction in Dc DRE_{LW} , thus a reduction in the Dc LW/SW DRE ratio and thus an amplification of the dust cooling effect at TOA. Say the other around, the effect of a high LST during the heatwave is to increase the amount of radiation which escapes to space. Since the radiation budget at SRF is nearly unchanged, the effect in the atmosphere is to reduce the heating of the atmosphere. In HW2 (blue lines in Figure 14a and b) the use of a decreased air temperature in the dust layer has an impact at both SRF and TOA and of opposite sign. The decrease of the air temperature has a direct impact on the gaseous transmittance. According to Dubuisson et al. (2004) and following the correlated k-distribution (Lacis and Oinas, 1991) used in GAME, a decrease in air temperature yields a decrease in the absorption coefficient. This will result in more LW radiation propagating upward in the atmosphere and thus in a reduction of the dust Dc DRE_{LW} in the dust layers and consequently at SRF (Figure 14a), and in an increase of the LW radiation reaching TOA (Figure 14b). Consequently, when the air temperature in the dust layer is reduced by 6 °C, DD DRE_{NET} is shifted towards negative values at SRF and towards positive values at TOA. Seen from the heatwave perspective, the effect of high temperatures in the dust layer during the heatwave is to reduce the dust net cooling effect at SRF and to amplify it at



TOA (decrease of the amount of radiation which escapes to space). These opposite variations will have the effect of reducing
 470 the atmospheric heating provoked by the dust particles. Finally, when both HW1 and HW2 scenarios are considered
 simultaneously (HW3), their effect on the daily DD DRE_{NET} and the daily Dc LW/SW DRE ratio combine (green lines in
 Figure 14a and b). At SRF DD DRE_{NET} is shifted towards negative values like in HW2 (in HW1 DD DRE_{NET} was nearly
 unchanged). At TOA HW1 and HW2 have opposite effects on DD DRE_{NET} in the second half of the episode when the 2019
 LST is larger than the climatological LST; consequently their effects compensate and DD DRE_{NET} in HW3 is practically
 475 unchanged with respect to the results of this study. Interestingly, in the first half of the episode when the 2019 LST is smaller
 than the climatological LST, both HW1 and HW2 produce a shift of DD DRE_{NET} at TOA towards positive values which result
 in a reduction of the dust net cooling effect in HW3. From the heatwave perspective, we conclude that the higher LST and the
 higher air temperature due to the heatwave that accompanied the dust episode 1) provoked a reduction of the dust net cooling
 effect at the surface, 2) left unchanged the dust net cooling effect at TOA, and 3) consequently reduced the dust net heating of
 480 the atmosphere.



485 **Figure 13.** (a) Daily Df and DD DRE_{NET} (left axis) and Daily Dc LW/SW DRE ratio (right axis) at SRF; (b) Daily Df and DD
 DRE_{NET} (left axis) and Daily Dc LW/SW DRE ratio (right axis) at TOA; (c) Daily mean LST. Site: BCN; Time period: 23-30 June.
 The shaded area represents the period during which both SRF and TOA Dc LW/SW DRE ratio cross the 100 % threshold.

4 Conclusions

This companion paper of Córdoba-Jabonero et al. (2021) estimates the temporal variation of the instantaneous and daily dust
 490 longwave and net direct radiative effect during an intense dust episode that occurred between 23 and 30 June, 2019, and
 coincided with a mega-heatwave. It also investigates the effect of the heatwave on the dust radiative effect. The radiative effect
 was calculated with the GAME radiative transfer model separately for the fine- and coarse-mode dust. The dust radiative
 properties in the longwave spectral range were calculated with a Mie code and particle microphysics from AERONET. The
 dust fine- and coarse-mode vertical distribution as well as the shortwave DRE were taken from the companion paper. Two



495




Figure 14. Daily DD DRE_{NET} (solid lines, left axis) and Daily Dc LW/SW DRE ratio (dotted lines, right axis) at (a) SRF and (b) TOA; (c) Daily mean LST. Site: BCN; Time period: 23-30 June. HW1: climatological LST; HW2: $T(\text{air})-6^\circ\text{C}$ in the dust layer; HW3: climatological LST and $T(\text{air})-6^\circ\text{C}$ in the dust layer. In the LST plot, the climatological surface temperature is also plotted.

500 sites are considered: Barcelona, Spain, (23-30 June) and Leipzig, Germany, (29-30 June) to estimate the impact of the dust transport on its DRE.

Independently of the atmospheric level where it is estimated, the instantaneous Df DRE_{LW} are low and do not exceed $+0.6\text{ W m}^{-2}$ in BCN and $+0.2\text{ W m}^{-2}$ in LPZ. The respective daily values do not exceed $+0.42$ and $+0.08\text{ W m}^{-2}$. In average over the whole event the contribution of Df daily DRE_{LW} to the total dust DRE_{LW} is not higher than 4 %. Most of the dust
 505 DRE_{LW} is thus produced by the coarse-mode particles. Instantaneous Dc DRE_{LW} at SRF in BCN reaches a maximum of $+11.4\text{ W m}^{-2}$ and the daily values a maximum of $+9.8\text{ W m}^{-2}$. In LPZ the instantaneous and daily values reach maxima at $+3.3$ and $+1.3\text{ W m}^{-2}$, respectively. Dc DRE_{LW} at TOA is smaller than at SRF which produces a negative (cooling) effect in the atmosphere. Df DRE_{LW} in BCN represents 6.7 (3.7) % at SRF (TOA) of its SW counterpart. The Dc LW/SW DRE ratio in BCN increases from $\sim 80\%$ at the beginning of the episode up to 170 % towards the end of it. Such an unusual tendency is attributed to increasing coarse-mode size and surface temperature along the episode. In the last four days of the episode Dc
 510 DRE_{LW} is larger than Dc DRE_{SW} (Dc LW/SW DRE ratio $> 100\%$) at both SRF and TOA. This is a singular result of this study which has the effect of reducing considerably the SW cooling.

The results of the dust net DRE are discussed in terms of daily values which are the magnitude that matters for assessing the effect of aerosols on the Earth-Atmospheric radiative budget. The fine mode DRE_{NET} is similar to DRE_{SW} since
 515 $DRE_{LW} \ll DRE_{SW}$. In BCN Dc DRE_{NET} at SRF is negative at the beginning of the event and positive afterwards (peak at $+3.1\text{ W m}^{-2}$). In LPZ Dc DRE_{NET} is slightly negative (mean of -0.25 W m^{-2}). At TOA the same tendency is observed in BCN (peak at $+3.1\text{ W m}^{-2}$). But in LPZ Dc DRE_{NET} is also positive at TOA (mean of $+0.50\text{ W m}^{-2}$). This stronger effect of the dust DRE_{LW} at TOA (vs. SRF) in LPZ is related with the quite high dust plume in LPZ ($> 3.3\text{ km}$). Overall, the total dust net DRE is small: -3.0 (-2.3) W m^{-2} at SRF (TOA) in BCN and -1.40 (-0.35) W m^{-2} in LPZ. On one occasion at SRF in BCN DD



520 DRE_{NET} is even positive (+0.9 W m⁻² on 28 June) indicating a total dust net warming at the surface, which contrasts with the “traditional” dust cooling effect usually observed in clear sky conditions.

As far as dust direct radiative efficiency is concerned, Df LW DREff is small so that Df net DREff is nearly equal to Df SW DREff. Dc LW DREff is positive which contributes to decrease significantly the net DREff with respect to the SW one. Overall, the total dust net DREff is: -54.1 (-37.3) W m⁻² τ⁻¹ at SRF (TOA) in BCN and -85.3 (-114.2) W m⁻² τ⁻¹ in LPZ. The higher dust direct radiative efficiency in LPZ (vs. BCN) is due to the SW component, already discussed in the companion paper. At TOA the difference between of DD net DREff between both sites is caused by the larger compensation of the LW component in BCN (vs. LPZ). Finally it should be noted that, in BCN at the event scale and for the total dust, the inclusion of the LW DRE in the calculation of the total net DREff reduces a factor 1.6 the SW DREff at SRF and BOA and a factor 1.8 in the atmosphere. These reduction factors significantly larger than 1 highlight the importance of considering the longwave component in studies focused on mineral dust radiative effects.

In order to evaluate the impact of the heatwave that accompanied the dust intrusion on the dust radiative effect, we perform a sensitivity study on the surface temperature and the air temperature in the dust layer, both linked to the heatwave and upon which the LW DRE strongly depends. Three scenarios are considered: 1) the LST is set to climatological values, 2) the air temperature in the dust layer is reduced of 6 °C, and 3) a combination of the first two scenarios. Our findings show that the increase of LST and air temperature in the dust layer caused by the heatwave 1) provoked a reduction of the dust net cooling effect at the surface, 2) left unchanged the dust net cooling effect at TOA, and 3) consequently reduced the dust net heating of the atmosphere. The situation at the surface is a vicious circle: the heatwave reduces the dust cooling effect which, in turn, may increase some critical variables associated to the heatwave (e.g., LST and air temperature). The effect of the heatwave on the dust radiative effect is reverse as it contributes to cool down the atmosphere. Since recent studies have warned that mega-heatwaves such as the one studied in this work might become more frequent in the future, the novel results presented in this paper call for more research on the subject.

Data availability. Part of the data used in this publication were obtained as part of the AERONET and MPLNET networks and are publicly available. For additional data or information please contact the authors.

Author Contributions. MS and CC-J designed the study and wrote the original draft paper. CC-J, MS and AA provided data. CC-J, MS and MALC performed data analysis with contributions from AA, AC, M-PZ, AR-G- and CM-P. All authors reviewed and edited the final version of the manuscript. All the authors agreed to the final version of the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgments. The MPLNET project is funded by the NASA Radiation Sciences Program and Earth Observing System. The MPLNET staff at NASA GSFC is warmly acknowledged for the continuous help in keeping the P-MPL systems and the data analysis up to date. We particularly thank E.J. Welton for providing the P-MPL unit in place at the Barcelona site.

Financial support. This research was funded by the Spanish Ministry of Science, Innovation and Universities (CGL2017-90884-REDT and PRX18/00137 “Salvador de Madariaga” programme), the Spanish Ministry of Science and Innovation (PID2019-104205GB-C21 and PID2019-103886RB-I00), the H2020 programme from the European Union (GA no. 654109, 778349, 871115 and 101008004), and both Units of Excellence “María de Maeztu” (MDM-2016-0600 and MDM-2017-0737) financed by the Spanish State Research Agency (AEI). MALC is supported by the INTA predoctoral contract programme.



References

- Abdelkader, M., Metzger, S., Mamouri, R. E., Astitha, M., Barrie, L., Levin, Z., and Lelieveld, J.: Dust–air pollution dynamics over the eastern Mediterranean, *Atmos. Chem. Phys.*, 15, 9173–9189, <https://doi.org/10.5194/acp-15-9173-2015>, 2015.
- 560 AEMET: June 2019 heatwave in the context of the climate crisis, AEMET blog, <https://aemetblog.es/2019/07/02/analisis-de-la-ola-de-calor-de-junio-de-2019-en-un-contexto-de-crisis-climatica/>, last access: 27 April, 2021.
- Athanasopoulou, E., Protonotariou, V., Papangelis, G., Tombrou, M., Mihalopoulos, N., and Gerasopoulos, E.: Long-range transport of Saharan dust and chemical transformations over the Eastern Mediterranean, *Atmospheric Environment*, 140, 592–604, <https://doi.org/10.1016/j.atmosenv.2016.06.041>, 2016.
- 565 Barragan, R., Romano, S., Sicard, M., Burlizzi, P., Perrone, M. R., and Comeron, A.: Estimation of mineral dust direct radiative forcing at the European Aerosol Research Lidar NETwork site of Lecce, Italy, during the ChArMEx/ADRIMED summer 2013 campaign: Impact of radiative transfer model spectral resolutions, *J. Geophys. Res. Atmos.*, 121, doi:10.1002/2016JD025016, 2016.
- 570 Barragan, R., Sicard, M., Totems, J., Léon, J. F., Dulac, F., Mallet, M., Pelon, J., Alados-Arboledas, L., Amodeo, A., Augustin, P., Boselli, A., Bravo-Aranda, J. A., Burlizzi, P., Chazette, P., Comerón, A., D’Amico, G., Dubuisson, P., Granados-Muñoz, M. J., Leto, G., Guerrero-Rascado, J. L., Madonna, F., Mona, L., Muñoz-Porcar, C., Pappalardo, G., Perrone, M. R., Pont, V., Rocadenbosch, F., Rodriguez-Gomez, A., Scollo, S., Spinelli, N., Titos, G., Wang, X., Zanmar Sanchez, R.: Spatio-temporal monitoring by ground-based and air- and space-borne lidars of a moderate Saharan dust event affecting southern Europe in June 2013 in the framework of the ADRIMED/ChArMEx campaign, *Air Qual Atmos Health*, doi: 10.1007/s11869-016-0447-7, 2017.
- 575 Barragan, R., Molero, F., Granados-Muñoz, M. J., Salvador, P., Pujadas, M. and Artíñano, B.: Feasibility of Ceilometers Data to Estimate Radiative Forcing Values: Application to Different Conditions around the COVID-19 Lockdown Period, *Remote Sensing*, 12(22), 3699, doi:10.3390/rs12223699, 2020.
- Barriopedro, D., Fisher, E., Luterbacher, J., Trigo, R.M., and García-Herrera, R.: The hot summer of 2010: redrawing the temperature record map of Europe. *Science* 322, 220. <https://doi.org/10.1126/science.1201224>, 2011.
- 580 Basu, R., and Samet, J.M.: Relation between elevated ambient temperature and mortality: a review of the epidemiologic evidence, *Epidemiol. Rev.*, 24, 190–202, 2002.
- Copernicus, Climate Change Service, <https://climate.copernicus.eu/record-breaking-temperatures-june>, last access: 27 April, 2021.
- 585 Chung, E.-S., Sohn, B.-J., and Schmetz, J.: Diurnal variation of outgoing longwave radiation associated with high cloud and UTH changes from Meteosat-5 measurements, *Meteorol. Atmos. Phys.*, 105, 109–119, doi: 10.1007/s00703-009-0041-8, 2009.
- Córdoba-Jabonero, C., Sicard, M., López-Cayuela, M.-Á., Ansmann, A., Comerón, A., Zorzano, M.-P., Rodríguez-Gómez, A., and Muñoz-Porcar, C.: Aerosol radiative impact during the summer 2019 heatwave produced partly by an inter-continental Saharan dust outbreak – Part 1: Short-wave dust direct radiative effect, *Atmos. Chem. Phys.*, 21, 6455–6479, <https://doi.org/10.5194/acp-21-6455-2021>, 2021.
- 590 Díaz, J., Carmona, R., Mirón, I.J., Ortiz, C., León, I., and Linares, C.: Geographical variation in relative risks associated with heat: update of Spain's heat wave prevention plan, *Environ. Int.*, 85, 273–283, 2015.
- di Sarra, A., Di Biagio, C., Meloni, D., Monteleone, F., Pace, G., Pugnaghi, S., and Sferlazzo, D.: Shortwave and longwave radiative effects of the intense Saharan dust event of 25–26 March, 2010, at Lampedusa (Mediterranean sea), *J. Geophys. Res.* 116, D23209, doi:10.1029/2011JD016238, 2011.
- 595 Dubuisson, P., Buriez, J. C., and Fouquart, Y.: High Spectral Resolution Solar Radiative Transfer in Absorbing and Scattering Media: Application to the Satellite Simulation, *J. Quant. Spectrosc. Ra.*, 55, 103–126, [https://doi.org/10.1016/0022-4073\(95\)00134-4](https://doi.org/10.1016/0022-4073(95)00134-4), 1996.



- Dubuisson, P.; Dessailly, D.; Vesperini, M.; Frouin, R.: Water Vapor Retrieval Over Ocean using Near-infrared Radiometry, *J. Geophys. Res. Atmos.*, 109, D19106, <https://doi.org/10.1029/2004JD004516>, 2004.
- 600 Dubuisson, P., Roger, J., Mallet, M., and Dubovik, O.: A Code to Compute the Direct Solar Radiative Forcing: Application to Anthropogenic Aerosols during the Escompte Experiment, in: Proc. International Radiation Symposium (IRS 2004) on Current Problems in Atmospheric Radiation, 23–28 August 2004, Busan, Korea, edited by: Fischer, H. and Sohn, B.-J., A. Deepak Publishing, Hampton, 127–130, 2006.
- 605 Dufresne, J., Gautier, C., and Ricchiuzzi, P.: Longwave scattering effects of mineral aerosols, *J. Atmos. Sci.* 59, 1959–1966, 2002.
- Freitas, S. C., Trigo, I., Macedo, J., Barroso, C., Silva, R., and Perdigao, R.: Land Surface Temperature from multiple geostationary satellites. *International Journal of Remote Sensing*, 34, 3051–3068, 2013.
- Granados-Muñoz, M. J., Sicard, M., Román, R., Benavent-Oltra, J. A., Barragán, R., Brogniez, G., Denjean, C., Mallet, M., 610 Formenti, P., Torres, B., and Alados-Arboledas, L.: Impact of mineral dust on shortwave and longwave radiation: evaluation of different vertically resolved parameterizations in 1-D radiative transfer computations, *Atmos. Chem. Phys.*, 19, 523–542, <https://doi.org/10.5194/acp-19-523-2019>, 2019a.
- Granados-Muñoz, M. J., Sicard, M., Papagiannopoulos, N., Barragán, R., Bravo-Aranda, J. A., and Nicolae, D.: Two-dimensional mineral dust radiative effect calculations from CALIPSO observations over Europe, *Atmos. Chem. Phys.*, 615 19, 13157–13173, <https://doi.org/10.5194/acp-19-13157-2019>, 2019b.
- Griffin, D.W.: Atmospheric movement of microorganisms in clouds of desert dust and implications for human health, *Clin. Microbiol. Rev.*, 20 (3), 459–477, 2007.
- Krekov, G. M.: Models of atmospheric aerosols, edited by: Jennings, S. G., *Aerosol Effects on Climate*, University of Arizona Press, Tucson, AZ, 9–72, 1993.
- 620 Kunst, A. E., Looman C. W., and Mackenbach, J. P.: Outdoor air temperature and mortality in the Netherlands: a time-series analysis, *Am. J. Epidemiol.*, 137, 331–341, 1993.
- Lacis, A.A., and Oinas, V.: A Description of the Correlated K Distribution Method for Modeling Nongray Gaseous Absorption, Thermal Emission, and Multiple Scattering in Vertically Inhomogeneous Atmospheres, *J. Geophys. Res. Atmos.*, 96, 9027–9063, <https://doi.org/10.1029/90JD01945>, 1991.
- 625 Ma, F., Yuan, X., Jiao, Y., and Ji, P.: Unprecedented Europe heat in June–July 2019: Risk in the historical and future context, *Geophysical Research Letters*, 47, e2020GL087809, <https://doi.org/10.1029/2020GL087809>, 2020.
- Meloni, D., Junkermann, W., Di Sarra, A., Cacciani, M., De Silvestri, L., Di Iorio, T., Estellés, V., Gómez-Amo, J. L., Pace, G., and Sferlazzo, D. M.: Altitude-resolved Shortwave and Longwave Radiative Effects of Desert Dust in the Mediterranean during the GAMARF Campaign: Indications of a Net Daily Cooling in the Dust Layer, *J. Geophys. Res.-* 630 *Atmos.*, 120, 3386–3407, <https://doi.org/10.1002/2014JD022312>, 2015.
- Osborne, S. R., Baran, A. J., Johnson, B. T., Haywood, J. M., Hesse, E., and Newman, S.: Short-wave and Long-wave Radiative Properties of Saharan Dust Aerosol, *Q. J. Roy. Meteor. Soc.*, 137, 1149–1167, <https://doi.org/10.1002/qj.771>, 2011.
- Osipov, S., Stenchikov, G., Brindley, H., and Banks, J.: Diurnal cycle of the dust instantaneous direct radiative forcing over the Arabian Peninsula, *Atmos. Chem. Phys.*, 15, 9537–9553, <https://doi.org/10.5194/acp-15-9537-2015>, 2015.
- 635 Papayannis, A., Amiridis, V., Mona, L., Tsaknakis, G., Balis, D., Bösenberg, J., Chaikovski, A., De Tomasi, F., Grigorov, I., Mattis, I., Mitev, V., Müller, D., Nickovic, S., Pérez, C., Pietruczuk, A., Pisani, G., Ravetta, F., Rizi, V., Sicard, M., Trickl, T., Wiegner, M., Gerding, M., Mamouri, R. E., D’Amico, G., and Pappalardo, G.: Systematic lidar observations of Saharan dust over Europe in the frame of EARLINET (2000–2002), *J. Geophys. Res.*, 113, D10204, doi:10.1029/2007JD009028, 2008.



- 640 Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Grill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40, 1002, <https://doi.org/10.1029/2000RG000095>, 2002.
- Querol X, Pérez N, Reche C, Ealo M, Ripoll A, Tur J, Pandolfi M, Pey J, Salvador P, Moreno T, and Alastuey A.: African dust and air quality over Spain: Is it only dust that matters? *Sci Total Environ.*, 10, 686:737-752, doi: 10.1016/j.scitotenv.2019.05.349, 2019.
- 645 Sicard, M., Bertolín, S., Mallet, M., Dubuisson, P., and Comerón, A.: Estimation of Mineral Dust Long-Wave Radiative Forcing: Sensitivity Study to Particle Properties and Application to Real Cases in the Region of Barcelona, *Atmos. Chem. Phys.*, 14, 9213-9231, <https://doi.org/10.5194/acp-14-9213-2014>, 2014a.
- Sicard, M., Bertolín, S., Muñoz, C., Rodríguez, A., Rocadenbosch, F., and Comerón, A.: Separation of aerosol fine- and coarse mode radiative properties: Effect on the mineral dust longwave, direct radiative forcing, *Geophys. Res. Lett.*, 41, 6978–6985, <https://doi.org/10.1002/2014GL060946>, 2014b.
- 650 Slingo, A., Wilderspin, R. C., and Brentnall, S. J.: Simulation of the diurnal cycle of outgoing longwave radiation with an atmospheric GCM, *Monthly Weather Review*, 115, 1451-1457, 1987.
- Sousa, P. M., Barriopedro, D., Ramos, A. M., García-Herrera, R., Espírito-Santo, F., and Trigo, R. M.: Saharan air intrusions as a relevant mechanism for Iberian heatwaves: The record breaking events of August 2018 and June 2019, *Weather and Climate Extremes*, 26, 100224, <https://doi.org/10.1016/j.wace.2019.100224>, 2019.
- 655 Stamnes, K., Tsay, S., Wiscombe, W., and Jayaweera, K.: Numerically Stable Algorithm for Discrete-Ordinate-Method Radiative Transfer in Multiple Scattering and Emitting Layered Media, *Appl. Optics*, 27, 2502–2509, <https://doi.org/10.1364/AO.27.002502>, 1988.
- 660 Tobías, A., Pérez, L., Díaz, J., Linares, C., Pey, J., Alastruey, A., and Querol, X.: Shortterm effects of particulate matter on total mortality during Saharan dust outbreaks: a Case-Crossover analysis in Madrid (Spain), *Sci. Total. Environ.*, 412–413, 386–389, 2011.
- Vautard, R., van Aalst, M., Boucher, O., Drouin, A., Haustein, K., Kreienkamp, F., van Oldenborgh, G. J., Otto, F. E. L., Ribes, A., Robin, Y., Schneider, M., Soubeyroux, J.-M., Stott, P., Seneviratne, S. I., Vogel, M. M., and Wehner, M.: Human contribution to the record-breaking June and July 2019 heatwaves in Western Europe, *Environmental Research Letters*, 15(9), 094077, 2020.
- 665 Volz, F.: Infrared optical constants of aerosols at some locations, *Appl. Optics*, 22, 23, 3690–3700, 1983.
- Xu, P., Wang, L., Liu, Y., Chen, W., and Huang, P.: The record-breaking heat wave of June 2019 in Central Europe, *Atmos Sci Lett.*, 21:e964, <https://doi.org/10.1002/asl.964>, 2020a.
- 670 Xu, W., Kuang, Y., Liang, L., He, Y., Cheng, H., Bian, Y., Tao, J., Zhang, G., Zhao, P., Ma, N., Zhao, H., Zhou, G., Su, H., Cheng, Y., Xu, X., Shao, M., and Sun, Y.: Dust-Dominated Coarse Particles as a Medium for Rapid Secondary Organic and Inorganic Aerosol Formation in Highly Polluted Air, *Environmental Science & Technology*, 54, 24, 15710-15721, doi: 10.1021/acs.est.0c07243, 2020b.
- 675 Xu, P., Wang, L., Huang, P., Chen, W.: Disentangling dynamical and thermodynamical contributions to the record-breaking heatwave over Central Europe in June 2019, *Atmospheric Research*, 252, 105446, <https://doi.org/10.1016/j.atmosres.2020.105446>, 2021.